1 Diallel Analysis for Morphological and Biochemical Traits in Tomato Cultivated

2 Under the Influence of Tomato Leaf Curl Virus

3 Prashant Kaushik[1] and Major Singh Dhaliwal[2]

[1] Instituto de Conservación y Mejora de la Agrodiversidad Valenciana, Universitat
Politècnica de València, Valencia 46022, Spain.
[2] Department of Vegetable Science, Punjab Agricultural University, Ludhiana 141004,
India.

Abstract: Eloquent information about the genetic basis of inheritance is important for any
breeding program. Therefore, a diallel study was conducted under the influence of tomato
leaf curl virus (TLCV) disease, using the eleven advanced lines of tomato. Firstly, the
information regarding percent disease index (PDI) was determined via artificial screening
with viruliferous whiteflies. Later, these lines were crossed in half diallel mating design to
produce fifty-five one-way hybrids. These hybrids and parental genotypes were
evaluated for seven morphological and three biochemical traits under open field
conditions. Using the Griffing approach (Method II and Model I) basis of inheritance of
traits were determined. Also, a Bayesian model was applied to the total yield descriptor.
Correlations data indicated that total yield was not correlated with any other trait. The
significant general combining ability (GCA) and specific combining ability (SCA)
values indicates exploitable genetic variation. The broad-sense heritability values were
larger than narrow-sense heritability, showing that selection will be efficient for the
improvement of these traits. Hybrid combinations H23, H42 and H49 can be
considered efficient for the selection of multiple traits, including yield. Overall, this study
provides a useful information regarding the genetics of important traits of tomato under
TLCV infestation.

Keywords: Tomato; TLCV; Combining ability; Resistance breeding; Hybrid
development; Diallel analysis; Griffing approach; Bayesian
27

28 1. Introduction

Tomato (*Solanum lycopersicum* L.) is among the most cultivated plants; hence, the
efforts of its genetic improvement dates back to last century and are still enduring through
traditional breeding and genomics-based approaches [1,2]. Tomatoes are well acclimatized
and breed to yield under extreme climatic conditions like drought and frost [3–5]. But,
insect pest and diseases are still big challenges for the successful production of tomatoes.
Worldwide, approximately around 146 viruses belonging to 33 different genera are reported
to infect tomato plant [6,7]. Among them, the genus *Begomovirus* causes huge economic
losses to tomato production. Belonging to this genus a DNA virus known as the tomato leaf
curl virus (TLCV) is a serious disease of tomato and its incidence, can easily result up to 90
percent yield loss to the tomato crop [8,9].
Tomato production especially of autumn season crop in Northern India and summer
season crop of Southern India is susceptible to the high incidence of tomato leaf curl virus
(TLCV) disease [10]. TLCV is transmitted by whitefly(*Bemisia tabaci* Genn.) in a
circulative and persistent manner [11]. Hitherto, in North India, tomato leaf curl New Delhi
virus a strain of TLCV reported from New Delhi region of India has an extensive
distribution and also infects several other vegetable crops e.g. Eggplant, Squash and

Pepper [12]. As a response to TLCV infection plant leaves shows symptoms like, curling
of leaf margins, shrinking and thickening of leaf surface. While the overall plant become
stunted in growth, with few and misshaped fruits[11]. Extensive efforts in the form of
phenotypic screening has been taken in order to identify resistant genotypes . In this
respect, the artificial cage inoculation using viruliferous whiteflies is the most competent
and reliable method to carry out the screening for TLCV disease. Although with artificial
screening, the plant gives a strong reaction response, than it might give under the field
conditions[13,14].

Tomatoes are a important source of nutraceuticals like vitamins (C, K, B6 etc.),
phenolic acids, and minerals (folate, manganese etc.). All of these are vital for human
health and body development [15,16]. Particularly, its fruits contain one of the important
dietary carotenoids known as lycopene, important for the prevention of chronic diseases
like breast, lung and prostate cancer [17–19]. The lycopene content of tomato varies based
on genotypes genetic makeup, cultivation environment, disease pressure, and genotype-by-
environment interactions [20]. The nutraceutical properties of tomato fruit have
industrialized the processing of tomatoes; commonly, tomatoes are processed as juice,
ketchup, paste, and sauce [21]. These biochemical aspects of the tomato fruit have become
an important goal of tomato breeding programs [22,23].

Hybrid development is a successful approach for vegetable improvement, especially
for solanaceous vegetables. Also, to chalk out a breeding strategy for successful cultivation
under TLCV infestation it is important to have information about the inheritance of traits
under the prevailing conditions. Therefore, estimation general combining ability (GCA) and
the specific combining ability (SCA) is important for genetic enhancement of the crop. But,
the amount of variation in GCA and SCA values not totally rest on gene effects besides it
also involves the gene structure of the parents involved [24]. Diallel matting design based
on general linear model framed by Griffing [25] is a popular choice and widely accepted
tool for identification of the hybrid combinations of interest in tomato and in other
members of Solanaceae [26–28]. Previously studies indicated that leaf characteristics and
foliar pubescence affect the feeding preferences of whiteflies [29,30]. Sometimes,
Griffing`s method is not adequate in case of missing data, imbalance, and outliers under the
situations where chances of bias are too high to avoid. This is especially true with
experiments carried out under disease pressure conditions [31,32].

The use of more rigorous Bayesian methods are used to overcome these limitations
[33,34]. The BayesDiallel approach uses a Markov chain Monte Carlo (MCMC) sampling
distribution which intern provides a greater flexibility, which further improves the
biological interpretability of results. Therefore, in this study, the parameter of the total yield
(kg/plant) was investigated with this Bayesian approach. Bayesian approach is not popular
among the plant breeding community because of calculation limitations and complexity of
statistics involved [35]. But, the different models based on Bayesian approach provides a
more vigorous and detailed analysis of highly variable and complex trait like yield. An
acquaintance of the genetics of important morphological traits under tomato leaf curl virus
infestation conditions will be helpful for carrying out efficient selection and breeding.

Therefore, the objectives of this study were to determine GCA, SCA, and heritability
of tomato genotypes crossed in a half diallel mating design. Further, the BayesDiallel
approach was used to provide a more comprehensive analysis of diallel data generated for
the total yield. By applying BayesDiallel approach first time on the total yield data of the
diallel cross under TLCV conditions along with Griffing's method, we aim to precisely
estimate the combining ability and the heritability estimates and to suggest a robust
approach aimed at tomato TLCV resistance breeding.

2. Materials and Methods

 2.1. Plant Material and Artificial Screening

Eleven advanced lines of tomato developed at the Punjab Agricultural University
(coordinates at 30º54'6.893" N 75º48'27.989" E), Ludhiana, India were artificially screened
for resistance to TLCV disease during 2011-12. The artificial screening of seedlings (~ 2-3
week old) was carried out by challenging 25 plants of each of the eleven genotype with
viruliferous whiteflies reared on the TLCV disease affected plant, the detailed method is
provided elsewhere [26]. The disease reaction of genotypes was scored on the scale, where
0–10% resistant (R), >10–30% moderately resistant (MR), >30-70% susceptible (S), and
>70-100% highly susceptible (HS) [36,37]. Further, the potential disease incidence (PDI)
was measured as the (number of infected plants/total number of plants) $\times$ 100. While, the
eleven genotypes were crossed in a diallel mating design during February-March of 2012
resulting in fifty-five one way F_1 hybrids. Thereafter, the first cross combination, H1 is
referred to as the first cross in the half diallel i.e. H1 (P1 $\times$ P2) and so on, until the last cross
as H55 (P10×P11) (Table 1). The 11 parental genotypes and 55 one-way hybrids were
evaluated under the whitefly infestation conditions during the August of 2012 in a
randomized complete block design. Each entry was replicated twice and each replication
accommodated 15 plants. Plant production practices were exercised as per the Package of
Practices (Anonymous) and no chemical treatment was used to control the whiteflies. Data
were recorded on 13 central plants leaving one plant on either side of the row.

114
115

Table 1. Representation of the cross or hybrid combinations (55 in total) developed as a result of half-diallel matting design.

	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11
P1		H1	H2	H3	H4	H5	H6	H7	H8	H9	H10
P2			H11	H12	H13	H14	H15	H16	H17	H18	H19
P3				H20	H21	H22	H23	H24	H25	H26	H27
P4					H28	H29	H30	H31	H32	H33	H34
P5						H35	H36	H37	H38	H39	H40
P6							H41	H42	H43	H44	H45
P7								H46	H47	H48	H49
P8									H50	H51	H52
P9										H53	H54
P10											H55
P11											

2.2. Morphological and Biochemical Data Analysis

Data for seven morphological and three biochemical traits were recorded and
inferences were made. Total yield (kg plant^{-1}) and marketable yield (kg plant^{-1}) were
determined on per plant basis i.e. by diving the harvest of all picking with the number of
plants. Fruit weight (g) was estimated as an average weight per fruit of the ten fruits sample
collected at the red ripe stage. Five fruits per replication were used to measure the
equatorial diameter, polar diameter and pericarp thickness. The polar diameter of cut fruits
was measured as the distance between the stalk end and the blossom end. Conversely, the
equatorial diameter was measured as the transversal distance of the fruit. Pericarp thickness
was measured from centre of the fruit. These fruit-based morphometric measurements were
recorded with the help of the Vernier caliper.

Three biochemical parameters estimated included dry matter, lycopene and total
soluble solids (TSS) using three samples for each replication. Each sample was constituted
of five red ripe fruits. Dry matter percentage was measured as the change of weight before
and after oven drying at 70ºC and was calculated based on the formula $100 \times$ (dry
weight / fresh weight). TSS content of fruits was determined with the help of a hand
refractometer (RA-130- KEM, Kyoto Electronics Manufacturing Co., Ltd., Kyoto, Japan).
The readings were recorded as ºBrix (0 to 32) at room temperature. Lycopene content was
determined by the method suggested elsewhere [38]. The optical densities of processed
extracts for the lycopene content were recorded at 505 nanometers (nm) using Spectronic
20 (Thermo Fisher Scientific, Waltham, Massachusetts, USA). The combining abilities
were estimated based on the Griffing's Method 2 (parental genotypes and one-way hybrids)
with Model 1 (fixed effect).[25] The diallel calculations were performed with the help of
pacakge AGD-R (analysis of genetic designs with the R software package) version 4.0 [39].
Pearson coefficients of correlation (r) and there P-values were calculated and plotted using
the packages corrplot[40]. The chart.Correlation() function within the
PerformanceAnalytics package was used to generate scatter plots and histograms along
with detailed information regarding correlations[41].

2.3. Bayesian Model-Based Analysis of Total Yield

The R package BayesDiallel was used for the Bayesian model-based analysis of total
yield data [34]. Briefly we used the "Bab" model in the BayesDillel package, this model
includes an overall effect of being inbred (B), an additive component (a), and a measure of
parent-specific inbred deviation (b). Also, the "Bab" model was applied using an MCMC
Gibbs sampler with five chains, 10,000 iterations, and burn-in of 1000 [35].

3. Results and Discussion

3.1. Artificial Screening

Artificial screening with viruliferous whiteflies provided a dissimilar percent of disease
incidence (PDI) values for each genotype (Figure 1). The minimum PDI% was recorded in
P4 (29.2%, MR), whereas the maximum was recorded with the P1 (100%, HS) (Figure 1).
All the eleven parental genotypes exhibited the TLCV symptoms after 60 days of artificial
inoculation with viruliferous whiteflies. On the TLCV disease scale, two lines showed the
mild resistance (MR), in contrast, three were highly susceptible (HS), while the rest were
susceptible (Figure 1).
Earlier studies have shown that as compared to a natural field environment screening of
genotypes for TLCV, artificial screening is more useful as it ensures a uniform disease
infection, and leave few chances for a susceptible plant to escape infection due to non-
preference and loss of whiteflies infectivity. But, 100% disease incidence is also common
with artificial inoculation method [13,42]. Artificial screening method using whiteflies is
commonly applied to find out the reaction of tomato lines to TLCV disease [43–45].
Previously, the cultivated accessions and the landraces of tomato exhibited a varying range
of the disease symptoms but lacked the complete resistance [46]. However, wild relatives of
tomato e.g. *S. chmielewskii*, *S. habrochaites*and *S. pimpinellifolium*were found resistant
even with artificial screening [14]. The differences in the results of field-based screening
method could also be due to the difference in virus strain, vector genotype or altered
feeding conditions [47, 48]

Figure 1. The percent disease incidence (PDI) reaction of parents to TLCV along with the grading scale (x-axis), where, HS (Highly Susceptible), MR (Moderately Resistant), and S (Susceptible).

3.2.*Variation in Parents and Hybrids*

The magnitude of the mean squares of genotypes indicated that there were significant differences among the genotypes for all morphological characters studied pointing out the presence of genetic variability (Table 2). Similarly, significant GCA and SCA effects for all measured traits were detected (P ≤ 0.01). The lowest values of GCA/SCA ratio was noted for polar diameter and TSS content (around 0.05) while the highest was recorded for the fruit weight (0.34). The remaining seven traits ranged from 0.10 to 0.16 (Table 2). This indicated that the non-additive gene effects were more prevalent for the characters under investigation. In general, the GCA variance was higher than that of SCA variance for the characters studied. The broad-sense heritability values were higher (above 0.9) than the narrow-sense heritability(Table 2), showing that selection between hybrids and varieties will be efficient for the improvement of these traits.

A higher magnitude of additive gene effects is useful for the development of pure-lines and to proceed with selection based breeding approaches. Non-additive gene effects are used for the development of hybrids. In previous studies presence of both additive and non-additive gene action was reported for most of the characters studied in a tomato. For example, additive gene action was reported to control yield and its component traits in tomato[49,50]. In contrast, nonadditive gene action was reported to control many traits in tomato [51,52]. Similarly, the additive and non-additive inheritance of biochemical traits like dry matter, lycopene, and TSS was also reported [53,54].

202 **Table 2.** Analysis of variance for GCA and SCA for the ten descriptors in tomato
203 under leaf curl virus conditions including GCA/SCA ratio, and narrow sense (h^2) and
204 broad sense heritabilities (H^2).

Source of Variation	Genotypes[a]	GCA[a]	SCA[a]	Error	GCA/SCA ratio	h^2	H^2
DF	65	10	55	65			
Dry matter	0.38***	0.66***	0.33***	0.11	0.16	0.22	0.95
Equatorial diameter	0.68***	0.87***	0.65***	0.01	0.10	0.17	0.97
Fruit weight	964.97***	2755.90**	639.35***	31.90	0.34	0.38	0.94
Locules	0.87***	1.48***	0.76***	0.02	0.15	0.22	0.96
Lycopene	1.58***	1.91***	1.51***	0.01	0.10	0.16	0.95
Marketable yield	0.55***	0.71***	0.52***	0.01	0.10	0.17	0.96
Pericarp thickness	0.03***	0.05***	0.03***	0.07	0.13	0.20	0.95
Polar diameter	0.83***	0.55***	0.88***	0.01	0.05	0.10	0.97
Total yield	0.56***	0.91***	0.50***	0.01	0.15	0.22	0.96
TSS	0.35***	0.28***	0.36***	0.01	0.06	0.10	0.94

[a]***, **, * indicate significant at p<0.001, p<0.01,or p<0.05,respectively.

3.3. Bayesian Model-Based Estimates and Predictions of Total Yield

Both GCA and SCA contributed for the total yield showing that both additive and
nonadditive effects were significant. However, GCA values for the total yield was higher
than the SCA values (Table 2). The predicted means further revealed this with the
BayesDiallel Bab model (Figure 2.).
The fixed and individual strain additive effects were more stable and less desperessed,
than the parent of origin and inbreeding effect (Figure 3). Justifying a large amount of GCA
component identified. The parental genotypes P1, P5 and P9 showed a negative parent of
origin and inbreeding effects (Figure 3). The observed vs expected values were found
different for some hybrid combinations (Figure 2). It could be attributed to the fact that
whiteflies have a preference for some genotypes. The virulence affected the degree and
plant reaction to the disease contributed to such effects [55,56].
Likewise, under TLCV disease pressure Bayesian approach helped in the
determination of heritable and non-heritable components influencing the total yield. This
precise determination can be used to determine the most appropriate breeding strategy for
maximising the genetic gain. Overall, this variance projection approach (VarP) is more
precise in providing the information about inheritance classes those will affect the future
experiments if these eleven parental genotypes are used again in the future [57]. The total
yield included the additive effect (VarP[a]=0.24) while, the effect of non-additive variance

was in the form of inbred penalty (VarP[B]=0.07), and the noise was 58.97 for the total
yield (Figure 4). Previously, in case of cross-pollinated carrots, it was found that the
influence of nonadditive variation was largely due to the overall inbred penalty (non-
additive effects), which contributed significantly to canopy height, shoot biomass, and root
biomass [35].
`

Figure 2. Total yield data of 55 hybrids and 11 parental genotypes in a half diallel.
On the left, of parental (P1 to P11), with crossed boxes indicating the missing data
(as half diallel was used) and the shaded box representing the values based on the
horizontal scale (below). The right side graph shows predictive means based on the
Bab diallel model on the scale of (-1.04 to 1.19).

Figure 3. Highest posterior density (HDP) intervals of parent-specific additive effects, parent of origin and inbreeding effect of the 11 parents studied.

Figure 4. Diallel variance projections the genetic architecture for total yield.

3.4. GCA and SCA estimates

Estimates of general combining ability are presented in Table 4. The estimates were
highly significant for all the characters studied for dry matter content. P2 expressed the
highest GCA effect (0.346. For equatorial diameter, P4 was the best general combiner with
the estimates of 0.273. Regarding fruit weight P5 (9.596) was the best general combiner
followed by P8 (8.453) and P7 (7.770) (Table 4). The genotype with the highest CGA
effects for the number of locules is P2 (0.355) followed by P8 (0.350) and P4 (0.261). The
genotypes with above average GCA effects for lycopene content included P3 (0.454), P6
(0.427) and P9 (0.213) (Table 4). The highest GCA effects for marketable yield of 0.188
were recorded in genotypes P2, followed by P11 and P10. For pericarp thickness P8 (0.115)
has the highest GCA effects. For polar diameter, highest GCA effects were observed in P7
(0.263). While for total yield P10 (0.215), P11 (0.164) and P6 (0.154) high GCA effects.
Likewise, the GCA effects of 0.195 and 0.152 were recorded for TSS content in the parents
P8 and P1, respectively.

Parent	Dry matter[a]	Equatorial diameter[a]	Fruit weight[a]	Locules[a]	Lycopene[a]	Marketable yield[a]	Pericarp thickness[a]	Polar diameter[a]	Total yield[a]	TSS[a]
P1	0.091^{***}	-0.201^{***}	-12.458^{***}	-0.022^{***}	-0.121^{***}	-0.333^{***}	$-0.009ns$	0.061^{*}	-0.366^{***}	0.152^{***}
P2	0.346^{***}	-0.181^{***}	6.773^{***}	0.355^{***}	-0.415^{***}	0.188^{***}	0.003^{ns}	$-0.020ns$	0.151^{***}	-0.048^{*}
P3	-0.105^{***}	-0.098^{***}	-15.122^{***}	-0.298^{***}	0.454^{***}	-0.133^{***}	-0.007^{ns}	-0.061^{*}	-0.206^{***}	0.010^{ns}
P4	-0.121^{***}	0.273^{***}	8.495^{***}	0.261^{***}	-0.086^{**}	0.048^{ns}	-0.002^{ns}	$0.025ns$	0.089^{**}	-0.168^{***}
P5	-0.067^{**}	0.137^{***}	9.596^{***}	-0.085^{**}	-0.061^{*}	-0.180^{***}	-0.016^{*}	0.177^{***}	-0.207^{***}	0.034^{ns}
P6	-0.178^{***}	0.041^{ns}	-0.690^{ns}	-0.273^{***}	0.427^{***}	0.161^{***}	-0.038^{***}	-0.257^{***}	0.154^{***}	-0.071^{**}
P7	0.021^{ns}	0.195^{***}	7.770^{***}	-0.089^{**}	0.106^{***}	0.012^{ns}	0.018^{*}	0.263^{***}	$0.032ns$	0.023^{ns}
P8	0.170^{***}	0.168^{***}	8.453^{***}	0.350^{***}	-0.169^{***}	0.018^{ns}	0.115^{***}	$-0.039ns$	$0.023ns$	0.195^{***}
P9	0.033^{ns}	-0.045^{ns}	4.867^{***}	0.126^{***}	0.213^{***}	-0.070^{**}	0.029^{***}	-0.178^{***}	-0.050^{*}	-0.011^{ns}
P10	-0.185^{***}	0.007^{ns}	$-0.081ns$	-0.120^{***}	-0.215^{***}	0.141^{***}	-0.035^{***}	0.060^{*}	0.215^{***}	-0.098^{***}
P11	-0.003^{ns}	-0.296^{***}	-17.605^{***}	-0.205^{***}	-0.132^{***}	0.148^{***}	-0.058^{***}	-0.032^{ns}	0.164^{***}	-0.019^{ns}

[a] ***, **, * indicate significant at $p<0.001$, $p<0.01$, or $p<0.05$, respectively.

The results of specific combining ability estimates are given in Table S1. The crosses
with the highest SCA effects for Dry matter content are H22 (1.343), H9 (0.986) and H17
(0.903). The highest SCA effects of 1.386, 1.179 and 1.097 were recorded in the crosses
H46, H35 and H23 respectively for the equatorial diameter. For fruit weight, the highest
positive SCA effects were observed in crosses H17 (26.728), H35(26.639) and H16
(23.633). For locule number crosses H34, H43 and H46 have the highest SCA effects of
1.551, 1.289 and 1.217 respectively. Similarly, the highest SCA estimates for the lycopene
content were recorded in the crosses H30 (1.724), H24 (1.617) and H14 (1.460) in that
order. For marketable yield crosses H49 (1.649), H23 (1.134) and H32 (0.918) recorded
the highest SCA values. Likewise, for pericarp thickness H29 (0.218), H42 (0.166) and
H25 (0.140) were significant. Crosses H3 (1.589), H34 (1.283) and H35 (1.005) possessed
the highest SCA values for the polar diameter. In case of total yield H49, H23 and H42
with values of 1.539, 1.157 and 0.917 recorded the highest SCA effects respectively. For
the TSS content crosses with highest SCA effects included H49 (1.017), H26 (0.961) and
H28 (0.928). Hybrid combinations, H23, H35, H42, H46 and H49 were found to be good
specific combiners for more than one trait (Figure 5). However, none of the hybrids
exhibited significant SCA effects for all the traits. The information about the SCA and
GCA is crucial for maximizing the genetic gain. In case of self-pollinated crops, SCA
effects generally do not contribute to the improvement of the trait. These results agree with
the results reported in the previous studies under stress and under natural conditions
[22,58–60].

Figure 5. Promising hybrid cross combinations identified based on SCA values,
under leaf curl virus infestation.

Correlations

A total of 14 correlations were found to be significant (P < 0.05) (Figure 6). One of the
correlation was absolute (0.966) that is between total yield and marketable yield. Locule
number is found to be correlated with fruit weight (0.43), equatorial diameter (0.32), dry
matter (0.31) and TSS (0.17). While the fruit weight was correlated with equatorial
diameter (0.54), locules (0.43), pericarp thickness (0.28) and polar diameter
(0.27). Previous research works conducted on tomato showed the similar results. In this
direction, a high correlation was noticed between fruit yield, fruit weight and pericarp
thickness [61,62]. Although it is worth mentioning that no correlation was found between
yield and any of the morphological and biochemical traits. Tomato plant yield under
TLCV pressure is independent of any morphological and biochemical trait. Likewise,

earlier it was shown that under salt stress, plants survival and yield are independent of each
other[63].

Figure 6. Pearson's correlation coefficients with significant values at p<0.001
(***), p<0.01 (**), or p<0.05 (*), respectively (upper diagonal) along with the
pattern of the distribution of data via scatter plots and histograms (lower diagonal)
(A). While all the significant Pearson's correlation coefficients at p<0.05 (B).

4. Conclusions

305 During the past decades, breeding for tomato leaf curl virus resistance has been a major
306 focus for the resistance breeding programs in tomato. Therefore, breeding efforts have been

made to combine significant resistance to TLCV with important fruit quality and yield traits. The diallel matting design is a popular choice as it helps in the identification of parents with good GCA effects and hybrids with good SCA effects. Additionally, it provides the important information on gene action and inheritance of the traits. In this study, we evaluated seven morphological and three biochemical traits of interest for tomato breeding under leaf curl virus pressure. The high diversity in the material was confirmed by GCA and SCA values for all traits. This showed the significance of both the additive and the non-additive effects in the inheritance of the traits evaluated. Also, we have dissected the inheritance of total yield using the Bayesian approach. It was shown that total yield was more dependent on additive variance than the non-additive variance. Overall, this information will be useful to design and develop breeding programs aiming to improve TLCV resistance along with a respectable combination of important traits. The moderately resistant and high yielding parents (P4 and P9) and hybrids (H23, H42, and H49) could be used for the resistance breeding in tomato .

Conflicts of Interest: The author declares no conflict of interest.

References

1. Kalloo, G. *Genetic Improvement of Tomato*; Springer Science & Business Media, 2012; ISBN 978-3-642-84275-7.
2. Víquez-Zamora, M.; Vosman, B.; van de Geest, H.; Bovy, A.; Visser, R. G.; Finkers, R.; van Heusden, A. W. Tomato breeding in the genomics era: insights from a SNP array. *BMC Genomics***2013**, *14*, 354, doi:10.1186/1471-2164-14-354.
3. Hossain, M. A.; Wani, S. H.; Bhattacharjee, S.; Burritt, D. J.; Tran, L.-S. P. *Drought Stress Tolerance in Plants, Vol 1: Physiology and Biochemistry*; Springer, 2016; ISBN 978-3-319-28899-4.
4. Jacobsen, S.-E.; Jensen, C. R.; Liu, F. Improving crop production in the arid Mediterranean climate. *Field Crops Research***2012**, *128*, 34–47, doi:10.1016/j.fcr.2011.12.001.
5. Jensen, C. R.; Battilani, A.; Plauborg, F.; Psarras, G.; Chartzoulakis, K.; Janowiak, F.; Stikic, R.; Jovanovic, Z.; Li, G.; Qi, X.; Liu, F.; Jacobsen, S.-E.; Andersen, M. N. Deficit irrigation based on drought tolerance and root signalling in potatoes and tomatoes. *Agricultural Water Management***2010**, *98*, 403–413, doi:10.1016/j.agwat.2010.10.018.
6. S.Kumar, G.; T.S, T. *Disease Problems In Vegetable Production*; Scientific Publishers, 2012; ISBN 978-93-86347-51-0.
7. Xu, C.; Sun, X.; Taylor, A.; Jiao, C.; Xu, Y.; Cai, X.; Wang, X.; Ge, C.; Pan, G.; Wang, Q.; Fei, Z.; Wang, Q. Diversity, Distribution, and Evolution of Tomato Viruses in China Uncovered by Small RNA Sequencing. *J Virol***2017**, *91*, 1-14, doi:10.1128/JVI.00173-17.
8. Picó, B.; Díez, M. J.; Nuez, F. Viral diseases causing the greatest economic losses to the tomato crop. II. The Tomato yellow leaf curl virus — a review. *Scientia Horticulturae***1996**, *67*, 151–196, doi:10.1016/S0304-4238(96)00945-4.
9. Saxena, S.; Tiwari, A. K. *Begomoviruses: Occurrence and Management in Asia and Africa*; Springer, 2017; ISBN 978-981-10-5984-1.
10. Gaikwad, K. A.; Cheema, D. S.; Sharma, A.; Dhaliwal, M. S. REACTIOOFELITETOMATO (*SOLANUMLYCOPERSICUM* L.) GERMPLASM AGAINST TOMATO LEAF CURL VIRUS DISEASE. *Acta Horticulturae***2011**, 157–161, doi:10.17660/ActaHortic.2011.914.28.
11. Moriones, E.; Praveen, S.; Chakraborty, S. Tomato Leaf Curl New Delhi Virus: An Emerging Virus Complex Threatening Vegetable and Fiber Crops. *Viruses***2017**, *9*, 264, doi:10.3390/v9100264.

12. Sharma, N.; Prasad, M. An insight into plant–Tomato leaf curl New Delhi virus
interaction. *Nucleus***2017**, *60*, 335–348, doi:10.1007/s13237-017-0224-5.
13. Picó, B.; Díez, M. J.; Nuez, F. Evaluation of whitefly-mediated inoculation techniques
to screen Lycopersiconesculentum and wild relatives for resistance to Tomato yellow
leaf curl virus. *Euphytica***1998**, *101*, 259–271, doi:10.1023/A:1018353806051.
14. Singh, R. K.; Rai, N.; Singh, M.; Saha, S.; Singh, S. N. Detection of tomato leaf curl
virus resistance and inheritance in tomato (*Solanum lycopersicum* L.). *The Journal of*
*Agricultural Science***2015**, *153*, 78–89, doi:10.1017/S0021859613000932.
15. Díez, M. J.; Nuez, F. Tomato. In *Vegetables II*; Handbook of Plant Breeding;
Springer, New York, NY, 2008; pp. 249–323 ISBN 978-0-387-74108-6.
16. Kaushik, P.; Andújar, I.; Vilanova, S.; Plazas, M.; Gramazio, P.; Herraiz, F. J.; Brar,
N. S.; Prohens, J. Breeding Vegetables with Increased Content in Bioactive Phenolic
Acids. *Molecules***2015**, *20*, 18464–18481, doi:10.3390/molecules201018464.
17. Arias, R.; Lee, T.-C.; Logendra, L.; Janes, H. Correlation of Lycopene Measured by
HPLC with the L^*, a^*, b^* Color Readings of a Hydroponic Tomato and the
Relationship of Maturity with Color and Lycopene Content. *J. Agric. Food*
*Chem.***2000**, *48*, 1697–1702, doi:10.1021/jf990974e.
18. Palozza, P.; Simone, R. E.; Catalano, A.; Mele, M. C. Tomato Lycopene and Lung
Cancer Prevention: From Experimental to Human Studies. *Cancers (Basel)***2011**, *3*,
2333–2357, doi:10.3390/cancers3022333.
19. Wei, M. Y.; Giovannucci, E. L. Lycopene, Tomato Products, and Prostate Cancer
Incidence: A Review and Reassessment in the PSA Screening Era. *J. Oncol***2012**,
p.271063, Available online: https://www.hindawi.com/journals/jo/2012/271063/.
20. Des Marais, D. L.; Hernandez, K. M.; Juenger, T. E. Genotype-by-Environment
Interaction and Plasticity: Exploring Genomic Responses of Plants to the Abiotic
Environment. *Annual Review of Ecology, Evolution, and Systematics***2013**, *44*, 5–29,
doi:10.1146/annurev-ecolsys-110512-135806.
21. Beckles, D. M. Factors affecting the postharvest soluble solids and sugar content of
tomato (Solanumlycopersicum L.) fruit. *Postharvest Biology and Technology***2012**, *63*,
129–140, doi:10.1016/j.postharvbio.2011.05.016.
22. Acharya, B.; Dutta, S.; Dutta, S.; Chattopadhyay, A. Breeding tomato for
simultaneous improvement of processing quality, fruit yield, and dual disease
tolerance. *International Journal of Vegetable Science***2018**, *0*, 1–17,
doi:10.1080/19315260.2018.1427648.
23. Panthee, D. R.; Perkins-Veazie, P.; Anderson, C.; Ibrahem, R. DiallelAnalysis for
Lycopene Content in the Hybrids Derived from Different Colored Parents in Tomato.
*American Journal of Plant Sciences***2015**, *06*, 1483, doi:10.4236/ajps.2015.69147.
24. Shattuck, v. I.; Christie, B.; Corso, C. Principles for Griffing's combining ability
analysis. *Genetica***1993**, *90*, 73–77, doi:10.1007/BF01435180.
25. Griffing, B. Concept of General and Specific Combining Ability in Relation toDiallel
Crossing Systems. *Aust. Jnl. Of Bio. Sci.***1956**, *9*, 463–493, doi:10.1071/bi9560463.
26. Hanson, P. M.; Licardo, O.; Hanudin; Wang, J.-F.; Chen, J. DiallelAnalysis of
Bacterial Wilt Resistance in Tomato Derived from Different Sources. *Plant*
*Disease***1998**, *82*, 74–78, doi:10.1094/PDIS.1998.82.1.74.
27. Kaushik, P.; Plazas, M.; Prohens, J.; Vilanova, S.; Gramazio, P. Diallel genetic
analysis for multiple traits in eggplant and assessment of genetic distances for
predicting hybrids performance. *PLOSONE***2018**, *13*, e0199943,
doi:10.1371/journal.pone.0199943.
28. Rêgo, E. R. do; Rêgo, M. M. do; Finger, F. L.; Cruz, C. D.; Casali, V. W. D. A diallel
study of yield components and fruit quality in chilli pepper (*Capsicum baccatum*).
*Euphytica***2009**, *168*, 275–287, doi:10.1007/s10681-009-9947-y.

29. Taggar, G. K.; Gill, R. S. Preference of whitefly, <Emphasis
Type="Italic">Bemisiatabaci,</Emphasis> towards black gram genotypes: Role of
morphological leaf characteristics. *Phytoparasitica***2012**, *40*, 461–474,
doi:10.1007/s12600-012-0247-z.
30. Toscano, L. C.; Jr, B.; Leal, A.; Maruyama, W. I. Nonpreference of whitefly for
oviposition in tomato genotypes. *Scientia Agricola***2002**, *59*, 677–681,
doi:10.1590/S0103-90162002000400009.
31. Wu, H. X.; Matheson, A. C. Analysis of half-diallel mating design with missing
crosses: theory and SAS program for testing and estimating GCA and SCA fixed
effects. *SilvaeGenetica***2000**, *49*, 130–137.
32. Xiang, B.; Li, B. A new mixed analytical method for genetic analysis of diallel data.
*Can. J. For. Res.***2001**, *31*, 2252–2259, doi:10.1139/x01-154.
33. Greenberg, A. J.; Hackett, S. R.; Harshman, L. G.; Clark, A. G. A Hierarchical
Bayesian Model for a Novel Sparse Partial Diallel Crossing Design. *Genetics***2010**,
185, 361–373, doi:10.1534/genetics.110.115055.
34. Lenarcic, A. B.; Svenson, K. L.; Churchill, G. A.; Valdar, W. A general Bayesian
approach to analyzing diallel crosses of inbred strains. *Genetics***2012**, *190*, 413–435,
doi:10.1534/genetics.111.132563.
35. Turner, S. D.; Maurizio, P. L.; Valdar, W.; Yandell, B. S.; Simon, P. W. Dissecting
the Genetic Architecture of Shoot Growth in Carrot (Daucus carota L.) Using a Diallel
Mating Design. *G3: Genes, Genomes, Genetics***2018**, *8*, 411–426,
doi:10.1534/g3.117.300235.
36. Banerjee, M. K.; Kalloo Inheritance of tomato leaf curl virus resistance in
Lycopersicon hirsutum f. *glabratum. Euphytica***1987**, *36*, 581–584,
doi:10.1007/BF00041506.
37. Sharma, O. P.; Mahajan, R.; Singh, S. Field reaction of some root-knot resistant
tomato varieties and hybrids to tobacco mosaic and tobacco leaf curl viruses. *Indian
Journal of Mycology and Plant Pathology***1984**, *14*, 148–150.
38. Maria Alda, L.; Gogoaşă, I.; Bordean, D.-M.; Gergen, I.; Alda, S.; Moldovan, C.;
Niţă, L. Lycopene content of tomatoes and tomato products. *J.
AgroalimentaryProcesses Technol.***2009**, *15*, 540-542.
39. Rodríguez, F.; Alvarado, G.; Pacheco, Á.; Crossa, J.; Burgueño, J. AGD-R (Analysis
of Genetic Designs with R for Windows) Version 4.0 2017.
40. Wei, T.; Simko, V.; Levy, M.; Xie, Y.; Jin, Y.; Zemla, J. *corrplot: Visualization of a
Correlation Matrix*; 2017;
41. Peterson, B. G.; Carl, P.; Boudt, K.; Bennett, R.; Ulrich, J.; Zivot, E.; Cornilly, D.;
Hung, E.; Lestel, M.; Balkissoon, K.; Wuertz, D. *PerformanceAnalytics: Econometric
Tools for Performance and Risk Analysis*; 2018;
42. Lapidot, M.; Ben-Joseph, R.; Cohen, L.; Machbash, Z.; Levy, D. Development of a
Scale for Evaluation of Tomato yellow leaf curl virus Resistance Level in Tomato
Plants. *Phytopathology***2006**, *96*, 1404–1408, doi:10.1094/PHYTO-96-1404.
43. Ambawat, S.; Sharma, P.; Yadav, N. R.; Yadav, R. C. MYB transcription factor genes
as regulators for plant responses: an overview. *Physiol Mol Biol Plants***2013**, *19*, 307–
321, doi:10.1007/s12298-013-0179-1.
44. Gaikwad, K. A.; Sharma, A.; Cheema, D. S.
MOLECULARDETECTIONANDCHARACTERIZATIONOFLEAFCURLVIRUSIN
FECTINGTOMATOINPUNJAB, INDIA. *Acta Horticulturae***2011**, 153–156,
doi:10.17660/ActaHortic.2011.914.27.
45. Lapidot, M.; Friedmann, M.; Lachman, O.; Yehezkel, A.; Nahon, S.; Cohen, S.;
Pilowsky, M. Comparison of Resistance Level to Tomato Yellow LeafCurlVirus

Among Commercial Cultivars and Breeding Lines. *Plant Disease***1997**, *81*, 1425–
1428, doi:10.1094/PDIS.1997.81.12.1425.

46. Pérez-De-Castro, A.; Campos, G.; Julián, O.; Dueñas, F.; Álvarez, M.; Martínez-
Zubiaur, Y.; Díez, M. J. Evaluation under diverse conditions of a differential host
reaction scale to Tomato yellow leaf curl virus in tomato.
*PhytopathologiaMediterranea***2017**, *56*, 50–61, doi:10.14601/Phytopathol_Mediterr-
18356.

47. Fereres, A.; Peñaflor, M. F. G. V.; Favaro, C. F.; Azevedo, K. E. X.; Landi, C. H.;
Maluta, N. K. P.; Bento, J. M. S.; Lopes, J. R. S. Tomato Infection by Whitefly-
Transmitted Circulative and Non-Circulative Viruses Induce Contrasting Changes in
Plant Volatiles and Vector Behaviour. *Viruses***2016**, *8*, 225, doi:10.3390/v8080225.

48. Channarayappa, C.; Shivashankar, G.; Muniyappa, v.; Frist, R. H. Resistance of
Lycopersicon species to Bemisiatabaci, a tomato leaf curl virus vector. *Can. J.
Bot.***1992**, *70*, 2184–2192, doi:10.1139/b92-270.

49. Farzane, A.; Nemati, H.; Aroiee, H.; Mirshamsi, A.; Vahdati, N. The Estimate of
Combining Ability and Heterosis for Yield and Yield Components in Tomato
(Lycopersiconesculentum Mill.). *J. Biol. Environ. Sci.***2012**, *6,* 129-134.

50. Shalaby, T. A. Mode of gene action, heterosis and inbreeding depression for yield and
its components in tomato (Solanumlycopersicum L.). *Scientia Horticulturae***2013**,
164, 540–543, doi:10.1016/j.scienta.2013.10.006.

51. Shankar, A.; Reddy, R.V.S.K.; Sujatha, M.; Pratap, M. Combining ability and gene
action studies for yield and yield contributing traits in tomato (*Solanumlycopersicum*
L.) Helix **2012**, *6*, 431 - 435.

52. Govindarasu, P.; Muthukrishnan, C. R.; Irulappan, I. Combining-ability for yield and
its components in tomato. *Scientia Horticulturae***1981**, *14*, 125–130,
doi:10.1016/0304-4238(81)90003-0.

53. Dufera, J. T.; Mohammed, W.; Hussien, S.; Kumar, V. Genetic Analysis for Fruit
Yield and Quality Attributes in Some Processing and Fresh Market Tomato
Genotypes. *International Journal of Vegetable Science***2018**, *24*, 227–235,
doi:10.1080/19315260.2017.1410873.

54. Kumar, R.; Srivastava, K.; Singh, N. P.; Vasistha, N. K.; Singh, R. K.; Singh, M. K.
Combining Ability Analysis for Yield and Quality Traits in Tomato
(Solanumlycopersicum L.). *Journal of Agricultural Science***2013**, *5*, 213-218,
doi:10.5539/jas.v5n2p213.

55. Bleeker, P. M.; Diergaarde, P. J.; Ament, K.; Guerra, J.; Weidner, M.; Schütz, S.; de
Both, M. T. J.; Haring, M. A.; Schuurink, R. C. The Role of Specific Tomato Volatiles
in Tomato-Whitefly Interaction. *Plant Physiol***2009**, *151*, 925–935,
doi:10.1104/pp.109.142661.

56. Maluta, N. K. P.; Fereres, A.; Lopes, J. R. S. Settling preferences of the whitefly
vector Bemisiatabaci on infected plants varies with virus family and transmission
mode. *EntomologiaExperimentalis et Applicata165*, 138–147, doi:10.1111/eea.12631.

57. Crowley, J. J.; Kim, Y.; Lenarcic, A. B.; Quackenbush, C. R.; Barrick, C. J.; Adkins,
D. E.; Shaw, G. S.; Miller, D. R.; de Villena, F. P.-M.; Sullivan, P. F.; Valdar, W.
Genetics of adverse reactions to haloperidol in a mouse diallel: a drug-placebo
experiment and Bayesian causal analysis. *Genetics* **2014**, *196*, 321–347,
doi:10.1534/genetics.113.156901.

58. Lunt, K. A.; Roux, C. Z.; Blokpoel, I.; Booyse, M. Inheritance of flavour in six tomato
(Lycopersiconesculentum) varieties. *South African Journal of Plant and Soil* **1999**, *16*,
113–117, doi:10.1080/02571862.1999.10634994.

59. Pandiarana, N.; Chattopadhyay, A.; Seth, T.; Shende, V. D.; Dutta, S.; Hazra, P.
Heterobeltiosis, potence ratio and genetic control of processing quality and disease

severity traits in tomato. *New Zealand Journal of Crop and Horticultural Science* **2015**, *43*, 282–293, doi:10.1080/01140671.2015.1083039.

60. Souza, L. M. de; Paterniani, M. E. A.; Melo, P. C. T. de; Melo, A. M. de Diallel cross among fresh market tomato inbreeding lines. *Horticultura Brasileira* **2012**, *30*, 246–251, doi:10.1590/S0102-05362012000200011.

61. Kumar, D.; Kumar, R.; Kumar, S.; Bhardwaj, M. L.; Thakur, M. C.; Kumar, R.; Thakur, K. S.; Dogra, B. S.; Vikram, A.; Thakur, A.; Kumar, P. Genetic Variability, Correlation and Path Coefficient Analysis in Tomato. *International Journal of Vegetable Science* **2013**, *19*, 313–323, doi:10.1080/19315260.2012.726701.

62. Souza, L. M. de; Melo, P. C. T.; Luders, R. R.; Melo, A. M. Correlations between yield and fruit quality characteristics of fresh market tomatoes. *Horticultura Brasileira* **2012**, *30*, 627–631, doi:10.1590/S0102-05362012000400011.

63. Rao, E. S.; Kadirvel, P.; Symonds, R. C.; Ebert, A. W. Relationship between survival and yield related traits in *Solanum pimpinellifolium* under salt stress. *Euphytica* **2013**, *190*, 215–228, doi:10.1007/s10681-012-0801-2.