1 **Coping with stress: morphological changes in response to environmental stimuli**

2 **in a fungal plant pathogen**

7 Carolina Sardinha Francisco[1], Xin Ma[1], Maria Manuela Zwyssig[1], Bruce A. McDonald[1],

8 Javier Palma-Guerrero[1*]

13 [1]Plant Pathology Group, Institute of Integrative Biology, ETH Zürich, 8092 Zürich,

14 Switzerland

19 *Corresponding author

20 E-mail: javier.palma@usys.ethz.ch

Abstract

During their life cycles, pathogens have to adapt to many biotic and abiotic environmental stresses to maximize their overall fitness. Morphological transitions are one of the least understood of the many strategies employed by fungal plant pathogens to adapt to constantly changing environments, even though different morphotypes may play important biological roles. We characterized the responses of the wheat pathogen *Zymoseptoria tritici* to a series of environmental stresses in order to understand the effects of changing environments on fungal morphology and adaptation. We found that all tested stresses induced morphological changes, but different responses were found among four different strains. A transcription analysis showed that morphogenesis and virulence factors are co-regulated. We discovered that *Z. tritici* forms chlamydospores and demonstrated that these structures are better able to survive extreme cold, heat and drought than other cell types. We also show that blastospores (the "yeast-like" form of the pathogen typically found only in laboratory conditions) can form from germinated pycnidiospores on the surface of wheat leaves, suggesting that this morphotype can play an important role in the natural history of *Z. tritici*. Our findings illustrate how changing environmental conditions can affect cellular morphology and lead to the formation of new morphotypes, with each morphotype having a potential impact on both pathogen survival and disease epidemiology.

Introduction

Fungi occupy a wide range of niches that can impose different environmental constraints on their growth, reproduction and survival. In response, fungi evolved the ability to detect and respond to different environmental stimuli to maximize their fitness [1]. One of the strategies that fungi employ to cope with diverse biotic and abiotic stresses is to change their morphology [2–4]. For fungal pathogens, different morphotypes can play different roles during the host-pathogen interaction to optimize overall pathogen fitness. For example, hyphae can be the most appropriate morphology to cross physical barriers, colonize host tissue [2,5], and escape harmful environments generated by host defences [2,6] while yeast cells (or "yeast-like" cells) may be a better morphology to enable dispersal among host niches or between hosts and increase the pathogen population size [2,7]. The majority of fungal morphotype transitions occur between hyphal and yeast growth forms. Fungi with this ability are called dimorphic. Usually, they grow as saprophytic molds that live on dead organic matter and many are facultative human pathogens, though dimorphism is also found in plant pathogens [2,8,9]. The morphological transition can be co-regulated with the expression of genes encoding virulence [2–5,7,10,11]. The induction of virulence genes is usually associated with a specific morphotype. For example, certain virulence genes are expressed only during the yeast-phase of growth in *Histoplasma capsulatum,* while other virulence genes are expressed only during hyphal growth in *Candida albicans* [5,6,12].

Some fungi can produce additional morphologies, including pseudohyphae and

chlamydospores. Fungi that can produce more than two morphotypes are called

pleomorphic. Pseudohyphae are distinguished from true hyphae by constrictions that

form at the septal junctions, leading to a loss in cytoplasmic continuity between mother

and daughter cells [13]. Though little is known about the ecological significance of

pseudohyphae and their role during host infection, it was suggested that this

morphotype facilitates foraging for scarce nutrients and increases motility in the host

[7,14]. Chlamydospores are spherical, thick-walled cells formed on the tips of

pseudohyphae or on distal parts of differentiated hyphae called suspensor cells [15]. In

contrast to pseudohyphae, chlamydospores are well-characterized and known to

operate as long-term survival structures that can resist many environmental stresses

[15–19]. Pleomorphism has been observed in fungal pathogens of humans [15,17,20]

and plants [18,21,22].

Zymoseptoria tritici is the most damaging pathogen of wheat in Europe [23,24] and an

important pathogen of wheat worldwide. *Z. tritici* is considered a dimorphic fungus that

grows either as blastospores (the "yeast-like" form) or as filamentous hyphae depending

on the culture conditions. Though the role of blastospores in nature was unclear until

now, hyphae are essential for penetrating wheat leaves through stomata [25,26]. The

stimuli that induce morphological changes and the cell signaling pathways involved in

morphological transitions remain largely unknown in this pathogen. Nutrient-rich media

and temperatures ranging from 15°C to 18°C tend to induce blastosporulation

(blastospore replication by budding) while nutrient-limited media and high temperatures

(from 25°C to 28°C) tend to promote the blastospore-to-hyphae transition in the *Z. tritici*

reference strain IPO323 [27,28]. Some genes affecting morphogenesis in *Z. tritici* were

identified and functionally characterized. Most of these belonged to the mitogen-

activated protein kinase (MAPK) or cAMP-dependent signaling pathways involved in

extracellular signal transduction, regulating many cellular processes, development and

virulence [27–37]. However, the transcriptome signatures associated with specific

morphotypes were not reported until now.

In this study we aimed to characterize how environmental changes affect morphology and

adaptation in a fungal plant pathogen. We found that *Z. tritici* changed its morphology in

response to all tested stresses, though distinct responses were observed for four different

strains sampled from the same geographical population. Transcriptional analyses showed

a co-regulation of mycelial growth with expression of virulence factors. We discovered that

Z. tritici forms chlamydospores both *in vitro* and *in planta*. We also found that

pseudohyphae will form in some environments. We showed that blastospores can form

by budding from pycnidiospores inoculated onto the surface of wheat leaves,

demonstrating that blastospores can be produced under natural conditions. Based on

these findings, we propose that morphological transitions in *Z. tritici* are a stress response

that allows the fungus to optimize fitness in a changing environment.

Results

High temperature induces different morphological responses among *Z. tritici* strains.

We exposed four different *Z. tritici* strains growing in yeast sucrose broth (YSB) at 27°C, a highly stressful temperature for *Z. tritici*. Three of the strains (1E4, 3D1, and 3D7) differentiated to filamentous growth at 27°C (Fig 1) as previously reported for the reference strain IPO323 [27,28]. The timing of the hyphal transition ranged from 8 hours after incubation (hai) for 1E4, to 24 hai for 3D1 and 3D7 (Figs 1B and 1C). Branching and hyphal elongation were frequently observed at 48 hai (Figs 1D-1F). Surprisingly, high temperature promoted the formation of two new morphologies, with appearances matching pseudohyphae and chlamydospores, in the 1A5 strain (Figs 1E and 1F). However, no hyphal growth was observed for this strain. Interestingly, chlamydospore-like cells were found for all strains at 144 hai (Fig 1F). Cellular differentiation from blastospores to pseudohyphae began in 1A5 at 24 hai, resulting in a series of conjoined elongated cells, characteristic of pseudohyphae [13] (Fig 1C). We also observed several swollen cells at 24 hai that we believe represent the initial differentiation from blastospores into chlamydospore-like cells (Fig 1C), but the complete differentiation occurred between 48 and 96 hai (Figs 1D and 1E). Chlamydospore-like cells were observed at the ends of pseudohyphae or distal parts of filamentous hyphae (Fig 1F). Chlamydospore-like cells that were detached from suspensor cells were also found (data not shown). Blastospores incubated in YSB at 18°C continued to replicate only as blastospores via blastosporulation and were used as controls (Fig 1A).

A nutrient-limited environment promotes filamentous growth.

Carbon- or nitrogen-depleted medium promoted hyphal growth in all four strains (Fig 2;

S1 Fig). In a carbon-depleted environment (Minimal medium (MM) without sucrose),

cellular differentiation occurred after 24 hours of incubation (Fig 2). At 24 hai, hyphal

formation was observed with the first hyphal branch formed terminally (1A5 and 3D1) or

laterally (1E4 and 3D7) (Fig 2B). Hyphal elongation and branching continued at 48 and

72 hai (data not shown). By 144 hai mycelial growth was predominant, with frequent

anastomosis events among hyphae (Fig 2C). In MM with sucrose, all strains grew as a

mixture of blastospores and hyphae, with blastospores budding from filamentous hyphae,

showing that the morphological transition is bidirectional (S1C Fig). Blastospores

incubated in YSB continued to replicate only as blastospores via blastosporulation (Fig

2A).

In nitrogen-depleted media (MM without ammonium nitrate), three strains (1A5, 3D1, and

3D7) initially formed hyphae from one apical end, generating the first and second hyphal

elongations within 72 hai (S1B Fig). 1E4 responded most quickly to a low nitrogen

environment, with hyphae forming at 24 hai (S1A Fig) and a network of filamentous

hyphae formed at 72 hai (S1B Fig). In regular MM all strains grew as a mixture of

morphotypes (S1C Fig). Interestingly, the hyphae induced under both carbon and nitrogen

limitation (Fig 2; S1 Fig) were thinner than those produced by high temperature (Fig 1).

Oxidative stress induces morphological changes in some strains.

To determine the effect of oxidative stress on the *Z. tritici* morphology we examined the

growth in presence of hydrogen peroxide (H_2O_2) at two different spore concentrations. At

a high initial cell density (10^5 blastospores/mL), the presence of H_2O_2 stimulated hyphal

growth in 1E4 up to a concentration of 5 mM (S2 Fig). 1A5, 3D1, and 3D7 continued to

replicate as blastospores, but there was a substantial reduction in blastospore formation

starting at 5 mM H_2O_2 (S2 Fig). To test whether a lower concentration of blastospores

would change sensitivity to oxidative stress, we inoculated flasks with 10^3

blastospores/mL. After 72 hours of incubation in 0.5 mM H_2O_2, strains 1E4 and 3D1

showed apical and lateral hyphal branching (S3 Fig), but both strains continued to form

blastospores. At 1 mM, blastospores from 1E4, 3D1, and 3D7 developed into a mixture of

pseudohyphae and regular hyphae (S3 Fig). The higher sensitivity at low cell densities

suggests that the exogenous H_2O_2 was more rapidly degraded by secreted

hydroperoxidase proteins (catalases, peroxidases and catalase-peroxidase) when cell

densities were 100X higher. The 1A5 strain did not switch to a different morphotype at

any H_2O_2 concentration, continuing to replicate as blastospores (S3 Fig). Starting at 2 mM

H_2O_2, none of the strains changed their morphology (S3 Fig) and blastosporulation ceased

(from 5 mM to 10 mM), suggesting a cytotoxic effect of H_2O_2 at these concentrations.

Inoculum size effect suggests that quorum sensing may affect morphology.

In order to evaluate the effect of the inoculum size on the morphological transition, we

incubated all four strains at five cell densities in two growth conditions shown to induce

hyphal growth. After 144 hours of incubation in C-depleted medium, we observed that

morphology was independent of cell density for all four strains up to a concentration of

10^5 blastospores/mL (S4A Fig), with most of the blastospores switching to hyphal growth

and forming mycelial colonies. However, when the initial cell density was $\geq 10^6$

blastospores/mL, we observed a substantial reduction in the blastospore-to-hyphae

transition and in hyphal length. No hyphae formed for 3D1 and 3D7 at initial cell densities

of 10^7 blastospores/mL, and they were clearly reduced for 1E4 and 1A5 (S4A Fig).

In YSB at 27°C we also found that initial inoculum size affected cell morphology (S4B

Fig), though there were more pronounced differences among strains in this environment.

At 144 hai, starting from the lowest initial cell density (10^3 blastospores/mL), we

observed pseudohyphae formation in 1A5, 1E4, and 3D1, and a mixture of

pseudohyphae and chlamydospore-like cells in 3D7. At an initial density of 10^4

blastospores/mL, the same phenotypes were found, except that 3D1 formed apical and

lateral hyphae. Intermediate initial cell densities (10^5 and 10^6 blastospores/mL)

promoted hyphal growth for 1E4, 3D1 and 3D7, but the 3D7 strain grew as a mixture of

mycelium and chlamydospore-like cells at 10^6 blastospores/mL. The increase of the

inoculum size to 10^7 blastospores/mL induced formation of chlamydospore-like

structures in 1E4 and 3D1, and a transition from blastospores to pseudohyphae and

chlamydospore-like cells in 3D7. 1A5 was not able to switch to hyphae under high

temperatures in any of the initial cell densities (S4B Fig). The overall pattern of cell

density dependence shown here suggests that a quorum sensing molecule may be

controlling the morphological transition, as has been reported for other pleomorphic

fungi [38–40].

Mycelial growth is co-regulated with virulence-related genes.

Aiming to identify genes associated with mycelial growth in nutrient-poor environments,

we compared the transcriptomes from blastospores and two different hyphal inducing

conditions for the four strains (S5Fig). The total number of sequencing reads and mapped

reads per strain under different growth conditions is given in S1 Table. A total of 4888,

5296, 5323, and 6003 genes were differentially expressed among all tested conditions in

1A5, 1E4, 3D1, and 3D7, respectively. On average, 24% and 25% of the genes were

upregulated during blastospore and mycelial growth, respectively.

The upregulated genes shared among all four strains for each morphology were selected

based on pairwise comparisons. Two distinct sets of 134 genes shared among the four

strains were identified in both comparisons. The first set is composed of genes that were

upregulated during blastospore growth: these genes were upregulated in YSB compared

to C-depleted medium, and were upregulated in YSB compared to 7dpi infection, FDR

≤0.01 (S6A Fig; S2 Table). The second set consists of genes that were upregulated during

mycelial growth: these were upregulated in C-depleted medium compared to YSB, and

were upregulated in wheat infection compared to YSB, FDR ≤0.01 (S6B Fig; S3 Table).

For both comparisons, the sets of genes that do not overlap represent genes with a strain-

specific expression profile.

Among the 134 shared genes that are significantly upregulated during blastospore growth,

20% are annotated as *hypothetical proteins* (Fig 3A; S2 Table). The remaining 107 genes

are divided into fourteen biological process categories, with *metabolism* being the most

represented term (44%). This category was composed mainly of non-secreted enzymes,

including oxidoreductases (n=11), dehydrogenases (n=8), hydrolases (n=7), and

transferases (n=9) (Fig 3A). No enrichment was found for any other category. However,

seven genes from other categories were already described in other fungi to be involved

in blastosporulation (S4 Table). As expected, genes related to vegetative blastospore

growth were highly expressed in nutrient-rich medium - the environment most conducive

for this morphotype (S6C Fig).

Among the 134 genes that were upregulated in all strains during mycelial growth, we

observed an enrichment of secreted proteins (32%) known to be associated with fungal

virulence, including *candidate effectors* (small secreted proteins - SSPs), *secreted*

proteases, *secreted glycosidases*, *secreted chloroperoxidases, cell wall hydrolases,*

hydrophobins, and *hydrophobic surface binding proteins* (Fig 3B; S3 Table). Overall,

these virulence-associated genes were upregulated in C-depleted medium and during

all stages of plant infection, albeit with a reduction in expression during the saprophytic

phase of infection (at 28 dpi). The peak of expression of these virulence-associated

genes preceded the first visual symptoms of infection, at 12 dpi for 1A5 and 3D7, and at

14 dpi for 1E4 and 3D1 (S6D Fig), as previously reported for some of these gene

categories [41]. These patterns suggest that mycelial growth is co-regulated with the

expression of genes relevant for virulence even in the absence of the host. We also

detected the upregulation of six genes related to cellular morphogenesis (S5 Table). The

transcription profiles of these genes showed an upregulation in both mycelial growth

conditions (C-depleted medium and plant infection at 7 dpi) with the peak of expression

occurring at 7 dpi for all four strains (S6E Fig).

Chlamydospore-like cells are produced by *Z. tritici*.

As mentioned above, we observed the formation of structures with the same

morphology as chlamydospores. These structures had never been previously reported

in *Z. tritici*. We aimed to determine whether these chlamydospore-like cells had

properties associated with chlamydospores in other fungi. Chlamydospores are known

to be survival structures that have thicker cell walls and a high content of lipid droplets

[42]. We stained the different morphological cell types of 1A5 with the chitin-binding dye

Calcofluor White (CFW) and the lipid droplets dye Nile Red (NR) to measure these

properties. The intensity of CFW fluorescence was higher in chlamydospore-like cells

(Fig 4A-4C) compared with blastospores, hyphae, and pseudohyphae that exhibited a

weaker fluorescence for CFW (Fig 4). NR staining revealed that chlamydospore-like

cells had a high content of lipid droplets, which was also observed for blastospores and

pseudohyphae. On the contrary, hyphae showed a lower lipid droplet content than

spores, with the exception of the hyphal branching zones (Fig 4) as has been reported

for other fungi [43]. We then examined the thickness of the cell walls for three cell types

using transmission electron microcopy (TEM). TEM showed that the chlamydospore-like

cells had thicker cell walls than blastospores and pycnidiospores (Fig 5). We also found

that pseudohyphae and the elongated suspensor cells, where chlamydospores are

usually attached to the surrounding mycelium, have a cell wall thickness similar to

pycnidiospores (Fig 5). No differences were observed for the chlamydospore-like cells

produced by the 1A5 and 3D7 strains. On average, chlamydospore-like cells produced

by *Z. tritici* had a cell wall thickness of ~460 nm, about four times thicker than the cell

walls found in blastospores and pycnidiospores. Despite the thickness of the cell wall,

we could not differentiate an extra cell wall layer as reported in chlamydospores from

other fungal species [44,45].

To determine if the chlamydospore-like cells are also produced during plant infection,

wheat plants were inoculated with pycnidiospores of all four strains expressing

cytoplasmic GFP and kept in a greenhouse chamber until pycnidia formation. Microscopic

examination of the spore solution harvested from 20-day old infected leaves showed the

presence of chlamydospore-like cells mixed among the pycnidiospores (S7 Fig).

The chlamydospore-like cells are metabolically active and highly stress tolerant.

The chlamydospore-like cells of 1A5 germinated and produced hyphae. After 12 hours of

incubation on water agar (WA) plates, we observed germ tubes emerging from the

chlamydospore-like cells, which led to the development of hyphae at 24 hai (Fig 6B and

6C). Blastospores used as controls initially germinated from one apical side, generating

the first and second germ tubes within 12 hai (Fig 6A). We inoculated wheat with

chlamydospore-like cells from GFP-tagged 1A5 to observe by confocal microscopy

whether these cells could germinate on the leaf surface. At 24 hpi, free chlamydospore-

like cells and chlamydospore-like cells still attached to a suspensor cell produced germ

tubes and hyphae on the surface of wheat leaves (Fig 6D). These results suggest that

chlamydospore-like cells may be able to cause infections.

We also investigated whether chlamydospore-like cells share other characteristics of

fungal resting spores, such as an ability to resist drought and temperature extremes.

Incubation for one day under drought stress reduced viability of blastospores and

pseudohyphae by 70%, while the chlamydospore-like cells had a reduction in cell viability

of only 23% (Fig 7A). These rates changed little after 3 days of drought stress. After 5

days of drought stress, the survival rates were 10%, 19%, and 49% for blastospores,

pseudohyphae and chlamydospore-like cells, respectively (Fig 7A). After 10 days of

drought stress, the chlamydospore-like cells exhibited a survival rate of 50%, significantly

higher than the other cell types (Fig 7A). In the cold stress experiment, only the

chlamydospore-like cells survived the cold shock, with no statistically significant difference

in the survival rates of 23%, 17%, and 19% after 1 min, 2 min and 3 min in liquid nitrogen,

respectively (Fig 7B). In the heat stress experiment, 30°C affected blastospores and

chlamydospore-like cells equally, but the 35°C treatment killed all the blastospores while

chlamydospore-like cells and pseudohyphae survived at rates of 12% and 6%,

respectively (Fig 7C). None of the morphotypes survived at 40°C (Fig 7C). For all tested

environmental stresses, chlamydospore-like cells were the morphotype that exhibited the

highest survival rate, suggesting that this structure exhibits ecological characteristics

typical for fungal resting spores that enable improved survival to different stresses

encountered in nature.

Germinating pycnidiospores produce blastospores on the surface of wheat leaves.

Pycnidiospores of a GFP-tagged 3D7 strain were inoculated onto wheat plants and

monitored using confocal microscopy. At 24 hai, we observed the expected pycnidiospore

germination and initial hyphal branching, but we also observed newly-budding

blastospores (Fig 8A). After 48 hours, there was an increase in the number of budding

points on germinated pycnidiospores, and an increase in the number of free blastospores

visible on the leaf (Fig 8B). This suggests that pycnidiospores of *Z. tritici* can germinate to

produce additional blastospores even before penetration of the wheat leaf, providing an

alternative mechanism for splash-dispersal of infective propagules and potentially

providing a substantial increase in the total inoculum load associated with an infection

cycle. At 48 hpi we also noticed many anastomosis events between pycnidiospores and

new blastospores (Fig 8B).

Discussion

Fungi evolved to perceive a wide array of environmental signals and reprogram their

cellular responses accordingly in order to maximize their fitness. Though much research

has been oriented around morphological changes and their associated transcriptional

pathways in response to different stressors in fungal pathogens of humans [10,15,46],

knowledge in this area is limited for plant pathogens. Our study shows how different

environmental stresses can affect cellular morphologies in the wheat pathogen *Z. tritici*.

We also demonstrate the phenotypic plasticity of this fungus in response to

environmental changes, as well as differential responses to the same environments

among strains. An unexpected outcome of this study was the discovery of two novel

morphotypes, one of which may have important epidemiological consequences.

We found that several environmental stresses can affect morphological transitions in

strain-specific ways for four natural Z. tritici strains sampled from the same Swiss

population. All four strains switched to hyphal growth in a nutrient-poor minimal medium.

Morphological changes induced by limited nutrients have also been described in other

dimorphic plant pathogenic fungi distributed across different phyla [9,47]. Nutrients are

known to be present on the leaf surface in very small quantities and are usually

surrounded by epicuticular waxes [48]. As a result, plant pathogens typically experience

nutritient limitations while growing on leaves. It is thought that secreted enzymes,

including plant cell wall degrading enzymes (PCWDEs) such as cutinases, are required

to obtain nutrients during this stage of development, but Z. tritici has few PCWDEs

compared to other fungi [49], suggesting that the perception of a nutrient-limited

environment may act as a stimulus to induce a switch to an exploratory morphotype

(hyphae) that can better explore the environment for food. We found a significant

increase in the frequency of anastomosis events during mycelium formation under

nutrient-limited conditions. Mycelial networks formed by anastomosis enable a shared

cytoplasmic flow that can more efficiently distribute water, nutrients, and signals within a

fungal colony [50,51]. Moreover, hyphal fusions were shown to be required for virulence

for some plant pathogens [52,53]. We hypothesize that hyphal anastomosis provides

benefits during epiphytic growth of germinating Z. tritici pycnidiospores prior to stomatal

penetration, with anastomosis facilitating colony establishment and increasing strain

fitness during colonization of the host.

The oxidative burst is a ubiquitous plant defense response to pathogen infection,

resulting in a rapid release of reactive oxygen species (ROS), including hydrogen

peroxide (H_2O_2) [54]. The ROS produced during plant defense were proposed to have

various functions, including activation of plant defense-related genes and production of

phytoalexins in addition to a direct antimicrobial effect that inhibits pathogen growth [55–

57]. Though many studies have demonstrated its antimicrobial effects, the effects of

H_2O_2 on morphological transitions remain largely unexplored. In *Candida albicans*, H_2O_2

induces hyphal differentiation in a dose-dependent manner [58]. Our results show that

blastospores of three out of four *Z. tritici* strains switch to hyphae and pseudohyphae

when exposed to H_2O_2 in a dose-dependent manner. A cell density dependence was

also found. The different cell density responses suggest that the H_2O_2 is more rapidly

metabolized by fungal hydroperoxidase proteins (catalases, peroxidases and catalase-

peroxidase), leading to lower toxicity at higher cell densities. Several genes encoding

peroxidases and catalase-peroxidases were identified in the genome of *Z. tritici* [59,60],

which may explain the observed phenotype. The inability of 1A5 to differentiate hyphae

from blastospores at any H_2O_2 concentration suggests that this strain either cannot

sense oxidative stress or that it possesses a more efficient antioxidant response that

can neutralize the effects of H_2O_2. Further experiments will be needed to test these

hypotheses.

Temperature is an important abiotic factor that strongly affects growth, reproduction, and

development in fungi [66,67]. Thermally dimorphic human fungal pathogens usually

undergo morphological transitions in response to host temperature [47,68]. *Z. tritici* is the

first plant pathogen shown to undergo morphological transitions after exposure to high

temperatures [28,66]. Our experiments showed that a change in temperature from 18°C

to 27°C triggered hyphal growth in three out of four tested *Z. tritici* strains. For these

three strains, an increase in hyphal elongation and formation of branches occured soon

after exposure to the higher temperature. The hyphae formed at a high temperature

appeared thicker than those formed under nutritient limitations and it is not clear if they

would be able to cause infections.

We used transcriptome sequencing to identify genes associated with mycelial growth in

Z. tritici. It was shown in other plant pathogenic fungi that starvation induces the

expression of virulence-related genes [69,70]. In *Z. tritici*, we found that genes relevant

for virulence, including secreted hydrolytic proteins and candidate effectors, were

upregulated in both hyphae-inducing conditions (carbon-depleted medium and *in*

planta), suggesting that genes controlling morphogenesis and virulence may be co-

regulated under nutrient-limited conditions. In *C. albicans,* genes related to hyphal

growth are co-regulated by transcription factors that affect expression of genes involved

in virulence [5,7]. However, it is important to note that not every virulence function is

connected to morphogenesis. The specific genes involved in the activation and

regulation of these morphological transitions remain unknown for *Z. tritici*.

An unexpected outcome of this study was the discovery of chlamydospore-like cells in *Z.*

tritici. The preponderance of evidence indicates that these cells are truly

chlamydospores. Their morphology matched the chlamydospores described in other

fungi: spherical, thick-walled cells with a high lipid content that form either intercalated or

at the ends of filamentous hyphae and occasionally on pseudohyphae [17,71]. We also

showed that these cells were able to survive several stresses that killed other cell types,

including dessication and high and low temperatures, consistent with the function of

chlamydospores as long-term survival structures in other fungi [16–18,72]. We conclude

that *Z. tritici* produces chlamydospores.

This is the first time that chlamydospores have been found in a plant pathogen that is

thought to mainly inhabit the phyllosphere. Though chlamydospores were especially

prominent in the 1A5 strain at high temperature, we observed chlamydospore production

in all four strains both *in vitro* and *in planta*. We hypothesize that the chlamydospores

produced by *Z. tritici* may contribute an important source of inoculum that can survive

between growing seasons and perhaps form a long-term spore-bank that can persist for

many years in wheat-growing regions. Chlamydospores may also contribute to long-

distance movement of the pathogen on infested straw or seeds. These properties could

impact the management of septoria tritici blotch (STB), but additional studies will be

needed to understand the roles played by chlamydospores during the STB disease cycle.

Another finding with epidemiological implications was to demonstrate that blastospores

can form from germinating pycnidiospores on the surface of wheat leaves. This

possibility was proposed more than 30 years ago [73,74], but was not followed up in

subsequent studies. The formation of blastospores during *in vitro* growth in nutrient-rich

medium is why *Z. tritici* has long been considered to be a dimorphic fungus, but until

now it was not clear if the blastospores could also form on leaf surfaces in nature. Our

experiments confirmed that blastospores are produced during a natural infection. It

remains unclear whether blastospores make a significant contribution to the

development of epidemics. Given their small size and increasing abundance over time,

we hypothesize that they may play an important role in the explosive increase in STB

that can occur following prolonged rainy periods that last for several days [73,75].

In conclusion, we demonstrate that a fungal plant pathogen responds morphologically to

diverse environmental stresses, and we propose distinct biological roles for different

morphotypes. We also illustrate the advantage of including different natural strains when

studying the response of a species to environmental stresses, as we found different

morphological responses among the tested strains. Our findings provide important new

insights into the *Z. tritici*-wheat pathosystem including: 1) the phenotypic plasticity of *Z.*

tritici in response to different environmental changes; 2) the reclassification of *Z. tritici* as

a pleomorphic fungus due to its ability to produce four distinct morphotypes; 3) the

discovery of chlamydospores that may help the fungus to survive stressful conditions; 4)

the formation of blastospores from germinating pycnidiospores on the surface of wheat

leaves, enabling a potentially significant increase in inoculum size. These findings position

Z. tritici as an excellent model organism to explore how morphological changes can affect

adaptation to different environments in fungal plant pathogens.

Materials and methods

Fungal strains, environmental stimuli, and growth conditions.

Four previously characterized *Zymoseptoria tritici* strains (ST99CH_1A5, ST99CH_1E4,

ST99CH_3D1, and ST99CH_3D7, abbreviated as 1A5, 1E4, 3D1, and 3D7,

respectively); [76,77], were used in this study. Strains expressing cytoplasmic GFP [78]

were provided by Andrea Sanchez-Vallet after being generated by Sreedhar Kilaru and

Gero Steinberg as described in Zala *et al* 2018 [79]. All strains were grown in YSB

(yeast extract 10 g/L and sucrose 10 g/L; pH 6.8). Depending on the tested

environment, we used either YSB or a defined salts medium (Minimal Medium – MM, pH

5.8; [80]. Each strain was stored in glycerol at -80°C until required and then recovered in

YSB medium incubated at 18°C for three days. For nutrient limitation experiments, the

cells were washed three times with sterile distilled water to remove any remaining

nutrients prior to placing them into nutrient-limited conditions. Cell concentrations were

determined by counting blastospores using a KOVA cell chamber system (KOVA

International Inc., USA).

Effect of temperature on growth morphology.

Blastospores of each strain were inoculated on YSB at a final concentration of 10^5

blastospores/mL. Three flasks were incubated at 18°C (as control) and three at 27°C for

four days. An aliquot was taken from each flask at 8, 24, 48, 96, and 144 hai to monitor

cell morphology by light microscopy using a Leica DM2500 microscope with LAS version

4.6.0 software. These experiments were repeated three times.

Effect of nutrient limitation on growth morphology.

MM was used to monitor the transition in growth morphologies under different nutrient limitations. The effect of carbon depletion was assessed by adding each strain at a final concentration of 10^5 blastospores/mL into either regular MM, which contains sucrose as the main carbon source, or into MM without sucrose (C-depleted medium) and incubated at 18°C. The cell morphology was observed by light microcopy every 24 hours until 144 hai. The effect of nitrogen depletion was determined using MM lacking ammonium nitrate, which is normally present in MM as the main nitrogen source. We compared the growth morphologies of cells growing in MM with and without ammonium nitrate by light microcopy during 96 hours of incubation at 18°C. The growth morphologies in both nutrient-limited conditions were compared to the morphologies found in the nutrient-rich YSB. Experiments were repeated three times for each nutrient environment.

Effect of oxidative stress on growth morphology.

H_2O_2 was added into YSB at different concentrations (0, 0.5, 1, 2, 5, and 10 mM) to induce different levels of exogenous oxidative stress. Each strain was added to a final concentration of 10^5 blastospores/mL and incubated at 18°C. Morphological changes and cellular morphologies were observed by light microscopy at 72 hai. To determine whether a lower concentration of blastospores promotes a higher susceptibility to oxidative stress, we performed the same experiment using a final concentration of 10^3 blastospores/mL. These experiments were repeated twice.

Effect of inoculum size on growth morphology.

Flasks containing YSB or C-depleted medium were inoculated with a final cell density of

10^7 blastospores/mL, followed by a 10X serial dilution to a concentration of 10^3

blastospores/mL. Flasks containing C-depleted medium were incubated at 18°C and

flasks with YSB were incubated at 27°C. Cell morphology was determined every 24 hours

using light microscopy over 144 hours of incubation. These experiments were repeated

three times.

RNA extraction, library preparation, and sequencing.

Flasks containing a final concentration of 10^5 blastospores/mL in YSB or C-depleted

medium were incubated at 18°C to induce blastosporulation or mycelial growth,

respectively. Total RNA was extracted by using TRIzol (Thermo Fisher Scientific,

Waltham, Massachusetts) following the manufacturer's instructions. RNA quality was

checked using a Qubit fluorometer (Thermo Fisher Scientific, Waltham, Massachusetts)

and a TapeStation 2200 (Agilent, Santa Clara, California). Truseq stranded mRNA kit

(Illumina Inc., San Diego, CA, USA) was used for library preparation, and ribosomal RNA

was depleted by polyA enrichment. The quality and quantity of the enriched libraries were

assessed using a Qubit fluorometer (Thermo Fisher Scientific) and TapeStation (Agilent).

Libraries were sequenced on the Illumina HiSeq 2500 at 4 x 100 bp paired end reads.

Transcriptome mapping and quantification.

The sequencing adapters and reads shorter than 15 bp were trimmed from raw Illumina

reads using Trimmomatic v.0.36 [81] with the following settings:

ILLUMINACLIP:TrueSeq3-PE.fa:2:30:10, LEADING:2 TRAILING:2,

SLIDINGWINDOW:4:15 MINLEN:15. Filtered reads of each Z. tritici strain for the most

common growth morphologies (blastospores and mycelial growth) were mapped against

their respective genomes using Tophat v.2.0.13 [82]. HTSeq-count v.0.6.1 was used to

calculate the gene counts [83]. Mapped reads with more than one reported alignment

were excluded from further analyses. We applied the TMM (trimmed mean of M values)

method implemented in the Bioconductor edgeR v.3.12.0 package to normalize the

gene counts and to calculate the differentially expressed genes under each condition

[84,85]. Only genes with average counts per million reads per sample >1 for at least one

growth condition were counted as expressed genes. The Benjamin-Hochberg false

discovery rate (FDR) correction was used to adjust p-values based on the Fisher exact

test.

Transcription profile analysis.

We generated the CPM (counts per million mapped reads) values based on TMM-

normalized library sizes obtained from the edgeR v.3.12.0 package [84]. Only genes

with adjusted p-values of FDR≤0.01 were included in the analysis. The published

transcriptomes of each strain during wheat infection at 7 dpi [41] were used for

comparative analyses (datasets in NCBI SRA SRP077418). Only genes that were

significantly upregulated during both wheat infection at 7 dpi and in C-depleted medium

compared to YSB were considered to be mycelial growth-related genes. These genes

were used in further analyses. Specific transcriptome signatures were identified based

on the differentially expressed genes that were shared among all four strains under each

growth condition, according to the *Z. tritici* reference genome annotations [86,87]. We

also used the published transcriptomes of different stages of wheat infection [41] to

interpret the expression profiles of some genes under both axenic and *in planta*

conditions. The $\log_2$-transformed CPM values of those genes were plotted using the R

package ggplot2 [88].

Fluorescence microscopy to quantify chitin and lipids in different morphotypes.

Blastospores of 1A5 were inoculated into YSB to a final concentration of 10^5

blastospores/mL and incubated at 18°C or 27°C to induce blastosporulation or

chlamydospore-like formation, respectively. 1A5 was also inoculated at the same

concentration into C-depleted medium and incubated at 18°C to induce hyphal growth.

After four days of incubation, the resulting cells were harvested and fixed with 70% (v/v)

ethanol for 30 minutes, followed by three washes with phosphate buffer saline (PBS). The

cells were stained with 1 μg/mL of CFW and 2.5 μg/mL of NR (both reagents from Sigma-

Aldrich Chemie Gmbh, Munich, Germany) for 15 minutes. The stained cells were viewed

with a Leica DM2500 fluorescence microscope with LAS version 4.6.0 software, using a

UV filter system for CFW consisting of a BP excitation filter at 340-380 nm and a long

pass emission filter (>425 nm), and a red filter system for NR consisting of a BP excitation

filter at 546/10 nm and an emission filter at 585/40 nm.

Transmission electron microscopy.

Cells of 1A5 and 3D7 were collected after 4 days of growth in YSB at 18°C or 27°C.

Pycnidiospores for each strain were harvested from infected wheat plants. Samples were

prepared using microwave-assisted fixation in 0.1 M cacodylate buffer (2.5%

glutaraldehyde and 2% paraformaldehyde), using a PELCO Biowave (Ted Pella,

California, USA), followed by embedding in 2% low melting temperature agarose

(SeaPlaque Agarose, FMC Corporation, USA). A second step of microwave-assisted

fixation was used after embedding in agarose. Before resin infiltration, the samples went

through two steps of post-fixation, first in osmium tetroxide (OsO4) in 0.1 M cacodylate

buffer and then in 1% uranyl acetate (UrAc), followed by a graded series of ethanol

washes for dehydration (25%, 50%, 75%, 90%, and 100%). Samples were infiltrated with

Spurr resin and polymerized at 60°C for 48 hours. Ultrathin sections (60 nm) were cut

using the Leica Ultracut FC6 (Leica Microsystem, Vienna) and then transferred to formvar

coated carbon grids, followed by post-staining with uranyl acetate and lead citrate.

Micrographs were collected using a Morgagni 268 transmission electron microscope

operating at 100 kV and using a CCD (1376x1032 pixel) camera.

Germination of chlamydospore-like cells and their survival under stress.

Chlamydospore-like cells and blastospores of the 1A5 strain were harvested from YSB

after 4 days of growth at 27°C or 18°C, respectively, and their concentrations were

adjusted to ~10^5 cells/mL. 10 μL was spotted onto WA (1% agar in water) and the plates

were incubated at 18°C. Germ tube formation was checked at 0, 12, and 24 hours after

inoculation by light microscopy.

To determine whether chlamydospore-like cells differ in their ability to survive

environmental stresses compared to blastospores and pseudohyphae, we exposed $8x10^3$

cells/mL of each morphotype to different stresses, including drought and extreme

temperatures.

For drought stress, 200 μL of each cell solution were spread onto Amersham Hybond™

N$^+$ nylon membranes (GE Healthcare Life Sciences, Chicago, Illinois) and dried in a sterile

Petri dish placed on a sterile bench in a laminar flow hood. The Petri dishes were then

placed into a sealed box containing anhydrous silica gel for 1, 3, 5, or 10 days. After the

drought stress, the membranes were placed onto YMA plates (4g/L yeast extract, 4g/L

malt extract, 4g/L sucrose, and 12g/L agar), incubated at 18°C and colony formation was

monitored. Control membranes containing each morphotype were transferred directly to

YMA plates without any drying treatments.

For cold stress, 300 μL aliquots of each cell solution were submerged into liquid nitrogen

for 1, 2, or 3 minutes, and 200 μL of the thawed cell solution were spread onto YMA plates.

Control samples were incubated at room temperature. For heat stress, 200 μL aliquots of

each morphotype were incubated for 24 hours at 30, 35, or 40°C, followed by plating on

YMA. Controls were kept at 18°C. YMA plates from control and treated samples were

incubated at 18°C for seven days. Survival rates for each cell type were estimated based

on the number of colonies formed after exposure to the stressful conditions compared to

the number of colonies formed by the controls.

Plant infection assays.

The susceptible wheat cultivar Drifter was used in all experiments. Plant growth

conditions were described earlier [89]. All infection assays used 16-day-old plants. For

all four strains, blastospore suspensions produced after 3 days of growth in YSB at 18°C

were filtered through two layers of cheesecloth and diluted to a final concentration of 10^6

blastospores/mL in 30 mL of sterile water supplemented with 0.1% (v/v) Tween 20.

Spore suspensions were applied to run-off using a sprayer and the plants were kept for

three days in sealed plastic bags, followed by 21 days in a greenhouse. Leaves with

pycnidia were harvested and transferred to a 50 ml Falcon tube containing sterile water

and gently agitated to harvest the pycnidiospores. Pycnidiospore suspensions were

adjusted to a final concentration of 10^6 pycnidiospores/mL and a new batch of plants

were inoculated as described above. The plants infected by pycnidiospores were used

to observe the formation of chlamydospore-like cells on wheat plants.

To determine whether chlamydospore-like cells produce germ tubes and initiate hyphal

growth after coming into contact with the host, we inoculated Drifter with chlamydospore-

like cells from the 1A5 strain tagged with cytoplasmic GFP. Production of chlamydospore-

like cells was induced using high temperature (YSB at 27°C) as described above. The

suspension of chlamydospore-like cells was prepared and concentrations were

standardized as described for blastospore infections. The inoculation procedure was the

same except the inoculated plants were kept in plastic bags for only 24 hours.

To observe formation of chlamydospore-like cells *in planta*, infected leaves from wheat

plants inoculated with pycnidiospores were harvested at 20 dpi. The infected leaves were

incubated in a humidity chamber for 24 hours, following by washes with sterile distilled

water to collect the released spores. The spore suspension was checked for the presence

of chlamydospore-like cells using the fluorescent microcope with a GFP filter consisting

of a BP excitation filter at 480/40 nm and a BP emission filter at 527/30 nm.

To observe blastospore formation on the surface of wheat leaves, we inoculated plants

using blastospores of the 3D7 strain tagged with cytoplasmic GFP. At 21 dpi, GFP-tagged

3D7 pycnidiospores were harvested and a new batch of plants was inoculated using these

pycnidiospores as described earlier. The inoculated plants were kept in plastic bags at

100% humidity until they were examined for blastospores at 24 and 48 hai.

Confocal microscopy of germinated pycnidiospores and chlamydospore-like cells.

Confocal images were obtained using a Zeiss LSM 780 inverted microscope with ZEN

Black 2012 software. An argon laser at 500 nm was used to excite GFP fluorescence and

chloroplast autofluorescence with an emission wavelength of 490-535 nm and 624-682

nm, respectively. Plants inoculated with GFP-tagged 1A5 chlamydospore-like cells were

checked at 24 hai to observe germination of the chlamydospore-like cells. The same

confocal settings were used to determine whether pycnidiospores produced blastospores

on the leaf surface. Wheat plants inoculated with GFP-tagged 3D7 pycnidiospores were

checked at 24 and 48 hai.

Acknowledgements

We acknowledge Gero Steinberg and Sreedhar Kilaru for generating the fluorescent *Z.*

tritici Swiss strains used in this study, and Andrea Sanchez-Vallet for providing the strains.

We thank Daniel Croll and Andrea Sanchez-Vallet for critical reading of the manuscript.

We also thank Andrea Sanchez-Vallet, Lukas Meile, Julien P. L. Alassimone, and Danilo

A. Pereira for discussions in the course of this work. RNA processing and sequencing

were performed in the Genetic Diversity Center (GDC) of ETH Zurich and the Genomics

Facility of ETH Basel, respectively. Transmission electron microscopy (TEM) and confocal

microscopy were performed at the Scientific Center for Optical and Electron Microscopy

(ScopeM) of ETH Zurich.

References

1. Brown AJP, Cowen LE, di Pietro A, Quinn J. Stress Adaptation. Microbiology

Spectrum. 2017;5. doi:10.1128/microbiolspec.funk-0048-2016

2. Lin X, Alspaugh AJ, Liu H, Harris S. Fungal morphogenesis. Cold Spring Harbor

perspectives in medicine. 2014;5: a019679. doi:10.1101/cshperspect.a019679

3. Mayer FL, Wilson D, Hube B. *Candida albicans* pathogenicity mechanisms. Virulence.

2013;4: 119–128. doi:10.4161/viru.22913

4. Wang L, Lin X. Morphogenesis in fungal pathogenicity: shape, size, and surface.

PLoS Pathogens. 2012;8: e1003027. doi:10.1371/journal.ppat.1003027

5. Kumamoto CA, Vinces MD. Contributions of hyphae and hypha- co- regulated genes

to *Candida albicans* virulence. Cellular Microbiology. 2005;7: 1546–1554.

doi:10.1111/j.1462-5822.2005.00616.x

6. Brand A. Hyphal growth in human fungal pathogens and its role in virulence.

International Journal of Microbiology. 2012;2012: 517529. doi:10.1155/2012/517529

7. Thompson DS, Carlisle PL, Kadosh D. Coevolution of morphology and virulence in

Candida species. Eukaryotic Cell. 2011;10: 1173–1182. doi:10.1128/ec.05085-11

8. Lübbehüsen TL, Nielsen J, McIntyre M. Morphology and physiology of the dimorphic

fungus *Mucor circinelloides* (syn. *M. racemosus*) during anaerobic growth. Mycological

Research. 2003;107: 223–230. doi:10.1017/s0953756203007299

9. Nadal M, García- Pedrajas MD, Gold SE. Dimorphism in fungal plant pathogens.

FEMS Microbiology Letters. 2008;284: 127–134. doi:10.1111/j.1574-6968.2008.01173.x

10. Klein BS, Tebbets B. Dimorphism and virulence in fungi. Current Opinion in

Microbiology. 2007;10: 314–319. doi:10.1016/j.mib.2007.04.002

11. Nigg M, Laroche J, Landry CR, Bernier L. RNAseq analysis highlights specific

transcriptome signatures of yeast and mycelial growth phases in the Dutch Elm disease

fungus *Ophiostoma* novo-*ulmi*. G3: Genes|Genomes|Genetics. 2015;5: 2487–2495.

doi:10.1534/g3.115.021022

12. Edwards JA, Chen C, Kemski MM, Hu J, Mitchell TK, Rappleye CA. *Histoplasma*

yeast and mycelial transcriptomes reveal pathogenic-phase and lineage-specific gene

expression profiles. BMC Genomics. 2013;14: 1–19. doi:10.1186/1471-2164-14-695

13. Veses V, Gow NA. Pseudohypha budding patterns of *Candida albicans*. Medical

Mycology. 2009;47: 268–75. doi:10.1080/13693780802245474

14. Gancedo JM. Control of pseudohyphae formation in *Saccharomyces cerevisiae*.

FEMS Microbiology Reviews. 2001;25: 107–123. doi:10.1111/j.1574-

976.2001.tb00573.x

15. Noble SM, Gianetti BA, Witchley JN. *Candida albicans* cell-type switching and

functional plasticity in the mammalian host. Nature Reviews Microbiology. 2017;15: 96–

108. doi:10.1038/nrmicro.2016.157

16. Couteaudier Y, Alabouvette C. Survival and inoculum potential of conidia and

chlamydospores of *Fusarium oxysporum* f.sp. *lini* in soil. Canadian Journal of

Microbiology. 1990;36: 551–556. doi:10.1139/m90-096

17. Lin X, Heitman J. Chlamydospore formation during hyphal growth in *Cryptococcus*

neoformans. Eukaryotic Cell. 2005;4: 1746–54. doi:10.1128/EC.4.10.1746-1754.2005

18. McNeel DJ, Kulkarni RK, Nickerson KW. Pleomorphism in *Ceratocystis ulmi*:

chlamydospore formation. Canadian Journal of Botany. 1983;61: 1349–1352.

doi:10.1139/b83-143

19. Son H, Lee J, Lee YW. Mannitol induces the conversion of conidia to

chlamydospore-like structures that confer enhanced tolerance to heat, drought, and UV

in *Gibberella zeae*. Microbiological Research. 2012;167: 608–615.

doi:10.1016/j.micres.2012.04.001

20. Miyaji M, Sano A, Sharmin S, Kamei K, Nishimura K. The role of chlamydospores of

Paracoccidioides brasiliensis. Nippon Ishinkin Gakkai Zasshi. 2003;44: 133–138.

doi:10.3314/jjmm.44.133

21. Hoes JA. Development of chlamydospore in *Verticillium nigrescens* and *Verticillium*

nubilum. Canadian Journal of Botany. 1971;49: 1836–1866. doi:10.1139/b71-263

22. Schade D, Walther A, Wendland J. The development of a transformation system for

the dimorphic plant pathogen *Holleya sinecauda* based on Ashbya gossypii DNA

elements. Fungal Genetics and Biology. 2003;40: 65–71. doi:10.1016/s1087-

1845(03)00064-1

23. Fones H, Gurr S. The impact of Septoria tritici Blotch disease on wheat: An EU

perspective. Fungal Genetics and Biology. 2015;79: 3–7. doi:10.1016/j.fgb.2015.04.004

24. Torriani SF, Melichar JP, Mills C, Pain N, Sierotzki H, Courbot M. *Zymoseptoria*

tritici: A major threat to wheat production, integrated approaches to control. Fungal

Genetics and Biology. 2015;79: 8–12. doi:10.1016/j.fgb.2015.04.010

25. Kema GHJ, Yu D, Rijkenberg FHJ, Shaw MW, Baayen RP. Histology of the

pathogenesis of *Mycosphaerella graminicola* in wheat. Phytopathology. 1996;86: 777–

786. doi:10.1094/Phyto-86-777

26. Steinberg G. Cell biology of *Zymoseptoria tritici*: Pathogen cell organization and

wheat infection. Fungal Genetics and Biology. 2015;79: 17–23.

doi:10.1016/j.fgb.2015.04.002

27. King R, Urban M, Lauder RP, Hawkins N, Evans M, Plummer A, et al. A conserved

fungal glycosyltransferase facilitates pathogenesis of plants by enabling hyphal growth

on solid surfaces. PLoS Pathogens. 2017;13: e1006672.

doi:10.1371/journal.ppat.1006672

28. Motteram J, Lovegrove A, Pirie E, Marsh J, Devonshire J, van de Meene A, et al.

Aberrant protein N- glycosylation impacts upon infection- related growth transitions of

the haploid plant- pathogenic fungus *Mycosphaerella graminicola*. Molecular

Microbiology. 2011;81: 415–433. doi:10.1111/j.1365-2958.2011.07701.x

29. Choi YE, Goodwin SB. Gene encoding a c-Type cyclin in *Mycosphaerella*

graminicola is involved in aerial mycelium formation, filamentous growth, hyphal

swelling, melanin biosynthesis, stress response, and pathogenicity. Molecular Plant-

Microbe Interactions. 2011;24: 469–477. doi:10.1094/MPMI-04-10-0090

30. Cousin A, Mehrabi R, Guilleroux M, Dufresne M, Lee T, Waalwijk C, et al. The MAP

kinase- encoding gene MgFus3 of the non- appressorium phytopathogen

Mycosphaerella graminicola is required for penetration and *in vitro* pycnidia formation.

Molecular Plant Pathology. 2006;7: 269–278. doi:10.1111/j.1364-3703.2006.00337.x

31. Mehrabi R, M'Barek S, van der Lee TA, Waalwijk C, Wit P de, Kema GHJ. Gα and

Gβ proteins regulate the cyclic AMP pathway that is required for development and

pathogenicity of the phytopathogen *Mycosphaerella graminicola*. Eukaryotic Cell.

2009;8: 1001–1013. doi:10.1128/ec.00258-08

32. Mehrabi R, Kema GHJ. Protein kinase A subunits of the ascomycete pathogen

Mycosphaerella graminicola regulate asexual fructification, filamentation, melanization

and osmosensing. Molecular Plant Pathology. 2006;7: 565–577. doi:10.1111/j.1364-

3703.2006.00361.x

33. Mehrabi R, van der Lee T, Kema GHJ. MgSlt2, a cellular integrity MAP kinase gene

of fungal wheat pathogen *Mycosphaerella graminicola*, is dispensable for penetration

but essential for invasive growth. Molecular Plant-Microbe Interactions. 2006;9: 389–

398. doi:10.1094/MPMI-19-0389

34. Gohari AM, Mehrabi R, Robert O, Ince I, Boeren S, Schuster M, et al. Molecular

characterization and functional analyses of ZtWor1, a transcriptional regulator of the

fungal wheat pathogen *Zymoseptoria tritici*. Molecular Plant Pathology. 2014;15: 394–

405. doi:10.1111/mpp.12102

35. Mohammadi N, Mehrabi R, Gohari AM, Goltapeh E, Safaie N, Kema GHJ. The ZtVf1

transcription factor regulates development and virulence in the foliar wheat pathogen

Zymoseptoria tritici. Fungal Genetics and Biology. 2017;109: 26–35.

doi:10.1016/j.fgb.2017.10.003

36. Yemelin A, Brauchler A, Jacob S, Laufer J, Heck L, Foster AJ, et al. Identification of

factors involved in dimorphism and pathogenicity of *Zymoseptoria tritici*. PLoS ONE.

2017;12: e0183065. doi:10.1371/journal.pone.0183065

37. Mehrabi R, Zwiers LH, de Waard MA de, Kema GHJ. MgHog1 regulates dimorphism

and pathogenicity in the fungal wheat pathogen *Mycosphaerella graminicola*. Molecular

Plant-Microbe Interactions. 2006;19: 1262–1269. doi:10.1094/mpmi-19-1262

38. Hornby JM, Jacobitz-Kizzier SM, McNeel DJ, Jensen EC, Treves DS, Nickerson KW.

Inoculum size effect in dimorphic fungi: extracellular control of yeast-mycelium

dimorphism in *Ceratocystis ulmi*. Applied and Environmental Microbiology. 2004;70:

1356–9.

39. Hornby JM, Jensen EC, Lisec AD, Tasto JJ, Jahnke B, Shoemaker R, et al. Quorum

sensing in the dimorphic fungus *Candida albicans* is mediated by farnesol. Applied and

Environmental Microbiology. 2001;67: 2982–2992. doi:10.1128/AEM.67.7.2982-

2992.2001

40. Wedge ME, Naruzawa ES, Nigg M, Bernier L. Diversity in yeast–mycelium

dimorphism response of the Dutch Elm disease pathogens: the inoculum size effect.

Canadian Journal of Microbiology. 2016;62: 525–529. doi:10.1139/cjm-2015-0795

41. Palma-Guerrero J, Ma X, Torriani SF, Zala M, Francisco CS, Hartmann FE, et al.

Comparative transcriptome analyses in *Zymoseptoria tritici* reveal significant differences

in gene expression among strains during plant infection. Molecular Plant-Microbe

Interactions. 2017; doi:10.1094/MPMI-07-16-0146-R

42. Böttcher B, Pöllath C, Staib P, Hube B, Brunke S. *Candida* species rewired hyphae

developmental programs for chlamydospore formation. Frontiers in Microbiology.

2016;7: 1697. doi:10.3389/fmicb.2016.01697

43. Bago B, Zipfel W, Williams RM, Jun J, Arreola R, Lammers PJ, et al. Translocation

and utilization of fungal storage lipid in the arbuscular mycorrhizal symbiosis. Plant

Physiology. 2002;128: 108–124. doi:10.1104/pp.010466

44. Fabry W, Schmid E, Schraps M, Ansorg R. Isolation and purification of

chlamydospores of *Candida albicans*. Medical Mycology. 2003;41: 53–58.

doi:10.1080/mmy.41.1.53.58

45. Sitton JW, Cook RJ. Comparative morphology and survival of chlamydospores of

Fusarium roseum "Culmorum" and "Graminearum." Phytopathology. 1981;71: 85–90.

doi:10.1094/Phyto-71-85

46. Biswas S, van Dijck P, Datta A. Environmental sensing and signal transduction

pathways regulating morphopathogenic determinants of *Candida albicans*. Microbiology

and Molecular Biology Reviews. 2007;71: 348–376. doi:10.1128/mmbr.00009-06

47. Gauthier GM. Dimorphism in fungal pathogens of mammals, plants, and insects.

PLoS Pathogens. 2015;11: e1004608. doi:10.1371/journal.ppat.1004608

48. Derridj S. Nutrients on the leaf surface. In : Morris CE, Nicot PC, Nguyen-The C,

editors. Aerial Plant Surface Microbiology. New York: Plenum Press; 1996. p. 25-42.

49. Ohm RA, Feau N, Henrissat B, Schoch CL, Horwitz BA, Barry KW, et al. Diverse

lifestyles and strategies of plant pathogenesis encoded in the genomes of eighteen

Dothideomycetes fungi. PLoS Pathogens. 2012;8: e1003037.

doi:10.1371/journal.ppat.1003037

50. Fleißner A, Sarkar S, Jacobson DJ, Roca GM, Read ND, Glass LN. The so locus is

required for vegetative cell fusion and postfertilization events in *Neurospora crassa*.

Eukaryotic Cell. 2005;4: 920–930. doi:10.1128/ec.4.5.920-930.2005

51. Glass NL, Rasmussen C, Roca MG, Read ND. Hyphal homing, fusion and mycelial

interconnectedness. Trends in Microbiology. 2004;12: 135–141.

doi:10.1016/j.tim.2004.01.007

52. Craven KD, Vélëz H, Cho Y, Lawrence CB, Mitchell TK. Anastomosis is required for

virulence of the fungal necrotroph *Alternaria brassicicola*. Eukaryotic Cell. 2008;7: 675–

683. doi:10.1128/ec.00423-07

53. Rosales RCP, di Pietro A. Vegetative hyphal fusion is not essential for plant infection

by *Fusarium oxysporum*. Eukaryotic Cell. 2008;7: 162–171. doi:10.1128/ec.00258-07

54. Shetty NP, Mehrabi R, Lütken H, Haldrup A, Kema GHJ, Collinge DB, et al. Role of

hydrogen peroxide during the interaction between the hemibiotrophic fungal pathogen

Septoria tritici and wheat. New Phytologist. 2007;174: 637–647. doi:10.1111/j.1469-

8137.2007.02026.x

55. Fones H, Preston GM. Reactive oxygen and oxidative stress tolerance in plant

pathogenic *Pseudomonas*. FEMS Microbiology Letters. 2012;327: 1–8.

doi:10.1111/j.1574-6968.2011.02449.x

56. Yang C, Li W, Cao J, Meng F, Yu Y, Huang J, et al. Activation of ethylene signaling

pathways enhances disease resistance by regulating ROS and phytoalexin production in

rice. The Plant Journal. 2017;89: 338–353. doi:10.1111/tpj.13388

57. Lamb C, Dixon RA. The oxidative burst in plant disease resistance. Annual Review

of Plant Physiology and Plant Molecular Biology. 1997;48: 251–275.

doi:10.1146/annurev.arplant.48.1.251

58. Nasution O, Srinivasa K, Kim M, Kim YJ, Kim W, Jeong W, et al. Hydrogen peroxide

induces hyphal differentiation in *Candida albicans*. Eukaryotic Cell. 2008;7: 2008–2011.

doi:10.1128/ec.00105-08

59. Levy E, Eyal Z, Hochmant A. Purification and characterization of a catalase-

peroxidase from the fungus *Septoria tritici*. Archives of Biochemistry and Biophysics.

1992;296: 321–327. doi:10.1016/0003-9861(92)90579-l

60. do Amaral AM, Antoniw J, Rudd JJ, Hammond-Kosack KE. Defining the predicted

protein secretome of the fungal wheat leaf pathogen *Mycosphaerella graminicola*. PLoS

ONE. 2012;7: e49904. doi:10.1371/journal.pone.0049904

61. Bignell E. The molecular basis of pH sensing, signaling, and homeostasis in fungi.

Advances in applied microbiology. 2012;79: 1–18. doi:10.1016/b978-0-12-394318-

7.00001-2

62. Ruiz-Herrera J, Leon CG, Guevara-Olvera L, Carabez-Trejo A. Yeast-mycelial

dimorphism of haploid and diploid strains of *Ustilago maydis*. Microbiology. 1995;141:

695–703. doi:10.1099/13500872-141-3-695

63. Richards A, Gow NA, Veses V. Identification of vacuole defects in fungi. Journal of

Microbiological Methods. 2012;91: 155–163. doi:10.1016/j.mimet.2012.08.002

64. Veses V, Gow NA. Vacuolar dynamics during the morphogenetic transition in

Candida albicans. FEMS Yeast Research. 2008;8: 1339–1348. doi:10.1111/j.1567-

1364.2008.00447.x

65. Edwards MC, Bowling DJF. The growth of rust germ tubes towards stomata in relation to pH gradients. Physiological and Molecular Plant Pathology. 1986;29: 185– 196. doi:10.1016/s0048-4059(86)80020-0

66. Lendenmann MH, Croll D, Palma-Guerrero J, Stewart EL, McDonald BA. QTL mapping of temperature sensitivity reveals candidate genes for thermal adaptation and growth morphology in the plant pathogenic fungus *Zymoseptoria tritici*. Heredity. 2016;116: 384. doi:10.1038/hdy.2015.111

67. Ponomarenko A, Goodwin SB, Kema GHJ. Septoria tritici Blotch (STB) of wheat. Plant Health Instructor. 2011; doi:10.1094/PHI-I-2011-0407-01

68. Nemecek JC, Wüthrich M, Klein BS. Global control of dimorphism and virulence in fungi. Science. 2006;312: 583–588. doi:10.1126/science.1124105

69. Coleman M, Henricot B, Arnau J, Oliver RP. Starvation-induced genes of the tomato pathogen *Cladosporium fulvum* are also induced during growth in planta. Molecular Plant-Microbe Interactions. 1997;10: 1106–1109. doi:10.1094/MPMI.1997.10.9.1106

70. Talbot NJ, Ebbole DJ, Hamer JE. Identification and characterization of MPG1, a gene involved in pathogenicity from the rice blast fungus *Magnaporthe grisea*. The Plant

Cell. 1993;5: 1575–1590. doi:10.1105/tpc.5.11.1575

71. Martin SW, Douglas LM, Konopka JB. Cell cycle dynamics and quorum sensing in

Candida albicans chlamydospores are distinct from budding and hyphal growth.

Eukaryotic Cell. 2005;4: 1191–1202. doi:10.1128/ec.4.7.1191-1202.2005

72. Oliveira RR, Aguiar BM, Tessmann DJ, Pujade-Renaud V, Vida JB. Chlamydospore

formation by *Corynespora cassiicola*. Tropical Plant Pathology. 2012;37: 415–418.

doi:10.1590/s1982-56762012000600006

73. Annone JG. Observacion de conidios secundarios de *Septoria tritici* Rob. ex Desm.

por meio de impresiones foliareas. In : Kohli MM, van Beuningen LT, editors. In :

Conferencia regional sobre la septoriosis del trigo. Centro International de Mejoramiento

de Maiz y Trigo (CIMMYT). Montevideo, 1990. p. 80-87.

74. Djerbi M. Épidémiologie du *Septoria tritici* Rob. et Desm. consevation et mode de

formation de l'inoculum primaire. Société Française de Phytopathologie. 1977; 91–101.

75. Jones GD, Lee NP. Production of secondary conidia by *Septoria tritici* in culture.

Transactions of the British Mycological Society. 1974;62: 212-213. doi:10.1016/s0007-

1536(74)80025-2

76. Zhan J, Linde CC, Jürgens T, Merz U, Steinebrunner F, McDonald BA. Variation for

neutral markers is correlated with variation for quantitative traits in the plant pathogenic

fungus *Mycosphaerella graminicola*. Molecular Ecology. 2005;14: 2683–2693.

doi:10.1111/j.1365-294x.2005.02638.x

77. Croll D, Zala M, McDonald BA. Breakage-fusion-bridge cycles and large insertions

contribute to the rapid evolution of accessory chromosomes in a fungal pathogen. PLoS

Genetics. 2013;9: e1003567. doi:10.1371/journal.pgen.1003567

78. Kilaru S, Schuster M, Studholme D, Soanes D, Lin C, Talbot NJ, Steinberg G. A

codon-optimazed green fluorescent protein for live cell imaging in *Zymoseptoria tritici*.

Fungal Genetics and Biology. 2015;79: 125–131.

79. Zala M, Mikaberidze A, Alassimone JPL, Muhammad A, McDonald BA, Sánchez-

Vallet A. Mixed infections alter virulence dynamics and reduce transmission in the wheat

pathogen *Zymoseptoria tritici*. in submission.

80. Vogel HJ. A convenient growth medium for Neurospora (medium N). Microbial

Genetics Bulletin. 1956;13: 42–43.

81. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics. 2014;30: 2114–2120. doi:10.1093/bioinformatics/btu170

82. Trapnell C, Pachter L, Salzberg SL. TopHat: discovering splice junctions with RNA-

Seq. Bioinformatics. 2009;25: 1105–1111. doi:10.1093/bioinformatics/btp120

83. Anders S, Pyl PT, Huber W. HTSeq—a Python framework to work with high-

throughput sequencing data. Bioinformatics. 2015;31: 166–169.

doi:10.1093/bioinformatics/btu638

1001

1002   84. Robinson MD, Oshlack A. A scaling normalization method for differential expression

1003   analysis of RNA-seq data. Genome Biology. 2010;11: 1–9. doi:10.1186/gb-2010-11-3-

1004   r25

1005

1006   85. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for

1007   differential expression analysis of digital gene expression data. Bioinformatics. 2010;26:

1008   139–140. doi:10.1093/bioinformatics/btp616

1009

1010   86. Grandaubert J, Bhattacharyya A, Stukenbrock EH. RNA-seq based gene annotation

1011   and comparative genomics of four fungal grass pathogens in the genus *Zymoseptoria*

1012   identify novel orphan genes and species-specific invasions of transposable elements.

1013   G3: Genes|Genomes|Genetics. 2015;5: 1323–1333. doi:10.1534/g3.115.017731

1014

1015   87. Plissonneau C, Hartmann FE, Croll D. Pangenome analyses of the wheat pathogen

1016   *Zymoseptoria tritici* reveal the structural basis of a highly plastic eukaryotic genome.

1017   BMC Biology. 2018;16: 5. doi:10.1186/s12915-017-0457-4

1018

1019   88. Wickham H. ggplot2 – Elegant Graphics for Data Analysis . 2nd ed. Springer-Verlag;

1020   2009. p. 260. doi:10.18637/jss.v077.b02

1021

1022   89. Meile L, Croll D, Brunner PC, Plissonneau C, Hartmann FE, McDonald BA, et al. A

1023   fungal avirulence factor encoded in a highly plastic genomic region triggers partial

1024   resistance to Septoria tritici Blotch. New Phytologist. 2018;219: 1048–1061.

1025   doi:10.1111/nph.15180

1026

1027

1028   **Supporting information**

1029

S1 Fig. Effect of nitrogen depletion stress on the cell morphology of four

***Zymoseptoria tritici* strains.** A nutrient-poor medium (MM without ammonium nitrate)

was used to monitor the growth form transition. (A) At 24 hai, only blastospores from the

1E4 strain switched to hyphal growth forming lateral filaments (black arrow). The rest of

the strains remain undifferentiated. Bars represent 10 µm. (B) At 72 hai, blastospores of

all tested *Z. tritici* strains incubated under nitrogen depletion undergo a blastospore-to-

hyphae transition. Branching and hyphal elongation were predominantly observed at this

time point. Bars represent 30 µm. (C) Regular MM containing ammonium nitrate induced

a mixture of morphotypes, such as blastospores, hyphae and budding blastospores from

filamentous hyphae. Bars represent 30 µm.

1040

S2 Fig. Effect of exogenous oxidative stress induced by H_2O_2 on the cell

morphology of four *Zymoseptoria tritici* strains using a high blastospore inoculum

density (10^5 blastospore/mL). A nutrient-rich medium (YSB) containing different

concentrations of H_2O_2 was used to incubate the strains under oxidative stress for 72

hours. Blastopores kept on YSB medium without addition of H_2O_2 continue to multiply as

budding blastospores via blastosporulation and were used as a control (A, B, C, D - bars

represent 50 μm). Despite the increases in H_2O_2 concentration in the medium, no

morphological changes were observed for 1A5 (E, I, M, Q – bars represent 50 μm), 3D1

(G, K – bars represent 50 μm; O and S – bars represent 10 μm), and 3D7 (H, L, P, T -

bars represent 50 μm). The 1E4 strain responded morphologically to H_2O_2 concentrations

at 0.5 mM, 1 mM, and 5 mM (F, J, N - bars represent 50 μm) undergoing the blastospore-

to-hyphae transition. As observed for other strains (Q, S, and T), blastospores of the 1E4

strain remained as undifferentiated structures at 10 mM (R - bars represent 50 μm).

1054

S3 Fig. Effect of exogenous oxidative stress induced by H_2O_2 on the cell

morphology of four *Zymoseptoria tritici* strains using a low blastospore inoculum

density (10^3 blastospore/mL). A nutrient-rich medium (YSB) containing different

concentrations of H_2O_2 was used to incubate the strains under oxidative stress for 96

hours. Blastopores kept on YSB medium without addition of H_2O_2 continued to multiply as

budding blastospores via blastosporulation and were used as a control. At 0.5 mM, only

the 1E4 and 3D1 strains showed hyphal branching and the initial formation of mycelium.

At 1 mM, blastospores from 1E4, 3D1, and 3D7 grew as a mixture of pseudohyphae

(asterisks) and filamentous hyphae (black arrows). At 2 mM, none of the strains changed

their morphology nor were they able to asexually reproduce via blastosporulation.

Blastospores of the 1A5 strain did not switch to different morphotypes in any of the tested

concentrations of H_2O_2. Bars represent 10 μm.

1067

S4 Fig. Cell morphology of four *Zymoseptoria tritici* strains after 144 hours of

incubation on two distinct hyphal induction media and at different initial cell

densities. (A) Blastospores incubated in carbon-depleted MM at 18°C demonstrated a

morphological independence of the inoculum size until the concentration of 10^5

blastospores/mL, where they were able to switch to hyphal growth and to form mycelial

colonies. At higher initial inoculum concentrations ($\geq 10^6$ blastospores/mL), a drastic

reduction of growth form transition and in hyphal length was observed for all tested strains.

(B) Blastospores incubated in YSB medium at 27°C showed significant differences among

the strains. At lower initial inoculum concentrations (10^3 and 10^4 blastospores/mL) only

pseudohyphae and chlamydospores formed. Intermediate inoculum concentrations (10^5

blastospores/mL and 10^6 blastospores/mL) promoted hyphal growth for the 1E4, 3D1 and

3D7 strains. A ten-fold increase in the initial cell density (10^7 blastospores/mL) favored

chlamydospore formation in all strains. 1A5 blastospores were not able to switch to

hyphae and undergo transitions to pseudohyphae and chlamydospores for all tested

concentrations. Bars represent 10 μm.

1083

S5 Fig. Morphological forms of four *Zymoseptoria tritici* strains grown in two distinct morphotype-induction conditions used for the RNA-Seq. Nutrient-rich medium (YSB) induces only blastosporulation. Nutrient-poor medium (MM) and the wheat leaf surface induced the blastospore-to-hyphae transition and mycelial growth in all tested strains. Wheat leaves infected with *Z. tritici* strains expressing cytoplasmic GFP (green color) were analyzed using confocal microscopy. The blue color corresponds to chlorophyll A detected from chloroplast autofluorescence. Bars represent 10 μm.

S6 Fig. Core transcriptome signatures of four *Zymoseptoria tritici* strains from two distinct growth phases. (A-B) Venn diagram of the overexpressed genes (FDR$\leq$0.01) for each strain grown under blastospore and mycelial growth conditions. (C) Graphic representing average expression (log2-transformed counts per million values) of seven genes related to blastosporulation during the blastospore growth phase among strains. (D) Graphic representing average expression (log2-transformed counts per million values) of 38 virulence-related genes during the mycelial growth phase. (E) Graphic representing average expression (log2-transformed counts per million values) of six genes related to cellular morphogenesis during the mycelial growth phase.

S7 Fig. Chlamydospore cells observed in a spore solution harvested from 20-day-old infected leaves. Plants were previously inoculated with pycnidiospores of four *Z. tritici* strains. Free chlamydospores expressing cytoplasmic GFP were frequently observed

1105   mixed with pycnidiospores in the spore solution of all tested strains. Bars represent 30

1106   µm. The proportion of chlamydospores was visibly lower compared to pycnidiospores.

1107

1108   **S1 Table.** Overview of the transcriptomic data of four *Zymoseptoria tritici* strains obtained

1109   from two distinct growth phases (blastospore *vs.* mycelium).

1110

1111   **S2 Table.** The shared 134 genes overexpressed during blastospore growth phase among

1112   four *Zymoseptoria tritici* strains.

1113

1114   **S3 Table.** The shared 134 genes overexpressed during mycelial growth phase among

1115   four *Zymoseptoria tritici* strains.

1116

1117   **S4 Table.** Expression values of seven genes overexpressed in blastospore growth phase

1118   and associated with blastosporulation.

1119

1120   **S5 Table.** Expression values of 39 genes overexpressed in mycelium growth phase and

1121   associated with virulence factors.

1122

1123   **S6 Table.** Expression values of six genes overexpressed in mycelium growth phase and

1124   associated with cellular morphogenesis.

1125

1126

Figures and Tables

Figure Captions

Fig 1. Effect of temperature on the cell morphology of four *Zymoseptoria tritici* strains. (A) Blastospores incubated on nutrient-rich YSB medium at 18°C (the control environment) multiply by budding as blastospores (white triangles). (B) The effect of high temperatures on *Z. tritici* morphology was observed by exposing the strains to 27°C. At 8 hours after incubation (hai) at the high temperature, blastospores of the 1E4 strain were faster to respond to the new environment by switching to filamentous growth. Black triangles point to hyphal branches. (C) At 24 hai, blastospores of the 3D1 and 3D7 strains began growing as hyphae. The 1A5 strain produced swollen cells (black arrows) and pseudohyphae (asterisks), without any evidence of hyphal differentiation. (D) At 48 hai, hyphal branches were abundant for the 1E4, 3D1, and 3D7 strains. The 1A5 strain continued to produce only swollen structures (black arrows). (E) At 96 hai, filamentous growth was observed for the 1E4, 3D1, and 3D7 strains. For the 1A5 strain, high temperature promoted the formation of two new morphotypes: pseudohyphae (asterisks) and chlamydospore-like cells (black arrows). (F) After 144 hours of incubation, the formation of chlamydospore-like cells was observed mainly at the tips of pseudohyphae for the 1A5 strain and at the distal ends of mycelial filaments for the remaining strains (1E4, 3D1, and 3D7) (black arrows). Bars represent 10 µm.

Fig 2. Effect of carbon limitation on the cell morphology of four *Zymoseptoria tritici*

strains. A nutrient-rich medium (YSB) and a carbon-depleted medium (Minimal Media

without sucrose) were used to monitor the growth form transitions. (A) Blastospores

incubated on YSB continued to multiply as budding blastospores and were used as

controls. (B) Under carbon limitation stress, the blastospores of all tested strains began

growing as hyphae after 24 hours of incubation (arrows). (C) At 144 hai, a mycelial growth

was predominant, with frequent vegetative cell fusion events (asterisks) among hyphae.

Bars represent 20 μm.

1156

1157 **Fig 3. Transcriptome signatures of blastospore and mycelial growth phases**

1158 **organized based on enrichment of molecular functional terms.** (A) Common

1159 upregulated genes during blastospore growth among the *Zymoseptoria tritici* strains. (B)

1160 Common upregulated genes for mycelial growth among the *Z. tritici* strains.

1161

Fig 4. Staining of cell wall chitin and lipid content in the different morphotypes of

Zymoseptoria tritici. Cell walls were stained with the chitin-binding dye Calcofluor White

(blue color) and the lipid droplets were stained with Nile Red (red color). Blastospores,

hyphae, and pseudohyphae showed a weaker fluorescence for Calcofluor White

compared to chlamydospores (A), including those formed terminally (B) or intercalarily (C)

via suspensor cells. A high accumulation of lipid droplets was observed in

chlamydospores, as well as in blastospores and pseudohyphae. Hyphae only showed a

high accumulation of lipid droplets in the hyphal branching zone. Bars represent 10 μm.

1170

Fig 5. Chlamydospore cells produced by *Zymoseptoria tritici* have thicker cell walls than pycnidiospores and blastospores. Black triangles point to a detaching daughter cell budding from a blastospore mother cell. Images were obtained by transmission electron microscopy. Two different magnifications are shown for each morphotype: upper panel scale bar = 2000 nm (low magnification), lower panel scale bar = 500 nm (high magnification).

Fig 6. Chlamydospores produced by *Zymoseptoria tritici* are able to germinate both *in vitro* and on the surface of wheat leaves. (A) Blastospores were used as a control. Apical germination and hyphal elongation were observed after 12 hours of incubation on water agar (WA) plates at 18°C. At 24 hai, blastosporulation was observed from the germinated blastospores. (B) Terminal chlamydospores formed via suspensor cells and (C) free-chlamydospores began to germinate after 12 h of incubation on water agar. Hyphal elongation was obvious by 24 h. Bars represent 10 μm. (D) Free-chlamydospores (upper image) and terminal chlamydospores (lower image) expressing GFP (green color) produced germ tubes on the surface of wheat leaves at 24 hours after infection. Blue color corresponds to the autofluorescence detected from chlorophyll A. Asterisks indicate the suspensor cells where chlamydospores are attached. Bars represent 8 μm.

Fig 7. Chlamydospores produced by *Zymoseptoria tritici* are more tolerant to stress than blastospores and pseudohyphae. Survival rate (%) was estimated based upon

the number of colonies formed after exposure to the stressful environment compared to

the number of colonies formed in the control environment after seven days. (A)

Blastospores, pseudohyphae and chlamydospore cells were kept for 1, 3, 5, and 10 days

in a sealed box containing anhydrous silica gel to produce a dry environment. This

simulated drought stress affected blastospore and pseudohyphae survival over time.

Chlamydospores were less affected and exhibited a survival rate of 50% after 10 days

under drought stress. (B) The same three morphotypes were subjected to a cold shock.

Blastospores, pseudohyphae, and chlamydospores were submerged in liquid nitrogen for

1, 2, and 3 minutes at -196 °C. Only chlamydospores survived this cold stress. (C) The

same three morphotypes were subjected to heat stress based on incubation for 24 hours

at 30°C, 35°C or 40°C. Blastospores and chlamydospores were equally affected at 30°C.

The incubation at 35°C killed all the blastospores, while chlamydospore survived at a 12%

rate, double the rate observed for pseudohyphae (6%). No colonies were observed after

incubation at 40°C. One star indicates that the adjusted p value is less than 0.05, two

stars indicates that the p value is less than 0.005, and three stars indicates that the p

value is less than 0.0005. ns indicates no significant difference.

1208

1209   **Fig 8. Germinated pycnidiospores of *Zymoseptoria tritici* produce new**

1210   **blastospores via blastosporulation on the surface of wheat leaves.** (A) At 24 hours

1211   after infection, pycnidiospores (asterisk) germinated and formed hyphal branches

1212   expressing GFP (green color). The germinated pycnidiospores produced new

1213   blastospores on the host leaves (arrow). (B) We observed a significant increase in budding

points on germinated pycnidiospores (arrows), as well as more free blastospores on the

leaf surface at 48 hours after infection. At this time point, we also observed several

vegetative cell fusion events between blastospores and pycnidiospores, which are

characterized by a tubular shaped bridge between the fusing cells (asterisks). Blue color

corresponds to autofluorescence detected from chlorophyll A. Bars represent 10 μm.

Accession numbers

All raw sequencing data were deposited into the NCBI Sequence Read Archive under the

accession number SRP152081.

Author contributions

Conceptualization: Carolina Sardinha Francisco, Javier Palma-Guerrero

Formal analysis: Carolina Sardinha Francisco, Xin Ma, Javier Palma-Guerrero.

Funding acquisition: Bruce A. McDonald

Investigation: Carolina Sardinha Francisco, Maria Manuela Zwyssig, Javier Palma-Guerrero.

Methodology: Carolina Sardinha Francisco, Javier Palma-Guerrero.

Project administration: Javier Palma-Guerrero

Resources: Bruce A. McDonald

Supervision: Javier Palma-Guerrero, Bruce A. McDonald

Validation: Carolina Sardinha Francisco

1235 **Writing – original draft:** Carolina Sardinha Francisco, Javier Palma-Guerrero, Bruce A.

1236 McDonald.

1237 **Writing – review & editing:** Carolina Sardinha Francisco, Javier Palma-Guerrero, Bruce

1238 A. McDonald.

1239

1240 **Financial disclosure statement**

1241 C.S.F. was supported by a PhD research fellowship from CAPES - Brazil (process n°

1242 002087/2015-04). The funders had no role in study design, data collection and analysis,

1243 decision to publish, or preparation of the manuscript.

1244

1245 **Competing interests**

1246 The authors have declared that no competing interests exist.

1247
1248

A. Upregulated genes during blastospore growth

- Cell wall protein (1%)
- Cytoskeleton (2%)
- Glycoside hydrolase (1%)
- Hypothetical protein (20%)
- Secondary metabolite (3%)
- Secretion system (1%)
- Others (13%)
- Cell wall hydrolase (2%)
- RNA processing (2%)
- Transporter (4%)
- Cell signalling (3%)
- Transcription factor (2%)
- Secreted protease (2%)
- Metabolism (44%)

B. Upregulated genes during hyphal growth

- Secreted chloroperoxidase (5%)
- Candidate effector (12%)
- Secreted glycosidase (4%)
- Hypothetical protein (13%)
- Membrane binding protein (4%)
- Metabolism (9%)
- Hydrophobic surface binding (2%)
- Secreted protease (7%)
- Hydrophobin (1%)
- Transcription factor (2%)
- Others (14%)
- Cell signaling (8%)
- Cell wall hydrolase (1%)
- Rna processing (7%)
- Transporter (11%)

Blastospore

Hyphae

Pseudohyphae

Chlamydospores

A

B

C

Pycnidiospore   **Blastospore**   **Chlamydospore**

	In vitro			*In planta*
	0h	**12h**	**24h**	**24h**