Studies of NH_4^+ and NO_3^- uptake ability of subalpine plants and resource-use strategy identified by their functional traits.

Legay N.[1,2*+], Grassein F. [3,4*], Arnoldi C. [1], Segura R. [3,4], Laîné P. [3,4], Lavorel S.[1], Clément J.C.[1,5].

* Shared first co-authorship
+ corresponding author: gabinpiton@gmail.com

[1] Laboratoire d'Ecologie Alpine, CNRS UMR 5553, Université Joseph Fourier, BP 53, 38041 Grenoble Cedex 09, France

[2] INSA Centre Val de Loire, Université de Tours, CNRS, UMR 7324 CITERES, 37200 Tours, France

[3] Université de Caen Basse-Normandie, UMR 950 Ecophysiologie Végétale, Agronomie et nutritions N, C, S, Esplanade de la Paix, CS 14032, 14032 CAEN Cedex 5, France

[4] INRA, UMR 950 Ecophysiologie Végétale, Agronomie et nutritions N, C, S, Esplanade de la Paix, CS 14032, 14032 CAEN Cedex 5, France

[5] CARRTEL, UMR 0042 INRA-Univ. Savoie Mont Blanc, FR-73376, Le Bourget du Lac, France

Running title: plant N uptake and functional strategy

SUMMARY:

The leaf economics spectrum (LES) is based on a suite of leaf traits related to plant functioning and ranges from resource-conservative to resource-acquisitive strategies. However, the relationships with root traits, and the associated belowground plant functioning such as N uptake, including nitrate (NO_3^-) and ammonium (NH_4^+), is still poorly known. Additionally, environmental variations occurring both in time and in space could uncouple LES from root traits. We explored, in subalpine grasslands, the relationships between leaf and root morphological traits for 3 dominant perennial grass species, and to what extent they contribute to the whole-plant economics spectrum. We also investigated the link between this spectrum and NO_3^- and NH_4^+ uptake rates, as well as the variations of uptake across four grasslands differing by the land-use history at peak biomass and in autumn. Although poorly correlated with leaf traits, root traits contributed to an economic spectrum at the whole plant level. Higher NH_4^+ and NO_3^- uptake abilities were associated with the resource-acquisitive strategy. Nonetheless, NH_4^+ and NO_3^- uptake within species varied between land-uses and with sampling time, suggesting that LES and plant traits are good, but still incomplete, descriptors of plant functioning. Soil environmental variations between grasslands and sampling times may also drive to some extent the NH_4^+ and NO_3^- uptake ability of species. Our results support the current efforts to build a more general framework including above- and below-ground processes when studying plant community functioning.

Keywords: Leaf and root traits, Nitrate and ammonium uptake, plant assimilation, Resource use strategy, Subalpine grasslands

Introduction

Functional traits have been widely used to describe different plant strategies. One major axis of specialisation involves a trade-off between conservation of resources in well protected and long lived tissues, and acquisition of resources by tissue with high use-efficiency and turn-over, and commonly referred as the leaf economic spectrum (LES, Wright *et al*. 2004). More specifically, species with an exploitative strategy share similar leaf attributes such as high specific leaf area (SLA) and nitrogen concentrations (LNC) that have been associated with short leaf life-span, high photosynthetic capacity as well as high decomposability (Reich 2014, Cornwell *et al.* 2008). And the conservation/acquisition strategies have been numerously linked to plant ecology, fast-growing exploitative species dominating in nutrient rich environment while the slow-growing conservative species are more common in poor or harsh conditions (Chapin 1980, Ordonez et al. 2009). Despite some evidences of a similar contribution of root traits to the plant strategy (Roumet et al. 2006, Freschet et al. 2010, Fort et al. 2013), the importance of root traits is less understood than for leaf traits, mainly because weak correlations between analogous leaf and root traits have been reported (Craine et al 2005, Tjoelker et al. 2005, Freschet et al. 2010), and also because root functioning is often overlooked compared to leaves in field conditions.

Nutrient uptake ability is one of the main functions provided by roots, and both physiological and morphological adjustments are involved at the root level to support changes in nutrient shoot demand (Hodge 2004, James et al. 2009). Nutrient uptake is indeed influenced both by anatomical properties of roots involved in the prospection such as specific root length, as well as physiological adjustments such as kinetic parameters like maximal uptake rate (Vmax, but see Bassirirad 2000). Among nutrients, nitrogen is one of the most well studied one, especially its inorganic forms (NO_3^- and NH_4^+) both for plants and for ecosystem functioning. However, rarely have morphological and physiological properties of root been assessed

simultaneously in field conditions, whereas some supports come from species grown in

standardized conditions (Maire et al. 2009, Grassein et al. 2015). NH_4^+ and NO_3^- uptake can

indeed be influenced by several environmental factors justifying the use of controlled

conditions to estimate uptake parameters in a comparative purpose. For example, NH_4^+ and

NO_3^- uptake has been reported to vary in response to temperature or pH (Garnett and

Smethurst 1999). Nevertheless, NH_4^+ and NO_3^- uptake ability also differs between species,

and is partially related to plant strategy and their functional traits (Grassein *et al.* 2015), but

these results need to be validated for plant grown in natural conditions. Finally, NH_4^+ and

NO_3^- transporters have two components: a constitutive component and a component induced

by the presence of NH_4^+ and NO_3^- in the soil solution. Thus, it is important to study

interspecific differences for NH_4^+ and NO_3^- uptake at a given site. Otherwise, it is difficult to

interpret differences as resulting from species differences.

Subalpine grasslands are subject to the combined effects of climate and anthropogenic factors,

both influencing N cycling and thus N availability for organisms (Bardgett *et al.* 2005, Legay

et al. 2013). Decreased management intensity favours plant species with resource

conservative traits (Quétier *et al.* 2007), which are usually associated with fungal-dominated

belowground communities (de Vries *et al.* 2012, Grigulis *et al.* 2013). Concomitantly, it slows

down N cycling (Zeller *et al.* 2000, Robson *et al.* 2010), favouring the accumulation of soil

ammonium (NH_4^+) rather than soil nitrate (NO_3^-) (Robson *et al.* 2007). Plants growing in such

variable conditions are likely to adjust their N uptake ability, as it has been shown for

functional traits (Quétier *et al.* 2007, Grassein et al. 2015).

In this study, we investigated the relationships between functional traits and inorganic N

(NH_4^+ and NO_3^-) uptake for three perennial grass species with contrasted leaf economic

strategies. Because soil inorganic NH_4^+ and NO_3^- availability and plant NH_4^+ and NO_3^- uptake

ability are likely to vary across seasons and in response to management (Jaeger *et al.* 1999,

Miller *et al.* 2009), we examined these relationships for individuals occurring in four

subalpine grasslands with different management and throughout the growing season, thereby

testing their temporal consistency. Estimating root NH_4^+ and NO_3^- uptake, and measuring

functional traits for leaves and roots, we tested the following hypotheses: (1) the root traits are

also contributing to the economics spectrum observed for leaf traits, (2) and more exploitative

species should be more efficient to take up both NH_4^+ and NO_3^-. (3) As functional traits are

influenced by environmental conditions, we hypothesise that NH_4^+ and NO_3^- uptake will be

influenced by environmental variations between grasslands, as well as during the growing

season, probably following NH_4^+ and NO_3^- availability, both in quantity and in quality

depending on the most abundant form.

Material and methods

Study site and species

The study site is located in the upper Romanche valley of the central French Alps between the

village of Villar d'Arêne and the Lautaret Pass (Table 1). The climate is subalpine with a

strong continental influence. Winters are cold and snowy, with monthly average minimum

temperatures of -15.9°C in February, maximum monthly average temperature of 23.8°C in

July, and mean annual precipitation of 956mm (unpublished data, sajf.ujf-grenoble.fr). The

growing season starts following snow melt in late April - early May and continues until late

September or October depending on the date of the first snow in autumn.

Given the hypothesis that NH_4^+ and NO_3^- uptake could be an important hard plant trait related

to resource use strategy (as suggested by soft structural and morphological traits) and to field

dominance, and due to the degree of precision chosen for NH_4^+ and NO_3^- uptake estimations

(see 2.2), a compromise was necessary regarding the number of species, grasslands and

replicates to be investigated. This sampling adjustment was required to conduct N uptake

estimations for all individuals in a brief enough time period so that most abiotic factors

remained as comparable as possible (soil moisture, temperature, radiation).

We chose three common and dominant grass species, *Dactylis glomerata* L., *Bromopsis*

erecta (Huds.) Fourr. (formerly *Bromus erectus* (Huds.))   and *Patzkea paniculata* (L.)

G.H.Loos (formerly *Festuca paniculata* (L.) Schinz & Thell.). All species are perennial,

arbuscular mycorrhizal non-dependent species and span a gradient from more exploitative (*D.*

glomerata) to more conservative (*F. paniculata*) resource use strategies (Grassein et al. 2015).

Four grasslands (Table 1), described in Quétier *et al.* (2007), were chosen for their contrasting

past and current management, and were similar to the grasslands studied by Robson *et al.*

(2007, 2010) : (i) Terraced Mown and Fertilized (TMF), (ii) Terraced Unmown not fertilized

but lightly grazed (TU), (iii) Un-terraced Mown grassland (UM) and (iv) Un-terraced

Unmown but lightly grazed grassland (UU), representing a gradient of decreasing

management intensity. To reflect field dominance patterns, *D. glomerata* was sampled in

TMF, *B. erectus* in TU, *F. paniculata* in UM, and all three species were sampled in UU where

they coexist, although *F. paniculata* was dominant (Table 1).

To assess NH_4^+ and NO_3^- uptake patterns over the growing season, the same sampling design

was repeated twice during 2010. At each date for each species and grassland, we sampled the

roots and soil (approximately: 25x25x25 cm) of five individuals (genetically distinct

individuals at least 2m apart). The first sampling corresponded to the peak biomass and

targeted flowering onset (just before anthesis), and the second sampling corresponded to

autumn after last management activities occurred. For *D. glomerata* in TMF and *B. erectus* in

TU, the two sampling dates were mid-June and mid-September. For *F. paniculata* in UM and

the three species in UU, the sampling dates were: early July and early September. These two

dates are called "Summer" and "Autumn" hereafter. As much as possible, species were

sampled at the same time the day to avoid any diurnal variation in N uptake (Gessler et al.

1998). In total, we have sampled 12 points (3 species*2 seasons*2 habitats per species).

Soil nitrogen pools

At each date and for each grassland, soil nitrogen concentrations were measured from six soil

cores (dimensions 4.5 cm Ø, 10 cm deep) kept on ice in the field and maintained at 4°C upon

return to the laboratory (within 2h). Soils were sieved through a 5.6-mm mesh to remove roots

and stones. A subsample of 10g fresh sieved soil was prepared for extraction of inorganic N in

0.5M K_2SO_4, and analysed using a colorimetric analyser (FS-IV autoanalyser (OI-Analytical,

College Station, TX, USA) (following Bowman *et al*. 2003) to measure soil concentrations of

ammonium (NH_4^+), nitrate (NO_3^-) and Total Dissolved Nitrogen (TDN). Soil aliquots were

used to determine soil water (7 days at 70°C) and soil organic matter contents (550°C during

4 hours). Finally, soil subsamples were air-dried to measure soil pH, or ground to a fine

powder for measurements of total carbon (C) and N contents using an elemental analyser

(FlashEA 1112, Thermo Fisher Scientific Inc., Waltham, MA, USA).

At each date, five individuals of each species, with roots and soil, were excavated from each

field, transferred within half an hour to the laboratory located at the Lautaret Pass (Station

Alpine Joseph Fourier) and kept at 4°C until the NO_3^- and NH_4^+ uptake rate measurements to

maintain the functional integrity of the roots. Living and young fine roots were washed with

deionised water, cut to 2-cm length and then, rinsed in 1mM $CaSO_4$ at 4°C for 3 min. The

NO_3^- and NH_4^+ uptake rates were measured during the first hour following plant harvest as

described by Louahlia *et al.* (2000). The optimal conditions for uptake measurements by

excised root determined by Lainé *et al.* (1993) were used in the present study.

Functional traits

Functional traits were measured for roots and leaves using standardised protocols (Perez-Harguindeguy *et al.* 2013). Two of the individual root sub-samples were used to estimate root dry matter content (RDMC), specific root length (SRL, Winrhizo® software, fresh length per unit of dry mass), and were further analysed to obtain root-^{15}N natural abundance and root nitrogen concentration (RNC, N mass per unit of dry mass). Specific leaf area (SLA, fresh area per unit of dry mass), leaf and root dry matter contents (LDMC and RDMC, dry mass per unit of fresh mass), leaf nitrogen concentration (LNC, N mass per unit of dry mass) were also measured.

Nitrogen uptake estimation: the "excised" roots method

Although measuring only a net N uptake, which is the result of influx and efflux, the direct measurement of N uptake using excised roots allows characterising the plant uptake kinetics for NO_3^- and NH_4^+ while controlling for the environmental variations. This makes it possible to compare different species at the cost to lose some of ecological meanings (Lucash *et al.* 2007). This method was thus applied to plants collected in the field. Root N uptake kinetics started within 60 min after excision, thereby avoiding the potential decline in N uptake ability reported to start after 3h (Louahlia *et al.* 2000). Nitrate and ammonium uptake by plants involved mainly the transport system called HATS (High Affinity Transport System). It contributes to N uptake at low to moderate concentrations of external N (<1mM) and saturates at 0.2-0.5 mM (Kronzucker *et al.* 1999, Min *et al.* 2000), which makes it the more likely system used by plants growing in natural and semi-natural ecosystems limited by N (Bassirirad 2000, Maire *et al.* 2009). The estimation of the maximum NH_4^+ and NO_3^- uptake rates by HATS requires a range of N concentrations below 1mM at which the Vmax can be reached depending on species (Grassein *et al.* 2015). Consequently, uptake was estimated from the accumulation of ^{15}N in root sub-samples incubated for one hour in a buffer solution

(pH = 5.5- following Leon *et al.* (1995)), containing a range of N concentrations (20, 50, 100,

250, 500 and 1000 µM). Six sub-samples were incubated in $K^{15}NO_3$ and the other six in

$(^{15}NH_4)_2SO_4$ with a ^{15}N excess of 99% atom. The two N forms were tested individually in

order to avoid possible interactions (Kronzucker *et al.* 1999). Solution volumes and fresh

weights were selected to avoid N depletion during the experiment. After 1h incubation, roots

were washed twice for one minute with a 1mM $CaSO_4$ at 4°C to stop any metabolic

processes. Roots were then dried at 60°C for 72h, ground to a fine powder and analysed by

IRMS at the University of Caen (Isoprime GV instruments, Stockport, UK) to obtain ^{15}N

Atom% and N concentrations.

Data analysis

Nitrogen Uptake Rate (NUR) was calculated for each concentration and each inorganic N

form (NH_4^+ and NO_3^-) using the ^{15}N increase in the root incubated compared to the non-

incubated control, and expressed by unit of time and dry mass ($nmolN.h^{-1}.g^{-1}$ of dry roots, see

Leon *et al.* 1995). The dependence of NUR on substrate concentration was fitted for each

individual and Hanes's relations (Michaelis transformation) was used to estimate the

maximum uptake rate (Vmax) defined as the maximum NUR for NH_4^+ and NO_3^- (Leon *et al.*

1995). Finally, the $NH_4^+:NO_3^-$ uptake ratio was calculated as the ratio between NH_4^+ Vmax

and NO_3^- Vmax.

A principal component analysis (PCA) was performed using all plant functional traits at the

individual level to describe their functional strategy based on leaf and root traits. To

investigate the relationships between functional traits of leaves and roots, and NH_4^+ and NO_3^-

uptake ability (hypothesis 1), we used Pearson correlation coefficients. Relationships between

the functional strategy and uptake of NH_4^+ and NO_3^- at the root level were tested using

regression analyses between the N uptake rates (Vmax) and the first PCA.

Comparisons of NH_4^+ and NO_3^- uptake rates for species (hypothesis 2), fields and date

(hypothesis 3) were conducted with ANOVA followed by Tukey tests to compare species and

grasslands. In details, the effects of sampling time and fields on plant traits within each

species were tested using two-ways ANOVA. Similarly, the effects of sampling time and

fields on maximal NH_4^+ and NO_3^- uptake rates within each species were tested using two-

ways ANOVA. The effects of fields and sampling time on $NH_4^+:NO_3^-$ ratio within each

species were tested using two-ways ANOVA. Then, we tested only in UU grasslands, the

species effect using one-way ANOVA. Finally, we used a two-ways ANOVA and Tukey post

hoc test to test soil parameters differences between fields and dates. Data were log-

transformed when necessary to achieve normality and heteroscedasticity. All statistical

analyses were performed using the software R 3.4.4, with multivariate analyses (PCA) being

performed using the package Ade4 (Dray & Dufour 2007).

Results

We observed large variations for leaf and root functional traits in spite of a restricted number

of species in our study (Table 2). The range of variation was similar to, and sometimes even

higher than the variability reported in Fort *et al.* (2013) for a larger set of species occurring in

a similar ecosystem, including *D. glomerata* and *B. erectus*. The PCA of functional traits

highlighted a first axis explaining 62.1% of the total variance (Fig. 1). The three species

differed significantly for their mean position along this axis (p=0.012), with positive values

for *D. glomerata* and negative values for *F. paniculata*. Positive values along this axis were

characterised by high SLA, LNC and SRL, and low LDMC. Among these, SLA and LNC

have been reported as major contributors to a resource economic spectrum establishing the

existence of a fundamental trade-off between plant features allowing resource capture and

those allowing resource conservation.

This functional axis was positively correlated to NH_4^+ and NO_3^- Vmax (Fig 2a and 2b) and

negatively to NH_4^+:NO_3^- uptake ratio (Fig 2c) indicating a more pronounced preference for

NH_4^+ at lower values of axis 1. Except RDMC, all traits taken separately were poorer

predictors of the NO_3^- and NH_4^+ maximum uptake rates than this functional axis, although the

first PCA axis was significantly correlated with all functional traits (Table 3).

In UU grassland, NH_4^+Vmax in summer was similar for the three species (Fig. 3a) but greater

for *D. glomerata* for NO_3^- Vmax (p <0.001, Fig. 3b). Vmax in autumn for both N forms was

lower for *F. paniculata* compared to the two other species (NH_4^+ p<0.05, NO_3^- p< 0.001).

Comparing the different grasslands within species, we observed reduced NO_3^- and NH_4^+

Vmax values in the UU grassland for *D. glomerata* (in summer and in autumn) and *B. erectus*

(summer) compared to the other grasslands. On the other hand, highest NH_4^+ Vmax for *B.*

erectus and *F. paniculata* were found in UU during the autumn. Illustrating the seasonal

variability, all species in the UU grassland had higher NH_4^+ maximum uptake rates in the

autumn than in the summer, as well as higher NO_3^- uptake for *B. erectus*. NH_4^+:NO_3^- uptake

ratio did not vary in time, but always showed higher values in the UU for the three species

compared to the other grasslands, and overall greater values for *F. paniculata* (Fig. 4).

Within species, a limited number of traits were significantly different between grasslands

(Table 2). We only observed significant differences in autumn, with highest LDMC in TU for

B. erectus, highest LDMC and RDMC in UU for *D. glomerata*, and highest LDMC but lowest

SRL for *F. paniculata* in UM. However, changes in response to the season were more

consistent among species and grasslands, with an increase of LDMC and a decrease of SLA in

autumn compared to the summer. We also observed higher LNC for *D. glomerata* and *F.*

paniculata during the summer than during the autumn in TMF and UM respectively, and

higher RDMC during the autumn for *D. glomerata* in UU.

Since all species occurred in the UU grasslands, we choose to focus on soil parameters from

this grassland. UU and UM only differ for SWC in autumn (Table 4), all other soil variables

were similar between these two grasslands, which had similar past land-use history (Table1).

UU had consistently higher SWC and SOM, and lower soil pH and CN ratio than TMF and

TU. All grasslands had similar soil NH_4^+ concentrations. During the summer, we observed

higher TDN and NH_4^+:NO_3^- soil ratio, and lower soil NO_3^- concentration in UU compared to

TMF and TU, but we did not find these differences in autumn.

Discussion:

Relationships between leaf and root traits

In the aim to find parallels between above and below-ground organs (e.g. Roumet et al. 2006),

several studies have investigated the relationships between analogous traits measured for

leaves and roots. While positive relationships have been reported for SLA *vs* SRL (Craine &

Lee 2003, Craine et al. 2005, Freschet et al. 2010), other studies have reported a lack of

relationships between SLA *vs* SRL (Craine et al. 2001, Tjoelker et al. 2005). In our study, we

did not find any relationships between SRL/SLA, LDMC/RDMC and LNC/RNC, and this

could be related to our limited number of species/replicates. Nevertheless, we observed trade-

offs at the leaf and root levels between traits, namely N concentration and dry matter content.

A similar spectra of resource use at the leaf and root levels has already been reported

(Freschet et al. 2010), despite a relatively low degree of relationships between analogous traits

belowground and aboveground. Different selective pressure for leaf and root traits as well as

specialisation for the acquisition of different resources (e.g. light *vs* nutrient) could explain

this absence of association between belowground and aboveground traits (Craine et al. 2005,

Liu et al. 2010), while the global strategy at the plant level could remain the same since high

efficiency for light or for nutrients could be related to the same physiological adaptation, as

pointed out previously for stress tolerance (Chapin 1980). Although we found that leaf

functional traits (LNC) can be correlated with root NH_4^+ and NO_3^- maximal uptake rate as

previously shown in other studies (Osone et al. 2008; Maire et al. 2009), here root traits

(RDMC, SRL) appeared to be more related to NH_4^+ and NO_3^- uptake rates (Rewald et al.

2014), even if deeper understanding of the relationship between root traits and nutrient

acquisition remains needed (Roumet et al. 2016).

Relationship between N maximum uptake rate (Vmax) and plant strategy

Our results showed that a stronger exploitative syndrome (higher SRL, SLA, LNC and lower

RDMC) was associated with higher Vmax for both inorganic N forms, rejecting the

hypothesis of a trade-off between maximum uptake rate of each N forms. Indeed, ammonium

toxicity has been reported for some plant species (review in Britto & Kronzucker 2002), as

well as negative interactions between the uptake of NH_4^+ and NO_3^- (Kronzucker et al. 1999),

and this could promote a trade-off in the acquisition of NH_4^+ and NO_3^- between species

(Maire et al. 2009). Here, we estimated NH_4^+ and NO_3^- uptake independently to avoid such

interactions during measurements, and our results did not support a trade-off but rather

suggest a synergistic uptake of both N forms. Provision of NO_3^- has been demonstrated to

alleviate the NH_4^+ toxicity (Britto & Kronzucker 2002), and even to favour NH_4^+ uptake. We

indeed observed higher uptakes for NH_4^+ than for NO_3^-, indicating a preference of all species

for NH_4^+, especially for individuals with a more conservative syndrome of traits. Besides,

more exploitative plants have a lower preference for NH_4^+ compared to more conservative

individuals, but expressed higher maximal uptake rates than more conservative individuals for

both N forms. Overall, changes in functional leaf traits related to a higher potential

photosynthesis efficiency and light capture appeared to be associated at the root level with

higher N maximal uptake rates for both N forms.

Nitrogen uptake variations in response to management and sampling dates

Nitrogen uptake rate is usually considered as a property of plant species, but little is

known about variation in within-species N uptake rates in grasslands with different land-use

history and at different times during the growing season. In our study, we observed that

nitrogen uptake rates could differ strongly for the same species in different grasslands (e.g. *B.*

erectus and *D. glomerata* in the UU grassland). On the other hand, the time of the year also

influenced the N uptake rates of all species, with for example an opposite response between

grasslands for *B. erectus* depending on the sampling date. Overall, grasslands were weakly

discriminated by functional traits, suggesting that other factors such as soil parameters may

explain the within species N uptake differences between grasslands.

Nitrogen uptake can vary depending on the amount of N available in the soil (Gavito

et al. 2001). Soil NH_4^+ concentration, the main N source taken up by plants in our study, was

similar in the four investigated grasslands, whereas a higher soil total dissolved N (TDN) was

measured in the UU grassland. Consequently, the lower N uptake rates observed in this

grassland cannot be explained by a lower N availability. As reported by previous studies,

subalpine grasslands can show the legacy effects of former management activities, leading to

slower N cycling (Zeller et al. 2000, Robson et al. 2007). Indeed, we observed lower pH and

higher soil water and organic matter contents in the UU grassland suggesting variations in N

cycling and in the quality of the available N, not only in its quantity (Garnett & Smethurst

1999, Robson et al. 2010). Supporting this hypothesis, we observed variations in soil NO_3^-

concentrations, and consequently soil $NH_4^+:NO_3^-$ ratio, between the studied grasslands.

Although we could not directly relate *in situ* soil parameters to N plant uptake estimated under

"controlled" conditions, we interestingly observed parallel changes for NH_4^+ uptake and soil

$NH_4^+: NO_3^-$ ratio in grasslands where individuals have been sampled. For example, both *B.*

erectus NH_4^+ uptake and soil NO_3^- concentration were lower during the summer and higher

during the autumn in TU than in UU.

Rarely investigated in natural ecosystems, experimental evidences on cultivated plants

have demonstrated the effects of soil $NH_4^+: NO_3^-$ ratio concentration on plant N uptake

(Errebhi & Wilcox 1990, Bar-Tal et al. 2001). Yet, the effects were largely species-dependent

and trade off were sometime reported between NH_4^+ and NO_3^- uptakes (Warncke & Barber

1973, Kronzucker et al. 1999, Maire et al. 2009). The preferential uptake for an inorganic N

form could also be influenced by environmental and physiological factors (Britto &

Kronzucker 2013). Our results did not support any trade-off in the intrinsic ability of plant

species to take up both N forms, even after removing possible environmental conditions or

interactions between N forms. Although we could not directly test for the relationship

between soil parameters and plant N uptake rates, differences between grasslands in the N

uptake within species highlight that management practices may have important effects on

plant N uptake, likely through N cycling changes and the quality of N pool available as

already pointed out by previous studies (Zeller et al. 2000, Robson et al. 2007). Other studies

have suggested that N preference could be dependent of the soil availability of the different N

forms (Näsholm et al. 2009, Stoelken et al. 2010). While our results partially supported this

hypothesis, with variation within species between different grassland, the different species

sampled in the same grassland showed differences in their $NH_4^+:NO_3^-$ uptake ratio, supporting

the hypothesis that this "preference" is partially related to the strategy of species, or at least to

species identity. But overall, more exploitative species with higher maximum uptake rates for

one N form is also likely to have high uptake rates for other N forms (Kastovska &

Santruckova 2011).

Nevertheless, the plant preference for N forms is a complex topic (Britto &

Kronzucker 2013), and careful considerations should be given to the environmental

conditions where the species occur. Since N cycling is controlled by a large set of parameters

such as pH, soil humidity, land-use, short and long-term variations in the predominant N

forms available for plants are to be expected. Under some conditions, plants can also take up

organic N (amino acids) directly and/or through fungi (Näsholm et al. 2000). While we

assumed that this source of N is of limited importance for our species in our relatively fertile

grasslands (Kahmen et al. 2009), a full understanding of the N preference, and discussion

about species coexistence through N forms sharing, would require careful investigations,

beyond the possibility in our study. Nonetheless, the variability we observed in the ratio of

uptake between the inorganic N forms suggested that, to some extent, plant physiology was

adjusted to match the soil conditions where species occurred. Yet, differences between

species with different strategies remains, with higher uptake for both forms associated with a

more exploitative strategy. We hypothesised that this should be also the case for organic N

sources (Kastovska and Santruckova 2011). Nevertheless, we acknowledge that this question

could be more important in harsh environments where organic N is relatively more abundant

as a N source for plants (Mozdzer et al 2014). Further investigations remain needed on the

variations of plant N uptake under field conditions, in link with potential variations in N

cycling in response to land-use or during the season (Robson et al. 2010, Legay et al. 2013).

Variations of N uptake ability during the growing season

Plant N uptake ability also varies during the growing season, with N uptake increasing

(Stahl et al. 2011) or decreasing (Jaeger et al. 1999) depending on the ecosystems

investigated. In the UU grassland, NH_4^+ uptake was higher for all species during the autumn

than during the summer, and the same was found for NO_3^- uptake by *B. erectus*. Plant activity

is usually considered to slow down during the autumn compared to the peak biomass in

summer, an assumption supported by higher LDMC and lower SLA for all species related to

the senescence of leaves. However, we did not observe any changes for root traits, suggesting

that roots could remain active during this time of the growing season, especially in the process

of resource storage, an important feature for subalpine/alpine plants (Kleijn et al. 2005).

Additionally, studies have reported an increase of grasslands N cycling rate in the autumn that

could be explained by more favourable soil conditions (first rains and mild temperature), and

associated with still active N uptake by plants as observed in our study (Miller et al. 2009,

Larsen et al. 2012). Despite the fact that only few soil parameters differed between the two

investigated seasons in the UU grassland, the N uptake increase in autumn was more likely a

site-dependent effect related to soil conditions (Miller et al. 2009, Stahl et al. 2011, Legay et

al. 2013) rather than a species response since they did show the same pattern in the other

grasslands. Yet, a multiple-year study remains necessary to better conclude on these seasonal

patterns.

Conclusions

By estimating inorganic root N uptake under controlled conditions from plants grown up

under field conditions, our results support the assumption that root and leaf functional traits

are associated with the ability of plants to acquire N. In particular, the observed pattern for

roots characteristics appeared similar to the one observed in the leaf economic spectrum, with

higher N uptake associated with more exploitative syndrome of traits. However, a weak

relationship between leaf and root traits suggests that leaf traits alone were insufficient to

predict N uptake. Additionally, N uptake varied within species during the growing season and

in response to local conditions, making root traits and soil parameters important features of

the relationships between plant functioning and grasslands N cycling.

Acknowledgements:

We wish to thank Marie-Paule Bataille for IRMS analyses. The authors thank the "Conseil

Régional de Basse-Normandie" for the funding of a postdoctoral position to FG. This study

was conducted as part of ERA-Net BiodivERsA project VITAL, ANR-08-BDVA-008.

References

Bardgett R.D., Bowman W.D., Kaufmann R., Schmidt S.K. (2005) A temporal approach to linking aboveground and belowground ecology. Trends in Ecology & Evolution **20**:634–641.

Bar-Tal A., Aloni B., Karni L., Rosenberg R. (2001) Nitrogen Nutrition of Greenhouse Pepper. II. Effects of Nitrogen Concentration and NO3: NH4 Ratio on Growth, Transpiration, and Nutrient Uptake. HortScience **36**:1252–1259.

Bassirirad H. (2000) Kinetics of nutrient uptake by roots: responses to global change. New Phytologist **147**:155–169.

Bowman W.D., Bahnj L., Damm M. (2003). Alpine landscape variation in foliar nitrogen and phosphorus concentrations and the relation to soil nitrogen and phosphorus availability. Arctic, Antarctic, and Alpine Research **35**: 144–149. https://doi.org/10.1657/1523-0430

Britto D.T., Kronzucker H.J. (2002) NH4+ toxicity in higher plants: a critical review. Journal of Plant Physiology **159**:567–584.

Britto D.T., Kronzucker H.J. (2013) Ecological significance and complexity of N-source preference in plants. Annals of Botany **112**:957–963.

Chapin F.S. (1980) The Mineral Nutrition of Wild Plants. Annual Review of Ecology and Systematics **11**:233–260.

Cornwell W.K., Cornelissen J.H.C., Amatangelo K., Dorrepaal E., Eviner V.T., Godoy O., Hobbie S.E., Hoorens B., Kurokawa H., Pérez-Harguindeguy N., Quested H.M., Santiago L.S., Wardle D.A., Wright I.J., Aerts R., Allison S.D., Van Bodegom P., Brovkin V., Chatain A., Callaghan T.V., Díaz S., Garnier E., Gurvich D.E., Kazakou E., Klein J.A., Read J., Reich P.B., Soudzilovskaia N.A., Vaieretti M.V., Westoby M. (2008) Plant species traits are the predominant control on litter decomposition rates within biomes worldwide. Ecology Letters **11**:1065–1071.

Craine J.M., Froehle J., Tilman D.G., Wedin D.A., Chapin I.I.I. (2001) The relationships among root and leaf traits of 76 grassland species and relative abundance along fertility and disturbance gradients. Oikos **93**:274–285.

Craine J.M., Lee W.G. (2003) Covariation in leaf and root traits for native and non-native grasses along an altitudinal gradient in New Zealand. Oecologia **134**:471–478.

Craine J.M., Lee W.G., Bond W.J., Williams R.J., Johnson L.C. (2005) Environmental constraints on a global relationship among leaf and root traits of grasses. Ecology **86**:12–19.

de Vries F.T., Manning P., Tallowin J.R.B., Mortimer S.R., Pilgrim E.S., Harrison K.A., Hobbs P.J., Quirk H., Shipley B., Cornelissen J.H.C., Kattge J., Bardgett R.D. (2012) Abiotic drivers and plant traits explain landscape-scale patterns in soil microbial communities. Ecology Letters **15**:1230–1239.

Dray S., Dufour A.-B. (2007) The ade4 Package: Implementing the Duality Diagram for Ecologists. Journal of Statistical Software **22** [online] URL: http://ideas.repec.org/a/jss/jstsof/22i04.html.

Errebhi M., Wilcox G.E. (1990) Tomato growth and nutrient uptake pattern as influenced by nitrogen form ratio. Journal of Plant Nutrition **13**:1031–1043.

Fort F., Jouany C., Cruz P. (2013) Root and leaf functional trait relations in Poaceae species: implications of differing resource-acquisition strategies. Journal of Plant Ecology **6**:211–219.

Freschet G.T., Cornelissen J.H.C., Van Logtestijn R.S.P., Aerts R. (2010) Evidence of the "plant economics spectrum" in a subarctic flora. Journal of Ecology **98**:362–373.

Garnett T.P., Smethurst P.J. (1999) Ammonium and nitrate uptake by Eucalyptus nitens: effects of pH and temperature. Plant and Soil **214**:133–140.

Gavito M.E., Curtis P.S., Mikkelsen T.N., Jakobsen I. (2001) Interactive effects of soil temperature, atmospheric carbon dioxide and soil N on root development, biomass and nutrient uptake of winter wheat during vegetative growth. Journal of Experimental Botany **52**:1913–1923.

Gessler A., Schneider S., Von Sengbusch D., Weber P., Hanemann U., Huber C., Rothe A., Kreutzer K., Rennenberg H. (1998) Field and laboratory experiments on net uptake of nitrate and ammonium by the roots of spruce (Picea abies) and beech (Fagus sylvatica) trees. New Phytologist **138**: 275–285. https://doi.org/10.1046/j.1469-8137.1998.00107.x

Grassein F., Lemauviel-Lavenant S., Lavorel S., Bahn M., Bardgett R.D., Desclos-Theveniau M., Laîné P. (2015) Relationship between functional traits and inorganic nitrogen acquisition among eight contrasting European grass species. Annals of Botany **115: 107-115.**

Grigulis K., Lavorel S., Krainer U., Legay N., Baxendale C., Dumont M., Kastl E., Arnoldi C., Bardgett R.D., Poly F., Pommier T., Schloter M., Tappeiner U., Bahn M., Clément J.-C. (2013) Relative contributions of plant traits and soil microbial properties to mountain grassland ecosystem services. Journal of Ecology **101**:47–57.

Hodge A. (2004) The plastic plant: root responses to heterogeneous supplies of nutrients. New Phytologist **162**: 9–24. https://doi.org/10.1111/j.1469-8137.2004.01015.x

Jaeger C.H., Monson R.K., Fisk M.C., Schmidt S.K. (1999) Seasonal partitioning of nitrogen by plants and soil microorganisms in an alpine ecosystem. Ecology **80**:1883–1891.

James J.J., Mangold J.M., Sheley R.L., Svejcar T. (2009) Root plasticity of native and invasive Great Basin species in response to soil nitrogen heterogeneity. Plant Ecology **202**: 211–220. https://doi.org/10.1007/s11258-008-9457-3

Kahmen A., Livesley S.J., Arndt S.K. (2009) High potential, but low actual, glycine uptake of dominant plant species in three Australian land-use types with intermediate N availability. Plant and Soil **325**:109–121.

Kastovska E., Santruckova H. (2011) Comparison of uptake of different N forms by soil microorganisms and two wet-grassland plants: A pot study. Soil Biology Biochemistry **43**: 1285–1291. https://doi.org/10.1016/j.soilbio.2011.02.021

Kleijn D., Treier U.A., Muller-Scharer H. (2005) The importance of nitrogen and carbohydrate storage for plant growth of the alpine herb Veratrum album. New Phytologist **166**: 565–575. https://doi.org/10.1111/j.1469-8137.2005.01321.x

Kronzucker H.J., Siddiqi M.Y., Glass A.D.M., Kirk G.J.D. (1999) Nitrate-Ammonium Synergism in Rice. A Subcellular Flux Analysis. Plant Physiology **119**:1041–1046.

Lainé P., Ourry A., Macduff J., Boucaud J., Salette J. (1993) Kinetic parameters of nitrate uptake by different catch crop species: effects of low temperatures or previous nitrate starvation. Physiologia Plantarum **88**:85–92.

Larsen K.S., Michelsen A., Jonasson S., Beier C., Grogan P. (2012) Nitrogen Uptake During Fall, Winter and Spring Differs Among Plant Functional Groups in a Subarctic Heath Ecosystem. Ecosystems **15**: 927–939. https://doi.org/10.1007/s10021-012-9555-x

Legay N., Grassein F., Robson T.M., Personeni E., Bataillé M.-P., Lavorel S., Clément J.-C. (2013) Comparison of inorganic nitrogen uptake dynamics following snowmelt and at peak biomass in subalpine grasslands. Biogeosciences **10**:7631-7645.

Léon M., Lainé P., Ourry A., Boucaud J. (1995) Increased uptake of native soil nitrogen by roots of Lolium multiflorum Lam. after nitrogen fertilization is explained by a stimulation of the uptake process itself. Plant and Soil **173**:197–203.

Liu G., Freschet G.T., Pan X., Cornelissen J.H.C., Li Y., Dong M. (2010) Coordinated variation in leaf and root traits across multiple spatial scales in Chinese semi-arid and arid ecosystems. New Phytologist **188**:543–553.

Louahlia S., Lainé P., Thornton B., Ourry A., Boucaud J. (2000) The role of N-remobilisation

and the uptake of NH4+ and NO3- by Lolium perenne L. in laminae growth following defoliation under field conditions. Plant and Soil **220**:175–187.

Lucash M.S., Eissenstat D.M., Joslin J.D., McFarlane K.J., Yanai R.D. (2007) Estimating nutrient uptake by mature tree roots under field conditions: challenges and opportunities. Trees **21**:593–603.

Maire V., Gross N., Da Silveira Pontes L., Picon-Cochard C., Soussana J. (2009) Trade-off between root nitrogen acquisition and shoot nitrogen utilization across 13 co-occurring pasture grass species. Functional Ecology **23**:668–679.

Miller A.E., Schimel J.P., Sickman J.O., Skeen K., Meixner T., Melack J.M. (2009) Seasonal variation in nitrogen uptake and turnover in two high-elevation soils: mineralization responses are site-dependent. Biogeochemistry **93**:253–270.

Min X.J., Siddiqi M.Y., Guy R.D., Glass A.D.M., Kronzucker H.J. (2000) A comparative kinetic analysis of nitrate and ammonium influx in two early-successional tree species of temperate and boreal forest ecosystems. Plant Cell Environment **23**: 321–328. https://doi.org/10.1046/j.1365-3040.2000.00546.x

Mozdzer T.J., McGlathery K.J., Mills A.L., Zieman J.C. (2014) Latitudinal variation in the availability and use of dissolved organic nitrogen in Atlantic coast salt marshes. Ecology **95**: 3293–3303. https://doi.org/10.1890/13-1823.1

Nasholm T., Huss-Danell K., Hogberg P. (2000). Uptake of organic nitrogen in the field by four agriculturally important plant species. Ecology **81**: 1155–1161. https://doi.org/10.2307/177188

Nasholm T., Kielland K., Ganeteg U. (2009) Uptake of organic nitrogen by plants. New Phytologist **182**: 31–48. https://doi.org/10.1111/j.1469-8137.2008.02751.x

Ordonez J.C., van Bodegom P.M., Witte J.-P.M., Wright I.J., Reich P.B., Aerts R. (2009) A global study of relationships between leaf traits, climate and soil measures of nutrient fertility. Global Ecology and Biogeography **18**: 137–149. https://doi.org/10.1111/j.1466-8238.2008.00441.x

Osone Y., Ishida A., Tateno M. (2008) Correlation between relative growth rate and specific leaf area requires associations of specific leaf area with nitrogen absorption rate of roots. New Phytologist **179**: 417–427. https://doi.org/10.1111/j.1469-8137.2008.02476.x

Pérez-Harguindeguy N., Díaz S., Garnier E., Lavorel S., Poorter H., Jaureguiberry P., Bret-Harte M. S., Cornwell W. K., Craine J. M., Gurvich D. E., Urcelay C., Veneklaas E. J., Reich P. B., Poorter L., Wright I. J., Ray P., Enrico L., Pausas J. G., de Vos A. C., Buchmann N., Funes G., Quétier F., Hodgson J. G., Thompson K., Morgan H. D., ter Steege H., van der Heijden M. G. A., Sack L., Blonder B., Poschlod P., Vaieretti M. V., Conti G., Staver A. C., Aquino S., Cornelissen J. H. C. (2013) New handbook for standardised measurement of plant functional traits worldwide. Australian Journal of Botany **61**: 167-234. doi: 10.1071/BT12225

Quétier F., Thébault A., Lavorel S. (2007) Plant traits in a state and transition framework as markers of ecosystem response to land-use change. Ecological Monographs **77**:33–52.

Reich P.B. (2014) The world-wide "fast–slow" plant economics spectrum: a traits manifesto. Journal of Ecology **102**:275–301.

Rewald B., Rechenmacher A., Godbold D.L. (2014) It's Complicated: Intraroot System Variability of Respiration and Morphological Traits in Four Deciduous Tree Species. Plant Physiology **166**: 736–745. https://doi.org/10.1104/pp.114.240267

Robson T.M., Baptist F., Clément J., Lavorel S. (2010) Land use in subalpine grasslands affects nitrogen cycling via changes in plant community and soil microbial uptake dynamics. Journal of Ecology **98**:62–73.

Robson T.M., Lavorel S., Clement J.-C., Roux X.L. (2007) Neglect of mowing and manuring leads to slower nitrogen cycling in subalpine grasslands. Soil Biology and

Biochemistry **39**:930–941.
Roumet C., Birouste M., Picon-Cochard C., Ghestem M., Osman N., Vrignon-Brenas S., Cao
K., Stokes A. (2016) Root structure-function relationships in 74 species: evidence of a
root economics spectrum related to carbon economy. New Phytologist **210**: 815–826.
https://doi.org/10.1111/nph.13828
Roumet C., Urcelay C., Díaz S. (2006) Suites of root traits differ between annual and
perennial species growing in the field. New Phytologist **170**:357–368.
Stahl V.M., Beyschlag W., Werner C. (2011) Dynamic niche sharing in dry acidic grasslands
-a 15N-labeling experiment. Plant and Soil **344**:389–400.
Stoelken G., Simon J., Ehlting B., Rennenberg H. (2010) The presence of amino acids affects
inorganic N uptake in non-mycorrhizal seedlings of European beech (Fagus sylvatica).
Tree Physiology **30**: 1118–1128. https://doi.org/10.1093/treephys/tpq050
Tjoelker M.G., Craine J.M., Wedin D., Reich P.B., Tilman D. (2005) Linking leaf and root
trait syndromes among 39 grassland and savannah species. New Phytologist **167**:493–
508.
Warncke D.D., Barber S.A. (1973) Ammmonium and Nitrate Uptake by Corn (Zea mays L.)
as Influenced by Nitrogen Concentration and NH+4 /NO-3 Ratio1. Agronomy Journal
65:950.
Wright I.J., Reich P.B., Westoby M., Ackerly D.D., Baruch Z., Bongers F., Cavender-Bares
J., Chapin T., Cornelissen J.H.C., Diemer M., Flexas J., Garnier E., Groom P.K.,
Gulias J., Hikosaka K., Lamont B.B., Lee T., Lee W., Lusk C., Midgley J.J., Navas
M.-L., Niinemets U., Oleksyn J., Osada N., Poorter H., Poot P., Prior L., Pyankov
V.I., Roumet C., Thomas S.C., Tjoelker M.G., Veneklaas E.J., Villar R. (2004) The
worldwide leaf economics spectrum. Nature **428**:821–827.
Zeller V., Bahn M., Aichner M., Tappeiner U. (2000) Impact of land-use change on nitrogen
mineralization in subalpine grasslands in the Southern Alps. Biology and Fertility of
Soils **31**:441–448.

Table 1: Description of the studied grasslands. Past and current land uses describe the former

and current management of these grasslands (see Quetier et al. 2006 for more information).

TMF: Terraced Mown and Fertilized, TU: Terraced Unmown not fertilized but lightly grazed,

TM: Unterraced Mown, UU: Unterraced Unmown but lightly grazed.

Field label	GPS coordinates	Sampled species	Past Land Use	Current Land Use	Elevation
TMF	45° 2'42.77"N 6°20'35.29"E	*Dactylis glomerata*	Arable rotation	fertilized hay meadow	1800m
TU	45° 2'24.74"N 6°21'38.15"E	*Bromus erectus*	Arable rotation	grazed pasture (sheep)	1840m
UM	45° 1'58.59"N 6°23'0.80"E	*Festuca paniculata*	Hay meadow	unfertilized hay meadow	1980m
UU	45° 1'55.76"N 6°23'1.57"E	*Festuca paniculata* *Bromus erectus* *Dactylis glomerata*	Hay meadow	grazed pasture (sheep)	1960m

Table 2: Mean values ± standard errors of leaf and root traits for each species, site and sampling time (n=5). For a given trait and species, statistically similar values have the same letter (Tukey post-hoc test). Bold values indicate the season with the highest trait values for a given species in a given grassland.

Species	Site	Season	SLA (mm².g⁻¹)	LDMC (mg.g⁻¹)	LNC (mg.g⁻¹)	SRL (m.g⁻¹)	RDMC (mg.g⁻¹)	RNC (mg.g⁻¹)
B. erectus	TU	Summer	**30.07 ± 0.8** ^{a}	277.6 ± 2.1 ^{c}	21.46 ± 0.2	249.86 ± 72.0	307.77 ± 22.4	6.06 ± 0.1
		Autumn	15.43 ± 1.1 ^{c}	**444.06 ± 16.8** ^{a}	12.09 ± 0.9	253.58 ± 34.3	303.5 ± 6.3	7.68 ± 0.4
	UU	Summer	**22.58 ± 0.7** ^{b}	305.06 ± 4.2 ^{c}	15.57 ± 0.8	328 ± 74.2	290.82 ± 21.7	7.17 ± 0.1
		Autumn	16.37 ± 1 ^{c}	**385.1 ± 11.5** ^{b}	19.28 ± 2.7	222.12 ± 75.4	288.32 ± 21.6	7.87 ± 0.5
D. glomerata	TMF	Summer	**34.06 ± 1.6** ^{a}	248.79 ± 17.7 ^{c}	**32.6 ± 3.4** ^{a}	426.22 ± 60.5	232.49 ± 20.5 ^{b}	7.28 ± 0.4
		Autumn	20.15 ± 2 ^{b}	**298.51 ± 2.5** ^{b}	19.41 ± 2.2 ^{b}	290 ± 48.6	249.09 ± 8 ^{b}	7.99 ± 0.3
	UU	Summer	26.17 ± 0.3 ^{ab}	265.78 ± 9.1 ^{bc}	**24.2 ± 1.4** ^{ab}	318.69 ± 63.8	254.08 ± 5.6 ^{b}	6.91 ± 0.5
		Autumn	24.34 ± 1.7 ^{b}	**389.69 ± 10.5** ^{a}	21.81 ± 2.5 ^{ab}	216.87 ± 44.7	**304.99 ± 8.8** ^{a}	7.16 ± 0.2
F. paniculata	UM	Summer	**23.26 ± 0.4** ^{a}	232.06 ± 0.6 ^{c}	**20.22 ± 1.9** ^{a}	195.85 ± 21.3 ^{ab}	300.1 ± 21.5	5.15 ± 0.4
		Autumn	8.78 ± 0.5 ^{b}	**433.87 ± 18.8** ^{a}	10.61 ± 1.4 ^{b}	153.37 ± 20.5 ^{b}	369.12 ± 21.7	4.32 ± 0.5
	UU	Summer	**21.36 ± 0.9** ^{a}	261.06 ± 15.6 ^{c}	14.52 ± 1.8 ^{ab}	166.21 ± 20.7 ^{ab}	380.91 ± 6.4	4.16 ± 0.1
		Autumn	11.76 ± 1.4 ^{b}	**369.71 ± 7.2** ^{b}	15.53 ± 3.6 ^{ab}	293.71 ± 54.5 ^{a}	366.84 ± 23.3	4.31 ± 0.1

Table 3. Pearson correlations between NH_4^+ and NO_3^- maximum uptake rates (Vmax), PCA axes, leaf and root traits. Significant values (p-value <0.05) are indicated in bold.

	$VmaxNH_4^+$	$VmaxNO_3^-$	axe1	axe2	ratio	SLA	LDMC	SRL	RDMC	LNC
$VmaxNO_3^-$	**0.93**									
axe1	**0.65**	**0.76**								
axe2	-0.28	-0.24	0.00							
ratio	-0.46	**-0.68**	**-0.64**	0.43						
SLA	0.37	0.53	**0.84**	0.38	-0.54					
LDMC	-0.17	-0.26	**-0.60**	**-0.74**	0.09	**-0.74**				
SRL	0.55	**0.70**	**0.81**	-0.14	-0.51	0.54	-0.34			
RDMC	**-0.68**	**-0.77**	**-0.91**	0.31	**0.75**	**-0.63**	0.37	**-0.71**		
LNC	**0.64**	**0.65**	**0.87**	0.10	-0.32	**0.69**	-0.51	**0.66**	**-0.74**	
RNC	**0.58**	**0.62**	**0.65**	**-0.70**	**-0.80**	0.34	0.08	0.50	**-0.83**	0.42

Table 4. Soil properties (mean values ± SE) for each grassland and at each sampling time. No significant difference values between sites at a given date are shown by the same letter (Tukey post-hoc test). Values in bold indicate the highest values when the considered soil parameter was significantly different between dates in a grassland. na: not available because of a sampling issue. nd: not detectable: under the level of sensitivity of the method; TDN: total dissolved nitrogen).

		TMF	TU	UM	UU
Soil Water	Summer	**22.33 ± 0.71** [b]	**18.33 ± 1.12** [c]	**na**	**34.89 ± 1.20** [a]
Content (%)	Autumn	11.44 ± 0.33 [b]	7.25 ± 0.88 [c]	13.58 ± 1.18 [b]	17.74 ± 1.54 [a]
pH	Summer	8.01 ± 0.04 [a]	8.03 ± 0.04 [a]	n.a.	**6.31 ± 0.05** [b]
	Autumn	7.98 ± 0.02 [a]	8.05 ± 0.04 [a]	5.85 ± 0.05 [b]	6.02 ± 0.08 [b]
Soil Organic	Summer	13.16 ± 0.48 [c]	14.39 ± 0.93 [b]	n.a.	18.42 ± 0.64 [a]
Matter (%)	Autumn	12.47 ± 0.65 [b]	11.38 ± 0.93 [b]	14.02 ± 0.43 [ab]	16.98 ± 1.29 [a]
C:N ratio	Summer	**14.91 ± 0.35** [a]	14.02 ± 0.64 [a]	n.a.	12.10 ± 0.15 [b]
	Autumn	13.46 ± 0.39 [a]	13.43 ± 0.64 [ab]	11.63 ± 0.14 [b]	11.74 ± 0.43 [b]
TDN ($\mu gN.g^{-1}$	Summer	20.83 ± 2.95 [b]	16.85 ± 1.33 [b]	n.a.	55.59 ± 12.15 [a]
soil)	Autumn	**46.12 ± 4.44** [a]	**46.13 ± 4.74** [a]	34.54 ± 1.21 [ab]	46.65 ± 6.04 [a]
NO_3^- content	Summer	3.87 ± 0.48 [ab]	**5.63 ± 0.64** [a]	n.a.	2.31 ± 0.53 [b]
($\mu gN.g^{-1}$ soil)	Autumn	2.99 ± 0.54 [a]	0.54 ± 0.10 [b]	n.d.	1.39 ± 0.66 [ab]
NH_4^+ content	Summer	11.19 ± 0.47 [a]	15.43 ± 1.82 [a]	n.a.	12.23 ± 1.77 [a]
($\mu gN.g^{-1}$ soil)	Autumn	10.07 ± 1.37 [a]	11.54 ± 1.55 [a]	6.97 ± 0.94 [a]	10.17 ± 1.92 [a]
NH_4^+:NO_3^- ratio	Summer	3.56 ± 0.65 [b]	2.97 ± 0.30 [b]	n.a.	7.85 ± 1.62 [a]
	Autumn	3.53 ± 0.36 [b]	**28.68 ± 10.7** [a]	n.d.	12.9 ± 6.4 [ab]

Figures legends:

Figure 1. Principal components analysis (PCA) of functional traits measured for the leaves and roots of three grass species (Be: *Bromus erectus*, Dg: *Dactylis glomerata* and Fp: *Festuca paniculata*), in each grassland with different management (UU: unterraced unmown, UM: unterraced mown, TMF: terraced mown and fertilized, TU: terraced unmown). SLA: Specific leaf area, LDMC: Leaf dry matter content, LNC: Leaf nitrogen content, SRL: Specific root length, RDMC: Root dry matter content, RNC: Root nitrogen content.

Figure 2: Relationships between the first axes of the PCA (fig 1) and Vmax for NH_4^+ (a), NO_3^- (b) and $NH_4^+:NO_3^-$ uptake ratio (c). The three relationships were significant (p-values<0.05) assuming a polynomial relationship of order=2, and the resulting R^2 are indicated on each graph.

Figure 3. Vmax (Maximal uptake rate) for NH_4^+ (a) and NO_3^- (b) of *D. glomerata*, *B. erectus* and *F. paniculata*. Within each combination of site and species, dates with the same letter had similar uptake parameters (Tukey post hoc test at 5% level, after an Anova with date as main effect). For each species, the significance of the differences between the two sites for uptake parameters were tested using a Student test, and stars indicate the dates at which the two sites differ significantly with a p-value<0.05.

Figure 4: NH_4^+:NO_3^- uptake ratio for the three species, in each site and at the two sampling times. The uptake ratio is unitless (ratio between NH_4^+Vmax and NO_3^-Vmax). Within each species, *or *** indicate significant site effects (p value < 0.05 and 0.001 respectively) within each species (two-ways ANOVA with site, date and the interaction as main effect). In the grassland (UU) where all species occurred, the differences between species and sampling time were tested using two-ways ANOVA. Similar letters connect species with similar values in the UU grassland at both sampling dates.