Spatial frequency tuning of perceptual learning and transfer in global motion

Jordi M Asher [1*], Vincenzo Romei[1,2], Paul B Hibbard[1],

1 University of Essex, Department of Psychology, Wivenhoe Park, United Kingdom, CO4 3SQ
2 Dipartimento di Psicologia and Centro studi e Ricerche in Neuroscienze Cognitive, Campus di Cesena, Università di Bologna 47521 Cesena, Italy

* jashera@essex.ac.uk

Abstract

Perceptual learning is typically highly specific to the stimuli and task used during training. However, recently it has been shown that training on global motion can transfer to untrained tasks, reflecting the generalising properties of mechanisms at this level of processing. We investigated if a) feedback was required for learning when using an equivalent noise global motion coherence task, and b) the transfer across spatial frequency of training on a global motion coherence task, and the transfer of this training to a measure of contrast sensitivity. For our first experiment two groups, with and without feedback, trained for ten days on a broadband global motion coherence task. Results indicated that feedback was a requirement for learning. For the second experiment training consisted of five days of direction discrimination on one of three global motion tasks (broadband, low or high frequency random-dot Gabors), with trial-by-trial auditory feedback. A pre- and post-training assessment was also conducted, consisting of all three types of global motion stimuli (without feedback) and high and low spatial frequency contrast sensitivity. We predicted that if learning involves low level processing, then based on the frequency specificity of the lower level areas, transfer would be contingent on the frequency of training. However, if learning involves a higher, global level of processing, more selective transfer would occur that matches the broadband tuning of the higher processing levels. Our training paradigm was successful at eliciting improvement in the trained tasks over the five days. However, post-training assessments found learning and transfer exclusively for the group trained on low spatial frequency global motion. This group exhibited increased sensitivity to low spatial frequency contrast, and an improvement for the broadband global motion condition. Our findings are consistent with perceptual learning which depends on the global stage of motion processing in V5.

Introduction

Perceptual learning has attracted much attention as a potential tool to aid recovery of lost visual function for clinical populations [1]. The success of perceptual training in amblyopia [2–4], presbyopia [4] and cortical damage [5–7] has demonstrated sensory plasticity in adulthood. This evidence contradicts the position that sensory development is restricted to a critical period early in life [8,9] and that the visual system is hard-wired in mature systems [10]. While it has repeatedly been established that

training can improve perceptual abilities [11], these benefits tend to be highly specific for both the perceptual features of the stimuli [12–14] and the behavioural task used in training [15]. This specificity severely limits the effectiveness of perceptual learning as a general therapeutic tool. Resolving the conditions under which learning is tied to the features and tasks used in training, and how much it can generalise to new tasks and stimuli, is imperative for understanding the mechanisms of perceptual learning [11,16].

Models of Perceptual Learning

An important theoretical question in perceptual learning concerns the location in the processing stream of the learning mechanism(s), for which there are, broadly speaking, two main positions. The first explains perceptual learning in terms of a change in the neurons that code for that feature [17,18]. Perceptual learning is typically highly specific for the features of the stimuli used for training, including their orientation [13,19,20], spatial frequency [11,13,21], direction of motion [15], retinal location [14,20,22–25] and the eye to which they are presented [14,23,24]. On the basis of this feature-specificity it has been argued that the underlying brain area responsible for the learning process must be the primary visual cortex [14] where the receptive fields of cells display this high degree of specificity. Gilbert et al., (2001) [18] suggest that this indicates that there is an element of plasticity early in cortical visual processing and consider that individual neurons at this level change over time [26]. One way perceptual learning could occur is by altering the receptive fields of sensory neurons to exploit correlations between their responses, via anti-Hebbian or Hebbian learning [18]. Anti-Hebbian learning has the aim of reducing the correlations between the responses of neurons, so as to produce an efficient encoding of sensory information [27–29]. Conversely, Hebbian learning, by accentuating the correlations between responses, might be beneficial in tasks such as global motion and form, in which it is important to segregate signal from background noise [18].

However, Mollon and Danilova (1996) [30] argue that the failure of perceptual learning to transfer to test stimuli that differ from those used in training cannot be used to infer unambiguously that learning occurs through a change in the representation of information in early visual areas. The specificity of learning, and lack of transfer, merely indicate that these early visual areas are involved in the learning process. Herzog and Fahle (1998) [31] suggest that learning is more complex than proposed by the visual representation model. They argue that learning is not driven purely by a set of stimuli in a bottom up manner, and that a model of learning also needs to include top down factors such as an internal estimation of one's performance and knowledge and understanding of the task. While later 'low-level' models [26] do consider top-down influences on learning, such as attention and feedback, they still argue that the change occurs at the level of the receptive field.

The second prominent position hypothesises that learning takes place at higher levels within the processing hierarchy [24,32–35]. Consistent with the lower-level models, Dosher and Lu (2010) [33] agree that Hebbian learning is one of the most important forms of perceptual learning. However, unlike models where the change occurs at the level of the receptive fields in V1, higher-level models propose that the repeated experience at V1 alters the connections between lower visual areas and the high level areas where the perceptual decision is then made [36]. Therefore, rather than a change in the sensory neuron, learning represents a change in the weights of connections between the input, sensory encoding layer and the classification layer at which the decision is made.

Many behavioural studies of perceptual learning are unable to distinguish between improvements that result from a change in sensory encoding, and those that result from a change in the connections between sensory and decision-making stages of processing.

While the failure of perceptual learning to transfer across retinal location, orientation and other stimulus dimensions implicates V1 in the learning process, this in itself cannot dissociate a change in the receptive fields of neurons from a change in their connections to subsequent processing stages [37]. Petrov et al., [36, 37] suggest using perceptual learning experiments in which the sensory representations required for two tasks are the same, but the decision stages differ, or where both sensory representations and the decision stage are common to both. Bejjanki et al., (2011) [38] propose a model that is computationally similar to the late theory models, however it illustrates how changing the population codes in the early sensory areas can create similar changes in response to those made by high-level re-weighting models. Bejjanki et al., (2011) [38] argue that the model captures the characteristics for perceptual learning in both behavioural and physiological changes, through sensory inputs improving the decision weights in the feed-forward connections, and improved probabilistic inference in early visual areas resulting from increased neural activity in the feedback network.

The effects of learning on the receptive field structure of visual neurons have been measured directly using single-cell recordings. Changes in orientation and spatial frequency tuning in V1 [39, 40] and V4 [41], and increased sensitivity to motion coherence in V5 [42], have all been found as a result of perceptual learning. Neuroimaging in humans observers has indicated a reorganisation of the visual cortex after training [43–45]. However it has been argued that the changes to the tuning of individual sensory neurons are not sufficient to account for the behavioural improvements achieved through training, and that the effects on populations of neurons, including the reductions in the correlation in noise between neural responses, need to be taken into account [46]. Dosher and Lu (1998) [32] suggested that learning may be explained as an improved ability to filter out external noise and reduce internal noise. Likewise, Gu et al., (2011) [46] predicted that by pooling sensory information across cells, noise would be reduced, efficiency improved and learning would occur. Noise levels were examined in macaques, by recording from individual pairs of neurons. Gu et al., (2011) [46] found that noise correlations were weaker for those trained on a perceptual discrimination task than those who were untrained. However, their results also showed that optimally decoding all neurons did not predict a meaningful change in learning performance. Gu et al., (2011) [46] concluded that the reduction in noise is also an unlikely explanation for the effects of training, and support the idea that perceptual learning may change the weighting of sensory inputs to a decision unit. It is generally judged that the changes to responses in sensory neurons are not sufficient to explain the improvements in task performance that are observed [33, 46].

Transfer

A priority for perceptual learning researchers is identifying learning that transfers. A number of recent studies have demonstrated transfer after training [47–49]. Xiao et al., (2008) [48] demonstrated a transfer of contrast discrimination to different retinal locations using a double-training protocol. Double-training involves overt training on a task-relevant feature in addition to exposure to a task-irrelevant feature [50]. Xiao et al., (2008) [48] trained observers using two simultaneous tasks; one task was to detect a relevant feature (contrast) at one retinal location, followed by detection of an irrelevant feature (orientation) at another location. They reported full transfer of contrast sensitivity to the untrained location. Similarly Zhang et al. (2010) [49] showed that observers who were trained to discriminate orientation and simultaneously exposed to a second orientation experienced a full transfer of improvement in orientation sensitivity as long as training preceded exposure, rather than being simultaneously presented. Exposure to the untrained orientation was not sufficient to lead to transfer. Zhang et al., (2010) [49] proposed a rule-based learning model to account for the transfer, where

higher level decision units learn and re-weight the V1 inputs. They proposed that the [110] absence of functional connections to the untrained orientation or location prevents any [111] potential reweighting. Double training, through exposure, activates the functional [112] connections at the new locations or orientations, enabling transfer. This suggests that [113] there may not be a straightforward correspondence between stimulus feature and the [114] neural loci involved in performing the task. Double-training has been a successful [115] paradigm in obtaining evidence of transfer [48,51,52] however, task difficulty during [116] training and confidence in decisions still play a vital role in the learning and transfer [117] process [53]. [118]

The Reverse Hierarchy Theory is an early theoretical model proposed to account for [119] the transfer of learning [54]. Hochstein and Ahissar (2002) [54] proposed that learning is [120] not a simple bottom up, stimulus driven process. Rather, learning is a top down [121] process, beginning at higher processing areas and filtering backwards to lower-level [122] areas, with the degree of transfer dependent on the characteristics of the receptive fields [123] involved in performing the training task. Hochstein and Ahissar (2002) [54] proposed [124] that transfer occurs as a result of modification of neurons found in the higher cortical [125] levels, where receptive fields generalise. Alternatively, specificity could arise from [126] changes in neurons where the receptive fields are localised, at the lower cortical levels. [127] Global processing may therefore play an important role in creating learning that [128] generalises to other tasks and stimuli [55]. If, as predicted by feature specificity, [129] learning is stimulus driven and reflects the tuning properties of the relevant receptive [130] fields, what specificity is expected for tasks that are not processed in early, highly [131] specific lower visual areas? For example, perceptual learning has been shown to improve [132] detection and discrimination in tasks using global form [56] and global motion [57–59]. [133] These kinds of tasks are known to be processed at higher levels of the visual cortex, in [134] areas V4 [56] and V5 [60,61], respectively. [135]

The visual system is typically represented as a hierarchy of cortical areas [62,63]. At [136] the lowest level, V1 and V2 represent simple visual dimensions, such as the position, [137] orientation and scale of local image features [12]. The receptive fields of cells in V1 are [138] small, only responding to a very restricted area of the visual field [64–66]. Furthermore, [139] physiological evidence from non-human animals has shown that V1 neurons are sharply [140] tuned to orientation [67,68], spatial frequency [67,69] and direction [63,70]. In contrast, [141] the receptive fields in higher cortical areas, such as those found in V4 and V5, are larger [142] than those of V1 and are less dependent on location, retinal size, viewpoint, and [143] lighting [64,69,71–75]. [144]

The Reverse Hierarchy Theory predicts that the feedback connections from higher [145] levels back to early visual processing areas may be involved in facilitating learning that [146] transfers, and that the key to understanding specificity and transfer lies in the hierarchy [147] of processing and the feedforward and feedback connections between them [54,55]. [148]

Global versus Local Processing [149]

The receptive fields of neurons in higher cortical areas integrate information to represent [150] global stimulus properties [18,76–79]. Global processes are investigated using stimuli or [151] tasks that can only be resolved through integration and segregation of coherent or [152] conflicting information [78–80]. For example, the perception of motion is hypothesised [153] to occur as a two stage process [77,81]. At the first stage, spatial frequency- and [154] orientation-tuned mechanisms in V1 encode the motion signals that occur locally within [155] the receptive fields of individual neurons [70,78]. However, to process more complex [156] motion, the ambiguous or conflicting first-stage signals need to be integrated over a [157] wider spatial area to provide a global representation of motion, and to unambiguously [158] estimate local velocity [82,83]. This process is hypothesised to occur higher in the visual [159] hierarchy in area V5 [77,81]. Single cell recordings from the visual cortex of non-human [160]

primates have identified the neural populations responsible for the two processing stages to be V1 and MT/V5, respectively [70]. The role that V5 plays in spatially integrating motion signals is well supported from non-human primate data, as well as neuroimaging studies in humans [60, 64, 84, 85]. Interestingly, the re-entrant connections from V5 to V1 are particularly extensive, suggesting an important role for top-down information in the encoding and interpretation of motion [86].

Global motion coherence is typically studied using random dot kinematograms, requiring observers to make a direction judgement from a stimulus comprised of a pattern of moving dots. A typical stimulus will often contain a proportion of signal dots moving in one direction and noise dots moving in random directions [15, 59, 87] (fig 1a). Difficulty is increased by reducing the ratio of signal-to-noise dots; the more noise dots the lower the coherence, and the more difficult task. In order to perceive a coherent global motion, observers need to integrate and segregate the motion signals over space and time [78]. Another method of investigating motion coherence is to use an equivalent noise paradigm [58, 88–90]. Rather than having distinct populations of signal and noise dots, all dots contribute both to the signal and the noise by drawing the direction of motion for each dot from a random distribution. Dots move the same distance between each frame and the direction travelled is independent of the directions of the other dots [90]. Difficulty is increased by manipulating the standard deviation of the distribution of directions presented, thus each dot contributes to the signal [89] (fig 1b).

Fig 1. Schematic of global motion stimuli where a) is a typical stimulus comprised of signal dots and noise dots (approximately 30% coherence). b) An equivalent noise stimulus where all dots contribute equally to the signal and the noise (drawn from a distribution of approximately 90° out of a possible 360°). In each case the correct answer would be that the dots are moving towards the right.

The local and global stages of motion processing differ in their spatial frequency tuning. Local motion detectors have receptive fields that are narrowly tuned for spatial frequency [67]. In contrast, receptive fields that process global motion have a much broader tuning. Cells higher in the processing cortical hierarchy, involved in the perception of global aspects of the image, thus generalise over stimulus parameters such as spatial frequency and location. Modification of these generalising receptive fields could therefore produce perceptual learning that also generalises over these stimulus parameters. On the other hand, specificity in perceptual learning arises from changes in lower-level cortical areas, where neurons show much more specific tuning for stimulus features. Since perceptual learning occurs for both global motion [58, 59] and form tasks [80, 91], this suggests that learning is not restricted to the initial encoding of information in V1, and can occur at higher levels of cortical processing. Learning that involves higher level global aspects of perception is of particular interest since it has the potential to produce more generalisable improvements in perception [55]. Establishing the conditions for and the extent to which global motion training generalises its learning in this way is an important step to understanding the global level of processing in

Perception of global motion   198

The role that V5 plays in the perception of global processes has been highlighted by   199
extra-cellular recordings [60, 92] and lesion studies [93, 94] in non-human primates.   200
Receptive fields in V5 can be up to tenfold larger than those in V1. V5 neurons have   201
large, broadly tuned receptive fields that sum the responses of a set of V1 neurons,   202
across space, orientation and spatial frequency [66]. More recently, Lui et al., (2007) [84] 203
showed that the majority of V5 neurons in marmoset monkeys have band-pass spatial   204
and temporal frequency tuning, with a preference for low spatial frequencies. Incoming 205
sensory information causing V5 to respond drives feedback projection from V5 to V1   206
that corresponds to the retinotopic locations around the stimulus location of the V1   207
cells [74]. Using a noise masking paradigm Amano et al., (2009) [76] investigated if the 208
specificity of local motion frequency tuning was preserved once pooling for the   209
perception of global motion has occurred. Stimuli were composed of either drifting   210
Gabors or plaid elements that were either signal or noise. Signal elements were defined 211
by a consistent drift speed and direction, and noise elements were defined by random   212
speeds and directions. Frequency tuning was determined by examining how noise   213
elements at one spatial frequency interfered with the extraction of the signal at another 214
frequency. Amano et al., (2009) [76] found that thresholds increased as the spatial   215
frequency of the noise elements was reduced, and concluded that this is indicative of a 216
broadband, low-pass tuning of motion pooling. The results obtained by Amano et al.,   217
(2009) [76] are consistent with those of Bex and Dakin (2002) [77], and suggest a global 218
level pooling for low spatial frequencies. Moreover, this is further supported by an fMRI 219
study [95] which found that while areas V1,V2,V3 and V4 showed bandpass spatial   220
frequency tuning, area V5+ exhibited an attenuated response for high spatial   221
frequencies, but no reduction in responses for low spatial frequencies. Amano et al.,   222
(2009) [76] propose that, within V5, there is a "motion-pooling mechanism" that   223
demonstrates a broadband, low-pass tuning. Pooling of visual sensory information is an 224
important step for integration and segregation and has also been found in studies   225
investigating transfer in binocular stereopsis [96]. The neural mechanisms of   226
stereoscopic vision are also known to be processed on multiple levels of the visual   227
hierarchy [97], and the pooling of information across spatial frequency mechanisms has 228
been proposed as an important step in estimating depth [98–101]. In the same way,   229
pooling in V5 for low frequencies is potentially an important step for integration and 230
segregation of coherent motion.   231

Learning mechanisms for global tasks   232

Training using global motion has been found to transfer between eyes [15, 59]; increase 233
sensitivity for detection and discrimination [5]; help recover some of the blind field for 234
cortically blind subjects [5]; and reduce contrast thresholds for drifting stimuli [58]. 235
Huxlin et al., (2009) [5] suggested that extensive training using global motion may result 236
in general improvements in visual sensitivity, and proposed that the improvements may   237
be a result of the feedback connections from V5 to V1. Huxlin et al., (2009) [5] explain 238
their findings using the Reverse Hierarchy Model of perceptual learning [54] and   239
propose that training may stimulate and reactivate "intact islands" of activity.   240
    Levi et al., (2014) [58] also found transfer of learning in a control population.   241
Transfer occurred from global motion to contrast sensitivity, a task known to be   242
processed in an early visual area where cells are narrowly tuned [67]. Levi et al.,   243
(2014) [58] suggested that improvements in their study may be due to the combination   244
of learning a global motion task, at a higher cortical level, while simultaneously being 245

exposed to high contrast, broad-frequency random dot stimuli. Levi et al., (2014) [58]   246
remarked that a striking result from their findings was the particular improvement in   247
direction discrimination for low spatial frequency stimuli, despite training with   248
broadband random dot stimuli. They consider these findings evidence for a broad   249
transfer of learning, across spatial frequency, in the primary visual areas, and that the   250
learning achieved through training on global tasks differs fundamentally from that from   251
local tasks.   252

Greater transfer of learning is proposed to occur through the re-weighting of   253
high-level perceptual units, which tend to generalise across stimulus properties to a   254
greater degree than low-level perceptual neurons [34, 58]. However, physiological   255
evidence of training on global motion tasks suggests that, as in training for the   256
discrimination of local image properties, learning changes the weighting between the   257
sensory neurons (in this case direction selective neurons in V5) and neurons involved in   258
the decision making process, rather than altering the sensory neuron's properties [102].   259
Given the differences in spatial and temporal frequency tuning of processing between the   260
local and global levels [60, 67, 84], it is important to determine the degree of specificity   261
for low-level stimulus properties when training with global stimuli. This allows us to   262
determine whether the degree of transfer is limited by the tuning properties of global   263
detectors, in the same way as for information encoded purely locally, and thus whether   264
the involvement of sensory mechanisms is the same for the two levels of processing.   265

Current study: Learning transfer from global motion   266

Levi et al., (2014) [58] found that after training on global motion tasks, with a   267
broadband frequency range, observers showed an increase in contrast sensitivity.   268
However, broad frequency stimuli contain both high and low frequencies, and contrast   269
detection is known to be associated with processing in early visual areas [21, 103, 104].   270
Therefore, if part of the learning mechanism for global motion operates at the early,   271
local processing stage, this may explain the transfer of learning they found, since the   272
broadband stimuli used would activate neurons with a range of spatial frequency tuning.   273
This would support the model of learning which asserts that learning changes involve   274
early, local receptive cells [17, 18, 26].   275

However, if the learning mechanism for global motion is located at higher levels,   276
where motion detectors have a preference for low spatial frequencies [76, 84], it might be   277
predicted that perceptual learning for global motion would show a broad transfer to   278
stimuli that contain low spatial frequency components. This would also be consistent   279
with the transfer of learning from global motion to contrast sensitivity for drifting   280
gratings found by Levi et al., (2014) [58]. The spatial frequency tuning of learning for   281
global motion may thus be used to determine whether this learning involves early, local   282
motion detectors, or is restricted to the global processing stage. In order to test this, we   283
focused on the spatial frequency tuning of learning, using a global motion coherence   284
task. Since the stimuli used by Levi et al., (2014) [58] were broadband, it was not   285
possible to determine the degree to which learning depends on the spatial frequency   286
content of their training stimuli. To address this, we used a similar learning paradigm,   287
but used test and training stimuli that contained (1) a broad range of spatial   288
frequencies; (2) only low spatial frequencies or (3) only high spatial frequencies.   289
Additionally, we included measures for contrast sensitivity for static low and high   290
spatial frequency targets. Furthermore, based on the predictions that learning and   291
transfer are most successful with a combination of easy and difficult   292
trials [47, 53, 105, 106] the stimulus levels chosen ensured a range from extremely easy   293
(high coherence) to very difficult (low coherence), with a high proportion of easy trials.   294
If training at the global level impacts learning at the early, local motion processing   295
stage, we may expect an improvement in the trained frequency, and some transfer   296

between low and broad frequencies, and high and broad frequencies, but no transfer between high and low frequencies (fig 2a). We might also expect frequency-tuned improvements in contrast sensitivity, mediated by the training on global motion stimuli. A second possibility is that learning depends on re-entrant connections from V5 to V1 [107], consistent with the reverse hierarchy model [55], but that this feedback is limited by the spatial frequency tuning of V5 mechanisms. Under this model, we would expect improvement in performance for low- and broad-frequency test stimuli, regardless of the frequency tuning of the stimuli used in training. Additionally, if the learning mechanism for global motion operates at the higher levels, between the directionally-tuned sensory neurons in V5 and the decision stage [36, 102] then a different pattern of transfer might be expected, with a pronounced effect for low frequencies, for which these mechanisms are tuned, more modest effects for broadband stimuli, and minimal transfer from high frequencies (fig 2b). Finally, if global motion training improves contrast sensitivity, we would expect to find some specificity to the trained frequency of global motion stimuli (fig 2c).

Fig 2. a) If transfer occurs at the local level within V1, improvement across frequencies for global motion would be predicted to occur for stimuli with shared properties. Training on Broad Frequency stimuli should improve performance on both High and Low Frequency stimuli. Training on Low frequency should result in some improvement for Broad Frequency stimuli, as would training on High Frequency stimuli. We would not expect any transfer of learning between Low and High frequencies b): If training for global motion transfers at the global level we predict that, as a result of the preference for low frequencies, a more selective transfer would occur c) Finally, if training on global motion improves contrast sensitivity we would expect that this would occur based on shared spatial frequency properties

The purpose of this study was to investigate the specificity of spatial frequency tuning in global motion after training and to determine the extent to which training with global motion stimuli is specific for the spatial frequency content of the training stimuli. In order to assess this, observers completed pre- and post-assessments in all frequencies of global motion, both frequencies of contrast sensitivity and a control task for orientation discrimination. The control task was added as a control for the possibility that training is simply an increased ability resulting from participating in psychophysical experiments. However, prior to collecting the data for this study we questioned whether trial-by-trial feedback was a requirement to learning.

Experiment 1: Is feedback necessary for learning?

The objective of this experiment was methodological in nature, in that we investigated the necessity of feedback for perceptual learning to occur for our specific stimuli. Performance feedback notifies a learner how accurate their performance is, and can be

generated internally (unsupervised learning) or provided by an external source [325] (supervised learning) [109]. External feedback often takes the form of an auditory or [326] visual cue that is presented after a trial to indicate a correct or incorrect response. [327] Behavioural perceptual learning studies have shown that using external feedback can [328] improve learning and increase efficiency [110,111]. On the other hand, internal feedback [329] is an internally generated benchmark to one's own performance on a task [109]. It has [330] been hypothesised that learning should occur without external feedback, if the [331] observer's internal confidence and accuracy are sufficiently high [22]. [332]

In a study conducted by Seitz et al., (2006) [112], learning did not occur without [333] feedback even when easy trials were presented. However, some studies have found that [334] learning occurs without external feedback [11,15,25,36,61,106]. This results in a [335] complicated pattern of empirical findings, and the specific nature of feedback and its [336] role in perceptual learning is unclear. Recently, Lu and Dosher (2012) [106] found that [337] interleaving high accuracy (easy) trials and low accuracy (difficult) trials resulted in [338] perceptual learning without the need for feedback even on difficult trials. Based on the [339] findings of Lu and Dosher (2012) [106] we predicted that we would find learning in both [340] conditions, as long as there were easy and difficult trials presented in an interleaved [341] method. As detailed in the following sections, our study found that learning did not [342] occur in the condition where no feedback was provided, even when easy trials were [343] presented. Learning was only found when feedback was presented. With this in mind [344] our design for experiment 2 included feedback during training, but no feedback when [345] testing. [346]

Methods and Materials: [347]

Observers [348]

24 observers were randomly and evenly assigned into a feedback or a no-feedback group. [349] All observers were employees or students from the University of Essex and self declared [350] as having normal or corrected-to-normal vision. All work was carried out in accordance [351] with the Code of Ethics of the World Medical Association (Declaration of Helsinki). [352] The study procedures were approved by the University of Essex University Ethics [353] Committee (JA1601). All observers gave informed written consent and were either paid [354] or received course credit for their participation. [355]

Apparatus [356]

Stimuli were generated and presented with Matlab 2015a using the Psychophysics [357] Toolbox extensions [113–115]. The broadband global motion stimuli were presented [358] using a 27" 2.7 Ghz IMac running OSX 1.9.5 iMac with a display resolution of 2560 x [359] 1440 pixels and a 60 Hz refresh rate. Viewing distance was 450mm, the stimuli [360] subtended a visual angle of 66.8°, and one pixel subtended 1.77 arc minutes. [361]

Stimuli and Procedure [362]

Global motion stimuli contained 100 Gaussian elements, each with a standard deviation [363] of 6.8 arc minutes. Elements were presented within a mid-grey rectangle measuring [364] 17.6° x 17.6° on a mid-grey background. Elements moved 5 pixels per frame, and each [365] element moved a fixed distance of 8.8° . Dots wrapped around the rectangle when [366] approaching the edges (fig 5a). Global motion stimuli were based on the task designed [367] by Williams and Sekuler (1984) [90]; dots move around the screen at 7 levels of [368] coherence (180°; 300°; 330°; 340°; 345°; 350°; 355°). Lower numbers depict higher [369] coherence, and higher numbers depict lower coherence and thus more randomness. At [370]

each level of coherence each dot moves a fixed distance within the defined arc of degree of coherence, while moving in a leftwards or rightwards direction (fig 3).

All tasks were presented as two-alternative-forced-choice (2AFC) decisions (left or right response), using the method of constant stimuli (MOCS). Observers responded by pressing the left or right arrow key associated with perceived left and right motion. When feedback was present, this was provided as an immediate auditory beep after each trial, a high pitched tone for a correct response (2000Hz for 10ms), and a low pitched tone for an incorrect response (200Hz for 40ms).

Fig 3. The random walk that creates the motion of dots with a 90°coherence. The arc defines the potential area of movement each dot could take at that coherence. The arrow indicates the actual trajectory and motion of the dot on each step. On the right; possible trajectories for the random walk for stimuli with 180°, 270°and 360°coherence.

General Statistical Methods

Moscatelli et al., (2012) [116] proposed using the Generalised Linear Mixed Effects Model (GLMM) for psychophysical data. The GLMM is an extension of the General Linear Model (GLM), and one that provides a more robust statistical analysis where the data contain irregular response distributions. The GLMM contains both fixed and random effects. The fixed component estimates are the effects of interest, in our study they are a) the day (or session) of testing and b) each level of the stimulus (coherence, contrast or orientation). The modelling of random effects assesses the differences between related groups (such as those from different observers) that allow inference to a larger population [117].

The linear component of the model is given by:

$$\eta = \alpha + \beta X \qquad (1)$$

where α and β are the intercept and the slope of the fixed effects parameters respectively. The link function converts the expected outcome variable (the proportion of correct responses, p) to the linear predictor [118]. Here, a logit function was used, adapted to take account of change performance, such that:

$$ln\frac{p - 0.5}{1 - p} = \alpha + \beta X \qquad (2)$$

Selecting the model: All response data were analysed using a GLMM conducted using Matlab (using the *fitglme* function), the logit link function and REMPL (Restricted maximum pseudo likelihood) to estimate the model parameters. Each model included observers as a random effects, and Time and Level and their interaction as fixed effects. Each model was assessed for goodness of fit, using random slopes or intercepts only, or their combination. Models (3a-3d) were compared using the likelihood ratio to test which model produced the lowest model fit statistics [116]. The Akaike information criterion (AIC) is one of the best known criteria to use in selecting a good model, where fitted values are most likely to be true values, thus providing a fit closest to reality [117]. The best regression model will provide the lowest value of AIC.

$$R \sim T + L + T \times L + (1|O) \tag{3a}$$
$$R \sim T + L + T \times L + (1 + T|O) \tag{3b}$$
$$R \sim T + L \ + T \times L + (1 + L|O) \tag{3c}$$
$$R \sim T + L + T \times L + (1 + L|O) + (1 + T|O) \tag{3d}$$

Where R is the observer response (proportion correct) T is time, the day or session of testing, L is the the stimulus level (degree of coherence, level of contrast, or degree of orientation) and O is the observer.

Interpreting the model The intercept is the proportion of correct responses when the linear model value is zero. An increase in intercept over time would reflect an overall increase in the number of correct responses. The slope relates to how the proportion of correct responses increases with the strength of the stimulus (degree of coherence, amount of contrast, or degree of orientation). Overall performance can also be summarised in terms of the 75% threshold, the point at which observers responded correctly on 75% of trials.

Results

To verify the spread of easy and difficult trials across the stimulus levels suggested by Liu et al. [106], accuracy for the first day was evaluated across all observers for each level of the stimulus and is reported in (Table 1). The accuracy across levels confirms that there is an even range of easy trials (85% and above) difficult trials (65% and below) with the balance around 75%.

Table 1. Accuracy at each level of the stimulus.

Stimulus Levels	Coherence °	Accuracy (%)
180	180	98.1
300	60	96.0
335	25	80.4
340	20	77.7
345	15	73.5
350	10	64.1
355	5	61.5

To identify if easy and difficult trials were sufficient for learning in the absence of trial-by-trial feedback, we analysed individuals' daily performance across the two training groups (Feedback or No Feedback). Individual responses from the 10 days' training were aggregated for each variable; observer, day and level of coherence. The response data were analysed independently for each group (Feedback or No Feedback)

using the method outlined above. Response was modelled as a fixed effect of day and coherence and an interaction between these two predictors. Results are presented in table 2. For the Feedback Group, there was a significant negative effect of day however there was a significant positive Day × Coherence interaction indicating there was a change in slope across the 10 days. The same analysis was undertaken for the No Feedback group, however, and there was a main effect of day and a significant negative interaction. The no feedback group performed worse after 10 days training. Performance over the 10 days is illustrated in fig 4, where the slope for the feedback group gets steeper over time, day 10 (blue) versus day 1 (red), along with the 75% discrimination threshold, which shifts leftwards, indicating a reduction in threshold between the start and end of training. Conversely, the group trained without feedback performs progressively worse over time and the 75% discrimination threshold shifts rightwards, indicating an increase in threshold.

Table 2. Fixed effects parameters for the Pilot Data

Training		β	SE	tStat	DF	pValue	LCI	UCI
No Feedback	Intercept (Day 1)	-1.595	0.1555	-10.258	836	2.456e-23	-1.90073	-1.2902
	Day (Change)	-0.0361	0.074	-0.4886	836	0.625	-0.1812	0.1089
	Coherence	0.07576	0.00207	36.598	836	8.65e-176 †	0.0717	0.0798
	Day × Coherence	-0.00075	0.00032	-2.30953	836	0.02116 *	-0.00139	-0.0001
Feedback	Intercept (Day 1)	-0.6099	0.140	-4.35	836	1.527e-05	-0.885	-0.335
	Day (Change)	-0.097	0.0076	-12.78	836	2.78e-34*	-0.112	-0.082
	Coherence	0.0292	0.0016	18.39	836	1.096e-63†	0.026	0.0324
	Day × Coherence	0.0087	0.0004	24.3234	836	3.052e-99*	0.00797	0.0094

Table notes :* indicates significant p value
and † performance increases significantly with increasing coherence

Fig 4. Mean thresholds (in degrees) over 10 days on global motion training for feedback (red) and no feedback (blue). Error bars represent ± SE.

Discussion

The purpose of the first experiment was to establish the necessity of performance feedback when undertaking a period of training on a task to discriminate the direction of global motion. Fahle and Edelman (1993) [22] predicted that internal reinforcement could act as the teaching signal when performance feedback was absent and when the confidence is high [53, 119]. Therefore, perceptual learning should occur when training procedures include a mixture of easy and difficult trials. This was the case for the study conducted by Lu and Dosher (2012) [106] where learning occurred for easy and difficult trials without feedback. For this specific task, consistent with Seitz et al., (2006) [112], our results found that feedback was a requirement for learning. After ten days of training the group who received feedback improved significantly, while the group receiving no feedback did not improve (worsened).

Feedback and Perceptual Learning

Our predictions were based on the findings by Lu and Dosher (2012) [106] who demonstrated that interleaving high accuracy (easy) trials and low accuracy (difficult) trials, resulted in perceptual learning without the need for feedback. Lu and Dosher (2012) [106] explain their findings using the Augmented Hebbian Re-weighting Model (AHRM) [33]. When external feedback is provided, the post-synaptic activation is shifted further in the correct direction, enforcing appropriate weight changes in the decision unit. However, when external feedback is absent the model acts in a similar manner to an unsupervised Hebbian model, using the observer's internal response. In this situation learning is dependent on the the level of difficulty of the task, and uses the observer's internal confidence to update the weights [106]. Where a task is easy, the weights still move, on average, in the correct direction [120]. For a difficult task, the correlation is weak and the process of updating the decision weights are slow and random, and learning does not occur.

The results of our study showed that learning did not occur without feedback, even when easy trials were presented. While our task did span the 65-85% accuracy range used by Lu and Dosher (2012) [106], there were some differences between the studies. Firstly, Lu and Dosher (2012) [106] used an adaptive staircase to track performance and stimuli were presented at either 85% and 65% accuracy throughout the experiment. Ours, on the other hand, did not include an adaptive staircase. Stimuli were randomly presented using MOCS. This method involves random stimulus selection from a predefined range of stimulus magnitudes [121]. In our experiment observers were randomly presented 1 of 7 threshold levels. Interleaving these levels may influence how decision weights are updated and reduce the observer's confidence in their judgements, for example by disrupting observers' meta-cognitive judgements of perceptual confidence [122] and thus their ability to selectively weight high-confidence trials in perceptual learning. [53]

The findings bear some similarity to those found by Seitz et al., (2006) [112], who also used MOCS. Seitz et al., (2006) [112] found that learning did not occur without feedback, even when easy trials were presented. Observers trained on one of two tasks, either to discriminate the direction of low luminance motion stimuli, or to discriminate the orientation of a bar that had been masked in spatial noise. After training, both groups who had received feedback showed an improvement, while the groups without feedback did not improve. Seitz et al., (2006) [112] note that many of the experiments investigating perceptual learning use adaptive staircase procedures, and even though easy trials were present, this difference may contribute to the difference in findings. Seitz et al., (2006) [112] propose that interleaving easy and difficult trials within the staircase may allow for "better bootstrapping" from easy to hard, compared to trials

that are randomly presented.

A second difference between the study conducted by Lu and Dosher (2012) [106] and ours was that while they used an orientation discrimination task, ours used a global motion coherence task. These tasks are known to be processed at different levels of the visual hierarchy. Learning without feedback has been obtained for local motion tasks. Ball and Sekuler (1982) [15] found that observers did not need feedback to improve on their direction discrimination task. In their study, observers were required to make a same/different judgement for two rapidly presented trials. In the "same" trials, motion took the same direction, and in the "different" trials the direction of motion varied by 3°. However, while the no feedback group did not receive trial-by-trial response feedback, they were rewarded with two cents for a correct response and had one cent deducted for each incorrect response, which may be construed as end of block feedback [112], which has been found to be as effective as trial-by-trial feedback [111].

Learning without feedback has also been found for global motion coherence tasks [61, 123]. However, none of these examples used the equivalent noise coherence task, but rather used the signal/noise coherence task. Huxlin et al., and Levi et al., [5, 58] using an equivalent noise method of global motion, also found perceptual learning, however they used an adaptive staircase for training. When investigating task-irrelevant learning, Watanabe et al., (2002) [124] found that task-irrelevant local motion improved passively, but did not find the same for task-irrelevant global motion. They suggest that this is indicative of the lower levels of the visual hierarchy being more receptive to modification, when attention is limited.

The results of this study ultimately provide us with evidence that to obtain learning during training, using the equivalent noise motion coherence used by Levi et al., (2014) [58], and presenting randomly these via MOCS, our procedure should include trial-by-trial feedback during the training process. Furthermore, since learning did not occur without feedback, no feedback would be provided during the pre- and post-assessment phases.

Experiment 2: Does learning transfer from global motion?

Methods and Materials

Observers

30 new observers, 12 male and 18 female, took part, randomly and evenly distributed across the three conditions, resulting in a total of 10 individuals in each condition.

Design and Apparatus

A series of baseline measures were completed for global motion direction discrimination, contrast sensitivity and orientation discrimination (pre-and post-training assessments). Assessment stimuli were presented on a VIEWPixx/3D 23.6 inch monitor with a display resolution of 1920 x 1080 pixels, with a 120 Hz refresh rate, using a Dell Precision T3610 PC running Windows 7. One pixel subtended 1.6 arc minutes and stimuli were viewed from a distance of 570mm. Head position for testing was stabilised using a chin rest. Following this, all observers undertook five consecutive days of global motion training in one of three spatial frequency groups (broad, high or low). Training stimuli were presented on a 19" monitor with a display resolution of 1980 x 1080 pixels and 60 Hz refresh rate, using a PC running Windows 7. One pixel subtended 1.7 arc minutes. Stimuli were viewed from a distance of 500mm. All stimuli were presented for 1 second.

Stimuli

Global Motion: Broadband stimuli were the same as previously described. For low-frequency stimuli, the elements were circularly symmetric Gabor patches. The standard deviation of the Gaussian window, σ, was 30.1 arc minutes, and the spatial frequency of luminance modulation, f, was 1 cycle/degree. For each element, the luminance profile was defined as a function of horizontal and vertical position (x, y) as:

$$d(x, y) = \sqrt{(x - x_0)^2 + (y - y_0)^2} \qquad (4)$$

$$L(d) = \frac{A}{\sigma\sqrt{2\pi}} \exp\frac{-d^2}{2\sigma^2} \cos 2\pi f d \qquad (5)$$

where x_0 and y_0 is the central position of the element, and A determines the contrast. Elements for the high frequency stimuli were defined in the same way, but had a standard deviation of 7.48 arc minutes and a spatial frequency, f, of 4 cycles/degree. For all stimuli, elements were initially uniformly randomly distributed within a region of 16.6°x 16.6° on the centre of the screen. A central black fixation dot was presented at all times when stimuli were not being displayed. Examples of the stimuli are shown in fig 5 a-c. Motion was created using the method detailed in fig 3. On each frame each dot moved a fixed distance of 8.5 arc minutes.

Fig 5. Global motion stimuli, at broad (a), low (b) and high spatial frequencies (c).

Pre- and Post-training Assessment

Global motion: Stimuli were identical to those described in the training session, with the following exceptions. The standard deviations of the elements were 6.4 arc minutes (broadband), 28.4 arc minutes (low frequency) and 7.0 arc minutes (high frequency). Stimuli were presented within a mid-grey rectangle measuring 15.9°x 15.9°, each element moved a fixed distance of 8 arc minutes.

Contrast Sensitivity: Stimuli were Gabor patches, with a spatial frequency of 1 cycle per degree (/°) or 4 cycles/°, presented in the centre of the screen on a mid-grey background, tilted either $\pm$ 20° (fig 6a&b). The Gaussian envelope of the Gabor stimulus had a standard deviation of 1.1°. 7 levels of contrast (0.05, 0.1, 0.15, 0.175, 0.2, 0.3, 0.4 % Michelson Contrast) were presented.

Orientation Discrimination: Gabor patches were presented for 100ms on a mid-grey background, measuring 7.9° x 7.9°, and presented 6.6° either above or below the central fixation point. Orientation was centred at zero, at which point lines were vertical. There was a total range of 1.8° difference in tilt, between +/- 0.9°, with 7 linearly spaced points within the range ($\pm$ 0, 0.67°, 1.3° or 2°). The spatial frequency of the Gabor patch was 1.85 /° and the standard deviation of the Gaussian envelope was 0.2° fig 6c. Contrast was fixed at 50% Michelson Contrast.

Fig 6. Gabors for a & b) contrast sensitivity for a) low spatial frequency- 1 cycle per degree, (b) high spatial frequency- 4 cycles per degree, and c) orientation discrimination which is tilted leftwards at 2° with mid spatial frequency- 2 cycles per degree

Procedure

Training on Global Motion

For each observer, training was undertaken at one spatial frequency only, totalling 420 trials daily for 5 continuous days. Feedback was provided after each trial.

Pre- and Post-Assessments

Measures were taken for global motion (high, broad and low spatial frequency), contrast sensitivity (high and low spatial frequency) and orientation discrimination (angle of tilt from the vertical) of an oriented Gabor patch. Responses were captured on the DataPixx response box for contrast sensitivity, and left and right arrows on the keyboard for global motion and orientation discrimination. The presentation order of trials was randomised by contrast and orientation or angle. There were 20 repetitions for each of the seven levels, for each condition. Testing was performed in a darkened room, before and after training.

Results: Experiment 2

Training Data

The number of correct responses from the daily training was calculated for each observer, for each day and level of coherence. Training data were analysed independently for the three groups trained on different frequencies (Broad, Low, High) and were modelled as a fixed effect of day and coherence and an interaction between these two predictors.

Results were analysed using the method previously described and are reported in tab 3. Performance improved for the broad and low trained groups and there was a significant positive coherence by day interaction found for both conditions. For the group trained with high spatial frequency stimuli there was a significant main effect of day. This suggests that while there was an significant increase in the total number of correct responses over the five days, this did not vary as a function of of coherence fig 3.

Table 3. Fixed effects parameters for the Training Data

Training		β	SE	tStat	DF	pValue	LCI	UCI
Broad	Intercept	-2.4642	0.74997	-3.2858	346	0.00111	-3.9393	-0.989
	Day	0.05161	0.0167	3.0869	346	0.00219*	0.0187	0.084
	Coherence	0.04848	0.0016	30.009	346	2.54e-98 †	0.0453	0.0517
	Day $\times$ Coherence	0.00271	0.000436	6.218	346	1.45e-09*	0.00185	0.00356
Low	Intercept	-1.081	0.5887	-1.8359	346	0.0672	-2.2388	0.0771
	Day	0.0421	0.120721	0.349	346	0.728	-0.19536	0.27952
	Coherence	0.0216	0.0012	18.1634	346	2.991e-52 †	0.0193	0.023981
	Day $\times$ Coherence	0.00331	0.0004	8.182	346	5.383e-15*	0.00251	0.0041
High	Intercept	-1.9579	0.1366	-14.336	346	6.56e-37	-2.227	-1.689
	Day	0.0894	0.02066	4.32719	346	1.98e-05*	0.048755	0.13001
	Coherence	0.0746	0.0026	28.3151	346	4.13e-92†	0.069	0.0798
	Day $\times$ Coherence	-0.0015	0.00079	-1.857	346	0.0642	-0.002995	8.61e-05

Table notes :* indicates significant p value
and † performance increases significantly with increasing coherence

Fig 7. Training results over the 5 days for broad, low and high spatial frequencies. The red and blue dotted lines show 75% threshold on days 1 and 5 respectively.

Pre- and Post-Assessment Analysis: Global Motion

Learning is often measured through monitoring performance at a particular threshold, which is expected to shift the psychometric function leftwards if performance is improved, 8A. This threshold describes performance in terms of accuracy as a function of the strength of the stimulus [125] which are usually positively correlated, and is

expected to reach asymptotic performance at the highest stimulus intensity [126]. 596
Inspection of the pre- and post-assessment data revealed that in some conditions 597
performance did not reach perfect accuracy, asymptoting at a proportion of correct 598
responses that was less than 1; this resulted in a poor psychometric fit of the observer 599
response data using the GLMM. To accommodate this, a nonlinear generalised mixed 600
effects model (NLME) was used to include an additional parameter in order to model 601
variability in the asymptotic performance at high signal levels, as has been applied in 602
other perceptual learning studies [127–129]. 603

Nonlinear regression analysis The nonlinear regression provides three measures to 604
assess a change in performance over time. Firstly, like the GLMM a leftward shift in the 605
curve indicates an improvement in threshold (8A). An increase in slope indicates an 606
increase in the rate at which performance increases with increase signal level (8B). 607
Finally a change in the asymptote indicates a significant change to the performance at 608
the highest level of stimulus intensity (8C). These changes are independent aspects of 609
the psychometric function fit, and may not necessarily always be congruent. For 610
example, it is possible to obtain an increase in one measure and a decrease (or no 611
change) in another. 612

Fig 8. Psychometric functions illustrating the three measures by which the non-linear regression provides evidence of a
change. **A:** A leftward shift of the function indicates a general increase across stimulus levels, that did not vary as stimulus
intensity increased. **B:** A steeper slope indicates an increase in the number of correct responses as stimulus intensity increases.
C: An upward shift of the asymptote indicates an increase in performance where stimulus intensity is at its highest.

Thus analysis of the pre- and post-assessment data was undertaken using a nonlinear 613
mixed effects regression with the *nlmefit* function in Matlab with the following model: 614

$$p = 0.5 + \frac{(A + A_d C)}{1 + e^{-(K + K_d S)(C - (C_0 + C_{0d} S))}} \tag{6}$$

where p is the proportion of correct responses, A determines the asymptotic level of 615
performance, K defines the slope and C_0 defines the threshold. A_d, K_d and C_{0d} 616
determine the change in these parameters after training, C is the coherence level, and S 617
is a dummy variable, taking on values of 0 or 1 for the pre- and post-training sessions, 618
respectively. 619

95% confidence limits were calculated using parametric bootstrap for 1000 simulated 620
experiments, with the same number of simulated observers and repetitions as the actual 621
experiment. Results for each condition are plotted in figs 9 to 11. 622

We proposed that should transfer occur at the local level within V1, improvement 623
across frequencies for global motion would be predicted to occur for stimuli with shared 624
frequency properties. Training on Broad Frequency stimuli should improve performance 625
on both High and Low Frequency stimuli. Training on Low frequency should result in 626

some improvement for Broad Frequency stimuli, as would training on High Frequency stimuli. We would not expect any transfer of learning between Low and High frequencies (see fig 13a). However, if transfer occurred at a global level, a more selective transfer would occur, mirroring the frequency tuning of V5 (see fig 13(b)).

Broad-frequency trained group

We predicted that should we obtain frequency specific transfer, then the Broad Frequency trained group should show some improvement in performance in the broad, low and high spatial frequency tests. The Broad trained group (fig 9) showed no consistent improvement in performance at post-testing. Most strikingly, when training and testing with **broad frequency** stimuli, there was no significant change in slope or threshold, and a significantly lower asymptote. For the **low frequency** test, no significant change was found for any of the three measures. Finally for the **high frequency** test, there was no change in slope, a significant reduction in asymptote, and a significant reduction in threshold. The increase in sensitivity at low signal levels in this condition was the only significant improvement found for the broad trained condition.

Low-frequency trained group:

The low-frequency trained group showed the greatest improvement and transfer (fig 10). For the **broad frequency** test stimuli, there was a general upward shift in the asymptote but an increase in threshold and no change to the slope. This shows reduced sensitivity at low signal levels, but an increase is performance at higher levels. For the **low frequency** test stimuli, a reduction in threshold and increase in asymptote show clear overall improvement in performance. For the **high frequency** test stimuli, a reduction in threshold also indicates a significant improvement in performance at lower stimulus coherence. The significantly shallower slope suggests that at post-assessment the proportion of correct responses decreased for higher stimulus intensities in comparison with pre-test performance. Finally there was no change to asymptotic performance which was close to one at the pre-assessment phase suggesting little for improvement at the highest levels of coherence.

High-frequency trained group:

The group trained with high spatial frequency stimuli showed no improvement in any of the three spatial frequency conditions, for any of the three measures of threshold, slope or asymptote (fig 11).

Transfer : Global Motion

In summary, only the low frequency trained provided reliable evidence for transfer to untrained motion conditions, with some improvement in performance for broadband test stimuli.

Fig 9. Top: Assessment results for the Broad Frequency trained condition on Broad, Low and High spatial frequency tests. Performance is presented as the proportion of correct responses, where blue circles show the results for the pre-assessment, and pink squares for the post assessment. 75% thresholds for pre and post are indicated by the green and orange vertical dotted lines respectively.

Bottom: Change statistics for the asymptote, threshold and slope, respectively. Plots show the median performance and the 95% confidence intervals for the change in performance between pre- and post-assessments. The red horizontal line at zero represents no change, confidence intervals crossing the zero line reflect no significant improvement. Points above the reference line show an improvement in performance and those below reflect a decrease in performance.

Pre and Post Analysis: Contrast Sensitivity   663

We predicted that, should contrast sensitivity improve as a result of training on global   664
motion, then it would depend on the frequency of training (fig reffig:trangm top). Data   665
were analysed for all but the same 2 observers who did not complete the final session   666
using the generalised linear effects model previously outlined. The measures of analysis   667
are the intercept and slope , and an interaction between the two predictors.   668

Fig 10. Top: Assessment results for the Low Frequency trained condition on Broad, Low and High spatial frequency tests. Performance is presented as proportion of correct responses, where blue (circles) show the results for the pre-assessment, and pink (squares) the post assessment. 75% thresholds for pre and post are indicated by the green and orange vertical dotted lines respectively.
Bottom: Change statistics for the asymptote, threshold and slope respectively. Plots show the median performance and the 95% confidence intervals for the change in performance between pre- and post-assessments. The red horizontal line at zero represents no change, confidence intervals crossing the zero line reflect no significant improvement. Points above the reference line show an improvement in performance and those below reflect a decrease in performance

Broad-frequency trained group: 669

The group trained with broad frequency global motion showed a significant decrease in 670
performance between pre- to post-training conditions, and a significant increase in slope 671
between pre- and post-training. This suggests that their performance decreased at lower 672
levels of the stimuli, while increasing at higher signal levels. Because of this ambiguity 673
we plotted 75% thresholds for pre and post performance (fig 12 top left). 674
Post-assessment thresholds shifted rightwards suggesting worse performance overall. 675

Fig 11. Top: Assessment results for the High Frequency trained condition on Broad, Low and High spatial frequency tests. Performance is presented as proportion of correct responses, where blue (circles) show the results for the pre-assessment, and pink (squares) the post assessment. 75% thresholds for pre and post are indicated by the green and orange vertical dotted lines respectively.
Bottom: Change statistics for the asymptote, threshold and slope respectively. Plots show the median performance and the 95% confidence intervals for the change in performance between pre- and post-assessments. The red horizontal line at zero represents no change, confidence intervals crossing the zero line reflect no significant improvement. Points above the reference line show an improvement in performance and those below reflect a decrease in performance

Low-frequency trained group:

The group trained with low frequency global motion showed a significant decrease in threshold, and a significant increase in slope. Again, we plotted 75% thresholds for pre and post performance (fig reffig:csplot middle left). In this instance the steeper slope, and leftward shift of the threshold, suggests improvement overall.

Table 4. Contrast Sensitivity Table

Training	Low Frequency	β	SE	tStat	DF	pValue	LCI	UCI
Broad	Threshold (Pre)	-4.0361	1.3136	-3.0725	129	0.0026	-6.6352	-1.437
	Threshold Change (Post)	-1.4324	0.39039	-3.6691	129	0.00035*	-2.2047	-0.66
	Contrast	33.009	9.4797	3.4821	129	0.00068†	14.253	51.765
	Session × Contrast	8.4892	2.2901	3.707	129	0.00031*	3.9582	13.02
Low	Threshold (Pre)	-1.9253	0.66009	-2.9167	129	0.00417	-3.2313	-0.61929
	Threshold Change (Post)	-1.6317	0.4122	-3.9585	129	0.00012*	-2.4472	-0.81612
	Contrast	11.338	3.203	3.5397	129	0.00056†	5.0004	17.675
	Session × Contrast	13.035	2.4795	5.2571	129	5.90e-07*	8.129	17.94
High	Threshold (Pre)	-3.5587	0.78832	-4.5143	136	1.3627e-05	-5.1177	-1.9998
	Threshold Change (Post)	-0.20622	0.45674	-0.45151	136	0.65234	-1.1095	0.69701
	Contrast	24.238	3.6543	6.6329	136	7.14e-10†	17.012	31.465
	Session × Contrast	3.5416	2.4152	1.4664	136	0.14485	-1.2346	8.3177
	High Frequency							
Broad	Threshold (Pre)	-3.1141	0.72157	-4.3158	129	3.14e-05	-4.5417	-1.6865
	Threshold Change (Post)	-0.39475	0.28081	-1.4057	129	0.1622	-0.95034	0.16084
	Contrast	17.372	2.3389	7.4273	129	1.34e-11†	12.744	21.999
	Session × Contrast	2.3333	1.5341	1.521	129	0.13072	-0.70195	5.3686
Low	Threshold (Pre)	-3.388	0.41928	-8.0805	129	4.01e-13	-4.2175	-2.5584
	Threshold Change (Post)	0.0001	0.25711	0.00045	129	0.99964	-0.509	0.50881
	Contrast	22.041	2.9883	7.3757	129	1.76e-11†	16.128	27.953
	Session × Contrast	-1.236	1.1863	-1.0418	129	0.29943	-3.5831	1.1112
High	Threshold (Pre)	-3.5827	0.78951	-4.5379	136	1.2371e-05	-5.144	-2.0214
	Threshold Change (Post)	-0.045188	0.26063	-0.17338	136	0.86261	-0.56059	0.47022
	Contrast	20.64	2.2427	9.203	136	5.496e-16†	16.205	25.075
	Session × Contrast	-0.7794	1.3781	-0.56554	136	0.57264	-3.5048	1.946

Table Notes:* indicates significant p value
and † performance increases significantly with increasing coherence

High-frequency trained group: 681

The high frequency trained group showed no significant change for any condition (fig 12   682
bottom left).   683

Fig 12. Results for Broad (top row), Low (middle) and high (bottom) trained groups in the for the contrast sensitivity pre- and post-assessments. Low spatial frequencies are shown in the left column and high spatial frequencies on the right. Performance is presented as proportion of correct responses as a function of Michelson Contrast. Blue (triangles) show the results for the pre-assessment, and pink (crosses) the post assessment. 75% thresholds for pre and post are indicated by the green and orange vertical lines respectively.

Pre and Post Analysis: Orientation Discrimination

The orientation discrimination task was added as an untrained control, for which we predicted no improvement. Data were analysed for all but 3 observers who did not complete the final session, using the generalised linear effects model previously outlined. The measures of analysis are the intercept and slope , and an interaction between the two variables. For analysis, the degrees of orientation ($\pm$ 0, 0.67°, 1.3° or 2°) were

Fig 13. Figures a & b at the top show the predicted transfer. The bottom figure illustrates the actual transfer found

converted to absolute values. Results are plotted in fig 14, and the fixed effects are listed in tab 5

Fig 14. Assessment results for all trained groups (broad, low and high) for the orientation discrimination control task. Performance is presented as proportion of correct responses, where blue (triangles) show the results for the pre-assessment, and pink (crosses) the post assessment. 75% thresholds for pre and post are indicated by the green and orange vertical lines respectively.

Table 5. Fixed Effects table for Orientation Discrimination

Training		β	SE	tStat	DF	pValue	LCI	UCI
Broad	Threshold (Pre)	-1.9841	0.79477	-2.4964	284	0.013	-3.5485	-0.41967
	Threshold Change (Post)	0.02189	0.55128	0.039723	284	0.96834	-1.0632	1.107
	Orientation	2.8444	0.35047	8.1161	284	1.47e-14†	2.1546	3.5343
	Session $\times$ Orientation	0.0025	0.10152	0.0245	284	0.981	-0.197	0.2023
Low	Threshold (Pre)	-1.5733	0.78192	-2.0121	284	0.045	-3.1124	-0.034191
	Threshold Change (Post)	-0.38185	0.58052	-0.65778	284	0.511	-1.5245	0.76081
	Orientation	2.3483	0.29959	7.8381	284	9.25e-14†	1.7586	2.938
	Session $\times$ Orientation	0.18935	0.097122	1.9496	284	0.0522	-0.0018	0.38052
High	Threshold (Pre)	-1.1825	0.37792	-3.1289	284	0.00194	-1.9263	-0.43858
	Threshold Change (Post)	-0.47101	0.1002	-4.7005	284	4.05e-06	-0.66824	-0.27377
	Orientation	2.3751	0.33682	7.0516	284	1.35e-11†	1.7121	3.0381
	Session $\times$ Orientation	0.28973	0.08728	3.3196	284	0.001*	0.11794	0.46153

Table notes : † performance increases significantly with increasing coherence

Neither the broad- or low-frequency trained group showed any significant change on this task. However, the high spatial frequency trained group showed a significant increase in threshold over session, and a significant increase in slope following training. The data shows that the 75% threshold at post assessment shift rightwards, suggesting worse performance overall.

Discussion

Having established the necessity of feedback for learning, the objective of the main study was to investigate the specificity of spatial frequency tuning in perceptual learning for global motion. We trained three groups of observers on a global motion task with stimuli tuned to three different spatial frequency ranges (Broad, Low, High). Our training results demonstrated that perceptual learning took place over the five days for all groups. In order to establish the degree to which learning generalised between the spatial frequency content of the stimuli, we compared the results of the pre- and post-training assessments for each condition of global motion. Analysis from the post-training assessment showed that unambiguous improvement was restricted to the low frequency trained group. Notably the broad trained group performed worse than they did at pre-assessment stage, and there was no change for the high trained group.

We predicted that if training on global motion improved contrast sensitivity at the local level it would be specific to the trained frequency, and restricted to low and broad frequency stimuli if transfer occurred at the global level. Pre- and post-training assessment for static contrast sensitivity found that there was a significant improvement for the low trained group on the low spatial frequency contrast condition. While the broad trained group showed some improvement, there was overall a decrease in threshold. No improvement was found for the high spatial frequency trained group, even on the high spatial frequency contrast sensitivity. Furthermore, no group showed improvement in sensitivity for the high spatial frequency contrast condition. The orientation discrimination task was included as a control condition and no improvement for any trained frequency was predicted or found. Our study supports that the perception of global motion shows a bias for low spatial frequencies and suggests the re-entrant connections from V5 to V1 play a crucial role in transfer.

Why is the study of global motion important?

Damage to the visual cortex, as a result of stroke or other brain injury can result in dramatic changes to connectivity between areas. It has been proposed that perceptual training may play a role in strengthening existing neural pathways and may even create novel connections [1–7]. While specificity is a limitation to the effectiveness of perceptual learning as a tool for therapy, there is evidence that this specificity is reduced for visually impaired populations [130–135]. Huxlin et al., (2009) [5] proposed that using global motion may tap into "islands of activity" within V1 through the feedback connections from V5 to V1. However, it is still unclear how much the brain is able to compensate for damage, and whether recovery involves building new connections or changes in the functional connectivity within the existing pathways [136]. Understanding how the cortical hierarchy is organised in terms of the nature of the feedforward and re-entrant pathways is central to developing theories of perception [137].

This study sought to explore the effects of training on global motion and its transfer to other spatial frequencies and tasks . The back projections from V5 to V1 are known to be crucial for the perception of motion [138], strongly represented [74, 86] and rapidly updated [138, 139], making this a theoretically plausible route for perceptual learning. Recently Romei et al., (2016) [107] published evidence suggesting that the re-entrant connections from V5 to V1 are malleable [107]. A novel dual-coil Transcranial Magnetic Stimulation (TMS) protocol aimed at inducing Hebbian plasticity [140] was employed to enhance the perception of coherent motion by targeting area V5 and V1 [107]. Thresholds were lower after TMS application, but critically dependent on the inter-pulse timing and direction of stimulation between V5 and V1. Motion perception was found to be strengthened for the re-entrant connection from V5 to V1, but not the feedforward connection from V1 to V5. Recently, a similar paired cortico-cortical TMS

protocol (ccPAS), Chiappini et al., [108] induced a direction-selective improvement in ₇₄₇
performance by combining subthreshold stimulation with the simultaneous presentation ₇₄₈
of direction-specific moving stimuli. This provides accumulating evidence that the ₇₄₉
re-entrant connections from direction-tuned neurons play a role in perceptual learning ₇₅₀
in global motion coherence tasks. Furthermore, recordings obtained from macaque ₇₅₁
monkeys have previously suggested that there is no delay for information processed in ₇₅₂
V5 to be fed back to lower areas [139]. Hupe et al., (2016) [139] recorded extracellular ₇₅₃
responses from neurons in V1, V2 and V3 after inactivating V5 and compared the time ₇₅₄
course of responses for moving and flashed stimuli. Their results indicated there was no ₇₅₅
delay for either case and Hupe et al., (2016) [139] proposed that the feedback from V5 is ₇₅₆
present prior to the bottom up information from the feedforward connections, however ₇₅₇
they were not able to test this hypothesis. More recently, Bridge et al., (2016) [136] ₇₅₈
found evidence of an independent motion pathway in human observers, using ₇₅₉
diffusion-weighted MRI (see [141] for more information), that suggests a direct ₇₆₀
ipsilateral link between the lateral geniculate nucleus (LGN) and V5. These findings ₇₆₁
question the straightforward hierarchical structure of cortical connectivity and the ₇₆₂
generally assumed flow of information within the visual system. ₇₆₃

Perception and learning of contrast stimuli ₇₆₄

Our starting point for the investigation was the results from Levi et al., (2014) [58] who ₇₆₅
found that learning generalised from high-level global motion training to low-level ₇₆₆
contrast detection tasks. The pathway for contrast detection and discrimination is ₇₆₇
believed to develop during childhood and adolescence, reaching its peak in adulthood, ₇₆₈
followed by a decline in late adulthood [142]. Early psychophysical studies supported ₇₆₉
this view, finding little perceptual learning for contrast-dependent stimuli [17]. However, ₇₇₀
later findings [143] suggested that improvement in contrast sensitivity was dependent on ₇₇₁
having enough training. Improvement in contrast sensitivity is now a well documented ₇₇₂
result [48, 49, 143–146] but has been found to be highly tuned to spatial frequency, ₇₇₃
specific for trained eye and retinal location of training, but not selective for ₇₇₄
orientation [21]. Sowden et al., (2002) [21] trained observers to detect the orientations of ₇₇₅
sine-wave gratings at varying contrasts. Results indicated that thresholds were reduced, ₇₇₆
and that this learning was tuned for spatial frequency and specific to the trained ₇₇₇
location, but not specific for the orientation of the grating. As a result, they suggested ₇₇₈
that the mechanism for contrast detection is located in a sub-population of cells in V1, ₇₇₉
such as those in $4C\alpha$, where cells are tuned for spatial frequency but not orientation [21]. ₇₈₀
  Since Levi et al., (2014) [58] reported the biggest improvement for low spatial ₇₈₁
frequency stimuli in drifting targets, this may suggest improvement was specific to the ₇₈₂
temporal and spatial features of the training stimuli. We had questioned if the direction ₇₈₃
selective cells in the area might account for the improvement in contrast sensitivity. ₇₈₄
Thus, our paradigm tested this for static frequency specific stimuli. Our results were ₇₈₅
consistent with those found by Levi et al., (2014) [58] in that improvement occurred for ₇₈₆
low frequency stimuli, and only for the groups who trained on broad and low frequency ₇₈₇
global motion. There was no improvement on the high spatial frequency (4 cylces per ₇₈₈
degree) contrast stimuli for any trained condition. ₇₈₉

Learning and transfer for global motion training ₇₉₀

Based on the evidence for the frequency tuning of V5 [76, 77, 84, 95], we predicted that ₇₉₁
we might have obtained greater improvements (and transfer) between broad and low ₇₉₂
spatial frequencies, but limited or no transfer for the high spatial frequency trained ₇₉₃
group. Furthermore, we hypothesised that if learning occurred in the lower, local levels ₇₉₄
of processing, we may find transfer between the spatial frequency specific training of ₇₉₅

global motion stimuli to contrast sensitivity. With the higher level pooling of low spatial 796
frequencies we expected that if transfer took place as a result of global motion, then we 797
would predict some transfer between those frequencies containing low spatial frequency 798
information, specifically between the low and the broad conditions. After the five day 799
training period, there was significant improvement for the groups trained for all spatial 800
frequencies 801

When assessing the post-training transfer to untrained global motion frequencies, the 802
low spatial frequency trained group was the only group to show clear improvement in 803
performance, and evidence of transfer to another condition. Although threshold was 804
worse for the broad test after training with low frequency stimuli, the asymptotic 805
performance was better. Interestingly, for the high spatial frequency test, there was a 806
significant improvement to the threshold although the slope was significantly shallower. 807
This suggests that the sensitivity for lower coherence levels increased. Performance at 808
the highest levels reached an asymptotic performance close to 1 at pre-assessment and 809
remained unchanged at the post-assessment stage. Finally the shallower slope suggests 810
a reduction in correct responses as a function of stimulus intensity. 811

The *high spatial frequency trained group* showed no improvement and no transfer to 812
any other spatial frequency. The most surprising results were those obtained from the 813
broad frequency trained group. Asymptotic performance was significantly worse at the 814
post test stage for their own trained frequency and the high frequency test, with no 815
significant change in the low frequency test. There was no improvement in the slopes for 816
any condition, and a small, but significant threshold improvement was found for the 817
high frequency test 818

What our findings suggest for the models of perceptual learning 819

The two main theories of visual perceptual learning suggest that learning either occurs 820
in the lower levels within the receptive fields of sensory neurons [17, 18, 26], or at a 821
higher more cognitive level [34] as a reweighting of the decision unit. We predicted that 822
should global motion training improve contrast detection, then based on the frequency 823
tuning of the early visual area, we would expect specificity for the spatial frequency of 824
training. In this case we would expect improvement on the low frequency contrast 825
stimuli for low and broad trained groups and for the high frequency contrast stimuli 826
improvement from the high and broad trained groups. Improvement was exclusive to 827
the low frequency contrast condition for the low trained group. The specificity of 828
transfer to low frequencies is consistent with V5 influencing processing at V1 through 829
feedback loops [107]. This remains compatible with the view that global motion 830
detectors pool for low frequency information. Suggesting that the re-entrant connections 831
to V1 after global motion training only update low frequency channels. 832

We predicted that should transfer occur between motion conditions it would be most 833
likely to occur either between groups that shared frequency content, or based on the 834
frequency turning of V5. The group trained on high spatial frequencies showed no 835
improvement and no transfer. Since high spatial frequency content has been shown to 836
be attenuated in V5 [95] this again suggests that improvements for the untrained tasks 837
are absent due to the updating of low frequency channels. The groups trained on low 838
and broad frequency motion conditions showed transfer to low spatial frequency contrast 839
detection, also supporting role of the back projections to low frequency channels. 840

The low frequency training condition also showed some improvement for the broad 841
motion test stimuli, with a significant increase in asymptotic performance. The low 842
frequency (and to a lesser part the broad) conditions will have experienced a high level 843
of correlation in the activity in frequency channels between the feedforward and 844
feedback connections. We speculate that this joint activity may be important for 845
perceptual learning. The improvement for the low frequency contrast detection 846

condition from the broad and low trained groups are consistent with this hypothesis. ⁸⁴⁷

Dosher and Lu (2010) [33] suggest that the reweighting of decision units is likely to ₈₄₈ be the dominant mechanism of learning. However, if as proposed by Dosher et al. ₈₄₉ learning occurred primarily as a result of an increase to the weight at a higher level ₈₅₀ decision making unit, we may have expected that the learning from the training, at the ₈₅₁ very least, be maintained at the assessment stage. However, our results found ₈₅₂ performance at post-assessment stage were not unambiguously better for all groups. ₈₅₃ The only difference between the exercises was that trial-by-trial feedback was present ₈₅₄ for the training phase and absent for the testing phase. Our pilot study found that, for ₈₅₅ our methods, learning occurs with feedback, but not without. Additionally, Herzog and ₈₅₆ Fahle (1997) [111]) suggested that once feedback is removed, performance plateaus ₈₅₇ around the level last obtained with feedback, so our expectation was that observers ₈₅₈ would maintain the improvement achieved for their trained frequency. This was the ₈₅₉ motivation for removing feedback during testing. ₈₆₀

We offer some speculative ideas that may help account for the general absence of ₈₆₁ improvement at post-assessment stage. There is emerging evidence that interleaving ₈₆₂ random stimuli may disrupt perceptual learning [147]. Yu et al., (2004) [143] found that ₈₆₃ when interleaving random (as when using MOCS) contrasted Gabor stimuli, learning ₈₆₄ did not occur, but did when only one contrast was presented at a time. Yu et al., ₈₆₅ (2004) [143] termed the interleaving of contrast stimuli as "contrast roving", and suggest ₈₆₆ that the effect of non-learning as a result of contrast roving implies that a temporally ₈₆₇ organised pattern of stimulus presentation may be required for perceptual learning to ₈₆₈ occur. Additionally, Kuai ee al., (2005) [145] found that temporal patterning was evident ₈₆₉ for motion direction discrimination. Observers were either trained using randomly ₈₇₀ varied trials or with a fixed temporal pattern of exposure (changing in a sequential ₈₇₁ clockwise direction). The first groups showed no improvement in discrimination, ₈₇₂ however the second group with the fixed temporal exposure did. Cong and Zhang ₈₇₃ (2014) [148] showed that a semantic tag associated with each stimulus presentation ₈₇₄ enabled significant learning. Sequential tags such as A,B,C,D and 1,2,3,4 which contain ₈₇₅ semantic and sequential identity information both resulted in significant learning. Cong ₈₇₆ and Zhang (2014) [148] suggest that tagging and temporal sequencing may help direct ₈₇₇ the system to switch attention to the neuronal set responsible to the specific stimulus. ₈₇₈ These are consistent with the experiments conducted by Seitz et al., (2006) [112], which ₈₇₉ as previously discussed found no learning without feedback using MOCS. ₈₈₀

There is some evidence that interleaving easy and difficult trials for perceptual ₈₈₁ training may only lead to a temporary improvement, that (a) does not persist and (b) ₈₈₂ disappears when easy trials are removed [149,150]. In a difficult task without feedback ₈₈₃ learning does not occur, however it does with feedback [106], since feedback improves ₈₈₄ detection for difficult tasks. Removing the feedback may be equivalent to removing the ₈₈₅ easy trials. Within a training session comprised of difficult trials, Lin and Dosher ₈₈₆ (2017) [105] presented observers a single block of mixed easy and difficult trials which ₈₈₇ led to rapid sustained improvement in accuracy within the block. However, after ₈₈₈ removing the easy trials and presenting only the difficult trials performance slowly ₈₈₉ returned to prior levels. In our study, improvement was sustained only while feedback ₈₉₀ was present, with the exception for the low frequency trained group. Since learning and ₈₉₁ transfer was evident for the low spatial frequency trained group, this may suggest an ₈₉₂ interaction between the amount of learning, and the benefit of feedback. In the broad ₈₉₃ and high spatial frequency trained groups, the amount of learning that took place was ₈₉₄ not sufficient to overcome the reduction in performance when feedback is removed. In ₈₉₅ contrast, learning for the low spatial frequency trained group was robust and persistent, ₈₉₆ and thus incurred less disruption by the absence of feedback. ₈₉₇

Ultimately, neither model alone accounts for the findings in this study and, while the ₈₉₈

two positions are pitched as competing theories, they may not be mutually exclusive. In an attempt to combine the two models, Solgi and Weng (2013) [151] models learning as a two-way process, (descending and ascending) that attributes training effects to the connections between the early and higher sensory areas, and also as an increased representation in the lower level. For transfer to occur, Solgi and Weng (2013) [151] argue there is an important role for the re-entrant connections from higher levels, our results are consistent with Solgi and Weng (2013) [151] and suggest that the re-entrant connections play a vital role in transfer.

Conclusions

Our results support the Reverse Hierarchy Theory that the feedback connections from higher levels back to early visual processing areas may be involved in facilitating learning that transfers [54, 55], and are consistent with models that suggest that the the re-entrant connections play a vital role in learning and transfer [38, 151].

Our first experiment suggested that for perceptual learning of an equivalent noise global motion task, presented using MOCS, feedback was a necessity. However after training with feedback, there was little evidence of a robust improvement when feedback was removed at the post-assessment stage. The only condition that showed a unambiguous learning and transfer were the low spatial frequency trained group. Our findings are in line with current research which finds global motion detectors pool for low spatial frequencies.

Our main experiment found robust learning and transfer for the group trained with low spatial frequency specific global motion gabors. This has provided evidence of frequency specific transfer from in global motion perception, and to a static untrained task. The frequency tuning of these results is consistent with perceptual learning which depends on the global stage of motion processing in V5.

Acknowledgments

This research was funded by a University of Essex Doctoral studentship, and grants from ESSEXLab and PsyPAG to J.A.

References

1. Campana G, Maniglia M. Editorial: Improving visual deficits with perceptual learning. Frontiers in Psychology. 2015;6(APR):1–3. doi:10.3758/BF03194845.

2. Hess RF, Hayes A, Field DJ. Contour integration and cortical processing. Journal of Physiology Paris. 2003;97(2-3):105–119. doi:10.1016/j.jphysparis.2003.09.013.

3. Levi DM, Li RW. Perceptual learning as a potential treatment for amblyopia: A mini-review. Vision Research. 2009;49(21):2535–2549. doi:10.1016/j.visres.2009.02.010.

4. Polat U. Making perceptual learning practical to improve visual functions. Vision Research. 2009;49(21):2566–2573. doi:10.1016/j.visres.2009.06.005.

5. Huxlin KR, Martin T, Kelly K, Riley M, Friedman DI, Burgin WS, et al. Perceptual relearning of complex visual motion after V1 damage in humans. The Journal of Neuroscience. 2009;29(13):3981–3991. doi:10.1523/JNEUROSCI.4882-08.2009.

6. Sahraie A, Trevethan CT, MacLeod MJ, Murray AD, Olson JA, Weiskrantz L. Increased sensitivity after repeated stimulation of residual spatial channels in blindsight. Proceedings of the National Academy of Sciences of the United States of America. 2006;103(40):14971–14976. doi:10.1073/pnas.0607073103.

7. Trevethan CT, Urquhart J, Ward R, Gentleman D, Sahraie A. Evidence for perceptual learning with repeated stimulation after partial and total cortical blindness. Advances in Cognitive Psychology. 2012;8(1):29–37. doi:10.2478/v10053-008-0099-8.

8. Sagi D. Perceptual learning in Vision Research. Vision Research. 2011;51(13):1552–1566. doi:10.1016/j.visres.2010.10.019.

9. Seitz AR, Watanabe T. A unified model for perceptual learning. Trends in Cognitive Sciences. 2005;9(7):329–334. doi:10.1016/j.tics.2005.05.010.

10. Hubel DH, Wiesel TN. The period of susceptibility to the physiological effects of unilateral eye closure in kittens. Journal of Physiology. 1970;206:419–436.

11. Fahle M. Perceptual learning: Specificity versus generalization. Current Opinion in Neurobiology. 2005;15(2):154–160. doi:10.1016/j.conb.2005.03.010.

12. Ahissar M, Hochstein S. Task difficulty and the specificity of perceptual learning. Letters to Nature. 1997;387(6631):401–406. doi:10.1038/387401a0.

13. Fiorentini A, Berardi N. Perceptual learning specific for orientation and spatial frequency. Nature. 1980;287:43–44. doi:10.1038/287043a0.

14. Karni A, Sagi D. Where practice makes perfect in texture discrimination: evidence for primary visual cortex plasticity. Proceedings of the National Academy of Sciences of the United States of America. 1991;88(11):4966–4970. doi:DOI 10.1073/pnas.88.11.4966.

15. Ball K, Sekuler R. A specific and enduring improvement in visual motion discrimination. Science. 1982;218(4573):697–698. doi:10.1126/science.7134968.

16. Lu ZL, Hua T, Huang CB, Zhou Y, Dosher BA. Visual perceptual learning. Neurobiology of learning and memory. 2011;95(2):145–51. doi:10.1016/j.nlm.2010.09.010.

17. Adini Y, Sagi D, Tsodyks M. Context-enabled learning in the human visual system. Nature. 2002;415(February):2–5.

18. Gilbert CD, Sigman M, Crist RE. The Neural Basis of Perceptual Learning Review. Neuron. 2001;31:681–697. doi:10.1016/S0896-6273(01)00424-X.

19. Fahle M. Specificity of learning curvature, orientation, and vernier discriminations. Vision Research. 1997;37(14):1885–1895. doi:10.1016/S0042-6989(96)00308-2.

20. Schoups AA, Vogels R, Orban GA. Human perceptual learning in identifying the oblique orientation: retinotopy, orientation specificity and monocularity. The Journal of physiology. 1995;483.3:797–810.

21. Sowden PT, Rose D, Davies IRL. Perceptual learning of luminance contrast detection: Specific for spatial frequency and retinal location but not orientation. Vision Research. 2002;42(10):1249–1258. doi:10.1016/S0042-6989(02)00019-6.

22. Fahle M, Edelman S. Long-term learning in vernier acuity: Effects of stimulus orientation, range and of feedback. Vision Research. 1993;33(3):397–412. doi:10.1016/0042-6989(93)90094-D.

23. Fahle M, Edelman S, Poggio T. Fast perceptual learning in hyperacuity. Vision Research. 1995;35(21):3003–3013. doi:10.1016/0042-6989(95)00044-Z.

24. Poggio T, Fahle M, Edelman S. Fast perceptual learning in visual hyperacuity. Massachusetts Inst of Tech Artfical Intelligence Lab. 1991;.

25. Shiu LP, Pashler H. Improvement in line orientation discrimination is retinally local but dependent on cognitive set. Perception & psychophysics. 1992;52(5):582–588. doi:10.3758/BF03206720.

26. Gilbert CD, Li W. Top-down influences on visual processing. Nature Reviews Neuroscience. 2013;14(5):350–363. doi:10.1038/nrn3476.

27. Barlow HB, Foldiak P. The computing neuron. Adaptation and decorrelation in the cortex. 1989; p. 54–72.

28. Simoncelli EP. Vision and the statistics of the visual environment. Current Opinion in Neurobiology. 2003;13(2):144–149. doi:10.1016/S0959-4388(03)00047-3.

29. Wei XX, Stocker AA. Efficient coding provides a direct link between prior and likelihood in perceptual Bayesian inference. Advances in Neural Information Processing Systems. 2012;25(May):1–9.

30. Mollon JD, Danilova MV. Three remarks on perceptual learning. Spatial vision. 1996;10(1):51–58. doi:10.1016/j.aqpro.2013.07.003.

31. Herzog MH, Fahle M. Modeling perceptual learning: difficulties and how they can be overcome. Biological cybernetics. 1998;78(2):107–117. doi:10.1007/s004220050418.

32. Dosher BA, Lu ZL. Perceptual learning reflects external noise filtering and internal noise reduction through channel reweighting. Proceedings of the National Academy of Sciences of the United States of America. 1998;95(23):13988–93. doi:10.1073/pnas.95.23.13988.

33. Dosher BA, Lu ZL. Hebbian Reweighting on Stable Representations in Perceptual Learning. Learning & Perception. 2009;1(1):37–58. doi:10.1556/LP.1.2009.1.4.

34. Dosher BA, Jeter PE, Liu J, Lu ZL. An integrated reweighting theory of perceptual learning. Proceedings of the National Academy of Sciences of the United States of America. 2013;110(33):13678–13683. doi:10.1073/pnas.1312552110.

35. Herzog MH, Fahle M. A recurrent model for perceptual learning. Journal of Optical Technology. 1999;66(9):836.

36. Petrov AA, Dosher BA, Lu ZL. Perceptual learning without feedback in non-stationary contexts: Data and model. Vision Research. 2006;46(19):3177–3197. doi:10.1016/j.visres.2006.03.022.

37. Petrov AA, Dosher BA, Lu ZL. The dynamics of perceptual learning: An incremental reweighting model. Psychological Review. 2005;112(4):715–743. doi:10.1037/0033-295X.112.4.715.

38. Bejjanki VR, Beck JM, Lu ZL, Pouget A. Perceptual learning as improved probabilistic inference in early sensory areas. Nature Neuroscience. 2011;14(5):642–50. doi:10.1016/S0733-8619(03)00096-3.

39. Schoups AA, Vogels R, Qian N, Orban G. Practising orientation identification improves orientation coding in V1 neurons. Nature. 2001;412(6846):549–553. doi:10.1038/35087601.

40. Ren Z, Zhou J, Yao Z, Wang Z, Yuan N, Xu G, et al. Neuronal basis of perceptual learning in striate cortex. Nature Scientific Reports. 2016;6:24769. doi:10.1038/srep24769.

41. Yang T, Maunsell JHR. The Effect of Perceptual Learning on Neuronal Responses in Monkey Visual Area V4. Journal of Neuroscience. 2004;24(7):1617–1626. doi:10.1523/JNEUROSCI.4442-03.2004.

42. Zohary E, Celebrini S, Britten KH, Newsome WT. Neuronal Plasticity that Underlies Improvement in Perceptual Performance. Science. 1994;263(5151):1289–1292.

43. Jehee JFM, Ling S, Swisher JD, van Bergen RS, Tong F. Perceptual Learning Selectively Refines Orientation Representations in Early Visual Cortex. Journal of Neuroscience. 2012;32(47):16747–16753. doi:10.1523/JNEUROSCI.6112-11.2012.

44. Shibata K, Watanabe T, Sasaki Y, Kawato M. Perceptual learning incepted by decoded fMRI neurofeedback without stimulus presentation. Science (New York, NY). 2011;334(6061):1413–5. doi:10.1126/science.1212003.

45. Furmanski CS, Schluppeck D, E, Ngel SA. Learning strengthens the response of primary visual cortex to simple patterns. Current Biology. 2014;14(7):573–578. doi:10.1016/j.

46. Gu Y, Liu S, Fetsch CR, Yang Y, Fok S, Sunkara A, et al. Perceptual learning reduces interneuronal correlations in macaque visual cortex. Neuron. 2011;71(4):750–761. doi:10.1016/j.neuron.2011.06.015.

47. Hung SCSC, Seitz AR. Prolonged Training at Threshold Promotes Robust Retinotopic Specificity in Perceptual Learning. Journal of Neuroscience. 2014;34(25):8423–8431. doi:10.1523/JNEUROSCI.0745-14.2014.

48. Xiao LQ, Zhang JY, Wang R, Klein SA, Levi DM, Yu C. Complete Transfer of Perceptual Learning across Retinal Locations Enabled by Double Training. Current Biology. 2008;18(24):1922–1926. doi:10.1016/j.cub.2008.10.030.

49. Zhang J, Zhang G, Xiao L, Klein SA, Levi DM, Yu C. Rule-Based Learning Explains Visual Perceptual Learning and Its Specificity and Transfer. Journal of Neuroscience. 2010;30(37):12323–12328. doi:10.1523/JNEUROSCI.0704-10.2010.

50. Watanabe T, Sasaki Y. Perceptual learning: Toward a comprehensive theory. Annual review of psychology. 2015;3(66):197–221. doi:10.1146/annurev-psych-010814-015214.Perceptual.

51. Wang R, Zhang JYJY, Klein SA, Levi DM, Yu C. Vernier perceptual learning transfers to completely untrained retinal locations after double training: a "piggybacking" effect. Journal of vision. 2014;14(13):12. doi:10.1167/14.13.12.

52. Cong Lj, Wang Rj, Yu C, Zhang JY. Perceptual learning of basic visual features remains task specific with Training-Plus-Exposure (TPE) training. Journal of Vision. 2016;16(3:13):1–9. doi:10.1167/16.3.13.

53. Talluri BC, Hung SC, Seitz AR, Seriès P. Confidence-based integrated reweighting model of task-difficulty explains location-based specificity in perceptual learning. Journal of vision. 2015;15(10):17. doi:10.1167/15.10.17.

54. Hochstein S, Ahissar M. Hierarchies and reverse hierarchies in the Visual System. Neuron. 2002;36(3):791–804. doi:10.1016/S0896-6273(02)01091-7.

55. Ahissar M, Hochstein S. The reverse hierarchy theory of visual perceptual learning. Trends in Cognitive Sciences. 2004;8(10):457–464. doi:10.1016/j.tics.2004.08.011.

56. Ostwald D, Lam JM, Li S, Kourtzi Z. Neural Coding of Global Form in the Human Visual Cortex. Journal of Neurophysiology. 2008;99(5):2456–2469. doi:10.1152/jn.01307.2007.

57. Aaen-Stockdale C, Hess RF. The amblyopic deficit for global motion is spatial scale invariant. Vision Research. 2008;48(19):1965–1971. doi:10.1016/j.visres.2008.06.012.

58. Levi A, Shaked D, Tadin D, Huxlin KR. Is improved contrast sensitivity a natural consequence of visual training? Journal of Vision. 2014;14(10):1158–1158. doi:10.1167/14.10.1158.

59. Ball K, Sekuler R. Direction-specific improvement in motion discrimination. Vision Research. 1987;27(6):953–965. doi:10.1016/0042-6989(87)90011-3.

60. Newsome WT, Park B, York N, Brook S. A Selective Impairment of Motion Perception Following Lesions of the Middle Temporal Visual Area (MT). The Journal of Neuroscience. 1988;8(6).

61. Vaina LM, Belliveau JW, Roziers EBd, Zeffiro TA. Neural systems underlying learning and representation of global motion. Proceedings of the National Academy of Sciences. 1998;95(21):12657–12662. doi:10.1073/pnas.95.21.12657.

62. Lennie P. Single units and visual cortical organization. Perception. 1998;27:889–935. doi:10.1068/p270889.

63. Livingstone M, Hubel DH. Segregation of Form, Color, Movement, and Depth: Anatomy, Physiology, and Perception. Science. 1988;240(4835):740–749.

64. Furlan M, Smith AT. Global Motion Processing in Human Visual Cortical Areas V2 and V3. Journal of Neuroscience. 2016;36(27):7314–7324. doi:10.1523/JNEUROSCI.0025-16.2016.

65. Lamme VAF. Recurrent corticocortical interactions in neural disease. Archives of Neurology. 2003;60(2):178–184. doi:10.1001/archneur.60.2.178.

66. Simoncelli EP, Heeger DJ. A Model of Neuronal Responses in Visual Area MT. Vision Research. 1998;38(5):743–761. doi:S0042698997001831 [pii].

67. DeValois RL, Albrecht DG, Thonrell LG. Spatial Frequency Selectivity of Cells in Macaque Visual Cortex. Vision research. 1982;22(5):545–559. doi:10.1016/0042-6989(82)90113-4.

68. Hubel DH, Wiesel TN. Receptive fields, binocular interaction and functional architecture in the cat's visual cortex. Journal of Physiology. 1962;160:106–154. doi:10.1523/JNEUROSCI.1991-09.2009.

69. Movshon JA, Thompson ID, Tolhurst DJ. Spatial and temporal contrast sensitivity of neurones in areas 17 and 18 of the cat visual cortex. J Physiol. 1978;283:101–120.

70. Movshon JA, Newsome WT. Visual response properties of striate cortical neurons projecting to area MT in macaque monkeys. Journal of Neuroscience. 1996;16(23):7733–7741. doi:10.1167/9.8.752.

71. Felleman DJ, Van Essen DC. Receptive field properties of neurons in area V3 of macaque monkey extrastriate cortex. Journal of neurophysiology. 1987;57(4):889–920.

72. Hubel DH, Wiesel TN. Receptive fields and functional acrhitecture in two nonstriate visual areas (18 and 19) of the cat. Journal of neurophysiology. 1965;28(2):229–289.

73. Mikami A, Newsome WT, Wurtz RH. Motion selectivity in macaque visual cortex. II. Spatiotemporal range of directional interactions in MT and V1. Journal of neurophysiology. 1986;55(6):1328–39.

74. Sillito AM, Cudeiro J, Jones HE. Always returning: feedback and sensory processing in visual cortex and thalamus. Trends in Neurosciences. 2006;29(6):307–316. doi:10.1016/j.tins.2006.05.001.

75. Zeki SM. Functional organization of a visual area in the posterior bank of the superior temporal sulcus of the rhesus monkey. The Journal of Physiology. 1974;236(3):549–573. doi:10.1113/jphysiol.1974.sp010452.

76. Amano K, Edwards M, Badcock DR, Nishida S. Spatial-frequency tuning in the pooling of one- and two-dimensional motion signals. Vision Research. 2009;49(23):2862–2869. doi:10.1016/j.visres.2009.08.026.

77. Bex PJ, Dakin SC. Comparison of the spatial-frequency selectivity of local and global motion detectors. Journal of the Optical Society of America A, Optics, image science, and vision. 2002;19(4):670–7. doi:10.1364/JOSAA.19.000670.

78. Burr D, Thompson P. Motion psychophysics: 1985-2010. Vision Research. 2011;51(13):1431–1456. doi:10.1016/j.visres.2011.02.008.

79. Nishida S. Advancement of motion psychophysics : Review 2001 – 2010. Science. 2011;11(2011):1–53. doi:10.1167/11.5.11.Introduction.

80. Garcia A, Kuai SG, Kourtzi Z. Differences in the time course of learning for hard compared to easy training. Frontiers in Psychology. 2013;4(MAR):1–8. doi:10.3389/fpsyg.2013.00110.

81. Braddick OJ. Segmentation versus integration in visual motion processing. Trends in Neurosciences. 1993;16(7):263–268. doi:10.1016/0166-2236(93)90179-P.

82. Adelson EH, Movshon JA. Phenomenal coherence of moving visual patterns. Nature. 1982;300(5892):523–525.

83. Wilson HR, Ferrera VP, Yo C. A Psychophysically Motivated Model for 2-Dimensional Motion Perception. Visual Neuroscience. 1992;9(1):79–97.

84. Lui LL, Bourne JA, Rosa MGP. Spatial and temporal frequency selectivity of neurons in the middle temporal visual area of new world monkeys (Callithrix jacchus). European Journal of Neuroscience. 2007;25(6):1780–1792. doi:10.1111/j.1460-9568.2007.05453.x.

85. Pasternak T, Merigan WH. Motion Perception following Lesions of the Superior Temporal Sulcus in the Monkey. Cerebral Cortex. 1994;May/June(4):247–259.

86. Rockland KS, Knutson T. Feedback Connections From Area Mt Of The Squirrel-Monkey To Areas V1 And V2. Journal Of Comparative Neurology. 2000;425(3):345–368.

87. Vaina LM, Sundareswaran V, Harris JG. Learning to ignore: psychophysics and computational modeling of fast learning of direction in noisy motion stimuli. Cognitive Brain Research. 1995;2(3):155–163. doi:10.1016/0926-6410(95)90004-7.

88. Dakin SC, Mareschal I, Bex PJ. Local and global limitations on direction integration assessed using equivalent noise analysis. Vision Research. 2005;45(24):3027–3049. doi:10.1016/j.visres.2005.07.037.

89. Tibber MS, Kelly MG, Jansari A, Dakin SC, Shepherd AJ. An inability to exclude visual noise in migraine. Investigative Ophthalmology and Visual Science. 2014;55(4):2539–2546. doi:10.1167/iovs.14-13877.

90. Williams DW, Sekuler R. Coherent global motion percepts from stochastic local motions. Vision Research. 1984;24(1):55–62. doi:10.1016/0042-6989(84)90144-5.

91. McGovern DP, Webb BS, Peirce JW. Transfer of perceptual learning between different visual tasks. Journal of Vision. 2012;12(11):4–4. doi:10.1167/12.11.4.

92. Britten KH, Shadlen MN, Newsome WT, Movshon JA. The analysis of visual motion: a comparison of neuronal and psychophysical performance. The Journal of Neuroscience. 1992;12(12):4745–4765. doi:10.1.1.123.9899.

93. Newsome WT, Britten KH, Movshon JA. Neuronal correlates of a perceptual decision. Nature. 1989;341(6237):52–54. doi:10.1038/341052a0.

94. Rudolph K, Pasternak T. Transient and permanent deficits in motion perception after lesions of cortical areas MT and MST in the macaque monkey. Cerebral Cortex. 1999;9(1):90–100. doi:10.1093/cercor/9.1.90.

95. Henriksson L, Nurminen L, Hyvarinen A, Vanni S, Hyvarinen A, Vanni S, et al. Spatial frequency tuning in human retinotopic visual areas. Journal of Vision. 2008;8(2008):1–13. doi:10.1167/8.10.5.Introduction.

96. Li RW, Tran TT, Craven AP, Leung TW, Chat SW, Levi DM. Sharpening coarse-to-fine stereo vision by perceptual learning: asymmetric transfer across the spatial frequency spectrum. Royal Society open science. 2016;3(1):150523. doi:10.1098/rsos.150523.

97. Parker AJ. Binocular depth perception and the cerebral cortex. Nature Reviews Neuroscience. 2007;8(5):379–91. doi:10.1038/nrn2131.

98. Fleet DJ, Wagner H, Heeger DJ. Neural Encoding of Binocular Disparity: Energy Models, Position Shifts and Phase Shifts. Vision Research. 1996;36(12):1839–1857.

99. Hibbard PB, Goutcher R, Hunter DW. Encoding and estimation of first- and second-order binocular disparity in natural images. Vision Research. 2016;120:108–120. doi:10.1016/j.visres.2015.10.016.

100. Smallman HS, MacLeod DIA. Spatial scale interactions in stereo sensitivity and the neural representation of binocular disparity. Perception. 1997;26(8):977–994. doi:10.1068/p260977.

101. Tanaka H, Ohzawa I. Neural Basis for Stereopsis from Second-Order Contrast Cues. Journal of Neuroscience. 2006;26(16):4370–4382. doi:10.1523/JNEUROSCI.4379-05.2006.

102. Law C, Gold JI. Neural correlates of perceptual learning in a sensory-motor, but not a sensory, cortical area. Nature Neuroscience. 2008;11(4):505–513. doi:10.1038/nn2070.

103. Hua T, Bao P, Huang CB, Wang Z, Xu J, Zhou Y, et al. Perceptual Learning Improves Contrast Sensitivity of V1 Neurons in Cats. Current Biology. 2010;20(10):887–894. doi:10.1016/j.cub.2010.03.066.

104. Yu Q, Zhang P, Qiu J, Fang F. Perceptual Learning of Contrast Detection in the Human Lateral Geniculate Nucleus. Current Biology. 2016;26(23):3176–3182. doi:10.1016/j.cub.2016.09.034.

105. Lin Z, Dosher BA, Lu ZL. Mixture of easy trials enables transient and sustained perceptual improvements through priming and perceptual learning. Scientific Reports. 2017;7(1):7421. doi:10.1038/s41598-017-06989-0.

106. Liu J, Lu ZL, Dosher BA. Mixed training at high and low accuracy levels leads to perceptual learning without feedback. Vision Research. 2012;61:15–24. doi:10.1016/j.visres.2011.12.002.

107. Romei V, Chiappini E, Hibbard PB, Avenanti A. Empowering Reentrant Projections from V5 to V1 Boosts Sensitivity to Motion. Current Biology. 2016;26(16):2155–2160. doi:10.1016/j.cub.2016.06.009.

108. Chiappini E,Silvanto, J, Hibbard PB, Avenanti A, Romei V. Strengthening functionally specific neural pathways with transcranial brain stimulation. Current Biology. 2018;26(28):2R735–R736. doi:10.1016/j.cub.2018.05.083.

109. Fahle M, Poggio T. Perceptual learning. MIT Press; 2002.

110. Dobres J, Watanabe T. Response feedback triggers long-term consolidation of perceptual learning independently of performance gains. Journal of Vision. 2012;12(8):9–9. doi:10.1167/12.8.9.

111. Herzog MH, Fahle M. The Role of Feedback in Learning a Vernier Discrimination Task. Vision Research. 1997;37(15):2133–2141. doi:http://dx.doi.org/10.1016/S0042-6989(97)00043-6.

112. Seitz AR, Nanez JE, Holloway S, Tsushima Y, Watanabe T. Two cases requiring external reinforcement in perceptual learning. Journal of Vision. 2006;6(9):966–973. doi:10.1167/6.9.9.

113. Brainard DH. The psychophysics toolbox. Spatial vision. 1997;10:433–436.

114. Kleiner M, Brainard D, Pelli D, Ingling A, Murray R, Broussard C, et al. What's new in Psychtoolbox-3. Perception. 2007;36(14):1.

115. Pelli DG. The VideoToolbox software for visual psychophysics: Transforming numbers into movies. Spatial vision. 1997;10(4):437–442.

116. Moscatelli A, Mezzetti M, Lacquaniti F. Modeling psychophysical data at the population-level: The generalized linear mixed model. Journal of Vision. 2012;12(11):26–26. doi:10.1167/12.11.26.

117. Agresti A. Categorical Data Analysis. 2nd ed. Wiley-Interscience; 2002.

118. Bolker BM, Brooks ME, Clark CJ, Geange SW, Poulsen JR, Stevens MHH, et al. Generalized linear mixed models: a practical guide for ecology and evolution. Trends in Ecology and Evolution. 2009;24(3):127–135. doi:10.1016/j.tree.2008.10.008.

119. Jeter PE, Dosher BA, Lu ZL. Task precision at transfer determines specificity of perceptual learning. Journal of Vision. 2009;9(3):1–13. doi:10.1167/9.3.1.Introduction.

120. Liu J, Dosher BA, Lu ZL. Modeling trial by trial and block feedback in perceptual learning. Vision Research. 2014;99:46–56. doi:10.1016/j.visres.2014.01.001.

121. Kingdom FAA, Prins N. Psychophysics: a practical introduction; 2010.

122. Zylberberg A, Roelfsema PR, Sigman M. Variance misperception explains illusions of confidence in simple perceptual decisions. Consciousness and Cognition. 2014;27(1):246–253. doi:10.1016/j.concog.2014.05.012.

123. Watanabe T, Náñez JE, Sasaki Y. Perceptual learning without perception. Nature. 2001;413(6858):844–848. doi:10.1038/35101601.

124. Watanabe T, Náñez JE, Koyama S, Mukai I, Liederman J, Sasaki Y. Greater plasticity in lower-level than higher-level visual motion processing in a passive perceptual learning task. Nature neuroscience. 2002;5(10):1003–1009. doi:10.1038/nn915.

125. Gold JI, Law CT, Connolly P, Bennur S. Relationships Between the Threshold and Slope of Psychometric and Neurometric Functions During Perceptual Learning: Implications for Neuronal Pooling. Journal of Neurophysiology. 2010;103(1):140–154. doi:10.1152/jn.00744.2009.

126. Swanson WH, Birch EE. Extracting thresholds from noisy psychophysical data. Perception & Psychophysics. 1992;51(5):409–422. doi:10.3758/BF03211637.

127. Ling S, Carrasco M. Sustained and transient covert attention enhance the signal via different contrast response functions. Vision Research. 2006;46(8-9):1210–1220. doi:10.1016/j.visres.2005.05.008.

128. Herrmann K, Montaser-Kouhsari L, Carrasco M, Heeger DJ. When size matters: Attention affects performance by contrast or response gain. Nature Neuroscience. 2010;13(12):1554–1561. doi:10.1038/nn.2669.

129. Donovan I, Carrasco M. Exogenous attention facilitates location transfer of perceptual learning. Journal of vision. 2015;15(10):1–16. doi:10.1167/15.10.11.doi.

130. Das A, Huxlin KR. New Approaches to Visual Rehabilitation for Cortical Blindness: Outcomes and Putative Mechanisms. The Neuroscientist. 2010;16(4):374–387. doi:10.1177/1073858409356112.

131. Das a, Tadin D, Huxlin KR. Beyond Blindsight: Properties of Visual Relearning in Cortically Blind Fields. Journal of Neuroscience. 2014;34(35):11652–11664. doi:10.1523/JNEUROSCI.1076-14.2014.

132. Hess RF, Mansouri B, Thompson B. A binocular approach to treating amblyopia: Antisuppression therapy. Optometry and Vision Science. 2010;87(9):697–704. doi:10.1097/OPX.0b013e3181ea18e9.

133. Huxlin KR. Perceptual plasticity in damaged adult visual systems. Vision Research. 2008;48(20):2154–2166. doi:10.1016/j.visres.2008.05.022.

134. Levi DM. Prentice award lecture 2011: Removing the brakes on plasticity in the amblyopic brain. Optometry and Vision Science. 2012;89(6):827–838. doi:10.1097/OPX.0b013e318257a187.

135. Melnick MD, Tadin D, Huxlin KR. Relearning to See in Cortical Blindness. The Neuroscientist. 2016;22(2):199–212. doi:10.1177/1073858415621035.

136. Bridge H, Thomas O, Jbabdi S, Cowey A. Changes in connectivity after visual cortical brain damage underlie altered visual function. Brain. 2008;131(6):1433–1444. doi:10.1093/brain/awn063.

137. Markov NT, Vezoli J, Chameau P, Falchier A, Quilodran R, Huissoud C, et al. Anatomy of hierarchy: Feedforward and feedback pathways in macaque visual cortex. Journal of Comparative Neurology. 2014;522(1):225–259. doi:10.1002/cne.23458.

138. Pascual-Leone A, Walsh V. Fast Backprojections from the Motion to the Primary Visual Area Necessary for Visual Awareness. Science. 2001;292(5516):510–512. doi:10.1126/science.1057099.

139. Hupé JM, James AC, Girard P, Lomber SG, Payne BR, Bullier J. Feedback connections act on the early part of the responses in monkey visual cortex. Journal of Neurophysiology. 2001;85(1):134–145. doi:11152714.

140. Romei V, Thut G, Silvanto J. Information-Based Approaches of Noninvasive Transcranial Brain Stimulation. Trends in Neurosciences. 2016;39(11):782–795. doi:10.1016/j.tins.2016.09.001.

141. Jones DK, Simmons A, Williams SCR, Horsfield MA. Non-invasive assessment of axonal fiber connectivity in the human brain via diffusion tensor MRI. Magnetic Resonance in Medicine. 1999;42(1):37–41.

142. Owsley C. Aging and vision. Vision Research. 2011;51(13):1610–1622. doi:10.1016/j.visres.2010.10.020.

143. Yu C, Klein SA, Levi DM. Perceptual learning in contrast discrimination and the (minimal) role of context. Journal of Vision. 2004;4(2004):169–182. doi:10.1167/4.3.4.

144. Adini Y, Wilkonsky A, Haspel R, Tsodyks M, Sagi D. Perceptual learning in contrast discrimination: the effect of contrast uncertainty. Journal of vision. 2004;4(2004):993–1005. doi:10.1167/4.12.2.

145. Kuai SG, Zhang JY, Klein SA, Levi DM, Yu C. The essential role of stimulus temporal patterning in enabling perceptual learning. Nature Neuroscience. 2005;8(11):1497–1499. doi:10.1038/nn1546.

146. Zhang JY, Kuai SG, Xiao LQ, Klein SA, Levi DM, Yu C. Stimulus coding rules for perceptual learning. PLoS Biology. 2008;6(8):1651–1660. doi:10.1371/journal.pbio.0060197.

147. Aberg KC, Herzog MH. Interleaving bisection stimuli - randomly or in sequence - does not disrupt perceptual learning, it just makes it more difficult. Vision Research. 2009;49(21):2591–2598. doi:10.1016/j.visres.2009.07.006.

148. Cong LJ, Zhang JY. Perceptual learning of contrast discrimination under roving: The role of semantic sequence in stimulus tagging. Journal of Vision. 2014;14(13):1–8. doi:10.1167/14.13.1.

149. Lin Z, Murray SO. Priming of awareness or how not to measure visual awareness. Journal of Vision. 2014;14(2014):1–17. doi:10.1167/14.1.27.

150. Lin Z, Murray SO. Automaticity of Unconscious Response Inhibition : Comment on Chiu and Aron (2014). Journal of Experimental Psychology: General. 2015;144(1):244–254.

151. Solgi M, Liu T, Weng J. A computational developmental model for specificity and transfer in perceptual learning. Journal of Vision. 2013;13:7. doi:10.1167/13.1.7.