Spatial frequency tuning of perceptual learning and transfer in global motion

Jordi M Asher [1*], Vincenzo Romei [1,2], Paul B Hibbard [1],

1 University of Essex, Department of Psychology, Wivenhoe Park, United Kingdom, CO4 3SQ
2 Dipartimento di Psicologia and Centro studi e Ricerche in Neuroscienze Cognitive, Campus di Cesena, Università di Bologna 47521 Cesena, Italy

* jashera@essex.ac.uk

Abstract

Perceptual learning is typically highly specific to the stimuli and task used during training. However, recently it has been shown that training on global motion can transfer to untrained tasks, reflecting the generalising properties of mechanisms at this level of processing. We investigated a) if feedback was required for learning when using an equivalent noise global motion coherence task, and b) the transfer across spatial frequency of training on a global motion coherence task, and the transfer of this training to a measure of contrast sensitivity. For our first experiment two groups, with and without feedback, trained for ten days on a broadband global motion coherence task. Results indicated that feedback was a requirement for learning. For the second experiment training consisted of five days of direction discrimination on one of three global motion tasks (broadband, low or high frequency random-dot Gabors), with trial-by-trial auditory feedback. A pre- and post-training assessment was also conducted, consisting of all three types of global motion stimuli (without feedback) and high and low spatial frequency contrast sensitivity. We predicted that if learning and transfer is cortically localised, then transfer would show specificity to the area processing the task (global motion). In this case, we would predict a broad transfer between spatial frequency conditions of global motion only. However, if transfer occurred as a result of backward generalisation, a more selective transfer would occur matching the low-pass broadband tuning of the area processing global motion. Our training paradigm was successful at eliciting improvement in the trained tasks over the five days. However, post-training transfer to trained or untrained tasks was only reported for the low spatial frequency trained group. This group exhibited increased sensitivity to low spatial frequency contrast, and an improvement for the broadband global motion condition. Our findings suggest that the feedback projections from global to local stages of processing play a role in transfer.

Introduction

Perceptual learning has attracted much attention as a potential tool to aid recovery of lost visual function for clinical populations [1]. The success of perceptual training in amblyopia [2–4], presbyopia [4] and cortical damage [5–7] has demonstrated sensory plasticity in adulthood. This evidence contradicts the position that sensory development is restricted to a critical period early in life [8,9] and that the visual system is hard-wired in mature systems [10]. While it has repeatedly been established that training can improve perceptual abilities [11], these benefits tend to be highly specific for both the perceptual features of the stimuli [12–14] and the behavioural task used in training [15]. This specificity severely limits the effectiveness of perceptual learning as a general therapeutic tool. Resolving the conditions under which learning is tied to the features and tasks used in training, and how much it can generalise to new tasks and stimuli, is imperative for understanding the mechanisms of perceptual learning [11,16]. Our study aims to identify the relative location of one of the mechanisms involved when learning direction discrimination for a global motion coherence task. We do this by evaluating the spatial frequency tuning of improvements in performance for trained and untrained tasks. For the purpose of this paper we dissociate between models and mechanisms of perceptual learning using two general categories i) the How and ii) the What and Where. Models of perceptual learning aim to resolve 'how' learning occurs. Understanding which mechanisms underlie this learning address the 'what' and 'where' (and possibly 'why') questions. In this way we consider a mechanism a feature or component of the learning model [17].

Models of Perceptual Learning - How

The 'how' is often the primary focus for research and there are, broadly speaking, two predominant models of perceptual learning. The first explains perceptual learning in terms of a change in the neurons that code for that feature [18,19]. The second position argues that learning is a result of the change in weights of readout between the sensory representation and the decision units [20–24]. The first position developed as a result of accumulating evidence that improvements were highly specific for the features of the stimuli used for training. Early studies identified that learning was specific for orientation [13,25,26], spatial frequency [11,13,27], direction of motion [15], retinal location [14,24,26,28–30] of stimuli and the eye to which they are presented [14,24,29]. On the basis of this feature-specificity it has been argued that the underlying brain area responsible for the learning process must be within the primary visual cortex [14] where the receptive fields of cells display a high degree of specificity. The implication of this position is a potential degree of plasticity in individual neurons within an area previously thought to be incapable of structural change [14,19,31]. Although the failure of perceptual learning to transfer across retinal location, orientation and other stimulus dimensions implicates V1, learning is far more complex than can be explained by the simplicity of the stimulus driven, bottom-up, visual representation

model [32]. A model of learning also needs to include top-down factors such as attention (see [31] for a review), confidence in decision making [33, 34] and knowledge and understanding of the task [35]. The second position proposes that repeated experience alters the connections between lower visual areas and the high level areas where the perceptual decision is then made [36]. Rather than a physical change in the sensory neuron, learning represents a change in the weights of connections between the input, sensory encoding layer and the classification layer at which the decision is made [20–24].

One consistent aspect across models, is that perceptual learning is likely to occur as a result of exploiting correlations between neural responses [19].

Behavioural studies of perceptual learning are unable to distinguish between learning as a result of changes in sensory encoding, or as a result of a reweighting of the connections between sensory and decision-making stages of processing. However, learning effects on the structure of receptive fields of visual neurons have been measured using single cell recordings. Changes in orientation and spatial frequency tuning in V1 [37, 38] and V4 [39], and increased sensitivity to motion coherence in V5 [40], have all been found as a result of perceptual learning. In addition, neuroimaging in human observers has indicated a reorganisation of the visual cortex after training [41–43]. However, this in itself cannot dissociate between learning models [32, 35, 44]. Thus, it is generally judged that physiological models are not sufficient to explain the improvements in task performance exclusively [21, 45].

Mechanisms of Perceptual Learning - What, Where, Why

In order to begin to unravel the 'what' , 'where' and 'why' questions, the behaviours of the system in question need to be understood, and the sensory output mapped to the behavioural output [17]. In this section we will consider some of the behaviours associated with the learning process.

Specificity

A hallmark behaviour of perceptual learning is its specificity to the task, and features, of training. Early psychophysical research investigated learning and transfer using lower-level visual tasks such as texture discrimination [14] and motion discrimination [46] that match the receptive fields of neurons in the lower visual processing areas. At the lowest level, V1 and V2 represent simple visual dimensions, such as the position, orientation and scale of local image features [12]. The receptive fields of cells in V1 are small and only respond to a very restricted area of the visual field [47–49]. Furthermore, V1 neurons are sharply tuned to orientation [50, 51], spatial frequency [50, 52] and direction [53, 54]. For example contrast detection, which is a low-level visual task, is proposed to depend on processing in V1. The pathway for contrast detection and discrimination is believed to develop during childhood and adolescence, reaching its peak in adulthood, followed by a decline in late adulthood [55]. Early psychophysical studies found little perceptual learning for contrast-dependent stimuli [18]. More recently, learning has been found to occur

as long as 'enough training' is provided [56]. The improvement in contrast $\hspace{2em}$ 88
sensitivity has been found to be highly tuned to spatial frequency, specific for $\hspace{1em}$ 89
the trained eye and retinal location of training, but not selective for $\hspace{3em}$ 90
orientation [27, 56–61]. Conversely, the receptive fields in higher cortical areas, $\hspace{0.5em}$ 91
such as those found in V4 and V5, are larger than those of V1 and are less $\hspace{1em}$ 92
dependent on location, retinal size, viewpoint, and lighting [47, 52, 62–66]. $\hspace{2em}$ 93

Transfer $\hspace{40em}$ 94

More recently, a number of studies have demonstrated transfer following training $\hspace{0.5em}$ 95
using a double-training protocol [59, 61, 67]. Double-training involves overt $\hspace{2em}$ 96
training on a task-relevant feature in addition to exposure to a task-irrelevant $\hspace{1em}$ 97
feature [68]. Contrast sensitivity, historically specific to the retinal location of of $\hspace{0.2em}$ 98
training was found to transfer between retinal locations when observers were $\hspace{1em}$ 99
trained on two simultaneous tasks [59]. Observers were required to detect a $\hspace{2em}$ 100
relevant feature (contrast) at one retinal location, followed by detection of an $\hspace{1em}$ 101
irrelevant feature (orientation) at another location, and a full transfer of contrast $\hspace{0.2em}$ 102
sensitivity to the untrained location was found [59]. Similarly, observers who $\hspace{1.5em}$ 103
were trained to discriminate orientation and simultaneously exposed to a second $\hspace{0.5em}$ 104
orientation experienced a full transfer of improvement in orientation sensitivity $\hspace{0.5em}$ 105
as long as training preceded exposure, rather than being simultaneously $\hspace{2em}$ 106
presented [61]. Zhang et al. (2010) [61] proposed a rule-based learning model to $\hspace{0.5em}$ 107
account for the transfer, where higher level decision units learn and re-weight $\hspace{1em}$ 108
the V1 inputs. They proposed that the absence of functional connections to the $\hspace{0.5em}$ 109
untrained orientation or location prevents any potential reweighting. Double $\hspace{1.5em}$ 110
training, through exposure, activates the functional connections at the new $\hspace{1.5em}$ 111
locations or orientations, enabling transfer. This suggests that there may not be $\hspace{0.5em}$ 112
a straightforward correspondence between stimulus features and the neural loci $\hspace{0.5em}$ 113
involved in performing the task. Double-training has been a successful paradigm $\hspace{0.3em}$ 114
in obtaining evidence of transfer [59, 69, 70], however task difficulty during $\hspace{1.5em}$ 115
training, and observer confidence still play a vital role in the learning and $\hspace{1.5em}$ 116
transfer process [34, 71]. An early theoretical model of transfer is the Reverse $\hspace{1em}$ 117
Hierarchy Theory [72]. This model proposes that transfer is a top-down process, $\hspace{0.3em}$ 118
and the degree of transfer is dependent on the characteristics of the receptive $\hspace{1em}$ 119
fields involved in performing the training task. Transfer occurs as a result of $\hspace{1.5em}$ 120
modification of neurons found in the higher cortical levels, where receptive fields $\hspace{0.2em}$ 121
generalise. In contrast, specificity, occurs at the lower cortical levels where the $\hspace{0.5em}$ 122
receptive fields are localised [72]. As established, learning is highly specific to $\hspace{1.5em}$ 123
the features of the stimulus and reflects the tuning properties of the relevant $\hspace{1em}$ 124
receptive fields. However, what specificity is expected for tasks that are not $\hspace{2em}$ 125
processed in early visual areas? For example, perceptual learning improves $\hspace{2em}$ 126
detection and discrimination in tasks using global form [73] and global $\hspace{3em}$ 127
motion [46, 74, 75]. These kinds of tasks are known to be processed at higher $\hspace{1.5em}$ 128
levels of the visual cortex, for example areas V3 [76], V4 [73] and V5 [77, 78]. $\hspace{1em}$ 129
The Reverse Hierarchy Theory predicts that the feedback connections from $\hspace{2em}$ 130
higher levels back to early visual processing areas may be involved in facilitating $\hspace{0.2em}$ 131
learning that transfers, and that the key to understanding specificity and $\hspace{2em}$ 132

transfer lies in the hierarchy of processing and the feedforward and feedback ₁₃₃
connections between them [33, 72]. Based on this, tasks that require higher level ₁₃₄
integration or segregation (global processing) may therefore play an important ₁₃₅
role in creating learning that generalises to other tasks and stimuli [33]. ₁₃₆

Local vs Global Processing ₁₃₇

The receptive fields of neurons in higher cortical areas integrate information to ₁₃₈
represent global stimulus properties [19, 79–82]. Global processes are investigated ₁₃₉
using stimuli or tasks that can only be resolved through integration and ₁₄₀
segregation of coherent or conflicting information [81–83]. Cells higher in the ₁₄₁
processing cortical hierarchy are involved in the perception of global aspects of ₁₄₂
an image and generalise across individual features such as spatial frequency and ₁₄₃
location. Based on the predictions of the Reverse Hierarchy Theory, modification ₁₄₄
of these generalising receptive fields may produce perceptual learning that also ₁₄₅
generalises over these stimulus parameters. Since perceptual learning occurs for ₁₄₆
both global motion [46, 75] and form tasks [83, 84], this suggests that learning is ₁₄₇
not restricted to the initial encoding of information in V1, and can occur at ₁₄₈
higher levels of cortical processing. Learning that involves higher level global ₁₄₉
aspects of perception is of particular interest since it has the potential to ₁₅₀
produce more generalisable improvements [33]. Huxlin et al. (2009) [5] ₁₅₁
suggested that extensive training using global motion may result in general ₁₅₂
improvements in visual sensitivity. They postulate that the improvements may ₁₅₃
be a result of the feedback connections from higher brain areas back to V1, ₁₅₄
where training may stimulate and reactivate "intact islands" of activity. ₁₅₅

Testing models of perceptual learning of global motion ₁₅₆

Paradigms to evaluate the transfer of learning based on features of the task may ₁₅₇
shed light on the location of the learning mechanisms involved. In order to ₁₅₈
behaviourally disassociate the location of a learning mechanism, Petrov et ₁₅₉
al., [36, 44] suggest using perceptual learning experiments in which the sensory ₁₆₀
representations required for two tasks are the same, but the decision stages ₁₆₁
differ, or where both sensory representations and the decision stage are common ₁₆₂
to both. Training using a global motion task has been found to transfer between ₁₆₃
eyes [15, 46]; increase sensitivity for detection and discrimination [5]; help ₁₆₄
recover some of the blind field for cortically blind subjects [5]; and reduce ₁₆₅
contrast thresholds for drifting stimuli [75]. Transfer of learning has also been ₁₆₆
reported in a control population [75], where a post-training improvement was ₁₆₇
found for contrast sensitivity. This is a particularly interesting result when ₁₆₈
taking into account that a globally processed trained task improved contrast ₁₆₉
detection, which is known to be processed in an early visual locations such as ₁₇₀
$4C\alpha$, where cells are tuned for spatial frequency but not orientation [27]. The ₁₇₁
success in obtaining improvements in untrained tasks/ features makes global ₁₇₂
motion an interesting task to investigate this further. ₁₇₃

Global Motion Processing

The perception of motion is hypothesised to occur as a two stage process [80,85]. At the first stage, spatial frequency- and orientation-tuned mechanisms in V1 encode the motion signals that occur locally within the receptive fields of individual neurons [54,81]. However, local and global stages of motion processing differ in their spatial frequency tuning. In order to process more complex motion, ambiguous or conflicting signals from the first stage need to be integrated over a wider spatial area to provide a global representation of motion and velocity [86,87].

A number of areas within the visual cortical hierarchy play a functional role in processing motion. Areas V2 and V4 have a role in processing moving orientation signals [88,89]. V3A also plays a role in several aspects of motion processing [62], with 76% of neurons being selective for orientation and 40% showing strong direction selectivity. However, evidence from lesion studies [90,91], extra-cellular recordings [77,92] and neuroimaging in humans [93] support area V5 as a brain area that is heavily involved in processing global motion [77,85,90–92,94]. Most neurons in V5 are strongly direction selective [93,95], and the evidence for the role it plays in spatially integrating motion signals is well supported by non-human primate data, and neuroimaging studies in humans [47,77,96,97]. Receptive fields in V5 can be up to tenfold larger than those in V1 [65], with broad spatial frequency and orientation tuning, allowing them to sum the responses of V1 neurons, across space, orientation and spatial and temporal frequency [49].

Spatial frequency tuning of global motion processing

Psychophysical studies have shown that global motion detectors have relatively broad spatial frequency tuning [80]. This is consistent with single-cell recordings from area V5 in marmoset monkeys that exhibit bandpass spatial and temporal frequency tuning, with a preference for low spatial frequencies [96]. Amano et al. (2009) [79] suggested that, within V5, there is a "motion-pooling mechanism" with broadband, low-pass tuning. Pooling of visual sensory information has also been found in studies investigating transfer in binocular stereopsis [98]. The neural mechanisms of stereoscopic vision are also known to be processed on multiple levels of the visual hierarchy [99], and the pooling of information across spatial frequency mechanisms has been proposed as an important step in estimating depth [100–103]. In the same way, pooling across low frequencies is likely an important step for integration and segregation of coherent motion. Furthermore, a recent fMRI study [104] reported that areas V5+ and V3A both exhibit an attenuated response for high spatial frequencies.

Global motion and the reverse hierarchy in practice

The Reverse Hierarchy Theory predicts that an increase in sensitivity at lower cortical areas, as a result of the feedback connections from higher cortical areas, would be dependent on the frequency tuning of the global motion detectors [72].

Feedback connections are argued to be fundamental to efficient cortical organisation, and of all the feedback systems terminating in V1, the connections from V5 to V1 have been found to cover the most territory [105]. Furthermore, these feedback (or re-entrant) connections terminate in different combinations within various layers throughout V1 [65, 105] and are rapidly updated [106, 107]. Recordings from macaque monkeys have suggested that there is almost no delay for information processed in V5 to be fed back to lower areas, and it has been proposed that the feedback from V5 is present prior to the bottom up information from the feedforward connections [106, 107]. This has recently been supported by evidence of an independent motion pathway in humans, finding a direct link from lateral geniculate nucleus (LGN) to V5 [108]. Projections are targeted retinotopically to V1 in locations lying within the V5 receptive field [65]. Thus, as V5 responses are influenced by the bottom-up responses from V1, the top-down responses from V5 shape the responses of the V1 cells which provide the sensory input [65]. Sillito et al. (2006) [65] suggest that the "iterative interaction" between the two stages could account for the selectivity at both levels. Backward projections are a theoretically plausible route for perceptual learning, and are consistent with the predictions from the Reverse Hierarchy Theory. Overall this suggests a theoretically important role for top-down information in obtaining transfer from global motion [105].

Transcranial Magnetic Stimulation (TMS) has been used to stimulate the re-entrant connections from V5 to V1, enhancing the perception of global motion [109]. Using a novel paired cortico-cortical TMS protocol (ccPAS) to induce Hebbian plasticity, observers' thresholds for motion detection were reduced when the feedback connections from V5 to V1 were stimulated. However, the improvement in perception was critically dependent on the timing and direction of stimulation. There was no change when the feedforward connections from V1 to V5 were stimulated [110]. This also suggests that these re-entrant connections are malleable [110]. Using a similar method, a direction-selective improvement was induced by pairing subthreshold stimulation with the simultaneous presentation of direction-specific moving stimuli [111]. This provides additional support for the accumulating evidence that the re-entrant connections from direction-tuned neurons in areas such as V5 play a role in perceptual learning in global motion coherence tasks.

Our Study

Given the differences in spatial and temporal frequency tuning of processing between the local and global levels [50, 77, 96], measuring the tuning of training for these dimensions allows us to understand the role of each level in global motion learning.

Prior to collecting the data for our main study, we also questioned whether trial-by-trial feedback was a requirement for learning. Therefore, we first investigated the necessity of feedback for perceptual learning to occur for our specific stimuli.

Experiment 1

The specific nature of feedback and its role in perceptual learning is unclear, and external feedback has been shown to improve learning and increase efficiency [112,113]. However, some studies have found that learning occurs without external feedback [11,15,30,36,78,114]. Recently, it has been found that interleaving high accuracy (easy) trials and low accuracy (difficult) trials resulted in perceptual learning without the need for feedback, even on difficult trials [114]. Based on these results we predicted that we should find learning in both conditions, as long as easy and difficult trials were interleaved. As detailed in the following sections, our study found that learning did not occur in the condition where no feedback was provided, even with easy trials. Learning only occurred for the feedback condition. With this in mind our design for experiment 2 included feedback during training, but no feedback when testing.

Experiment 2 - Main Study

The purpose of this study was to extend the pre- and post-training measures captured by Levi et al. (2015) [75]. Thus, as well as testing for transfer to contrast sensitivity, we included measures of transfer to the trained and untrained global motion frequencies.

Transfer from global motion could involve a combination of reweighting that includes the global motion mechanisms in higher areas such as V5, and the re-entrant connections from global mechanisms to local mechanisms. Differences in the spatial frequency tuning at the local and global stage mean that these models make different predictions for the transfer of learning across spatial frequency, and different psychophysical tasks (Fig. 1).

Predictions

Cortically localised global reweighting: If transfer relies on global motion mechanisms

In this case learning, and transfer of learning, for global motion would reflect the broad, approximately low-pass spatial frequency tuning of global motion detectors in areas such as V5 and V3/V3A. There would thus be most transfer when the training and test stimuli contain low spatial frequency components (robust transfer is predicted from low frequency to low frequency conditions; modest transfer from low to broad frequency stimuli, and from broad to low and broad stimuli, and an unlikely but possible transfer from broad to high and high to broad.). No transfer to contrast sensitivity would be predicted, since learning would result in changes in weightings between direction-tuned global motion mechanisms, and higher-level decision stages.

Backward generalisation: If transfer relies on feedback projections from global to local stages of processing

If transfer occurs broadly across tasks and levels in a top down manner, as predicted by the Reverse Hierarchy Theory [33], we would predict transfer to be restricted by the frequency tuning of the global motion detectors, but to transfer more broadly across tasks. We would expect most transfer when both training and test stimuli contained low spatial frequency components (robust transfer from low to low frequency stimuli; modest transfer from low to broad, and broad to low and broad to broad, with an unlikely but possible transfer from high to broad, and broad to high). We would also expect transfer to contrast sensitivity, and for this to show the same low-pass spatial frequency tuning. This would also be consistent with the transfer found by Levi et al. (2015) [75].

Finally, a mid frequency orientation discrimination condition was included as a control task. Since none of the motion training stimuli contain any orientation information, we predicted no transfer to occur from any trained condition or either level of processing.

Methods and Materials:

Observers

24 observers for Experiment 1 and 30 (new) observers for Experiment 2 were randomly and evenly assigned into groups. For experiment 1 there was a feedback and a no-feedback group. For experiment 2 groups were categorised by the spatial frequency of training (broad, low, high). All observers were employees or students from the University of Essex and self declared as having normal or corrected-to-normal vision. All work was carried out in accordance with the Code of Ethics of the World Medical Association (Declaration of Helsinki). The study procedures were approved by the University of Essex Ethics Committee (JA1601). All observers gave informed written consent and were either paid or received course credit for their participation.

Experiment 1:

Apparatus

Stimuli were generated and presented with Matlab 2015a using the Psychophysics Toolbox extensions [115–117]. The broadband global motion stimuli were presented using a 27" 2.7 Ghz iMac running OSX 1.9.5 iMac with a display resolution of 2560 x 1440 pixels and a 60 Hz refresh rate. Viewing distance was 450mm, the stimuli subtended a visual angle of 66.8°, and one pixel subtended 1.77 arc minutes.

Stimuli and Procedure

Global motion stimuli contained 100 Gaussian elements, each with a standard deviation of 6.8 arc minutes (see Fig. 2a). Elements were presented within a

Fig 1. (a) **Expected improvement from training:** We would expect robust improvement on frequency specific motion stimuli over the 5 days training, and for this to be reflected at the post-assessment as an improvement compared to the pre-assessment results.

Predictions of transfer (b) **Cortically localised global reweighting:** In this case we may predict that transfer from frequency specific global motion would reflect the broad, approximately low-pass spatial frequency tuning of global motion detectors in areas such as V5 and V3/V3A. There would thus be most transfer when both training and test stimuli contained low spatial frequency components

(c) **Backward generalisation:** Should transfer occur as a result of the backward projections from V5 to V1 (as predicted by the Reverse Hierarchy Theory [33]) we would therefore also expect robust transfer from low to low frequency stimuli; modest transfer from low to broad, and broad to low and broad to broad, with an unlikely but possible transfer from high to broad, and broad to high. Importantly, should there be transfer to contrast sensitivity we would expect this to reflect the low-pass spatial frequency tuning of global areas.

mid-grey rectangle measuring 17.6° x 17.6° on a mid-grey background. Elements moved 5 pixels per frame, and each element moved a fixed distance of 8.8°. Dots wrapped around the rectangle when approaching the edges.

Global motion stimuli were based on the task designed by Williams and Sekuler (1984) [118]; dots moved around the screen at 7 levels of coherence (180°; 300°; 330°; 340°; 345°; 350°; 355°). Lower numbers depict higher

Fig 2. Global motion stimuli, at broad (a), low (b) and high spatial frequencies (c). For all stimuli the spatial frequency and speed of motion were held constant.

coherence, and higher numbers depict lower coherence and thus more randomness. At each level of coherence each dot moved a fixed distance within the defined arc of degree of coherence, while moving in a leftwards or rightwards direction (see Fig. 3). All tasks were presented as two-alternative-forced-choice (2AFC) decisions (left or right response), using the method of constant stimuli (MOCS). Observers responded by pressing the left or right arrow key associated with perceived leftwards or rightwards motion. When feedback was present, this was provided as an immediate auditory beep after each trial, a high pitched tone for a correct response (2000Hz for 10ms), and a low pitched tone for an incorrect response (200Hz for 40ms). Each day 40 repetitions of each level of coherence was presented for a total of 280 trials.

Experiment 2:

Design and Apparatus

A series of baseline measures were completed for global motion direction discrimination, contrast sensitivity and orientation discrimination (pre-and post-training assessments). Assessment stimuli were presented on a VIEWPixx/3D 23.6 inch monitor with a display resolution of 1920 x 1080 pixels, with a 120 Hz refresh rate, using a Dell Precision T3610 PC running Windows 7. One pixel subtended 1.6 arc minutes and stimuli were viewed from a distance of 570mm. Head position for testing was stabilised using a chin rest. Following this, all observers undertook five consecutive days of global motion training in one of three spatial frequency groups (broad, high or low). Training stimuli were presented on a 19" monitor with a display resolution of 1980 x 1080 pixels and 60 Hz refresh rate, using a PC running Windows 7. One pixel subtended 1.7 arc minutes. Stimuli were viewed from a distance of 500mm. All stimuli were presented for 1 second.

Training Stimuli

Global Motion: Broadband stimuli were the same as previously described. For low-frequency stimuli, the elements were circularly symmetric Gabor patches. The standard deviation of the Gaussian window, σ, was 30.1 arc minutes, and

Fig 3. Equivalent noise global motion stimuli a) The random walk that creates the motion of dots with a 90° arc (for descriptive purposes only). The arc defines the potential area of movement each dot could take at that level. The higher the degree of arc, the lower the coherence. The arrow indicates the actual trajectory and motion of the dot on each step. b) Potential trajectories for the random walk for levels with 180°, 270° 255°and 360° arc. c) An schematic example of an equivalent noise global motion stimulus. All dots contribute equally to the signal and the noise (drawn from a distribution between 0° and 360 °). Each dot moves independently within its arc, however in general the dots are moving rightwards.

the spatial frequency of luminance modulation, f, was 1 cycle/degree. For each element, the luminance profile was defined as a function of horizontal and vertical position (x, y) as:

$$d(x, y) = \sqrt{(x - x_0)^2 + (y - y_0)^2} \tag{1}$$

$$L(d) = \frac{A}{\sigma\sqrt{2\pi}} \exp \frac{-d^2}{2\sigma^2} \cos 2\pi f d \tag{2}$$

where x_0 and y_0 is the central position of the element, and A determines the contrast. Elements for the high frequency stimuli were defined in the same way, but had a standard deviation of 7.48 arc minutes and a spatial frequency, f, of 4 cycles/degree. For all stimuli, the spatial frequency and the speed of motion were held constant. Initially all elements were uniformly and randomly distributed within a region of 16.6°x 16.6° on the centre of the screen. A central black fixation dot was presented at all times when stimuli were not being displayed. Examples of the stimuli are shown in see Fig. 2 a-c. Motion was created using the method detailed in Fig. 3. On each frame each dot moved a fixed distance of 8.5 arc minutes.

Pre- and Post-training Stimuli

1. *Global motion:* Stimuli were identical to those described in the training session, with the following exceptions. The standard deviations of the

elements were 6.4 arc minutes (broadband), 28.4 arc minutes (low   387
frequency) and 7.0 arc minutes (high frequency). Stimuli were presented   388
within a mid-grey rectangle measuring 15.9°x 15.9°, each element moved a   389
fixed distance of 8 arc minutes.   390

2. *Contrast Sensitivity:* Stimuli were Gabor patches, with a spatial frequency   391
   of 1 cycle per degree (/°) or 4 cycles/°, presented in the centre of the   392
   screen on a mid-grey background, tilted either ± 20° (see Fig. 4a&b). The   393
   Gaussian envelope of the Gabor stimulus had a standard deviation of 1.1°.   394
   7 levels of contrast (0.05, 0.1, 0.15, 0.175, 0.2, 0.3, 0.4 % Michelson   395
   Contrast) were presented.   396

3. *Orientation Discrimination:* Gabor patches were presented for 100ms on a   397
   mid-grey background, measuring 7.9° x 7.9°, and presented 6.6° either   398
   above or below the central fixation point. Orientation was centred at zero,   399
   at which point lines were vertical. There was a total range of 1.8°   400
   difference in tilt, between ± 0.9°, with 7 linearly spaced points within the   401
   range. The spatial frequency of the Gabor patch was 1.85 cycles/° and the   402
   standard deviation of the Gaussian envelope was 0.2° (see Fig. 4c).   403
   Contrast was fixed at 50% Michelson Contrast.   404

Fig 4. Stimuli: The Gabors used for a) low spatial frequency contrast
sensitivity - 1 cycle per degree, b) high spatial frequency- 4 cycles per degree,
and c) orientation discrimination which is tilted leftwards at 2° with mid spatial
frequency- 2 cycles per degree

Procedure   405

Training on Global Motion   406

For each observer, training was undertaken at one spatial frequency only,   407
totalling 420 trials daily for 5 continuous days. Feedback was provided after   408
each trial.   409

Pre- and Post-Assessments   410

Measures were taken for global motion (high, broad and low spatial frequency),   411
contrast sensitivity (high and low spatial frequency) and orientation   412

discrimination (angle of tilt from the vertical) of an oriented Gabor patch. 413
Responses were captured on the DataPixx response box for contrast sensitivity, 414
and left and right arrows on the keyboard for global motion and orientation 415
discrimination. The presentation order of trials was randomised for direction 416
and coherence (global motion), orientation and contrast (contrast sensitivity) or 417
orientation (orientation discrimination). There were 20 repetitions for each of 418
the seven levels, for each condition. Testing was performed in a darkened room, 419
before and after training. 420

General Statistical Methods 421

Moscatelli et al. (2012) [119] proposed using the Generalised Linear Mixed 422
Effects Model (GLMM) for psychophysical data. The GLMM is an extension of 423
the General Linear Model (GLM) that provides a more robust statistical analysis 424
where the data contain irregular response distributions. The GLMM contains 425
both fixed and random effects. The fixed component estimates are the effects of 426
interest, in our study they are a) the day (or session) of testing and b) each level 427
of the stimulus (coherence, contrast or orientation). The modelling of random 428
effects assesses the differences between related groups of data points (such as 429
those from different observers) that allow inference to a larger population [120]. 430
The linear component of the model is given by: 431

$$\eta = \alpha + \beta X \tag{3}$$

where α and β are the intercept and the slope of the fixed effects parameters 432
respectively. The link function converts the expected outcome variable (the 433
proportion of correct responses, p), to the linear predictor [121]. Here, a logit 434
function was used, adapted to take account of chance performance, such that: 435

$$ln\frac{p - 0.5}{1 - p} = \alpha + \beta X \tag{4}$$

Selecting the model: All response data were analysed using a GLMM 436
conducted using Matlab (using the *fitglme* function), the logit link function and 437
REMPL (restricted maximum pseudo likelihood) to estimate the model 438
parameters. Each model included observers as a random effect, and Time and 439
Level and their interaction as fixed effects. Each model was assessed for 440
goodness of fit, using random slopes or intercepts only, or their combination. 441
Models (5a-5d) were compared to test which model produced the lowest model 442
fit statistics [119]. The Akaike information criterion (AIC) is one of the best 443
known criteria to use in selecting a good model, where fitted values are most 444
likely to be true values, thus providing a fit closest to reality [120]. The best 445
regression model will provide the lowest value of AIC. 446

$$R \sim T + L + T \times L + (1|O) \tag{5a}$$
$$R \sim T + L + T \times L + (1 + T|O) \tag{5b}$$
$$R \sim T + L \ + T \times L + (1 + L|O) \tag{5c}$$
$$R \sim T + L + T \times L + (1 + L|O) + (1 + T|O) \tag{5d}$$

R is the observer response (proportion correct) T is time, the day or session of testing, L is the stimulus level (degree of coherence, level of contrast, or degree of orientation) and O is the observer.

Interpreting the model The intercept is the proportion of correct responses when the linear model value is zero. An increase in intercept over time reflects an overall increase in the number of correct responses. The slope relates to how the proportion of correct responses increases with the strength of the stimulus (degree of coherence, level of contrast, or degree of orientation). Overall performance can also be summarised in terms of the 75% threshold, the point at which observers responded correctly on 75% of trials.

Results

Feedback and perceptual learning: Experiment 1

To verify the spread of easy and difficult trials across the stimulus levels suggested by Liu et al. [114], accuracy for the first day was evaluated across all observers for each level of the stimulus and is reported in (Table 1). The accuracy across levels confirms that there is an even distribution of easy trials (85% and above) and difficult trials (65% and below) with the balance around 75%.

Table 1. **Accuracy at each level of the stimulus.**

Stimulus Levels	Coherence °	Accuracy (%)
180	180	98.1
300	60	96.0
335	25	80.4
340	20	77.7
345	15	73.5
350	10	64.1
355	5	61.5

To identify if this distribution of easy and difficult trials was sufficient for learning in the absence of trial-by-trial feedback, we analysed individuals' daily performance across the two training groups (Feedback or No Feedback). Individual responses from the 10 days' training were aggregated for each variable; observer, day and level of coherence. The response data were analysed independently for each group (Feedback or No Feedback) using the method

outlined above. Response was modelled as a fixed effect of day and coherence and an interaction between these two predictors (see Table 2). For the Feedback Group, there was a significant negative effect of day however there was a significant positive day × coherence interaction indicating there was a change in slope across the 10 days. The same analysis was undertaken for the No Feedback group, and found a main effect of day with a significant negative interaction. The no feedback group performed worse after 10 days training (see Fig. 5(b)). Performance over the 10 days is illustrated in Fig. 5(a), where the slope for the feedback group gets steeper over time, day 10 (blue) versus day 1 (red), along with the 75% discrimination threshold, which shifts leftwards, indicating a reduction in threshold between the start and end of training. Conversely, the group trained without feedback performs progressively worse over time and the 75% discrimination threshold shifts rightwards, indicating an increase in threshold.

Table 2. Fixed effects parameters for the Pilot Data

Training		β	SE	tStat	DF	pValue	LCI	UCI
No Feedback	Intercept (Day 1)	-1.595	0.1555	-10.258	836	2.456e-23	-1.90073	-1.2902
	Day (Change)	-0.0361	0.074	-0.4886	836	0.625	-0.1812	0.1089
	Coherence	0.07576	0.00207	36.598	836	8.65e-176 †	0.0717	0.0798
	Day × Coherence	-0.00075	0.00032	-2.30953	836	0.02116 *	-0.00139	-0.0001
Feedback	Intercept (Day 1)	-0.6099	0.140	-4.35	836	1.527e-05	-0.885	-0.335
	Day (Change)	-0.097	0.0076	-12.78	836	2.78e-34*	-0.112	-0.082
	Coherence	0.0292	0.0016	18.39	836	1.096e-63†	0.026	0.0324
	Day × Coherence	0.0087	0.0004	24.3234	836	3.052e-99*	0.00797	0.0094

Table notes :* indicates significant p value
and † performance increases significantly with increasing coherence

Learning and transfer of global motion: Experiment 2

The number of correct responses from the daily training was calculated for each observer, for each day and level of coherence. Training data were analysed independently for the three groups trained on different frequencies (Broad, Low, High) and were modelled as a fixed effect of day and coherence and an interaction between these two predictors.

Results were analysed using the method previously described (see Table 3). Performance improved for the broad and low trained groups and there was a significant positive coherence-by-day interaction found for both conditions. For the group trained with high spatial frequency stimuli there was a significant main effect of day. This shows that there was a significant increase in the total number of correct responses over the five days (see Fig. 3). These results thus show an improvement in performance during the training phase for all three spatial frequency conditions.

Fig 5. Mean thresholds (in degrees) over 10 days on global motion training for feedback (a) and no feedback (b) groups. The red and blue dotted lines show 75% threshold on days 1 and 10 respectively. 75% threshold decreased for the feedback group, and increased for the no feedback group.

Fig 6. Training results over the 5 days for broad, low and high spatial frequencies. The red and blue dotted lines show 75% threshold on days 1 and 5 respectively.

Table 3. Fixed effects parameters for the Training Data

Training		β	SE	tStat	DF	pValue	LCI	UCI
Broad	Intercept	-2.4642	0.74997	-3.2858	346	0.00111	-3.9393	-0.989
	Day	0.05161	0.0167	3.0869	346	0.00219*	0.0187	0.084
	Coherence	0.04848	0.0016	30.009	346	2.54e-98 †	0.0453	0.0517
	Day × Coherence	0.00271	0.000436	6.218	346	1.45e-09*	0.00185	0.00356
Low	Intercept	-1.081	0.5887	-1.8359	346	0.0672	-2.2388	0.0771
	Day	0.0421	0.120721	0.349	346	0.728	-0.19536	0.27952
	Coherence	0.0216	0.0012	18.1634	346	2.991e-52 †	0.0193	0.023981
	Day × Coherence	0.00331	0.0004	8.182	346	5.383e-15*	0.00251	0.0041
High	Intercept	-1.9579	0.1366	-14.336	346	6.56e-37	-2.227	-1.689
	Day	0.0894	0.02066	4.32719	346	1.98e-05*	0.048755	0.13001
	Coherence	0.0746	0.0026	28.3151	346	4.13e-92†	0.069	0.0798
	Day × Coherence	-0.0015	0.00079	-1.857	346	0.0642	-0.002995	8.61e-05

Table notes :* indicates significant p value
and † performance increases significantly with increasing coherence

Pre- and Post-Assessment Analysis: Global Motion

Learning is often measured through monitoring performance at a particular
threshold, which is expected to shift the psychometric function leftwards if
performance is improved, (see Fig. 7(a)). The psychometric function describes
performance in terms of accuracy as a function of the strength of the
stimulus [122], which are usually positively correlated, and it is expected to
reach asymptotic performance at the highest stimulus intensity [123]. Inspection
of the pre- and post-assessment data revealed that in some conditions
performance did not reach perfect accuracy, asymptoting at a proportion of
correct responses that was less than 1; this resulted in a poor psychometric fit of
the observer response data using the GLMM. To accommodate this, a nonlinear
generalised mixed effects model (NLME) was used to include an additional
parameter in order to model variability in the asymptotic performance at high
signal levels, as has been applied in other perceptual learning studies [124–126].

Nonlinear regression analysis

The nonlinear regression provides three measures to assess a change in
performance over time. Firstly, like the GLMM a leftward shift in the curve
indicates an improvement in threshold (7(a)). An increase in slope indicates an
increase in the rate at which performance increases with increase signal level
(7(b)). Finally a change in the asymptote indicates a significant change to the
performance at the highest level of stimulus intensity (7(c)). These changes are
independent aspects of the psychometric function fit, and may not necessarily be
congruent. For example, it is possible to obtain an increase in one measure and
a decrease (or no change) in another.

Analysis of the pre- and post-assessment data was undertaken using a
nonlinear mixed effects regression with the *nlmefit* function in Matlab with the
following model:

Fig 7. Psychometric functions illustrating the three measures by which the non-linear regression provides evidence of a change. (a) A leftward shift of the function indicates a general increase across stimulus levels, that did not vary as stimulus intensity increased. (b) A steeper slope indicates an increase in the number of correct responses as stimulus intensity increases. (c) An upward shift of the asymptote indicates an increase in performance where stimulus intensity is at its highest. However, in this example there has been no shift of the 75% correct detection threshold.

$$p = 0.5 + \frac{(A + A_d C)}{1 + e^{-(K + K_d S)(C - (C_0 + C_{0d} S))}} \tag{6}$$

where p is the proportion of correct responses, A determines the asymptotic level of performance, K defines the slope and C_0 defines the threshold. d is the post-training change in performance, where A_d, K_d and C_{0d} determine the change in asymptote, slope and threshold respectively. C is the coherence level, and S is a dummy variable, taking on values of 0 (pre) or 1 (post) training sessions.

95% confidence limits were calculated using parametric bootstrap for 1000 simulated experiments, with the same number of simulated observers and repetitions as the actual experiment. Results for each condition are plotted in Figs. 8 to 10.

Predictions

We predicted that should transfer occur cortically, this would be located at the global motion processing level, and transfer across frequencies would most likely occur for stimuli with frequency properties that reflect the broadband low-pass frequency tuning of global motion detectors (see Fig 11(a)). Firstly, we predicted no condition would show transfer to contrast sensitivity, as these are processed in different cortical locations. For global motion, we predicted;

- For the broad frequency trained condition; (i) robust learning on their own trained frequency (ii) moderate transfer to low spatial frequency global motion, and (iii) an unlikely but possible transfer to the high frequency condition.

- For the low frequency trained condition; (i) robust learning on their own

Fig 8. Top: Assessment results for the Broad Frequency trained condition on Broad, Low and High spatial frequency tests. Performance is presented as the proportion of correct responses, where blue circles show the results for the pre-assessment, and pink squares for the post-assessment. 75% thresholds for pre and post are indicated by the green and orange vertical dotted lines respectively. **Bottom:** Change statistics for the asymptote, threshold and slope, respectively. Plots show the median performance and the 95% confidence intervals for the change in performance between pre- and post-assessments. The red horizontal line at zero represents no change, confidence intervals crossing the zero line reflect no significant improvement. Points above the reference line show an improvement in performance and those below reflect a decrease in performance.

Fig 9. Top: Assessment results for the Low Frequency trained condition on Broad, Low and High spatial frequency tests. Performance is presented as proportion of correct responses, where blue (circles) show the results for the pre-assessment, and pink (squares) the post-assessment. 75% thresholds for pre- and post-assessments are indicated by the green and orange vertical dotted lines respectively.

Bottom: Change statistics for the asymptote, threshold and slope respectively. Plots show the median performance and the 95% confidence intervals for the change in performance between pre- and post-assessments. The red horizontal line at zero represents no change, confidence intervals crossing the zero line reflect no significant improvement. Points above the reference line show an improvement in performance and those below reflect a decrease in performance.

Fig 10. Top: Assessment results for the High Frequency trained condition on Broad, Low and High spatial frequency tests. Performance is presented as proportion of correct responses, where blue (circles) show the results for the pre-assessment, and pink (squares) the post-assessment. 75% thresholds for pre and post are indicated by the green and orange vertical dotted lines respectively. **Bottom:** Change statistics for the asymptote, threshold and slope respectively. Plots show the median performance and the 95% confidence intervals for the change in performance between pre- and post-assessments. The red horizontal line at zero represents no change, confidence intervals crossing the zero line reflect no significant improvement. Points above the reference line show an improvement in performance and those below reflect a decrease in performance.

trained frequency and (ii) robust transfer to broad spatial frequency global ₅₄₈ motion. ₅₄₉

- For the high frequency trained condition we predicted; (i) moderate ₅₅₀ learning on their own trained frequency and (ii) an unlikely but possible ₅₅₁ transfer to the broad frequency condition. ₅₅₂

Conversely, should transfer occur as a result of backwards generalisation ₅₅₃ using the re-entrant connections from global motion processing areas to V1 we ₅₅₄ would predict that transfer would reflect the pooling of spatial frequencies and ₅₅₅ likely attenuation of high spatial frequencies. ₅₅₆

- For the broad frequency trained condition; (i) robust learning on their own ₅₅₇ trained frequency (ii) moderate transfer to low spatial frequency global ₅₅₈ motion, (iii) a possible but unlikely transfer to high spatial frequency ₅₅₉ global motion and (iv) a moderate transfer to low spatial frequency ₅₆₀ contrast sensitivity ₅₆₁

- For the low frequency trained condition; (i) robust learning on their own ₅₆₂ trained frequency and (ii) robust transfer to broad spatial frequency global ₅₆₃ motion, and (iii) robust transfer to low spatial frequency contrast ₅₆₄ sensitivity ₅₆₅

- For the high frequency trained condition we predicted; (i) moderate ₅₆₆ learning on their own trained frequency and (ii) an unlikely but possible ₅₆₇ transfer to the broad frequency condition, and (iii) no transfer to any ₅₆₈ frequency of contrast sensitivity ₅₆₉

Transfer to global motion: broad-frequency trained group ₅₇₀

We had predicted robust to moderate improvement for the broad trained group, ₅₇₁ however this group showed no consistent transfer to trained or untrained ₅₇₂ conditions (see Fig. 8). The post-assessment on the **trained task** found no ₅₇₃ significant change in slope *(proportion correct as a function of frequency)* or ₅₇₄ threshold *(coherence required to obtain 75% correct)*, and a significantly lower ₅₇₅ asymptote *(proportion of correct responses at the highest coherence)*. Assessing ₅₇₆ transfer to **untrained tasks**, there was no significant change for any measure ₅₇₇ (intercept, slope or asymptote) for the **low frequency** test. Finally for the ₅₇₈ **high frequency** test, there was no change in slope, a significant reduction in ₅₇₉ asymptote (indicating worse performance at higher coherence), and a small but ₅₈₀ significant reduction in threshold. The increase in the proportion of correct ₅₈₁ responses at the lower coherence levels was the only significant improvement for ₅₈₂ this group, however neither of the other two measures were consistent with an ₅₈₃ improvement. ₅₈₄

Transfer to global motion: low-frequency trained group: ₅₈₅

The low-frequency trained group was the only condition to exhibit transfer to ₅₈₆ trained and untrained motion conditions (see Fig. 9). In the post-assessment on ₅₈₇

Fig 11. (a) Predictions of learning and transfer as shown in Fig. 1. We predicted a) that if transfer was local to the processing mechanism transfer would occur for global motion conditions with shared spatial frequency properties. (b) Performance improved for all groups during the five days training, with robust improvement for the low and broad trained groups, and moderate improvement for the high frequency group. However, at the post-assessment stage, when tested without the presence of trial by trial feedback, neither the broad nor high groups reflected a change from the pre-assessment measures. Improvement was restricted to the low trained group. (c) When assessing transfer to untrained frequencies, only the low spatial frequency group showed evidence of transfer, and performance improved for broad frequency motion. In addition, the low spatial frequency trained group was the only group to show an improvement on contrast sensitivity and this was exclusive to the low frequency contrast condition.

the **trained task**, while there was no significant change to the slope *(proportion* ₅₈₈
correct as a function of frequency) there was significant reduction in threshold ₅₈₉
(coherence required to obtain 75% correct), and a significantly higher asymptote ₅₉₀
(proportion of correct responses at the highest coherence). Assessing transfer to ₅₉₁
untrained tasks, there was a modest improvement for the **broad frequency** ₅₉₂
test reflected by the was a significant increase in asymptotic performance. ₅₉₃
However, there was an increase in threshold and no change to the slope. This ₅₉₄
shows reduced sensitivity at low signal levels, but an increase is performance at ₅₉₅
higher levels. Finally, the **high frequency** test stimuli show a small reduction ₅₉₆
in threshold but a significantly shallower slope. Suggestive of increased ₅₉₇
performance at lower coherence but decreased performance at higher stimulus ₅₉₈
coherence. There was no change in asymptotic performance, and overall no ₅₉₉
evidence of transfer. ₆₀₀

Transfer to global motion: high-frequency trained group: ₆₀₁

The group trained with high spatial frequency stimuli showed no transfer to any ₆₀₂
trained or untrained motion condition for any of the three measures (see ₆₀₃
Fig. 10). ₆₀₄

Transfer to global motion: Participant level variability ₆₀₅

To investigate the lack of transfer to the trained task, a participant level plot ₆₀₆
(see Fig.12) illustrates the change in number of correct responses between pre- ₆₀₇
and post-training. This was calculated by subtracting the pre-assessment scores ₆₀₈
from the post-assessment scores. Changes above the dotted line at 0 show ₆₀₉
improved performance, while those below the line show worsened performance. ₆₁₀
The spread of scores for the broad/broad and broad/high condition show that ₆₁₁
one observer performed particularly poorly at higher coherence levels, but not ₆₁₂
persistently so. Therefore, the lack of overall improvement for the broad trained ₆₁₃
group is unlikely to be as a result of an outlier. However, the significantly worse ₆₁₄
performance in asymptotic performance may, to some degree, be as a result of ₆₁₅
this one individual. ₆₁₆

Transfer to global motion: Summary ₆₁₇

In summary, only the low frequency trained group provided reliable evidence for ₆₁₈
transfer to trained and untrained motion conditions, with a moderate transfer to ₆₁₉
the broadband frequency condition. ₆₂₀

Fig 12. Change scores (the difference between the number correct on the pre- and post-assessments) are plotted for each individual. Each observer is illustrated by a colour grouped by training and test condition (broad, low and high spatial frequency motion). Of particular interest is the broad trained group, which had a decline in performance post training. One observer performed worse at higher coherence levels after training, which explains the decline in asymptotic performance for the broad trained group. However, the overall spread of data suggest that on average post-assessment results did not differ significantly.

Transfer to contrast sensitivity 621

We predicted that, should contrast sensitivity improve as a result of training on global motion it could only occur as a result of backward generalisation, as the two tasks are processed in cortically separate locations. We predicted transfer would be to limited conditions containing low frequencies, namely the broad and low trained groups.

Data were analysed for all but the same 2 observers who did not complete the final session using the generalised linear effects model previously outlined. The measures of analysis are the intercept and slope, and an interaction between the two predictors.

Table 4. Contrast Sensitivity Table

Training	Low Frequency	β	SE	tStat	DF	pValue	LCI	UCI
Broad	Intercept (Pre)	-4.0361	1.3136	-3.0725	129	0.0026	-6.6352	-1.437
	Intercept (Post)	-1.4324	0.39039	-3.6691	129	0.00035*	-2.2047	-0.66
	Contrast	33.009	9.4797	3.4821	129	0.00068†	14.253	51.765
	Session × Contrast	8.4892	2.2901	3.707	129	0.00031*	3.9582	13.02
Low	Intercept (Pre)	-1.9253	0.66009	-2.9167	129	0.00417	-3.2313	-0.61929
	Intercept (Post)	-1.6317	0.4122	-3.9585	129	0.00012*	-2.4472	-0.81612
	Contrast	11.338	3.203	3.5397	129	0.00056†	5.0004	17.675
	Session × Contrast	13.035	2.4795	5.2571	129	5.90e-07*	8.129	17.94
High	Intercept (Pre)	-3.5587	0.78832	-4.5143	136	1.3627e-05	-5.1177	-1.9998
	Intercept (Post)	-0.20622	0.45674	-0.45151	136	0.65234	-1.1095	0.69701
	Contrast	24.238	3.6543	6.6329	136	7.14e-10†	17.012	31.465
	Session × Contrast	3.5416	2.4152	1.4664	136	0.14485	-1.2346	8.3177

Training	High Frequency	β	SE	tStat	DF	pValue	LCI	UCI
Broad	Intercept (Pre)	-3.1141	0.72157	-4.3158	129	3.14e-05	-4.5417	-1.6865
	Intercept (Post)	-0.39475	0.28081	-1.4057	129	0.1622	-0.95034	0.16084
	Contrast	17.372	2.3389	7.4273	129	1.34e-11†	12.744	21.999
	Session × Contrast	2.3333	1.5341	1.521	129	0.13072	-0.70195	5.3686
Low	Intercept (Pre)	-3.388	0.41928	-8.0805	129	4.01e-13	-4.2175	-2.5584
	Intercept (Post)	0.0001	0.25711	0.00045	129	0.99964	-0.509	0.50881
	Contrast	22.041	2.9883	7.3757	129	1.76e-11†	16.128	27.953
	Session × Contrast	-1.236	1.1863	-1.0418	129	0.29943	-3.5831	1.1112
High	Intercept (Pre)	-3.5827	0.78951	-4.5379	136	1.2371e-05	-5.144	-2.0214
	Intercept (Post)	-0.045188	0.26063	-0.17338	136	0.86261	-0.56059	0.47022
	Contrast	20.64	2.2427	9.203	136	5.496e-16†	16.205	25.075
	Session × Contrast	-0.7794	1.3781	-0.56554	136	0.57264	-3.5048	1.946

Table Notes:* indicates significant p value
and † performance increases significantly with increasing coherence

Broad-frequency trained group: 631

For low spatial frequency contrast there was a a significant decrease in intercept *(fewer correct responses at the lowest coherence)*, but significant increase in slope *(an increase in the proportion of correct responses as a function of frequency).*

Fig 13. Contrast sensitivity results for Broad (top row), Low (middle) and High (bottom) trained groups for pre- and post-assessments. Low spatial frequencies are shown in the left column and high spatial frequencies on the right. Performance is presented as the proportion of correct responses as a function of Michelson Contrast. Blue (triangles) show the results for the pre-assessment, and pink (crosses) the post-assessment. 75% thresholds for pre and post are indicated by the green and orange vertical lines respectively.

Because of this ambiguity we plotted 75% thresholds for pre and post performance (see Fig. 13 top left). Post-assessment thresholds shifted rightwards suggesting worse performance overall. 635 636 637

Low-frequency trained group: 638

The group trained with low frequency global motion showed a significant decrease in intercept, and a significant increase in slope. Again, we plotted 75% thresholds for pre and post performance (Fig. 13 left middle). In this instance the steeper slope, and leftward shift of the threshold, suggests improvement overall. 639 640 641 642 643

Fig 14. Orientation discrimination results for all trained groups (broad, low and high) for pre- and post-assessments. Performance is presented as proportion of correct responses, where blue (triangles) show the results for the pre-assessment, and pink (crosses) the post-assessment. 75% thresholds for pre and post are indicated by the green and orange vertical lines respectively.

High-frequency trained group:

The high frequency trained group showed no significant change for any condition (see Fig. 13 bottom left).

Pre and Post Analysis: Orientation Discrimination

This task was added as an untrained control, for which we predicted no improvement, as no orientation information was present in any of the training stimuli. Data were analysed for all but 3 observers who did not complete the final session, using the generalised linear effects model previously outlined. The measures of analysis are the intercept and slope , and an interaction between the two variables. For analysis, the orientation levels were converted to absolute values. Results are plotted in Fig. 14, and the fixed effects are listed in tab 5. Neither the broad- or low-frequency trained group showed any significant change on this task. However, the high spatial frequency trained group showed a significant decrease in intercept over session, and a significant increase in slope following training. However, the 75% threshold at post-assessment shift rightwards, suggesting worse performance overall.

Discussion

Damage to the visual cortex, as a result of stroke or other brain injury can result in dramatic changes to connectivity between areas. It has been proposed that

Table 5. Fixed Effects table for Orientation Discrimination

Training		β	SE	tStat	DF	pValue	LCI	UCI
Broad	Intercept (Pre)	-1.9841	0.79477	-2.4964	284	0.013	-3.5485	-0.41967
	Intercept (Post)	0.02189	0.55128	0.039723	284	0.96834	-1.0632	1.107
	Orientation	2.8444	0.35047	8.1161	284	1.47e-14†	2.1546	3.5343
	Session $\times$ Orientation	0.0025	0.10152	0.0245	284	0.981	-0.197	0.2023
Low	Intercept (Pre)	-1.5733	0.78192	-2.0121	284	0.045	-3.1124	-0.034191
	Intercept (Post)	-0.38185	0.58052	-0.65778	284	0.511	-1.5245	0.76081
	Orientation	2.3483	0.29959	7.8381	284	9.25e-14†	1.7586	2.938
	Session $\times$ Orientation	0.18935	0.097122	1.9496	284	0.0522	-0.0018	0.38052
High	Intercept (Pre)	-1.1825	0.37792	-3.1289	284	0.00194	-1.9263	-0.43858
	Intercept (Post)	-0.47101	0.1002	-4.7005	284	4.05e-06	-0.66824	-0.27377
	Orientation	2.3751	0.33682	7.0516	284	1.35e-11†	1.7121	3.0381
	Session $\times$ Orientation	0.28973	0.08728	3.3196	284	0.001*	0.11794	0.46153

Table notes : † performance increases significantly with increasing coherence

perceptual training may play a role in strengthening existing neural pathways and may even create novel connections [1–7]. While specificity is a limitation to the effectiveness of perceptual learning as a tool for therapy, there is evidence that this specificity is reduced for visually impaired populations [127–132]. Huxlin et al. (2009) [5] proposed that using global motion may tap into "islands of activity" within V1 through the feedback connections from V5 to V1. However, it is still unclear how much the brain is able to compensate for damage, and whether recovery involves building new connections or changes in the functional connectivity within the existing pathways [133]. Understanding how the cortical hierarchy is organised in terms of the nature of the feedforward and re-entrant pathways is central to developing theories of perception [134].

Feedback and Perceptual Learning: Experiment 1

The purpose of the first experiment was to establish the necessity of performance feedback when undertaking a period of training on a task to discriminate the direction of global motion. Fahle and Edelman (1993) [28] predicted that internal reinforcement could act as the teaching signal when performance feedback was absent and when the confidence is high [34, 135]. Therefore, perceptual learning should occur when training procedures include a mixture of easy and difficult trials. This was the case for the study conducted by Liu et al. (2012) [114] where learning occurred for easy and difficult trials without feedback. For this specific task, consistent with Seitz et al. (2006) [136], our results found that feedback was a requirement for learning. After ten days of training the group who received feedback improved significantly, while the performance for the no feedback group deteriorated.

Our predictions were based on the findings of Liu et al. (2012) [114] who demonstrated that interleaving high accuracy (easy) trials and low accuracy (difficult) trials, resulted in perceptual learning without the need for feedback.

Liu et al. (2012) [114] explain their findings using the Augmented Hebbian   690
Re-weighting Model (AHRM) [21]. When external feedback is provided, the   691
post-synaptic activation is shifted further in the correct direction, enforcing   692
appropriate weight changes in the decision unit. However, when external   693
feedback is absent the model uses the observer's internal response. In this   694
situation learning is dependent on the level of difficulty of the task, and uses the   695
observer's internal confidence to update the weights [114]. Where a task is easy,   696
the weights still move, on average, in the correct direction [137]. For a difficult   697
task, the neural signal is weak and a clear indication of the appropriate changes   698
that are required is absent. As a result the process of updating the decision   699
weights is ineffective, and learning does not occur.   700

The results of our study showed that learning did not occur without feedback,   701
even when easy trials were presented. While our task spanned the 65-85%   702
accuracy range used by Liu et al. (2012) [114], there were some differences   703
between the studies. Firstly, Liu et al. (2012) used an adaptive staircase to   704
track performance and stimuli were presented at either 85% or 65% accuracy,   705
and at no other levels, throughout the experiment. Ours, on the other hand, did   706
not include an adaptive staircase. Stimuli were randomly presented using   707
MOCS. This method involves random stimulus selection from a predefined range   708
of stimulus magnitudes [138]. In our experiment observers were randomly   709
presented 1 of 7 threshold levels. Interleaving these levels may influence how   710
decision weights are updated and reduce the observer's confidence in their   711
judgements, for example by disrupting observers' meta-cognitive judgements of   712
perceptual confidence [139] and thus their ability to selectively weight   713
high-confidence trials in perceptual learning [34].   714

Our findings bear some similarity to those found by Seitz et al. (2006) [136],   715
who also used MOCS. They found that learning did not occur without feedback,   716
even when easy trials were presented. Observers trained on one of two tasks,   717
either to discriminate the direction of low luminance motion stimuli, or to   718
discriminate the orientation of a bar that had been masked in spatial noise.   719
After training, both groups who had received feedback showed an improvement,   720
while the groups without feedback did not improve. Seitz et al. (2006) [136]   721
note that many of the experiments investigating perceptual learning use   722
adaptive staircase procedures, and even though easy trials were present, this   723
difference may contribute to the difference in findings. Seitz et al. (2006) [136]   724
propose that interleaving easy and difficult trials within the staircase may allow   725
for "better bootstrapping" from easy to hard, compared to trials that are   726
randomly presented.   727

A second difference between the study conducted by Liu et al. (2012) and   728
ours, was the type of task used. We used a global motion coherence task and   729
not an orientation discrimination task [114]. These tasks are known to be   730
processed at different levels of the visual hierarchy. Learning without feedback   731
has been obtained for local motion tasks. Ball and Sekuler (1982) [15] found   732
that observers did not need feedback to improve on their direction   733
discrimination task. In their study, observers were required to make a   734
same/different judgement for two rapidly presented trials. In the "same" trials,   735

motion took the same direction, and in the "different" trials the direction of motion varied by 3°. However, while the no feedback group did not receive trial-by-trial response feedback, they were rewarded with two cents for a correct response and had one cent deducted for each incorrect response, which may be construed as end of block feedback [136], which has been found to be as effective as trial-by-trial feedback [113].

Learning without feedback has also been found for global motion coherence tasks [78,140]. However, none of these examples used the equivalent noise coherence task, but rather used the ratio of signal-to-noise coherence task. Levi et al. [75], using an equivalent noise method of global motion, also found perceptual learning, however they used an adaptive staircase for training paired with trial-by-trial feedback. When investigating task-irrelevant learning, Watanabe et al. (2002) [141] found that task-irrelevant local motion improved passively, but did not find the same for task-irrelevant global motion. They suggest that this is indicative of the lower levels of the visual hierarchy being more receptive to modification, when attention is limited. It was unexpected for the group trained without feedback that performance would deteriorate, and highlights interesting questions about motivation and performance over time as confidence and attention declines. The results of this study ultimately provide us with evidence that to obtain learning during training, using the equivalent noise motion coherence used by Levi et al. (2015) [75], and presenting stimuli randomly using MOCS, our procedure should include trial-by-trial feedback during the training process. Furthermore, since learning did not occur without feedback, no feedback would be provided during the pre- and post-assessment phases.

Learning and Transfer of Global Motion: Experiment 2

Having established the necessity of feedback for learning, the objective of the main study was to investigate the specificity of spatial frequency tuning in perceptual learning for global motion. We trained three groups of observers on a global motion task with stimuli tuned to three different spatial frequency ranges (Broad, Low and High) and performed pre- and post-training assessments for all frequencies and high and low spatial frequency contrast detection.

Summary of Training

Our training results demonstrated that after the five day training period on a frequency specific global motion coherence task with trial-by-trial feedback, there was significant improvement in performance for all trained groups.

Pre- and Post-Assessment predictions

Based on the evidence for the frequency tuning of V5 [79,80,96,104], we predicted that we might have obtained greater improvements (and transfer) between broad and low spatial frequencies, but limited or no transfer for the high spatial frequency trained group. Furthermore, we predicted that should

transfer occur as a result of the reweighting of the cortically localised global 777
motion mechanisms, then any transfer would reflect the broad, approximately 778
low-pass spatial frequency tuning of global motion detectors. There would thus 779
be most transfer when both training and test stimuli contained low spatial 780
frequency components, and a moderate improvement for those trained or tested 781
with broad frequency components. Since contrast discrimination is not 782
processed in the same cortical location as global motion we predicted no transfer 783
to contrast sensitivity would occur from any trained condition. 784

If transfer occurs as a result of backward generalisation, making use of the 785
re-entrant (or feedback) connections as postulated by the Reverse Hierarchy 786
Theory [33, 72], we predicted transfer to show specificity to the frequency tuning 787
of the global motion processing areas. Given the broadband low-pass frequency 788
tuning of the motion processing areas, such as V5 and V3A, we predicted robust 789
to moderate improvement where stimuli contain low spatial frequency 790
components (learning to transfer from low to low; modest transfer of learning 791
from low to broad, and broad to low and broad). In addition, we predicted 792
frequency-specific improvement in contrast sensitivity with the broad trained 793
group showing the most transfer, and improvements reflecting the spatial 794
frequency tuning of the training stimuli. 795

Comment on the predictions 796

There is the possibility that transfer relies on forward connections from local to 797
global motion mechanisms. However, given the tight frequency tuning at the 798
local level and the well evidenced, highly specific characteristic of area V1, we 799
would not make a prediction of transfer across frequencies occurring as a result 800
of low level feed-forward mechanisms. However if learning did rely on the 801
connections from local or global levels in a hierarchical fashion, we would expect 802
frequency specific improvement on the matching spatial frequency motion 803
condition (low to low, high to high, and broad to low and high). However, any 804
transfer at the level of global motion, would depend on the global level 805
mechanism so would again reflect the tuning properties of the global detectors. 806

Pre- and Post-Assessment findings 807

Our analysis of the pre- and post-assessment data found most support for 808
transfer as a result of backwards generalisation, however not all our predictions 809
were supported. 810

Firstly, while not an explicit prediction, we had expected that the groups 811
trained on their specific frequency conditions would naturally show improved 812
performance for their trained frequency. However, when analysing the pre- and 813
post-training motion results only the group trained using low spatial frequency 814
Gabors displayed a significant learning, i.e. 'transfer to the trained task'. 815
Notably, the broad trained group performed worse than they did at 816
pre-assessment stage, and there was no change for the high trained group. 817
Furthermore the only evidence of transfer to untrained conditions, was from the 818
low frequency trained group on the broad test condition. Pre- and post-training 819

assessment for contrast sensitivity found that there was a significant ₈₂₀
improvement exclusively for the low trained group on the low spatial frequency ₈₂₁
contrast condition. No further improvement was found for any other trained or ₈₂₂
tested frequency. Finally, the orientation discrimination control task showed no ₈₂₃
improvement from any trained frequency. ₈₂₄

Evaluating the transfer from global motion ₈₂₅

This study explored the effects of training on global motion and its transfer to ₈₂₆
other spatial frequencies and tasks. When assessing the post-training transfer to ₈₂₇
trained and untrained global motion frequencies, the *low spatial frequency* ₈₂₈
trained group was the only group to 'transfer to trained task' and transfer to ₈₂₉
another condition. Although the 75% threshold was worse for the broad test ₈₃₀
after training, the asymptotic performance was significantly better. Interestingly, ₈₃₁
for the high spatial frequency test, there was a significant improvement to the ₈₃₂
threshold although the slope was significantly shallower. This suggests that the ₈₃₃
sensitivity for lower coherence levels increased. Performance at the highest levels ₈₃₄
reached an asymptotic performance close to 1 at pre-assessment and remained ₈₃₅
unchanged at the post-assessment stage. Finally the shallower slope suggests a ₈₃₆
reduction in correct responses as a function of stimulus intensity. ₈₃₇

The *high spatial frequency trained group* showed no improvement and no ₈₃₈
transfer to any other spatial frequency. ₈₃₉

The most surprising results were those obtained from the *broad frequency* ₈₄₀
trained group. Asymptotic performance was significantly worse at the post test ₈₄₁
stage for their own trained frequency and the high frequency test, with no ₈₄₂
significant change in the low frequency test. There was no improvement in the ₈₄₃
slopes for any condition, and a small but significant improvement was found in ₈₄₄
75% threshold for the high frequency test. Viewing individual performance ₈₄₅
revealed that the reduction in asymptotic performance may be accounted for by ₈₄₆
an outlier, however the outlier was unlikely to account for the overall absence of ₈₄₇
improvement in performance across all levels and measures, as there was no ₈₄₈
clear improvement evident for the other observers. ₈₄₉

Comparisons to Levi et al. ₈₅₀

Since Levi et al. (2015) reported improved contrast detection for low frequency ₈₅₁
drifting targets, we had questioned if the improvement was specific to the ₈₅₂
temporal and spatial features of the training stimuli. Half the cells in layer $4C\alpha$ ₈₅₃
of V1, where improvement in contrast has been argued to take effect [27], are ₈₅₄
tuned for direction of motion [142]. It might be expected therefore that ₈₅₅
improvement in contrast sensitivity would be limited to moving stimuli. ₈₅₆

Levi et al. (2015) evaluated three groups. The first trained on static, ₈₅₇
cardinally oriented Gabors, and made a horizontal or vertical direction ₈₅₈
judgement. The second trained on high intensity broadband frequency random ₈₅₉
dot coherence task, also making a direction judgement. The final group was ₈₆₀
untrained but undertook pre- and post-measures. We focus directly on the ₈₆₁

findings from the second training group which formed the basis for our initial ₈₆₂
enquiry. ₈₆₃

Training ₈₆₄

Levi et al. trained observers using a 3 up 1 down staircase, and provided ₈₆₅
auditory feedback on every trial. After 10 days training (300 trials daily), they ₈₆₆
reported an improvement in global motion perception. There was no specific pre- ₈₆₇
or post-measure for global motion integration (without trial-by-trial feedback). ₈₆₈
Improvement as a result of training was assessed by calculating the difference ₈₆₉
between performance on the first and last days of training. Our training was ₈₇₀
delivered using MOCS over 5 days with 420 trials (across 7 defined levels). To ₈₇₁
assess improvement across the training days we used a mixed-effects regression ₈₇₂
that includes all the responses over the 5 days [119]. Our training results are ₈₇₃
fully consistent with the findings of Levi et al.; training on a global motion ₈₇₄
integration task improves when training is paired with trial-by-trial feedback. ₈₇₅

Pre and post-measures ₈₇₆

As previously stated Levi et al., did not undertake pre- and post-measures for ₈₇₇
the trained task. Thus while our post-assessment for the broadband trained ₈₇₈
group showed that once feedback was removed the task was performed at prior ₈₇₉
levels, we are unable to make a direct comparison between their study and ours ₈₈₀
on this finding. Levi et al., tested four measures of transfer to contrast ₈₈₁
sensitivity; (i) orientation discrimination of static, horizontal or vertical ₈₈₂
non-flickering Gabors at varying spatial frequency (ii) orientation discrimination ₈₈₃
of moving horizontal or vertical gratings (temporal frequency of 10 hz) of ₈₈₄
varying spatial frequency (iii) direction discrimination (left/right) of moving ₈₈₅
gratings (temporal frequency of 10 hz) of varying spatial frequency and finally ₈₈₆
(iv) direction discrimination (left/right) of moving gratings (varying temporal ₈₈₇
frequency) and a spatial frequency held constant at 2 cycles per degree (medium ₈₈₈
frequency). ₈₈₉

For (i) static ordinal Gabors, the group showed no improvement at any ₈₉₀
spatial frequency and the pre- and post-measures were no different from the ₈₉₁
untrained group. Only the moving gratings displayed an improvement in ₈₉₂
contrast sensitivity and these were predominantly found for low spatial ₈₉₃
frequency gratings. ₈₉₄

The result from (i) is the most similar to our static contrast detection ₈₉₅
condition. Like Levi et al., our group trained on broadband global motion ₈₉₆
stimuli displayed no improvement for any tested spatial frequency for static ₈₉₇
contrast detection. ₈₉₈

Since Levi et al. (2015) [75] reported the biggest improvement for low spatial ₈₉₉
frequency stimuli exclusively in moving targets, this may suggest improvement ₉₀₀
was specific to the temporal and spatial features of the training stimuli. In ₉₀₁
contrast, our results suggest that when trained on low frequency global motion, ₉₀₂
robust learning and transfer occurred for stimuli that contained low frequency ₉₀₃

elements, including to a static contrast sensitivity task. Both of these findings ₉₀₄
are consistent with the low-pass frequency tuning of V5. ₉₀₅

What our findings suggest for the mechanisms of perceptual learning ₉₀₆ ₉₀₇

Specificity ₉₀₈

We predicted that should global motion training improve contrast detection, ₉₀₉
given the frequency tuning of the early visual areas, we would expect specificity ₉₁₀
to the spatial frequency of training. Improvement was restricted to the low ₉₁₁
frequency contrast condition for the low trained group. The specificity of ₉₁₂
transfer to low frequencies is consistent with the low-pass frequency tuning of ₉₁₃
global motion areas such as V5 and V3/V3A influencing processing at V1 ₉₁₄
through feedback loops [109]. These results suggest however that there is some ₉₁₅
specificity for low frequency elements. This remains compatible with the view ₉₁₆
that global motion detectors pool via low frequency information. This may ₉₁₇
suggest that the re-entrant connections to V1, after training on global motion, ₉₁₈
only update low frequency channels. ₉₁₉

Transfer ₉₂₀

We predicted that if transfer occurred as a result of backward generalisation, ₉₂₁
then improvement would reflect the frequency tuning of the global motion ₉₂₂
detectors. The group trained on high spatial frequencies showed no improvement ₉₂₃
and no transfer. Since the representation of high spatial frequency content is ₉₂₄
attenuated in V5 [104], this may explain the lack of improvement in any ₉₂₅
post-assessment from the high or broad frequency trained groups. This may ₉₂₆
suggest any stimuli containing a high frequency element are attenuated for the ₉₂₇
purpose of the backward projections to V1 from global motion processing areas. ₉₂₈
   The broad trained group performed worse overall for almost all ₉₂₉
post-assessments. For the low spatial frequency contrast detection, there was ₉₃₀
evidence of a significantly steeper slope, however (a change in the rate of ₉₃₁
increase in proportion of correct responses with contrast), however this did not ₉₃₂
lead to a robust improvement, and the average detection threshold was worse. ₉₃₃
   Transfer was only found for the low frequency trained group. This group ₉₃₄
showed improvement in low spatial frequency contrast detection, a significant ₉₃₅
increase in asymptotic performance for the broad frequency global motion test, ₉₃₆
and also improved on their own task. This may suggest that after training on ₉₃₇
global motion only the low frequency channels are updated. The low frequency ₉₃₈
trained group will have experienced a high level of correlation in the activity in ₉₃₉
frequency channels between the feedforward and feedback connections. This ₉₄₀
joint activity may be important for perceptual learning, and is consistent with ₉₄₁
the proposal that there is an iterative interaction between the global motion ₉₄₂
processing areas and V1, since feedback from higher to lower areas engage the ₉₄₃
connections that mirror the features of the stimulus [65]. ₉₄₄

What our findings suggest for the models of perceptual learning 945

The two main theories of visual perceptual learning suggest that learning either 946
occurs through physiological changes to sensory neurons [18,19,31], or as a 947
reweighting [22] of the decision units. Our findings are not able to distinguish 948
between learning as a result of a physiological change or a higher-level 949
computational reweighting. However neither model is able to account for the 950
breadth of learning (and non-learning) and transfer within our results including 951
the general absence of improvement and the occasional decline in performance at 952
the post-assessment stage. The only difference between these components of the 953
experiment was that trial-by-trial feedback was present for the training phase 954
and absent for the testing phase. Our pilot study found that, for our methods, 955
learning occurs with feedback, but not without. Additionally, Herzog and Fahle 956
(1997) [113] suggested that once feedback is removed, performance plateaus 957
around the level last obtained with feedback, so our expectation was that 958
observers would maintain the improvement achieved for their trained frequency. 959
Neither model alone accounts for the ambiguous performance at the 960
post-assessment stage or the decline in performance by the broad trained group. 961
On the other hand, the specificity to low spatial frequencies is consistent with 962
both models, given the frequency tuning of the global motion detectors. 963

While the two positions are pitched as competing theories, they may not be 964
mutually exclusive. In an attempt to combine the two models, Solgi and Weng 965
(2013) [143] model learning as a two-way process (descending and ascending). 966
The model attributes training effects to the reweighting of connections between 967
early and higher sensory areas, but crucially also as a result of increased 968
neuronal activity representing the trained feature [143]. For transfer to occur, 969
Solgi and Weng (2013) [143] argue there is an important role for the re-entrant 970
connections from higher levels. Similarly, Bejjanki et al. (2011) [144] propose a 971
model that is computationally similar to the reweighting models, however it 972
illustrates how changing the population codes in the early sensory areas can 973
create similar changes in response to those made by high-level re-weighting 974
models. Bejjanki et al. (2011) [144] argue that their model captures the 975
characteristics for perceptual learning for both behavioural and physiological 976
changes. Sensory inputs improve the decision weights in the feed-forward 977
connections, and improved probabilistic inference in early visual areas as result 978
of the increased neural activity in the feedback network. This is compatible with 979
our findings that suggest there the high level of consistency experienced for the 980
group trained and tested on the low frequency conditions. Ultimately, a model 981
of perceptual learning needs to account for multiple cognitive influences. 982
Maniglia and Seitz (2018) [145] argue that perceptual learning is not a detached 983
or isolated process, instead multiple mechanisms react and interact to produce 984
learning. 985

Afterthoughts, Speculation and Unresolved Points 986

We offer some speculative ideas that may help account for the general absence of 987
improvement at the post-assessment stage. There is emerging evidence that 988

interleaving random stimuli may disrupt perceptual learning [146]. Yu et al. 989
(2004) [56] found that when interleaving random contrasted Gabor stimuli, such 990
as when using MOCS, learning did not occur, but did when only one contrast 991
was presented at a time. Yu et al. (2004) [56] termed the interleaving of contrast 992
stimuli as "contrast roving", and suggest that the effect of non-learning as a 993
result of contrast roving implies that a temporally organised pattern of stimulus 994
presentation may be required for perceptual learning to occur. Additionally, 995
Kuai et al. (2005) [58] found that temporal patterning was evident for motion 996
direction discrimination. Observers were either trained using randomly varied 997
trials or with a fixed temporal pattern of exposure (changing in a sequential 998
clockwise direction). The first group showed no improvement in discrimination, 999
however the second group with the fixed temporal exposure did. Cong and 1000
Zhang (2014) [147] showed that a semantic tag associated with each stimulus 1001
presentation enabled significant learning. Sequential tags such as A,B,C,D and 1002
1,2,3,4 which contain semantic and sequential identity information both resulted 1003
in significant learning. Cong and Zhang (2014) [147] suggest that tagging and 1004
temporal sequencing may help direct the system to switch attention to the 1005
neuronal set responsible to the specific stimulus. These are consistent with the 1006
experiments conducted by Seitz et al. (2006) [136], which as previously 1007
discussed found no learning without feedback using MOCS. 1008

In addition, there is also evidence that interleaving easy and difficult trials 1009
during perceptual training may only lead to a temporary improvement, that (a) 1010
does not persist over time and (b) disappears when easy trials are 1011
removed [148, 149]. In a difficult task without feedback learning does not occur, 1012
however it does with feedback [114]. Removing the feedback may be equivalent 1013
to removing the easy trials, and therefore reducing confidence, which is known 1014
to be an important factor in learning and perception [71]. Lin and Dosher 1015
(2017) [150] presented observers with a single block of mixed easy and difficult 1016
trials which led to rapid improvement within the block. However, after removing 1017
the easy trials and presenting only the difficult trials performance slowly 1018
returned to prior levels. This is consistent with findings that internal 1019
reinforcement could act as the teaching signal when confidence is high [34, 135]. 1020

In our study, improvement in the trained task was sustained only while 1021
feedback was present. At the post-assessment stage, the combination of a 1022
difficult task, the interleaved coherence levels and the general low decision 1023
confidence in the absence of feedback may have inhibited 'transfer to the trained 1024
task' and, like Lin and Dosher (2017), performance returned to baseline. It is 1025
thus a possibility that the amount of learning for the broad and high spatial 1026
frequency trained groups was insufficient to overcome the reduction in 1027
performance when feedback was removed. 1028

An exception to this was the improvement in performance and 'transfer to 1029
the trained task', for the low frequency trained group. Since learning and 1030
transfer to trained task were evident for the low spatial frequency trained group, 1031
this may suggest an interaction between the amount of learning, and the benefit 1032
of feedback. Furthermore, the improvement by the low frequency group over the 1033
five training days shows the biggest move from baseline performance, and thus 1034

potentially benefited from more learning, which was robust, persistent and thus incurred less disruption by the absence of feedback.

Conclusions

Our results support the early ideas from Reverse Hierarchy Theory that the feedback connections from higher levels back to early visual processing areas may be involved in facilitating learning that transfers [33, 72], and are consistent with models that suggest that the re-entrant connections play a vital role in learning and transfer [143, 144].

Our first experiment suggested that for perceptual learning of an equivalent noise global motion task, presented using MOCS, feedback was a necessity. However after training with feedback, there was little evidence of a robust improvement when feedback was removed at the post-assessment stage. The only condition that showed unambiguous learning and transfer was the low spatial frequency trained group. Our findings are in line with current research which finds global motion detectors pool for low spatial frequencies.

Our main experiment found robust learning and transfer for the group trained with low spatial frequency specific global motion gabors. This has provided evidence of frequency specific transfer from a higher-level global motion task to a low-level static untrained local task. The frequency tuning of these results is consistent with perceptual learning which depends on the global stage of motion processing.

Finally, the frequency tuning of these results suggest that the mechanism of transfer of global motion is dependent on backward generalisation, and specific to the frequency tuning of the global motion detectors. Transfer exhibited a high specificity towards low spatial frequencies, consistent with the attenuation found in the processing of global motion.

Our study supports the idea that the perception of global motion shows a bias for low spatial frequencies and suggests the backward projections and the re-entrant connections from global motion processing areas to V1 play a crucial role in transfer to trained and untrained tasks.

Acknowledgments

This research was funded by a University of Essex Doctoral studentship, and grants from ESSEXLab and PsyPAG to J.A.

References

1. Campana G, Maniglia M. Editorial: Improving visual deficits with perceptual learning. Frontiers in Psychology. 2015;6(APR):1–3. doi:10.3758/BF03194845.

2. Hess RF, Hayes A, Field DJ. Contour integration and cortical processing. Journal of Physiology Paris. 2003;97(2-3):105–119. doi:10.1016/j.jphysparis.2003.09.013.

3. Levi DM, Li RW. Perceptual learning as a potential treatment for amblyopia: A mini-review. Vision Research. 2009;49(21):2535–2549. doi:10.1016/j.visres.2009.02.010.

4. Polat U. Making perceptual learning practical to improve visual functions. Vision Research. 2009;49(21):2566–2573. doi:10.1016/j.visres.2009.06.005.

5. Huxlin KR, Martin T, Kelly K, Riley M, Friedman DI, Burgin WS, et al. Perceptual relearning of complex visual motion after V1 damage in humans. The Journal of Neuroscience. 2009;29(13):3981–3991. doi:10.1523/JNEUROSCI.4882-08.2009.

6. Sahraie A, Trevethan CT, MacLeod MJ, Murray AD, Olson JA, Weiskrantz L. Increased sensitivity after repeated stimulation of residual spatial channels in blindsight. Proceedings of the National Academy of Sciences of the United States of America. 2006;103(40):14971–14976. doi:10.1073/pnas.0607073103.

7. Trevethan CT, Urquhart J, Ward R, Gentleman D, Sahraie A. Evidence for perceptual learning with repeated stimulation after partial and total cortical blindness. Advances in Cognitive Psychology. 2012;8(1):29–37. doi:10.2478/v10053-008-0099-8.

8. Sagi D. Perceptual learning in Vision Research. Vision Research. 2011;51(13):1552–1566. doi:10.1016/j.visres.2010.10.019.

9. Seitz AR, Watanabe T. A unified model for perceptual learning. Trends in Cognitive Sciences. 2005;9(7):329–334. doi:10.1016/j.tics.2005.05.010.

10. Hubel DH, Wiesel TN. The period of susceptibility to the physiological effects of unilateral eye closure in kittens. Journal of Physiology. 1970;206:419–436.

11. Fahle M. Perceptual learning: Specificity versus generalization. Current Opinion in Neurobiology. 2005;15(2):154–160. doi:10.1016/j.conb.2005.03.010.

12. Ahissar M, Hochstein S. Task difficulty and the specificity of perceptual learning. Letters to Nature. 1997;387(6631):401–406. doi:10.1038/387401a0.

13. Fiorentini A, Berardi N. Perceptual learning specific for orientation and spatial frequency. Nature. 1980;287:43–44. doi:10.1038/287043a0.

14. Karni A, Sagi D. Where practice makes perfect in texture discrimination: evidence for primary visual cortex plasticity. Proceedings of the National Academy of Sciences of the United States of America. 1991;88(11):4966–4970. doi:DOI 10.1073/pnas.88.11.4966.

15. Ball K, Sekuler R. A specific and enduring improvement in visual motion discrimination. Science. 1982;218(4573):697–698. doi:10.1126/science.7134968.

16. Lu ZL, Hua T, Huang CB, Zhou Y, Dosher BA. Visual perceptual learning. Neurobiology of learning and memory. 2011;95(2):145–51. doi:10.1016/j.nlm.2010.09.010.

17. Zednik C. Mechanisms in cognitive science. In: Glennan S, Illari P, editors. The Routledge Handbook of Mechanisms and Mechanical Philosophy. Pylyshyn 1980. Oxon: Routledge; 2016. p. 389–400.

18. Adini Y, Sagi D, Tsodyks M. Context-enabled learning in the human visual system. Nature. 2002;415(February):2–5.

19. Gilbert CD, Sigman M, Crist RE. The Neural Basis of Perceptual Learning Review. Neuron. 2001;31:681–697. doi:10.1016/S0896-6273(01)00424-X.

20. Dosher BA, Lu ZL. Perceptual learning reflects external noise filtering and internal noise reduction through channel reweighting. Proceedings of the National Academy of Sciences of the United States of America. 1998;95(23):13988–93. doi:10.1073/pnas.95.23.13988.

21. Dosher BA, Lu ZL. Hebbian Reweighting on Stable Representations in Perceptual Learning. Learning & Perception. 2009;1(1):37–58. doi:10.1556/LP.1.2009.1.4.

22. Dosher BA, Jeter PE, Liu J, Lu ZL. An integrated reweighting theory of perceptual learning. Proceedings of the National Academy of Sciences of the United States of America. 2013;110(33):13678–13683. doi:10.1073/pnas.1312552110.

23. Herzog MH, Fahle M. A recurrent model for perceptual learning. Journal of Optical Technology. 1999;66(9):836.

24. Poggio T, Fahle M, Edelman S. Fast perceptual learning in visual hyperacuity. Massachusetts Inst of Tech Artfical Intelligence Lab. 1991;.

25. Fahle M. Specificity of learning curvature, orientation, and vernier discriminations. Vision Research. 1997;37(14):1885–1895. doi:10.1016/S0042-6989(96)00308-2.

26. Schoups AA, Vogels R, Orban GA. Human perceptual learning in identifying the oblique orientation: retinotopy, orientation specificity and monocularity. The Journal of physiology. 1995;483.3:797–810.

27. Sowden PT, Rose D, Davies IRL. Perceptual learning of luminance contrast detection: Specific for spatial frequency and retinal location but not orientation. Vision Research. 2002;42(10):1249–1258. doi:10.1016/S0042-6989(02)00019-6.

28. Fahle M, Edelman S. Long-term learning in vernier acuity: Effects of stimulus orientation, range and of feedback. Vision Research. 1993;33(3):397–412. doi:10.1016/0042-6989(93)90094-D.

29. Fahle M, Edelman S, Poggio T. Fast perceptual learning in hyperacuity. Vision Research. 1995;35(21):3003–3013. doi:10.1016/0042-6989(95)00044-Z.

30. Shiu LP, Pashler H. Improvement in line orientation discrimination is retinally local but dependent on cognitive set. Perception & psychophysics. 1992;52(5):582–588. doi:10.3758/BF03206720.

31. Gilbert CD, Li W. Top-down influences on visual processing. Nature Reviews Neuroscience. 2013;14(5):350–363. doi:10.1038/nrn3476.

32. Mollon JD, Danilova MV. Three remarks on perceptual learning. Spatial vision. 1996;10(1):51–58. doi:10.1016/j.aqpro.2013.07.003.

33. Ahissar M, Hochstein S. The reverse hierarchy theory of visual perceptual learning. Trends in Cognitive Sciences. 2004;8(10):457–464. doi:10.1016/j.tics.2004.08.011.

34. Talluri BC, Hung SC, Seitz AR, Seriès P. Confidence-based integrated reweighting model of task-difficulty explains location-based specificity in perceptual learning. Journal of vision. 2015;15(10):17. doi:10.1167/15.10.17.

35. Herzog MH, Fahle M. Modeling perceptual learning: difficulties and how they can be overcome. Biological cybernetics. 1998;78(2):107–117. doi:10.1007/s004220050418.

36. Petrov AA, Dosher BA, Lu ZL. Perceptual learning without feedback in non-stationary contexts: Data and model. Vision Research. 2006;46(19):3177–3197. doi:10.1016/j.visres.2006.03.022.

37. Schoups AA, Vogels R, Qian N, Orban G. Practising orientation identification improves orientation coding in V1 neurons. Nature. 2001;412(6846):549–553. doi:10.1038/35087601.

38. Ren Z, Zhou J, Yao Z, Wang Z, Yuan N, Xu G, et al. Neuronal basis of perceptual learning in striate cortex. Nature Scientific Reports. 2016;6:24769. doi:10.1038/srep24769.

39. Yang T, Maunsell JHR. The Effect of Perceptual Learning on Neuronal Responses in Monkey Visual Area V4. Journal of Neuroscience. 2004;24(7):1617–1626. doi:10.1523/JNEUROSCI.4442-03.2004.

40. Zohary E, Celebrini S, Britten KH, Newsome WT. Neuronal Plasticity that Underlies Improvement in Perceptual Performance. Science. 1994;263(5151):1289–1292.

41. Jehee JFM, Ling S, Swisher JD, van Bergen RS, Tong F. Perceptual Learning Selectively Refines Orientation Representations in Early Visual Cortex. Journal of Neuroscience. 2012;32(47):16747–16753. doi:10.1523/JNEUROSCI.6112-11.2012.

42. Shibata K, Watanabe T, Sasaki Y, Kawato M. Perceptual learning incepted by decoded fMRI neurofeedback without stimulus presentation. Science (New York, NY). 2011;334(6061):1413–5. doi:10.1126/science.1212003.

43. Furmanski CS, Schluppeck D, E, Ngel SA. Learning strengthens the response of primary visual cortex to simple patterns. Current Biology. 2014;14(7):573–578. doi:10.1016/j.

44. Petrov AA, Dosher BA, Lu ZL. The dynamics of perceptual learning: An incremental reweighting model. Psychological Review. 2005;112(4):715–743. doi:10.1037/0033-295X.112.4.715.

45. Gu Y, Liu S, Fetsch CR, Yang Y, Fok S, Sunkara A, et al. Perceptual learning reduces interneuronal correlations in macaque visual cortex. Neuron. 2011;71(4):750–761. doi:10.1016/j.neuron.2011.06.015.

46. Ball K, Sekuler R. Direction-specific improvement in motion discrimination. Vision Research. 1987;27(6):953–965. doi:10.1016/0042-6989(87)90011-3.

47. Furlan M, Smith AT. Global Motion Processing in Human Visual Cortical Areas V2 and V3. Journal of Neuroscience. 2016;36(27):7314–7324. doi:10.1523/JNEUROSCI.0025-16.2016.

48. Lamme VAF. Recurrent corticocortical interactions in neural disease. Archives of Neurology. 2003;60(2):178–184. doi:10.1001/archneur.60.2.178.

49. Simoncelli EP, Heeger DJ. A Model of Neuronal Responses in Visual Area MT. Vision Research. 1998;38(5):743–761. doi:S0042698997001831 [pii].

50. DeValois RL, Albrecht DG, Thonrell LG. Spatial Frequency Selectivity of Cells in Macaque Visual Cortex. Vision research. 1982;22(5):545–559. doi:10.1016/0042-6989(82)90113-4.

51. Hubel DH, Wiesel TN. Receptive fields, binocular interaction and functional architecture in the cat's visual cortex. Journal of Physiology. 1962;160:106–154. doi:10.1523/JNEUROSCI.1991-09.2009.

52. Movshon JA, Thompson ID, Tolhurst DJ. Spatial and temporal contrast sensitivity of neurones in areas 17 and 18 of the cat visual cortex. J Physiol. 1978;283:101–120.

53. Livingstone M, Hubel DH. Segregation of Form, Color, Movement, and Depth: Anatomy, Physiology, and Perception. Science. 1988;240(4835):740–749.

54. Movshon JA, Newsome WT. Visual response properties of striate cortical neurons projecting to area MT in macaque monkeys. Journal of Neuroscience. 1996;16(23):7733–7741. doi:10.1167/9.8.752.

55. Owsley C. Aging and vision. Vision Research. 2011;51(13):1610–1622. doi:10.1016/j.visres.2010.10.020.

56. Yu C, Klein SA, Levi DM. Perceptual learning in contrast discrimination and the (minimal) role of context. Journal of Vision. 2004;4(2004):169–182. doi:10.1167/4.3.4.

57. Adini Y, Wilkonsky A, Haspel R, Tsodyks M, Sagi D. Perceptual learning in contrast discrimination: the effect of contrast uncertainty. Journal of vision. 2004;4(2004):993–1005. doi:10.1167/4.12.2.

58. Kuai SG, Zhang JY, Klein SA, Levi DM, Yu C. The essential role of stimulus temporal patterning in enabling perceptual learning. Nature Neuroscience. 2005;8(11):1497–1499. doi:10.1038/nn1546.

59. Xiao LQ, Zhang JY, Wang R, Klein SA, Levi DM, Yu C. Complete Transfer of Perceptual Learning across Retinal Locations Enabled by Double Training. Current Biology. 2008;18(24):1922–1926. doi:10.1016/j.cub.2008.10.030.

60. Zhang JY, Kuai SG, Xiao LQ, Klein SA, Levi DM, Yu C. Stimulus coding rules for perceptual learning. PLoS Biology. 2008;6(8):1651–1660. doi:10.1371/journal.pbio.0060197.

61. Zhang J, Zhang G, Xiao L, Klein SA, Levi DM, Yu C. Rule-Based Learning Explains Visual Perceptual Learning and Its Specificity and Transfer. Journal of Neuroscience. 2010;30(37):12323–12328. doi:10.1523/JNEUROSCI.0704-10.2010.

62. Felleman DJ, Van Essen DC. Receptive field properties of neurons in area V3 of macaque monkey extrastriate cortex. Journal of neurophysiology. 1987;57(4):889–920.

63. Hubel DH, Wiesel TN. Receptive fields and functional acrhitecture in two nonstriate visual areas (18 and 19) of the cat. Journal of neurophysiology. 1965;28(2):229–289.

64. Mikami A, Newsome WT, Wurtz RH. Motion selectivity in macaque visual cortex. II. Spatiotemporal range of directional interactions in MT and V1. Journal of neurophysiology. 1986;55(6):1328–39.

65. Sillito AM, Cudeiro J, Jones HE. Always returning: feedback and sensory processing in visual cortex and thalamus. Trends in Neurosciences. 2006;29(6):307–316. doi:10.1016/j.tins.2006.05.001.

66. Zeki SM. Functional organization of a visual area in the posterior bank of the superior temporal sulcus of the rhesus monkey. The Journal of Physiology. 1974;236(3):549–573. doi:10.1113/jphysiol.1974.sp010452.

67. Hung SC, Seitz AR. Prolonged Training at Threshold Promotes Robust Retinotopic Specificity in Perceptual Learning. Journal of Neuroscience. 2014;34(25):8423–8431. doi:10.1523/JNEUROSCI.0745-14.2014.

68. Watanabe T, Sasaki Y. Perceptual learning: Toward a comprehensive theory. Annual review of psychology. 2015;3(66):197–221. doi:10.1146/annurev-psych-010814-015214.Perceptual.

69. Wang R, Zhang JYJY, Klein SA, Levi DM, Yu C. Vernier perceptual learning transfers to completely untrained retinal locations after double training: a "piggybacking" effect. Journal of vision. 2014;14(13):12. doi:10.1167/14.13.12.

70. Cong Lj, Wang Rj, Yu C, Zhang JY. Perceptual learning of basic visual features remains task specific with Training-Plus-Exposure (TPE) training. Journal of Vision. 2016;16(3:13):1–9. doi:10.1167/16.3.13.

71. Asher JM, Hibbard PB. First- and second-order contributions to depth perception in anti-correlated random dot stereograms. Scientific Reports. 2018;8(14120):1–20. doi:10.1038/s41598-018-32500-4.

72. Hochstein S, Ahissar M. Hierarchies and reverse hierarchies in the Visual System. Neuron. 2002;36(3):791–804. doi:10.1016/S0896-6273(02)01091-7.

73. Ostwald D, Lam JM, Li S, Kourtzi Z. Neural Coding of Global Form in the Human Visual Cortex. Journal of Neurophysiology. 2008;99(5):2456–2469. doi:10.1152/jn.01307.2007.

74. Aaen-Stockdale C, Hess RF. The amblyopic deficit for global motion is spatial scale invariant. Vision Research. 2008;48(19):1965–1971. doi:10.1016/j.visres.2008.06.012.

75. Levi A, Shaked D, Tadin D, Huxlin KR. Is improved contrast sensitivity a natural consequence of visual training? Journal of Vision. 2015;14(10):1158–1158. doi:10.1167/14.10.1158.

76. Chen N, Cai P, Zhou T, Thompson B, Fang F. Perceptual learning modifies the functional specializations of visual cortical areas. Proceedings of the National Academy of Sciences. 2016;113(20):5724–5729. doi:10.1073/pnas.1524160113.

77. Newsome WT, Park B, York N, Brook S. A Selective Impairment of Motion Perception Following Lesions of the Middle Temporal Visual Area (MT). The Journal of Neuroscience. 1988;8(6).

78. Vaina LM, Belliveau JW, Roziers EBd, Zeffiro TA. Neural systems underlying learning and representation of global motion. Proceedings of the National Academy of Sciences. 1998;95(21):12657–12662. doi:10.1073/pnas.95.21.12657.

79. Amano K, Edwards M, Badcock DR, Nishida S. Spatial-frequency tuning in the pooling of one- and two-dimensional motion signals. Vision Research. 2009;49(23):2862–2869. doi:10.1016/j.visres.2009.08.026.

80. Bex PJ, Dakin SC. Comparison of the spatial-frequency selectivity of local and global motion detectors. Journal of the Optical Society of America A, Optics, image science, and vision. 2002;19(4):670–7. doi:10.1364/JOSAA.19.000670.

81. Burr D, Thompson P. Motion psychophysics: 1985-2010. Vision Research. 2011;51(13):1431–1456. doi:10.1016/j.visres.2011.02.008.

82. Nishida S. Advancement of motion psychophysics : Review 2001 – 2010. Science. 2011;11(2011):1–53. doi:10.1167/11.5.11.Introduction.

83. Garcia A, Kuai SG, Kourtzi Z. Differences in the time course of learning for hard compared to easy training. Frontiers in Psychology. 2013;4(MAR):1–8. doi:10.3389/fpsyg.2013.00110.

84. McGovern DP, Webb BS, Peirce JW. Transfer of perceptual learning between different visual tasks. Journal of Vision. 2012;12(11):4–4. doi:10.1167/12.11.4.

85. Braddick OJ. Segmentation versus integration in visual motion processing. Trends in Neurosciences. 1993;16(7):263–268. doi:10.1016/0166-2236(93)90179-P.

86. Adelson EH, Movshon JA. Phenomenal coherence of moving visual patterns. Nature. 1982;300(5892):523–525.

87. Wilson HR, Ferrera VP, Yo C. A Psychophysically Motivated Model for 2-Dimensional Motion Perception. Visual Neuroscience. 1992;9(1):79–97.

88. An X, Gong H, Qian L, Wang X, Pan Y, Zhang X, et al. Distinct Functional Organizations for Processing Different Motion Signals in V1, V2, and V4 of Macaque. Journal of Neuroscience. 2012;32(39):13363–13379. doi:10.1523/JNEUROSCI.1900-12.2012.

89. Li P, Zhu S, Chen M, Han C, Xu H, Hu J, et al. A Motion direction preference map in monkey V4. Neuron. 2013;78(2):376–388. doi:10.1016/j.neuron.2013.02.024.

90. Newsome WT, Britten KH, Movshon JA. Neuronal correlates of a perceptual decision. Nature. 1989;341(6237):52–54. doi:10.1038/341052a0.

91. Rudolph K, Pasternak T. Transient and permanent deficits in motion perception after lesions of cortical areas MT and MST in the macaque monkey. Cerebral Cortex. 1999;9(1):90–100. doi:10.1093/cercor/9.1.90.

92. Britten KH, Shadlen MN, Newsome WT, Movshon JA. The analysis of visual motion: a comparison of neuronal and psychophysical performance. The Journal of Neuroscience. 1992;12(12):4745–4765. doi:10.1.1.123.9899.

93. Rees G, Friston K, Koch C. A direct quantitative relationship between the functional properties of human and macaque V5. Nature Neuroscience. 2000;3(7):716–723. doi:10.1038/76673.

94. Cowey A, Campana G, Walsh V, Vaina LM. The role of human extra-striate visual areas V5/MT and V2/V3 in the perception of the direction of global motion: A transcranial magnetic stimulation study. Experimental Brain Research. 2006;171(4):558–562. doi:10.1007/s00221-006-0479-6.

95. Hedges JH, Gartshteyn Y, Kohn A, Rust NC, Shadlen MN, Newsome WT, et al. Dissociation of neuronal and psychophysical responses to local and global motion. Current Biology. 2011;21(23):2023–2028. doi:10.1016/j.cub.2011.10.049.

96. Lui LL, Bourne JA, Rosa MGP. Spatial and temporal frequency selectivity of neurons in the middle temporal visual area of new world monkeys (Callithrix jacchus). European Journal of Neuroscience. 2007;25(6):1780–1792. doi:10.1111/j.1460-9568.2007.05453.x.

97. Pasternak T, Merigan WH. Motion Perception following Lesions of the Superior Temporal Sulcus in the Monkey. Cerebral Cortex. 1994;May/June(4):247–259.

98. Li RW, Tran TT, Craven AP, Leung TW, Chat SW, Levi DM. Sharpening coarse-to-fine stereo vision by perceptual learning: asymmetric transfer across the spatial frequency spectrum. Royal Society open science. 2016;3(1):150523. doi:10.1098/rsos.150523.

99. Parker AJ. Binocular depth perception and the cerebral cortex. Nature Reviews Neuroscience. 2007;8(5):379–91. doi:10.1038/nrn2131.

100. Fleet DJ, Wagner H, Heeger DJ. Neural Encoding of Binocular Disparity: Energy Models, Position Shifts and Phase Shifts. Vision Research. 1996;36(12):1839–1857.

101. Hibbard PB, Goutcher R, Hunter DW. Encoding and estimation of first- and second-order binocular disparity in natural images. Vision Research. 2016;120:108–120. doi:10.1016/j.visres.2015.10.016.

102. Smallman HS, MacLeod DIA. Spatial scale interactions in stereo sensitivity and the neural representation of binocular disparity. Perception. 1997;26(8):977–994. doi:10.1068/p260977.

103. Tanaka H, Ohzawa I. Neural Basis for Stereopsis from Second-Order Contrast Cues. Journal of Neuroscience. 2006;26(16):4370–4382. doi:10.1523/JNEUROSCI.4379-05.2006.

104. Henriksson L, Nurminen L, Hyvarinen A, Vanni S, Hyvarinen A, Vanni S, et al. Spatial frequency tuning in human retinotopic visual areas. Journal of Vision. 2008;8(2008):1–13. doi:10.1167/8.10.5.Introduction.

105. Rockland KS, Knutson T. Feedback Connections From Area Mt Of The Squirrel-Monkey To Areas V1 And V2. Journal Of Comparative Neurology. 2000;425(3):345–368.

106. Hupé JM, James AC, Girard P, Lomber SG, Payne BR, Bullier J. Feedback connections act on the early part of the responses in monkey visual cortex. Journal of Neurophysiology. 2001;85(1):134–145. doi:11152714.

107. Pascual-Leone A, Walsh V. Fast Backprojections from the Motion to the Primary Visual Area Necessary for Visual Awareness. Science. 2001;292(5516):510–512. doi:10.1126/science.1057099.

108. Bridge H, Stagg CJ, Near J, Lau Ci, Zisner A, Cader MZ. Altered neurochemical coupling in the occipital cortex in migraine with visual aura. Cephalalgia. 2015;35(11):1025–1030. doi:10.1177/0333102414566860.

109. Romei V, Chiappini E, Hibbard PB, Avenanti A. Empowering Reentrant Projections from V5 to V1 Boosts Sensitivity to Motion. Current Biology. 2016;26(16):2155–2160. doi:10.1016/j.cub.2016.06.009.

110. Romei V, Thut G, Silvanto J. Information-Based Approaches of Noninvasive Transcranial Brain Stimulation. Trends in Neurosciences. 2016;39(11):782–795. doi:10.1016/j.tins.2016.09.001.

111. Chiappini E, Silvanto J, Hibbard PB, Avenanti A, Romei V. Strengthening functionally specific neural pathways with transcranial brain stimulation. Current Biology. 2018;28(13):R735–R736. doi:10.1016/j.cub.2018.05.083.

112. Dobres J, Watanabe T. Response feedback triggers long-term consolidation of perceptual learning independently of performance gains. Journal of Vision. 2012;12(8):9–9. doi:10.1167/12.8.9.

113. Herzog MH, Fahle M. The Role of Feedback in Learning a Vernier Discrimination Task. Vision Research. 1997;37(15):2133–2141. doi:http://dx.doi.org/10.1016/S0042-6989(97)00043-6.

114. Liu J, Lu ZL, Dosher BA. Mixed training at high and low accuracy levels leads to perceptual learning without feedback. Vision Research. 2012;61:15–24. doi:10.1016/j.visres.2011.12.002.

115. Brainard DH. The psychophysics toolbox. Spatial vision. 1997;10:433–436.

116. Kleiner M, Brainard D, Pelli D, Ingling A, Murray R, Broussard C, et al. What's new in Psychtoolbox-3. Perception. 2007;36(14):1.

117. Pelli DG. The VideoToolbox software for visual psychophysics: Transforming numbers into movies. Spatial vision. 1997;10(4):437–442.

118. Williams DW, Sekuler R. Coherent global motion percepts from stochastic local motions. Vision Research. 1984;24(1):55–62. doi:10.1016/0042-6989(84)90144-5.

119. Moscatelli A, Mezzetti M, Lacquaniti F. Modeling psychophysical data at the population-level: The generalized linear mixed model. Journal of Vision. 2012;12(11):26–26. doi:10.1167/12.11.26.

120. Agresti A. Categorical Data Analysis. 2nd ed. Wiley-Interscience; 2002.

121. Bolker BM, Brooks ME, Clark CJ, Geange SW, Poulsen JR, Stevens MHH, et al. Generalized linear mixed models: a practical guide for ecology and evolution. Trends in Ecology and Evolution. 2009;24(3):127–135. doi:10.1016/j.tree.2008.10.008.

122. Gold JI, Law CT, Connolly P, Bennur S. Relationships Between the Threshold and Slope of Psychometric and Neurometric Functions During Perceptual Learning: Implications for Neuronal Pooling. Journal of Neurophysiology. 2010;103(1):140–154. doi:10.1152/jn.00744.2009.

123. Swanson WH, Birch EE. Extracting thresholds from noisy psychophysical data. Perception & Psychophysics. 1992;51(5):409–422. doi:10.3758/BF03211637.

124. Ling S, Carrasco M. Sustained and transient covert attention enhance the signal via different contrast response functions. Vision Research. 2006;46(8-9):1210–1220. doi:10.1016/j.visres.2005.05.008.

125. Herrmann K, Montaser-Kouhsari L, Carrasco M, Heeger DJ. When size matters: Attention affects performance by contrast or response gain. Nature Neuroscience. 2010;13(12):1554–1561. doi:10.1038/nn.2669.

126. Donovan I, Carrasco M. Exogenous attention facilitates location transfer of perceptual learning. Journal of vision. 2015;15(10):1–16. doi:10.1167/15.10.11.doi.

127. Das A, Huxlin KR. New Approaches to Visual Rehabilitation for Cortical Blindness: Outcomes and Putative Mechanisms. The Neuroscientist. 2010;16(4):374–387. doi:10.1177/1073858409356112.

128. Das a, Tadin D, Huxlin KR. Beyond Blindsight: Properties of Visual Relearning in Cortically Blind Fields. Journal of Neuroscience. 2014;34(35):11652–11664. doi:10.1523/JNEUROSCI.1076-14.2014.

129. Hess RF, Mansouri B, Thompson B. A binocular approach to treating amblyopia: Antisuppression therapy. Optometry and Vision Science. 2010;87(9):697–704. doi:10.1097/OPX.0b013e3181ea18e9.

130. Huxlin KR. Perceptual plasticity in damaged adult visual systems. Vision Research. 2008;48(20):2154–2166. doi:10.1016/j.visres.2008.05.022.

131. Levi DM. Prentice award lecture 2011: Removing the brakes on plasticity in the amblyopic brain. Optometry and Vision Science. 2012;89(6):827–838. doi:10.1097/OPX.0b013e318257a187.

132. Melnick MD, Tadin D, Huxlin KR. Relearning to See in Cortical Blindness. The Neuroscientist. 2016;22(2):199–212. doi:10.1177/1073858415621035.

133. Bridge H, Thomas O, Jbabdi S, Cowey A. Changes in connectivity after visual cortical brain damage underlie altered visual function. Brain. 2008;131(6):1433–1444. doi:10.1093/brain/awn063.

134. Markov NT, Vezoli J, Chameau P, Falchier A, Quilodran R, Huissoud C, et al. Anatomy of hierarchy: Feedforward and feedback pathways in macaque visual cortex. Journal of Comparative Neurology. 2014;522(1):225–259. doi:10.1002/cne.23458.

135. Jeter PE, Dosher BA, Lu ZL. Task precision at transfer determines specificity of perceptual learning. Journal of Vision. 2009;9(3):1–13. doi:10.1167/9.3.1.Introduction.

136. Seitz AR, Nanez JE, Holloway S, Tsushima Y, Watanabe T. Two cases requiring external reinforcement in perceptual learning. Journal of Vision. 2006;6(9):966–973. doi:10.1167/6.9.9.

137. Liu J, Dosher BA, Lu ZL. Modeling trial by trial and block feedback in perceptual learning. Vision Research. 2014;99:46–56. doi:10.1016/j.visres.2014.01.001.

138. Kingdom FAA, Prins N. Psychophysics: a practical introduction. Amsterdam: Elsevier; 2010.

139. Zylberberg A, Roelfsema PR, Sigman M. Variance misperception explains illusions of confidence in simple perceptual decisions. Consciousness and Cognition. 2014;27(1):246–253. doi:10.1016/j.concog.2014.05.012.

140. Watanabe T, Náñez JE, Sasaki Y. Perceptual learning without perception. Nature. 2001;413(6858):844–848. doi:10.1038/35101601.

141. Watanabe T, Náñez JE, Koyama S, Mukai I, Liederman J, Sasaki Y. Greater plasticity in lower-level than higher-level visual motion processing in a passive perceptual learning task. Nature neuroscience. 2002;5(10):1003–1009. doi:10.1038/nn915.

142. Hawken MJ, Parker AJ, Lund JS. Laminar organization and contrast sensitivity of direction-selective cells in the striate cortex of the Old World monkey. Journal of Neuroscience. 1988;8(10):3541–3548.

143. Solgi M, Liu T, Weng J. A computational developmental model for specificity and transfer in perceptual learning. Journal of Vision. 2013;13:7. doi:10.1167/13.1.7.

144. Bejjanki VR, Beck JM, Lu ZL, Pouget A. Perceptual learning as improved probabilistic inference in early sensory areas. Nature Neuroscience. 2011;14(5):642–50. doi:10.1016/S0733-8619(03)00096-3.

145. Maniglia M, Seitz AR. Towards a whole brain model of Perceptual Learning. Current Opinion in Behavioral Sciences. 2018;20:47–55. doi:10.1016/j.cobeha.2017.10.004.

146. Aberg KC, Herzog MH. Interleaving bisection stimuli - randomly or in sequence - does not disrupt perceptual learning, it just makes it more difficult. Vision Research. 2009;49(21):2591–2598. doi:10.1016/j.visres.2009.07.006.

147. Cong LJ, Zhang JY. Perceptual learning of contrast discrimination under roving: The role of semantic sequence in stimulus tagging. Journal of Vision. 2014;14(13):1–8. doi:10.1167/14.13.1.

148. Lin Z, Murray SO. Priming of awareness or how not to measure visual awareness. Journal of Vision. 2014;14(2014):1–17. doi:10.1167/14.1.27.

149. Lin Z, Murray SO. Automaticity of Unconscious Response Inhibition : Comment on Chiu and Aron (2014). Journal of Experimental Psychology: General. 2015;144(1):244–254.

150. Lin Z, Dosher BA, Lu ZL. Mixture of easy trials enables transient and sustained perceptual improvements through priming and perceptual learning. Scientific Reports. 2017;7(1):7421. doi:10.1038/s41598-017-06989-0.