A comprehensive suite of measures to quantify cell shape and actin

organization

Running title: Shape and texture measures

Authors: Elaheh Alizadeh[1], Wenlong Xu[1], Jordan Castle[2], Jacqueline Foss[3,4], Ashok

Prasad[1,3,**] .

1- Department of Chemical and Biological Engineering, 2- Department of Biology, 3-School of Biomedical

Engineering, 4- Department of Mechanical engineering, Colorado State University, Fort Collins, CO, 80523.

**Corresponding author

KEY WORDS: Morphometrics, shape quantification, texture quantification, cell

shape, Cytoskeleton, Cancer.

Summary statement

14 We develop a set of quantitative measures for studying changes in the shape and

15 texture of cells, and experimentally validate their performance on osteosarcoma cells

16 perturbed with different cytoskeletal drugs.

Abstract

A number of recent studies have shown that cell shape and cytoskeletal texture can be
used as sensitive readouts of the physiological state of the cell. However, utilization of
this information requires the development of quantitative measures that can describe
relevant aspects of cell shape. In this paper we develop a set of quantitative measures,
some used previously and others that are new, that have desirable properties for shape
and texture quantification of cells. These measures, broadly classifiable into the
categories of textural, spreading and irregularity measures, are tested by using them to
discriminate between osteosarcoma cell lines treated with different cytoskeletal drugs.
We find that even though specific classification tasks often rely on a few measures, these
are not the same between all classification tasks, thus requiring the use of the entire suite
of measures for classification and discrimination. We provide detailed descriptions of the
measures, as well as codes to implement them. Image based quantitative analysis has the

potential to become a new field of biological data ("image-omics"), providing
quantitative insight into cellular processes.
32

Introduction

The shape of a cell spread on a substrate is determined by the balance between the

internal and external forces exerted on the cell boundary. The cell exerts forces and

responds to external forces, from the extra-cellular matrix (ECM) or from neighboring

cells, with the help of molecular motors and the cellular cytoskeleton, which is thus the

ultimate determinant of cell shape (Fletcher & Mullins 2010a),(Pollard & Cooper 2009).

The cytoskeleton is a complex network, made of three major kinds of filaments -- f-actin,

microtubules and intermediate filaments -- that form a cross-linked dynamic meshwork in

the cytoplasm, providing shape and structure to the cell (Fletcher & Mullins

2010b),(Ingber et al. 1994). The most dynamic constituent of the cytoskeleton, which is

especially important in force generation and motility, is the filamentous actin (f-actin)

network(Stricker et al. 2010). The f-actin network is directly involved in the formation of

lamellipodia and filopodia through polymerization of f-actin against the cell

membrane(Pollard & Borisy 2003). A third kind of cellular protrusions, blebs, are a result

of the cortical actin network detaching from the cell membrane(Charras et al. 2008), and

the convex shapes of adherent cells have been shown to result from myosin-II driven

actin contractility(Bischofs et al. 2008). Cellular adhesion is maintained by an actin

network that is often organized into contractile bundles called stress fibers(Pellegrin &

Mellor 2007), which are classified into four types, dorsal and ventral fibers, transverse

arcs and the perinuclear actin cap(Tojkander et al. 2012; Levy et al. 2010). Dorsal stress

fibers are stabilizing structures associated with the cell edge, while the transverse arcs

and the ventral fibers are major contractile structures, and the perinuclear actin cap

regulates the shape of the nucleus(Tojkander et al. 2012).

The f-actin network is also deeply involved in force generation, force sensing and

mechanotransduction. Contractile forces generated by myosin motors within cytoskeletal

networks, membrane extension caused by actin polymerization, changes in osmotic

pressure by opening of water or ion channels are examples of internal forces that play a

role in shape of a cell. External forces leading to shape changes are applied through

neighboring cells or ECM (Paluch & Heisenberg 2009). Actin filaments can generate

and also resist mechanical stresses and cell deformation. But they can also eventually

reorganize and change their structure, thereby sometimes relaxing external stresses.

Different mechanical properties of the cell cytoskeleton and ECM will lead to different

shapes for the cell.

Thus the f-actin network is primarily responsible for the shape acquired by the

adherent cell. It follows that the structure of the f-actin network must be related to the

global shape of the cell, though the exact relation between the two is likely to be complex

and non-linear.

Cell shape has been investigated as a marker and predictor of cellular phenotype

and behavior. In our previous work, we used Zernike moments and geometric parameters

as a measure of cell shape to distinguish between high metastatic and low metastatic

osteosarcoma cancer cell lines with 99% accuracy(Alizadeh et al. 2016; Lyons et al.

2016). Other groups have also reported that cell shape can predict tumor grade (Sailem &

Bakal 2017), changes in the nuclear/cytoplasmic ratio of NFkB (Sero et al. 2015), and

YAP (Yes-associated protein), (Sero & Bakal 2017), chemosensitivity of human colon

cancer cell lines(Pasqualato et al. 2013) ,(Pasqualato et al. 2012), differences in motility

(Keren et al. 2008), forms of motility (Cooper et al. 2015), the progress of the Epithelial

to Mesenchymal transition (EMT) (Ren et al. 2015) and the differentiation of Human

Mesenchymal Stem Cells (hMSCs) into osteoblasts (Matsuoka et al. 2013).

Given the importance of the actin cytoskeleton in determining the shape of the cell

and the nucleus however, the structure and organization of the f-actin network may

provide additional information that can improve the prediction of cell physiology and

better distinguish between cells in different states. There is a case to be made, therefore,

of including measures of actin organization into studies of cell shape and its relation with
cell states, especially as actin staining is often used as the primary means to determine the
shape of the cell. Actin staining also provides textural information, which is directly
related with actin structure, and as we show below, addition of this textural information
significantly improves the discrimination between cell types.

The importance of cell and nuclear morphology and cytoskeletal organization as a
tool for understanding and predicting cell behavior raises the need to appropriately
quantify the shape and cytoskeletal texture of a cell. Good quantitative measures, which
we call cellular morphometrics, should be capable of capturing biologically important
differences between different experimental conditions. However, not all measures will be
optimal for each comparison, and thus one will have to begin with a larger set of
quantitative measures and discard uninformative ones if needed. Furthermore, good
measures must not show significant differences between technical replicates in the same
experiment. Based on these arguments we chose to test and develop cellular
morphometrics on cells treated with pharmacological modulators of the cytoskeleton.
These quantitative measures also meet the criteria proposed for good morphometrics by
Pincus et. al. (Pincus & Theriot 2007), i.e., that shape measures should possess fidelity,
capture biologically important variation and be meaningful and interpretable.

Experimental Methods
For this study DUNN and DLM8 osteosarcoma cancer cells were used. The DLM8 line is
derived from the DUNN cell line with selection for metastasis(Asai et al. 1998).
Therefore, DLM8 is closely related to DUNN except for degree of its invasiveness. Both
cell lines were a gift from Dr. Douglas Thamm (Colorado State University, CO, USA).
They were cultured on glass detergent washed and air dried (GDA) substrates with
standard culture conditions of $37^{o}C$ and 5% carbon dioxide concentration in Dulbecco's
Modified Eagle Medium (DMEM) (Sigma Alrdich). DMEM was supplemented with
10% EquaFETAL Fetal Bovine Serum (Atlas Biologicals) and 100 Units/ml penicillin

with 100 μg/ml streptomycin (Fisher Scientific-Hyclone). After 45 hours of culture, cells

were incubated with different cytoskeletal drugs with description, conditions, and

vendors listed in the **Table 1** for 3 hours. Then the cells were washed and fixed with 4%

paraformaldehyde. Finally, they were fluorescently stained for nuclei (DAPI from

Molecular probes) and actin (Acti-stain 488 phalloidin from Cytoskeleton, Inc). All the

drugs were dissolved in Dimethyl sulfoxide (DMSO) (Fisher BioReagents™) and to drop

its effect on cell shape and actin structure control study was also treated with DMSO with

the same molarity as other drugs. Then the cells were imaged using fluorescent

microscopy. Representative images of each cell line treated with these drugs are shown in

Figure 1.

Image processing

In order to strike a balance between the throughput and the accuracy of the image

processing, the image processing is fully supervised by the operator to reduce the number

of artifacts and also automated as much as possible to speed-up the processing. The

image processing code is stream-lined into a consecutive step-by-step workflow which

consists of four steps as follows. 1) A graphic user interface (GUI) enables the binary

thresholding of the actin and nucleus images. While a thresholding value is suggested

automatically by Otsu's method, users can easily adjust the thresholding value using a

slide bar in the GUI by visually checking the original image and the thresholded image

displayed side-by-side in the GUI. 2) Cell declustering in the thresholded images is done

using an optimized template of the open-source software CellProfiler (Kamentsky et al.

2011). 3) The outputs from the CellProfiler are then visually examined by the operator

and corrections can be made if necessary using the modules functionalized into this step.

To facilitate the following analyses, each cell is centered and saved separately into one

1024x1024 image. Except for CellProfiler, all the other image processing codes were

programmed in-house using Matlab (Mathworks) and are available upon requests. Also

available are the detailed protocol on how to configure the image processing codes and

the CellProfiler template used for cell declustering.

Data analysis methods

Quantifying changes in multidimensional shape space

Each cell can be represented as a point in multidimensional shape space. For ease

of analysis this vector can be projected down to a lower-dimensional space of the first

few principal components (PCs). Differences between different conditions or different

cell lines can be represented by p-value between their shape distributions for each PC

space. While doing multiple comparisons, we will often pick the most informative

principal component for our analysis from among the first four PCs. This is done in the

following manner. We pick the PC whose worst case, i.e. largest p-value among all

comparisons, is better (i.e. smaller p-value) than that of any other PC, so that it is the best

single measure for distinguishing between all the comparisons. In other words, for each

principal component maximum p-value between all comparisons is calculated as Eqn 1.

$$\text{MaxP}_{(PC1)} = \text{Max } \{\text{p-value}_{(PC1,\ 1)}\ ,\ \dots,\ \text{p-value}_{(PC1,\ n)}\} \qquad (1)$$

152   Where p-value $_{(PC1,\ n)}$ is the p-value for n^{th} comparison in the first principal component.

153   Then from the first four PCs, the one that has the smallest MaxP, maximum p-value

154   between different comparisons, is chosen. In each shape category, we choose the best

155   principal component by this criteria, and will describe them as the Primary Principal

156   Components (PPC) of each shape quantification measure or category, for each analysis.

Pearson correlation

Pearson correlation coefficient is calculated between features as shown in Eqn 2.

Here r is Pearson correlation coefficient, x_i is the feature x for i^{th} sample, y_i is the feature y

for i^{th} sample, μ_x is the average value of the feature x for all of the samples, and μ_y is the

average value of the feature y for all of the samples. This coefficient is calculated for all

14 cell lines and drug combinations (2 cell lines x 7 drugs) and the averaged coefficient is

recorded. The results are shown in heat map plots in **Figure S.** 1. In these plots the

diagonal elements are correlation between features with themselves, so they are all 1.

Also, Pearson correlation coefficient is symmetric so each heat map plot will be

symmetric as well.

$$r = \frac{\sum_{i=1}^{n}(x_i - \mu_x)(y_i - \mu_y)}{\sqrt{\sum_{i=1}^{n}(x_i - \mu_x)^2}\sqrt{\sum_{i=1}^{n}(y_i - \mu_y)^2}} \qquad (2)$$

167   **Results**

168   **Developing Shape quantifiers**

The actin structure is obtained from pixel intensities of labelled actin, which we can

represent by a greyscale image (**Figure 2A**). These greyscale images of cells, or images

of stained nuclei, are converted into a binary image (**Figure 2B**) and the coordinates of

the edges are recorded (**Figure 2C**), yielding the cell boundary. To extract features of the

morphology from this numerical data, we introduce three classes of shape measures,

described as textural, spreading, and irregularity classes. Textural measures are divided

into three subcategories called band based measurements, gray scale fractal dimension,

and gray scale measures which are calculated based on the intensity plots of actin, **Figure**

2A. Spreading measures, which include measures based on a Zernike moment

representation of shape as well as subcategories involving basic geometric parameters

such as area and perimeter, are extracted from the binary representation of the

cell(**Figure 2B**), its convex hull, or a similar image of the nuclei. Irregularity measures

include parameters such as the waviness and roughness, which use information regarding

the pixels at the boundary of the cell, **Figure 2C**. Each of these shape categories have

been described below.

Textural measures

Band based measurements

186   This parameter is sensitive to changes in the radial symmetry of the distribution of

actin filaments. We divide the image into 10 equally spaced concentric regions (Δr)

around the center of mass of the image, called bands, as shown in **Figure S. 2**. Five

quantifiers are used to measure the differences in actin distribution between these bands.

First, average intensity for each band is calculated. Then indices of the bands which have

the lowest or the highest average intensity and their value are recorded. The last

measurement is what is called above average adjusted intensity of the bands which is

formulated in **Table S**. 1 along with other band based measures(J H Slater et al. 2015).

Band based measures are especially important in characterizing the actin

distribution of the cells treated with Cytochalasin D, where actin has very unique

symmetrical distribution. In these cells, dense foci are formed around the nucleus. The

bands located in the central region of the cells are void of actin. At the outer bands, short

linear actin structures aligned along the radius are observed.

Gray scale fractal dimension

Fractal dimension (FD) is a measure of roughness of objects, and can be applied to

the characterization of texture in engineered and natural images. For planar images the

fractal dimension lies between 2 and 3. There are many methods to calculate this

measure, and we chose the box counting method for its inherent simplicity. In this

method the binary image is first covered with an evenly spaced grid with side length of e.

Then, the number of boxes which cover the fractal image are counted. This process is

repeated by decreasing the side length and the FD is calculated based on Eqn 3

$$\text{FD}=\lim_{e\to 0} \frac{\log N(e)}{\log\left(\frac{1}{e}\right)} \tag{3}$$

Here e is the size of each box in the grid and the variable N ($\square$) is the number of the

boxes which contain the fractal.

We calculate FD based on a binarization of the greyscale image using edge

detection methods to identify actin voids. This is done using four different edge detection

methods in Matlab. To provide an example, binarized images of a cell from DUNN cell

line treated with Cytochalasin D using these four edge detection methods are shown in

Figure S. 3. We found that different edge detection methods pick up different aspects of

the actin structure, and so for every cell their gray scale image is binarized using these

four methods, and their FD calculated (Costa 2013).

Other gray scale measures

Haralik et, al introduced a procedure for quantifying the texture of satellite images

based on the spatial relation between the gray tone of neighboring pixels in an image.

They used these measures for classifying the satellite images(Haralick et al. 1973). In this

method the gray level co-occurrence matrix (GLCM, sometimes called Spatial-

Dependence Matrix) is calculated for gray scale images. For an image that has intensities

of 1, 2, …, g, the co-occurrence matrix is a g x g matrix such that its ijth element is the

number of the times that a pixel in the image has intensity equal to "i" and the pixel at a

pre-defined distance $\vec{d}$ (which we choose to be 1 pixel) from it has intensity of j. An

example of a 4 pixels x 4 pixels image with 5 gray tone levels is shown in **Figure S.** 4**.**

In a GLCM matrix when there are very few dominant transitions in gray tone of an

image in neighboring pixels, the matrix will have small value for all of the entries. Each

diagonal element of a GLCM matrix represents the number of the times that gray tone

does not change in the neighboring pixel. Large diagonal elements imply that the image

is homogenous. Large numbers in the far upper right and far lower left of the matrix

implies large transitions in intensity and high contrast in an image. After calculating the

GCLM matrix, shown for the example image in **Figure S.** 4**C**, 23 different measures are

calculated to quantify the texture in an image. The list of measures are tabulated in **Table**

S. 2, (Haralick et al. 1973), (Soh et al. 1999), (Uppuluri 2010). For example one of the

parameters is the contrast. For interpretation of this measure, it is useful to keep in mind

that the contrast in an image is proportional to the changes in gray tone, n=|i-j|, so the far

upper right and far lower left which have bigger value of n will have bigger contribution

to the contrast parameter. For homogenous images, the diagonal entries will be large, and

will have a bigger contribution to the homogeneity measure.

In this study we calculate the GLCM matrix for the vector $\vec{d}$ equal to 1 pixel in

magnitude and with directions of 0^o, 45^o, 90^o, and 125^o and then the average value for

each parameter is reported. This process yields rotation invariant measures.

Spreading measures

In this class of shape quantifiers, the binary image of cell and nuclei is used. These

measures include geometric parameters for nuclei and cell, convex hull measures and

Zernike moments for cells.

Geometric measures

Parameters like area, perimeter, major axis of fitted ellipse, minor axis of fitted

ellipse, and their ratio are examples of geometric measures. All geometric measures used

in this study to quantify cell and nuclei's shapes are listed in **Table S4** and **Table S5**.

Convex Hull measurements

A convex polygon is a closed polygon such that the connecting line between any

pair of points inside the polygon, lies completely inside the polygon. **Figure S. 5A** and **B**

shows an example of a convex and non-convex polygon. The convex hull of a 2D shape

is the smallest convex polygon that encloses the whole shape (Welzl & Gärtner 2014).

An example of the convex hull of a 2D cell shape is shown in **Figure S. 5C**. Convex hull

geometric measures used in this study are listed in **Table S6.**

Zernike moments

To calculate Zernike moments, the image of the cell is projected to the Zernike

Polynomial basis function and the coefficients, called Zernike moments, are used as

descriptor of cell shape. Zernike moments are complex numbers whose magnitude is

rotation invariant, and can be made displacement invariant by moving the cell to the

center of the imaging frame such that the center of mass of the image falls on the

geometric center of the frame. The procedure has been discussed extensively previously

(Alizadeh et al. 2016),and Zernike moments have been used previously in other types of

biological image recognition (Tahmasbi et al. 2011).

Irregularity of boundary measures

In addition to cell spreading, irregularities in the boundary also carries information

about the state of the cell. For example an irregular border could arise due to a large

number of filopodia in the cell, which is a signature of a highly dynamic cytoskeleton. A

highly contractile cell may retract from focal adhesions at the boundary creating many

membrane protrusions that increase variability. To quantify irregularity of the cell

boundary, the 2D boundary of the cell is used as shown in **Figure 2C**. We use the pixel

positions that mark the boundary of the cell to calculate waviness, which estimates the

periodic variation in the boundary, and roughness, which measures the non-periodic

variation in the boundary, as discussed below.

Waviness measures

Using a Fourier series, a signal can be expanded in terms of linear combinations of

orthogonal basis functions of sines and cosines with increasing frequencies. In the

Fourier series expansion, it is assumed that the input signal is periodic. The outline of a

cell is a closed curve, hence it is counted as periodic signal. The boundary of the cell

could be represented in Cartesian coordinates, x and y, or polar coordinates, ρ and θ.

Regardless of coordinate selection, it will lead to two independent signals which can be

separately written as linear combination of cos and sin basis functions as Eqn 4.

$$f(x) = C_{x,0} + \sum_{n=1}^{nPixel} A_{x,n} \cos(wnx) + B_{y,n} \sin(wnx) \qquad (4)$$

Where nPixel is the number of the pixels in the boundary of a cell and $\frac{w}{2\pi}$ is

fundamental frequency which is equal to $\frac{1}{2\pi * nPixel}$. The variable n is an integer number

and n*w is the nth frequency in the decomposition. Variables A_n and B_n are amplitudes

of n^{th} frequency, $C_{x,0}$ is the mean of the signal, and $f(x)$ is the input signal.

If we use all the frequencies, we can reconstruct the cell. But the number of the

descriptors will be too high. There is a tradeoff between the number of frequencies (shape

descriptors) used and the accuracy of the reconstruction. We are interested in quantifying

shape without dealing with high number of parameters. Since the amplitude decreases

with increasing frequency, we can filter higher frequencies and just use lower frequencies

to decrease the number of descriptors used for the analysis. Here, with qualitative

analysis of reconstructed cells with different frequency we decided to use only the first 35

frequencies as descriptor of the cell. Reconstruction of shape of a few cells using these 35

frequencies is excellent, as shown in **Figure 3.**

However, a remaining issue for the Fourier coefficients is that they are not

rotationally invariant. We can construct a rotationally invariant measure by using Eqn 5,

which removes the phase difference in the Fourier expansion.

$$C_n = \sqrt{A_n^2 + B_n^2} \qquad (5)$$

This also reduces the number of parameters by half. Even though reconstruction of

cell shape will not be possible with the rotationally invariant measures, our results show

that they are nevertheless useful parameters for distinguishing between cells with

different degree of variations in the radius. We will refer to these parameters as Waviness

parameters. Interestingly, for Fourier parameters, all the features show low correlation

coefficient with each other (~0.35), suggesting that each coefficient carries mostly

independent information. This observation was expected since we use orthogonal basis

functions.

Roughness

As explained earlier, in the Fourier series decomposition of shape, higher

frequencies have small amplitude and we neglect them. However, we can still obtain

information about small amplitude variations from those high frequency terms. Following

the method of Villanueva et. al., (Villanueva 2015) to account for high frequency

measures, we reconstruct the cell with the 35 Fourier components and subtract it from the

original signal. Now we can calculate the roughness measures from the remainder of the

signal. These parameters are also highly correlated with each other and are listed in
Table S. 7.

Validating performance of developed measures on osteosarcoma cells perturbed with different cytoskeletal drugs.

Drugs which directly perturb microtubules, myosin-II or actin leave a unique signature on actin structure which is quantifiable by textural measures.

We found that the most dramatic and unique effects on cell shape and texture arise
from drugs that directly perturb microtubules or actin filaments. As shown in **Fig. 1**,
control cells acquire a polygonal shape on surfaces. Depolymerizing actin using
Cytochalasin D leads to rounded cells where the actin has re-organized into alternating
concentric rings of high and low density. Within each ring the actin is structured in very
unique radial streaks! Increasing the depolymerization rate of microtubules using
Nocodazole leads to a small rounded morphology. Inhibiting myosin II activity and
decreasing cell's contractility using Blebbistatin leads to increase in irregularities in the
cell boundary. These changes in shape and structure of a cell are similar for both DUNN
and DLM8 cell lines. Changes in structure and morphology of the cells using other drugs
are subtle and not easily identifiable by eye.

To quantify changes in shape and structure of the cells perturbed with different
drugs, gray scale and binary image of the cells along with information of the boundary of
the cells were used to measure all features in the 9 shape categories detailed above for
each cell. **Table 2** demonstrates quantified changes in all the measures for all the drugs.
As demonstrated in this table the changes in textural measures for all the drugs are
significant. This means that all the drugs that perturbed actin either directly, or indirectly
through perturbing other cytoskeletal components, lead to significant changes in actin
cytoskeleton organization.

Actin reorganization changes the spreading measures for both the cell and the nucleus

341
342

To explore changes in 2D shape of a cell and nuclei by changing actin distribution
we compare the PPC of the geometric measures of the cell and nuclei. As shown in **Table**
2 perturbing actin significantly changes cell geometric measures other than DLM8 cell
line treated with Jasplakinolide. Interestingly changes in actin structure not only changes
cell geometric measures, but it also changes nuclei geometric measures for all conditions
other than both cell lines treated with Blebbistatin and DUNN cell line treated with
Jasplakinolide and PP2 drugs.

Zernike moments and Convex Hull parameters are similar in some respects.
Although their quantification method is very different, as shown in reconstruction of the
image of cells using Zernike moments (**Figure S. 6**) both measures ignore the
irregularities and fine fluctuations in the boundary. As demonstrated in **Table 2** changes
in the PPC for both measures are not distinguishable for DUNN cell lines treated with
PP2, and both cell lines treated with FAKI 14. Moreover, hull geometric measures do not
change significantly for DLM8 cells treated with Cytochalasin-D and DUNN cells treated
with Jasplakinolide. In addition, Zernike moments do not change significantly for the
DLM8 cells treated with Blebbistatin and DUNN cells treated with Cytochalasin-D. All
other drugs lead to distinguishable changes in convex hull and Zernike moment
measures.

Since many cell shape parameters are expected to be correlated with each other,
we performed a correlation analysis of all the measures within each shape quantification
category, which are shown in **Figure S.** 7. These results indicate that Zernike moments,
cell geometric measures, nuclei geometric measures, and convex hull geometric measures
are highly correlated with each other.

Cytoskeletal reorganization leads to changes in irregularities of a cell's boundary

As shown in **Table 2**, waviness measures for all the drug conditions, except for DUNN cell lines treated with Jasplakinolide or PP2 and both cell lines treated with FAKI 14, changes significantly with respect to controls. In addition, the roughness measures change significantly for both cell lines treated with Blebbistatin, Cytochalasin-D, FAKI 14, and Nocodazole. In both cell lines, Jasplakinolide does not change the roughness measure significantly, which is also the case for the DUNN cell lines treated with PP2.

Different categories of shape quantifiers represent partly non-redundant shape information

While we have shown that each of our major categories and subcategories serve as good shape measures, in that they can be used to look for interpretable shape changes in different drug conditions, it is not clear whether we need all of them to represent shape. In order to estimate the degree of redundant information carried by the different shape categories, we calculated the Pearson correlation coefficient between all the features from different shape categories (shown in **Table S8**). In general, features from two different shape categories are weakly correlated (<0.4) except for Convex Hull and Cell Geometric features which are highly correlated with each other. There are a few other specific exceptions for which the features are also highly correlated. They are as follows. Zernike moments with $n<16$ and $m=0$ are highly correlated with Area. The correlation coefficient between Cell Area and Zernike Moment (0_0) is 1. It decreases with increasing order till it is almost zero for Zernike Moment (22_0), then it becomes negative and increases in magnitude (**Figure S.** 7). The coefficient C_0 from waviness features has a correlation of 1 with the Mean Cell Radius. It is also highly correlated with Cell Area (0.93) and Convex Hull area. Since the Fourier decomposition is based on a radial representation of the cell shape, C_0 is a measure of the average cell radius, and should be expected to show these high correlations. Fractal Dimension measures also have high correlation with Cell Geometric, Nuclei Geometric, Gray Scale measures, and Zernike moments with $m=0$

and n<17. However, apart from these specific cases, the relatively high number of weak

correlations between different shape categories implies that these shape categories

contain non-redundant information about cell shape. Thus, quantitative shape analysis

should ideally be carried out with representations from all of these shape categories in

order for most efficient discrimination between different experimental conditions.

Discussion:

In this paper we introduce and provide the tools to quantify cell shape and

cytoskeletal structure based on two dimensional images of cell morphology and actin

structure. The Matlab toolbox used to process the images and quantify the shape and

structure of a cell is shared in GitHub repository to be used by others

(https://github.com/Elaheh-Alizadeh/Quantifiction-of-shape-and-structure),

(https://github.com/Wenlong-Xu/Image_Processing_Cell_Shape). This work is among

the few that quantifies structure of the actin. Using statistically significant differences

between experimental conditions in principal component space as our measure, we

explore the capacity of these quantitative measures in capturing biologically important

information. We perturb the cytoskeleton of the cells with different drugs and explore

their effect on cell shape and its structure. We first use textural measures to verify that

actin structure of the cell changes in the cells treated by cytoskeletal drugs and then we

explore changes in the cell's and nuclei's 2D shape and irregularities of the cell boundary

accompanied by changes in actin structure. Here we showed that most of the drugs used

in this study directly or indirectly lead to significant changes in actin structure. Then we

explored effect of changes in structure of the cell on irregularities of cell boundary, nuclei

spreading, and cell spreading. In most of the cases changes in actin structure are

accompanied with significant changes in irregularities of cell boundary, and cell and

nuclei spreading. The results show that textural measures and spreading measures are

related but their relation is not simple, and the two classes of measures carry non-

redundant information. It is worth mentioning that although we implemented textural

measures to quantify actin structure, it can be used to quantify other cell structures as

well. These shape quantification methods presented in this paper will prove useful for

computer aided diagnostics in diseases such as cancer that are associated with

cytoskeletal perturbations, assessment of qualitative cellular changes in different

experimental conditions, and for mechanistic understanding of the determination of cell

shape. Textural measures introduced in this paper can also be used to quantify the texture

of image of slices of tumors rather than single cells. The advantage of using these

measures is that the results are more consistent, with a lower probability of human error

and inconsistency. Furthermore these methods are fast and do not require expensive

training.

Acknowledgement:

We gratefully acknowledge support from an NSF CAREER grant (PHY-1151454) to A.

P.

List of Abbreviations:

Dimethyl Sulfoxide (DMSO)

Dulbecco's Modified Eagle Medium (DMEM)

Epithelial to Mesenchymal transition (EMT)

Extra-cellular Matrix (ECM)

Glass Detergent washed and Air dried (GDA)

Gray Level Co-occurrence Matrix (GLCM)

Human Mesenchymal Stem Cells (hMSCs)

Primary Principal Components (PPC)

Principal Components (PCs)

Figure 1: Representative images of DUNN and DLM8 osteosarcoma cancer cell lines with different drug treatments. Blue color represents nuclei and green color represents actin cytoskeleton.

Figure 2: Representation of cell shape. A. Gray scale images using labeled actin. Intensity of each pixel is recorded and represented as a greyscale intensity plot. B. 2D outline: Position of each pixel in the boundary is recorded in polar or Cartesian coordinate.

Cell boundary
FS_Reconstruction

Figure 3: Reconstruction of the cells using Fourier decomposition. The red line is the actual boundary of the cell and black line is the reconstruction of the cell using the first 35 terms in the Fourier series expansion.

458   **Table 1: Description of the drugs used to perturb cytoskeleton.**

Drug	Blebbistatin	Cytochalasin D	Jasplakinolide	FAKI 14	Nocodazole	PP2
Target	Inhibits non-muscle myosin II activity	Inhibits actin polymerization	1. Promotes actin polymerization 2. stabilizes actin filaments	Inhibits focal adhesion kinase	Depolymerizing microtubules	Inhibits Src kinase
Application in cancer therapy	Inhibits cancer invasion(Nelson et al. 2011)	Used as chemotherapeutic agent[(Trendowski, Zoino, et al. 2015; Trendowski, Mitchell, et al. 2015).	Reduces lung metastases of systemic Lewis lung carcinoma(Takeuchi et al. 1998).	Showed anti-metastatic, anti-neoplastic and anti-angiogenic properties (Kang et al. 2015; Gabarra-Niecko et al. 2003; McLean et al. 2003; Hecker & Gladson 2003).	Has chemotherapeutic properties by arresting cells in mitosis(Endo et al. 2010).	Prevention of cancer metastasis(Nam et al. 2002).
Effect	1.Reduces F-actin contractility 2.Increase in cell	Decreases cell contractility	Stabilizes F-actin(Posey & Bierer 1999)	Reduces cell motility and focal adhesion	Cell shapes which lack tails and long protrusions	1. Induces strong cell-cell contact. 2.Enhances E-cadherin/catenin

	spreading and 3.Decreases cell motility(Sero et al. 2015)			turnover(Schlaepfer et al. 1999).	and decreased cell asymmetry(Rodionov et al. 1993).	expression which are strongly associated with the actin cytoskeleton(Nam et al. 2002).
Concentration (uM)	10	1	0.01	1	5	1
Vendor	Santa Cruz Biotechnology	Calbiochem	Calbiochem	Santa Cruz Biotechnology	Sigma Alrdich	Calbiochem

Table 2: Significance of changes in primary principal component (PPC) of
morphometric classes for different drugs according to T-test analysis. The p-value of
the null hypothesis that the experimental condition has the same mean as the control is
reported for each class of morphometrics, based on the PPC. There are significant
changes (p-value < 1E-3) in some actin textural measures (Band-Based, Fractal
Dimension and Gray scale measures), spreading measures (Cell Geometric, Nuclei
Geometric, Hull Geometric and Zernike moment) and Irregularity measures (Waviness
and Roughness) for every experimental condition. Sample size is 300 cells for each cell
line and drug condition other than DLM8 FAKI 14 which has a sample size of 200. In
this table, P-values<1E-9 are highlighted in dark green, 1E-9 <P-values<1E-6 are
highlighted in light green, and 1E-6 <P-values<1E-3 are highlighted in yellow.

Comparisons: Control vs	Cell Line	Band Based	Fractal Dimension	Gray Scale	Cell Geometric	Nuclei	Hull Geometric	Zernike Moment	Waviness	Roughness
Blebbistatin	DUNN	1.00E-07	2.60E-14	0.008	1.60E-78	0.006	3.10E-14	3.00E-21	2.00E-35	2.70E-45
	DLM8	3.00E-08	6.30E-45	6.00E-09	6.00E-25	0.004	7.90E-08	0.0018	2.00E-12	5.30E-16
Cytochalasin D	DUNN	3.00E-50	2.10E-31	2.00E-23	3.50E-21	2.00E-22	2.50E-06	0.0092	5.00E-10	3.30E-27
	DLM8	4.00E-31	2.10E-13	1.00E-23	4.50E-15	1.00E-18	0.05285	1.00E-05	3.00E-07	8.20E-07
FAKI 14	DUNN	2.00E-09	5.16E-31	1.00E-23	5.20E-09	2.00E-04	0.04184	0.1194	0.1398	2.20E-12
	DLM8	4.00E-04	6.60E-22	2.00E-21	1.50E-06	2.00E-06	0.07864	0.0997	0.3125	7.80E-07
Jasplakinolide	DUNN	2.00E-04	2.10E-13	9.00E-05	1.30E-08	0.001	0.02999	4.00E-05	0.0036	0.051
	DLM8	0.003	2.80E-09	2.00E-04	0.00207	5.00E-04	1.20E-05	0.0005	0.0001	0.02117
Nocodazole	DUNN	6.00E-13	5.60E-31	3.00E-25	1.30E-28	1.00E-04	4.60E-10	9.00E-07	1.00E-12	3.30E-16
	DLM8	0.001	9.80E-14	4.00E-07	1.20E-14	6.00E-06	3.00E-09	6.00E-14	3.00E-21	5.30E-13
PP2	DUNN	2.00E-21	1.40E-11	2.00E-10	2.50E-06	0.003	0.03602	0.0243	0.0089	0.00387
	DLM8	4.00E-07	0.00035	3.00E-09	0.00043	8.00E-10	2.70E-09	4.00E-13	4.00E-11	0.00078

Bibliography

Alizadeh, E. et al., 2016. Measuring systematic changes in invasive cancer cell shape
using Zernike moments. *Integr. Biol.*, 8(11), pp.1183–1193. Available at:
http://xlink.rsc.org/?DOI=C6IB00100A [Accessed January 8, 2017].

Asai, T. et al., 1998. Establishment and characterization of a murine osteosarcoma cell
line (LM8) with high metastatic potential to the lung. *International Journal of*
Cancer, 76(3), pp.418–422.

Bischofs, I.B. et al., 2008. Filamentous network mechanics and active contractility
determine cell and tissue shape. *Biophysical journal*, 95(7), pp.3488–96. Available
at: http://www.ncbi.nlm.nih.gov/pubmed/18599642 [Accessed January 5, 2017].

Charras, G.T. et al., 2008. Life and times of a cellular bleb. *Biophysical Journal*, 94(5),
pp.1836–1853.

Cooper, S. et al., 2015. Apolar and polar transitions drive the conversion between
amoeboid and mesenchymal shapes in melanoma cells. *Molecular biology of the*
cell,   26(22),   pp.4163–70.   Available   at:
http://www.ncbi.nlm.nih.gov/pubmed/26310441 [Accessed February 14, 2017].

Costa, A., 2013. Hausdorff (Box-Counting) Fractal Dimension. Available at:
https://www.mathworks.com/matlabcentral/fileexchange/30329-hausdorff--box-
counting--fractal-dimension.

De, J. & Villanueva, 2015. DESIGN AND IMPLEMENTATION OF A
CALCULATION PROGRAM IN MATLAB PLATFORM FOR EVALUATION OF
ROUGHNESS.   Available   at:
https://upcommons.upc.edu/bitstream/handle/2117/85844/Master's Thesis JdV.pdf.

Endo, K. et al., 2010. Nocodazole induces mitotic cell death with apoptotic-like features
in Saccharomyces cerevisiae. *FEBS Letters*, 584(11), pp.2387–2392. Available at:

http://www.sciencedirect.com/science/article/pii/S0014579310002991.

Fletcher, D.A. & Mullins, R.D., 2010a. Cell mechanics and the cytoskeleton. *Nature*,
463(7280), pp.485–92. Available at:
http://www.nature.com/doifinder/10.1038/nature08908 [Accessed August 21, 2017].

Fletcher, D.A. & Mullins, R.D., 2010b. Cell mechanics and the cytoskeleton. *Nature*,
463(7280), pp.485–492.

Gabarra-Niecko, V., Schaller, M.D. & Dunty, J.M., 2003. FAK regulates biological
processes important for the pathogenesis of cancer. *Cancer metastasis reviews*,
22(4), pp.359–74. Available at: http://www.ncbi.nlm.nih.gov/pubmed/12884911
[Accessed June 22, 2017].

Haralick, R., SHANMUGAM, K. & ITS'HAK, D., 1973. Textural Features for Image
classification. *IEEE Transactions on Systems, Man, and Cybernetics*, SMC-3(6),
pp.610–621.

Hecker, T.P. & Gladson, C.L., 2003. Focal adhesion kinase in cancer. *Frontiers in
bioscience : a journal and virtual library*, 8, pp.s705-14. Available at:
http://www.ncbi.nlm.nih.gov/pubmed/12700131 [Accessed June 22, 2017].

Ingber, D.E., Sun, Z. & Betensky, H., 1994. E , Cytoskeletal ~ Ec ~ an ~ Cs ~ Control in
Angiogene. *Science*, 7(12).

Kamentsky, L. et al., 2011. Improved structure, function and compatibility for
cellprofiler: Modular high-throughput image analysis software. *Bioinformatics*,
27(8), pp.1179–1180.

Kang, C.G. et al., 2015. Rho-associated kinase signaling is required for osteopontin-
induced cell invasion through inactivating cofilin in human non-small cell lung
cancer cell lines. *Bioorganic & medicinal chemistry letters*, 25(9), pp.1956–60.
Available at: http://linkinghub.elsevier.com/retrieve/pii/S0960894X15002206

[Accessed June 22, 2017].

Keren, K. et al., 2008. Mechanism of shape determination in motile cells. *Nature*,
453(7194), pp.475–480.

Levy, J.R. et al., 2010. Cell adhesion: integrating cytoskeletal dynamics and cellular
tension. *Molecular Cell*, 11(9), pp.633–643.

Lyons, S.M. et al., 2016. Changes in cell shape are correlated with metastatic potential in
murine and human osteosarcomas. *Biology Open*.

Matlab, Properties of gray-level co-occurrence matrix. Available at:
https://www.mathworks.com/help/images/ref/graycoprops.html?searchHighlight=glc
m&s_tid=doc_srchtitle.

Matsuoka, F. et al., 2013. Morphology-Based Prediction of Osteogenic Differentiation
Potential of Human Mesenchymal Stem Cells. *PLoS ONE*, 8(2).

McLean, G.W., Avizienyte, E. & Frame, M.C., 2003. Focal adhesion kinase as a potential
target in oncology. *Expert opinion on pharmacotherapy*, 4(2), pp.227–34. Available
at: http://www.tandfonline.com/doi/full/10.1517/14656566.4.2.227 [Accessed June
22, 2017].

Nam, J. et al., 2002. Src family kinase inhibitor PP2 restores the E-cadherin/catenin cell
adhesion system in human cancer cells and reduces cancer metastasis. *Clinical*
Cancer Research, 8(7), pp.2430–6. Available at:
http://www.ncbi.nlm.nih.gov/pubmed/12114449.

Nelson, T. et al., 2011. The myosin II inhibitor blebbistatin more potently inhibits 3-D
cell invasion than 2-D cell migration in a human metastatic tumor cell. *Cancer*
Research, 71(18 Supplement), pp.C1--C1. Available at:
http://cancerres.aacrjournals.org/content/71/18_Supplement/C1.

Paluch, E. & Heisenberg, C.-P., 2009. Biology and Physics of Cell Shape Changes in
Development. *Current Biology*, 19(17), pp.R790–R799. Available at:
http://linkinghub.elsevier.com/retrieve/pii/S0960982209014511 [Accessed August
21, 2017].

Pasqualato, A. et al., 2012. Quantitative shape analysis of chemoresistant colon cancer
cells: Correlation between morphotype and phenotype. *Experimental Cell Research*,
318(7), pp.835–846. Available at: http://dx.doi.org/10.1016/j.yexcr.2012.01.022.

Pasqualato, A. et al., 2013. Shape in migration: Quantitative image analysis of migrating
chemoresistant HCT-8 colon cancer cells. *Cell Adhesion and Migration*, 7(5),
pp.450–459.

Pellegrin, S. & Mellor, H., 2007. Actin stress fibres. *Journal of Cell Science*, 120(20),
pp.3491–3499. Available at: http://jcs.biologists.org/content/120/20/3491.abstract.

Pincus, Z. & Theriot, J.A., 2007. Comparison of quantitative methods for cell-shape
analysis. *Journal of Microscopy*, 227(2), pp.140–156. Available at:
http://doi.wiley.com/10.1111/j.1365-2818.2007.01799.x [Accessed March 19, 2017].

Pollard, T.D. & Borisy, G.G., 2003. Cellular Motility Driven by Assembly and
Disassembly of Actin Filaments Cellular Motility Driven by Assembly and
Disassembly of Actin Filaments. *Cell*, 112(4), pp.453–465.

Pollard, T.D. & Cooper, J.A., 2009. Actin, a Central Player in Cell Shape and Movement.
Science, 326(5957). Available at:
http://science.sciencemag.org/content/326/5957/1208.full [Accessed August 20,
2017].

Posey, S.C. & Bierer, B.E., 1999. Actin stabilization by jasplakinolide enhances
apoptosis induced by cytokine deprivation. *The Journal of biological chemistry*,
274(7), pp.4259–4265.

Ren, Z.-X. et al., 2015. Suitable parameter choice on quantitative morphology of A549 cell in epithelial - mesenchymal transition. *Bioscience reports*, pp.1–7. Available at: http://www.ncbi.nlm.nih.gov/pubmed/25899751.

Rodionov, V.I. et al., 1993. Microtubule-dependent Control of Cell Shape and Pseudopodial Activity Is Inhibited by the Antibody to Kinesin Motor Domain. , 123(6), pp.1811–1820. Available at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2290860/pdf/jc12361811.pdf.

Sailem, H.Z. & Bakal, C., 2017. Identification of clinically predictive metagenes that encode components of a network coupling cell shape to transcription by image-omics. *Genome research*, 27(2), pp.196–207. Available at: http://www.ncbi.nlm.nih.gov/pubmed/27864353 [Accessed February 14, 2017].

Schlaepfer, D.D., Hauck, C.R. & Sieg, D.J., 1999. Signaling through focal adhesion kinase. *Progress in biophysics and molecular biology*, 71(3–4), pp.435–78. Available at: http://www.ncbi.nlm.nih.gov/pubmed/10354709 [Accessed June 22, 2017].

Sero, J.E. et al., 2015. Cell shape and the microenvironment regulate nuclear translocation of NF-κB in breast epithelial and tumor cells. *Molecular systems biology*, 11(3), p.790. Available at: http://www.ncbi.nlm.nih.gov/pubmed/26148352 [Accessed February 28, 2017].

Sero, J.E. & Bakal, C., 2017. Multiparametric Analysis of Cell Shape Demonstrates that β-PIX Directly Couples YAP Activation to Extracellular Matrix Adhesion. *Cell Systems*, 4(1), p.84–96.e6. Available at: http://linkinghub.elsevier.com/retrieve/pii/S2405471216304100 [Accessed February 14, 2017].

Slater, J.H. et al., 2015. Recapitulation and Modulation of the Cellular Architecture of a User-Chosen Cell of Interest Using Cell-Derived, Biomimetic Patterning. *ACS Nano*,

9(6), pp.6128–6138. Available at: https://www.ncbi.nlm.nih.gov/pubmed/25988713.

Slater, J.H. et al., 2015. Recapitulation and Modulation of the Cellular Architecture of a

User-Chosen Cell of Interest Using Cell-Derived, Biomimetic Patterning. *ACS Nano*,

9(6), pp.6128–6138. Available at:

http://pubs.acs.org/doi/abs/10.1021/acsnano.5b01366 [Accessed February 23, 2017].

Soh, L., Tsatsoulis, C. & Member, S., 1999. Texture Analysis of SAR Sea Ice Imagery

Using Gray Level Co-Occurrence Matrices Texture Analysis of SAR Sea Ice

Imagery Using Gray Level Co-Occurrence Matrices. *Ieee Transactions on*

Geoscience and Remote Sensing, 37(2), pp.780–795. Available at:

http://citeseerx.ist.psu.edu/viewdoc/download;jsessionid=3CCA532A27525D5E952

3557CD011F596?doi=10.1.1.224.3585&rep=rep1&type=pdf%5Cnhttp://citeseerx.is

t.psu.edu/viewdoc/summary?doi=10.1.1.224.3585.

Stricker, J., Falzone, T. & Gardel, M., 2010. Mechanics of the F-actin Cytoskeleton.

Journal of Biomechanics, 43(1), pp.247–253.

Tahmasbi, A., Saki, F. & Shokouhi, S.B., 2011. Classification of benign and malignant

masses based on Zernike moments. *Computers in Biology and Medicine*, 41(8),

pp.726–735.

Takeuchi, H. et al., 1998. Jasplakinolide: interaction with radiation and hyperthermia in

human prostate carcinoma and Lewis lung carcinoma. *Cancer chemotherapy and*

pharmacology, 42(6), pp.491–496.

Tojkander, S., Gateva, G. & Lappalainen, P., 2012. Actin stress fibers - assembly,

dynamics and biological roles. *Journal of Cell Science*, 125(8), pp.1855–1864.

Available at: http://jcs.biologists.org/cgi/doi/10.1242/jcs.098087.

Trendowski, M., Mitchell, J.M., et al., 2015. Chemotherapy with cytochalasin congeners

in vitro and in vivo against murine models. *Investigational New Drugs*, 33(2),

pp.290–299. Available at: http://www.ncbi.nlm.nih.gov/pubmed/25563824
[Accessed June 22, 2017].

Trendowski, M., Zoino, J.N., et al., 2015. Preparation, In Vivo Administration, Dose-
Limiting Toxicities, and Antineoplastic Activity of Cytochalasin B. *Translational*
Oncology, 8(4), pp.308–317. Available at:
http://www.ncbi.nlm.nih.gov/pubmed/26310377 [Accessed June 22, 2017].

Uppuluri, A., 2010. GCLM feature calculation. Available at:
https://www.mathworks.com/matlabcentral/fileexchange/22354-glcm-features4-m--
vectorized-version-of-glcm-features1-m--with-code-changes-?s_tid=prof_contriblnk.

Welzl, E. & Gärtner, B., 2014. Computational Geometry. , pp.16–29. Available at:
https://www.ti.inf.ethz.ch/ew/Lehre/CG13/index.html.

 SUPPLEMENTARY INFORMATION

Figure S. 1 Pearson correlation coefficient within each shape category.

Within each shape category features are highly correlated other than band based,

waviness (Fourier), and some of the nuclei measures. Each pixel is the average of

pearson correleation for 14 cases, 7drugs x 2 cell lines.

644 **Figure S. 2: Band based division of textural image of a cell.** Cell is divided into 10

645 equally spaced radial bands. All the measures calculated based on this representation is

646 listed in **Table S**. 1.

Figure S. 3: Matlab's Edging methods used to binerize gray scale image of actin.

Four different methods of Roberts, Sobel, Canny, and zerocross are used in this paper.

Figure S. 4: Calculation of GCLM matrix. A) A cartoon of a 4pixels x 4pixels image.

B) The image has 5 gray tones as shown. C) Numerical representation of image based on

gray tone is shown. D) GCLM Matrix is 5 x 5 element and its ijth element is number of

the times which pixel with intensity of j is in distance   from the pixels with intensity of

i.   is an orbitrary vector but should be defined in advance. In this paper   is difined to

have length of one and it is defined in 4 directions of 0^0, 45^0, 90^0, and 135^0. Gray scale

measures are calculated based on four directions and their average is reported.

Figure S. 5: Convex Hull. A cartoon of a A) convex and B) non-convex polygon. A

convex polygon is a polygon which any line connecting two points inside the polygon

falls inside the polygon. C. Convex hull surrounding a cell.

Figure S. 6: Reconstruction of cell images for two different shaped cells. A. Original
images. B.   Reconstructed images using Zernike moments up to order (from left to the
right) 10, 20, 30, 40, 50.

Figure S. 7: Pearson correlation coefficient between Zernike Moments with m=0 and Cell Area. Correlation is maximum and positive with Zernike moment of 0_0, it decreases with increasing the order and it is almost 0 for Zernike moment of 22_0 and it is negative for higher orders and increases in magnitude. Each data point is the average of pearson correleation for 14 cases, 7drugs x 2 cell lines.

684 **Table S. 1: Band based measurements used in the paper**(John H. Slater et al. 2015).

Parameter		Formula
Max band intensity	Intensity of the band with maximum average intensity	$\text{Max} \left(\{ I_{B_1}, ..., I_{B_{10}} \} \right)$
Max band index	Location of the band with maximum average intensity	The n such that $I_n = \text{Max} \left(\{ I_{B_1}, ..., I_{B_{10}} \} \right)$
Minimum band intensity	Intensity of the band with minimum average intensity	$\text{Min} \left(\{ I_{B_1}, ..., I_{B_{10}} \} \right)$
Index of minimum band intensity	Location of the band with minimum average intensity	The m such that $I_m = \text{Min} \left(\{ I_{B_1}, ..., I_{B_{10}} \} \right)$
Above average adjusted intensity	Adjusted, weighted sum of average actin intensities greater than the mean intensity.	$\dfrac{\sum_{k=1}^{10} s\left(I_{B_k}\right) I_{B_k}}{\text{Max} \left(\{ I_{B_1}, ..., I_{B_{10}} \} \right) \sum_{k=1}^{10} s\left(I_{B_k}\right)}$ Where $S(I_{B_k}) =$ $\left\{ \begin{matrix} 1 & if\ I_{B_k} \geq mean\left(\{ I_{B_1}, ..., I_{B_{10}} \}\right) \\ 0 & otherwise \end{matrix} \right\}$

686   **Table S. 2: Gray scale measures used in this paper** (Matlab n.d.)**,** (Haralick et al.

687   1973)**,** (Soh et al. 1999)**,** (Uppuluri 2010)**.** See **Supplementary Table S. 3** for the

688   parameters used in this table.

Parameter	Formula
Autocorrelation	$\displaystyle\sum_{i=1}^{N_g}\sum_{j=1}^{N_g} ij\, p(i,j)$
Contrast_M(Matlab)	$\displaystyle\sum_{i=1}^{N_g}\sum_{j=1}^{N_g} \lvert i-j\rvert^2\, p(i,j)$
Correlation_M(Matlab)	$\displaystyle\sum_{i=1}^{N_g}\sum_{j=1}^{N_g} \frac{(i-\mu i)(j-\mu j)p(i,j)}{\sigma_i\sigma_j}$
Correlation_P	$\displaystyle\frac{\sum_{i=1}^{N_g}\sum_{j=1}^{N_g}(ij)p(i,j)-\mu_x\mu_y}{\sigma_x\sigma_y}$
Cluster Standard deviation	$\displaystyle\sum_{i=1}^{N_g}\sum_{j=1}^{N_g}\big(i+j-\mu_x \\ -\mu_y\big)^2 p(i,j)$
Cluster Prominence	$\displaystyle\sum_{i=1}^{N_g}\sum_{j=1}^{N_g}\big(i+j-\mu_x \\ -\mu_y\big)^4 p(i,j)$
Cluster Shade	$\displaystyle\sum_{i=1}^{N_g}\sum_{j=1}^{N_g}\big(i+j-\mu_x \\ -\mu_y\big)^3 p(i,j)$

Dissimilarity	$$\sum_{i=1}^{N_g}\sum_{j=1}^{N_g}\{|i-j|p(i,j)\}$$
Energy	$$\sum_{i=1}^{N_g}\sum_{j=1}^{N_g}p(i,j)^2$$
Entropy	$$-\sum_{i=1}^{N_g}\sum_{j=1}^{N_g}p(i,j)\log(p(i,j))$$
Homogeneity_M(Matlab)	$$\sum_{i=1}^{N_g}\sum_{j=1}^{N_g}\frac{p(i,j)}{1+|i-j|}$$
Homogeneity_P	$$\sum_{i=1}^{N_g}\sum_{j=1}^{N_g}\frac{p(i,j)}{1+(i-j)^2}$$
Maximum probability	$$MAX_{i,j}\,p(i,j)$$
Sum squares variance	$$\sum_{i=1}^{N_g}\sum_{j=1}^{N_g}(i-\mu)^2\,p(i,j)$$
Sum_average	$$\sum_{i=2}^{2N_g}i\,p_{x+y}(i)$$
Sum variance	$$\sum_{i=2}^{2N_g}(i-\text{SumEntropy})^2\,p_{x+y}(i)$$
Sum Entropy	$$-\sum_{i=2}^{2N_g}p_{x+y}(i)\log(p_{x+y}(i))$$

Difference Variance	$$\sum_{i=0}^{N_g-1} i^2 p_{x-y}(i)$$
Difference Entropy	$$-\sum_{i=0}^{N_g-1} p_{x-y}(i) \log\left(p_{x-y}(i)\right)$$
Information Measure of Correlation1	$$\frac{Entropy - HXY1}{\max\{HX, HY\}}$$
Information measure Correlation 2	$$\sqrt{\left(1 - e^{-2(HXY2-HXY)}\right)}$$
Inverse Difference Normalized(INN) (Matlab)	$$\sum_{i=1}^{N_g}\sum_{j=1}^{N_g} \frac{p(i,j)}{1 + \frac{\|i-j\|}{N_g}}$$
Inverse Difference Moment Normalized(Matlab)	$$\sum_{i=1}^{N_g}\sum_{j=1}^{N_g} \frac{p(i,j)}{1 + \frac{(i-j)^2}{N_g^2}}$$

691 **Table S. 3: Parameters used in the definitions of gray scale measures in the**

692 **Supplementary Table S. 2.**

Parameter	Formula
P(i,j)	ijth entry of co-occurance probability matrix
N_g	Highest intensity of gray level
μ	Mean value of p(i,j)
$p_x(i)$	$\sum_{j=1}^{N_g} p(i,j)$
$p_y(i)$	$\sum_{i=1}^{N_g} p(i,j)$
μ_x	$\sum_{i=1}^{N_g}\sum_{j=1}^{N_g} i\, p(i,j)$
μ_y	$\sum_{i=1}^{N_g}\sum_{j=1}^{N_g} j\, p(i,j)$
σ_x^2	$\sum_{i=1}^{N_g}\sum_{j=1}^{N_g} (i-\mu_x)^2\, p(i,j)$
σ_y^2	$\sum_{i=1}^{N_g}\sum_{j=1}^{N_g} (i-\mu_y)^2\, p(i,j)$
$p_{x+y}(k)$	$\sum_{i=1}^{N_g}\sum_{j=1}^{N_g} p(i,j)$

	Where i+j=k, k=2,..,$2N_g$
$p_{x-y}(k)$	$$\sum_{i=1}^{N_g}\sum_{j=1}^{N_g} p(i,j)$$ Where i-j=k, k=0,.., N_g-1
HX	$$-\sum_{i=1}^{N_g} p_x(i)\log\left(p_x(i)\right)$$
HY	$$-\sum_{i=1}^{N_g} p_y(i)\log\left(p_y(i)\right)$$
HXY1	$$-\sum_{i=1}^{N_g}\sum_{j=1}^{N_g} p(i,j)\log\left(p_x(i)p_y(j)\right)$$
HXY2	$$-\sum_{i=1}^{N_g}\sum_{j=1}^{N_g} p_x(i)p_y(j)\log\left(p_x(i)p_y(j)\right)$$

694   **Table S. 4: Geometric parameters.**

Parameter	Description	Unit
Cell area	Area of the cell	Pixel^2
Cell perimeter	Perimeter of the cell	Pixel
Cell major axis	The major axis of an ellipse drawn around the cell	Pixel
Cell minor axis	The minor axis of an ellipse drawn around the cell	Pixel
Cell circularity (Smoothness)	$\dfrac{4\pi\ Area}{Perimeter^2}$	Unit-less
Cell aspect Ratio	$\dfrac{Cell\ Major\ axis}{Cell\ Minor\ axis}$	Unit-less
Cell roundness	$\dfrac{4\ Area}{\pi\ Cell\ major\ axis\ ^2}$	Unit-less
Cell solidity	$\dfrac{Convex\ hull\ area}{Cell\ area}$	Unit-less
Max cell Radius	Maximum radius of the cell	Pixel
Cell radius ratio	$\dfrac{Max\ radius\ of\ the\ cell}{Min\ radius\ of\ the\ cell}$	Unit-less
Mean Cell Radius	The mean of all radii drawn from the hull's centroid to an exterior point	Pixel
CV Cell Radius	The relative variation of radii drawn from the cell's center to an exterior point.	Unit-less

	Given by the Standard of Deviation of all Radii divided by the mean of all radii	
Max Span	The maximum distance from one point on the convex hull to another	Pixel

697   **Table S. 5: Nuclei geometric parameters.**

Parameter	Description	Unit
Nuc area	Area of the nucleus	Pixel^2
Nuc perimeter	Perimeter of the nucleus	Pixel
Nuc major	Major axis of an ellipse drawn around the nucleus	Pixel
Nuc minor	Minor axis of an ellipse drawn around the nucleus	Pixel
Nuc circularity	$\dfrac{4\pi\;Nuc\;\;area}{Nuc\;perimeter^2}$	Unit-less
Nuc aspect ratio	$\dfrac{Nuc\;major}{Nuc\;minor}$	Unit-less
Nuc roundness	$\dfrac{4\;Nuc\;area}{\pi\;Nuc\;major^2}$	Unit-less
Nuc solidity	$\dfrac{Area\;of\;convex\;hull\;of\;nuclei}{Nuc\;area}$	Unit-less
Distance nuc to cell centroid	Distance between the centroids of cell and nucleus	Pixel
Nuc to cell orientation	Angle between the distance nuc to cell centroid and cell major axis	Degree

700 **Table S. 6: Convex hull geometric parameters.**

Parameter	Description	Unit
Hull area	Area of the convex hull	Pixel^2
Hull perimeter	Perimeter of the convex hull.	Pixel
Ratio of perimeter of Hull to cell	Ratio of hull perimeter to cell perimeter.	Unit-less
Hull circularity	$\dfrac{4\pi\ Hull\ area}{Hull\ perimeter^2}$	Unit-less
Max hull Radius	Maximum distance from centroid of Hull to an exterior point on the hull.	Pixel
Hull radius ratio	Maximum/minimum radius of hull.	Unit-less
Mean hull radius	The mean of all radii drawn from the hull's centroid to an exterior point.	Pixel
CV hull radius	$\dfrac{STD\ hull\ radius}{Mean\ hull\ radius}$	Unit-less
Bounding circle diameter	The diameter of the bounding circle drawn around the cell.	Pixel
Max circle to hull radius	The maximum distance from the center of the bounding circle to an edge of the convex hull.	Pixel

Circle to hull radius ratio	$\dfrac{\text{Max circle to hull radius}}{\text{Min circle to hull radius}}$	Unit-less
Mean circle to hull radius	The mean of all radii drawn from the circle's center to the hull	Pixel
CV_Circle2Hull_Radius	$\dfrac{STD(\text{ all radii drawn from the circle's center to the hull })}{\text{Mean circle to hull radius}}$	Unit-less

703 **Table S. 7: Roughness measurements.**

Parameter	Formula	Units
Mean of absolute values	$\mathrm{Ra} = \dfrac{1}{n} \sum\limits_{i=1}^{n} |y_i|$	Pixel
Root Mean Squared	$\mathrm{Ra} = \sqrt{\dfrac{1}{n} \sum\limits_{i=1}^{n} y_i^2}$	Pixel
Maximum height of the profile	$R_{yt} = R_p - R_v$	Pixel
Kurtosis	$R_{Ku} = \dfrac{1}{n R_q^4} \sum\limits_{i=1}^{n} y_i^4$	Pixel^4