Suppressive waves disambiguate the representation of long-range apparent

motion in awake monkey V1.

Chemla S[1], Reynaud A[1,3], di Volo M[2], Zerlaut Y[2,4], Perrinet L[1], Destexhe A[2,5] & Chavane F[1]*

[1] Institut de Neurosciences de la Timone (INT), UMR 7289 CNRS & Aix-Marseille Université, 27 Bd Jean

Moulin, 13385 Marseille Cedex 05, France

[2] Unité de Neurosciences, Information et Complexité , Centre National de la Recherche Scientifique, FRE

3693. 1 Avenue de la terrasse, 91198 Gif sur Yvette, France

[3] present address: McGill University, McGill Vision Research, Department of Ophthalmology, Montreal,

Quebec, Canada

[4] present address: Neural Coding laboratory, Center for Neuroscience and Cognitive Systems @UniTn,

Istituto Italiano di Tecnologia, Corso Bettini 31, 38068 Rovereto, Italy

[5] European Institute for Theoretical Neuroscience, 74 Rue du Faubourg Saint-Antoine, Paris, France

*Corresponding author: frederic.chavane@univ-amu.fr, phone: (33)491324033

Abstract

The "apparent motion" illusion is evoked when stationary stimuli are successively flashed in spatially separated positions. It depends on the precise spatial and temporal separations of the stimuli. For large spatiotemporal separation, the long-range apparent motion (lrAM), it remains unclear how the visual system computes unambiguous motion signals. Here we investigated whether intracortical interactions within retinotopic maps could shape a global motion representation at the level of V1 population in response to a lrAM. In fixating monkeys, voltage-sensitive dye imaging revealed the emergence of a spatio-temporal representation of the motion trajectory at the scale of V1 population activity, shaped by systematic backward suppressive waves. We show that these waves are the expected emergent property of a recurrent gain control fed by the horizontal intra-cortical network. Such non-linearities explain away ambiguous correspondence problems of the stimulus along the motion path, preformating V1 population response for an optimal read-out by downstream areas.

Introduction

When two stationary stimuli are successively flashed in spatially separated positions, it generates the so-called "apparent motion" illusion (Wertheimer 1912). This illusion, well characterized in psychophysics (Burr and Thompson 2011), depends on the spatio-temporal characteristics of the stimulus, being called "short-range" vs "long-range" apparent motion (lrAM) for spatial separation below or above 0.25° and temporal separation below or above 80 ms respectively (Braddick 1980). In psychophysics, intrinsic differences were reported between these two types of apparent motion, however, there is some debate whether it is underlined by same or different process (Cavanagh and Mather 1989). In physiology, while we have a clear idea on the neuronal processing generating direction-selective neuronal response to short-range apparent motion stimuli (Mikami, Newsome, and Wurtz 1986b), we still have a poor

understanding of how the visual system process lrAM. This is probably because the spatial separation

between individual strokes of the lrAM extend beyond the typical extent of receptive fields in the early

visual system, at least in primates. In the case of the lrAM, psychophysicists have long highlighted the

necessity to have a process, such as the "reviewing process" (Kahneman, Treisman, and Gibbs 1992),

that will link the transient apparitions of stimuli in different spatial and temporal positions in order to

generate a coherent motion percept of a single object, hereby solving the problem of "phenomenal

identity" (Ternus 1926) or "correspondence" (Ullman 1978). Downstream areas with large receptive

fields are a natural expected integration unit for such extended spatiotemporal input. Indeed, it has

been recently shown in human that the feedback from MT to V1 plays an important role in the

processing of lrAM (Wibral et al. 2009; Muckli et al. 2002; Vetter, Grosbras, and Muckli 2015), as well as

evidences of downstream activation along the ventral stream (Zhuo et al. 2003). However, it is still

unclear whether and how the "reviewing" process, needed to keep track of the object identity along the

motion trajectory, can be achieved within these receptive fields.

As suggested from fMRI experiments in human, the neuronal processing within V1 could

participate in formatting the representation of lrAM (Muckli et al. 2005). The extended precise

retinotopic map in V1 makes it indeed an ideal platform for representing and disambiguating, at the

level of the neuronal population, the trajectory of the apparent motion illusion, a representation that

could be read-out by downstream areas (Mumford 1991; Lee et al. 1998). In particular, V1 has the

highest resolution (Lee et al. 1998) to achieve the interactions in space and time needed to link the

individual strokes of the apparent motion (Lee et al. 1998; Adelson and Bergen 1985). In such context,

intra-cortical and inter-cortical connectivity would be the natural substrate to underlie the necessary

spatio-temporal interactions (Deco and Roland 2010; Muller et al. 2018). Importantly, these two

networks have intrinsically different spatio-temporal properties, the inter-cortical network operating

over very large extent but with poor spatial and temporal resolution (Angelucci et al. 2002; Stetter

2002), and the intra-cortical network has a more limited extent but with high spatial and temporal

resolution (Muller et al. 2014; Bringuier et al. 1999; Bullier 2001). Furthermore, they constitute the vast

majority of synaptic contacts in the cortex, the feedback accounting for less than 20% and the intra-

cortical connectivity contributing to 80% of the number of neuronal contacts, while the feedforward less

than 1% (Markov et al. 2011). Such connectivity seems therefore like a good candidate to link transient

spatio-temporal events (Muller et al. 2018). It was indeed shown, in the anesthetized cat, to shape

visual information for a dynamic representations of sequences of static stimuli (Jancke et al. 2004;

Gerard-Mercier et al. 2016) through non-linear gain controls of the feedforward input (Reynaud,

Masson, and Chavane 2012). However, it is still unclear whether and how the cortico-cortical

interactions could participate to shape the representation of lrAM within V1 retinotopic map in the

awake monkey.

To answer this question, we used optical imaging of voltage-sensitive dyes (VSDI) in the awake

fixating monkey, to measure how V1 neuronal population integrates a two-stroke lrAM that

overreached individual neuronal receptive field size. In response to a single stroke, activity in V1

propagates in space and time, as already documented (A. Grinvald et al. 1994; Slovin et al. 2002; Sato,

Nauhaus, and Carandini 2012; Bringuier et al. 1999; Muller et al. 2014), with spatial and temporal

constants that cover about 3 mm and 80 ms. In response to the lrAM of various spatio-temporal

separations, we observed that the cortical response systematically deviates from the linear prediction

and generates a wave of suppression that is initiated right at the second stimulus onset and propagates

to suppress the residual response to the first stimulus. A computational model was developed to

understand the potential origin of such suppressive waves. The model shows that two ingredients are

necessary to explain suppressive waves: the higher gain of inhibitory cells, and the shunting effect of the

associated synaptic conductances. Using a spatio-temporal decoding approach, we demonstrate that

such suppression waves explain away ambiguous representation of stimulus position along the apparent

motion trajectory. These waves thus preformat V1 population response for an unambiguous

representation of the lrAM. Using an opponent motion energy approach, we demonstrate that this

results in an optimal encoding of the stimulus velocity that could be easily read-out by downstream

areas.

Results

Characterizing the mesoscopic spatio-temporal impulse response function

Two-step apparent motion sequences of various spatio-temporal characteristics (Fig 1, A and B) were

presented to two behaving monkeys involved in a fixation task. The primary visual cortical response was

measured at the level of the population using voltage-sensitive dye imaging (Amiram Grinvald and

Hildesheim 2004; Chemla and Chavane 2010a). In response to a local stimulus (0.25° in diameter)

presented for 100 ms in two different visual positions (separated vertically by 1° or 2°), activity arises at

the retinotopic representation of these two positions and then spreads laterally over millimeters of

cortical surface (Fig.1C: lower position, Fig. 1D: upper position) (A. Grinvald et al. 1994; Reynaud,

Masson, and Chavane 2012; Muller et al. 2014). V1 activity is hereby reaching positions in space and

time well beyond 1° and 50ms. As a consequence, the evoked spread covers a large cortical extent that

can reach the representation of the other stimulus in space and beyond the inter-stimulus interval in

time. The space- and time- constants of our responses were systematically quantified on the two

monkeys and for the three stimulus durations we used (10, 50 and 100ms) on a 2D spatio-temporal (ST)

map (Fig. 2A). To produce these ST maps, cortical activity was averaged within the apparent-motion

trajectory (*dotted rectangle at frame 216 ms* in Fig. 1, C-G) to provide a unique spatial cortical dimension

(ordinate in Fig. 2A). First, we extracted the space-constant of a gaussian spatial fit for all time points

(see Fig. 2A, right-side of the maps). In both monkeys and across 19 sessions overall, the space-constant

increased from 1.6 +/- 0.5 mm at response onset to reach a maximum of 3.3 +/- 0.2 mm, independent of

the stimulus duration and monkeys (Fig. 2B, no significant difference observed between all stimuli

durations, t-test with p>0.01). The time-constants of the response time-course at the central

representation of the stimulus were measured using two halve gaussian functions fits (see Fig. 2A,

below the maps). In both monkeys, the time-constant at response onset was on average 23.6+/- 17.2

ms for all stimuli durations (except for monkey BR with a mean value of 44.5 +/- 14.5 ms for 100 ms

stimuli, see blue histogram in Fig. 2E), and 80 +/- 43.6 ms for response offset (Fig. 2F, no significant

difference observed between all stimuli durations, t-test with p>0.01). Lastly, we also extracted the

speed at which the response spreads across the cortical surface (see Fig. 2A, slanting lines) and obtained

a distribution with peak values of about 0.26 +/- 0.14 m/s, similar across monkeys and stimulus

durations (t-test with p>0.01), and similar to what has been observed in different species and states

(Slovin et al. 2002; Sato, Nauhaus, and Carandini 2012; Bringuier et al. 1999; Reynaud, Masson, and

Chavane 2012; Muller et al. 2014). This analysis showed that the spatio-temporal integrative properties

of the primary visual cortex are mostly independent of stimulus duration and are able to cover a large

spatial (3mm) and temporal (100ms) extent, bridging the cortical representation between our individual

stimuli in space and time.

Figure 1: Experimental protocol and time-sequence of the cortical response to the long-range apparent motion

*(lrAM). **A:** Two-step lrAM stimuli are presented to two awake fixating monkeys in their bottom left visual field,*

*while recording in their right visual cortex using VSDI. **B:** Spatio-temporal characteristics of lrAM stimuli, i.e.*

duration (DUR) , interstimulus interval (ISI) and spatial interval (SI), were varied to cover a [5-66.6]°/s range of

*speed. **C-E:** Cortical representation of evoked VSDI activity as a function of time, in response to respectively, a 100*

ms local stimulus in the down position, another one in the up position, and the sequence of these two stimuli (ISI =

50 ms and SI = 1°). The cortical area imaged is shown at upper left. The edge of the image color codes the

retinotopic borders as represented in A such as the vertical meridian (magenta), eccentricities (green and blue).

Scale bar: 2 mm; A: anterior, P: posterior, M: medial, L: lateral. Time in milliseconds after stimulus onset is shown at

*the top, while stimulation time is drawn at the bottom of each row (black lines). **F:** Activity pattern predicted by the*

linear combination in space and time of the response to stimulus 1 (row C) and the response to stimulus 2 (row D).

***G:** Suppression pattern obtained by subtracting the observed apparent motion response (row E) and the linear*

prediction (row F). Red contours delimit amplitude activity above a certain threshold: 1   ‰ in panels C-F and -

0.5‰ in panel G.

The evoked response to the lrAM is shaped by a suppressive wave

We next asked whether such lateral interactions contribute to shape the evoked population response to

the temporal succession of these two stimuli. For that purpose we measured the cortical population

response to a two-stroke upward apparent motion sequence (Fig. 1E). Such temporal sequence

generates a propagation of activity starting at the cortical representation of the first stimulus (S1) and

moving to the cortical representation of the second stimulus (S2), a cortical correlate of the illusory

motion (Jancke et al. 2004). The observed pattern of activity departs from the pattern predicted by a

simple linear summation of the lower and upper stimuli (Fig. 1F). If we subtract the observed (Fig. 1E)

and the linear predicted responses (Fig. 1F), two deviations from non-linearities are observed. First, a

suppression emerges at response onset and at the cortical representation of S2 (compare 1D and 1G at

frame 216ms). The suppression then gradually propagates over the cortical surface towards the

representation of S1 (Fig. 1G). We can hypothesize that the evoked activities by the two stimuli

composing the lrAM sequence interact together to generate this dynamic pattern of suppression. Since

the suppression is observed at the onset time of the response to S2, it has to be due to the activity

dynamics generated by S1 interacting with the integration of S2. However, the propagation of

suppression from the representation of S2 towards the representation of S1 is probably due to the

activity dynamics evoked by S2 interacting with the residual activity evoked by S1. Therefore, the

suppression wave could likely be the result of multiple interactions (e.g bidirectional) between the

activities evoked by the stimulus sequence.

Figure 2: Spatio-temporal characteristics of cortical responses to a local stimuli. A-C:*Spatio-temporal*

representations (ST) of the evoked cortical response to, respectively, 10 ms (A, red) , 50 ms (B, purple) and 100 ms

(C, blue) local stimuli. To produce the ST representation, we averaged spatial data along the stimulus trajectory

(rectangle in frame 216ms, Fig1C-G). For each spatial point, the temporal data were fitted to a combination of two

half Gaussians, as illustrated for one specific point in space (horizontal white line on the ST diagram) below the ST

maps. Similarly, for each time frame, the spatial data were fitted to a Gaussian function as shown on the right side

of each ST map for one specific point in time (vertical white line). ***D:*** *Space-constant of the Gaussian spatial fit*

(sigma parameter) plotted as a function of time for the three considered durations (10 ms in red, 50 ms in magenta

and 100 ms in blue) and for the two monkeys (top: monkey WA, bottom: monkey BR). ***E:*** *Histograms of time-*

constant at response onset (on) estimated from the temporal fit of the response for the three considered

durations and the two monkeys. ***F:*** *Histograms of time-constant at response offset (off) estimated from the*

temporal fit of the response for the three considered durations and the two monkeys. ***G:*** *Histograms of cortical*

speed of propagation estimated by linear regression on response latency (stairs-step contours, slanting lines and

slope of the linear regression) for the three considered durations and the two monkeys.

The suppressive wave is systematically observed

To better investigate how spreads of evoked activity and suppression shape the representation of lrAM,

we first show ST representations of examples taken for both monkeys and three stimuli speeds. The

example of Figure 1 is shown in Figure 3A (6.6°/s). In these ST representations, we can observe a clear

propagation of activity in response to a local stimulus (slanting lines in Fig. 3, A and B) that is remarkably

similar across both monkeys (Fig. 3, A and B, *first rows*) and speeds (three columns respectively for

6.6°/s, 10°/s and 33.3°/s, as shown in Fig. 2F). The ST representation of non-linearities (lower rows)

recentered on S2 onset, shows that suppression first appears at the cortical representation of S2 and at

S2 response onset, and then propagates towards the representation of S1, at a similar speed than the

one observed for the evoked activity to the first stimulus (Fig. 3, A and B, *second rows, slanting lines*). In

both monkeys and the three examples shown, this suppression propagates in a direction opposite to the

apparent motion sequence, from S2 to S1 representations. Functionally it results in silencing the residual

activity generated by S1.

Figure 3: The apparent motion stimulus induces a systematic suppression wave. *Spatio-temporal representation*

of VSDI responses to two-stroke apparent motion stimuli for three different speed (6.6°/s, 10°/s and 33.3 °/s) and

*two animals (**A:** monkey WA, **B:** monkey BR). The upper rows of A and B represent the observed response and the*

lower rows the non-linearities of the response (observed - linear prediction). Estimates of speed propagation are

reported on each ST diagram (black stairs-step are contours at threshold level, slanting lines are the slope of the

linear regression). Similar values are observed for both the observed activity and the non-linearities.

The suppressive wave propagates at the same speed and with same extent as the evoked spread

This suppressive wave was systematically observed for all two-stroke lrAM conditions tested (see

Fig.1B). This can be seen in the ST evoked response (centered on the onset of S1) and nonlinearities

(centered on the onset of S2) averaged across all conditions and sessions for both monkeys (Fig. 4A). To

better understand the origin of the suppression dynamics, and its dependence on stimulus conditions,

we characterized its spatio-temporal properties. First, we measured the onset of the apparition of the

suppression at S2 position. The latency of the observed suppression was the same as the latency of the

activity evoked by S2 alone (Fig. 4B, respectively 39.5 +/- 2.0 ms vs. 38.6 +/- 1.6 ms for monkey WA and

36.6 +/- 1.8 ms vs. 36.9 +/-2.1 ms for monkey BR, non-significantly different, t-test with p = 0.77 and p =

0.35 respectively for WA and BR). However, the suppression resulted in significantly delaying the

response onset evoked by S2 when presented within the apparent motion sequence (54.2 +/- 2.0 ms

and 68.3 +/- 5.3 ms for WA and BR respectively, Fig. 4B). Then, we quantified the spatial extent of the

suppression (σ of a Gaussian fit, Fig. 4C). In all conditions, the spatial extent of the suppression was of

about 2.8 mm (2.49 +/- 0.14 mm for WA and 3.08 +/- 0.18 mm for BR), similar and non significantly

different than the spatial extent of the evoked response (2.99 +/- 0.11 mm and 2.41 +/- 0.17 mm for WA

and BR respectively). Thus the suppressive wave starts at similar latency and covers similar spatial

extent. We next characterized the speed of propagation of activity (Fig. 4D black) and suppression (Fig.

4D blue), plotted as a function of stimulus speed. Remarkably, on both monkeys, the observed cortical

speeds were identical for both the propagation of activity and the suppression and completely

independent of the lrAM speed (0.28 +/- 0.26 m/s and 0.27 +/- 0.4 m/s respectively for WA and 0.21 +/-

0.15 m/s and 0.27 +/- 0.2 m/s respectively for BR). However, from the ST plots in Figure 3, we noticed

that the suppression does not seem to spread but rather propagates as a wave (Muller et al. 2014,

2018). To probe for this hypothesis we thus compared the dynamics of the response peak position (μ of

a Gaussian fit). In a spread, typically, the response peak will not move in space, as observed for evoked

response (Fig. 4E, the peak spatial position is not changing with time, slope of -1.3×10^{-5} +/- 1.1x10-4 m/s

and 1.6×10^{-4} +/- 3.4×10^{-4} m/s for WA and BR respectively), whereas in a wave it will follow the onset

spatial displacement, which is what we found for the suppression (Fig. 4E, the peak moves from position

2 to position 1, negative slope of -0.05 +/- 0.007 m/s and -0.034 +/- 0.005 m/s for WA and BR

respectively). Altogether, our results show that the suppression is initiated at response onset, have

similar spatial extent and propagation speed as the activity evoked response. Furthermore, although

evoked activity are waves hidden by spatial averaging (Muller et al. 2014), the suppression is still seen as

a wave in the averaged data. This strongly suggests that the suppression is likely to be mediated by the

same general process generating the propagation of evoked activity, most probably the intra-cortical

horizontal network (Muller et al. 2014). If the suppression is generated along the propagation of activity,

one prediction is that it should decrease in strength with spatial and temporal separation between the

two stimuli composing the lrAM. This is indeed what was observed, the suppression strength decreases

as a function of stimulus onset asynchrony and spatial separation (Fig. 4F, t-statistics on the slope of the

linear regression gives t = -0.92 with p=0.18 and t = -6.3 with p = 3.6x10-6, respectively for a spatial

interval of 1$^{\mathrm{o}}$ and 2$^{\mathrm{o}}$ (WA); t = -1.2 with p=0.12 and t = -1.6 with p = 0.05 (BR)).

Figure 4: The suppressive wave has the same properties as the evoked intra-cortical propagation.

A: *Spatio-temporal VSDI activity (top row) and non-linearities (bottom row) averaged across all lrAM speed*

conditions and centered on stimulus 1 (S1, top row) or stimulus 2 (S2, bottom row) onset, for both monkeys

(columns). ***B:*** *Boxplot of latency estimates comparing the onset of activity evoked by S2 alone ("evoked" condition),*

the response onset evoked by S2 when embedded in the lrAM sequence ("lrAM" condition) and the onset of the

suppression at S2 position ("suppr" condition). Boxplots illustrate median, 25 and 75% quartiles and minimum and

maximum of the distributions across all lrAM speed conditions, for the two monkeys (black WA, gray BR). ***C:*** *Boxplot*

of space-constants (parameter ***σ*** *of a Gaussian spatial fit) comparing the evoked response and the suppression, for*

the two monkeys. ***D:*** *For each condition in both monkeys (columns), we estimated the speed of propagation of the*

VSDI (black) and the non-linearity (blue). The upper row shows frequency histograms and the lower row these

speeds as a function of the speed of the lrAM stimulus. ***E:*** *Boxplot of the response peak propagation speed (slope of*

the linear regression on the parameter ***μ*** *of a Gaussian spatial fit) comparing the evoked response and the*

suppression, for both monkeys. ***F:*** *Suppression strength (normalized to the maximal response activity) as a function*

of stimulus onset asynchrony and spatial interval (open circle for SI = 1º, open square for SI = 2º), for both monkeys

(black WA, gray for BR).

The suppressive wave can be the result of a dynamic gain control

What can be the origin of such suppressive wave? Since inhibitory intra-cortical axons have more limited

spatial extent (Buzás et al. 2001), and that feedback from higher areas are excitatory (Salin and Bullier

1995), we can hypothesize that is does not result from a simple net inhibition, but rather as a byproduct

of the excitatory/inhibitory balance (Tsodyks et al. 1997; Ozeki et al. 2009). Indeed, as demonstrated

using center-surround stimulations, the suppressive wave can be the result of a simple dynamic input

normalization fed by propagation along the horizontal network (Reynaud, Masson, and Chavane 2012).

To determine the possible mechanisms generating the observed suppression, we used a mean-field

model designed to reproduce accurately VSDI (Zerlaut et al. 2018). In this model, it was assumed that

each pixel of the VSDI represents the average Vm of two populations of interacting neurons, excitatory

regular-spiking (RS) neurons, and inhibitory fast-spiking (FS) neurons (Chemla and Chavane 2010b).

Arranging this model into a spatially extended interconnected populations of RS-FS cells (Fig. 5A, see

Methods) allows to simulate the propagating waves observed in awake monkey under VSDI. The great

advantage of such model is to explicitly take into account conductance-based interactions (COBA) as

well as a different gain between excitation and inhibition. These ingredients are often neglected as they

introduce difficulties in mathematical tractability of mean field models (Landau et al. 2016; Vogels,

Rajan, and Abbott 2005). Nevertheless, these features are biologically relevant and, as we show here,

are actually the main elements determining waves suppression. Examples of two independent waves

are shown in Fig. 5B (upper row). When the two stimuli are presented in succession (see Fig. 5B lower

left) the observed response shows a suppression (Fig. 5B lower right), whose values are quantitatively

similar to those of experimental data (suppression of around 50% of the response max). Such

suppressive effect was robustly observed across a wide range of the parameters space. The first

parameter that was found to strongly affect the suppression is the ongoing spontaneous activity of the

system pre-stimulus. As we report in Fig. 5C (COBA model, red dots), the suppression decreases when

the spontaneous activity of the system increases (see example marked by a circle, Fig 5D). Moreover,

two further mechanisms were necessary to explain this suppressive effect. First, inhibitory cells need to

have a higher gain than excitatory cells. When the gain of FS cells was reduced (see inset of Fig. 5C) to

have a gain closer to the one of RS cells, the suppression effect was strongly affected (blue dots in Fig.

5C, example marked by a square in Fig. 5D). Accordingly, increasing FS cell gain (cyan dots in Fig. 5C,

example marked by a pentagon in Fig. 5D) increases the suppression strength. Second, the interaction

between excitatory and inhibitory inputs needed to occur through conductances-based mechanisms.

Indeed, when using a current-based (CUBA) model (see Methods), we mostly observed facilitation (black

triangles in Fig. 5C) that do not appear to propagate (see example marked by a triangle, Fig. 5D). While

we do not exclude that such suppression may be observed in current-based synapses, it is clear from

these data that the non-linearity of voltage dependent synapses induces a strong suppression in VSDI

signal. The suppression can thus be explained by the mesoscopic combination of the nonlinearity of

conductance interactions and the differential gain of excitatory and inhibitory cells.

Figure 5: A computational model to investigate the possible origin of the suppressive wave.

A: Mean-field model of excitatory and inhibitory neurons distributed on the cortical trajectory of the stimulus with

horizontal connectivity (longer for excitatory than inhibitory neurons). B: Model ST response to the first stimulus

(upper left), the second (upper right), the apparent motion sequence (lower left) and the non-linearities normalized

to the maximal response over space and time of the response to single stimuli (lower right). The input has an

amplitude v0=20 Hz. C: Amount of suppression/facilitation as a function of the spontaneous excitatory firing rate.

Colored dot stand for different interneurons gain (see inset), while black triangles stand for the Current-based

(CUBA) model, that shows little suppression but facilitation. The input has an amplitude v0=10 Hz. D:

Representative ST suppressive/facilitative patterns as marked in C by different geometric shapes (circle, square,

pentagon, triangle). The star in C corresponds to the model parameters used for obtaining the suppressive pattern

shown in B.

The function of the suppressive wave is to explain away ambiguous representations

What can be the function of the suppressive wave? Here we propose that it will shape an unambiguous

representation of motion along the apparent-motion trajectory. Indeed, silencing the cortical

representation of the initial stimulus when the second stimulus is being processed will have as a

consequence to represent only one stimulus at a time, hereby improving motion representation by

explaining away ambiguous position representation (problem of "phenomenal identity") (Ternus 1926).

To quantify such hypothesis, we developed a simple algorithm to decode, at every instant, what is the

most probable stimulus position that evoked the observed cortical spatial profile out of four categories:

no stimulus, S1, S2, or joint S1 & S2. We used the ST representations of the evoked activity to the

apparent motion sequence (Fig. 6A) and used the linear prediction (Fig. 6B) as a control. The decoding

was computed using the joint probability that the spatial profile observed at one point in time (white

profile) is drawn from the spatial profile observed during blank (first row, black), S1 (second row, red),

S2 (third row, blue), or the joint S1 & S2 (last row, green). In the example shown in figure 6, we apply

this decoding method to the activity evoked by a 6.6°/s two stroke apparent motion stimulus (Fig.6A).

When S1 is presented (red), the probability that the spatial profile of the evoked response will be similar

to the blank distribution is quickly dropping from 1 to 0 and the probability that the evoked response

will be decoded as being evoked by S1 alone is jumping from 0 to 1 very rapidly (in 10ms). When S2 is

presented (at time 50ms) there is a sharp and rapid transition from the evoked activity being decoded as

S1 to S2 (blue) in about 50ms. However, the probability that the evoked activity is evoked by S1 & S2 at

the same time (green) is only increased moderately (peaking at 0.5) and transiently. In contrast, when

we apply the same approach to the linear prediction (Fig.6B), while the beginning of the decoding is the

same (two first rows), as expected, when S2 appears, the evoked activity is ambiguously decoded as

being attributed to S2 or S1 & S2 conjointly with similar probability (around 0.5).

Figure 6: A dynamic decoding of stimulus position: Principle. *The decoding of stimulus position on ST maps, here*

*taking the example of the activity evoked by a 6.6 °/s lrAM stimulus shown in **A** or the activity pattern predicted by*

*the linear combination in space and time of the responses to both individual stimuli in **B**. The decoding consists in*

evaluating the probabilities that the spatial profile observed at each point in time (white contours in A and B) is

similar to one of the four spatial profiles shown on the left column: Blank (first row, black profile), S1 (second row,

red profile), S2 (third row, blue profile), and the joint S1 & S2 (last row, green profile). Each profile was computed by

averaging the corresponding ST response in a 50ms-window around the time of maximum response and

normalized. The four color-coded probabilities are respectively plotted as a function of time (time 0 corresponds to

*the onset of S2) for the lrAM response (**column A**) and for the linear prediction (**column B**). Compared to the linear*

prediction, the actual signal is more rapidly decoded, revealing a likely function of the suppressive wave:

disambiguating stimulus position representation.

We applied this approach to all speeds and sessions in both monkeys (Figure 7A&B), for spatial interval

of 1°, differentiated across the different inter-stimulus intervals (ISI). We separated these conditions

because, when S2 appears, the residual activity in response to S1 will be less important for long ISI (the

offset time constant being of the order of 80 ms). In both monkeys and for ISI <= 50ms, the averaged

results confirm the individual example shown in Figure 6: the evoked activity results in a sharp and clear

transition from the representation of S1 to the representation of S2, with only transient increase of the

representation of S1 & S2 conjointly. In comparison, the linear prediction always leads to an ambiguous

representation that cannot tease apart the probability that the evoked activity is coming from S2 alone

or S1 & S2 together (blue and green curves merging together). For an ISI >= 100ms, in contrast, the

evoked activity resembles more the linear prediction, as expected.

Figure 7: A dynamic decoding of stimulus position: Application to all lrAM speeds and sessions.

A: Color-coded probabilities (same as Figure 6) for the observed lrAM response (first row) and its corresponding linear prediction (second row) for monkey WA, averaged across three ISI categories: ISI < 25 ms (left column), ISI = 50 (central column) and ISI > 100 ms (right column). B: Application of the decoding algorithm to all the data of monkey BR. C: Explaining away index (see text and methods) computed as the probability of detecting joint S1 & S2 in the observed response minus the probability of detecting joint S1 & S2 in the linear prediction, from monkey WA data shown in panel A. D: Explaining away index from monkey BR data shown in panel B.

To quantify the effect of explaining away ambiguous positional representations during lrAM stimulations, we calculated an index by subtracting the probability of detecting joint S1&S2 in the observed and the linear prediction for both monkeys, $I_{E.A.} = P_{S1\&S2}^{obs} - P_{S1\&S2}^{pred}$ (Fig. 7C&D), and both stimuli spatial intervals (SI) of 1 and 2° (first and second rows respectively). In all conditions but the long SI and long ISI, a systematic decrease of the index was observed. This reveals a dynamic effect of explaining away the ambiguous representation of S1&S2. Importantly, in both monkeys and practically all conditions (ISIs and stimulus separation), we observed two peaks in the index decrease. They

correspond to the bidirectional interactions occurring for each of the two evoked waves. The first peak

corresponds to the effect of delaying response onset to S2 (by propagating activity from S1 to S2) , and

the second peak corresponds to a shortening of the representation of S1 (by propagating activity from

S2 to S1). Importantly, this calculation revealed two further phenomena that are expected because of

the propagation delay and spatial extent. First, the timing of the second peak is delayed when going

from 1 to 2° spatial separation. Second, the general amplitude of the decrease diminishes from short to

longer ISI.

Unambiguous representation for optimal encoding of velocity in V1

Disambiguating the cortical population representation of the lrAM could promote an accurate

encoding of direction-selective motion signals for an optimal read-out by downstream area. To test

whether the measured cortical response encodes an accurate direction-selective signal, we applied

opponent motion energy filters directly to V1 population responses (Adelson and Bergen 1985). Indeed,

direction selectivity in MT is well described and captured by motion energy models (Adelson and Bergen

1985; Rust et al. 2006). Such an approach is generally developed to model MT receptive field from a

spatio-temporal input image. The rationale here is to apply the same processing directly to V1

population responses that feed downstream areas such as MT or V4. This is justified by the fact that the

cortical extent imaged here (~ 9mm, corresponding to 3°, see (Dow et al. 1981; Van Essen, Newsome,

and Maunsell 1984)) actually corresponds to the V1 cortical extent converging to a MT or V4 neuron at

our recorded eccentricity (3°, see Albright and Desimone 1987; Gattass, Sousa, and Gross 1988). Since

we record VSD responses that represent both sub- and supra-thresholds activities (Chemla and Chavane

2010b), we first processed our ST maps through a non-linearity to account for the VSD to spike rate

transformation (Chen, Palmer, and Seidemann 2012) (Fig 8A). The resulting ST maps were convolved

with a set of spatio-temporal filters covering a wide range of speeds and scales. For a given value of

filter speed and scale, we squared and summed the convolution from filters in quadrature, and

subtracted the resulting phase-independent measure of local motion energy for opposite directions (ie.

MEu - MEd) to obtain the opponent motion energy response (OME, Fig. 8A). We thereby obtained the

opponent motion energy for all speeds, scales and directions. For each position on the ST map, we could

hence extract the filter velocity for which the opponent motion energy is maximal, that we represented

for both monkeys, and different velocities (10°/s upward in monkey1, Fig. 8B and -33°/s downward in

monkey 2, Fig. 8C). In this representation, the color hue represents the velocity of the filter yielding a

maximal opponent motion energy and the color intensity its amplitude (as a fraction of the maximum

evoked fluorescence response). The contour of the evoked response is overlaid in white to ease

comparison. The same analysis on the corresponding linear predictions serves as a control. For all the

conditions we explored, we then extracted the values of the optimal velocity within a ST region of

interest (between S1 and S2's centers and from 10 to 200 ms after stimulus 2 onset) and represented

them as a function of the AM speed for both monkeys (Fig. 8D and 8E). Our results show that the ST

response, disambiguated through the suppressive wave, is indeed generating a direction selective

motion energy for a speed that is well correlated with the stimulus speed. In other words, intra-cortical

non-linear interactions in V1 promote an unambiguous optimal encoding of velocity-selective motion

signal along the apparent motion path.

Figure 8: Encoding of direction-selective motion signal. A: *Application of the opponent motion energy model*

(Adelson & Bergen, 1985) to the ST representation of cortical response to an upward 10° /s AM sequence shown at

the top. The first step consists in convolving the ST data with a set of oriented ST filters. Phase-independency is

obtained by squaring and summing the outputs of quadrature pair of filters, while motion opponency is obtained by

subtracted the two oriented motion energies (OME = MEu-MEd). The maximal energy values for each ST filter are

plotted as a function of speed (in °/s) and scale (in mm). The energy values resulting from the same computation

applied on the linear prediction and the non-linearities for this AM sequence are respectively shown at the bottom

left and right. ***B:*** *ST representation of the opponent motion energies computed in panel A. For each ST position, the*

filter velocity for which the energy was maximal is represented as different color hue. The amplitude of the energy

is coded as color intensity. For comparison, the result for the corresponding linear prediction is shown below. ***C:***

Same than B for monkey BR, for another AM sequence condition (33.3° /s downward motion). ***D:*** *Filter speed that*

generated the strongest OME within a ST region of interest (see Methods) as a function of the actual lrAM speed

for monkey WA. The color and size of the dots (upward motion conditions) and squares (downward motion

conditions) code for the value of the direction-selectivity index (DI). ***E:*** *Same than D for monkey BR.*

23.

Discussion

We showed that intra-cortical interactions are playing a key role in shaping the sensory representation

of the long-range apparent motion within the retinotopic map of V1 in awake monkeys. Our results

demonstrate that intra-cortical propagation encompasses large spatial and temporal distances allowing

to link information between stimuli presented in distal spatial positions (spatial constant of about 3 mm,

equivalent to 1°, and time constant of about 80 ms). Interestingly, above these values, the apparent

motion illusion gradually fades out (Kolers 1972; Cavanagh and Mather 1989). In response to a two-

stroke lrAM sequence, we observe a clear displacement of activity on the cortical surface that deviates

from the linear prediction in two aspects. First, the initial stimulus suppresses and delays the response

to the second stimulus. Then, a suppressive wave is evoked by the second stimulus that strongly and

rapidly attenuates the residual activity evoked by the first stimulus. The spatio-temporal characteristics

of the suppression showed similar spatial constant and similar propagation speed as what was observed

for the evoked activity, independent of the speed of the apparent motion stimulus. However, the

suppression propagated as a true wave in direction of the initial stimulus position, even at the trial-

averaged level, an observation that departs from what we observed in the evoked activity (Muller et al.

2014). We propose that the suppression arises from a simple gain-control mechanisms pooling

feedforward and horizontal inputs (Reynaud, Masson, and Chavane 2012). To demonstrate this, we used

a conductance based mean-field model developed to account for VSD dynamics (Zerlaut et al. 2018).

This model shows that the observed suppression can be explained by nonlinear conductance

interactions, combined with the different gain of excitatory and inhibitory cells. A decoding approach

demonstrates that the suppressive wave acts as explaining away the ambiguous representation allowing

to represent only one stimulus at a time in the cortex. Using opponent motion analysis applied to the

24.

population response, we demonstrate that such unambiguous representation allows V1 to encode

accurately the velocity signal of the lrAM that could support the read-out process from downstream

areas.

Suppression and normalization as generic operations in the visual system

The dynamics of the suppression is seen here as a central and key mechanism by which the input is

shaped and normalized by V1 populations. When more than one stimulus is present in a visual scene,

suppressive interactions between the feedforward-driven activities is what is traditionally reported,

such as the well documented surround suppression (Blakemore and Tobin 1972; Angelucci et al. 2002;

Cavanaugh, Bair, and Movshon 2002). This suppression is generally attributed to be an emergent

property of the divisive normalization computation (Carandini and Heeger 2011). Importantly, we have

shown that this normalization process is dynamic and propagate from the representation of the stimulus

surround towards the representation of the center (Reynaud, Masson, and Chavane 2012). Adding a

new lateral input (mostly excitatory at long-distance) is therefore resulting in a decrease from the linear

prediction, a paradoxical inhibitory effect (Tsodyks et al. 1997) well captured by Stabilized Supralinear

Networks (Ozeki et al. 2009). Similar suppression was also seen in response to the line-motion stimulus

(Jancke et al. 2004), however, in that stimulus conditions, it was preceded by a transient facilitation. The

main difference with our paradigm is that, in the line-motion condition, the second stimulus, a bar, is

providing a feedforward activation all along the trajectory of the evoked wave. In the apparent motion,

the interactions involve only cortical interactions at positions that do not receive any feedforward input.

This may explain the differences observed with the line-motion stimulus. We believe that dynamic non-

linear interactions subtended by intra-cortical network acts as a general gain control shaping the

representation of visual stimulus in space and in time.

Modeling the suppressive waves

Possible mechanisms underlying the observed suppressive effects were investigated using a spatially extended computational model. We found that the model can reproduce the observed suppression, provided two mechanisms are present: excitatory and inhibitory cells have a different gain, with a higher gain for inhibition, and excitatory and inhibitory synaptic inputs must combine through conductance-based interactions. Although these two mechanisms are well known, they are usually neglected in mean-field models because they represent a mathematical difficulty. The classic mean-field models with linear (current-based) interactions and uniform gain in all cells, fail to reproduce the suppressive effect of propagating waves, and thus the present model can be considered as a step towards biologically more realistic mean-field models. Hence, by constructing a realistic mean-field model, we could demonstrate that this suppression wave is an expected byproduct of the known anatomy and does not need to be expressed solely by pure inhibition. This computational approach demonstrates how excitatory and inhibitory propagation of activity along horizontal network can dynamically change the cortical gain control resulting in the emergence of the observed suppression dynamics.

Backward suppression to keep track of object identity along the apparent motion path

This suppression can help to represent unambiguously one object at a time on the cortical surface, as our decoding model suggests. This means that the lateral interactions can link the transient spatio-temporal events while keeping track of the object moving along the trajectory. This could be a first mechanism involved in solving the correspondence problem (Ullman 1978). This problem, first introduced by Ternus as a problem of phenomenal identity (Ullman 1978; Ternus 1926), explicit the fact we need to keep track of the identity of an object in movement, and, in the case of multiple objects present at each time frame, a problem of correspondence may occur. The literature clearly show that

the correspondence is solved through spatio-temporal coherence more than shape or color consistency

(Kahneman, Treisman, and Gibbs 1992). The correspondence, called "reviewing" by Kahneman et al.

(1992) was proposed by these authors to *"operate(...) backward, (...) select(...) only a single item, and*

(...) is guided mainly by the features that control the unity and continuity of an object over time, but not

by the shape, color, or content of the target." We believe that the mechanisms of backward suppression

demonstrated here is an elementary and preliminary form of this reviewing process, explaining away

ambiguities in the representation of the object trajectory, that will evidently necessitate further

processing downstream the visual system. For instance, what we documented here could explain the

ability of our visual system to detect objects based solely on the coherence of their spatio-temporal

trajectory. In their seminal work, Watamaniuk and collaborators (1995) indeed showed that a single dot

following a temporally coherent trajectory can be detected against a background of dots following a

random walk, the only difference between signal and noise dots movement being their spatio-temporal

coherence (Watamaniuk, McKee, and Grzywacz 1995). Computational studies suggested that this ability

to detect coherent trajectories necessitates propagation of information in retinotopic reference frames

(Perrinet and Masson 2012), in full accordance with our results.

Local vs Global motion processing

The processing that we describe here clearly departs from classical motion integration documented in

short-range apparent motion using random-dot kinetogram (Mikami, Newsome, and Wurtz 1986b, [a]

1986) In these stimuli, motion occurs and is evenly distributed within a stationary aperture typically

covering a receptive field, and motion is extracted locally through motion energy detectors (Majaj,

Carandini, and Movshon 2007; Pack et al. 2006). Simple L-NL hierarchical models account very well for

the selective properties of neurons in V1 and MT in response to such kind of drifting or RDK stimuli (Rust

et al. 2006; Carandini et al. 2005). However, there should be intrinsic differences in the processes

involved in integrating local drifting motion vs global trajectory motion of a single object. Indeed,

Hedges and collaborators (2011) have showed that MT receptive fields are only sensitive to local motion

presented within stationary aperture, totally independent of the direction of long-range trajectory

simulation in which these local motion stimuli are embedded (Hedges et al. 2011). We have very limited

understanding of the processing actually required to extract motion information along a trajectory. The

experiments of Watamaniuk and colleagues show that this processing cannot be simply integrated from

large receptive field of downstream areas (Watamaniuk, McKee, and Grzywacz 1995). Here we suggest

that the visual system can simply encodes the trajectory at mesoscopic level within retinotopic map.

Encoding the motion trajectory in the retinotopic map for optimal read-out

The suppressive wave we documented decreases the residual activity evoked by the first stimulus,

hereby shaping the dynamic response within the retinotopic map of V1 that could be read out as motion

information by a downstream area. V4 or MT neurons have receptive fields whose retinotopic size

encompasses the cortical region we imaged in this study. As shown by our read-out analysis (Fig. 8),

those neurons will be able to simply detect this population-encoded direction selective motion

information through motion energy detectors (Adelson and Bergen 1985). This signifies that V1 intra-

cortical interactions would preformat the population representation of long-range apparent motion for

an optimal read-out by downstream areas (Adelson and Bergen 1985; Mumford 1991, 1992). One

intriguing consequence is that encoding of motion signal at the level of the population could be

operated without specific extraction of motion signal at the level of local V1 neuronal receptive fields.

Indeed, neurons with non-optimal direction preference or no direction selectivity could still participate

into this population response by small variations of their response that would occur at the right moment

depending on their position in the retinotopic space. In other words, V1 would have the possibility to

encode multiple motion signals in parallel at local and global level. These results are in accordance with

human fMRI experiments that showed that V1 is actively involved in the network that processes and

represents the perceived illusory lrAM (Muckli et al. 2005).

lrAM along ventral and dorsal streams, feedback vs horizontal propagation

In the visual cortex of the ferret, it was shown using VSDI, that lrAM induces feedback propagation of

differential activity from area 21 down to area 17 (Roland et al. 2006). Similarly, using stimuli that could

span a much large visual scale (16.5° spatial separation) and systematically larger cortical separations, it

was suggested that human MT complex feedbacks on early visual cortices to process long-range

apparent motion (Wibral et al. 2009; Vetter, Grosbras, and Muckli 2015). Areas on the ventral stream

(LOC) seems to be also implicated in processing such stimuli (Zhuo et al. 2003). Ventral stream areas

may actually be well suited since they will process the information about object through strong

feedback interactions with V1 (Poort et al. 2012) and are as well strongly involved in motion processing

(Roe et al. 2012; Ferrera, Rudolph, and Maunsell 1994). The experiment from Hedges et al. (2011)

indeed suggested that MT may not be the most appropriate area, at least in non-human primates, for

extracting motion along a lrAM trajectory. It is important to consider though that, in all these studies,

there are important difference in the spatial and a temporal scales of the lrAM has been presented that

may affect the relative weight of intra-cortical and feedback mechanisms processing this information

between and within the different visual areas (see Discussion in Reynaud, Masson, and Chavane 2012).

Conclusion

As recently proposed by Muller et al. (Muller et al. 2018), traveling waves within and between cortical

areas can provide an advantageous framework for dynamic computations that will influence neuronal

processing. However, in this review, it was also noted that there is a lack of evidence for a functional

role of these waves. Here we show that two discrete stimuli generating the long-range apparent motion

illusion, will induce multiple wave interactions resulting in propagation of suppression in a direction

opposite to that of the stimuli. Such suppression shapes the stimulus and helps the visual system to

keep track of the stimulus position along the motion trajectory, resulting in a precise encoding of

velocity information at a very early stage of processing. We believe that our work has revealed a first

elementary step in processing lrAM signals that will need further integration in downstream areas and

feedback controls. Further work is therefore needed to understand which areas, if any, is reading-out

the population representation of motion trajectory on V1 retinotopic map and the relative role of intra-

and inter-cortical interactions.

Materials and Methods

The experiments were conducted on two male rhesus macaque monkeys (macaca mulatta, aged 14 and

11 years old respectively for monkey WA and monkey BR) over a period of three years. The

experimental protocols had been previously approved by the local Ethical Committee for Animal

Research (approval A10/01/13, official national registration 71-French Ministry of Research) and all

procedures complied with the French and European regulations for Animal Research as well as the

Guidelines from the Society for Neuroscience.

Surgical preparation and VSDI protocol. The monkeys were chronically implanted with a head-holder

and a recording chamber located above the V1 and V2 cortical areas of the right hemisphere. After full

recovery, the monkeys were trained to perform foveal fixation of a small red target presented over

different static and moving backgrounds for up to 2-3s, with their head fixed. Once a good fixation

behavior was achieved, a third surgery was performed. The dura was removed surgically over the

recording aperture (18mm diameter) and a silicon-made artificial dura was inserted under aseptic

conditions to allow for a good optical access to the cortex over the whole period of weekly recordings.

Before each recording session conducted in awake animal, the cortical surface was stained with the

Voltage Sensitive Dye (VSD) RH-1691 (Optical Imaging ©) with the following procedure: The optical

chamber was open, artificial dura-mater was removed and cortical surface was cleaned under strict

sterile conditions. The dye solution was prepared in artificial cerebrospinal fluid (aCSF) at a

concentration of 0.2 mg/ml, and filtered through a 0.2µm filter. The recording chamber was filled with

this solution and closed for three hours, corresponding to the time lapse needed for a correct cortical

staining. The chamber was then rinsed thoroughly with filtered aCSF to remove any supernatant dye.

Before imaging, the artificial dura was placed back in position and the chamber was closed with

transparent agar and cover glass. Experimental control, data collection and eye position monitoring

were performed by the ReX software (NEI-NIH) running under the QNX operating system (Hays et al.,

1982). During each trial, the cortex was illuminated at 630 nm using epi-illumination and we recorded

optical signals high-pass filtered at 665 nm during 999ms with a Dalstar camera (512x512 pixels

resolution, frame rate of 110 Hz) driven by the Imager 3001 system (Optical Imaging ©). The beginning

of both online behavioral control and image acquisition were heartbeat-triggered. The surgical

preparation and VSD imaging protocol have been described elsewhere (Reynaud, Masson, and Chavane

2012; Muller et al. 2014).

Behavioral task and visual stimulation. Monkeys were trained for a simple fixation task. For each

experimental trial, the monkeys were required to fixate a central red dot within a precision window of

1°x1°. When correct fixation was achieved, the next heartbeat, detected with a pulse oximeter (Nonin

8600V), triggered the beginning of the acquisition window. A visual stimulus appeared 100 ms after this

trigger after which a blank screen was presented, ending the trial. Each trial ran for 700 ms. If the

monkey had maintained fixation up to the end of the acquisition period, a reward (fruit compote drop)

was given. Otherwise, the trial was canceled, an alert sound was delivered and the procedure was re-

initiated. The visual stimuli were computed on-line using VSG2/5 libraries and were displayed on a 22"

CRT monitor at a resolution of 1024x768 pixels. Refresh rate was set to 100Hz. Viewing distance was of

57cm. Luminance values were linearized by mean of a look-up table. We used Gaussian blobs with

standard deviation (controlling the spatial width) of 0.5°. They were presented at different positions,

located at 0.5° or 2° on the left of the vertical meridian respectively for monkey WA and monkey BR, and

between 1.5° and 4.5° below the horizontal meridian. We used different stimulus durations, 10 ms(1

frame), 50ms or 100ms and different interstimulus intervals (ISI) for the two-stroke apparent motion

stimulations (from 20 to 100 ms). All stimuli (single blobs of different durations, lrAM sequences and

two blank conditions i.e. where no visual stimulus) were randomly interleaved with an inter-trial interval

of 8 seconds for dye bleaching prevention.

Data analysis. Stacks of images were stored on hard-drives for offline analysis. The analysis was carried

on with Matlab R2014a (The MathWorks Inc. ©) using the Optimization, Statistics and Signal Processing

Toolboxes. VSD evoked responses to each stimulus were computed in three successive basic steps. First,

the recorded value at each pixel was divided by the average value before stimulus onset ("frame 0

division") to remove slow stimulus-independent fluctuations in illumination and background

fluorescence levels. Second, this value was subsequently subtracted by the value obtained for the blank

condition ("blank subtraction") to eliminate most of the noise due to heartbeat and respiration . Third a

linear detrending of the time series was applied to remove residual slow drifts induced by dye bleaching.

Spatio-temporal representation (ST data). For each time frame, activity was averaged across the x-

dimension within the apparent-motion trajectory (e.g. *dotted rectangle at frame 216 ms* in Fig. 1, C-G)

to provide a unique spatial cortical dimension as a function of time.

Latency estimation. Response latency was defined as the point in time at which the signal derivative

crossed a threshold set a 2.57 times (99% confidence) the SD of its baseline computed during a 100-ms-

long window right before stimulus onset.

Speed estimation. Within the ST representation, the speed of activity propagation was estimated by

computing the slope of the linear regression between each latency estimate as a function of the cortical

distance in the ST representation

Data Fitting. For extracting the space and time constants of the VSD responses, we fitted the ST data in

space (for each time frame) to a Gaussian function of the form:

$$F(x) \ = \ k \, e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

where σ, k and μ respectively denote the width (as the standard deviation), the amplitude and the

spatial position of the Gaussian. We use the slope of the linear regression of $\mu(t)$ for quantifying the

displacement of the response peak (see Fig. 4E).

In time (for each spatial point), the data was fitted to the combination of two halve Gaussian functions:

$$F(t) \ = \ F_{11}(t) \ + \ F_{12}(t)$$

$$F_{11}(t) \ = \ k_1 e^{-\frac{(t-t_c)^2}{2\tau_{on}^2}} . (t \ \leq t_c) \text{ and } F_{12}(t) \ = \ k_2 \, e^{-\frac{(t-t_c)^2}{2\tau_{off}^2}} . (t \ > t_c)$$

where τ_{on} and τ_{off} denote the time-constants of each half Gaussian, while k_1, k_2 and t_c are respectively

their peak to peak amplitudes and the time of their common center.

Statistical Procedure. We used a two-sample t-test procedure to test whether or not the distributions of

the VSD response properties (i.e. space-constant, time-constants, latencies and cortical speed) were

independent of stimulus duration or lrAM speed. $p < 0.01$ is considered significant.

33.

673 ***Mean-field computational model.*** We consider a spatially extended ring model where every node of the

674 ring represents the network activity of a large population of excitatory regular spiking (RS) and inhibitory

675 fast spiking (FS) cells (see Fig. 5A). We consider Adaptive Exponential integrate and fire (AdExp) neurons

676 evolving according to the following differential equations :

$$c_m \frac{dv}{dt} = g_L(E_L - v) + \Delta e^{(\frac{v-v_{th}}{\Delta})} - w + I_{syn}$$

$$\tau_w \frac{dw}{dt} = -w + b\,\tau_w \sum_k \delta\,(t - t_k) + a\,(v - E_L)$$

where $c_m = 100$pF is the membrane capacity, v is the voltage of the neuron and, whenever $v > v_{th} =$

-50mV at times t_k, v is reset to its resting value $v_{rest} = -50$mV. The leak term has a conductance

$g_L = 10$nS and a reversal potential $E_L = -65$mV. The exponential term has a different strength for RS

and FS cells, i.e. $\Delta = 2$mV ($\Delta = 0.5$mV) for excitatory (inhibitory) cells. Inhibitory neurons do not have

adaptation (a=b=0) while excitatory neurons have an adaptive dynamics with $a = 4$nS, b=40 nS and

$\tau_w = 500$ms . The synaptic current can be expressed as:

$$I_{syn} = Q_E(E_E - v)S_E + Q_I(E_I - v)S_I$$

where $S_{E/I} = \sum_{pre} \theta\,(t - t_{pre,E/I})e^{\frac{t-t_{pre,E/I}}{\tau_{E,I}}}$ is the postsynaptic current due to all presynaptic

excitatory/ Inhibitory neurons spiking at time $\tau_{pre,E/I}$ and θ is the Heaviside function. The reversal

potentials are $E_E = 0$mV and $E_I = -80$mV, the synaptic decays are equal for excitatory and inhibitory

cells, $\tau_{E,I} = 5$ms. The quantal conductances are $Q_E = 1$nS and $Q_I = 5$nS. We then consider a random

network with p=5% of connectivity and 80% of excitatory neurons.

The activity of the network is simulated using a mean field model, shown capable of quantitatively

predicting the stationary activity of the network and its response to an external stimuli (Zerlaut et al.

2018). All together, the dynamical equations for the spatially extended ring model read :

$$T\frac{\partial r_E(x,t)}{\partial t} = -r_E(x,t) + F_E(r_{drive} + r_{aff}(x,t) + \int_R dy \, G_E(x-y)r_E(y,t - \frac{|x-y|}{v_c}), r_I)$$

$$T\frac{\partial r_I(x,t)}{\partial t} = -r_I(x,t) + F_I(r_{drive} + r_{aff}(x,t) + \int_R dy \, G_I(x-y)r_E(y,t - \frac{|x-y|}{v_c}), r_I)$$

where $r_{E/I}(x,t)$ is the population rate of excitatory/Inhibitory cells at the space-time position (x,t),

$r_{aff}(x,t)$ is the excitatory afferent input targeting both excitatory and inhibitory populations and $G_{E/I}$ is

the spatial connectivity in between subpopulations that we chose as Gaussian of width $I_{exc} = 5$mm

(excitation) and $I_{inh} = 2.5$mm (inhibition). Moreover, $v_c = 300$mm/s is the axonal conduction speed,

r_{drive} an external time/space constant external drive and T=5ms is the decay time of population rate.

The functions $F_{E,I}$ are the transfer functions of excitatory/inhibitory neurons and are calculated

according to a semi-analytical tool as in Zerlaut et al. (Zerlaut et al. 2018) through an expansion in

function of the three statistics of neurons voltage, i.e. its average μ_V, its standard deviation σ_V and its

autocorrelation time τ_V:

$$F = \frac{1}{\tau_V} Erfc(\frac{v_{thr}^{eff} - \mu_V}{\sigma_V})$$

701   where *Erfc* is the error function and the effective threshold v_{thr}^{eff} is expressed as a first order expansion

702   with some fitting coefficients in function of $(\mu_V, \sigma_V, \tau_V)$. More details on this procedure can be found in

703   Zerlaut et al. (Zerlaut et al. 2018). The values $(\mu_V, \sigma_V, \tau_V)$ are calculated from shot-noise theory (Daley

704   and Vere-Jones 2007). Introducing the following quantities:

$$\mu_{G_E} = r_E K_E \tau_E Q_E$$

$$\sigma_{G_E} = Q_E \sqrt{\frac{r_E K_E \tau_E}{2}}$$

$$\mu_{G_I} = r_I K_I \tau_I Q_I$$

$$\sigma_{G_I} = Q_I \sqrt{\frac{r_I K_I \tau_I}{2}}$$

$$\mu_G = \mu_{G_E} + \mu_{G_I} + g_L$$

$$\tau_m = \frac{c_m}{\mu_G}$$

$$U_s = \frac{Q_s}{\mu_G}(E_s - \mu_V)$$

705   where $K_{E/I}$ is the amount synapses related to pre-synaptic excitatory/inhibitory neurons (we consider a

706   network of N=10000 neurons inside each node of the ring), we obtain the following equations for the

707   voltage moments:

$$\mu_V = \frac{\mu_{G_E}E_E + \mu_{G_I}E_I + g_L E_L}{\mu_G}$$

$$\sigma_V = \sqrt{\sum_s K_s r_s \frac{(U_s \tau_s)^2}{2(\tau_m + \tau_s)}}$$

$$\tau_V = \frac{\sum_s K_s r_s\,(U_s \tau_s)^2}{\sum_s K_s r_s\,(U_s \tau_s)^2/(\tau_m + \tau_s)}$$

708   The afferent input has the following form:

$$r_{aff}(x,t) = A \cdot \frac{1}{2\sqrt{\sigma_{inp}}} e^{-(\frac{x-x_0}{\sqrt{2}\sigma_{inp}})^2} \cdot \left(H(t - t_0)e^{-(\frac{t-t_0}{\sqrt{2}\tau_1})^2} + H(t_0 - t)e^{-(\frac{t-t_0}{\sqrt{2}\tau_2})^2} \right)$$

709   where A is the input amplitude, (x_0, t_0) the stimulus location. And H the heaviside function. The spatial

710   extension of the stimuli is $\sigma_0 = 3.5$mm, the time rise $\tau_1 = 15$ms and the decay time $\tau_2 = 90$ms.

711   The time delay in between stimulus 1 and stimulus 2 is $\Delta_t = 100$ms (if not stated differently) and the

712   spatial distance $\Delta_x = 7$mm. The VSDI signal is calculated as follows :

$$\frac{\delta_V}{V} = \frac{\mu_V - \mu_V^0}{\mu_V^0}$$

713   where μ_V^0 is the average voltage pre-stimuli.

715   *CUBA model :*

716   The current based model is obtained by considering the following synaptic coupling :

$$I_{syn} = Q_E^{CU} S_E + Q_I^{CU} S_I$$

717   where $Q_E^{CU} = 0.03pA$ and $Q_I^{CU} = -0.15pA$ are the coupling with excitatory and inhibitory neurons . The

718   rest of the parameters are the same. The voltage of the neurons is calculated accordingly, i.e.

$$\mu_V = \frac{r_E K_E \tau_E Q_E^{CU} + r_I K_I \tau_I Q_I^{CU} + E_L}{G_L}$$

Also in this case we use the same methodology to estimate the neurons transfer function as done for

the COBA model.

Different FS gain :

In order to modify the gain of FS cells we manually change the transfer function $F_I(r_E, r_I)$.In practice, for

any r_I we calculate the value r_E^* for which TF changes convexity. This gives us the slope

$\sigma_r = \frac{dF(r_E, r_I)}{dr_E}(r_E^*, r_I)$ and the maximal value F_{max} that we estimate calculating F for very high rates

(typically $r_E = 200Hz$).   We then use the following function :

$$F_I(r_E, r_I) = 2\text{max} \cdot \frac{1}{1 + e^{-(\frac{r_E - r_E^*}{\sigma_r})}}$$

where we recall that $r_E^*$and v_r change in function of r_I. This permits us to have a sigmoidal form of the

transfer function F. In order to change its slope we use a factor γ that scales the slope which becomes

then $\gamma\sigma_r$. In Fig. 4 we use γ equal to 1.2 or 0.8.

Decoding Model. The algorithm for the decoding model used in Figures 6 and 7 is detailed here. First,

the ST data (i.e. space-time matrix) were whitened (i.e. spatially decorrelated and scaled) by applying a

ZCA transformation. The whitening matrix was computed from the eigen-decomposition of the

covariance matrix of the blank data. Next, the four spatial profiles (blank, stimulus 1, stimulus 2 and joint

stimulus 1 and 2) were computed by averaging the corresponding ST response in a 50 ms-window

around the time of maximum response and then normalized. The decoding of any ST data (e.g. the

observed activity evoked by a 6.6 °/s two stroke apparent motion stimulus "*obs*" or its linearly predicted

pattern "*pred*") thus consisted in evaluating the likelihood that the spatial profile observed at one point

in time of the data $A(x,t)$was best correlated with one of the four spatial profiles $S_j$with $j \in \{1:4\}$). This

comes down to calculating the four probability $P_j(t)$of the form:

$$P_j(t) \ = e^{-\frac{1}{2\sigma_j} \Sigma_x (\frac{A(x,t)}{||A(x,t)||} - \frac{S_j(x)}{||S_j(x)||})^2}$$

741   where σ_j is the averaged standard deviation of the residual activity between $A(x,t)$ and $S_j(x)$.

742   Then, we defined the explaining away index as the probability of detecting joint S1&S2 in the observed

743   P_4^{obs} or $P_{S1\&s2}^{obs}$ minus the probability of detecting joint S1&S2 in the linear prediction P_4^{pred} or $P_{S1\&s2}^{pred}$ as

744   follows:

745
$$I_{E.A.} = P_{S1\&s2}^{obs} - P_{S1\&s2}^{pred}$$

Opponent motion energy model. . To extract motion information from the population responses, we

used the opponent motion energy model developed by (Adelson and Bergen 1985). Briefly, this model

consists of combining quadrature pairs of spatial and temporal filters to obtain oriented spatio-temporal

filters (i.e. Gabors) tuned in spatial frequency. The ranges of spatial and temporal frequencies were

chosen so that the speed (i.e. FT/FS) of the resulting ST filters varies from 2 to 70 °/s and the scale (i.e.

1/FS) from 0.2 to 6 mm. It resulted in 64 (FS,FT) couples representing 8 different speeds and scales. For

each couple, we obtained two filters tuned for upward motion and two filters tuned for downward

motion. The outputs of quadrature pairs of such filters are then squared and summed to give a phase-

independent measure of local motion energy for both directions (i.e MEu and MEd values). Lastly, the

opponent motion stage computes the difference between the oriented opposite energies (i.e. OME

values). Note that before applying the OME model, the ST data were first normalized and passed

through a non-linearity to account for the VSD to spike rate transformation as proposed by (Chen,

Palmer, and Seidemann 2012):

$$R_{SU} = k(R_{VSDI})^N$$

where R_{SU} and R_{VSDI} are respectively the average firing rate and the average normalized VSDI response, k

is a constant and N is an exponent. Here we took $k = 10$ and $N = 3.8$.

Finally, for each ST position on the map, we could extract the velocity of the filter that generated the

strongest OME and provide a ST velocity map representation (Fig. 8C-D) with velocity and amplitude as

color hue and color intensity respectively. We then averaged the optimal velocity within a ST region of

interest, spatially between S1 and S2's center positions and in time from 10 to 200 ms after stimulus 2

onset, to report a single value of filter speed for each AM speed condition (Fig. 8D-E). The direction-

selectivity index is given by:

$$DI = \frac{V_{OME} - \min(V_{OME})}{\max(V_{OME})}$$

where V_{OME} is to the amplitude of the OME.

Acknowledgments

The authors are thankful to Guillaume Masson, Andrew Meso, Eero Simoncelli, Dirk Jancke, Yves

Fregnac, Cyril Monier, Lyle Muller and Tony Movshon for fruitful discussions during different phases of

this work. They are also grateful to Marc Martin, Frédéric Barthélemy, Ivan Balansard and Luc Renaud

for their assistance regarding experiments.

The authors acknowledge funding from the European Community (FET grants FACETS FP6-015879 and

BrainScaleS FP7-269921), from la Fondation de l'œil (IA) and from the French National Research Agency

(ANR Trajectory, ANR-15-CE37-0011-01, and ANR Horizontal V1, ANR-17-CE37-0006-02).

Competing interests

The authors declare no competing interests.

References

Adelson, E. H., and J. R. Bergen. 1985. "Spatiotemporal Energy Models for the Perception of Motion." *Journal of the Optical Society of America. A, Optics and Image Science* 2 (2): 284–99.

Albright, T. D., and R. Desimone. 1987. "Local Precision of Visuotopic Organization in the Middle Temporal Area (MT) of the Macaque." *Experimental Brain Research. Experimentelle Hirnforschung. Experimentation Cerebrale* 65 (3): 582–92.

Angelucci, Alessandra, Jonathan B. Levitt, Emma J. S. Walton, Jean-Michel Hupe, Jean Bullier, and Jennifer S. Lund. 2002. "Circuits for Local and Global Signal Integration in Primary Visual Cortex." *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 22 (19): 8633–46.

Blakemore, C., and E. A. Tobin. 1972. "Lateral Inhibition between Orientation Detectors in the Cat's Visual Cortex." *Experimental Brain Research. Experimentelle Hirnforschung. Experimentation Cerebrale* 15 (4): 439–40.

Braddick, O. J. 1980. "Low-Level and High-Level Processes in Apparent Motion." *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 290 (1038): 137–51.

Bringuier, V., F. Chavane, L. Glaeser, and Y. Frégnac. 1999. "Horizontal Propagation of Visual Activity in the Synaptic Integration Field of Area 17 Neurons." *Science* 283 (5402): 695–99.

Bullier, J. 2001. "Integrated Model of Visual Processing." *Brain Research. Brain Research Reviews* 36 (2-3): 96–107.

Burr, David, and Peter Thompson. 2011. "Motion Psychophysics: 1985–2010." *Vision Research* 51 (13): 1431–56.

Buzás, P., U. T. Eysel, P. Adorján, and Z. F. Kisvárday. 2001. "Axonal Topography of Cortical Basket Cells in Relation to Orientation, Direction, and Ocular Dominance Maps." *The Journal of Comparative Neurology* 437 (3): 259–85.

Carandini, Matteo, Jonathan B. Demb, Valerio Mante, David J. Tolhurst, Yang Dan, Bruno A. Olshausen, Jack L. Gallant, and Nicole C. Rust. 2005. "Do We Know What the Early Visual System Does?" *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 25 (46): 10577–97.

Carandini, Matteo, and David J. Heeger. 2011. "Normalization as a Canonical Neural Computation." *Nature Reviews. Neuroscience* 13 (1): 51–62.

Cavanagh, P., and G. Mather. 1989. "Motion: The Long and Short of It." *Spatial Vision* 4 (2-3): 103–29.

Cavanaugh, James R., Wyeth Bair, and J. Anthony Movshon. 2002. "Selectivity and Spatial Distribution of Signals from the Receptive Field Surround in Macaque V1 Neurons." *Journal of Neurophysiology* 88 (5): 2547–56.

Chemla, S., and F. Chavane. 2010a. "Voltage-Sensitive Dye Imaging: Technique Review and Models." *Journal of Physiology, Paris* 104 (1-2): 40–50.

———. 2010b. "A Biophysical Cortical Column Model to Study the Multi-Component Origin of the VSDI Signal." *NeuroImage* 53 (2): 420–38.

Chen, Yuzhi, Chris R. Palmer, and Eyal Seidemann. 2012. "The Relationship between Voltage-Sensitive Dye Imaging Signals and Spiking Activity of Neural Populations in Primate V1." *Journal of Neurophysiology* 107 (12): 3281–95.

Daley, D. J., and David Vere-Jones. 2007. *An Introduction to the Theory of Point Processes: Volume II:*
General Theory and Structure. Springer Science & Business Media.
Deco, Gustavo, and Per Roland. 2010. "The Role of Multi-Area Interactions for the Computation of
Apparent Motion." *NeuroImage* 51 (3): 1018–26.
Dow, B. M., A. Z. Snyder, R. G. Vautin, and R. Bauer. 1981. "Magnification Factor and Receptive Field Size
in Foveal Striate Cortex of the Monkey." *Experimental Brain Research. Experimentelle*
Hirnforschung. Experimentation Cerebrale 44 (2): 213–28.
Ferrera, V. P., K. K. Rudolph, and J. H. Maunsell. 1994. "Responses of Neurons in the Parietal and
Temporal Visual Pathways during a Motion Task." *The Journal of Neuroscience: The Official Journal*
of the Society for Neuroscience 14 (10): 6171–86.
Gattass, R., A. P. Sousa, and C. G. Gross. 1988. "Visuotopic Organization and Extent of V3 and V4 of the
Macaque." *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 8 (6):
1831–45.
Gerard-Mercier, Florian, Pedro V. Carelli, Marc Pananceau, Xoana G. Troncoso, and Yves Frégnac. 2016.
"Synaptic Correlates of Low-Level Perception in V1." *The Journal of Neuroscience: The Official*
Journal of the Society for Neuroscience 36 (14): 3925–42.
Grinvald, A., E. E. Lieke, R. D. Frostig, and R. Hildesheim. 1994. "Cortical Point-Spread Function and Long-
Range Lateral Interactions Revealed by Real-Time Optical Imaging of Macaque Monkey Primary
Visual Cortex." *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 14
(5 Pt 1): 2545–68.
Grinvald, Amiram, and Rina Hildesheim. 2004. "VSDI: A New Era in Functional Imaging of Cortical
Dynamics." *Nature Reviews. Neuroscience* 5 (11): 874–85.
Hedges, James H., Yevgeniya Gartshteyn, Adam Kohn, Nicole C. Rust, Michael N. Shadlen, William T.
Newsome, and J. Anthony Movshon. 2011. "Dissociation of Neuronal and Psychophysical
Responses to Local and Global Motion." *Current Biology: CB* 21 (23): 2023–28.
Jancke, Dirk, Frédéric Chavane, Shmuel Naaman, and Amiram Grinvald. 2004. "Imaging Cortical
Correlates of Illusion in Early Visual Cortex." *Nature* 428 (6981): 423–26.
Kahneman, D., A. Treisman, and B. J. Gibbs. 1992. "The Reviewing of Object Files: Object-Specific
Integration of Information." *Cognitive Psychology* 24 (2): 175–219.
Kolers, Paul A. 1972. "THEORIES OF APPARENT MOTION." In *Aspects of Motion Perception*, 172–86.
Landau, Itamar D., Robert Egger, Vincent J. Dercksen, Marcel Oberlaender, and Haim Sompolinsky. 2016.
"The Impact of Structural Heterogeneity on Excitation-Inhibition Balance in Cortical Networks."
Neuron 92 (5): 1106–21.
Lee, T. S., D. Mumford, R. Romero, and V. A. Lamme. 1998. "The Role of the Primary Visual Cortex in
Higher Level Vision." *Vision Research* 38 (15-16): 2429–54.
Majaj, Najib J., Matteo Carandini, and J. Anthony Movshon. 2007. "Motion Integration by Neurons in
Macaque MT Is Local, Not Global." *The Journal of Neuroscience: The Official Journal of the Society*
for Neuroscience 27 (2): 366–70.
Markov, N. T., P. Misery, A. Falchier, C. Lamy, J. Vezoli, R. Quilodran, M. A. Gariel, et al. 2011. "Weight
Consistency Specifies Regularities of Macaque Cortical Networks." *Cerebral Cortex* 21 (6): 1254–72.
Mikami, A., W. T. Newsome, and R. H. Wurtz. 1986a. "Motion Selectivity in Macaque Visual Cortex. II.
Spatiotemporal Range of Directional Interactions in MT and V1." *Journal of Neurophysiology* 55 (6):
1328–39.
———. 1986b. "Motion Selectivity in Macaque Visual Cortex. I. Mechanisms of Direction and Speed
Selectivity in Extrastriate Area MT." *Journal of Neurophysiology* 55 (6): 1308–27.
Muckli, Lars, Axel Kohler, Nikolaus Kriegeskorte, and Wolf Singer. 2005. "Primary Visual Cortex Activity
along the Apparent-Motion Trace Reflects Illusory Perception." *PLoS Biology* 3 (8): e265.
Muckli, Lars, Nikolaus Kriegeskorte, Heinrich Lanfermann, Friedhelm E. Zanella, Wolf Singer, and Rainer

Goebel. 2002. "Apparent Motion: Event-Related Functional Magnetic Resonance Imaging of
Perceptual Switches and States." *The Journal of Neuroscience: The Official Journal of the Society for*
Neuroscience 22 (9): RC219.
Muller, Lyle, Frédéric Chavane, John Reynolds, and Terrence J. Sejnowski. 2018. "Cortical Travelling
Waves: Mechanisms and Computational Principles." *Nature Reviews. Neuroscience* 19 (5): 255–68.
Muller, Lyle, Alexandre Reynaud, Frédéric Chavane, and Alain Destexhe. 2014. "The Stimulus-Evoked
Population Response in Visual Cortex of Awake Monkey Is a Propagating Wave." *Nature*
Communications 5 (April): 3675.
Mumford, D. 1991. "On the Computational Architecture of the Neocortex. I. The Role of the Thalamo-
Cortical Loop." *Biological Cybernetics* 65 (2): 135–45.
———. 1992. "On the Computational Architecture of the Neocortex. II. The Role of Cortico-Cortical
Loops." *Biological Cybernetics* 66 (3): 241–51.
Ozeki, Hirofumi, Ian M. Finn, Evan S. Schaffer, Kenneth D. Miller, and David Ferster. 2009. "Inhibitory
Stabilization of the Cortical Network Underlies Visual Surround Suppression." *Neuron* 62 (4): 578–
92.
Pack, Christopher C., Bevil R. Conway, Richard T. Born, and Margaret S. Livingstone. 2006.
"Spatiotemporal Structure of Nonlinear Subunits in Macaque Visual Cortex." *The Journal of*
Neuroscience: The Official Journal of the Society for Neuroscience 26 (3): 893–907.
Perrinet, Laurent U., and Guillaume S. Masson. 2012. "Motion-Based Prediction Is Sufficient to Solve the
Aperture Problem." *Neural Computation* 24 (10): 2726–50.
Poort, Jasper, Florian Raudies, Aurel Wannig, Victor A. F. Lamme, Heiko Neumann, and Pieter R.
Roelfsema. 2012. "The Role of Attention in Figure-Ground Segregation in Areas V1 and V4 of the
Visual Cortex." *Neuron* 75 (1): 143–56.
Reynaud, Alexandre, Guillaume S. Masson, and Frédéric Chavane. 2012. "Dynamics of Local Input
Normalization Result from Balanced Short- and Long-Range Intracortical Interactions in Area V1."
The Journal of Neuroscience: The Official Journal of the Society for Neuroscience 32 (36): 12558–69.
Roe, Anna W., Leonardo Chelazzi, Charles E. Connor, Bevil R. Conway, Ichiro Fujita, Jack L. Gallant,
Haidong Lu, and Wim Vanduffel. 2012. "Toward a Unified Theory of Visual Area V4." *Neuron* 74 (1):
12–29.
Roland, Per E., Akitoshi Hanazawa, Calle Undeman, David Eriksson, Tamas Tompa, Hiroyuki Nakamura,
Sonata Valentiniene, and Bashir Ahmed. 2006. "Cortical Feedback Depolarization Waves: A
Mechanism of Top-down Influence on Early Visual Areas." *Proceedings of the National Academy of*
Sciences of the United States of America 103 (33): 12586–91.
Rust, Nicole C., Valerio Mante, Eero P. Simoncelli, and J. Anthony Movshon. 2006. "How MT Cells
Analyze the Motion of Visual Patterns." *Nature Neuroscience* 9 (11): 1421–31.
Salin, P. A., and J. Bullier. 1995. "Corticocortical Connections in the Visual System: Structure and
Function." *Physiological Reviews* 75 (1): 107–54.
Sato, Tatsuo K., Ian Nauhaus, and Matteo Carandini. 2012. "Traveling Waves in Visual Cortex." *Neuron*
75 (2): 218–29.
Slovin, Hamutal, Amos Arieli, Rina Hildesheim, and Amiram Grinvald. 2002. "Long-Term Voltage-
Sensitive Dye Imaging Reveals Cortical Dynamics in Behaving Monkeys." *Journal of Neurophysiology*
88 (6): 3421–38.
Stetter, Martin. 2002. "The Early Visual System of Macaque Monkeys." In *Exploration of Cortical*
Function, 23–45. Springer, Dordrecht.
Ternus, Josef. 1926. "Experimentelle Untersuchungen über Phänomenale Identität." *Psychologische*
Forschung 7 (1): 81–136.
Tsodyks, M. V., W. E. Skaggs, T. J. Sejnowski, and B. L. McNaughton. 1997. "Paradoxical Effects of
External Modulation of Inhibitory Interneurons." *The Journal of Neuroscience: The Official Journal*

of the Society for Neuroscience 17 (11): 4382–88.

Ullman, S. 1978. "Two Dimensionality of the Correspondence Process in Apparent Motion." *Perception* 7
(6): 683–93.

Van Essen, D. C., W. T. Newsome, and J. H. Maunsell. 1984. "The Visual Field Representation in Striate
Cortex of the Macaque Monkey: Asymmetries, Anisotropies, and Individual Variability." *Vision*
Research 24 (5): 429–48.

Vetter, Petra, Marie-Helene Grosbras, and Lars Muckli. 2015. "TMS over V5 Disrupts Motion Prediction."
Cerebral Cortex 25 (4): 1052–59.

Vogels, Tim P., Kanaka Rajan, and L. F. Abbott. 2005. "NEURAL NETWORK DYNAMICS." *Annual Review of*
Neuroscience 28 (1): 357–76.

Watamaniuk, S. N., S. P. McKee, and N. M. Grzywacz. 1995. "Detecting a Trajectory Embedded in
Random-Direction Motion Noise." *Vision Research* 35 (1): 65–77.

Wertheimer, Max. 1912. "Experimentelle Studium Uber Das Sehen von Bewegung." *Zeitschrift Fur*
Psychologie 61 (3): 161–265.

Wibral, Michael, Christoph Bledowski, Axel Kohler, Wolf Singer, and Lars Muckli. 2009. "The Timing of
Feedback to Early Visual Cortex in the Perception of Long-Range Apparent Motion." *Cerebral Cortex*
19 (7): 1567–82.

Zerlaut, Yann, Sandrine Chemla, Frederic Chavane, and Alain Destexhe. 2018. "Modeling Mesoscopic
Cortical Dynamics Using a Mean-Field Model of Conductance-Based Networks of Adaptive
Exponential Integrate-and-Fire Neurons." *Journal of Computational Neuroscience* 44 (1): 45–61.

Zhuo, Yan, Tian Gang Zhou, Heng Yi Rao, Jiong Jiong Wang, Ming Meng, Ming Chen, Cheng Zhou, and Lin
Chen. 2003. "Contributions of the Visual Ventral Pathway to Long-Range Apparent Motion." *Science*
299 (5605): 417–20.