Title: Simultaneous multi-area recordings suggest a novel hypothesis about how attention improves performance

Authors: Douglas A. Ruff* and Marlene R. Cohen

Affiliations: Department of Neuroscience and Center for the Neural Basis of Cognition, University of Pittsburgh, Pittsburgh, Pennsylvania 15260; Corespondence to: ruffd@pitt.edu

Abstract - Visual attention dramatically improves subjects' ability to see and also modulates the responses of neurons in every known visual and oculomotor area, but whether those modulations can account for perceptual improvements remains unclear. We measured the relationship between populations of visual neurons, oculomotor neurons, and behavior, which was inconsistent with all published hypotheses about how attention improves perception. Our results support the novel hypothesis that attention reshapes the visual representation of attended stimuli to more effectively influence the premotor neurons that guide decisions without explicitly changing the readout or decision process. Our results suggest a path toward understanding the neural underpinnings of perception and cognition in health and disease by analyzing neuronal responses in ways that are constrained by behavior and interactions between brain areas.

Main Text – Each of the huge number of psychophysical and physiological studies of visual attention show that attention profoundly affects subjects' perceptual abilities and also modulates the responses of populations of neurons at every stage of visual and oculomotor processing (*1*), Despite these oft replicated observations, whether any of the observed neuronal modulations can account for the improvements in psychophysical performance remains unknown. Three hypotheses have dominated the literature (Figure 1A): that attention 1) improves visual information coding (*2*, *3*), 2) improves communication between visual and oculomotor brain

areas (*4*), or 3) reduces uncontrolled fluctuations in the animals' cognitive state, thereby

improving performance and rendering observed effects of attention on response variability as

epiphenomena (*5*). Here we show that none of these hypotheses account for observed perceptual

improvements and we provide evidence for a novel hypothesis: that the well-known effects of

attention on firing rates and shared response variability in visual cortex reshape the

representation of attended stimuli such that they more effectively drive downstream neurons and

guide decisions without explicitly changing the weights relating sensory responses to

downstream neurons or behavior (Figure 1B). We assessed each hypothesized mechanism by

recording simultaneously from groups of neurons in two stages of visuomotor processing: area

MT, which encodes motion information (*6*) and the superior colliculus (SC), where neuronal

responses are either visual, oculomotor, or intermediate (*7-9*) and have been hypothesized to be

involved in computing perceptual decisions (*10*). Constraining our analyses by the animals'

behavior and the simultaneous recordings from both areas allowed us to reject each prior

hypothesis and greatly clarified the relationship between attention, neuronal responses and

behavior.

We compared evidence for and against three hypothesized attention mechanisms using neuronal

responses collected while two rhesus monkeys performed a widely studied motion direction

change-detection task (Figure 1C; *11, 12, 13*). The animals' performance was greatly affected

(Figure 1D) by a cue instructing them to shift spatial attention between a stimulus within the

same or opposite hemifield as the joint receptive fields of several dozen neurons that were

recorded on multielectrode probes in MT (Figure 1E, red points) and the SC (blue points). MT

and the SC represent different stages of perceptual decision-making and therefore provide the

opportunity to evaluate each hypothesized attention mechanism. MT contributes to motion

perception (*6, 14*). The SC is thought to play many roles in visually guided tasks (*7-9*), decision-

making (*10, 15*) and attention(*16*).

Figure 1

53
Figure 1. Hypotheses and methods. (A) Schematics describing predominant hypotheses about
links between attention, visual cortical activity, and behavior. The left plot depicts MT
population responses to two visual stimuli plotted along two dimensions in population response
space (e.g. the first two principal components; see Methods) and a readout dimension which
represents the visual information that is communicated to neuronal populations involved in
planning behavior during the uncued condition. The insets depict projections of the population
responses onto the readout dimension. Hypothesis 1 is that the MT representations of the two
stimuli become more easily distinguishable (e.g. by separating the distributions of responses to
the two stimuli). In this scenario, the distributions of projections along even a suboptimal
readout axis may also be more separable. Hypothesis 2 suggests that attention changes the
communication or readout dimensions, such that projections of responses to the two stimuli are
more separable. Hypothesis 3 proposes that attention-related changes in visual cortex are
epiphenomenal. Instead, it suggests that attention reduces uncontrolled variability in the
animals' internal states, which might produce less variable neuronal population responses and
therefore more separable projections onto the readout dimensions. (B) Our new hypothesis:
attention reshapes population responses so they are better aligned with relatively static readout
dimensions. This alignment could be a direct result of widely observed attention-related changes
in firing rates and response variability. (C) Direction change-detection task with cued attention.

The three predominant attention hypotheses make different predictions about how attention

should affect MT and the SC in our task. The first (information coding) hypothesis predicts that

attention improves the motion direction information encoded in MT. The second

(communication) hypothesis posits that attention improves the stimulus information that is

shared between MT and the SC. The third (internal states) hypothesis suggests that attention

reduces uncontrolled fluctuations in the animals' cognitive states, allowing them to focus on their

psychophysical task. In this scenario, the widely observed attention-related reduction in response

variability in visual cortex (*11, 13, 17*) would be an epiphenomenon of decreased variability in

cognitive states.

Past studies have evaluated these hypotheses by analyzing the responses of individual neurons or

pairs of neurons, which typically lack the statistical power to reveal a strong link to behavior.

Using our data set, we replicated the observations that have been used as evidence in favor of all

three hypotheses. Consistent with previous studies evaluating the information coding hypothesis

(*2, 3, 18*), we found that attention increased the trial-averaged responses of neurons in both MT

and the SC (Supplemental Figure 1A and B). Consistent with studies evaluating both the

information coding and internal states hypotheses (*5*), we found that attention decreased the

extent to which the trial to trial fluctuations in neuronal responses to repeated presentations of

the same stimulus are shared between pairs of MT neurons(*11, 13, 17*) (quantified as the average

spike count or noise correlation, or r_{SC}(*19*); Supplementary Figure 1C). Consistent with studies

evaluating the communication hypothesis, attention increases correlated variability between the

two areas (*12, 20*) (Supplemental Figure 1C). This attention-related increase was weakly

dependent on the visual responsivity of SC neurons (Supplemental Figure 2).

Most previous studies aim to connect the observed attention-related changes in single neurons or

pairs of neurons to behavior via an explicit or implicit model. Recordings from neuronal

populations provide the opportunity to do so directly. We reasoned that analyzing the

relationship between populations of simultaneously recorded neurons in multiple brain areas

with the animals' behavior would provide insight into the relative importance of each

hypothesized mechanism.

The internal states hypothesis provides a challenge for the approach of linking populations of

visual neurons with behavior because it proposes that there is no link between attention-related

changes in MT or other visual areas and performance. Instead, it posits that attention limits mind

wandering, and the changes in performance and neuronal responses simply reflect improved

stability in internal states (*5*). There are two reasons this hypothesis seems unlikely to account for

our data. First, the effects of spatial attention are spatially specific (e.g. correlated variability

increases in one hemisphere while decreasing in the other, even when neurons in the two

hemispheres are simultaneously recorded(*11*)), meaning that reductions in the variability of

global cognitive processes like arousal and motivation are unlikely to account for the attention-

related changes in visual cortex. Further, it is not obvious how reductions in fluctuations in

internal states could account for the attention-related increases in firing rates observed in spatial

attention studies like ours (Supplementary Figure 1), let alone the more complex firing rate

changes associated with feature attention (*21*).

This hypothesis can also be addressed using a population-analysis approach, by using the

responses of MT and SC neurons to attempt to quantify the variability in the animals' internal

states. We reasoned that fluctuations in internal states would 1) often occur at timescales longer

than the 400-600 ms between stimulus presentations in our task and 2) affect the covariability of

neurons in many brain areas, including both MT and the SC. We performed principal

components analysis on population responses to the identical visual stimuli that occurred before

the direction change on each trial (e.g. stimulus A in Figure 1; see Methods) in each attention

condition. Because the only variability in those population responses is internally generated, the

first principal component (PC) represents the axis of greatest shared variability in the population

of neurons in each brain area.

Figure 2

136
137   *Figure 2. Attention has opposite effects on slow fluctuations in neuronal population responses*
138   *within and across areas. (A, B) Autocorrelations between projections onto the first PCs of*
139   *population responses to repeated presentations of the same visual stimulus in (A) MT, and (B)*
140   *the SC. The x-axis plots time lag in units of stimulus presentations (400-600 ms; see Methods).*
141   *(C) Cross correlation between projections onto the first PCs in MT and the SC (same data and*

The autocorrelation functions of projections onto the first PC show that there is indeed response

variability in each area that fluctuates slowly and is reduced by attention (Figures 2A, B), which

is not as readily observable by computing noise correlations between pairs of neurons

(Supplementary Figure 1C). However, the cross-correlation of projections onto the first PC in

MT and the SC showed a qualitatively different time course than the autocorrelation function.

Furthermore, not only did attention not reduce the covariability of these signals between these

two areas (and presumably brain wide), attention *increased* the slow variability that is shared

between areas (Figure 2C and Supplementary Figure 1C). These results are in conflict with the

idea that the attention-related decrease in covariability within each area is a byproduct of a

decrease in uncontrolled fluctuations in internal states, because such a decrease should be brain-

wide.

Figure 3

Figure 3 – Schematic of our decoding procedure. We used linear regression to find the weights (second column) that best relate the **first ten principal components of the MT population's response** (left) to the visual stimulus (Stimulus decoder; top row), the animal's choice (Choice decoder; middle row), or the projections of the responses of the population of simultaneously recorded SC neurons (SC decoder; bottom row). We assessed the performance of each decoder by decoding stimulus information from MT responses on a separate set of trials using each set of weights (right column). See methods for detailed decoding and cross validation procedures.

To investigate the remaining hypotheses, we determined whether attention affects the amount of

stimulus information that can be decoded from the population of MT neurons using linear

decoders that are optimized to a) dissociate between the original and changed stimuli (Stimulus

decoder in Figure 3), b) predict the animals' choices (whether or not they made an eye

movement; Choice decoder), or c) predict the activity of the population of SC neurons we

recorded (SC decoder).

The information coding hypothesis posits that attention improves the stimulus information that

could be gleaned by an optimal stimulus decoder. However, attention did not significantly affect

the performance of an optimal decoder in our data set, even when we used a decoder optimized

separately for each attention condition (Figure 4A, left bars). Furthermore, theoretical work

suggests that the effects of attention on the stimulus information that can be decoded from small

neuronal populations are likely to be even more minimal for larger populations (*22-24*).

The communication hypothesis posits that attention improves communication between visual and

decision areas, meaning that it should change the weights relating MT responses to either

behavior or SC responses. We found that attention had larger effects on the stimulus information

that is related to the animals' choices on individual trials (Figure 4A, middle bars) or that is

shared with the SC (Figure 4A, right bars). However, this could arise from either a weight

change (Figure 1A) or a change within MT, to better align MT responses with static readout

weights (Figure 1B).

Figure 4

Figure 4. Effects of attention on the stimulus information that can be decoded from small populations of MT neurons. (A) Ability of a cross-validated linear decoder to distinguish the original from changed stimuli for each decoder. Error bars represent SEM. The effect of attention was significant for the Choice and SC decoders (paired t-tests, p<.05) but not for the Stimulus decoder (p=0.28). The effects of attention on the Choice and SC decoders were greater than for the stimulus decoder (paired t-tests, p<.05), but not significantly different from each other (p=0.21). (B) Weight swapping analysis demonstrates that decoding performance was typically better using the MT responses from the cued condition and the choice decoder weights from the uncued condition (y-axis) than using the MT responses from the uncued condition and the Choice decoder weights from the cued condition (x-axis; paired t-test, p<.05). (C) Same, using the weights from the SC decoder (paired t-test, p<.05). (D) The reshaping of the MT representation of the attended stimulus can be accomplished as a result of attention-related changes in response variability (e.g. noise correlations). The amount of stimulus information that can be decoded using a single choice decoder whose weights are determined from data from both attention conditions is indistinguishable for the cued data and data constructed using the mean responses from the uncued condition and the residuals from the cued condition (paired t-test, p=0.84).

We found no evidence that attention changes weights relating MT responses to SC responses or behavior. Both the Choice and SC decoders gleaned more stimulus information from MT responses in the attended than unattended condition, even when we used the weights computed in the opposite attention condition for which they were calculated (Figures 4B and 4C). Together, these neuronal population analyses that use the animals' behavior and the activity of downstream neurons to assess the hypothesized attention mechanisms

reveal that none of the three previously hypothesized mechanisms provides a satisfactory account

of the observed attention-related behavioral improvement.

Our data support the hypothesis that attention reshapes the representation of attended stimuli to

more effectively guide behavior (Figure 1B). In this scenario, the critical changes are in visual

cortex. However, this does not result in a large improvement in the stimulus information that can

be gleaned by an optimal stimulus decoder. Instead, the modulated neuronal activity in MT better

aligns with the readout dimensions using relatively static weights.

How could a reshaping of the representation of an attended stimulus be implemented? The

simplest mechanism would make use of the signatures of attention that have been observed for

many years (e.g. changes in firing rate gain or noise correlations(*2*)). We showed recently that

these factors can be readily changed by altering the balance of inhibition to excitation(*25*).

We investigated the possibility that these simple response changes can account for the attention-

related improvement in the stimulus information decoded using the choice decoder in two stages.

First, to verify the prediction of the weight-swapping analyses (Figures 4B and 4C), we

constructed a single choice decoder for both attention conditions and determined that it captured

the attention-related improvement in decoded stimulus information (compare the blue and yellow

bars in Figure 4D). Second, we used those same weights to decode stimulus information from

population responses constructed using the mean rates from the uncued condition but the

residuals from the cued condition. We found that by simply using residuals (which incorporate

both response variability that is private to each neuron and that which is shared between neurons)

from the cued condition was enough to completely account for the attention-related improvement

in decoded stimulus information (Figure 4D).

These results suggest that simple mechanisms like changing response variability are sufficient to

reshape the representation of attended stimuli so that they more effectively influence behavior.

The idea that changing correlated variability better aligns sensory responses to a fixed readout is

consistent with our recent observation that in this task, monkeys' choices are well-aligned with

the axis in population space that explains the most correlated noise (13). One exciting possibility

is that the correlated variability axis represents the fixed readout dimension, perhaps because it is

well-positioned to decode the motion direction of the broad set of stimuli that animals encounter

outside the limited environment of most laboratory tasks(18). If so, reducing noise correlations

and increasing firing rate gains would improve the stimulus information projected along that

readout axis (following the intuitions in (26)).

By combining psychophysics and multi-neuron, multi-area recordings, our study provides a

framework for understanding the neuronal mechanisms underlying a wide variety of sensory,

cognitive, and motor processes. For example, attention-related changes in gains and correlations

have been observed for many years. The key insight that they may improve performance by

aligning the responses of populations of visual neurons to fixed readout dimensions was only

possible using population analyses constrained by the animals' behavior rather than focusing on

one or two neurons.

In the future, it will be interesting to use the same approach to determine whether a similar

mechanism can account for behavioral changes associated with other cognitive processes (e.g.

feature attention) that might seem more likely to change the weights relating stimulus

information to downstream neurons or behavior. Further, many neuropsychiatric disorders

(including disorders of attention, Autism, and schizophrenia) are thought to involve changes in

the same computations thought to underlie attention (*27*). An exciting possibility is that these

changes might be identified and potential therapies evaluated in animal models using the

combination of behavioral evaluation and multi-neuron, multi-area recordings that we described

here.

References:
1.   M. Carrasco, Visual attention: the past 25 years. *Vision Res* **51**, 1484-1525 (2011).
2.   J. H. R. Maunsell, Neuronal Mechanisms of Visual Attention. *Annu Rev Vis Sci* **1**, 373-391
(2015).
3.   J. H. Reynolds, L. Chelazzi, Attentional modulation of visual processing. *Annual Review of*
Neuroscience **27**, 611-647 (2004).
4.   P. Fries, Rhythms for Cognition: Communication through Coherence. *Neuron* **88**, 220-
235 (2015).
5.   A. S. Ecker, G. H. Denfield, M. Bethge, A. S. Tolias, On the Structure of Neuronal
Population Activity under Fluctuations in Attentional State. *J Neurosci* **36**, 1775-1789
(2016).
6.   R. T. Born, D. C. Bradley, Structure and function of visual area MT. *Annual review of*
neuroscience **28**, 157-189 (2005).
7.   R. H. Wurtz, J. E. Albano, Visual-motor function of the primate superior colliculus.
Annual review of neuroscience **3**, 189-226 (1980).
8.   D. L. Sparks, Conceptual issues related to the role of the superior colliculus in the control
of gaze. *Current opinion in neurobiology* **9**, 698-707 (1999).
9.   N. J. Gandhi, H. A. Katnani, motor functions of the superior colliculus. *Annual Review of*
Neuroscience **34**, 205-231 (2011).
10.   G. D. Horwitz, W. T. Newsome, Separate Signals for Target Selection and Movement
Specification in the Superior Colliculus. *Science* **284**, 1158-1161 (1999).
11.   M. R. Cohen, J. H. R. Maunsell, Attention improves performance primarily by reducing
interneuronal correlations. *Nature Neuroscience* **12**, 1594-1600 (2009).
12.   D. A. Ruff, M. R. Cohen, Attention Increases Spike Count Correlations between Visual
Cortical Areas. *Journal of Neuroscience* **36**, 7523-7534 (2016).
13.   A. M. Ni, D. A. Ruff, J. J. Alberts, J. Symmonds, M. R. Cohen, Learning and attention
reveal a general relationship between population activity and behavior. *Science* **359**,
463-465 (2018).
14.   A. J. Parker, W. T. Newsome, Sense and the single neuron: probing the physiology of
perception. *Annual review of neuroscience* **21**, 227-277 (1998).
15.   G. D. Horwitz, W. T. Newsome, Target selection for saccadic eye movements: prelude
activity in the superior colliculus during a direction-discrimination task. *Journal of*
neurophysiology **86**, 2543-2558 (2001).
16.   R. J. Krauzlis, L. P. Lovejoy, A. Zénon, Superior Colliculus and Visual Spatial Attention.
Annual Review of Neuroscience **36**, 165-182 (2013).
17.   J. F. Mitchell, K. A. Sundberg, J. H. Reynolds, Spatial attention decorrelates intrinsic
activity fluctuations in macaque area V4. *Neuron* **63**, 879-888 (2009).
18.   D. A. Ruff, A. M. Ni, M. R. Cohen, Cognition as a Window into Neuronal Population
Space. *Annu Rev Neurosci* **41**, 77-97 (2018).
19.   M. R. Cohen, A. Kohn, Measuring and interpreting neuronal correlations. *Nature*
Neuroscience **14**, 811-819 (2011).
20.   D. A. Ruff, M. R. Cohen, A normalization model suggests that attention changes the
weighting of inputs between visual areas. *Proc Natl Acad Sci U S A* **114**, E4085-E4094
(2017).

21. S. Treue, J. C. Martinez-Trujillo, Feature-based attention influences motion processing
gain in macaque visual cortex. *Nature* **399**, 575-579 (1999).
22. R. Moreno-Bote *et al.*, Information-limiting correlations. *Nature Neuroscience* **17**, 1410-
1417 (2014).
23. I. Kanitscheider, R. Coen-Cagli, A. Pouget, Origin of information-limiting noise
correlations. *Proceedings of the National Academy of Sciences of the United States of*
America **112**, E6973-6982 (2015).
24. A. Kohn, R. Coen-cagli, I. Kanitscheider, A. Pouget, Correlations and Neuronal Population
Information. *Annual Review of Neuroscience* **39**, 237-256 (2016).
25. C. Huang *et al.*, Circuit models of low dimensional shared variability in cortical networks.
bioRxiv **217976**, (2017).
26. B. B. Averbeck, P. E. Latham, A. Pouget, Neural correlations, population coding and
computation. *Nature Reviews Neuroscience* **7**, 358-366 (2006).
27. M. Carandini, D. J. Heeger, Normalization as a canonical neural computation. *Nature*
Reviews Neuroscience **13**, 51-62 (2012).
342
343

Materials and Methods

Methods Summary

344
345
346
The subjects were two adult male rhesus monkeys (*Macaca mulatta*, 8 and 9 kg). All animal

procedures were approved by the Institutional Animal Care and Use Committees of the

University of Pittsburgh and Carnegie Mellon University. Using linear 24 channel moveable

probes (Plexon), we simultaneously recorded extracellular activity from direction-selective

neurons in area MT and neurons in the superior colliculus that responded either visually, prior to

a saccade, or both. Before beginning the experiment, we searched for neurons in both areas that

had overlapping spatial receptive fields (Figure 1E) as determined by mapping with both drifting

gratings and a delayed saccade task. The monkeys performed a direction change-detection task

that commenced upon fixation of a central spot (Figure 1C). Two drifting Gabor stimuli, whose

direction was selected to drive the recorded population of MT neurons well, flashed on and off

until the direction of one stimulus changed at a random, unsignaled time. The monkeys signaled

detection of the change by making a saccade to the changed stimulus within 450 ms of its onset.

The location of the stimulus change was cued using instruction trials prior to each block of trials

and the cue was valid 80% of the time. On the other 20% of trials, the change happened at the

uncued location. In order to encourage fixation on longer trials, catch trials, where no stimulus

changed direction and monkeys were rewarded for maintaining fixation, were intermixed. We

analyzed spiking activity in response to all visual stimuli except the first stimulus in each trial.

Materials and Methods

The subjects were two adult male rhesus monkeys (*Macaca mulatta*, 8 and 9 kg). All animal

procedures were approved by the Institutional Animal Care and Use Committees of the

University of Pittsburgh and Carnegie Mellon University.

We presented visual stimuli using custom software (written in MATLAB using the

Psychophysics Toolbox (*28, 29*) on a CRT monitor (calibrated to

linearize intensity; 1024 768 pixels; 120 Hz refresh rate) placed 54 cm from the animal. We

monitored eye position using an infrared eye tracker (Eyelink 1000; SR Research) and recorded

eye position and pupil diameter (1000 samples/s), neuronal responses (30,000 samples/s), and

the signal from a photodiode to align neuronal responses to stimulus presentation times (30,000

samples/s) using hardware from Ripple.

Behavioral Task

As previously described (*11*), a trial began when the monkey fixated a small, central spot within

a 1.25° per side, square fixation window in the center of a video display while two peripheral

Gabor stimuli (one overlapping the receptive fields of the recorded neurons, the other in the

opposite visual hemifield) synchronously flashed on (for 200 ms) and off (for a randomized

period between 200-400 ms) until, at a random, unsignaled time, the direction of one of the

stimuli changed from that of the preceding stimuli (Figure 1C). The monkey received a liquid

reward for making a saccade to the stimulus that changed. Attention was cued in blocks of 50-

100 trials, and alternated between blocks where attention was cued to either the left or the right

stimulus. In each block, the direction change occurred at the cued stimulus on 80% of trials, and

at the uncued stimulus in 20% of trials (all uncued changes used either the middle or largest

orientation change, Figure 1D). Catch trials, where no stimulus changed direction and the

monkey was rewarded for maintaining fixation, were randomly intermixed throughout each

block. Psychometric data were fit with Weibull functions. Before recording commenced, the

monkeys were extensively trained to have stable thresholds across a range of spatial locations (3-

6 months). During recording, the size, location, spatial frequency and direction of the Gabor

stimuli were selected to drive the recorded units well and the direction of all of the stimuli prior

to the orientation change was constant throughout a recording session.

Electrophysiological Recordings

We simultaneously recorded extracellularly from single units and sorted multiunit clusters (the

term "unit" refers to either). The dataset consisted of a total of 306 responsive MT units and 345

responsive SC units total (36-58 units per session, mean 20 in MT, 24 in the SC for Monkey HO;

36-53 units per session, mean 21 in MT, 22 in SC for Monkey ST) in both MT and the SC in the

right hemisphere using moveable, linear 24-channel V-probes (Plexon; inter-electrode spacing in

MT = 50μm, SC = 100μm). We presented visual stimuli and tracked eye position as previously

described (*12*). The data presented are from 6 days of recording for Monkey HO and 9 days of

recording for Monkey ST. Each day consisted of multiple blocks of the attention task (Figure 1C;

mean 1015 of trials for Monkey HO, 745 for Monkey ST) preceded by receptive field mapping

using a delayed saccade task and direction tuning during passive fixation.

Data Analysis

All spike sorting was done offline manually using Offline Sorter (version 3.3.5; Plexon). We

based our analyses on both single units and multiunit clusters and use the term "unit" to refer to

either. All neuronal analyses prior to Figure 3 used spike count responses between 50-250 ms

after stimulus onset to account for visual latencies in the two areas. To remove response

contamination from eye movements during change stimuli, data presented in the decoding

analyses in Figure 3 and 4 used shorter response windows. Responses to both unchanged and

changed stimuli were measured from 50-185 ms after stimulus onset for monkey HO and 50-220

ms for monkey ST. These times were selected based on the distribution of each animal's reaction

times, with the goal of maximizing the number of trials that could be included in the analyses.

Trials with reaction times that began during those windows were excluded. To minimize the

impact of adaptation on our results, we did not analyze the first stimulus presentation in each

trial. We only analyzed a recorded MT unit if its stimulus-driven firing rate was 10% higher than

its firing rate as measured in the 100 ms prior to the onset of the first stimulus. We only analyzed

a recorded SC unit if its stimulus-driven firing rate was 10% higher than its firing rate as

measured in the 100 ms prior to the onset of the first stimulus or if its response during a 100 ms

epoch prior to a saccade on correct trials to the contralateral side was 10% larger than that same

baseline. Stimulus presentations during which a microsaccade was detected were excluded from

analyses (*12, 30*)).

For firing rate analyses in Supplemental Figure 1A and B, attention indices were calculated using

average spike counts on the stimulus presentation prior to correct detections of the intermediate

change amount depending on whether attention was directed into or out of the receptive fields of

the recorded neurons using the formula $(attend_{in} - attend_{out})/(attend_{in} + attend_{out})$. Significance of

individual units was determined by a paired t-test ($p < 0.05$).

Noise correlations

We defined the correlated variability of each pair of simultaneously recorded units (quantified as

spike count correlation or r_{SC} (*19*)) as the Pearson correlation coefficient between the responses

of the two units to repeated presentations of the same stimulus. This measure of r_{SC} represents

noise correlations rather than signal correlations because the responses used in this analysis were

always to an identical visual stimulus. For Supplemental Figure 1C, we included responses from

stimulus presentations 2 though 10 from trials that ended with either a hit, miss or correct catch

trial and that were immediately followed by the maintenance of fixation and continuation of the

trial. We z-scored responses as a function of the stimulus presentation number in each trial and

then pooled data across stimulus presentations before calculating noise correlations. Results did

not qualitatively change if we did not perform this z-score procedure. For Supplemental Figure

1D, we included data from all stimulus presentations prior to the change stimulus (except the

first) and sorted them depending on what the behavioral outcome was on the subsequent stimulus

presentation. Pairs of units that were recorded on the same electrode were not included in

correlation analyses. The data presented in Supplementary Figures 1C consisted of 3,285 MT

pairs, 3,948 SC pairs and 6,934 between area pairs.

Slow timescale auto- and cross-correlations

To test the hypothesis that attention affects uncontrolled fluctuations in internal states, we

created a procedure to identify slow fluctuations in population responses. We plotted the

responses of the populations of simultaneously recorded MT or SC neurons to the stimuli before

the motion direction change (excluding the first stimulus) in a high dimensional space in which

each neuron's response comprises one axis. We performed principal components analysis (PCA)

on this cloud of points. Because the visual stimuli in this subset of the data were all identical, the

first principal component (PC) represents the axis that captures the most variance in non-

stimulus related population responses. We searched for slow fluctuations in internal states by

measuring the auto- and cross-correlations in projections onto this first PC in each area.

Decoding

We focused our decoding analyses on trials in which the third largest (middle) direction change

occurred, because changes of that magnitude occurred in both attention conditions. We used the

decoding strategy schematized in Figure 3. We began by constructing four matrices in each

attention condition: 'MT responses' (a # MT neurons x 2*# trials matrix of MT responses to the

stimuli before the direction change and the changed stimulus on the relevant trials), 'motion

direction' (a 1 x 2*# trials vector of zeros for the stimulus before the change, referred to as

'original', and ones for the changed stimulus, referred to as 'change'), 'choice' (a 1 x 2*# trials

vector of zeros for stimulus presentations on which the animal did not make an eye movement,

referred to as 'no saccade', and ones when the animal made an eye movement, referred to as

'saccade'), and 'SC responses' (a # SC neurons x 2*# trials matrix of SC responses to the stimuli

before the direction change and the changed stimulus on the relevant trials).

We cross validated by holding out the two stimulus presentations (for the original and changed

stimuli) from one trial at a time to perform the rest of our analyses. To reduce the number of

weights we needed to fit and therefore improve our confidence in the weights we did fit, we

performed PCA on the MT and SC responses to find the first 10 PCs in each area. We then

performed linear regression to find the weight vectors (for the Stimulus and Choice decoders) or

weight matrices (for the SC decoder) that related projections along the first ten MT PCs plus a

vector of ones to 'motion direction', 'choice', or projections along the first 10 SC PCs in each

attention condition.

We assessed the stimulus information in each decoder (Figure 4) by multiplying projections of

MT responses to the original and changed stimuli from the held-out trial by the fitted weights

and either determining whether those weighted sums correctly classified the stimuli as original or

changed (Stimulus and Choice decoders) or whether a linear classifier correctly classified those

stimulus presentations on the basis of the predicted SC responses (SC decoder). For the decoding

analysis in Figure 4D, we took a similar approach to the previously described choice decoder,

except that we combined data from both the cued and uncued conditions to calculate decoding

weights. We then decomposed the responses of the population responses to each stimulus in each

attention conditions into mean responses and residuals (R=M+S, where R is the number of

neurons x number of trials matrix of spike count responses to one stimulus in one attention

condition, M is a matrix of mean responses for each neuron, and S is the matrix of residuals). We

tested the hypothesis that attention-related changes in the residuals account for the improvement

in stimulus information used to guide behavior by decoding stimulus information from responses

created by using the mean responses from the uncued condition and residuals from the cued

condition.

501
502

Supplemental Figure 1

503
Supplemental Figure 1. Effects of attention on common analyses of individual units and pairs of
units (A) Attention increases firing rates in MT, quantified as the difference in firing rates in the
different attention conditions divided by the sum. Units with significant differences in average
responses for the two conditions are specified by black bars (t-test, p<.05). This distribution
(mean = 0.04, median = 0.04) is significantly different from zero (Wilcoxon signed rank test, p<
10^{-21}). (B) Same as A, for SC data. This distribution (mean = 0.073, median = 0.05) is significantly
different from zero (Wilcoxon signed rank test, p< 10^{-43}). (C) Within and between area noise
correlations calculated from spike counts during stimulus presentations that preceded successful
maintenance of fixation from trials that ended with either a hit or miss or were a successful
catch trial. Attention decreases average correlations within MT (Wilcoxon signed rank test, p<
10^{-12}), not in the SC (Wilcoxon signed rank test, p=0.8) and increases them between the two
areas (Wilcoxon signed rank test, p< 10^{-41}). Error bars are standard error of the mean. (D)
Within and between area noise correlations calculated from spike counts that immediately
preceded different behavioral outcomes during cued trials. Misses and false alarms are
associated with higher correlations within MT (t-test, p<10^{-3}) and SC (t-test, p<10^{-3}) but not
between the two areas (t-test, p=0.23). Error bars are standard error of the mean.
520

Supplemental Figure 2

521
Supplemental Figure 2. Relationship between SC responses during different task epochs and
attention-related correlation changes with MT. (A) Schematic of task timing depicts the three
100ms epochs used to count spikes in SC units. The baseline period began 100ms before the first
stimulus appeared, which is after stable fixation had been acquired. The stimulus period was
shifted 30 ms after the appearance of the visual stimulus, to account for the earliest visual
latencies observed in the SC. The delay period began 100ms after the first stimulus turned off
and always ended prior to the onset of the second stimulus. (B) Attention-related changes in
MT-SC r_{SC} plotted against the difference between each SC unit's response during the stimulus
and baseline periods. There are multiple MT-SC correlation differences measured for each SC
unit. Correlations between MT and SC were calculated using the same data and methods as
Supplementary Figure 1C (Pearson correlation, rho=0.087, p< 10^{-12}). Isolines depicting the decile
boundaries are overlaid over the individual data points. (C) Similar to B, but data are now sorted
by the difference between each SC unit's response during the stimulus and delay periods
(Pearson correlation, rho=0.092, p< 10^{-14}). (D) Similar to B, but data are now sorted by the
difference between each SC unit's response during the delay and baseline periods (Pearson
correlation, rho=-0.042, p< 10^{-4}).
538
539
540