Original Research:

The taxonomic and functional biogeographies of phytoplankton and zooplankton communities

across boreal lakes

Nicolas F. St-Gelais[*], Richard J. Vogt, Paul A. del Giorgio and Beatrix E. Beisner

Groupe de Recherche Interuniversitaire en Limnologie et en Environnement Aquatique (GRIL)
and Department of Biological Science, University of Québec at Montréal, Montréal, QC,
Canada.

Corresponding author: Nicolas F. St-Gelais (nicolas.fstgelais@gmail.com)
[*]Current address : Groupe de Recherche Interuniversitaire en Limnologie et en Environnement
Aquatique (GRIL), Department of Biological Sciences, Université de Montréal, Montréal,
Quebec, Canada

Orcid ID (NFS): 0000-0002-6721-5191
OrcidID (BEB): 0000-0001-6972-6887

Keywords: plankton; community; functional traits; Canada; water quality; habitat;

biogeography; morphometry; boreal

Abstract

Strong trophic interactions link primary producers (phytoplankton) and consumers (zooplankton)

in lakes. However, the influence of such interactions on the biogeographical distribution of the

taxa and functional traits of planktonic organisms in lakes has never been explicitly tested. To

better understand the spatial distribution of these two major aquatic groups, we related the

distributions of their taxa and functional traits across boreal lakes (104 for zooplankton and 48

for phytoplankton) to a common suite of environmental and spatial factors. We directly tested

the degree of coupling in their taxonomic and functional distributions across the subset of

common lakes. Phytoplankton functional composition responded mainly to properties related to

water quality, while zooplankton composition responded more strongly to lake morphometry.

Overall, the spatial distributions of phytoplankton and zooplankton were coupled at taxonomic

and functional levels but after controlling for the effect of environmental drivers (water quality

and morphometry) and dispersal limitation, no residual coupling could be attributed to trophic

interactions. The lack of support for the role of trophic interactions as a driver coupling the

distribution of plankton communities across boreal lakes indicates that taxon-specific and

functional trait driven ecological interactions may not modulate large-scale spatial patterns of

phytoplankton and zooplankton in a coordinated way. Our results point to community structuring

forces beyond the phytoplankton-zooplankton trophic coupling itself, and which are specific to

each trophic level: fish predation for zooplankton and resources for phytoplankton.

Introduction

The composition of ecological communities is the results of multiple assembly processes (Kraft and Ackerly 2014) occurring at different spatial scales (Declerck et al. 2011). At the local spatial scale, species interactions (e.g. competition, predation, mutualisms) are considered to be strong an assembly processes that structure ecological communities (Diamond 1975). However, the relative influence of such local species interactions on the larger-scale distributions of organisms within and across regions is not well understood (Gotelli et al. 2010, Wiens 2011).

In lakes, planktonic consumers (zooplankton) and primary producers (phytoplankton) interact strongly, mainly via trophic relationships (Porter 1977, Sterner 1989), and studies on trophic cascades have emphasized how changes at one trophic level can affect entire food webs (Carpenter et al. 1985). While the top-down and bottom-up effects have typically been studied for their influence on standing biomass, they also influence community composition (McQueen et al. 1989, Ghadouani et al. 2003). However, whether these strong local trophic interactions affect the larger-scale distribution of zooplankton and phytoplankton in lake metacommunities at regional scales is still unresolved.

At the regional scale, the effect of habitat on the taxonomic biogeography of zooplankton and phytoplankton is well studied (see Keller and Pitblado 1989, Pinel-Alloul et al. 1990, 1995, O'Brien et al. 2004, Soininen et al. 2011, Stomp et al. 2011). The taxonomic composition of both phytoplankton and zooplankton are known to respond strongly to lake nutrient concentrations (Keller and Pitblado 1989, Pinel-Alloul et al. 1990, Soininen et al. 2011), as well as to other environmental variables such as lake colour (dissolved organic carbon content, Wissel et al. 2003) and lake morphometry (e.g. lake depth, Pinel-Alloul et al. 1990). On the other hand, very few studies have examined the joint distributions of phytoplankton and zooplankton and

thus, the importance of the trophic link between, phytoplankton and zooplankton, as a potential

constraining factor of their respective taxonomic distributions on a landscape, has never been

directly tested. Soininen and colleagues (2007) compared the spatial concordance of distance-

decay relationships of phytoplankton and zooplankton taxa across ponds, finding that they were

significantly stronger for zooplankton than for phytoplankton, but only across drainage basins

and not within basins. However, they did not directly assess the joint taxonomic distributions of

the different plankton groups.

Although zooplankton exhibit prey selectivity (Vanderploeg 1981, Paffenhöfer 1984),

individual taxa often feed on many different phytoplankton species (Knisely and Geller 1986).

Zooplankton grazing thus reflects selection for certain functional traits (e.g. size, shape) that are

common to a number of phytoplankton species, more than for specific taxonomic groups per se.

Thus, if trophic interactions between phytoplankton and zooplankton are strongly constrained by

functional traits of both plankton groups, then joint biogeographical concordances should be

more evident for functional traits than for strict taxonomic classification. A temporal analogue of

this idea is the repeatable seasonal succession observed in many temperate lakes and captured

effectively by the PEG model, which characterizes the dynamic effects of such trophic

interactions by referring to specific functional groups of phytoplankton and zooplankton

(Sommer et al. 1986, 2012). If functional trophic interactions are equally spatially important, and

repeatable, across a landscape, one would expect any biogeographical concordance between

phytoplankton and zooplankton to be more observable using a functional trait lens than a

taxonomic one. Functional traits currently available for lake plankton account mechanistically

for trophic interaction potential related to food web responses, including resource availability

(phytoplankton), feeding behaviours (zooplankton), and predator evasion (phytoplankton and

zooplankton, Barnett et al. 2007, Hébert et al. 2016).

In addition to trophic interactions, spatial coupling between phytoplankton and

zooplankton, even at the functional trait level, could also arise from similar responses to

environmental drivers (habitat selection). For example, the taxonomic distributions of both

phytoplankton and zooplankton are known to respond independently to lake nutrient

concentrations (Keller and Pitblado 1989, Pinel-Alloul et al. 1990, Soininen et al. 2011), which

could result in coupled distributions at the landscape scale. Beyond lake nutrient status, however,

expectations for common responses of both plankton groups to gradients of other environmental

variables such as lake colour (dissolved organic carbon content), lake size, and stratification are

not evident. Furthermore, when considering a suite of environmental variables, different variable

types do not necessarily have the same spatial structure: water quality variables generally having

stronger spatial structure than do those related to lake morphometry (Lapierre et al. 2015).

Consequently, while there is good reason to expect that each plankton group will respond to

environmental variation, such responses are unlikely to be coordinated across different

environmental variable types. In fact, this lack of coordination could disrupt factors like trophic

interactions or nutrients that are likely to promote joint distributions. Proximate (water quality)

variables with their strong spatial structure across landscapes should therefore favour greater

plankton biogeographical concordance, than will less structured habitat variables characterizing

lake morphometry.

In addition to environmental gradients, dispersal limitation is also known to structure

phytoplankton and zooplankton communities (Heino et al. 2015). Previous work has indicated

that phytoplankton generally respond to environmental factors while zooplankton are more

dispersal-limited (Beisner et al. 2006, De Bie et al. 2012, Padial et al. 2014). Such differential

response patterns could also interfere with a strong biogeographical coupling of the groups. In

fact, differential relative responses of phytoplankton and zooplankton to environmental factors

and distances between lakes, make it necessary to control for their respective effects before

assessing the importance of trophic interactions on the biogeographical coupling of the two

plankton groups.

Here, we conducted the first comprehensive study that directly considers the relative

importance of trophic interactions between planktonic organisms and relates these to the effects

of environmental conditions and dispersal (landscape space) to affect the coupling of their

biogeographical distributions. Understanding of the importance of trophic interactions at the

landscape-scale has implications for the impact of broad-scale changes in zooplankton or

phytoplankton composition (e.g. Jeziorski et al. 2015) on aquatic food webs. To investigate

whether the signature of trophic interactions could be captured at a regional scale, we assessed

the coupling between the taxonomic and functional distribution of lake zooplankton and

phytoplankton communities.By using a stepwise framework, we first accounted for the effect of

habitat selection and dispersal, and then assessed whether residual variation in the observed

coupling between species and traits could be attribution to trophic interactions. This approach

also allowed us to investigate the environmental and spatial factors influencing each group

(phytoplankton or zooplankton) separately as well. Our study covers a large biogeographical

range (longest distance between two lakes is 1 228 km) and examines more than 100 lakes

clustered in three regions that together characterize geological and environmental variation in the

boreal belt across one of Canada's largest provinces (Québec). Because trophic interactions are

mediated by functional traits, we expected to observe a stronger residual coupling between the

distribution of zooplankton and phytoplankton traits compared to the taxonomical coupling.

Furthermore, we expected the coupling to be even stronger when only traits specific to trophic

interactions between zooplankton and phytoplankton were considered (e.g. pigment type, feeding

strategy).

Materials and Methods

Study lakes and sampling

Crustacean zooplankton samples were collected from 104 lakes with a low level of

anthropogenic disturbance, within three environmentally (Figure 1, Table 1) and geologically

(Roy 2012) distinct regions of Quebec, Canada, during the years 2010 (Abitibi, May to October),

2011 (Chicoutimi, June to October) and 2012 (Schefferville, July to August). Integrated

zooplankton samples were collected from the deepest point of each lake using a conical plankton

net (110 µm mesh, 0.30 m mouth diameter), equipped with a flow meter (General Oceanics,

USA), hauled vertically from 1 m above the sediments to the surface. Zooplankton samples were

anaesthetized using carbonated water and were preserved in 75% (final concentration) ethanol.

In a subset of 48 lakes, the phytoplankton community was also simultaneously sampled over the

photic zone using a flexible PVC sampler tube and an integrated subsample (250 ml) was

preserved in Lugol's solution.

Crustacean zooplankton were identified at the species level (but aggregated for analyses

at the genus level to correspond to the phytoplankton data), using an inverted microscope (50-

400X) and individuals were counted until a total of 200 individuals had been enumerated. For

each taxon present in a lake, the length of 20 mature individuals was measured and biomass by

taxon was estimated using length-dry-mass regressions (McCauley 1984, Culver et al. 1985).

Phytoplankton were enumerated at the genus level using the Ütermohl method on an inverted

microscope at 400X magnification. Phytoplankton biomass was estimated from biovolume

computed using cell and colony length measurements and corresponding geometric forms

(Hillebrand et al. 1999). We also measured key limnological variables to characterize the lake

and catchment environments. We used a multiparameter sonde (YSI, Yellow Springs

Instruments, OH, USA) to measure pH (at 0.5m depth) and temperature (at 0.5m depth intervals,

then averaged over the water column). Water samples were collected at 0.5m depth to measure

the concentration of chlorophyll *a* (Chl-*a*), total phosphorus (TP), total nitrogen (TN), dissolved

organic carbon (DOC) and coloured dissolved organic matter (CDOM). Chl-*a* was extracted

with 90% hot ethanol and absorption was measured spectrophotometrically before and after

acidification to account for phaeophytin (Lorenzen 1967, Nush 1980); TP was measured from

water samples using the molybdenum-blue method following persulfate digestion (Cattaneo and

Prairie 1995); TN was measured using nitrates after persulfate digestion; DOC concentration was

measured on an O.I. Analytical (Texas, USA) TIC/TOC using 0.45 μm filtered water after

sodium persulfate digestion; CDOM was measured using a UV/Vis UltroSpec 2100

spectrophotometer (Biochrom, Cambridge, UK) at 440 nm. Missing values (see Table 1) were

imputed using an approach based on random forest (missForest R package, Stekhoven and

Bühlmann 2012, NRMSE : 0.038). Lake depth was measured at sampling point using a Portable

Water Depth Sounder Gauge (Cole-Parmer). Lake area was derived using ArcGIS V10 software

(ESRI Inc., Redland, CA, USA) and catchment slope was estimated using a Digital Elevation

Model (Canadian Digital Elevation Data). To visualize environmental differences between the

three regions, we used a principal component analysis (PCA) using the *rda* function (vegan R

package, Oksanen et al. 2015). Finally, because our sampling was discontinuous on the

landscape we used the Euclidean distance between lakes to characterize the effect of dispersal

limitation on the distribution of taxa and functional traits.

Functional trait composition and diversity

Given that our objective was to test for a significant coupling between adjacent trophic

groups, we selected functional traits (Table 2) that explicitly characterize the grazing interaction

between phytoplankton and zooplankton, as well as their interaction with other trophic levels in

the food web. For phytoplankton, we selected traits that define how they interface with resources

(i.e. nutrients and light) and the zooplankton grazers (motility, edibility, colony formation). For

zooplankton, we focused on traits that define how they consume phytoplankton (i.e. feeding type

and trophic group). Phytoplankton trait values were obtained from a literature review (see

(Longhi and Beisner 2010 for details) and included (i) capacity for N-fixation, (ii) silica demand,

(iii) capacity for mixotrophy, (iv) pigment composition (v) cell motility and (vi) edibility to

zooplankton (>35 μm linear dimension) and (vii) tendency to form colonies. Crustacean

zooplankton traits values were also obtained from a literature review (see Barnett et al. 2007))

and included (i) feeding type and (ii) trophic group. For both zooplankton and phytoplankton, we

also used the average individual biomass of each taxon as an integrative functional trait of body

size (Brown et al. 2004, Litchman et al. 2013) related both to resource acquisition and grazer

avoidance (Table 2).

Statistical analyses

To visualize how phytoplankton and zooplankton taxa and traits were distributed on the

landscape, we used the percentage of lakes in which they occurred. We tested for differences in

taxonomic or functional composition between the three regions using the constrained ordination

technique Canonical Analysis of Principal Coordinates (*CAP* BiodiversityR package; Anderson

and Willis 2003). Using the CAP leave-one-out allocation success (% correct, Anderson and

Willis 2003) we assessed the distinctiveness of regional composition using the proportion of

correct allocation, which can be interpreted as the strength of the compositional differences

between regions. Prior to the CAP ordination, taxonomic composition was Hellinger-transformed

to reduce the effect of double zeros (Legendre and Gallagher 2001) and Bray-Curtis distance was

used for the CAP ordination.

To test whether the distributions of phytoplankton and zooplankton communities were

coupled across the landscape, we used a hierarchical framework (Figure 2). First, we tested for

significant coupling between the composition of the two groups at taxonomic and functional trait

levels using a Procrustes analysis (Mardia et al. 1980) on the subset of lakes (48) for which both

phytoplankton and zooplankton were sampled. Specifically, we tested the degree of concordance

between the PCA ordinations of phytoplankton and zooplankton taxonomic and functional

community compositions. For this analysis, species composition was transformed using the

Hellinger transformation (Legendre and Legendre 1998, Legendre and Gallagher 2001), and

functional trait composition was transformed using a logit transformation. We tested the

significance of the Procrustes statistic using a permutation procedure (9999 simulation, *protest*

function in vegan R package, Oksanen et al. 2015). To further investigate whether the coupling

could be explained by a similar response of the two groups to environmental variation or by

dispersal limitation (see Figure 2) we also tested for a significant coupling, using Procrustes

analysis, after controlling for environmental factors (water quality characteristics and

morphometry) and space (using between-lake distance based on latitude and longitude

coordinates).

To directly evaluate the relative importance of environmental and spatial variables as drivers

of the taxonomic and functional composition of phytoplankton and zooplankton, we used

distance-based redundancy analysis for taxonomic composition (dbRDA, Legendre and

Legendre 1998) and multiple regression for each functional trait; both followed by variation

partitioning (Borcard et al. 1992). We separated the environmental variables into two groups;

water quality (WQ with Chl-*a*, TP, TN, DOC, water color, pH, temperature) and morphometry

(M with lake area and depth). Prior to the RDAs and multiple regressions, we used a stepwise

selection (based on AIC) of variables within each group of environmental variables. We

identified shared variation between the two groups of environmental variables (WQ+M), and

between environmental variables and space (S) as: WQ+S and M+S to determine whether

environmental variables were spatially structured. Finally, to visualize the relationship between

phytoplankton and zooplankton taxa and functional traits, we used an RDA with all taxa and all

the functional traits combined.

Results

<u>Biogeographical patterns</u>

We observed important environmental differences between the three regions (Figure 1, Table 1).

The first principal component (PC) mainly differentiated the Abitibi and Chicoutimi regions

from the Schefferville region and represented differences that reflect latitude between the

regions: lakes in Abitibi and Chicoutimi being warmer, darker (higher *CDOM*) with higher

concentrations of chlorophyll *a*, dissolved organic carbon (*DOC*) and nutrients (*TN* and *TP*). The

second PC axis differentiated the Chicoutimi and Abitibi regions and was mainly related to

morphometric differences: lakes in the Abitibi region were shallower, with flatter catchments

and also with higher nutrient concentrations. Lakes in the Schefferville region were spread across

the second PC axis indicating that lake morphometry and catchment characteristics were highly

variable in this region.

The average percent occurrence (Figure 3a) was 24% for phytoplankton taxa (median

15%) and 21% for zooplankton taxa (Figure 3b, median 8%). Four phytoplankton taxa

Mallomonas (90%), *Cryptomonas* (90%), *Dinobryon* (82%) and *Asterionella* (63%), were

observed in more than 60% of lakes. For zooplankton *Leptodiaptomus* (89%), *Daphnia* (87%),

Bosmina (84%) and *Holopedium* (64%) had occurrences greater than 60%. Of the 56

phytoplankton taxa, 34 (61%) were observed in all three regions, while across all 27 zooplankton

taxa, 10 (37%) were observed in all regions. The differences in taxonomic composition between

regions were significant for both groups, but were less pronounced for phytoplankton (% correct

= 69%, p=0.01, Figure 4a) than for zooplankton (76%, p=0.01, Figure 4b).

For both phytoplankton and zooplankton, all functional traits were present in all three

regions. All phytoplankton functional traits occurred in more than 90% of lakes, with the

exception of two traits associated with cyanobacteria: presence of a vacuole for motility (63% of

lakes) and the potential to fix nitrogen (58% of lakes). The occurrence of zooplankton functional

traits ranged between 18% for carnivores and omnivore-herbivores, and 100% for herbivores,

with the average occurrence of zooplankton traits being 64% (median 75%). Functional

composition between regions did not differ for either phytoplankton, or zooplankton (% correct =

35%, p=0.69 and 49%, p=0.15 respectively, Figures 3c and 3d).

In the taxonomic Procrustes analyses we found a significant correlation between

phytoplankton and zooplankton taxa (Table 3), indicating a taxonomic coupling between the

distribution of the two groups. However, the correlation was not significant after controlling for

water quality, morphometry and space together, suggesting that the observed taxonomic coupling

cannot be attributed purely to trophic interactions but rather to both groups responding to the

same environmental and spatial drivers. On the other hand, using functional traits, there was a

significant coupling between the plankton groups only when traits related to trophic interactions

between phytoplankton and zooplankton (i.e. without phytoplankton resource acquisition traits;

including only motility, colonial and biovolume; Table 2) were used, but not when all traits were

considered (Table 3). The significant trait correlation did not remain after controlling for water

quality, morphometry or space (independently or together), indicating that the observed

correlation was again likely induced by one of these drivers and not by trophic interactions

directly.

Factors related to community composition

The RDA model explained 4% of the variation in phytoplankton taxonomic composition.

For functional composition , up to 27% of variation in the biomass proportion of the different

traits was explained in the multiple regressions (Figure 5), but no variation was associated with

the distribution of mixotrophy, non-motility, nor biovolume. We observed no direct effect of

spatial factors on phytoplankton functional trait variation, but a shared component between water

quality and spatial factors indicating that the water quality variables driving the distribution of

phytoplankton traits were spatially structured. Water quality variables were most consistently

explanatory factors of the phytoplankton functional traits for which variation was explained (blue

bars; Figure 5). The exceptions were for the flagellated trait, which responded to lake

morphometry, and the mixed pigment trait for which variation was shared between water quality

and morphometry. After forward selection, the RDA of phytoplankton taxonomic composition

was constrained by *TP*, *CDOM*, *pH* and lake area (Figure 6a). Based on the first axis, differences

in taxonomic composition could be mainly explained by lake nutrient status, pH and coloured

carbon content (*CDOM*). Similarly, functional composition also responded to nutrient status and

carbon content as well as lake temperature and chlorophyll *a* concentration according to the first

PC axis (Figure 6c). Functional traits related specifically to cyanobacteria (Pigment blue, N

fixation and Motility-V) were positively associated with the first axis, while traits (Pigment and

mixed brown, Si requiring, Motility-F) related to other key taxonomic groups (including

chrysophytes, cryptophytes and diatoms) were negatively associated. In the taxonomic and

functional RDA, the first two axes were significant.

For zooplankton, the global RDA models explained 11% of the variation in taxonomic

composition (Figure 5), and between 6% and 32% of the variation in the proportion of different

functional traits (Figure 5). Similarly, zooplankton variation was shared between water quality

and space, but spatial factors also independently explained a significant portion of the variation

in zooplankton taxonomic composition, B-filtration, D-filtration and size. For taxonomic

composition and most functional traits, a large portion of the variation was attributable to lake

morphometry (yellow bars; Figure 5). However, for most traits, and taxonomic composition,

some variation was either shared or explained independently by water quality.

In the subsequent RDA of zooplankton taxonomic composition, constrained by

morphometric and water quality variables (Figure 6b), variables related to lake productivity

(Chl*a*, *TP* and *TN*), carbon content (*DOC*) and temperature loaded on the first axis, while

variables relate to lake morphometry (lake depth and area) loaded more strongly on the second

axis. For functional traits (Figure 6d), the first axis was mainly related to lake depth, with the

second being related to lake area. The raptorial feeding trait was highly correlated with deep

lakes and C-filtration with shallow lakes. B-filtration was positively correlated with lake area and

size was negatively correlated with lake area while D-filtration was related to large and deep

lakes and Stationary suspension to small and shallow lakes. In the taxonomic and functional

RDA, the first two axes were significant.

Discussion

We observed a significant coupling between the landscape distributions of phytoplankton

and zooplankton at both taxonomic and functional levels. However, after taking into account the

coupling explained by environmental filtering and limitations to dispersal, we showed that in

boreal lakes, no residual coupling between phytoplankton and zooplankton could be attributed to

trophic interactions directly. Overall, the distribution of phytoplankton and zooplankton taxa and

traits was strongly influenced by water quality and lake morphometry and to a lesser extent by

dispersal limitation and these emerge as the main drivers coupling the distribution of

phytoplankton and zooplankton across boreal lakes. Moreover, phytoplankton and zooplankton

taxonomic composition displayed strong regional differences across the entire studied landscape,

but not when considered through a functional trait lens. Thus, although we observed important

taxonomic differences between regions, these differences did not translate into functional

differences between regions, despite the large environmental gradients covered.

Regional differences in composition and environment

Significant differences in phytoplankton and zooplankton taxonomic composition between the

three regions indicate a role for environmental factors or dispersal in influencing distributions at

the regional scale. On the other hand, functional composition did not differ between regions,

indicating that environmental variation or dispersal distances could not prevent any traits from

being distributed across the entire landscape. However, the biogeographical overlap in traits

does not mean that functional composition was similar across all lakes. Environmental factors

related to water quality and lake morphometry explained a significant amount of variation in

relative biomass of most functional traits across both plankton groups (Figure 5), indicating that

the control of functional composition acts at sub-regional scales. We also observed that control

of plankton taxonomic and functional composition by these lake characteristics was far more

important than was the effect of dispersal limitation. Finally, comparing across plankton groups,

dispersal limitation was more important for zooplankton than phytoplankton, supporting

previous work (Beisner et al. 2006, De Bie et al. 2012), but here now also including functional

traits.

Divergent responses of phytoplankton and zooplankton to their environment

Although both plankton groups responded strongly to environmental factors, the specific

variable types accounting for the most variation in taxonomic and functional composition

differed between phytoplankton and zooplankton. Phytoplankton taxonomic and functional trait

compositions responded most consistently and strongly to water quality (i.e. their proximal

environment). This is to be expected based on previously established strong relationships

between composition and lake nutrient status for broad taxonomic groups, groups which were

reflected in our functional (pigment) categorization (Watson et al. 1997). On the other hand,

zooplankton taxonomic and functional composition both responded more strongly to lake

morphometry. However,some water quality effects were also evident for zooplankton, indicating

overall an integrated response of zooplankton functional and taxonomic composition to their

proximal environment (water quality) and habitat characteristics (lake morphometry).

Zooplankton response to water quality could either occur as a direct effect, or as an

indirect bottom-up response to changes in phytoplankton community structure. For the more

prevalent response of zooplankton to lake morphometry, this reflects variation in habitat

characteristics, such as differences in lake physics (e.g. thermal stratification) or lake depth. We

consider it most likely that this morphometry effect arose indirectly, operating via the trophic

influence of fish predation, as previously observed (O'Brien et al. 2004). Specifically, lake depth

influences fish community composition (Jackson and Harvey 1989),with larger volume lakes

tending to have longer food chains (Post et al. 2000), thereby modulating the trophic cascade

effect on zooplankton through planktivore fish feeding (Carpenter et al. 1985). The variation in

zooplankton taxonomic and functional trait composition related to lake morphometry could thus

result from local variation in fish composition (data which we did not have). Further evidence

comes from the larger proportions of D-filtration in larger, deeper lakes: in larger lakes the

presence of an extra trophic level (total of 4- levels) decreases fish zooplanktivory, resulting in

reduced top-down pressure on large herbivorous cladocera, which are preferred prey for fish

(Christoffersen et al. 1993). Repercussions throughout the community were observable in the

ordination biplots (Figure 6d), with reductions in D-filtration being related to an increased

proportion of stationary suspension herbivory, dominated by calanoid copepods. Also, the

relative biomass of the C-filtration group was negatively related to lake depth, consistent with

the fact that most species within this functional feeding type are littoral species and shallow lakes

contain greater proportion of habitats that are littoral.

 Uncoupled biogeographies of phytoplankton and zooplankton

378   Using the common currency of functional traits, the biogeography of phytoplankton and

379 zooplankton were not coupled after controlling for environment and space. Even by only

380 including phytoplankton traits reflecting trophic interactions (grazer avoidance traits) no

381 coupling could be attributed to interactions between functional groups. Because no coupling

382 between phytoplankton and zooplankton was observed after controlling for either water quality,

383 morphometry or space, it suggests that one of these is the main driver of the observed coupling.

384 Because water quality was an important driver for most phytoplankton, as well as zooplankton

385 traits (see Figure 5), we can hypothesize that water quality was the main driver of the coupling

386 between phytoplankton and zooplankton traits. The lack of support for a coupling induced by

387 trophic interactions is surprising given the success of other functionally-based models to describe

388 coupled plankton dynamics through time (e.g. PEG model). It suggests that trophic interactions

389 are not consistent enough at the landscape level to result in a functional coupling of

390 phytoplankton and zooplankton.

391   Taxonomically, we also observed a significant coupling between phytoplankton and

392 zooplankton distribution and one that was stronger than observed functionally. The coupling

393 between phytoplankton and zooplankton taxonomy only became non-significant after controlling

394 for the combined effects of water quality, morphometry and space (Table 3), suggesting that the

395 observed coupling resulted from a combined effect of these three groups of drivers across boreal

396 lakes. While the greatest decrease in coupling was observed after controlling for space (Table 3),

397 spatial factors could not explain any independent variation in the distribution of phytoplankton

398 taxa (Figure 5). However, as most of the variation explained by water quality in the distribution

399 of phytoplankton and zooplankton (taxonomic) was spatially structured (light blue bars in

Figure 5) the observed spatial impact is likely not a result of dispersal limitation, but rather the

result of water quality variables that are spatially structured. Nevertheless, the coupling we

observed at the taxonomic level is consistent with Bowman and colleagues (2008) who observed

a spatial concordance between the community composition of phytoplankton and zooplankton in

eight Canadian lakes.

It is also important to acknowledge that the coupling between plankton groups that we

attribute to environmental factors could mask the effect of trophic interactions. For example, the

concordant taxonomic response to phosphorus (*TP*, Figure 6 a and b), could indicate that

phosphorus enrichment affecting phytoplankton triggers a response in the zooplankton, as would

be expected with changes in edibility of phytoplankton along this nutrient gradient (e.g. Watson

et al. 1997). Furthermore, for a large suite of environmental variables, spatial structure is likely

to vary differently across variable types, with for example, water quality variables generally

having stronger spatial structure than those related to lake morphometry (Lapierre et al. 2015).

This difference in spatial structuring could disrupt factors like trophic interactions or nutrients

that are likely to promote joint distributions. Proximate (water quality) variables with their strong

spatial structure across landscapes should therefore favour greater plankton biogeographical

concordance than would less structured habitat variables characterizing lake morphometry.

Unfortunately, mechanistic links such as these are difficult to verify with observational studies

like ours at these spatial scales, and would require smaller-scale experimental study to fully

verify.

The goal of including functional traits was to assess whether biogeographical coupling

would be more easily observed using a more mechanistic aggregation of organismal

differentiation than is done by pure taxonomy. To this end, we selected functional traits that are

directly representative of interactions between phytoplankton and zooplankton and in resource

acquisition. A lack of coupled biogeographies, after controlling for environmental variables was

observed. Hence, for the suite of feeding related functional traits used in this study, we found no

support for direct reciprocal influences of zooplankton and phytoplankton community structures

across the landscape. Overall, our study points to other food web components such as fish

predation and resource availability as independent drivers of each group's biogeography .

Overall, although phytoplankton and zooplankton are known to share strong trophic connections

in individual lakes (Porter 1977, Sterner 1989), which constitutes the basis of the main pathway

for matter and energy transfer in aquatic environments, these patterns do not seem to constrain

their joint biogeographical distributions. This result has implications for the predictability (or a

lack thereof) regarding large-scale change in plankton taxonomic or functional compositions

across lake landscapes with the spread of invasive species or the northward migration of species

with climate warming.

Acknowledgments

We thank Monique Arianne Resende and Marilyne Robidoux for help in the lab and field and

Roy Nahas for the GIS work. Funding from the Fonds de recherche du Québec - Nature et

Technologies (FRQNT) strategic clusters: Groupe de recherche interuniversitaire en limnologie

et en environnement aquatique (GRIL) and the Québec Centre for Biodiversity Science (QCBS)

is gratefully acknowledged. BEB and PDG acknowledge funding from the Natural Sciences and

Engineering Research Council of Canada (NSERC) in the form of Discovery Grants and a

Research Chair in Carbon Biogeochemistry of Boreal Aquatic Systems (CarBBAS) co-funded by

Hydro-Québec to PDG.

References

Anderson, M. and Willis, T. 2003. Canonical analysis of principal coordinates: a useful method of constrained ordination for ecology. - Ecology 84: 511–525.

Barnett, A. J. et al. 2007. Functional diversity of crustacean zooplankton communities: towards a trait-based classification. - Freshwater Biology 52: 796–813.

Beisner, B. E. et al. 2006. The role of environmental and spatial processes in structuring lake communities from bacteria to fish. - Ecology 87: 2985–91.

Borcard, D. et al. 1992. Partialling out the spatial component of ecological variation. - Ecology 73: 1045–1055.

Bowman, M. F. et al. 2008. Temporal and spatial concordance in community composition of phytoplankton, zooplankton, macroinvertebrate, crayfish, and fish on the Precambrian Shield. - Canadian Journal of Fisheries and Aquatic Sciences 65: 919–932.

Brown, J. et al. 2004. Toward a metabolic theory of ecology. - Ecology 85: 1771–1789.

Carpenter, S. R. et al. 1985. Cascading trophic interactions and lake productivity. - BioScience 35: 634–639.

Cattaneo, A. and Prairie, Y. T. 1995. Temporal variability in the chemical characteristics along the Riviere de l'Achigan: How many samples are necessary to describe stream chemistry? - Can. J. Fish. Aquatic Sci. 52: 828–835.

Christoffersen, K. et al. 1993. Potential role of fish predation and natural populations of zooplankton in structuring a plankton community in eutrophic lake water. - Limnol. Oceanogr. 38: 561–573.

Culver, D. A. et al. 1985. Biomass of Freshwater Crustacean Zooplankton from Length–Weight Regressions. - Canadian Journal of Fisheries and Aquatic Sciences 42: 1380–1390.

De Bie, T. et al. 2012. Body size and dispersal mode as key traits determining metacommunity structure of aquatic organisms. - Ecology Letters 15: 740–747.

Declerck, S. a. J. et al. 2011. Scale dependency of processes structuring metacommunities of cladocerans in temporary pools of High-Andes wetlands. - Ecography 34: 296–305.

Diamond, J. M. 1975. Assembly of species communities. - Ecology and evolution of communities 342: 444.

Ghadouani, A. et al. 2003. Effects of experimentally induced cyanobacterial blooms on crustacean zooplankton communities. - Freshwater Biology 48: 363–381.

Gotelli, N. J. et al. 2010. Macroecological signals of species interactions in the Danish avifauna. - PNAS 107: 5030–5035.

Hébert, M.-P. et al. 2016. A compilation of quantitative functional traits for marine and

freshwater crustacean zooplankton. - Ecology 97: 1081–1081.

Heino, J. et al. 2015. Metacommunity organisation, spatial extent and dispersal in aquatic systems: Patterns, processes and prospects. - Freshwater Biology 60: 845–869.

Hillebrand, H. et al. 1999. Biovolume calculation for pelagic and benthic microalgae. - Journal of Phycology 35: 403–424.

Jackson, D. A. and Harvey, H. H. 1989. Biogeographic associations in fish assemblages: local vs. regional processes. - Ecology 70: 1472–1484.

Jeziorski, A. et al. 2015. The jellification of north temperate lakes. - Proceedings of the Royal Society of London B: Biological Sciences in press.

Keller, W. and Pitblado, J. R. 1989. The distribution of crustacean zooplankton in Northern Ontario, Canada. - Journal of Biogeography 16: 249–259.

Knisely, K. and Geller, W. 1986. Selective feeding of four zooplankton species on natural lake phytoplankton. - Oecologia 69: 86–94.

Kraft, N. J. B. and Ackerly, D. D. 2014. Assembly of Plant Communities. - In: Monson, R. K. (ed), Ecology and the Environment. The Plant Sciences. Springer New York, pp. 67–88.

Lapierre, J. F. et al. 2015. Climate and landscape influence on indicators of lake carbon cycling through spatial patterns in dissolved organic carbon. - Global Change Biology 21: 4425–4435.

Legendre, P. and Legendre, L. 1998. Numerical ecology. - Elsevier Science.

Legendre, P. and Gallagher, E. D. 2001. Ecologically meaningful transformations for ordination of species data. - Oecologia 129: 271–280.

Litchman, E. et al. 2013. Trait-based approaches to zooplankton communities. - Journal of Plankton Research 35: 473–484.

Longhi, M. L. and Beisner, B. E. 2010. Patterns in taxonomic and functional diversity of lake phytoplankton. - Freshwater Biology 55: 1349–1366.

Lorenzen, C. J. 1967. Determination of chlorophyll and pheo-pigments: spectrophotometric equations. - Limnology and Oceanography 12: 343–346.

Mardia, K. et al. 1980. Multivariate analysis. - Academic press.

McCauley, E. 1984. The estimation of the abundance and biomass of zooplankton in samples. - A manual on methods for the assessment of secondary productivity in fresh waters. Blackwell Scientific, Oxford, UK: 228–265.

McQueen, D. J. et al. 1989. Bottom-up and top-down impacts on freshwater pelagic community structure. - Ecological Monographs 59: 289–309.

Nush, E. A. 1980. Comparation of different methods for clorophyll and pheopigment

determination. - Archiv für Hydrobiologie 14: 14–36.

O'Brien, J. W. et al. 2004. Physical, Chemical, and Biotic Effects on Arctic Zooplankton Communities and Diversity. - Limnology and Oceanography 49: 1250–1261.

Oksanen, J. et al. 2015. vegan: Community Ecology Package. R package version 2.3-0.

Padial, A. A. et al. 2014. Dispersal ability determines the role of environmental, spatial and temporal drivers of metacommunity structure. - PLoS ONE 9: 1–8.

Paffenhöfer, G. A. 1984. Food ingestion by the marine planktonic copepod Paracalanus in relation to abundance and size distribution of food. - Marine Biology 80: 323–333.

Pinel-Alloul, B. et al. 1990. Phytoplankton in Quebec Lakes: Variation with Lake Morphometry, and with Natural and Anthropogenic Acidification. - Canadian Journal of Fisheries and Aquatic Sciences 47: 1047–1057.

Pinel-Alloul, B. et al. 1995. Spatial and environmental components of freshwater zooplankton structure. - Ecoscience: 1–19.

Porter, K. G. 1977. The plant-animal interface in freshwater ecosystems. - American Scientist 65: 159 – 170.

Post, D. M. et al. 2000. Ecosystem size determines food-chain length in lakes. - Nature 405: 1047–1049.

Roy, G. 2012. The Great Geological Domains of Québec.

Soininen, J. et al. 2007. Neutrality, niches, and determinants of plankton metacommunity structure across boreal wetland ponds. - Ecoscience 14: 146–154.

Soininen, J. et al. 2011. Disentangling the spatial patterns in community composition of prokaryotic and eukaryotic lake plankton. - Limnology and Oceanography 56: 508–520.

Sommer, U. et al. 1986. The PEG-model of seasonal succession of planktonic events in fresh waters. - Archiv für Hydrobiologie 106: 433–471.

Sommer, U. et al. 2012. Beyond the Plankton Ecology Group (PEG) Model: Mechanisms Driving Plankton Succession. - Annual Review of Ecology, Evolution, and Systematics 43: 429–448.

Stekhoven, D. J. and Bühlmann, P. 2012. Missforest-Non-parametric missing value imputation for mixed-type data. - Bioinformatics 28: 112–118.

Sterner, R. W. 1989. The role of grazers in phytoplankton succession. - In: Plankton ecology. Springer, pp. 107–170.

Stomp, M. et al. 2011. Large-scale biodiversity patterns in freshwater phytoplankton. - Ecology 92: 2096–2107.

Vanderploeg, H. A. 1981. Seasonal Particle-Size Selection by Diaptomus sicilis in Offshore

Lake Michigan. - Canadian Journal of Fisheries and Aquatic Sciences 38: 504–517.

Watson, S. B. et al. 1997. Patterns in Phytoplankton Taxonomic Composition Across Temperate Lakes of Differing Nutrient Status. - Limnology and Oceanography 42: 487–495.

Wiens, J. J. 2011. The niche, biogeography and species interactions. - Philosophical transactions of the Royal Society of London. Series B, Biological sciences 366: 2336–2350.

Wissel, B. et al. 2003. Effects of water color on predation regimes and zooplankton assemblages in freshwater lakes. - Limnology and Oceanography 48: 1965–1976.

Table 1: Means and standard error (SE, in parenthesis) of environmental variables: water quality (WQ), and morphometry (M) with the number of missing values (#NA) and the transformation used.

Variable name (abbreviation)	Category	# NA	Units	Transformation	Regional averages (Standard Error)					
					Abitibi		Chicoutimi		Schefferville	
					Phyto.	Zoo.	Phyto.	Zoo.	Phyto.	Zoo.
Chlorophyll a (*Chl-a*)	WQ	0	ug/L	$\log_{10}$	3.1 (0.7)	3.7 (0.9)	1.9 (0.2)	2.0 (0.1)	0.9 (0.1)	0.8 (0.1)
Total Phosphorus (TP)	WQ	0	ug/L	$\log_{10}$	17.0 (3)	18.5 (3.5)	9.8 (0.9)	10.1 (0.6)	7.3 (0.7)	7.7 (0.6)
Total Nitrogen (TN)	WQ	0	mg/L	$\log_{10}$	0.3 (0.05)	0.3 (0.04)	0.2 (0.02)	0.2 (0.01)	0.2 (0.02)	0.2 (0.01)
Dissolved organic carbon (DOC)	WQ	2	mg/L	$\log_{10}$	12.4 (1.4)	12.4 (1.3)	7.5 (0.7)	7.7 (0.4)	3.8 (0.4)	4.0 (0.2)
Colored dissolved organic matter (CDOM)	WQ	1	1/m	$\log_{10}$	3.2 (0.7)	3.4 (0.8)	3.9 (0.5)	4.0 (0.4)	1.0 (0.2)	1.1 (0.1)
pH	WQ	1	-	-	7.4 (0.3)	7.5 (0.3)	6.9 (0.1)	7.0 (0.1)	6.8 (0.2)	7.1 (0.1)
Water temperature (0.5 m) (T_{water})	WQ	1	ºC	-	21.8 (1.1)	20.6 (1.0)	20.0 (0.5)	19.8 (0.3)	14.2 (0.2)	15.0 (0.2)
Maximum depth (Zmax)	M	0	m	$\log_{10}$	5.7 (1.4)	5.7 (1.2)	8.2 (1.5)	9.4 (1.1)	7.3 (1.5)	6.8 (0.9)
Lake Area (Area)	M	0	m^2	$\log_{10}$	5.89 (0.3)	5.7 (0.3)	6.1 (0.3)	6.2 (0.2)	6.0 (0.3)	6.1 (0.2)
Number of lakes					11	13	16	44	21	46

Table 2: Phytoplankton and zooplankton functional traits used in this study.

	Category	Traits	Values
Phytoplankton	Resource acquisition	Nitrogen fixation	yes/no
		Si requirement	yes/no
		Mixotroph	yes/no
		Pigments	Green
			Blue
			Brown
			Mixed
	Grazer avoidance	Motility	None
			F(Flagellated)
			V(Vacuole)
		Biovolume	numerical
		Colonial	yes/no
Zooplankton	Resource acquisition	Trophic Position	Carnivore
			Omnivore-Carnivore
			Omnivore
			Omnivore-Herbivores
			Herbivores
			Immature
		Feeding Type	B (Bosmina)-filtration
			C (Chydorus)-filtration
			D (Daphnia)-filtration
			S (Sidae)-filtration
			Stationary Suspension
			Raptorial

Table 3: Procrustes rotation analysis of species and the trait dataset correlation coefficients for phytoplankton (Phyto) and zooplankton (Zoo) after controlling (|) for water quality (WQ), morphometry (M) and space (S). Where * = 0.01<p<0.05, ** = 0.001<p<0.01,*** p<0.001, ns = non-significant.

	Control for	m12 squared	Correlation
Taxonomic	$Zoo_{taxo} \leftrightarrow Phyto_{taxo}$	0.85	*0.38****
	|wq	0.86	*0.37***
	|m	0.86	*0.37****
	|S	0.89	*0.34***
	|wq+m+S	0.92	ns
All traits	$Zoo_{funct} \leftrightarrow Phyto_{funct}$	0.92	ns
	|wq	0.92	ns
	|m	0.9	ns
	|S	0.92	ns
	|wq+m+S	0.93	ns
Without phytoplankton resource acquisition traits	$Zoo_{funct} \leftrightarrow Phyto_{funct}$	0.92	0.28*
	|wq	0.95	ns
	|m	0.94	ns
	|S	0.94	ns
	|wq+m+S	0.94	ns

Figure 1: (a) Map of sampled lakes across three regions of boreal Quebec, Canada. (b) Lakes

environmental principal component analysis (PCA) based on water quality and

morphometric variables where the different symbols represent the three sampled

regions. . Abbreviations are as follows total phosphorus (TP); total nitrogen (TN);

dissolved organic carbon (DOC); chlorophyll a (*Chl-a*); coloured dissolved organic

matter (CDOM); water temperature (Twater); lake maximum depth (Zmax); lake area

(Area).

Figure 2: Conceptual framework used to test for a significant coupling between zooplankton and phytoplankton communities using taxonomy and functional traits, while controlling for environmental factors (water quality WQ, morphometry M) and dispersal limitation (space S).

Figure 3: Ranked occurrence (percent of all lakes) of (a) phytoplankton and (b) zooplankton taxa.Bars and dots were coloured by Pigment trait type for phytoplankton and the Feeding strategy trait for zooplankton.

Figure 4: Two-dimensional scatter plot of canonical axes of the CAP ordinations for (a) phytoplankton and (b) zooplankton taxonomic composition and their respective functional trait compositions (c, d). Symbols represent the different regions. Strong regional differences occurred for taxonomic composition of phytoplankton (a, p=0.01) and zooplankton (b, p=0.01), but were not significant for functional trait composition (c and d).

Figure 5: Variation explained (adjusted R^2) by environmental and spatial factors for phytoplankton (left panel) and zooplankton (right panel) taxonomic composition and each functional trait. Only regression and RDA models that were globally significant are displayed. The variation that was independently explained by a group of variables is represented if significant, and shared variation represents the sum of variation shared between any of the groups of variables, represented with a +. Abbreviations are as follows : morphometry (M) ; water quality (WQ) ; space (S).

Figure 6: RDA ordination triplots of the (a, c) phytoplankton and (b,d) zooplankton

composition classified by (a,b) taxonomy and (c,d) functional (c, d) traits. The RDA

was constrained by environmental variables related to water quality (in blue) and by

variables related to lake morphometry (in yellow). Taxa and functional trait response

variables are represented by crosses and lakes by dots.s. In the taxonomic RDAs (a, b),

only taxa with loadings over 0.20 are displayed. Abbreviations are as follows total

phosphorus (TP); total nitrogen (TN); dissolved organic carbon (DOC); chlorophyll a

(*Chl-a*); coloured dissolved organic matter (CDOM); water temperature (T_{water}); lake

maximum depth (Zmax); lake area (Area).