The soil microbial food web revisited with metatranscriptomics - predatory

***Myxobacteria* as keystone taxon?**

Sebastian Petters[1], Andrea Söllinger[1,2], Mia Maria Bengtsson[1], Tim Urich[1,*]

[1]Institue of Microbiology, Center for Functional Genomics of Microbes, University of

Greifswald, Greifswald, Germany

[2]Department of Ecogenomics and Systems Biology, University of Vienna, Vienna,

Austria

* corresponding author

email: tim.urich@uni-greifswald.de

Abstract

Trophic interactions in the microbial food web of soils are crucial for nutrient and

carbon cycling. Traditionally, protozoa are considered the major micropredators of

bacteria in soil. However, some prokaryotes, such as *Myxobacteria* and *Bdellovibrio*

are also famous for bacterivorous life style. Until recently, it was impossible to assess

the abundance of pro- and eukaryotic micropredators in soils simultaneously. Using a

metatranscriptomic three-domain profiling of small subunit ribosomal RNA we

investigated the abundance of bacterivores in 28 datasets from eleven European

mineral and organic soils of different climatic zones. In all soils, *Myxobacteria*

comprised a significant proportion from 4 – 19% of prokaryotic 16S rRNA transcripts

and more than 60% of all bacterivores in most soils. *Haliangiaceae* and

Polyangiaceae were most abundant, while the name-giving *Myxococcaceae* were

barely present. Other bacterial predators like *Bdellovibrio* were low abundant. Also

Protozoan micropredator 18S rRNA transcripts, e.g. from *Cercozoa, Amoebozoa and*

Ciliophora, were on average less abundant, especially in mineral soils. *Nematodes*

were even less abundant. In addition, we applied a longitudinal approach to identify

bacterivores during beech litter colonisation. Here, *Myxobacteria* showed prey-

dependent, protozoa-like community dynamics during colonisation. Thus, their broad

prey range and high abundance suggests a major influence of *Myxobacteria* on

structuring the prokaryotic community composition in soil, and might warrant their

classification as keystone taxon. Our results suggest the presence of an ecologically

important "bacterial loop" in soil food webs, independent of protozoa and nematodes.

Introduction

Predation of predators on prey is a key process in structuring community composition in ecosystems and in maintaining high biodiversity. Predator - prey interactions and dynamics among animals and consequences for ecosystem functioning have been studied extensively since the early days of ecology. While less visible and thus less acknowledged, predation is not foreign to the microbial world. Eukaryotic as well as prokaryotic microorganisms are known to prey on other microorganisms in marine, aquatic and terrestrial habitats as part of the microbial food web (Clarholm, 1985; Azam et al., 1983).

Protozoa are traditionally considered the main microbial predators of bacteria, a notion that stems from the fact that, unlike in bacteria, where it is somewhat "exotic", predation is a common lifestyle among protozoa. Predatory protozoa are known from both aquatic and soil environments and have been considered a key-component of the "microbial loop" responsible for the remineralisation of nutrients (Bonkowski, 2004; Clarholm, 1985). Whereas protozoa in the aquatic system have been well characterised, both in terms of their identity and population size, research in soils has been much more hampered, since no adequate molecular tools have been available for a long time, cultivation is often difficult, and direct microscopic observations are impossible (Geisen et al., 2015).

Much fewer prokaryotic species are considered predatory, although a predatory lifestyle in prokaryotes probably evolved prior to its development in eukaryotes. Several bacterial predators have been identified, with more and more taxa exhibiting a predatory lifestyle being recognized recently. These include *Myxobacteria*, *Lysobacter*, *Bdellovibrio* and like organisms (BLO), *Vampirococcus*, and

Dapterobacter, among others (Reichenbach, 1999). Especially the *Myxobacteria*, with

their 'wolf pack hunting' strategy, are known micropredators since more than 70 years

ago and have been isolated from soils world-wide (Keane & Berleman, 2016;

Reichenbach, 1999).

It has been until recently impossible to assess bacterial and protist community

composition with the same methodology. Although PCR amplicon approaches

enabled the study of both groups separately, a direct comparison of their relative

abundances was not possible due to the absence of universal primers that would tackle

all groups without bias. However, these obstacles are avoided when applying random

hexamer-primed reverse transcription as in metatranscriptomics approaches that target

SSU rRNA of organisms from all three domains of life (Urich et al., 2008).

Furthermore, these rRNA transcripts are indicative of ribosomes and thus are likely

derived from metabolically active cells and can be considered markers for living

biomass. The generated cDNA fragments originate from different regions of the SSU

rRNA molecule unlike PCR primed specific sites, and are therefore insensitive to the

presence of introns or primer mismatches, when PCR primers are applied.

We have recently used this PCR-free metatranscriptomics approach to reveal the

diversity of the active soil protist communities within five different natural soil

systems in Europe, including forest, grassland and peat soils as well as beech litter

(Geisen et al., 2015).

Here we have focused on other groups of microbial predators - predatory bacteria. We

have assessed the relative abundance of SSU rRNAs from bacterial groups known to

exhibit a predatory lifestyle in these soils. Metatranscriptomics enabled the direct

comparison of SSU rRNA transcripts from bacterial and protozoan micropredators

and revealed that potentially predatory bacteria, especially *Myxobacteria*, were

abundantly detected in all soils, while protozoa abundances were much more variable.

The underlying causes and consequences for our perception of microbial predation in

soils are discussed and an alternative model of the soil microbial loop is put forward.

Material and Methods

Data acquisition

The investigated metatranscriptomes had been obtained from different previous

studies on a range of European soils (Table 1). These included 4 samples from organic

peatland, 3 samples from organic floodplain, 3 samples from gleic fluvisol, 3 samples

from mineral grassland, 2 samples from organic forest litter, 4 samples from mineral

forest soil, and 3 samples each from 3 different mineral shrubland soils. RNA, cDNA

and sequences were obtained as previouly described (Beulig et al., 2016; Epelde et al.,

2015; Geisen et al., 2015; Tveit et al., 2013; Urich et al., 2008).

Furthermore, metatrascriptomic data were obtained from four different beech litter

types (K, A, O and S), which had been incubated with the same microbial community

in mesocosms (see Wanek et al., 2010 for details of the experimental setup). Litter

samples were taken at three time points: after two weeks and after three and six

months after inoculation, flash-frozen in liquid nitrogen and stored at -80 °C. RNA

was extracted and double-stranded cDNA was prepared as previously described

(Urich et al., 2008). 454 pyrosequencing was performed at the Norwegian Sequencing

center, CEFS, University of Oslo (Norway). Raw sequence data were submitted to the

NCBI Sequence Read Archive (SRA) under the accession number SRP134247.

Bioinformatic analysis

Raw sequence datasets were filtered to a minimum length of 200 nucleotides and a

minimum mean quality score of 25 using prinseq-lite (Schmieder & Edwards, 2011).

SSU rRNA sequences were identified via SortMeRNA (Kopylova et al., 2012).

USEARCH (Edgar, 2010) was used to randomly subsample datasets to a maximum of

50 000 - 100 000 sequences. The datasets were mapped against the CREST database

silva123.1 by blastn (Altschul et al., 1990; Lanzén et al., 2012). The obtained blastn

files were taxonomically analysed using MEGAN (Huson et al., 2011, min score 155;

top percent 2.0; min support 1). The number of SSU rRNA reads of the investigated

organisms was normalized in MEGAN to the total number of read counts.

Investigated taxa with predatory lifestyle were *Myxococcales*, *Bdellovibrionales*,

Lysobacter, *Dapterobacter*, *Vampirococcus*, *Amoebozoa*, *Cercozoa*, *Ciliophora*,

Foraminifera, *Euglenozoa*, *Heterolobosea*, and *Nematoda*. Different Nematoda taxa

were not investigated separately. The read counts of the analysed bacterial

micropredators where substructed from the total bacterial SSU rRNA resulting in prey

bacterial rRNA. The read counts of each analysed bacterivorous group were then

normalized to the prey bacterial SSU rRNA reads.

Results for organic, excluding mofette (MO) samples, and mineral soils were tested

for differentially expressed sequences with the R package edgeR (McCarthy et al.,

2012; functions glmFit and glmLRT), using the non-normalized total read counts

MEGAN file.

Results

Abundance of bacterivores in soil microbiomes

We screened the SSU (16S and 18S) rRNA fraction of 28 soil metatranscriptome

datasets obtained from eleven different soils across Europe (Table 1) for bacterivorous

pro- and eukaryotes. *Myxococcales* SSU rRNA reads comprised a high proportion of

prey bacterial SSU rRNAs, ranging from 3.5 to 18.9% (9% on average), and higher

than all other investigated bacterivores (Figure 1a). Their highest proportion in

relation to bacteria was detected in peat soils. Additionally, an organic fluvisol and a

beach litter layer showed *Myxococcales* abundances above 10%. The latter came up as

the only exception in the pattern, i.e. here the Protozoa were the most abundant

bacterivorous group. Overall, SSU rRNAs of protozoa were the second most abundant

(Figure 1a). Like the *Myxococcales* they were generally more abundant in organic

soils than in mineral soils. The only two cases where their proportion was above 10%

of prey bacterial SSU rRNA reads were a peatland and a forest litter sample. While

Myxococcales abundance never dropped below 3.4%, protozoa abundance was much

lower in mineral soils (down to 0.7%). The third most abundant group were the

Nematoda (Figure 1a). They showed greater variation in abundance compared to the

aforementioned taxa, especially in organic soils, where they showed both their highest

abundance (8.5%), namely in the forest litter horizon, and also their lowest abundance

(< 0.1%), which occurred in the suboxic mofette soil. This was the only sampling site,

where their abundance dropped below 0.1% of the prey bacterial SSU rRNA reads.

The only other soils which showed *Nematoda* abundances above 1% were the organic

peatland samples and the mineral Rothamsted soil. All mineral soils showed fractions

of *Nematoda* SSU rRNAs within 0.1 – 1%. The *Bdellovibrionales* comprised even

lower SSU rRNA abundances. Similar to the aforementioned, highest relative

abundance of *Bdellovibrionales* was observed in organic soils (0.9% in peatland soil).

We did not detect *Vampirococcus* and *Dapterobacter* in any of the investigated

samples. *Lysobacter* comprised the lowest SSU rRNA abundances of all detected

micropredators, namely 0.07% or lower.

Looking at the composition of potential prey bacteria revealed that SSU rRNA from

gram-negative bacteria comprised approximately 80%, while SSU rRNA from gram-

positive were approximately 20%. The latter were slightly more abundant in mineral

soils (supplementary figure S1).

Myxococcales dominate bacterivorous taxa

Comparing all investigated bacterivorous groups, the *Myxococcales* were highest

abundant in every sampling site, except for the forest litter (Figure 1b). In fact, in nine

of the eleven sites, including all mineral sampling sites, the proportion of

Myxococcales SSU rRNAs was more than 60% of all micropredators. In the forest

litter their proportion of the bacterivorous groups was below 30%. Correspondingly,

the protozoa were the most abundant group in that site, comprising up to more than

half of all bacterial predators. However, in all the other sampled sites, the proportion

of the protozoa was below 40%, in three cases even below 20%. Those were namely

the organic mofette samples as well as the mineral Rothamsted site, and mine M from

Spain, where the lowest percentage of all micropredators was observed. All of the

sampling sites had *Nematoda* SSU rRNA below 20% Figure 1b). Their highest

proportions occurred in the organic forest litter samples. Moreover, the only other two

sites where their proportions were above 10%, were Rothamsted, where they were

even more than the protozoa, and mineral mine H. All other sites showed proportions

below 10%, with the lowest proportions in samples from mofette. The

Bdellovibrionales were below 10% of micropredators in all sampling sites

(Figure 1b).

Community composition of Myxobacteria

We analysed the community composition of *Myxococcales* in more detail (Figure 2a).

The most dominant family was *Haliangiaceae, followed by Polyangiaceae* and

Blrii41, a family level group in the SILVA taxonomy that is currently devoid of

cultured representatives. These three together comprised more than 2/3 of

Myxobacteria SSU rRNAs in all but one site. *Haliangiaceae* and *Polyangiaceae* were

more abundant in mineral soils, while Blrii41 was more characteristic for organic

soils. The name-giving family *Myxococcaceae*, which is comprised, among others, of

the most frequently isolated genera *Myxococcus* and *Corallococcus*, was barely

present.

Protozoa community composition

As previously found (Geisen et al., 2015), *Amoebozoa*, *Cercozoa* and *Ciliophora* were

the three most abundant protist groups (Figure 2b). While *Amoebozoa* and *Cercozoa*

dominated in mineral soils, the *Ciliophora* were most abundant in organic soils. The

remaining predatory groups *Foraminifera*, *Euglenozoa*, and *Heterolobosea* accounted

for low abundances on average. The mofette reference (MR) samples were an

exception, with *Foraminifera* comprising more than 20% of protists.

Dominance of Myxobacteria among bacterivores in mineral soils

We compared the average micropredator abundance (normalized to the prey bacteria)

between mineral and organic soils (excluding mofette samples) based on SSU rRNA

reads (Figure 3). *Lysobacter* data are not shown due to low abundances. Remarkably,

micropredator SSU rRNAs comprised 32.3% of prey SSU rRNAs in organic soils, as

compared to only 7.9% in mineral soils. Although concomitantly lower in abundance

in mineral soils, *Myxococcales* comprised the highest micropredator proportions in

both soil types (16.1% in organic vs. 5.7% in mineral soil). While protozoa were

almost equally abundant in organic soil, they comprised approx. 1/4 of *Myxobacteria*

in mineral soils. In fact, the percentage of *Myxococcales* within the bacterivores was

remarkably higher in mineral soils, i.e. 72% compared to 50% in organic soils. Thus,

the decrease in abundance of *Myxococcales* in mineral soils was not as strong as seen

in the other micropredators.

To statistically verify the observed differences in abundance between organic and

mineral soils, we tested the data for differentially expressed SSU rRNAs of

micropredators. While the protozoa ($p < 0.01$), *Lysobacter* ($p < 0.01$), and

Bdellovibrionales ($p = 0.02$) were significantly differently abundant between organic

and mineral soils, no significant differences were detected for *Myxococcales*

($p = 0.31$) and *Nematoda* ($p = 0.78$). This supports the observed phenomenon, where

the *Myxococcales* remained dominant in mineral soils, while the SSU rRNA

abundances of other investigated micropredators significantly decreased in mineral

soils.

Temporal dynamics of bacterivores during community succession

It has been shown that *Myxobacteria* can have a saprotrophic life style next to

bacterivory (reviewed in Reichenbach, 1999). We therefore analysed micropredator

dynamics in a litter colonisation experiment. In a longitudinal experiment four types

of sterilized beech litter differing in their C:N:P ratio were colonized by the same

microbial community taken from beech forest soil (Supplementary Table 1 in Wanek

et al., 2010). Metatranscriptome data were obtained from three time points over the

course of six months. SSU rRNA abundances of protozoa, *Nematoda*, and

Myxococcales, as well as total bacteria and fungi were assessed to investigate their

temporal dynamics (Figure 4). Microbiomes on litters K and S, which had a higher

nitrogen content, were strongly dominated by fungal SSU rRNAs as compared to

bacterial rRNAs, while litters A and O, which had lower nitrogen content, had higher

proportions of bacterial reads. The fungal:bacterial ratio stayed rather constant for

each litter type over time. SSU rRNAs of bacterivores generally increased in relative

abundance over time, especially from two weeks to three months. Remarkably, the

bacterivores (including *Myxococcales*) appeared earlier and in higher relative

abundance in litters with more prey bacteria. With few exceptions, protozoa

comprised the most abundant predator of bacteria, with *Myxococcales* and *Nematoda*

being second and third most abundant respectively. It appeared that after three months

a rather stable predator:prey ratio had established.

Discussion

Metatranscriptomics-enabled holistic assessment of soil micropredators

It has been until recently impossible to assess soil bacterial and protist community

composition with the same methodology. Although PCR amplicon approaches

enabled the study of both groups separately, a direct comparison of their relative

abundances was not possible due to the absence of universal primers that would tackle

all groups without bias. The rRNA fraction of metatranscriptomics data enables broad

three-domain community profiling of abundant bacteria, archaea, and eukaryotes via

rRNA (Urich et al., 2008). This innovative approach has constantly been developed

further and recently come to maturation due to lower-cost NGS sequencing

technologies and bioinformatic tools (e.g. (Bengtsson et al., 2018; Schwab et al.,

2014; Tveit et al., 2015)). Using this approach, we have recently created a first

molecular census of active protists in soils (Geisen et al., 2015). The study showed

that protozoa usually not detected with general PCR primers such as *Amoebozoa* and

Foraminifera are abundantly present and provided the most comprehensive picture of

active protist communities in soils to date. The strength of rRNA transcripts for

comparatively unbiased views into community composition and assessment of unseen

microbial diversity has recently gained popularity (e.g., Karst et al., 2018).

Predatory Myxobacteria as key-stone taxon in mineral soils?

In all investigated soils the *Myxobacteria* comprised a significant proportion of the

overall bacterial SSU rRNA transcripts. This confirms a recent PCR / 16S rRNA gene

based survey where *Myxobacteria* also comprised a substantial fraction (4.1%; Zhou

et al., 2014). In our study, lower micropredator abundances were detected in mineral

soils as compared to organic soils, which may be due to less available carbon resulting

in lower prey cell density in mineral soils. However, given the predatory traits of most

myxobacterial taxa, their high relative abundance hints towards an important role in

the microbial food web of soils. Moreover, with only one exception, the

Myxobacteria exhibited the highest abundances when compared to all other

bacterivores.

Traditionally, protists are considered to be the dominant group preying on bacteria

(e.g. Geisen et al., 2016; Trap et al., 2016). In contrast to this, our data suggest an

importance, possibly even dominance of *Myxococcales*. In fact, *Myxococcales*

comprised approx. 3/4 of all micropredators in mineral soils. Possibly, the smaller

pore sizes in mineral soils provide restricted access of protists to their bacterial prey,

as compared to the smaller *Myxococcales*. Thus, microorganisms inhabiting non-

continuous capillary pores could be protected from predation by Protozoa and

Nematoda, but not from the similarly-sized *Myxobacteria*. The prokaryotes inhabiting

the organic soil horizons, with unprotected macro-pore space, would in turn be

subjected to higher grazing pressure. The *Myxobacteria* and protozoa exhibit

fundamentally different predation strategies, with the much smaller *Myxobacteria*

being famous for their social 'wolf-pack' hunting combined with the secretion of lytic

enzymes, as compared to the larger phagotrophic protozoa (Reichenbach, 1999). The

more similar cell size of *Myxobacteria* and prey bacteria could thus favour

myxobacterial predation in mineral soils with small pores. Given the broad prey range

of *Myxobacteria*, their high abundance in soils suggests a major influence on

structuring the prokaryotic community composition, and might warrant their

classification as key-stone taxon.

Interestingly, the majority of *Myxobacteria* was not related to the well-studied and

easy-to-isolate *Myxococcaceae*, but belonged to families with a less well

characterised prey spectrum, such as *Haliangiaceae*, *Kofleriaceae*, and

Polyangiaceae (see Figure 2a), similar to findings of Zhou et al. (2014). The data in

this study hint toward biases in culturability within the *Myxococcale*s. In fact, one

large family-level group abundant in organic soils, represented in the SILVA database

by clone Blrii41, is currently without any cultured representatives.

A bacterial loop within the microbial loop

The soil survey could not give direct proof of whether the *Myxobacteria* (or any

presumed micropredator) actually showed bacterivorous behavior *in situ*. In a

colonisation experiment with sterilised beech litter we compared their succession with

other bacterivores, such as the protozoa. In general, the abundance of potential

bacterivores was positively associated with abundance of prey bacteria, and increased

over time indicating a developing food web during litter colonisation. *Myxococcales*

and protozoa abundances developed similarly over time. This observation hints at

Myxococcales indeed having a predominantly predatory and not saprotrophic lifestyle,

thus rendering the *Myxococcales* a prominent predatory taxon feeding on other

bacteria.

There is direct *in situ* evidence for myxobacterial bacterivory from RNA-stable

isotope probing studies. The hallmark study of Lueders and colleagues

(2006) introduced the use of isotopically labelled prey bacteria to target the general

diversity of micropredators in soil *in situ* and follow the carbon flow through the

bacterial channel of the soil food web. The authors reported the detection of labelled

sequences related to *Myxococcus*, *Lysobacter,* and *Bacteroidetes*. However, due to the

limited technology at that time, they were not able to assess the contribution of

predatory protozoa. Another shortcoming was the use of *E. coli* cells as prey because

the survival of *E. coli* cells added to soil is rather low. In a more recent follow-up

study with labelled native soil bacteria, Zhang and Lueders (2017) provided evidence

for niche partitioning between bacterial and eukaryotic micropredators in soil, driven

by the soil compartment. Interestingly, the *Myxobacteria* preyed on both gram-

positive and gram-negative bacteria. Like in their previous study, the relative

contribution of pro- and eukaryotic micropredators could not be assessed due to

methodological limitations. Another recent study also included fungi and bacteria

(Kramer et al., 2016), and indicated that the commonly accepted split of energy

channels does not exist.

The dominance of *Myxobacteria*, especially in mineral soils, suggests their important

role in the soil microbial food web (Figure 5). The so-called microbial loop in soil

(Bonkowski, 2004) is important for the remineralisation of nutrients, where especially

protozoa and *Nematoda* feed on bacteria and by this set free nutrients, which are in

turn provided for the bacteria as well as plants (Coleman, 1994). Our observations

hint at the presence of an ecologically important 'bacterial loop', especially in mineral

soils, within the prokaryotes and independent of protozoa, that has been overlooked

until today (Figure 5). As a consequence, bacterial micropredators might not only be

important for shaping microbial communities but might also prove to be important for

the recycling of nutrients in soils, as it has been shown for protozoa (Bonkowski,

2004; Koller et al., 2013), and thus potentially for nutrient and carbon cycling.

Although the *Myxococcales* order comprised a significantly high proportion of

bacterial SSU rRNA in all sites, differences at family-level were observed among the

soils. Nevertheless, the two most abundant families were the *Polyangiaceae* and

Haliangiaceae. The high dominance of the latter may well be due to the

Haliangiaceae taxon being clustered together with the *Kofleriaceae* taxon in the

current SILVA database. Interestingly the name-giving group, the *Myxococcaceae,*

comprised a very low proportion of all *Myxococcales*. The *Myxococcaceae* are known

to be easily cultivable from a variety of environmental samples. Our data show that

other, less well characterised families are in fact much more abundant in soil. Thus,

efforts should be undertaken to investigate their biology and in particular their prey

spectrum.

Methodological considerations

The micropredator abundance data in this study are derived from the abundance of

SSU rRNA in metatranscriptomes. This does not reflect organismic abundance but is

rather a proxy of living biomass (Urich & Schleper, 2011). In fact, several factor need

to be taken into account when comparing the SSU rRNA from different pro- and

eukaryotic organisms. Results of various studies suggest differences in RNA contents

per biomass (1) between organisms and (2) between growth phases, respectively. The

ribosome density in prokaryotic cells is generally considered higher than in

eukaryotes. However, few data are available to our knowledge. The RNA content of

E. coli was determined to be 15.7% of dry mass (dm) (Stouthamer & van

Leeuwenhoek, 39,545-565, 1973), of *Bacillus subtilis* between 8.5% and 14% dm^{-1}

(Tempest et al., 1968), *Saccaromyces cerevisiae* 23% dm^{-1} (Parada & Acevedo, 1983),

Aspergillus 5.9% (Carlsen et al., 2000) and *Penicillium chrysogenum* between 5% and

8% (Henriksen et al., 1996). Furthermore, prokaryotic cells in an exponential growth

phase are known to contain more RNA than cells in stationary phase (e.g. Tempest et

al., 1968). Preliminary data (Petters & Urich, unpublished) hint to correction factors

to be applied when comparing rRNA based abundances from metatranscriptomes

between pro- and eukaroytes. Nevertheless, a recent study showed that rRNA

correlates better with cell counts than ribosomal RNA genes (Giner et al., 2016).

Thus, our metatranscriptomics data identify the predatory *Myxobacteria* as important

players in the midst of the soil food web and suggests a prominent role in the soil

microbial loop in particular.

Acknowledgements

Andreas Richter, Wolfgang Wanek and colleagues (University of Vienna) are thanked

for providing beech litter samples. Daniela Teichmann and Sylvia Klaubauf

(University of Vienna) are acknowledged for excellent technical assistance in RNA

extraction. We thank Ave Tooming-Klunderud, Lex Nederbragt and others at the

Norwegian High-Throughput Sequencing Centre (University of Oslo) for 454

pyrosequencing. Andrea Söllinger acknoledges funding from the University of Vienna

(uni:docs) and the OeAD (Austrian agency for international mobility and cooperation

in education, science and research; Marietta-Blau-Fellowship).

Authors' contributions. The study was designed by TU. Data analysis was

performed by SP and TU, supported by AS and MB. The manuscript was written by

SP and TU, assisted by all co-authors.

References

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic

local alignment search tool. *Journal of Molecular Biology*, *215*(3), 403–10.

http://doi.org/10.1016/S0022-2836(05)80360-2

Azam, F., Fenchel, T., Field, J., Gray, J., Meyer-Reil, L., & Thingstad, F. (1983). The
Ecological Role of Water-Column Microbes in the Sea. *Marine Ecology*
Progress Series, *10*, 257–263. http://doi.org/10.3354/meps010257

Bengtsson, M. M., Wagner, K., Schwab, C., Urich, T., & Battin, T. J. (2018). Light
availability impacts structure and function of phototrophic stream biofilms across
domains and trophic levels. *Molecular Ecology*, *27*(14), 2913–2925.
http://doi.org/10.1111/MEC.14696

Beulig, F., Urich, T., Nowak, M., Trumbore, S. E., Gleixner, G., Gilfillan, G. D., …
Küsel, K. (2016). Altered carbon turnover processes and microbiomes in soils
under long-term extremely high CO_2 exposure. *Nature Microbiology*, *1*(2),
15025. http://doi.org/10.1038/nmicrobiol.2015.25

Bonkowski, M. (2004). Protozoa and plant growth: The microbial loop in soil
revisited. *New Phytologist*. http://doi.org/10.1111/j.1469-8137.2004.01066.x

Carlsen, M., Spohr, A. B., Nielsen, J., & Villadsen, J. (1996). Morphology and
physiology of an α-amylase producing strain of Aspergillus oryzae during batch
cultivations. *Biotechnology and Bioengineering*, *49*(3), 266–276.
http://doi.org/10.1002/(SICI)1097-0290(19960205)49:3<266::AID-
BIT4>3.0.CO;2-I

Choma, M., Bárta, J., Šantrůčková, H., & Urich, T. (2016). Low abundance of
Archaeorhizomycetes among fungi in soil metatranscriptomes. *Scientific*
Reports, *6*(1), 38455. http://doi.org/10.1038/srep38455

Clarholm, M. (1985). Interactions of bacteria, protozoa and plants leading to
mineralization of soil nitrogen. *Soil Biology and Biochemistry*, *17*(2), 181–187.
http://doi.org/10.1016/0038-0717(85)90113-0

Coleman, D. C. (1994). The microbial loop concept as used in terrestrial soil ecology
studies. *Microbial Ecology*, *28*(2), 245–250. http://doi.org/10.1007/BF00166814

Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST.
Bioinformatics, *26*(19), 2460–2461. http://doi.org/10.1093/bioinformatics/btq461

Epelde, L., Lanzén, A., Blanco, F., Urich, T., & Garbisu, C. (2015). Adaptation of soil
microbial community structure and function to chronic metal contamination at an
abandoned Pb-Zn mine. *FEMS Microbiology Ecology*, *91*(1), 1–11.
http://doi.org/10.1093/femsec/fiu007

Geisen, S., Koller, R., Hünninghaus, M., Dumack, K., Urich, T., & Bonkowski, M.
(2016). The soil food web revisited: Diverse and widespread mycophagous soil
protists. *Soil Biology and Biochemistry*, *94*, 10–18.
http://doi.org/10.1016/j.soilbio.2015.11.010

Geisen, S., Rosengarten, J., Koller, R., Mulder, C., Urich, T., & Bonkowski, M.
(2015). Pack hunting by a common soil amoeba on nematodes. *Environmental*
Microbiology, *17*(11), 4538–4546. http://doi.org/10.1111/1462-2920.12949

Geisen, S., Tveit, A. T., Clark, I. M., Richter, A., Svenning, M. M., Bonkowski, M., &
Urich, T. (2015). Metatranscriptomic census of active protists in soils. *ISME*
Journal, *9*(10), 2178–2190. http://doi.org/10.1038/ismej.2015.30

Giner, C. R., Forn, I., Romac, S., Logares, R., de Vargas, C., & Massana, R. (2016).
Environmental sequencing provides reasonable estimates of the relative
abundance of specific picoeukaryotes. *Applied and Environmental Microbiology*,
(May), AEM.00560-16. http://doi.org/10.1128/AEM.00560-16

Henriksen, C. M., Christensen, L. H., Nielsen, J., & Villadsen, J. (1996). Growth
energetics and metabolic fluxes in continuous cultures of Penicillium
chrysogenum. *Journal of Biotechnology*, *45*(2), 149–64. Retrieved from
http://www.ncbi.nlm.nih.gov/pubmed/9147448

Huson, D. H., Mitra, S., Ruscheweyh, H.-J., Weber, N., & Schuster, S. C. (2011).
Integrative analysis of environmental sequences using MEGAN4. *Genome*
Research, *21*(9), 1552–60. http://doi.org/10.1101/gr.120618.111

Karst, S. M., Dueholm, M. S., McIlroy, S. J., Kirkegaard, R. H., Nielsen, P. H., &
Albertsen, M. (2018). Retrieval of a million high-quality, full-length microbial
16S and 18S rRNA gene sequences without primer bias. *Nature Biotechnology*,
36(2), 190–195. http://doi.org/10.1038/nbt.4045

Keane, R., & Berleman, J. (2016). The predatory life cycle of Myxococcus xanthus.
Microbiology (United Kingdom). http://doi.org/10.1099/mic.0.000208

Koller, R., Rodriguez, A., Robin, C., Scheu, S., & Bonkowski, M. (2013). Protozoa
enhance foraging efficiency of arbuscular mycorrhizal fungi for mineral nitrogen
from organic matter in soil to the benefit of host plants. *New Phytologist*, *199*(1),
203–211. http://doi.org/10.1111/nph.12249

Kopylova, E., Noé, L., & Touzet, H. (2012). SortMeRNA: fast and accurate filtering
of ribosomal RNAs in metatranscriptomic data. *Bioinformatics (Oxford,*
England), *28*(24), 3211–7. http://doi.org/10.1093/bioinformatics/bts611

Kramer, S., Dibbern, D., Moll, J., Huenninghaus, M., Koller, R., Krueger, D., …
Kandeler, E. (2016). Resource partitioning between bacteria, fungi, and protists
in the detritusphere of an agricultural soil. *Frontiers in Microbiology*, *7*(SEP), 1–
12. http://doi.org/10.3389/fmicb.2016.01524

Lanzén, A., Jørgensen, S. L., Huson, D. H., Gorfer, M., Grindhaug, S. H., Jonassen, I., … Urich, T. (2012). CREST--classification resources for environmental sequence tags. *PloS One*, *7*(11), e49334. http://doi.org/10.1371/journal.pone.0049334

Lueders, T., Kindler, R., Miltner, A., Friedrich, M. W., & Kaestner, M. (2006). Identification of bacterial micropredators distinctively active in a soil microbial food web. *Applied and Environmental Microbiology*, *72*(8), 5342–5348. http://doi.org/10.1128/AEM.00400-06

McCarthy, D. J., Chen, Y., & Smyth, G. K. (2012). Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation. *Nucleic Acids Research*, *40*(10), 4288–4297. http://doi.org/10.1093/nar/gks042

Parada, G., & Acevedo, F. (1983). On the relation of temperature and RNA content to the specific growth rate inSaccharomyces cerevisiae. *Biotechnology and Bioengineering*, *25*(11), 2785–2788. http://doi.org/10.1002/bit.260251120

Reichenbach, H. (1999, February). The ecology of the myxobacteria. *Environmental Microbiology*. Wiley/Blackwell (10.1111). http://doi.org/10.1046/j.1462-2920.1999.00016.x

Schmieder, R., & Edwards, R. (2011). Quality control and preprocessing of metagenomic datasets. *Bioinformatics*, *27*(6), 863–864. http://doi.org/10.1093/bioinformatics/btr026

Schwab, C., Berry, D., Rauch, I., Rennisch, I., Ramesmayer, J., Hainzl, E., … Urich, T. (2014). Longitudinal study of murine microbiota activity and interactions with the host during acute inflammation and recovery. *ISME Journal*, *8*(5), 1101–1114. http://doi.org/10.1038/ismej.2013.223

Tempest, D. W., Dicks, J. W., & Ellwood, D. C. (1968). Influence of growth condition on the concentration of potassium in Bacillus subtilis var. niger and its possible relationship to cellular ribonucleic acid, teichoic acid and teichuronic acid. *The Biochemical Journal*, *106*(1), 237–43. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/4976492

Trap, J., Bonkowski, M., Plassard, C., Villenave, C., & Blanchart, E. (2016). Ecological importance of soil bacterivores for ecosystem functions. *Plant and Soil*, *398*(1–2), 1–24. http://doi.org/10.1007/s11104-015-2671-6

Tveit, A., Schwacke, R., Svenning, M. M., & Urich, T. (2013). Organic carbon transformations in high-Arctic peat soils: key functions and microorganisms. *The ISME Journal*, *7*(2), 299–311. http://doi.org/10.1038/ismej.2012.99

Tveit, A. T., Urich, T., Frenzel, P., & Svenning, M. M. (2015). Metabolic and trophic interactions modulate methane production by Arctic peat microbiota in response

to warming. *Proceedings of the National Academy of Sciences*, *112*(19), E2507–
E2516. http://doi.org/10.1073/pnas.1420797112

Urich, T., Lanzén, A., Qi, J., Huson, D. H., Schleper, C., & Schuster, S. C. (2008).
Simultaneous assessment of soil microbial community structure and function
through analysis of the meta-transcriptome. *PloS One*, *3*(6), e2527.
http://doi.org/10.1371/journal.pone.0002527

Urich, T., & Schleper, C. (2011). The "Double-RNA" Approach to Simultaneously
Assess the Structure and Function of a Soil Microbial Community BT - … and
Complementary Approaches. In … *and Complementary Approaches* (pp. 587–
596). John Wiley & Sons, Inc. Retrieved from
http://doi.wiley.com/10.1002/9781118010518.ch64%5Cnpapers2://publication/d
oi/10.1002/9781118010518.ch64

Wanek, W., Mooshammer, M., Blöchl, A., Hanreich, A., & Richter, A. (2010).
Determination of gross rates of amino acid production and immobilization in
decomposing leaf litter by a novel15N isotope pool dilution technique. *Soil*
Biology and Biochemistry, *42*(8), 1293–1302.
http://doi.org/10.1016/j.soilbio.2010.04.001

Zhang, L., & Lueders, T. (2017). Micropredator niche differentiation between bulk
soil and rhizosphere of an agricultural soil depends on bacterial prey. *FEMS*
Microbiology Ecology, *93*(9). http://doi.org/10.1093/femsec/fix103

Zhou, X. W., Li, S. G., Li, W., Jiang, D. M., Han, K., Wu, Z. H., & Li, Y. Z. (2014).
Myxobacterial community is a predominant and highly diverse bacterial group in soil
niches. *Environmental Microbiology Reports*, *6*(1), 45–56.
http://doi.org/10.1111/1758-2229.12107

Table 1. Context data for relevant sampling sites (changed from Choma *et al.*, 2016)

Site	Peatland soil "Knudsenheia"	Peatland soil "Solvatn"	Mofette	Mofette reference	Rothamsted grassland	Rotböhl	Forest Litter	Forest Soil	Mine L	Mine M	Mine H
Abbreviation	PsK	PsS	MO	MR	RS	RB	FL	FS	MiL	MiM	MiH
Location	Ny-Ålesund, Norway (Svalbard)	Ny-Ålesund, Norway (Svalbard)	Hartoušov, Czech Republic	Hartoušov, Czech Republic	Rothamsted, United Kingdom	Darmstadt, Germany	Vienna woods, Austria	Vienna woods, Austria	Coto Txomin, Spain	Coto Txomin, Spain	Coto Txomin, Spain
Climatic zone	Arctic	Arctic	Temperate	Temperate	Temperate	Temperate	Temperate	Temperate	Temperate	Temperate	Temperate
Biome	Fen wet land	Fen wet land	Floodplain	Floodplain	Grassland	Grassland	Temperate deciduous forest	Temperate deciduous forest	Shrubland	Shrubland	Shrubland
Dominant vegetation	Mosses	Mosses	*Filipendula ulmaria*	*Deschampsia cespitosa, Eriophorum vaginatum*	N.A.	N.A.	*Fagus sylvatica*	*Fagus sylvatica*	*Ulex europaeus*	*Festuca rubra*	*Festuca rubra*
Substrate type / Horizon	Organic peat (Top layer)	Organic peat (Top layer)	Organic soil	Gleic fluvisol	Mineral soil	Mineral soil	Litter horizon	Mineral soil (A horizon)	Mineral soil	Mineral soil	Mineral soil
pH	7.3	7.6	4.7	5.3	4.9	7.1	N.A.	4.5-5.1	3.9	5.6	5.9
Moisture (% soil dry weight)	1010	900	N.A	N.A.	33	32	18	43-64	52	49	30
# of replicates	2	2	3	3	2	1	2	4	3	3	3
Sampling time	August 2009	August 2009	July 2013	July 2013	July 2009	January 2006	May 2008	May 2008	March 2011	March 2011	March 2011
Sequencing method	454 GS FLX Titanium	454 GS FLX Titanium	Illumina HiSeq 2500	Illumina HiSeq 2500	454 GS FLX Titanium	454 GS 20	454 GS FLX	454 GS FLX	Illumina HiSeq 2000	Illumina HiSeq 2000	Illumina HiSeq 2000

Figure 1. Screening of pro- and eukaryotic micropredators. (A) Fraction of major

identified micropredator SSU rRNA normalized to SSU rRNA of prey bacteria. (B)

Fraction of major identified micropredator SSU rRNA normalized to total

micropredator SSU rRNA. Error bars show standard deviation of replicates. For sites

see Table 1.

Figure 2. Screening of *Myxococcales* and protozoa taxa. (A) Fraction of identified

Myxococcales SSU rRNA normalized to overall *Myxococcales* SSU rRNA. (B)

Fraction of protozoa SSU rRNA normalized to total protozoa SSU rRNA. For sites

see Table 1.

Figure 3. Comparison of organic and mineral soils. Fraction of major identified micropredator SSU rRNA normalized to SSU rRNA of prey bacteria. Average in organic soils (excluding MO samples) on the left; average in mineral soils in the right. Area of boxes resembles abundance of SSU rRNA. Numbers show proportions [%] of prey bacterial SSU rRNA. *Lysobacter* data are not shown due to low abundances.

 Figure 4. Colonisation of beech litter. Fraction of major identified micropredator,

 fungi, and prey bacteria SSU rRNA. Area of boxes resembles abundance of SSU

 rRNA. *Lysobacter* data are not shown due to low abundances. For litter types see

 Wanek *et al.*, 2010.

541 **Figure 5.** Simplified soil microbial loop. Left: Traditional microbial loop with
542 separate roles of prokaryotic and eukaryotic organisms. Right: Microbial loop
543 containing a bacterial loop independent of eukaryotic organisms. Straight arrows:
544 links between trophic levels. Bent arrows: provision of nutrients.