1 **Natural variation in *C. elegans* arsenic toxicity is explained by differences in**
2 **branched chain amino acid metabolism**
3

4 Stefan Zdraljevic[1,2], Bennett W. Fox[3], Christine Strand[5], Oishika Panda[3,4], Francisco J. Tenjo[3,4],
5 Shannon C. Brady[1,2], Tim A. Crombie[2], John G. Doench[5], Frank C. Schroeder[3], and Erik C.
6 Andersen[1,2,6] *
7

1. Interdisciplinary Biological Sciences Program, Northwestern University, Evanston, IL 60208,
USA
2. Department of Molecular Biosciences, Northwestern University, Evanston, IL 60208, USA
3. Boyce Thompson Institute and Department of Chemistry and Chemical Biology, Cornell
University, Ithaca, NY 14853, USA
4. The Buck Institute for Research on Aging, Novato, CA 94945, USA (Current address)
5. Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA
6. Robert H. Lurie Comprehensive Cancer Center of Northwestern University, Chicago, IL
60611, USA
* Corresponding author
18

Erik C. Andersen
Assistant Professor of Molecular Biosciences
Northwestern University
Evanston, IL 60208, USA
Tel: (847) 467-4382
Fax: (847) 491-4461
Email: Erik.Andersen@Northwestern.edu
26

Stefan Zdraljevic, stefanzdraljevic2018@u.northwestern.edu, ORCID 0000-0003-2883-4616
Bennett William Fox, bwf7@cornell.edu, ORCID 0000-0002-9749-3491
Christine Strand, cstrand@broadinstitute.org, ORCID 0000-0001-5783-6667
Oishika Panda, OPanda@buckinstitute.org, ORCID 0000-0002-1217-8231
Francisco J. Tenjo, fjt38@cornell.edu, ORCID 0000-0003-4848-916X
Shannon C. Brady, shannonbrady2014@u.northwestern.edu, ORCID 0000-0002-3043-1544
Tim A. Crombie, tcrombie@northwestern.edu, ORCID 0000-0002-5645-4154
John G. Doench, jdoench@broadinstitute.org, ORCID 0000-0002-3707-9889
Frank C. Schroeder, fs31@cornell.edu, ORCID 0000-0002-4420-0237
Erik Andersen, erik.andersen@northwestern.edu, ORCID 0000-0003-0229-9651

Abstract

We find that variation in the *dbt-1* gene underlies natural differences in *Caenorhabditis elegans* responses to the toxin arsenic. This gene encodes the E2 subunit of the branched-chain α-keto acid dehydrogenase (BCKDH) complex, a core component of branched-chain amino acid (BCAA) metabolism. We causally linked a non-synonymous variant in the conserved lipoyl domain of DBT-1 to differential arsenic responses. Using targeted metabolomics and chemical supplementation, we demonstrate that differences in responses to arsenic are caused by variation in iso-branched chain fatty acids. Additionally, we show that levels of branched chain fatty acids in human cells are perturbed by arsenic treatment. This finding has broad implications for arsenic toxicity and for arsenic-focused chemotherapeutics across human populations. Our study implicates the BCKDH complex and BCAA metabolism in arsenic responses, demonstrating the power of *C. elegans* natural genetic diversity to identify novel mechanisms by which environmental toxins affect organismal physiology.

Introduction

An estimated 100 million people are currently at risk of chronic exposure to arsenic, a toxic metalloid that can be found in the environment [1]. The high prevalence of environmental arsenic and the severe toxicity associated with exposure has made it the number one priority for the United States Agency for Toxic Substances and Disease Registry (https://www.atsdr.cdc.gov/SPL/). Inorganic trivalent arsenic As(III) compounds, which include arsenic trioxide (As_2O_3), are the most toxic forms of environmental arsenic [2,3]. In humans, As(III) is detoxified by consecutive methylation events, forming dimethylarsenite (DMA) [4,5]. However, this methylation process also creates the highly toxic monomethylarsenite (MMA) intermediate, so ratios of DMA to MMA determine levels of arsenic toxicity. Both MMA and DMA are produced from As(III) via the arsenic methyltransferase (AS3MT) [6]. Interestingly, individuals from human subpopulations that inhabit high arsenic environments have higher DMA/MMA ratios than individuals from low-arsenic environments. The elevated DMA/MMA ratio in these individuals is associated with natural differences in the *AS3MT* gene [7–9], which shows signs of strong positive selection. These results suggest that a more active AS3MT enzyme in these human subpopulations makes more DMA and enables adaptation to elevated environmental arsenic levels [6]. Importantly, population-wide differences in responses to environmental arsenic cannot be explained solely by variation in *AS3MT*, indicating that other genes must impact arsenic toxicity.

Despite its toxicity, arsenic trioxide has been used as a therapeutic agent for hundreds of years. Most recently, it was introduced as a highly effective cancer chemotherapeutic for the treatment of acute promyelocytic leukemia (APL) [10–13]. Hematopoietic differentiation and apoptosis in APL patients is blocked at the level of promyelocytes by the Promyelocytic Leukemia/Retinoic Acid Receptor alpha fusion protein caused by a t(15;17) chromosomal translocation [14,15]. Arsenic trioxide has been shown to directly bind a cysteine-rich region of the RING-B box coiled-coil domain of PML-RARα, which causes the degradation of the oncogenic fusion protein [16,17]. The success of arsenic trioxide (Trisenox®) has spurred its use in over a hundred clinical trials in the past decade [18]. Despite these successes, individual differences in the responses to arsenic-based treatments, including patient-specific dosing regimens and side effects, limit the full therapeutic benefit of this compound [19]. Medical practitioners require knowledge of the molecular mechanisms for how arsenic causes toxicity to provide the best individual-based therapeutic benefits.

Studies of the free-living roundworm *Caenorhabditis elegans* have greatly facilitated our understanding of basic cellular processes [20–23], including a number of studies that show that the effects of arsenic are similar to what is observed in mammalian model systems and humans. These effects include mitochondrial toxicity [24,25], the generation of reactive oxygen species (ROS) [26], genotoxicity [27], genome-wide shifts in chromatin structure [28], reduced lifespan [26], and the induction of the heat-shock response [29]. However, these studies were all performed in the genetic background of the standard *C. elegans* laboratory strain (N2). To date, 152 *C. elegans* strains have been isolated from various locations around the world [30–32], which contain a largely unexplored pool of genetic diversity much of which could underlie adaptive responses to environmental perturbations [33].

95   We used two quantitative genetic mapping approaches to show that a major source of variation in *C. elegans* responses to arsenic trioxide is caused by natural variation in the *dbt-1* gene, which encodes an essential component of the highly conserved branched-chain α-keto acid dehydrogenase (BCKDH) complex [34]. The BCKDH complex is a core component of branched-chain amino acid (BCAA) catabolism, which has not been previously implicated in arsenic responses. Furthermore, we show that a single missense variant in DBT-1(S78C), located in the highly conserved lipoyl-binding domain, underlies phenotypic variation in response to arsenic. Using targeted and untargeted metabolomics and chemical rescue experiments, we show that differences in wild isolate responses to arsenic trioxide are caused by differential synthesis of mono-methyl branched chain fatty acids (mmBCFA), metabolites with a central role in development [20]. These results demonstrate the power of using the natural genetic diversity across the *C. elegans* species to identify mechanisms by which environmental toxins affect physiology.

Results

Natural variation on chromosome II underlies differences in arsenic responses

We quantified arsenic trioxide sensitivity in *C. elegans* using a high-throughput fitness assay that utilizes the COPAS BIOSORT [35,36]. In this assay, three L4 larvae from each strain were sorted into arsenic trioxide or control conditions. After four days of growth, we quantified various attributes of populations that relate to the ability of *C. elegans* to grow in the presence of arsenic trioxide or control conditions (see Materials and Methods). To determine an appropriate concentration of arsenic trioxide for mapping experiments, we performed dose-response experiments on four genetically diverged isolates of *C. elegans*: N2, CB4856, JU775, and DL238 (Figure 1-source data 1). We focused on brood size and progeny length traits because they are the most direct measures of animal responses to environmental perturbations. We defined brood size as the total number of objects detected by the BIOSORT and progeny length by the median of populations of specific *C. elegans* strains. When compared to control conditions, all four strains produced fewer progeny at all arsenic trioxide concentrations, and the lowest concentration at which we observed a significant reduction in brood size for all strains was 1 mM (Figure 1-figure supplement 1A). We estimated broad-sense heritability (H^2) for the brood size trait in 1 mM arsenic trioxide to be (0.65) (Figure 1-source data 2) and the strain effect to be 0.48 (partial omega squared, $\omega_p{}^2$, Figure 1-source data 3), indicating that this trait has a large genetic component and a large strain effect. In addition to reduced brood sizes, we observed that the progeny of animals exposed to arsenic trioxide were shorter in length than the progeny of animals grown in control conditions (Figure 1-figure supplement 1B), which indicates an arsenic-induced developmental delay (H^2 = 0.15 - 0.49; Figure 1-source data 2 and $\omega_p{}^2$ = 0.02 - 0.60; Figure 1-source data 3, depending on the animal length summary statistic used). We found that CB4856 animals produced approximately 16% more offspring that were on average 20% larger than the other three strains when treated with 1 mM arsenic trioxide, suggesting that the CB4856 strain was more resistant to arsenic trioxide than the other three strains. To reduce the number of traits for subsequent analyses, we performed principal component analysis on all measured traits, which transforms the data to orthogonal axes that capture the most phenotypic variance (Figure 1-source data 1 and 4-6; See Materials and Methods). For 1 mM arsenic trioxide, we estimated the

broad-sense heritability (H^2) of the first principal component to be 0.12 (Figure 1-source data 2)
with an effect size of 0.16 ($\omega_p{}^2$) (Figure 1-source data 3). It is not surprising that the first principal
component explained a large percentage (70%) of the total phenotypic variance because many
of the summary statistic traits are highly correlated (Figure 1-figure supplement 2A-D; Figure 1-
source data 7). We noted that the first principal component (PC1) was most strongly influenced
by traits related to animal length, as indicated by the loadings (Figure 1-figure supplement 2A-D;
Figure 1-source data 3 and 6), suggesting that PC1 is a biologically relevant trait (Figure 1-figure
supplement 1C). Furthermore, because we observe a large range of effect sizes and broad-sense
heritability estimates across measured traits (Figure 1-figure supplement 3), we focused our
analyses on the PC1 trait for subsequent experiments.
The increased arsenic trioxide resistance of CB4856 compared to N2 motivated us to
perform linkage mapping experiments with a panel of recombinant inbred advanced intercross
lines (RIAILs) that were previously constructed through ten generations of intercrossing between
an N2 derivative (QX1430) and CB4856 [35]. To capture arsenic trioxide-induced phenotypic
differences, we exposed a panel of 252 RIAILs to 1 mM arsenic trioxide and corrected for growth
differences among RIAILs in control conditions and assay-to-assay variability using linear
regression (Figure 1-source data 8; see Materials and Methods). We performed linkage mapping
on processed traits and the eigenvector-transformed traits (principal components or PCs)
obtained from PCA that explained 90% of the variance in the processed trait set (Materials and
Methods). The rationale of this approach was to minimize trait fluctuations that could be caused
by only measuring the phenotypes of one replicate per RIAIL strain. In agreement with our
observations from the dose-response experiment, we found that PC1 captures 70.9% of the total
measured trait variance and is strongly influenced by correlated animal length traits (Figure 1-
figure supplement 4; Figure 1-source data 9-10). Linkage mapping analysis of the PC1 trait
revealed that arsenic trioxide-induced phenotypic variation is significantly associated with genetic
variation on the center of chromosome II (Figure 1A-figure supplement 5; Figure 1-source data
11). An additional four quantitative trait loci (QTL) were significantly associated with variation in
arsenic responses on chromosomes I, V, and X (Figure 1A-figure supplement 5). Consistent with
the loadings of PC1, we determined that PC1 is highly correlated with the both brood size and
animal length traits (Figure 1B), suggesting that PC1 captures RIAIL variation in these traits. To
further support this relationship to interpretable biological significance, we found that the animal
length and brood size traits map to the same region on the center of chromosome II (Figure 1-
figure supplement 6; Figure 1-source data 11). The QTL on the center of chromosome II explains
32.97% of the total RIAIL phenotypic variation for the PC1 trait, which accounts for 61.7% of the
total phenotypic variation that can be explained by genetic factors (H^2= 0.53) (Figure 1-figure
supplement 7; Figure 1-source data 12). Taken together, the five QTL identified by mapping the
PC1 trait account for 48.4% of the total RIAIL variation, corresponding to 91.3% of the total
phenotypic variation that can be explained by genetic factors. However, we did not account for
errors in genomic heritability estimates. In addition to the PC1, brood size, and animal length
traits, 69 additional measured traits map to the same genomic region (Figure 1-figure supplement
8; Figure 1-source data 11). This result was expected given correlation structure of the measured
traits (Figure 1-figure supplement 4). The PC1 QTL confidence interval spans from 7.04 to 8.87
Mb on chromosome II. This QTL is completely encompassed by the brood size (6.06-9.30 Mb)
and animal length (6.82-8.93 Mb) QTL confidence intervals (Figure 1-figure supplement 6).

However, each of these QTL confidence intervals span genomic regions greater than 1.5
megabases and contain hundreds of genes that vary between the N2 and CB4856 strains.
Therefore, we constructed near-isogenic lines (NILs) to isolate and narrow each QTL in a
controlled genetic background.

**Figure 1: A large-effect QTL on the center of chromosome II explains differences in arsenic
trioxide response between N2 and CB4856**

(A) Linkage mapping plots for the first principal component trait in the presence of 1000 µM arsenic trioxide
is shown. The significance values (logarithm of odds, LOD, ratio) for 1454 markers between the N2 and
CB4856 strains are on the y-axis, and the genomic position (Mb) separated by chromosome is plotted on
the x-axis. The associated 1.5 LOD-drop confidence intervals are represented by blue boxes. (B) The
correlation between brood size (blue; r^2 = 0.23, p-value = 2.4E-15) or animal length (pink; r^2 = 0.72, p-value
= 7E-70) with the first principal component trait. Each dot represents an individual RIAIL's phenotype, with
the animal length and brood size phenotype values on the x-axis and the first principal component
phenotype on the y-axis. (C) Tukey box plots of near-isogenic line (NIL) phenotype values for the first
principal component trait in the presence of 1000 µM arsenic trioxide is shown. NIL genotypes are indicated
below the plot as genomic ranges. The N2 trait is significantly different than the CB4856 and NIL traits
(Tukey HSD p-value < 1E-5).

202    To split the large QTL confidence intervals in half, we introgressed genomic regions from
the CB4856 strain on the left and right halves of the confidence intervals into the N2 genetic
background. In the presence of arsenic trioxide, both of these NILs recapitulated the parental
CB4856 PC1 phenotype (Figure 1C; Figure 1-source data 13), and had similar brood sizes and
progeny of similar length as the CB4856 parental strain (Figure 1-figure supplement 9; Figure 1-
source data 13). Furthermore, we show that similar to the RIAIL phenotypes, many of the
measured traits were highly correlated (Figure 1-figure supplement 10A; Figure 1-source data 14)
and contributed similarly to the PC1 trait (Figure 1-figure supplement 10B; Figure 1-source data
15). Importantly, the PC1 trait was highly correlated with the brood size and animal length
phenotypes (Figure 1-figure supplement 11). The phenotypic similarity of these NILs to the

CB4856 parental strain suggested that the two NILs might share an introgressed region of the CB4856 genome. To identify this shared introgressed region, we performed low-coverage whole-genome sequencing of the NIL strains and defined the left and right bounds of the CB4856 genomic introgression to be from 5.75 to 8.02 Mb and 7.83 to 9.66 Mb in the left and right NILs, respectively (Figure 1-source data 16). The left and right NILs recapitulate 88.6% and 91.6% of the effect size difference between N2 and CB4856 as measured by Cohen's F, respectively [37], which exceeds our observations the linkage mapping results where the QTL on chromosome II explained 61.7% of the total phenotypic variation in the RIAIL population. This discrepancy was observed likely because the NILs are a more homogenous genetic background and the experiment was performed at higher replication than the linkage mapping. Taken together, these results suggested that genetic differences between N2 and CB4856 within 7.83 to 8.02 Mb on chromosome II conferred resistance to arsenic trioxide.

In parallel to the linkage-mapping approach described above, we performed a genome-wide association (GWA) mapping experiment by quantifying the responses to arsenic trioxide for 86 wild *C. elegans* strains (Figure 2-source data 1) [30]. Consistent with previous experiments, the measured traits that most strongly influence PC1 are representative of animal length, as indicated by the loadings (Figure 2-figure supplement 1; Figure 2-source data 2-3). In agreement with the results from the linkage mapping approach, PC1 differences among the wild isolates mapped to a QTL on the center of chromosome II that spans from 7.71 Mb to 8.18 Mb (Figure 2A; Figure 2-source data 4-5). We noted that the brood size trait did not map to a significant QTL with the GWA mapping approach. To address this discrepancy, we estimated genomic broad- and narrow-sense heritability (H^2; h^2) for all of the wild isolate measured and principal component traits (Figure 2-figure supplement 2; Figure 2-source data 6). Broad-sense heritability for the brood size trait was approximately 0.05 for the wild isolates tested, which explains why we were unable to detect a significant QTL associated with this trait. By contrast, the PC1 trait H^2 was 0.16 when estimated using an additive-only model and 0.25 (h^2 = 0.11) when we incorporated the possibility for epistatisis (See Materials and Methods; Figure 2-source data 6). The marker found to be most correlated with the PC1 trait from GWA mapping (II:7,931,252), explains 84.6% (epistasis model) of the total heritable phenotypic variation. Of the 64 measured and principal component traits we mapped, 26 were significantly correlated with genetic variation on the center of chromosome II (Figure 2-figure supplement 3; Figure 2-source data 7). Notably, the CB4856 strain, which was one of the parents used to construct the RIAIL panel used for linkage mapping, had the non-reference genotype at the marker most correlated with PC1 (Figure 2B), suggesting that the same genetic variant(s) might be contributing to differential arsenic trioxide response among the RIAIL and wild isolate populations.

To fine map the PC1 QTL, we focused on variants from the *C. elegans* whole-genome variation dataset [31] that are shared among at least five percent of the 86 wild isolates exposed to arsenic trioxide. Under the assumption that the linkage and GWA mapping QTL are caused by the same genetic variation, we only considered variants present in the CB4856 strain. Eight markers within the QTL region are in complete linkage disequilibrium with each other and are most correlated with the first principal component trait (Figure 2C-figure supplement 4; Figure 2-source data 8). Only one of these markers is located within an annotated gene (*dbt-1*) and is predicted to encode a cysteine-to-serine variant at position 78 (C78S). Although it is possible that the causal variant underlying differential arsenic trioxide response in the *C. elegans* population is

256 an intergenic variant, we focused on the DBT-1(C78S) variant as a candidate to test for an effect
257 on arsenic response.

Figure 2: Variation in *C. elegans* wild isolates responses to arsenic trioxide maps to the center of chromosome II

Figure 2: Variation in *C. elegans* wild isolates responses to arsenic trioxide maps to the
center of chromosome II
(A) A manhattan plot for the first principal component in the presence of 1000 μM arsenic trioxide is shown.
Each dot represents an SNV that is present in at least 5% of the assayed wild population. The genomic
position in Mb, separated by chromosome, is plotted on the x-axis and the $-log_{10}(p)$ for each SNV is plotted
on the y-axis. SNVs are colored red if they pass the genome-wide Bonferroni-corrected significance
threshold, which is denoted by the gray horizontal line. SNVs are colored pink if they pass the genome-
wide eigen-decomposition significance threshold, which is denoted by the dotted gray horizontal line. The
genomic region of interest surrounding the QTL is represented by a cyan rectangle. (B) Tukey box plots of
phenotypes used for association mapping in (A) are shown. Each dot corresponds to the phenotype of an
individual strain, which is plotted on the y-axis. Strains are grouped by their genotype at the peak QTL
position (red SNV from panel A, ChrII:7,931,252), where REF corresponds to the allele from the reference
N2 strain. The N2 (orange) and CB4856 (blue) strains are highlighted. (C) Fine mapping of the chromosome
II region of interest (cyan region from panel A, 7.71 - 8.18 Mb) is shown. Each dot represents an SNV
present in the CB4856 strain. The association between the SNV and first principal component is shown on
the y-axis and the genomic position of the SNV is shown on the x-axis. Dots are colored by their SnpEff
predicted effect.
275

A cysteine-to-serine variant in DBT-1 contributes to arsenic response variation
The *C. elegans dbt-1* gene encodes the E2 component of the branched-chain α-keto acid
dehydrogenase complex (BCKDH) [34]. The BCKDH complex is a core component of branched-
chain amino acid (BCAA) catabolism and catalyzes the irreversible oxidative decarboxylation of

amino acid-derived branched-chain α-ketoacids [38]. The BCKDH complex belongs to a family of
α-ketoacid dehydrogenases that include pyruvate dehydrogenase (PDH) and α-ketoglutarate
dehydrogenase (KGDH) [39]. All three of these large enzymatic complexes include a central E2
component that is lipoylated at one critical lysine residue (two residues in PDH). The function of
these enzymatic complexes depends on the lipoylation of these lysine residues [39,40]. In
C. elegans, the putative lipoylated lysine residue is located at amino acid position 71 of DBT-1,
which is in close proximity to the C78S residue that we found to be highly correlated with arsenic
trioxide resistance.
To confirm that the C78S variant in DBT-1 contributes to differential arsenic trioxide
responses, we used CRISPR/Cas9-mediated genome editing to generate allele-replacement
strains by changing the C78 residue in the N2 strain to a serine and the S78 residue in the CB4856
strain to a cysteine. When treated with arsenic trioxide, the N2 DBT-1(S78) allele-replacement
strain recapitulated 75.5% of the phenotypic difference between the CB4856 and N2 strains as
measured with the first principal component, as measured by Cohen's F [37] (Figure 3; Figure 1-
source data 13). Similarly, the CB4856 DBT-1(C78) allele-replacement strain recapitulated 63.9%
of the total phenotypic difference between the two parental CB4856. The degree to which the
allele-replacement strains recapitulated the difference in the first principal component between
the N2 and CB4856 strains matched our observations from the linkage mapping experiment,
where the chromosome II QTL explained 61.7% of the total phenotypic variation in the RIAIL
population. This result suggested that the majority of heritable variation in arsenic trioxide
response was explained by the DBT-1(C78S) allele. We obtained similar results for the progeny
length trait (Figure 3-figure supplement 1; Figure 1-source data 13), which is likely because of the
high level of correlation between the animal length and first principal component trait (Figure 1-
figure supplement 10; Figure 3-figure supplement 2; Figure 1-source data 10-11). However, when
considering brood size, the N2 DBT-1(C78S) allele-replacement strain produced an intermediate
number of progeny in the presence of arsenic trioxide relative to the parental N2 and CB4856
strains. And the CB4856 DBT-1(S78C) allele-replacement strain produced fewer offspring than
both parental strains (Figure 3-figure supplement 1; Figure 1-source data 13). These results
suggested that additional genetic variants between the N2 and CB4856 strains might interact with
the DBT-1(C78S) allele to affect different aspects of physiology. Nevertheless, these results
functionally validated that the DBT-1 C78S variant underlies differences in physiological
responses to arsenic trioxide.

Figure 3: The DBT-1(C78S) variant contributes to arsenic trioxide responses
Tukey box plots of the first principal component generated by PCA on allele-replacement strain phenotypes measured by the COPAS BIOSORT 1000 μM arsenic trioxide exposure are shown (N2, orange; CB4856, blue; allele replacement strains, gray). Labels correspond to the genetic background and the corresponding residue at position 78 of DBT-1 (C for cysteine, S for serine). All pair-wise comparisons are significantly different (Tukey HSD, *p*-value < 1.42E-5).

Arsenic trioxide inhibits the DBT-1 C78 allele
Mono-methyl branched chain fatty acids (mmBCFA) are an important class of molecules that are produced via BCAA catabolism [20,34,41,42]. The production of mmBCFA requires the BCKDH, fatty acid synthase (FASN-1), acetyl-CoA carboxylase (POD-2), fatty acyl elongases (ELO-5/6), β-ketoacyl dehydratase (LET-767), and acyl CoA synthetase (ACS-1) [20,34,41,43–45]. Strains that lack functional *elo-5*, *elo-6*, or *dbt-1* produce less C15ISO and C17ISO mmBCFAs, arrest at the L1 larval stage, and can be rescued by supplementing the growth media with C15ISO or C17ISO [20,34,41] (Figure 4A).

      Because DBT-1 is involved in BCAA catabolism, we hypothesized that the DBT-1(C78S)-dependent difference in progeny length between the N2 and CB4856 strains after arsenic trioxide treatment might be caused by differential larval (L1) arrest through depletion of downstream mmBCFAs. To test this hypothesis, we quantified the abundance of the monomethyl-branched (ISO) and straight-chain (SC) forms of C15 and C17 in the N2, CB4856, and allele-swap genetic backgrounds. We measured the metabolite levels in staged L1 animals and normalized the detected amounts of C15ISO and C17ISO relative to the abundances of C15SC and C17SC, respectively, to mitigate confounding effects of differences in developmental rates that could result from genetic background differences or arsenic trioxide exposure. Generally, the ratios of C15ISO/C15SC and C17ISO/C17SC were reduced in

Figure 4: Differential production of mmBCFA underlies DBT-1(C78)-mediated sensitivity to arsenic trioxide.

(A) A simplified model of BCAA catabolism in *C. elegans*. The BCKDH complex, which consists of DBT-1, catalyzes the irreversible oxidative decarboxylation of branched-chain ketoacids. The products of these breakdown can then serve as building blocks for the mmBCFA that are required for developmental progression. (B) The difference in the C15ISO/C15SC (left panel) or C17ISO/C17SC (right panel) ratios between 100 µM arsenic trioxide and control conditions is plotted on the y-axis for three independent replicates of the CB4856 and CB4856 allele swap strains and six independent replicates of the N2 and N2 allele swap strains. The difference between the C15 ratio for the CB4856-CB4856 allele swap comparison is significant (Tukey HSD *p*-value = 0.0427733), but the difference between the C17 ratios for these two strains is not (Tukey HSD *p*-value = 0.164721). The difference between the C15 and C17 ratios for the N2-N2 allele swap comparisons are both significant (C15: Tukey HSD *p*-value = 0.0358; C17: Tukey HSD *p*-value = 0.003747). (C) Tukey box plots median animal length after arsenic trioxide or arsenic trioxide and 0.64 µM C15ISO exposure are shown (N2, orange; CB4856, blue; allele replacement strains, gray). Labels correspond to the genetic background and the corresponding residue at position 78 of DBT-1 (C for cysteine, S for serine). Every pair-wise strain comparison is significant except for the N2 DBT-1(S78) - CB4856 comparisons (Tukey's HSD *p*-value < 1.43E-6)*.*

arsenic-treated animals relative to controls (Figure 4B; Figure 4-source data 1-3). However, arsenic trioxide treatment had a 7.6-fold stronger effect on the C15ISO/C15SC ratio in N2, which naturally has the C78 allele, than on the N2 DBT-1(S78) allele swap strain. This difference suggests that the DBT-1(C78) allele is more strongly inhibited by arsenic trioxide (0.04 to 0.004, Tukey HSD p-value = 0.0358, n = 6). Similarly, we observed a 6.6-fold arsenic-induced reduction in the C17ISO/C17SC ratio when comparing the N2 DBT-1(C78) and N2 DBT-1(S78) strains (Tukey HSD p-value = 0.003747, n = 6). When comparing the CB4856 DBT-1(S78) and CB4856 DBT-1(C78) strains, we observed a 2.8-fold lower C15ISO/C15SC ratio (Tukey HSD p-value = 0.0427733, n = 3) and 1.5-fold lower C17ISO/C17SC ratio (Tukey HSD p-value = 0.164721, n = 3) in the in the CB4856 DBT-1(C78) strain. We noted that the C17ISO/straight-chain ratio difference was not significantly different between the two CB4856 genetic background strains. However, we observed a significant arsenic-induced decrease in raw C17ISO production in the CB4856 DBT-1(C78) strain (Tukey HSD p-value = 0.029) and no significant difference in the CB4856 DBT-1(S78) strain (Tukey HSD p-value = 0.1) (Figure 4-figure supplement 1). Importantly, these DBT-1(C78S) allele-specific reductions in ISO/straight-chain ratios were not driven by arsenic-induced differences in straight-chain fatty acids (Figure 4-figure supplement 2). These results explained the majority of the physiological differences between the N2 and CB4856 strains in the presence of arsenic trioxide (Figure 3) and suggested that the DBT-1(C78) allele was inhibited by arsenic trioxide more strongly than DBT-1(S78). Taken together, the differential reduction in branched-chain fatty acids likely underlies the majority of physiological differences between the sensitive and resistant *C. elegans* strains.

In addition to arsenic-induced differences in branched chain fatty acid production, we observed significant differences in branched/straight-chain ratios between the parental and allele swap strains when L1 larval animals were grown in control conditions (Figure 4-figure supplement 3; Figure 4-source data 1-3). Strains with the DBT-1(C78) had higher ISO/SC ratios relative to strains with the DBT-1(S78) for the C17 (CB4856 DBT-1(C78): Tukey HSD p-value = 0.0342525, n = 3; N2 DBT-1(C78): Tukey HSD p-value = 0.0342525, n = 6) and C15 ratios (CB4856: Tukey HSD p-value = 0.0168749, n = 3; N2: Tukey HSD p-value = 0.1239674, n = 6) (Figure 4-figure supplement 3; Figure 4-source data 1-3). We noted that the C15ISO/straight-chain ratio was not significantly different when comparing the N2 and N2 allele swap strain, but the direction of effect matched our other observations and we saw significant differences in C15ISO levels (N2-C15ISO DBT-1(C78): Tukey HSD p-value = 0.0265059, n = 6, Figure 4-figure supplement 4; Figure 4-source data 1-3). Importantly, the DBT-1 allele-specific differences in the fatty acid ratio and ISO measurements were not driven by differences in straight-chain fatty acids (Figure 4-figure supplement 4). However, we did not observe the same effect of the DBT-1(C78S) allele at the young adult life stage (Figure 4-figure supplement 5; Figure 4-source data 4). Taken together, these results suggest that the DBT-1(C78) allele produces more branched chain fatty acids than the DBT-1(S78) allele, but this effect was dependent on the developmental stage of the animals.

To test the hypothesis that differential arsenic-induced depletion of branched-chain fatty acids in strains with the DBT-1(C78S) causes physiological differences in growth, we tested if mmBCFA supplementation could rescue the effects of arsenic trioxide-induced toxicity. We exposed the parental and the DBT-1 allele-replacement strains to media containing arsenic trioxide alone, C15ISO alone, or a combination of arsenic trioxide and C15ISO. In agreement with previous experiments, the PC1 trait was highly correlated with animal length traits (Supplementary

Figure 4-figure supplement 6-7; Figure 4-source data 5-7). When grown in arsenic-trioxide alone
the average phenotypic difference between N2 DBT-1(C78) and N2 DBT-1(S78) allele was 3.39
units for the PC1 trait. C15ISO supplementation of the arsenic growth media caused a 56.3%
rescue of the allele-specific effect in the N2 genetic background (Figure 4C-figure supplement 8).
Similarly, when CB4856 DBT-1(C78) animals were supplemented with C15ISO, the allele-specific
PC1 phenotypic difference was reduced by 35.9% when compared to the difference between the
CB4856 DBT-1(C78) and CB4856 DBT-1(S78) strains in arsenic trioxide alone (Figure 4C-figure
supplement 8). By contrast, CB4856 DBT-1(S78) and N2 DBT-1(S78) phenotypes were
unaffected by C15ISO supplementation in arsenic trioxide media. Collectively, these data support
the hypothesis that the cysteine/serine variant in DBT-1 contributes to arsenic sensitivity in
C. elegans by reducing ISO fatty acid biosynthesis, and the DBT-1(C78) variant can be partially
rescued by supplementation with mmBCFAs.
420
Arsenic exposure increases mmBCFA production and favors a cysteine allele in human
DBT1
To test whether our results from *C. elegans* translate to human variation in arsenic sensitivity, we
tested the role of DBT1 variation on arsenic trioxide responses and mmBCFA biosynthesis in
human cells. The human DBT1 enzyme contains a serine at position 112 that corresponds to the
C78 residue in *C. elegans* DBT-1 (Figure 5A-figure supplement 1). Using CRISPR/Cas9, we
edited batch cultures of 293T cells to generate a subset of cells with DBT1(S112C). These cells
were exposed to arsenic trioxide or control conditions and we monitored changes in the fraction
of cells carrying the DBT1(C112) allele. We found that arsenic exposure caused a 4% increase
in the fraction of cells that contained the DBT1(C112) allele (Figure 5B, Fisher's exact test, *p*-
value < 1.9E-16; Figure 5-source data 1-2). Though the human DBT1 does not vary within the
human population at S112, two residues in close spatial proximity to S112 do vary among
individuals in the population (R113C and W84C) (Figure 5A) [46]. To test the effects of these
residues on arsenic trioxide sensitivity, we performed the same editing and arsenic selection
procedure described above. Over the course of the selection experiment, cells with DBT1(W84C)
and DBT1(R113C) increased by 2% and 1%, respectively (Figure 5B). Therefore, it appears that
all three missense variants caused a slight increase in fitness in batch-edited human cell cultures
exposed to arsenic. To determine if branched-chain fatty acid production was altered by arsenic
exposure, as we found in *C. elegans*, we measured mmBCFA production in unedited 293T cells
in arsenic and mock-treated cultures. We found that overall fatty acid production was markedly
reduced in arsenic-treated cultures. In contrast to our observations in *C. elegans*, straight-chain
fatty acids were more drastically reduced than ISO fatty acids (Figure 5-source data 3). These
results are consistent with our observations that arsenic does not have as strong of an effect on
C. elegans strains with the DBT-1 S78 allele. Additionally, the DBT1(C112) allele, similar to the
C. elegans DBT-1(C78) allele, was associated with higher production of mmBCFAs.

Figure 5: Protective effect of cysteine residues in human DBT1.
(A) Alignment of *C. elegans* DBT-1 and *H. sapiens* DBT1. The residues tested for an arsenic-specific effect are indicated with arrows - W84C (pink), S112C (blue), and R113C (black). The lysine that is post-translationally modified with a lipoid acid is highlighted in red. (B) The percent increase of edited human cells that contain the W84C, S112C, or R113C amino acid change in DBT1 in the presence 5 μM arsenic trioxide relative to control conditions are shown. The number of reads in 5 μM arsenic trioxide for all replicates are significantly different from control conditions (Fisher's exact test, *p*-value < 0.011).

Discussion

In this study, we characterized the effects of *C. elegans* natural genetic variation on physiological responses to the pervasive environmental toxin arsenic trioxide. Though the effects of this toxin have been extensively studied in a variety of systems [2,3,39,47,48], recent evidence from human population studies have revealed local adaptations within region-specific subpopulations [6,8,9,49]. Our investigation into the natural variation in *C. elegans* responses to arsenic trioxide led to the discovery of a novel mechanism by which this compound could elicit toxicity. We show that arsenic trioxide differentially inhibits two natural alleles of the E2 domain of the BCKDH complex, which is encoded by the *dbt-1* gene. Specifically, strains with the DBT-1(C78) allele are more sensitive to arsenic trioxide than strains carrying the DBT-1(S78). Furthermore, we show that the increased sensitivity of the DBT-1(C78) allele is largely explained by differences in the production of mmBCFAs (Figure 4B-C), which are critical molecules for developmental progression beyond the first larval stage. Arsenic is thought to inhibit the activity of both the pyruvate dehydrogenase (PDH) and the α-ketoglutarate (KGDH) dehydrogenase complexes through interactions with the reduced form of lipoate [39], which is the cofactor for the E2 domain of these complexes. Like the PDH and KGDH complexes, the E2 domain of BCKDH complex

requires the cofactor lipoate to perform its enzymatic function [50–52]. The inhibition of DBT-1 by arsenic trioxide could involve three-point coordination of arsenic by the C78 thiol and the reduced thiol groups of the nearby lipoate. However, based on the crystal structure (PDB:1Y8N), the atomic distance between the C78 thiol group and the thiol groups from the lipoylated lysine is ~32 Å, which might be too large a distance for coordinating arsenic (Figure 5A-figure supplement 1) [53]. Alternatively, arsenic trioxide could inhibit DBT-1(C78) through coordination between the thiol groups of C78 and C65 (~8.5 Å) (Figure 5A-figure supplement 1). Analogous thiol-dependent mechanisms have been proposed for the inhibition of other enzymes by arsenic [48]. Despite structural similarities and a shared cofactor, no evidence in the literature indicates that BCKDH is inhibited by arsenic trioxide, so these results demonstrate the first connection of arsenic toxicity to BCKDH E2 subunit inhibition.

Multiple sequence alignments show that cysteine residues C65 and C78 of *C. elegans* DBT-1 correspond to residues S112 and C99 of human DBT1 (Figure 5A). Though DBT1 does not vary at position 112 within the human population, two residues (R113C and W84C) in close spatial proximity do [46]. We hypothesized that cysteine variants in DBT1 would sensitize human cells to arsenic trioxide. However, we found that the cysteine variants (W84C, S112C, and R113C) proliferated slightly more rapidly than the parental cells in the presence of arsenic. Notably, a growing body of evidence suggests that certain cancer cells upregulate components involved in BCAA metabolism, and this upregulation promotes tumor growth [54,55]. Perhaps the increased proliferation of human cell lines that contain the DBT1 C112 allele (Figure 5) is caused by increased activity of the BCKDH complex. It is worth noting that human cell lines grown in culture do not have the same strict requirements for mmBCFA, and the requirements for different fatty acids are variable among diverse immortalized cell lines [56,57]. Furthermore, in *C. elegans*, the developmental defects associated with *dbt-1* loss-of-function can be rescued by neuronal-specific expression of *dbt-1* [34], suggesting that the physiological requirements of mmBCFA in *C. elegans* depend on the coordination of multiple tissues that cannot be recapitulated with cell-culture experiments. These results further highlight the complexity of arsenic toxicity, as well as the physiological requirements for BCAA within and across species and could explain the discrepancy between the physiological effects we observed in *C. elegans* and human cell-line experiments. Given that arsenic trioxide has become the standard-of-care for treating APL [58] and is gaining traction in treating other leukemias, it will be important to further explore the effects of arsenic on BCAA metabolism and cancer growth.

The C78 allele of DBT-1 is likely the derived allele in the *C. elegans* population because all other organisms have a serine at the corresponding position. The loss of the serine allele in the *C. elegans* population might have been caused by relaxed selection at this locus, though this hypothesis is difficult to test because of the effects of linked selection and decreased recombination in chromosome centers. It is hypothesized that the *C. elegans* species originated from the Pacific Rim and that the ancestral state more closely resembles the CB5846 strain than the N2 strain [30,59]. The CB4856 strain was isolated from the Hawaiian island of Oʻahu [60], where environments could have elevated levels of arsenic in the soil from volcanic activity, the farming of cane sugar, former construction material (canec) production facilities, or wood treatment plants (Hawaii.gov). It is possible that as the *C. elegans* species spread across the globe into areas with lower levels of arsenic in the soil and water, the selective pressure to maintain high arsenic tolerance was relaxed and the cysteine allele appeared. Alternatively,

higher mmBCFA levels at the L1 larval stage in strains with the DBT-1(C78) allele (Figure 4-figure
supplement 3-4) might cause faster development in certain conditions, although we did not
observe allele-specific growth differences in laboratory conditions. Despite these clues
suggesting selection in local environments, the genomic region surrounding the *dbt-1* locus does
not show a signature of selection as measured by Tajima's D [61] (Figure 2-figure supplement 5;
Figure 2-source data 9), and the strains with the DBT-1 S78 allele show no signs of geographic
clustering (Figure 2-figure supplement 6; Figure 2-source data 10). Our study suggests that
C. elegans is a powerful model to investigate the molecular mechanisms for how populations
respond to environmental toxins.
530
531 **Materials and methods**
532
533 **Key Resources Table:**

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
gene (*Caenorhabditis elegans*)	*dbt-1*	NA	Wormbase:WB Gene00014054	
strain, strain background (N2)	N2 DBT-1(S78)	This paper	Andersen_Lab: ECA581	N2::dbt-1(ean15[C78S])
strain, strain background (CB4856)	CB4856 DBT-1(C78)	This paper	Andersen_Lab: ECA590	CB4856::dbt-1(ean34[S78C])
strain, strain background (N2)	Left NIL; CB4856>N2 (II:5.75-8.02 Mb)	This paper	Andersen_Lab: ECA414	eanIR188[ChrII:5.75 - 8.02Mb]
strain, strain background (N2)	Right NIL; CB4856>N2 (II:7.83-9.66 Mb)	This paper	Andersen_Lab: ECA434	eanIR208[ChrII:7.83 - 9.66Mb]
sequence-based reagent	NIL Fd primer	This paper	Andersen_Lab: oECA609	tttcacacaaaccatgcgct
sequence-based reagent	NIL Rv primer	This paper	Andersen_Lab: oECA610	actcgtctgctgggtattct
sequence-based reagent	NIL Fd primer	This paper	Andersen_Lab: oECA611	tgtcttcgcacctttactcg
sequence-based reagent	NIL Rv primer	This paper	Andersen_Lab: oECA612	cattcaagtcagggcgtatcc
sequence-based reagent	Genotype C78S Edit	This paper	Andersen_Lab: oECA1163	GAAGGAATTGCCGAAGTTCAGGTTAAG
sequence-based reagent	Genotype C78S Edit	This paper	Andersen_Lab: oECA1165	CCGTCATCTCCACAAAAAGCTTTATCTCTC
sequence-based reagent	*dbt-1* gRNA	This paper	Andersen_Lab: crECA97	CCATCTCCTGTAGATACGAC
sequence-based reagent	N2 *dbt-1* repair oligo	This paper	Andersen_Lab: oECA1542	CTTCCAGGTACGTGAAAGAAGGAGATACGATTTCGCAGTTCGATAAAGTCT GTGAAGTGCAAAGTGATAAAGCAGCAGTAACCATCTCCAGTAGATACGACG GAATTGTCAAAAAATTGTAAGTTTCTTCCTAA
sequence-based reagent	CB4856 *dbt-1* repair oligo	This paper	Andersen_Lab: oECA1543	TTAGGAAGAAACTTACAATTTTTTGACAATTCCGTCGTATCTACAGGAGATG GTTACTGCTGCTTTATCGCTTTGCACTTCACAGACTTTATCGAACTGCGAAAT CGTATCTCCTTCTTTCACGTACCTGGAAG
sequence-based reagent	*dpy-10* repair oligo	10.1534/g enetics.11 4.166389	Andersen_Lab: crECA37	CACTTGAACTTCAATACGGCAAGATGAGAATGACTGGAAACCGTACCGCAT GCGGTGCCTATGGTAGCGGAGCTTCACATGGCTTCAGACCAACAGCCTAT
sequence-based reagent	*dpy-10* gRNA	10.1534/g enetics.11 4.166389	Andersen_Lab: crECA36	GCTACCATAGGCACCACGAG

sequence-based reagent	Human gRNA S112C and R113C	This paper	Guide_1 used in RDA_74	TCCATCATAACGACTAGTGA
sequence-based reagent	S112C repair template	This paper	1192 DBT1-repair-S112C	ATAGCATCTGTGAAGTTCAAAGTGATAAAGCTTCTGTTACAATCACTTGTCG TTATGATGGAGTCATTAAAAAACTCTATT
sequence-based reagent	R113C repair template	This paper	1193 DBT1-repair-R113C	ATAGCATCTGTGAAGTTCAAAGTGATAAAGCTTCTGTTACAATCACTAGTTG TTATGATGGAGTCATTAAAAAACTCTATT
sequence-based reagent	Fwd PCR C	This paper	1188 DBT1-PCR-C	TtgtggaaaggacgaaacaccgAGAAGGAGATACAGTGTCTCAGT
sequence-based reagent	Fwd PCR D	This paper	1189 DBT1-PCR-D	TtgtggaaaggacgaaacaccgTGTCTCAGTTTGATAGCATCTGTG
sequence-based reagent	Human gRNA W84C	This paper	Guide_2 used in RDA_75	TCTTTTAGGTATGTAAAAGA
sequence-based reagent	W84C repair template	This paper	1195 DBT1-repair-W84C-v2	GACTGTTTCCATAAAAGTGTCTCATTTCTTTTTCTTTTAGTTATGTGAAGGAA GGAGATACAGTGTCTCAGTTTGATAGCAT
sequence-based reagent	Fwd PCR A	This paper	1186 DBT1-PCR-A	TtgtggaaaggacgaaacaccgGCATGGCATTTACATCCTTAATATGAT
sequence-based reagent	Fwd PCR B	This paper	1187 DBT1-PCR-B	TtgtggaaaggacgaaacaccgCCTTAATATGATCTGTACTTATGACTGTTT
sequence-based reagent	Rev PCR 1	This paper	1190 DBT1-PCR-Rev1	TctactattctttcccctgcactgtCTACTAATGGCTTCCCCACAT
sequence-based reagent	Rev PCR 2	This paper	1191 DBT1-PCR-Rev2	TctactattctttcccctgcactgtCAATACCTTTTAAAGCTTCCGTTTCTAT
transfected construct (*Homo Sapiens*)	S112C and R113C Cas9-sgRNA plasmid	This paper	p1054	
transfected construct (*Homo Sapiens*)	W84C Cas9-sgRNA plasmid	This paper	p1052	

Strains

Animals were fed the bacterial strain OP50 and grown at 20°C on modified nematode growth
medium (NGM), containing 1% agar and 0.7% agarose to prevent burrowing of the wild isolates
[62]. For each assay, strains were grown for five generations with no strain entering starvation or
encountering dauer-inducing conditions [63]. Wild *C. elegans* isolates used for genome-wide
association and recombinant inbred advanced intercross lines (RIAILs) used for linkage mapping
have been described previously [31,32,35]. Strains constructed for this manuscript are listed
above in the Key Resources Table.

High-throughput fitness assay

We used the high-throughput fitness assay (HTA) described previously [35]. In short, strains are
passaged for four generations before bleach-synchronization and aliquoted to 96-well microtiter
plates at approximately one embryo per microliter in K medium [62]. The final concentration of
NaCl in the K medium for the genome-wide association (GWA) and linkage mapping assays was
51 mM. For all subsequent experiments the final NaCl concentration was 10.2 mM. The following
day, hatched and synchronized L1 animals were fed HB101 bacterial lysate (Pennsylvania State
University Shared Fermentation Facility, State College, PA, [64]) at a final concentration of 5
mg/ml and grown for two days at 20°C. Next, three L4 larvae were sorted using a large-particle
flow cytometer (COPAS BIOSORT, Union Biometrica, Holliston, MA) into microtiter plates that
contain HB101 lysate at 10 mg/ml, K medium, 31.25 µM kanamycin, and either arsenic trioxide
dissolved in 1% water or 1% water alone. The animals were then grown for four days at 20°C.
The resulting populations were treated with sodium azide (50 mM) prior to being measured with

the BIOSORT. All raw experimental data can be found on FigShare
().
559

Calculation of fitness traits for genetic mappings

Phenotype data generated using the BIOSORT were processed using the R package *easysorter*,
which was specifically developed for processing this type of data [65]. Briefly, the function
read_data, reads in raw phenotype data and runs a support vector machine to identify and
eliminate bubbles. Next, the *remove_contamination* function eliminates any wells that were
identified as contaminated prior to scoring population parameters. This analysis results in
processed BIOSORT data where each observation is for a given strain corresponds to the
measurements for an individual animal. However, the phenotypes we used for mapping and
follow-up experiments are summarized statistics of populations of animals in each well of a 96-
well plate. The *sumplate* function was used to generate summary statistics of the measured
parameters for each animal in each well. These summary statistics include the 10th, 25th, 50th,
75th, and 90th quantiles for time of flight (TOF), animal extinction (EXT), and three fluorescence
channels (Green, Yellow, and Red), which correspond to animal length, optical density, and ability
to pump fluorescent beads, respectively. Measured brood sizes (n) are normalized by the number
of animals that were originally sorted into each well (norm.n). For mapping experiments, a single
well replicate for each strain is summarized using the *sumplate* function. For follow-up
experiments, multiple replicates for each strain indicated by a unique plate, well, and column were
used. After summary statistics for each well are calculated, we accounted for differences between
assays using the *regress(assay=TRUE)* function in the *easysorter* package. Outliers in the GWA
and linkage mapping experiments were identified and eliminated using the *bamf_prune* function
in *easysorter*. For follow-up experiments that contained multiple replicates for each strain, we
eliminated strain replicates that were more than two standard deviations from the strain mean for
each condition tested. Finally, arsenic-specific effects were calculated using the
regress(assay=FALSE) function from *easysorter*, which accounts for strain-specific differences in
growth parameters present in control conditions.
585

Principal component analysis of processed BIOSORT measured traits

The COPAS BIOSORT measures individual animal length (TOF), optical density (EXT),
fluorescence (green, yellow, and red). We use these data to calculate the total number of animals
in a well and then normalize by the number of animals initially sorted into the well (brood size). All
of these measurements were then summarized using the *easysorter* package to generate various
summary statistics of each measured parameter, including five distribution quantiles and
measures of dispersion [35]. Because fluorescent beads were only used for linkage and GWA
mapping experiments, we eliminated all of these traits prior to PCA analysis for all non-mapping
experiments. Additionally, we removed all summary statistic traits related to data dispersion (IQR,
variance, and coefficient of variation) for all experiments. Prior to principal component analysis
(PCA), HTA phenotypes were scaled to have a mean of zero and a standard deviation of one
using the *scale* function in R. PCA was performed using the *prcomp* function in R [66].
Eigenvectors were subsequently extracted from the object returned by the *prcomp* function.
599

Arsenic trioxide dose-response assays

All dose-response experiments were performed on four genetically diverged strains (N2, CB4856,
DL238, and JU775) in technical quadruplicates prior to performing GWA and linkage mapping
experiments (Figure 1-source data 1). Animals were assayed using the HTA, and phenotypic
analyses were performed as described above. The arsenic trioxide concentration for GWA and
linkage mapping experiments was chosen based on an observable effect for animal length and
brood size phenotypes in the presence of arsenic.

Heritability calculations

For dose response experiments, broad-sense heritability (H^2) estimates were calculated using
the *lmer* function in the lme4 package with the following linear mixed model (phenotype ~1 +
(1|strain)) [67]. H^2 was then calculated as the fraction of the total variance that can be explained
by the random component (strain) of the mixed model. Prior to estimating H^2, we removed outlier
replicates that we defined as replicates with values greater than two standard deviations away
from the mean phenotype. Outliers were defined on a per-trait and per-strain basis. Heritability
estimates for dose response experiments are in Figure 1-source data 2.

Heritability estimates for the linkage mapping experiment were calculated using two
approaches. In both approaches, we used the previously described RIAIL genotype matrix to
compute relatedness matrices [35]. In the first approach, a variance component model using the
R package regress was used to estimate the fraction of phenotypic variation explained by additive
and epistatic genetic factors, H^2, or just additive genetic factors, h^2 [68,69], using the formula (y
~ 1, ~ZA+ZAA), where y is a vector of RIAIL phenotypes, ZA is the additive relatedness matrix,
and ZAA is the pairwise-interaction relatedness matrix. The additive relatedness matrix was
calculated as the correlation of marker genotypes between each pair of strains. In addition, a two-
component variance model was calculated with both an additive and pairwise-interaction effect.
The pairwise-interaction relatedness matrix was calculated as the Hadamard product of the
additive relatedness matrix.

The second approach utilized a linear mixed model and the realized additive and epistatic
relatedness matrices [70–73]. We used the *mmer* function in the sommer package with the
formula (y ~ A+E) to estimate variance components, where y is a vector of RIAIL phenotypes, A
is the realized additive relatedness matrix, and E is the epistatic relatedness matrix. This same
approach was used to estimate heritability for the GWA mapping phenotype data, with the only
difference being that we used the wild isolate genotype matrix described below. Heritability
estimates for RIAIL and wild isolate data are in Figure 1-source data 12 and Figure 2-source data
6, respectively.

Effect size calculations for dose response assay

We first fit a linear model with the formula (phenotype ~ strain) for all measured and principal
component traits for each concentration of arsenic trioxide using the *lm* R function. Next, we
extracted effect sizes using the *anova_stats* function from the sjstats R package [74]. Effect sizes
for dose responses are in Figure 1-source data 3.

Linkage mapping

A total of 262 RIAILs were phenotyped in the HTA described previously for control and arsenic
trioxide conditions [35,36]. The phenotype and genotype data were entered into R and scaled to

have a mean of zero and a variance of one for linkage analysis (Figure 1-source data 8).
Quantitative trait loci (QTL) were detected by calculating logarithm of odds (LOD) scores for each
marker and each trait as $-n(ln(1-r^2)/2ln(10))$, where r is the Pearson correlation coefficient
between RIAIL genotypes at the marker and phenotype trait values [75]. The maximum LOD
score for each chromosome for each trait was retained from three iterations of linkage mappings
(Figure 1-source data 11). We randomly permuted the phenotype values of each RIAIL while
maintaining correlation structure among phenotypes 1000 times to estimate the significance
threshold empirically. The significance threshold was set using a genome-wide error rate of 5%.
Confidence intervals were defined as the regions contained within a 1.5 LOD drop from the
maximum LOD score [76].
655
Near-isogenic line (NIL) generation
NILs were generated by crossing N2xCB4856 RIAILs to each parental genotype. For each NIL,
eight crosses were performed followed by six generations of selfing to homozygose the genome.
Reagents used to generate NILs are detailed in the Key Resources Table. The NILs responses
to 1000 µM arsenic trioxide were quantified using the HTA described above (Figure 1-source data
13). NIL whole-genome sequencing and analysis was performed as described previously [77]
(Figure 1-source data 16).
663
Genome-wide association mapping
Genome-wide association (GWA) mapping was performed using phenotype data from 86
C. elegans isotypes (Figure 2-source data 1). Genotype data were acquired from the latest VCF
release (Release 20180527) from CeNDR that was imputed as described previously [32]. We
used BCFtools [78] to filter variants that had any missing genotype calls and variants that were
below 5% minor allele frequency. We used PLINK v1.9 [79,80] to LD-prune the genotypes at a
threshold of $r^2 < 0.8$, using *--indep-pairwise 50 10 0.8*. This resulting genotype set consisted of
59,241 markers that were used to generate the realized additive kinship matrix using the *A.mat*
function in the *rrBLUP* R package [70] (Figure 2-source data 5). These markers were also used
for genome-wide mapping. However, because these markers still have substantial LD within this
genotype set, we performed eigen decomposition of the correlation matrix of the genotype matrix
using *eigs_sym* function in Rspectra package [81]. The correlation matrix was generated using
the *cor* function in the correlateR R package [82]. We set any eigenvalue greater than one from
this analysis to one and summed all of the resulting eigenvalues [83]. This number was 500.761,
which corresponds to the number of independent tests within the genotype matrix. We used the
GWAS function in the rrBLUP package to perform genome-wide mapping with the following
command: *rrBLUP::GWAS(pheno = PC1, geno = Pruned_Markers, K = KINSHIP, min.MAF =*
0.05, n.core = 1, P3D = FALSE, plot = FALSE). To perform fine-mapping, we defined confidence
intervals from the genome-wide mapping as +/- 100 SNVs from the rightmost and leftmost
markers above the Bonferroni significance threshold. We then generated a QTL region of interest
genotype matrix that was filtered as described above, with the one exception that we did not
perform LD pruning. We used PLINK v1.9 to extract the LD between the markers used for fine
mapping and the peak QTL marker identified from the genome-wide scan. We used the same
command as above to perform fine mapping, but with the reduced variant set. The workflow for
performing GWA mapping can be found on https://github.com/AndersenLab/cegwas2-nf. All trait

mapping results can be found on FigShare
(https://figshare.com/articles/GWA_results_for_all_traits_mapped_in_manuscript/7458932).
691

Generation of *dbt-1* allele replacement strains

Allele replacement strains were generated using CRISPR/Cas9-mediated genome editing, using the co-CRISPR approach [84] with Cas9 ribonucleoprotein delivery [85]. Alt-R^{TM} crRNA and tracrRNA were purchased from IDT (Skokie, IL). tracrRNA (IDT, 1072532) was injected at a concentration of 13.6 µM. The *dpy-10* and the *dbt-1* crRNAs were injected at 4 µM and 9.6 µM, respectively. The *dpy-10* and the *dbt-1* single-stranded oligodeoxynucleotides (ssODN) repair templates were injected at 1.34 µM and 4 µM, respectively. Cas9 protein (IDT, 1074182) was injected at 23 µM. To generate injection mixes, the tracrRNA and crRNAs were incubated at 95ºC for five minutes and 10ºC for 10 minutes. Next, Cas9 protein was added and incubated for five minutes at room temperature. Finally, repair templates and nuclease-free water were added to the mixtures and loaded into pulled injection needles (1B100F-4, World Precision Instruments, Sarasota, FL). Individual injected P_0 animals were transferred to new 6 cm NGM plates approximately 18 hours after injections. Individual F_1 rollers were then transferred to new 6 cm plates to generate self-progeny. The region surrounding the desired S78C (or C78S) edit was then amplified from F_1 rollers using primers oECA1163 and oECA1165. The PCR products were digested using the *Sfc*I restriction enzyme (R0561S, New England Biolabs, Ipswich, MA). Differential band patterns signified successfully edited strains because the N2 S78C, which is encoded by the CAG codon, creates an additional *Sfc*I cut site. Non-Dpy, non-Rol progeny from homozygous edited F_1 animals were propagated. If no homozygous edits were obtained, heterozygous F_1 progeny were propagated and screened for the presence of the homozygous edits. F_1 and F_2 progeny were then Sanger sequenced to verify the presence of the proper edited sequence. The phenotypes of allele swap strains in control and arsenic trioxide conditions were measured using the HTA described above (Figure 1-source data 13). PCA phenotypes for allele-swap strains were generated the same way as described above for GWA mapping traits and are located in Figure 1-source data 13.

Rescue with 13-methyltetradecanoic acid

Strains were grown as described for a standard HTA experiment. In addition to adding arsenic trioixde to experimental wells, we also added a range of C15iso (13-methyltetradecanoic acid, Matreya Catalog # 1605) concentrations to assay rescue of arsenic effects (Figure 4-source data 5).

Growth conditions for metabolite profiling

For L1 larval stage assays, chunks (~1 cm) were taken from starved plates and placed on multiple fresh 10 cm plates. Prior to starvation, animals were washed off of the plates using M9, and embryos were prepared by bleach synchronization. Approximately 40,000 embryos were resuspended in 25 ml of K medium and allowed to hatch overnight at 20ºC. L1 larvae were fed 15 mg/ml of HB101 lysate the following morning and allowed to grow at 20ºC for 72 hours. We harvested 100,000 embryos from gravid adults by bleaching. These embryos were hatched overnight in 50 ml of K medium in a 125 ml flask. The following day, we added arsenic trioxide to a final concentration of 100 µM and incubated the cultures for 24 hours. After 24 hours, we added

HB101 bacterial lysate (2 mg/ml) to each culture. Finally, we transferred the cultures to 50 ml conical tubes, centrifuged the cultures at 3000 RPM for three minutes to separate the pellet and supernatant. The supernatant and pellets from the cultures were frozen at -80ºC and prepared for analysis. For young adult stage assays, 45,000 animals per culture were prepared as described above but in S medium, at a density of three animals per microliter, and fed HB101 lysate (5 mg/mL). These cultures were shaken at 200 RPM, 20ºC in 50 mL Erlenmeyer flasks for 62 h. For harvesting, we settled 15 mL of cultures for 15 minutes at room temperature and then pipetted the top 12 mL of solution off of the culture. The remaining 3 mL of animal pellet was washed with 10 mL of M9, centrifuged at 1000 g for one minute, and then the supernatant removed. This wash was repeated once more with M9 and again with water. The final nematode pellet was snap frozen in liquid nitrogen.

Nematode extraction

Pellets were lyophilized 18-24 hours using a VirTis BenchTop 4K Freeze Dryer until a chalky consistency was achieved. Dried pellets were transferred to 1.5 mL microfuge tubes and dry pellet weight recorded. Pellets were disrupted in a Spex 1600 MiniG tissue grinder after the addition of three stainless steel grinding balls to each sample. Microfuge tubes were placed in a Cryoblock (Model 1660) cooled in liquid nitrogen, and samples were disrupted at 1100 RPM for two cycles of 30 seconds. Each sample was individually dragged across a microfuge tube rack eight times, inverted, and flicked five times to prevent clumping. This process was repeated two additional rounds for a total of six disruptions. Pellets were transferred to 4 mL glass vials in 3 mL 100% ethanol. Samples were sonicated for 20 minutes (on/off pulse cycles of two seconds at power 90 A) using a Qsonica Ultrasonic Processor (Model Q700) with a water bath cup horn adaptor (Model 431C2). Following sonication, glass vials were centrifuged at 2750 RCF for five minutes in an Eppendorf 5702 Centrifuge using rotor F-35-30-17. The resulting supernatant was transferred to a clean 4 mL glass vial and concentrated to dryness in an SC250EXP Speedvac Concentrator coupled to an RVT5105 Refrigerated Vapor Trap (Thermo Scientific). The resulting powder was suspended in 100% ethanol according to its original dry pellet weight: 0.01 mL 100% ethanol per mg of material. The suspension was sonicated for 10 minutes (pulse cycles of two seconds on and three seconds off at power 90 A) followed by centrifugation at 20,817 RCF in a refrigerated Eppendorf centrifuge 5417R at 4°C. The resulting supernatant was transferred to an HPLC vial containing a Phenomenex insert (cat #AR0-4521-12) and centrifuged at 2750 RCF for five minutes in an Eppendorf 5702 centrifuge. The resulting supernatant was transferred to a clean HPLC vial insert and stored at -20°C or analyzed immediately.

Mass spectrometric analysis

Reversed-phase chromatography was performed using a Dionex Ultimate 3000 Series LC system (HPG-3400 RS High Pressure pump, TCC-3000RS column compartment, WPS-3000TRS autosampler, DAD-3000 Diode Array Detector) controlled by Chromeleon Software (ThermoFisher Scientific) and coupled to an Orbitrap Q-Exactive mass spectrometer controlled by Xcalibur software (ThermoFisher Scientific). Metabolites were separated on a Kinetex EVO C18 column, 150 mm x 2.1 mm, particle size 1.7 μm, maintained at 40°C with a flow rate of 0.5 mL/min. Solvent A: 0.1% ammonium acetate in water; solvent B: acetonitrile (ACN). A/B gradient started at 5% B for 30 seconds, followed by a linear gradient to 95% B over 13.5 minutes, then a

linear gradient to 100% B over three minutes. 100% B was maintained for one minute. Column
was washed after each run with 5:1 isopropanol:ACN, flow rate of 0.12 mL/min for five minutes,
followed by 100% ACN for 2.9 minutes, a linear gradient to 95:5 water:ACN over 0.1 minutes, and
then 95:5 water:ACN for two minutes with a flow rate of 0.5 mL/min. A heated electrospray
ionization source (HESI-II) was used for the ionization with the following mass spectrometer
parameters: spray voltage: 3 kV; capillary temperature: 320°C; probe heater temperature: 300°C;
sheath gas: 70 AU; auxiliary gas flow: 2 AU; resolution: 240,000 FWHM at m/z 200; AGC target:
5e6; maximum injection time: 300 ms. Each sample was analyzed in negative and positive modes
with m/z range 200-800. Fatty acids and most ascarosides were detected as [M-H]$^-$ ions in
negative ionization mode. Peaks of known abundant ascarosides and fatty acids were used to
monitor mass accuracy, chromatographic peak shape, and instrument sensitivity for each sample.
Processed metabolite measures can be found in Figure 4-source data 1-4 [86].
789
790 **Statistical analyses**
791 All *p*-values testing the differences of strain phenotypes in the NIL, allele-replacement, and
792 C15ISO experiments were performed in R using the *TukeyHSD* function with an ANOVA model
793 with the formula (*phenotype ~ strain*). *p*-values of individual pairwise strain comparisons are
794 reported in each figure legend.
795
CRISPR-Cas9 gene editing in human cells
Gene-editing experiments were performed in a single parallel culture experiment using human
293T cells (ATCC) grown in DMEM with 10% FBS. On day zero, 300,000 cells were seeded per
well in a six-well plate format. The following day, two master mixes were prepared: a) LT-1
transfection reagent (Mirus) was diluted 1:10 in Opti-MEM and incubated for five minutes; b) a
DNA mix of 500 ng Cas9-sgRNA plasmid (Supplementary File 1-2) with 250 pmol repair template
oligonucleotide was diluted in Opti-MEM in a final volume of 100 µL. 250 µL of the lipid mix was
added to each of the DNA mixes and incubated at room temperature for 25 minutes. Following
incubation, the full 350 µL volume of DNA and lipid mix was added dropwise to the cells. These
six-well plates were then centrifuged at 1000 x g for 30 minutes. After six hours, the media on the
cells was replaced. For the next six days, cells were expanded and passaged as needed. On day
seven, one million cells were taken from each set of edited and unedited cells and placed into
separate T75s with either media-only or 5 µM arsenic-containing media. Days seven to fourteen,
arsenic and media-only conditions were maintained at healthy cell densities. Days fourteen to
eighteen, arsenic exposed cell populations were maintained off arsenic to allow the populations
to recover prior to sequencing. Media-only conditions were maintained in parallel. On day
eighteen, all arsenic and media-only conditions were pelleted for genomic DNA extraction.
813
Analysis of CRISPR-Cas9 editing in human cells
Genomic DNA was extracted from cell pellets using the QIAGEN (QIAGEN, Hilden, Germany)
Midi or Mini Kits based on the size of the cell pellet (51183, 51104) according to the
manufacturer's recommendations. DBT1 loci were first amplified with 17 cycles of PCR using a
touchdown protocol and the NEBnext 2x master mix (New England Biolabs M0541). The resulting
product served as input to a second PCR, using primers that appended a sample-specific barcode
and the necessary adaptors for Illumina sequencing. The resulting DNA was pooled, purified with

SPRI beads (A63880, Beckman Coulter, Brea, CA), and sequenced on an Illumina MiSeq with a 300-nucleotide single-end read with an eight nucleotide index read. For each sample, the number of reads exactly matching the wild-type and edited DBT1 sequence were determined (Figure 5-source data 1).

Preparing human cells for Mass Spectroscopy

Mass spectroscopy experiments used human 293T cells (ATCC) grown in DMEM with 10% FBS. On day zero, 150,000 cells were seeded into 15 cm tissue cultures dishes with 15 mL of either 2.5 µM arsenic or no arsenic media. Each condition had five replicates. On day three, the no arsenic cells were approaching confluence and required passaging. Arsenic conditions were at ~30% confluence and received a media change. On day seven, both conditions were near confluence, media was removed, and plates were rinsed with ice cold PBS, remaining liquid removed. Plates were frozen at -80ºC before processing for mass spectrometric analysis. Cells were scraped off the plates with PBS and pelleted in microfuge tubes. Cell pellets were lyophilized 18-24 hours using a VirTis BenchTop 4K Freeze Dryer and extracted in 100% ethanol using the same sonication program as described for nematode extraction. Following sonication, samples were centrifuged at 20,817 RCF in a refrigerated Eppendorf centrifuge 5417R at 4 °C. Clarified supernatant was aliquoted to a new tube and concentrated to dryness in an SC250EXP Speedvac Concentrator coupled to an RVT5105 Refrigerated Vapor Trap (Thermo Scientific). The resulting material was suspended in .1 mL 100% ethanol and analyzed by LC-MS as described. Metabolite measurements can be found in Figure 5-source data 3.

Tajima's D calculation

We used the VCF corresponding to CeNDR release 20160408 (https://elegansvariation.org/data/release/20160408) to calculate Tajima's D. Tajima's D was calculated using the *tajimas_d* function in the *cegwas* package using default parameters (window size = 500 SNVs, sliding window distance = 50 SNVs, outgroup = N2) (Figure 2-source data 9). Isolation locations of strains can be found in Figure 2-source data 10.

Funding:

This work was supported by a National Institutes of Health R01 subcontract to ECA (GM107227), the Chicago Biomedical Consortium with support from the Searle Funds at the Chicago Community Trust, a Sherman-Fairchild Cancer Innovation Award to ECA, and an American Cancer Society Research Scholar Grant to ECA (127313-RSG-15-135-01-DD), along with support to SZ from the Cell and Molecular Basis of Disease training grant (T32GM008061) and The Bernard and Martha Rappaport Fellowship. JGD is a Merkin Institute Fellow and is supported by the Next Generation Fund at the Broad Institute of MIT and Harvard. FCS, BWF, OP, and FJT were supported by R01 GM088290. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgments:

The authors would like to thank Samuel Rosenberg for assistance on early mappings of drug sensitivities, Mudra Hegde of the Broad Institute for assistance with sequence analysis, and members of the Andersen laboratory for critical reading of this manuscript.

References

1. Ravenscroft P, Brammer H, Richards K. Arsenic pollution : a global synthesis [Internet]. Chichester, U.K.: Wiley-Blackwell; 2009. Available: http://www.worldcat.org/oclc/214285927?referer=xid

2. Ratnaike RN. Acute and chronic arsenic toxicity. Postgrad Med J. 2003;79: 391–396. doi:10.1136/pmj.79.933.391

3. Mandal BK, Suzuki KT. Arsenic round the world: a review. Talanta. 2002;58: 201–235. doi:10.1016/S0039-9140(02)00268-0

4. Khairul I, Wang QQ, Jiang YH, Wang C, Naranmandura H. Metabolism, toxicity and anticancer activities of arsenic compounds. Oncotarget. 2017;8: 23905–23926. doi:10.18632/oncotarget.14733

5. Stýblo M, Drobná Z, Jaspers I, Lin S, Thomas DJ. The role of biomethylation in toxicity and carcinogenicity of arsenic: a research update. Environ Health Perspect. 2002;110 Suppl 5: 767–771. Available: https://www.ncbi.nlm.nih.gov/pubmed/12426129

6. Schlebusch CM, Gattepaille LM, Engström K, Vahter M, Jakobsson M, Broberg K. Human adaptation to arsenic-rich environments. Mol Biol Evol. 2015;32: 1544–1555. doi:10.1093/molbev/msv046

7. Chung C-J, Hsueh Y-M, Bai C-H, Huang Y-K, Huang Y-L, Yang M-H, et al. Polymorphisms in arsenic metabolism genes, urinary arsenic methylation profile and cancer. Cancer Causes Control. 2009;20: 1653–1661. doi:10.1007/s10552-009-9413-0

8. Fujihara J, Fujii Y, Agusa T, Kunito T, Yasuda T, Moritani T, et al. Ethnic differences in five intronic polymorphisms associated with arsenic metabolism within human arsenic (+3 oxidation state) methyltransferase (AS3MT) gene. Toxicol Appl Pharmacol. 2009;234: 41–46. doi:10.1016/j.taap.2008.09.026

9. Gomez-Rubio P, Meza-Montenegro MM, Cantu-Soto E, Klimecki WT. Genetic association between intronic variants in AS3MT and arsenic methylation efficiency is focused on a large linkage disequilibrium cluster in chromosome 10. J Appl Toxicol. 2010;30: 260–270. doi:10.1002/jat.1492

10. Chen GQ, Shi XG, Tang W, Xiong SM, Zhu J, Cai X, et al. Use of arsenic trioxide (As2O3) in the treatment of acute promyelocytic leukemia (APL): I. As2O3 exerts dose-dependent dual effects on APL cells. Blood. 1997;89: 3345–3353. Available: https://www.ncbi.nlm.nih.gov/pubmed/9129041

11. Antman KH. Introduction: the history of arsenic trioxide in cancer therapy. Oncologist. 2001;6 Suppl 2: 1–2. doi:10.1634/theoncologist.6-suppl_2-1

12. Murgo AJ. Clinical trials of arsenic trioxide in hematologic and solid tumors: overview of the National Cancer Institute Cooperative Research and Development Studies. Oncologist. 2001;6 Suppl 2: 22–28. doi:10.1634/theoncologist.6-suppl_2-22

13. Emi N. Arsenic Trioxide: Clinical Pharmacology and Therapeutic Results. Chemotherapy for Leukemia. Springer, Singapore; 2017. pp. 221–238. doi:10.1007/978-981-10-3332-2_13

14. de Thé H, Chomienne C, Lanotte M, Degos L, Dejean A. The t(15;17) translocation of acute
promyelocytic leukaemia fuses the retinoic acid receptor alpha gene to a novel transcribed
locus. Nature. 1990;347: 558–561. doi:10.1038/347558a0

15. Grignani F, Valtieri M, Gabbianelli M, Gelmetti V, Botta R, Luchetti L, et al. PML/RAR alpha
fusion protein expression in normal human hematopoietic progenitors dictates myeloid
commitment and the promyelocytic phenotype. Blood. 2000;96: 1531–1537. Available:
https://www.ncbi.nlm.nih.gov/pubmed/10942402

16. Zhang X-W, Yan X-J, Zhou Z-R, Yang F-F, Wu Z-Y, Sun H-B, et al. Arsenic trioxide
controls the fate of the PML-RARalpha oncoprotein by directly binding PML. Science.
2010;328: 240–243. doi:10.1126/science.1183424

17. Tomita A, Kiyoi H, Naoe T. Mechanisms of action and resistance to all-trans retinoic acid
(ATRA) and arsenic trioxide (As2O 3) in acute promyelocytic leukemia. Int J Hematol.
2013;97: 717–725. doi:10.1007/s12185-013-1354-4

18. Hoonjan M, Jadhav V, Bhatt P. Arsenic trioxide: insights into its evolution to an anticancer
agent. J Biol Inorg Chem. 2018;23: 313–329. doi:10.1007/s00775-018-1537-9

19. Zeidan AM, Gore SD. New strategies in acute promyelocytic leukemia: moving to an
entirely oral, chemotherapy-free upfront management approach. Clin Cancer Res. 2014;20:
4985–4993. doi:10.1158/1078-0432.CCR-13-2725

20. Kniazeva M, Crawford QT, Seiber M, Wang C-Y, Han M. Monomethyl branched-chain fatty
acids play an essential role in Caenorhabditis elegans development. PLoS Biol. 2004;2:
E257. doi:10.1371/journal.pbio.0020257

21. Luz AL, Godebo TR, Smith LL, Leuthner TC, Maurer LL, Meyer JN. Deficiencies in
mitochondrial dynamics sensitize Caenorhabditis elegans to arsenite and other
mitochondrial toxicants by reducing mitochondrial adaptability. Toxicology. 2017;387: 81–
94. doi:10.1016/j.tox.2017.05.018

22. Spracklin G, Fields B, Wan G, Becker D, Wallig A, Shukla A, et al. The RNAi Inheritance
Machinery of Caenorhabditis elegans. Genetics. 2017;206: 1403–1416.
doi:10.1534/genetics.116.198812

23. Watson E, MacNeil LT, Arda HE, Zhu LJ, Walhout AJM. Integration of Metabolic and Gene
Regulatory Networks Modulates the C. elegans Dietary Response. Cell. Elsevier; 2013;153:
253–266. doi:10.1016/j.cell.2013.02.050

24. Luz AL, Meyer JN. Effects of reduced mitochondrial DNA content on secondary
mitochondrial toxicant exposure in Caenorhabditis elegans. Mitochondrion. 2016;30: 255–
264. doi:10.1016/j.mito.2016.08.014

25. Luz AL, Godebo TR, Bhatt DP, Ilkayeva OR, Maurer LL, Hirschey MD, et al. From the
Cover: Arsenite Uncouples Mitochondrial Respiration and Induces a Warburg-like Effect in
Caenorhabditis elegans. Toxicol Sci. 2016;152: 349–362. doi:10.1093/toxsci/kfw093

26. Schmeisser S, Schmeisser K, Weimer S, Groth M, Priebe S, Fazius E, et al. Mitochondrial
hormesis links low-dose arsenite exposure to lifespan extension. Aging Cell. 2013;12: 508–
517. doi:10.1111/acel.12076

27.   Wyatt LH, Luz AL, Cao X, Maurer LL, Blawas AM, Aballay A, et al. Effects of methyl and
inorganic mercury exposure on genome homeostasis and mitochondrial function in
Caenorhabditis elegans. DNA Repair . 2017;52: 31–48. doi:10.1016/j.dnarep.2017.02.005

28.   Large EE, Xu W, Zhao Y, Brady SC, Long L, Butcher RA, et al. Selection on a Subunit of
the NURF Chromatin Remodeler Modifies Life History Traits in a Domesticated Strain of
Caenorhabditis elegans. PLoS Genet. Public Library of Science; 2016;12: e1006219.
doi:10.1371/journal.pgen.1006219

29.   Wang Y, Ezemaduka AN, Li Z, Chen Z, Song C. Joint Toxicity of Arsenic, Copper and
Glyphosate on Behavior, Reproduction and Heat Shock Protein Response in
Caenorhabditis elegans. Bull Environ Contam Toxicol. 2017;98: 465–471.
doi:10.1007/s00128-017-2042-5

30.   Andersen EC, Bloom JS, Kruglyak L, Félix M-A, Ghosh R, Gerke JP, et al. Chromosome-
scale selective sweeps shape *Caenorhabditis elegans* genomic diversity. Nat Genet.
Nature Publishing Group, a division of Macmillan Publishers Limited. All Rights Reserved.;
2012;44: 285–290. doi:10.1038/ng.1050

31.   Cook DE, Zdraljevic S, Tanny RE, Seo B, Riccardi DD, Noble LM, et al. The Genetic Basis
of Natural Variation in Caenorhabditis elegans Telomere Length. Genetics. 2016;204: 371–
383. doi:10.1534/genetics.116.191148

32.   Cook DE, Zdraljevic S, Roberts JP, Andersen EC. CeNDR, the Caenorhabditis elegans
natural diversity resource. Nucleic Acids Res. 2016; doi:10.1093/nar/gkw893

33.   Zdraljevic S, Andersen EC. Natural diversity facilitates the discovery of conserved
chemotherapeutic response mechanisms. Curr Opin Genet Dev. 2017;47: 41–47.
doi:10.1016/j.gde.2017.08.002

34.   Jia F, Cui M, Than MT, Han M. Developmental Defects of Caenorhabditis elegans Lacking
Branched-chain α-Ketoacid Dehydrogenase Are Mainly Caused by Monomethyl Branched-
chain Fatty Acid Deficiency. J Biol Chem. 2016;291: 2967–2973.
doi:10.1074/jbc.M115.676650

35.   Andersen EC, Shimko TC, Crissman JR, Ghosh R, Bloom JS, Seidel HS, et al. A Powerful
New Quantitative Genetics Platform, Combining *Caenorhabditis elegans* High-Throughput
Fitness Assays with a Large Collection of Recombinant Strains. G3 . Genetics Society of
America; 2015;5: g3.115.017178–920. doi:10.1534/g3.115.017178

36.   Zdraljevic S, Strand C, Seidel HS, Cook DE, Doench JG, Andersen EC. Natural variation in
a single amino acid substitution underlies physiological responses to topoisomerase II
poisons. PLoS Genet. 2017;13: e1006891. doi:10.1371/journal.pgen.1006891

37.   Cohen J. Statistical Power Analysis for the Behavioral Sciences [Internet]. Routledge; 2013.
Available: https://market.android.com/details?id=book-2v9zDAsLvA0C

38.   Adeva-Andany MM, López-Maside L, Donapetry-García C, Fernández-Fernández C, Sixto-
Leal C. Enzymes involved in branched-chain amino acid metabolism in humans. Amino
Acids. 2017;49: 1005–1028. doi:10.1007/s00726-017-2412-7

39.   Bergquist ER, Fischer RJ, Sugden KD, Martin BD. Inhibition by methylated organo-

arsenicals of the respiratory 2-oxo-acid dehydrogenases. J Organomet Chem. 2009;694:
973–980. doi:10.1016/j.jorganchem.2008.12.028

40. Reed LJ, Hackert ML. Structure-function relationships in dihydrolipoamide acyltransferases.
J Biol Chem. 1990;265: 8971–8974. Available:
https://www.ncbi.nlm.nih.gov/pubmed/2188967

41. Kniazeva M, Euler T, Han M. A branched-chain fatty acid is involved in post-embryonic
growth control in parallel to the insulin receptor pathway and its biosynthesis is feedback-
regulated in C. elegans. Genes Dev. 2008;22: 2102–2110. doi:10.1101/gad.1692008

42. Baugh LR. To Grow or Not to Grow: Nutritional Control of Development During
Caenorhabditis elegans L1 Arrest. Genetics. 2013;194: 539–555.
doi:10.1534/genetics.113.150847

43. Watts JL, Ristow M. Lipid and Carbohydrate Metabolism in Caenorhabditis elegans.
Genetics. 2017;207: 413–446. doi:10.1534/genetics.117.300106

44. Entchev EV, Schwudke D, Zagoriy V, Matyash V, Bogdanova A, Habermann B, et al. LET-
767 is required for the production of branched chain and long chain fatty acids in
Caenorhabditis elegans. J Biol Chem. 2008;283: 17550–17560.
doi:10.1074/jbc.M800965200

45. Zhu H, Shen H, Sewell AK, Kniazeva M, Han M. A novel sphingolipid-TORC1 pathway
critically promotes postembryonic development in Caenorhabditis elegans. Elife. 2013;2:
e00429. doi:10.7554/eLife.00429

46. Forbes SA, Bhamra G, Bamford S, Dawson E, Kok C, Clements J, et al. The Catalogue of
Somatic Mutations in Cancer (COSMIC). Curr Protoc Hum Genet. 2008;Chapter 10: Unit
10.11. doi:10.1002/0471142905.hg1011s57

47. Paul S, Banerjee N, Chatterjee A, Sau TJ, Das JK, Mishra PK, et al. Arsenic-induced
promoter hypomethylation and over-expression of ERCC2 reduces DNA repair capacity in
humans by non-disjunction of the ERCC2-Cdk7 complex. Metallomics. 2014;6: 864–873.
doi:10.1039/c3mt00328k

48. Shen S, Li X-F, Cullen WR, Weinfeld M, Le XC. Arsenic binding to proteins. Chem Rev.
2013;113: 7769–7792. doi:10.1021/cr300015c

49. Li J, Packianathan C, Rossman TG, Rosen BP. Nonsynonymous Polymorphisms in the
Human AS3MT Arsenic Methylation Gene: Implications for Arsenic Toxicity. Chem Res
Toxicol. 2017;30: 1481–1491. doi:10.1021/acs.chemrestox.7b00113

50. Pettit FH, Yeaman SJ, Reed LJ. Purification and characterization of branched chain alpha-
keto acid dehydrogenase complex of bovine kidney. Proc Natl Acad Sci U S A. 1978;75:
4881–4885. Available: https://www.ncbi.nlm.nih.gov/pubmed/283398

51. Heffelfinger SC, Sewell ET, Danner DJ. Identification of specific subunits of highly purified
bovine liver branched-chain ketoacid dehydrogenase. Biochemistry. 1983;22: 5519–5522.
doi:10.1021/bi00293a011

52. Yeaman SJ. The 2-oxo acid dehydrogenase complexes: recent advances. Biochem J.

1989;257: 625–632. Available: https://www.ncbi.nlm.nih.gov/pubmed/2649080

53. Kato M, Chuang JL, Tso S-C, Wynn RM, Chuang DT. Crystal structure of pyruvate
dehydrogenase kinase 3 bound to lipoyl domain 2 of human pyruvate dehydrogenase
complex. EMBO J. 2005;24: 1763–1774. doi:10.1038/sj.emboj.7600663

54. Burrage LC, Nagamani SCS, Campeau PM, Lee BH. Branched-chain amino acid
metabolism: from rare Mendelian diseases to more common disorders. Hum Mol Genet.
2014;23: R1–8. doi:10.1093/hmg/ddu123

55. Tönjes M, Barbus S, Park YJ, Wang W, Schlotter M, Lindroth AM, et al. BCAT1 promotes
cell proliferation through amino acid catabolism in gliomas carrying wild-type IDH1. Nat
Med. 2013;19: 901–908. doi:10.1038/nm.3217

56. Hughes-Fulford M, Chen Y, Tjandrawinata RR. Fatty acid regulates gene expression and
growth of human prostate cancer PC-3 cells. Carcinogenesis. 2001;22: 701–707.
doi:10.1093/carcin/22.5.701

57. Agostini M, Silva SD, Zecchin KG, Coletta RD, Jorge J, Loda M, et al. Fatty acid synthase
is required for the proliferation of human oral squamous carcinoma cells. Oral Oncol.
2004;40: 728–735. doi:10.1016/j.oraloncology.2004.01.011

58. Coombs CC, Tavakkoli M, Tallman MS. Acute promyelocytic leukemia: where did we start,
where are we now, and the future. Blood Cancer J. 2015;5: e304. doi:10.1038/bcj.2015.25

59. Thompson OA, Snoek LB, Nijveen H, Sterken MG, Volkers RJM, Brenchley R, et al.
Remarkably Divergent Regions Punctuate the Genome Assembly of the Caenorhabditis
elegans Hawaiian Strain CB4856. Genetics. 2015;200: 975–989.
doi:10.1534/genetics.115.175950

60. Hodgkin J, Doniach T. Natural variation and copulatory plug formation in Caenorhabditis
elegans. Genetics. 1997;146: 149–164. Available:
https://www.ncbi.nlm.nih.gov/pubmed/9136008

61. Tajima F. Statistical method for testing the neutral mutation hypothesis by DNA
polymorphism. Genetics. 1989;123: 585–595. Available:
https://www.ncbi.nlm.nih.gov/pubmed/2513255

62. Boyd WA, Smith MV, Freedman JH. Caenorhabditis elegans as a model in developmental
toxicology. Methods Mol Biol. Totowa, NJ: Humana Press; 2012;889: 15–24.
doi:10.1007/978-1-61779-867-2_3

63. Andersen EC, Bloom JS, Gerke JP, Kruglyak L. A variant in the neuropeptide receptor npr-
1 is a major determinant of *Caenorhabditis elegans* growth and physiology. 2014;10:
e1004156. doi:10.1371/journal.pgen.1004156

64. García-González AP, Ritter AD, Shrestha S, Andersen EC, Yilmaz LS, Walhout AJM.
Bacterial Metabolism Affects the C. elegans Response to Cancer Chemotherapeutics. Cell.
2017;169: 431–441.e8. doi:10.1016/j.cell.2017.03.046

65. Shimko TC, Andersen EC. COPASutils: an R package for reading, processing, and
visualizing data from COPAS large-particle flow cytometers. PLoS One. Public Library of

Science; 2014;9: e111090. doi:10.1371/journal.pone.0111090

66.   Team RC. R: A language and environment for statistical computing [Internet]. Vienna,
Austria; 2014. 2017.

67.   Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models using lme4
[Internet]. arXiv [stat.CO]. 2014. Available: http://arxiv.org/abs/1406.5823

68.   Bloom JS, Kotenko I, Sadhu MJ, Treusch S, Albert FW, Kruglyak L. Genetic interactions
contribute less than additive effects to quantitative trait variation in yeast. Nat Commun.
2015;6: 8712. doi:10.1038/ncomms9712

69.   David Clifford PM. The regress function. R News. 2006: 6–10. Available: https://cran.r-
project.org/web/packages/regress/regress.pdf

70.   Endelman JB. Ridge Regression and Other Kernels for Genomic Selection with R Package
rrBLUP. The Plant Genome Journal. Crop Science Society of America; 2011;4: 250–256.
doi:10.3835/plantgenome2011.08.0024

71.   Covarrubias-Pazaran G. Genome-Assisted Prediction of Quantitative Traits Using the R
Package sommer. PLoS One. 2016;11: e0156744. doi:10.1371/journal.pone.0156744

72.   Su G, Christensen OF, Ostersen T, Henryon M, Lund MS. Estimating additive and non-
additive genetic variances and predicting genetic merits using genome-wide dense single
nucleotide polymorphism markers. PLoS One. 2012;7: e45293.
doi:10.1371/journal.pone.0045293

73.   Endelman JB, Jannink J-L. Shrinkage estimation of the realized relationship matrix. G3 .
2012;2: 1405–1413. doi:10.1534/g3.112.004259

74.   Lüdecke D. sjstats: Statistical Functions for Regression Models [Internet]. 2018.
doi:10.5281/zenodo.1442812

75.   Bloom JS, Ehrenreich IM, Loo WT, Lite T-LV, Kruglyak L. Finding the sources of missing
heritability in a yeast cross. Nature. Nature Publishing Group; 2013;494: 1–6.
doi:10.1038/nature11867

76.   Broman KW, Wu H, Sen S, Churchill GA. R/qtl: QTL mapping in experimental crosses.
Bioinformatics. 2003;19: 889–890. Available:
http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=12724300&retmod
e=ref&cmd=prlinks

77.   Brady S, Evans K, Bloom J, Tanny R, Cook D, Giuliani S, et al. Common loci underlie
natural variation in diverse toxin responses [Internet]. bioRxiv. 2018. p. 325399.
doi:10.1101/325399

78.   Li H. A statistical framework for SNP calling, mutation discovery, association mapping and
population genetical parameter estimation from sequencing data. Bioinformatics. 2011;27:
2987–2993. doi:10.1093/bioinformatics/btr509

79.   Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, et al. PLINK: a tool
set for whole-genome association and population-based linkage analyses. Am J Hum
Genet. 2007;81: 559–575. doi:10.1086/519795

80. Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. Second-generation
PLINK: rising to the challenge of larger and richer datasets. Gigascience. 2015;4: 7.
doi:10.1186/s13742-015-0047-8

81. Qiu Y. RSpectra [Internet]. Github; Available: https://github.com/yixuan/RSpectra

82. Bilgrau AE. correlateR [Internet]. Github; 2018. Available:
https://github.com/AEBilgrau/correlateR

83. Li J, Ji L. Adjusting multiple testing in multilocus analyses using the eigenvalues of a
correlation matrix. Heredity . 2005;95: 221–227. doi:10.1038/sj.hdy.6800717

84. Kim H, Ishidate T, Ghanta KS, Seth M, Conte D, Shirayama M, et al. A co-CRISPR strategy
for efficient genome editing in Caenorhabditis elegans. Genetics. 2014;197: 1069–1080.
doi:10.1534/genetics.114.166389

85. Paix A, Folkmann A, Rasoloson D, Seydoux G. High Efficiency, Homology-Directed
Genome Editing in Caenorhabditis elegans Using CRISPR-Cas9 Ribonucleoprotein
Complexes. Genetics. 2015;201: 47–54. doi:10.1534/genetics.115.179382

86. Artyukhin AB, Zhang YK, Akagi AE, Panda O, Sternberg PW, Schroeder FC. Metabolomic
"Dark Matter" Dependent on Peroxisomal β-Oxidation in Caenorhabditis elegans. J Am
Chem Soc. 2018;140: 2841–2852. doi:10.1021/jacs.7b11811

1118

Supplementary Figures:

Figure 1-figure supplement 1: Arsenic trioxide dose response of four diverged *C. elegans* strains

Arsenic trioxide concentration in μM is plotted on the x-axis and the (A) normalized brood size, (B) median progeny length, or (C) the first principal component are plotted on the y-axis. For panels A-B, the y-axis values represent individual phenotypic measurements subtracted from the mean value in 0 μM arsenic trioxide. At least 15 replicates for each strain and condition are represented by Tukey box plots. Box plots are colored by strain (CB4856:blue, DL238:teal, JU775:pink, and N2:orange).

Figure 1-figure supplement 2: Trait correlations and principal component loadings of
arsenic trioxide dose response
The top panels for A-E represent the trait correlations of the measured traits. The bottom panels for A-E represent
the contribution of each trait to the principal components that explain 90% of the total variance in the experiment for
each condition: (A) Water, (B) 250 µM, (C) 500 µM, (D) 1000 µM, and (E) 2000 µM of arsenic trioxide. For each
plot, the tile color corresponds to the value corresponding to trait Pearson's correlation coefficient for the top panels
and principal component loading value for the bottom panels, where yellow colors represent higher values.

Figure 1-figure supplement 3: Effect size and broad-sense heritability estimates for the arsenic trioxide dose response

Each panel corresponds to a concentration of arsenic trioxide concentration, which is indicated above the plot panel. The x-axis represents the partial omega squared (ω_p^2) effect-size estimate and the y-axis represent the broad-sense heritability estimate (H^2). Each dot represents a different measured or principal component trait. The three traits discussed throughout the manuscript are highlighted — red: first principal component, pink: animal length, and blue: brood size.

Figure 1-figure supplement 4: Trait correlations and principal component loadings of
linkage mapping experiment
(A) The trait Pearson's correlation coefficient of the assay- and control-regressed measured traits. (B) The
contribution of each measured trait to the principal components that explain 90% of the total variance in the linkage
mapping experiment, which was performed at 1000 μM. For each plot, the tile color corresponds to the value, where
yellow colors represent higher values.

1152
Figure 1-figure supplement 5: RIAIL phenotypes from the linkage mapping experiment
Tukey box plots of the first principal component (A), assay- and control-regressed brood sizes and (B) median
animal lengths of the N2 and CB4856 RIAIL panel after exposure to arsenic trioxide. Each dot corresponds to the
phenotype for a single RIAIL. The RIAILs are separated by the N2 (orange) or CB4856 (blue) genotype at each
QTL detected by linkage mapping.

Figure 1-figure supplement 6: Linkage mapping results for brood size, animal length, and
the first principal component
Linkage mapping plots for regressed brood size (teal), median animal length (pink), and the first principal
component (black) in the presence of 1000 μM arsenic trioxide are shown. The significance values (logarithm of
odds, LOD, ratio) for 1454 markers between the N2 and CB4856 strains are on the y-axis, and the genomic position
(Mb) separated by chromosome is plotted on the x-axis. The associated 1.5 LOD-drop confidence intervals are
represented by teal, pink, and boxes for brood size, median animal length, and the first principal component
respectively.

Figure 1-figure supplement 7: Genomic-heritability estimates of linkage mapping traits
The genomic broad (H^2)- and narrow (h^2)-sense heritability estimates calculated using the expectation (E(A)) of the realized relatedness matrix or the realized relatedness matrix (A). Each dot represents a measured or principal component trait. Dots are colored black if that trait mapped to the center of chromosome II and red if no QTL was detected on the center of chromosome II. The brood size, animal length, and first principal component traits are marked for clarity.

1172 **Figure 1-figure supplement 8: Linkage mapping QTL summary**
1173 All QTL identified by linkage mapping are shown. Traits are labeled on the y-axis and the genomic position
1174 in Mb is plotted on the x-axis. Triangles represent the peak QTL position and bars represent the associated
1175 1.5-LOD drop QTL confidence interval. Triangles and bars are colored based on the associated LOD score,
1176 where red colors correspond to higher LOD values.
1177

1178 **Figure 1-figure supplement 9: NIL recapitulation of chromosome II QTL**
1179 Tukey box plots of near-isogenic line (NIL) phenotype values for the brood size (top panel) and animal length
1180 (bottom panel) traits in the presence of 1000 μM arsenic trioxide are shown. NIL genotypes are indicated below the
1181 plot as genomic ranges. The N2 brood sizes and progeny lengths are significantly different from all other strains
1182 (Brood size: Tukey HSD p-value < 1.56E-7; Animal Length: Tukey HSD p-value < 3.0E-14).

Figure 1-figure supplement 10: Trait correlations and principal component loadings of NIL and allele-swap recapitulation experiment

(A) The trait Pearson's correlation coefficient of the assay- and control-regressed measured traits. (B) The contribution of each measured trait to the principal components that explain 90% of the total variance in the NIL and allele swap-recapitulation experiment, which was performed at 1000 µM. For each plot, the tile color corresponds to the value, where yellow colors represent higher values.

Figure 1-figure supplement 11: Brood size and animal length are correlated with the first principal component for the NIL recapitulation experiment

The correlation between brood size (blue) or animal length (pink) with the first principal component trait for the NIL recapitulation experiment. Each dot represents an individual NIL or parental strain replicate phenotype, with the animal length and brood size phenotype values on the x-axis and the first principal component phenotype on the y-axis.

Figure 2-figure supplement 1: Trait correlations and principal component loadings of GWA mapping experiment

(A) The trait Pearson's correlation coefficient of the assay- and control-regressed Measured traits. (B) The contribution of each measured traits to the principal components that explain 90% of the total variance in the GWA mapping experiment, which was performed at 1000 µM. For each plot, the tile color corresponds to the value, where yellow colors represent higher values.

Figure 2-figure supplement 2: Genomic-heritability estimates of GWA mapping traits

The genomic broad (H^2)- and narrow (h^2)-sense heritability estimates calculated using the the realized relatedness matrix. Each dot represents a measured or principal component trait. The three traits discussed throughout the manuscript are highlighted — red: first principal component, pink: animal length, and blue: brood size.

Figure 2-figure supplement 3: GWA mapping QTL summary
All QTL identified by GWA mapping are shown. Traits are labeled on the y-axis and the genomic position in Mb is plotted on the x-axis. Triangles represent the peak QTL position and bars represent the associated QTL region of interest. Triangles and bars are colored based on the associated significance value, where red colors correspond to higher significance values.

Figure 2-figure supplement 4: Fine-mapping of the chromosome II QTL identified by GWA mapping

Fine mapping of the chromosome II region of interest (cyan region from panel A, 7.71 - 8.18 Mb) is shown. Each dot represents an SNV present in the phenotyped population. SNVs present in the CB4856 strain are shown in the left panel and SNVs present in other phenotyped strains, but REF in CB4856, are shown in the right panel. The association between the SNV and first principal component is shown on the y-axis and the genomic position of the SNV is shown on the x-axis. Dots are colored by their SnpEff predicted effect.

Figure 3-figure supplement 1: The DBT-1 C78S variant underlies arsenic trioxide sensitivity
in _C. elegans._
Tukey box plots of residual animal length (top panel) and median animal length (bottom panel) after exposure to
1000 µM arsenic trioxide are shown (N2, orange; CB4856, blue; allele replacement strains, gray). Labels correspond
to the genetic background and the corresponding residue at position 78 of DBT-1 (C for cysteine, S for serine).
Every pair-wise strain comparison is significant except for the N2 DBT-1(S78) - CB4856 comparison from animal
length (Brood size: Tukey's HSD p-value < 1.55E-5; Animal length: p-value << 1E-7)**.**

Figure 3-figure supplement 2: Brood size and animal length are correlated with the first principal component for the allele-swap recapitulation experiment

The correlation between brood size (blue) or animal length (pink) with the first principal component trait for the allele-swap recapitulation experiment. Each dot represents an individual allele-swap or parental strain replicate phenotype, with the animal length and brood size phenotype values on the x-axis and the first principal component phenotype on the y-axis.

Figure 4-figure supplement 1: Raw abundance of C17ISO for CB4856 and CB4856 allele swap.

The raw abundance of C17ISO is plotted on the y-axis for three independent replicates of the CB4856 and CB4856 allele swap strains exposed to control (teal) or 100 μM arsenic trioxide (pink) conditions. The difference between CB4856 swap mock and arsenic conditions is significant (Tukey HSD *p*-value = 0.029), but the difference between CB4856 mock and arsenic conditions is not (Tukey HSD *p*-value = 0.10).

Figure 4-figure supplement 2: Straight-chain fatty acids are not affected by arsenic trioxide
The difference in raw C15SC (left panel) or C17SC (right panel) abundances between 100 μM arsenic trioxide and control conditions is plotted on the y-axis for three independent replicates of the CB4856 and CB4856 allele swap strains and six independent replicates of the N2 and N2 allele swap strains. There are no significant differences when comparing the abundances between parental and allele-swap strains.

Figure 4-figure supplement 3: C15ISO and C17ISO to strait-chain ratios in control conditions

The C15ISO/C15SC (left panel) or C17ISO/C17SC (right panel) ratios in control conditions are plotted on the y-axis for three independent replicates of the CB4856 and CB4856 allele swap strains and six independent replicates of the N2 and N2 allele swap strains. The C15 and C17 ratios for the CB4856-CB4856 allele swap comparison are significant (C15: Tukey HSD p-value = 0.0168749; C17: Tukey HSD p-value = 0.0342525). The difference between the C17 ratio for the N2-N2 allele swap comparison is significant (Tukey HSD p-value = 0.0044667), but the difference in the C15 ratio is not significant (Tukey HSD p-value = 0.1239674).

Figure 4-figure supplement 4: Strains with the DBT-1(C78) allele produce more branched chain fatty acids in the L1 larval stage in control conditions.

Branched chain (left panel) and straight chain (right panel) fatty acid measurements in L1 animals are represented on the y-axis. There are significant differences in abundances when comparing all parental and allele-swap strains for C15ISO and C17ISO chain fatty acids (CB4856-C15ISO DBT-1(C78): Tukey HSD p-value = 0.0036201, n=3; N2-C15ISO DBT-1(C78): Tukey HSD p-value = 0.0265059, n=6; CB4856-C17ISO DBT-1(C78): Tukey HSD p-value = 0.0086572, n=3; N2-C17ISO DBT-1(C78): Tukey HSD p-value = 0.0022501, n=6). Conversely, we observe no

1257   significant differences in straight chain fatty production except for C17n production in the CB4856 background
1258   (CB4856-C15n DBT-1(C78): Tukey HSD *p*-value = 0.0787388, n=3; N2-C15n DBT-1(C78): Tukey HSD *p*-value =
1259   0.5817993, n=6; CB4856-C17n DBT-1(C78): Tukey HSD *p*-value = 0.0086572, n=3; N2-C17n DBT-1(C78): Tukey
1260   HSD *p*-value = 0.35827, n=6).

Figure 4-figure supplement 5: Young adult C15ISO and C17ISO to strait-chain ratios in control conditions

The C15ISO/C18SC (top panel) or C17ISO/C18SC(bottom panel) ratios of Young adult animals in control conditions are plotted on the y-axis for six independent replicates for all strains. There are no significant differences when comparing the abundances between parental and allele-swap strains.

Figure 4-figure supplement 6: Trait correlations and principal component loadings of C15ISO rescue experiment

(A) The trait Pearson's correlation coefficient of the assay- and control-regressed measured traits. (B) The contribution of each measured trait to the principal components that explain 90% of the total variance in the GWA mapping experiment, which was performed at 1000 µM. For each plot, the tile color corresponds to the value, where yellow colors represent higher values.

Figure 4-figure supplement 7: Brood size and animal length are correlated with the first principal component for the C15ISO rescue experiment

The correlation between brood size (blue) or animal length (pink) with the first principal component trait for the C15ISO rescue experiment. Each dot represents an individual allele-swap or parental strain replicate phenotype, with the animal length and brood size phenotype values on the x-axis and the first principal component phenotype on the y-axis.

Figure 4-figure supplement 8: Complete results from C15iso rescue experiment
Tukey box plots median animal length after C15ISO, arsenic trioxide, or arsenic trioxide and 0.64 μM C15ISO
exposure are shown (N2, orange; CB4856, blue; allele replacement strains, gray). Labels correspond to the genetic
background and the corresponding residue at position 78 of DBT-1 (C for cysteine, S for serine). Every pairwise
strain comparison is significant except for the N2 DBT-1(S78) - CB4856 comparisons (Tukey's HSD *p*-value <
1.43E-6).

Figure 5-figure supplement 1: Three-dimensional homology model of *C. elegans* DBT-1
A three-dimensional homology model of *C. elegans* DBT-1 (black) aligned to human pyruvate
dehydrogenase lipoyl domain (PDB:1Y8N) is shown. The C78 residue that confers resistance to arsenic
trioxide is highlighted in orange, and the C65 residue is highlighted in purple. The *C. elegans* K71 residue
is highlighted in red, and the human lipoylated lysine is highlighted in green.
1289

Figure 2-figure supplement 5: Tajima's _D_ across the arsenic trioxide QTL confidence
interval
Divergence, as measured by Tajima's D, is shown across the arsenic trioxide QTL confidence interval (II:7,430,000-
8,330,000). The whole-genome SNV data set [24,49] was used for Tajima's D calculations. Window size for the
calculations was 500 SNVs with a 10 SNV sliding window size. The vertical red line marks the position of the _dbt-1_
locus.

Figure 2-figure supplement 6: The worldwide distribution of the DBT-1(C78S) allele
Cysteine (REF) is shown in orange and serine (ALT) is shown in blue. Latitude and longitude coordinates
of sampling locations were used to plot individual strains on the map.

Supplementary Files:

Figure 1-source data 1: Arsenic dose response trait data for four strains in 0, 250, 500, 1000, or 2000 μM arsenic, labelled as water, arsenic1, arsenic2, arsenic3, arsenic4 in the Condition column. (Used to generate Supplementary Figure 1A-B)

Figure 1-source data 2: Broad-sense heritability estimates from arsenic dose response data. Estimates were calculated using the lmer function of the lme4 R package by fitting the mixed effect model equation lmer(value ~ (1|strain)), where value is the trait value and strain is the strain name. (Used to generate Supplementary Figure 3)

Figure 1-source data 3: Arsenic dose response estimates of effect sizes estimated by fitting the linear model: trait value ~ strain. (Used to generate Supplementary Figure 3)

Figure 1-source data 4: Arsenic dose response loadings of principal components (PCs) for the PCs that explain up to 90% of the total variance in the trait data. Condition names correspond to those in Supplementary File 1. (Used to generate Supplementary Figure 2A-D)

Figure 1-source data 5: Arsenic dose response principal component (PC) eigenvectors for the PCs that explain up to 90% of the total variance in the data set. PCA was performed all arsenic concentrations together in order to look at the relative PC value for each concentration. Condition names correspond to those in Supplementary File 1. (Used to generate Supplementary Figure 1C)

Figure 1-source data 6: Arsenic dose response loadings of principal components (PCs) for the PCs that explain up to 90% of the total variance in the trait data. PCA was performed all arsenic concentrations together in order to look at the relative PC value for each concentration.

Figure 1-source data 7: Arsenic dose response trait correlations where each row corresponds to the Pearson correlation coefficient for two traits. Condition names correspond to those in Supplementary File 1. (Used to generate Supplementary Figure 1A-D)

Figure 1-source data 8: RIAIL phenotype data used for linkage mapping. (Used to generate Figure 1B and Supplementary Figure 5)

Figure 1-source data 9: RIAIL trait correlations where each row corresponds to the Pearson correlation coefficient for two traits. (Used to generate Supplementary Figure 4A)

Figure 1-source data 10: RIAIL loadings of principal components (PCs) for the PCs that explain up to 90% of the total variance in the trait data. (Used to generate Supplementary Figure 4B)

Figure 1-source data 11: Results from linkage mapping experiment. (Used to generate Figure 1A, Supplementary Figure 6, and Supplementary Figure 8)

1343 **Figure 1-source data 12**: Genomic heritability estimates from RIAIL phenotype data. (Used to
1344 generate Supplementary Figure 7)
1345

1346 **Figure 1-source data 13**: Phenotypes of NILs and CRISPR allele swap strains in the presence
1347 of 1000 µM arsenic trioxide after correcting for strain differences in control conditions. (Used to
1348 generate Figure 1C, Figure 3, Supplementary Figure 9, and Supplementary Figure 16)
1349

1350 **Figure 1-source data 14**: NIL and CRISPR allele swap trait correlations. (Used to generate
1351 Supplementary Figure 10A)
1352

1353 **Figure 1-source data 15**: NIL and CRISPR allele swap trait loadings of principal components
1354 (PCs) for the PCs that explain up to 90% of the total variance in the trait data. (Used to generate
1355 Supplementary Figure 10B)
1356

1357 **Figure 1-source data 16**: NIL genotypes generated from whole-genome sequencing.
1358

1359 **Figure 2-source data 1**: All wild-isolate traits used for genome-wide association mapping.
1360

1361 **Figure 2-source data 2**: Wild isolate trait loadings of principal components (PCs) for the PCs that
1362 explain up to 90% of the total variance in the trait data. (Used to generate Supplementary Figure
1363 12B)
1364

1365 **Figure 2-source data 3**: Wild isolate trait correlations. (Used to generate Supplementary Figure
1366 12A).
1367

1368 **Figure 2-source data 4**: GWA mapping results for PC1. (Used to generate Figure 1A-B)
1369

1370 **Figure 2-source data 5**: Genotype matrix used for genome-wide mapping.
1371

1372 **Figure 2-source data 6**: Genomic heritability estimates of wild isolate traits. (Used to generate
1373 Supplementary Figure 13)
1374

1375 **Figure 2-source data 7**: All QTL identified by GWA mapping (Used to generate Supplementary
1376 Figure 14).
1377

1378 **Figure 2-source data 8**: Fine-mapping results for PC1. (Used to generate Figure 1C and
1379 Supplementary Figure 15)
1380

1381 **Figure 4-source data 1**: Metabolite measurements for the CB4856 and CB4856 allele swap
1382 strains (Used for Figure 4B and Supplementary Figures 18-20)
1383

1384 **Figure 4-source data 2**: Processed metabolite measurements for the CB4856 and CB4856 allele
1385 swap strains (Used for Figure 4B and Supplementary Figures 18-20)
1386

1387 **Figure 4-source data 3**: Metabolite measurements for the N2 and N2 allele swap strains (Used
1388 for Figure 4B and Supplementary Figures 18-20)

1389

1390 **Figure 4-source data 4**: Metabolite measurements for N2, CB4856, and both allele-swap strains
1391 at the L4 larval stage. (Used to generate Supplementary Figure 21)

1392

1393 **Figure 4-source data 5**: Processed phenotype data for the C15ISO rescue experiment (Used to
1394 generate Figure 4C)

1395

1396 **Figure 4-source data 6**: Trait correlations for the for the C15ISO rescue experiment (Used to
1397 generate Supplementary Figure 22A)

1398

1399 **Figure 4-source data 7**: C15ISO rescue trait loadings of principal components (PCs) for the PCs
1400 that explain up to 90% of the total variance in the trait data. (Used to generate Supplementary
1401 Figure 2B)

1402

1403 **Figure 5-source data 1**: Human cell line read data for CRISPR swap experiment in 293T
1404 cells.(Used to generate Figure 5B)

1405

1406 **Figure 5-source data 2**: Results from Fisher's exact test of human cell line read data for CRISPR
1407 swap experiment in 293T cells.

1408

1409 **Figure 5-source data 3**: Metabolite measurements from human cell line experiments.

1410

1411 **Figure 2-source data 9**: Tajima's D of GWA mapping confidence interval. (Used to generate
1412 Supplementary Figure 26)

1413

1414 **Figure 2-source data 10**: Isolation locations of strains used in GWA mapping. (Used to generate
1415 Supplementary Figure 27)

1416

1417 **Supplementary File 1**: Plasmid used for editing human cells with the S112C and R113C edits.

1418

1419 **Supplementary File 2**: Plasmid used for editing human cells with the W84C edit.

 Format of Supplementary Files:

1421

1422 **Figure 1-source data 1:** Arsenic dose response trait data for four strains in 0, 250, 500, 1000,
1423 or 2000 μM arsenic, labelled as water, arsenic1, arsenic2, arsenic3, arsenic4 in the Condition
1424 column.

1425

Column	Description
Strain	Strain name
Condition	Arsenic concentration
Trait	Trait name
Value	Trait value

1426

1427 **Figure 1-source data 2:** Broad-sense heritability estimates from arsenic dose response data.
1428 Estimates were calculated using the lmer function of the lme4 R package by fitting the mixed
1429 effect model equation lmer(value ~ (1|strain)), where value is the trait value and strain is the
1430 strain name.

Column	Description
DR_H2	Broad-sense heritability estimate
Strain_Effect	Variance of random effect (strain)
Residual_Var	Residual variance from mixed model
Coef_Gv	Coefficient of genetic variation - 100*(sqrt(genetic variance component)/response mean)
Coef_Rv	Coefficient of environmental variation - 100*(sqrt(environmental variance component)/response mean)
condition	Arsenic Concentration
trait	Trait name

1431

1432 **Figure 1-source data 3:** Arsenic dose response estimates of effect sizes estimated by fitting
1433 the linear model: trait value ~ strain. Estimates were performed for pairs of strains and for all
1434 strains. When all strains were used for the estimates, the strain1 and strain2 column values are
1435 "ALL". Condition names correspond to those in SuppData1.

Column	Description
term	ANOVA model term - only Strain effect is shown
power	Power to detect the effect at 0.05 significance threshold, based off the

	Cohen's F effect size
sumsq	Sum of squares from ANOVA
meansq	Mean squares (sumsq/df)
df	Degrees of freedom in ANOVA model
statistic	F statistic
p.value	P-value from ANOVA model
etasq	η^2 estimate of effect size
partial.etasq	Partial η^2 estimate of effect size
omegasq	ω^2 estimate of effect size
partial.omegasq	Partial ω^2 estimate of effect size
cohens.f	Cohen's F estimate of effect size
condition	Arsenic concentration
trait	Trait name
strain1	Strain name
strain2	Strain name

1436

1437 **Figure 1-source data 4:** Arsenic dose response loadings of principal components (PCs) for the
1438 PCs that explain up to 90% of the total variance in the trait data. Condition names correspond to
1439 those in SuppData1

Column	Description
Trait	Trait name
PC	Principal component
loading	Loading value
Condition	Arsenic concentration

1440

1441 **Figure 1-source data 5:** Arsenic dose response principal component (PC) eigenvectors for the
1442 PCs that explain up to 90% of the total variance in the data set. Condition names correspond to
1443 those in SuppData1.

Column	Description

Strain	Strain name
Condition	Arsenic Concentration
Trait	Principal component trait name
Value	Principal component eigenvectors

1444

1445 **Figure 1-source data 6:** Arsenic dose response loadings of principal components (PCs) for the
1446 PCs that explain up to 90% of the total variance in the trait data. PCA was performed all arsenic
1447 concentrations together in order to look at the relative PC value for each concentration

Column	Description
Trait	Trait name
PC	Principal component
loading	Loading value

1448

1449 **Figure 1-source data 7:** Arsenic dose response trait correlations where each row corresponds
1450 to the Pearson correlation coefficient for two traits. Condition names correspond to those in
1451 SuppData1.

Column	Description
trait_b	Trait name
Condition	Arsenic concentration
trait_a	Trait name
trait_cor	Pearson correlation coefficient between trait_a and trait_b

1452

1453 **Figure 1-source data 8:** RIAIL phenotype data used for linkage mapping

Column	Description
Condition	Condition name - all correspond to 1000 µM arsenic trioxide
Strain	Strain name
Trait	Trait name
Value	Trait value

1454

1455 **Figure 1-source data 9:** RIAIL trait correlations where each row corresponds to the Pearson
1456 correlation coefficient for two traits.

Column	Description

trait_b	Trait name
trait_a	Trait name
trait_cor	Pearson correlation coefficient between trait_a and trait_b

1457

1458 **Figure 1-source data 10:** RIAIL loadings of principal components (PCs) for the PCs that
1459 explain up to 90% of the total variance in the trait data.

Column	Description
Trait	Trait name
PC	Principal Component
loading	Principal Component loading value

1460

1461 **Figure 1-source data 11:** Results from linkage mapping experiment.

Column	Description
marker	Genotypic marker name
chr	Chromosome on which the marker is located
pos	Genomic position at which the marker is located, in bp (WS245)
trait	Toxin response measured by the BIOSORT in toxin.trait format
lod	LOD score indicating the strength of correlation between genotype at the marker and phenotype of RIAILs
threshold	GWER-derived LOD score above which a LOD score is considered significant
iteration	Numerical value indicating the number of fsearch() iterations at which the given LOD score was identified, where each iteration takes the highest LOD score of the previous iteration as a cofactor before performing the mapping
var_exp	For the highest significant LOD score per iteration, the amount of RIAIL phenotypic variation that can be explained by genotype at the peak marker
eff_size	Coefficient of a linear model between genotype and phenotype indicating the effect size of a QTL
ci_l_marker	Genotypic marker indicating the left boundary of a 95% confidence interval around a QTL peak marker
ci_l_pos	Position, in bp, across the chromosome indicating the left boundary of

	a 95% confidence interval around a QTL peak marker
ci_r_marker	Genotypic marker indicating the right boundary of a 95% confidence interval around a QTL peak marker
ci_r_pos	Position, in bp, across the chromosome indicating the right boundary of a 95% confidence interval around a QTL peak marker

1462

Figure 1-source data 12: Genomic heritability estimates from RIAIL phenotype data. Two
estimates are provided that both utilize a linear mixed effect model with the equation $y = X\beta +$
$Zu + \epsilon$ to estimate heritability. The estimates differ in their formulation for the strain additive
relatedness matrix, realized relatedness matrices correct for allele frequencies and the
expectation matrix does not. For both estimates, the Hadamard product of the additive
relatedness matrix was used to calculate the epistatic relatedness matrix.

Column	Description
trait	Trait name
H2_realized	Broad-sense heritability estimate using the realized kinship matrix (K), where $K = ZZ'/2\Sigma p_i(1-p_i)$. Where p_i corresponds to allele frequency.
h2_realized	Narrow-sense heritability estimate using the realized kinship matrix (K), where $K = ZZ'/2\Sigma p_i(1-p_i)$. Where p_i corresponds to allele frequency.
H2_expecation	Broad-sense heritability estimate using the non-realized kinship matrix (K), where K is the correlation of marker genotypes
h2_expecation	Narrow-sense heritability estimate using the non-realized kinship matrix (K), where K is the correlation of marker genotypes
QTL_VE	Sum of variance explained by all QTL for the given trait
QTL_II	TRUE/FALSE if the QTL corresponding to II:7296342 was identified
QTL_II_VE	Variance explained by the chromosome II:7296342 QTL, if detected

1469

1470 **Figure 1-source data 13:** Phenotypes of NILs and CRISPR allele swap strains in the presence
1471 of 1000 µM arsenic trioxide after correcting for strain differences in control conditions.

Column	Description
Strain	Strain name
Condition	Condition - all correspond to 1000 µM arsenictrioxide
u_id	Unique strain replicate id
Trait	Trait name

| Value | Trait value |

Figure 1-source data 14: NIL and CRISPR allele swap trait correlations, where each row corresponds to the Pearson correlation coefficient for two traits. All traits correspond to those in 1000 μM arsenic after correcting for strain differences in control conditions

Column	Description
trait_b	Trait name
trait_a	Trait name
trait_cor	Pearson correlation coefficient

1476

Figure 1-source data 15: NIL and CRISPR allele swap trait loadings of principal components (PCs) for the PCs that explain up to 90% of the total variance in the trait data. All traits correspond to those in 1000 μM arsenic after correcting for strain differences in control conditions

Column	Description
Trait	Trait name
PC	Principal component
loading	Loading value

1481

Figure 1-source data 16: NIL genotypes generated from whole-genome sequencing.

Column	Description
chrom	Chromosome
start	Start of region
end	End of region
sample	Strain name
gt	Genotype: 1=N2, 2=CB4856
supporting_sites	Number of sites supporting genotype call

sites	Total sites in region
DP	Average depth for all sites
switches	Number of genotype changes between sites
CIGAR	The run-length encoding of the genotypes called. B3A1B8 represents 3 B-parent calls, 1A-parent call, and 8 B-parent calls

1483

1484 **Figure 2-source data 1:** Wild isolate phenotype data in the presence of 1000 µM arsenic
1485 trioxide after correcting for growth in control conditions. There is a column for each trait.

Column	Description
Strain	Strain name
trait..1	Trait values
trait..n	Trait values

1486

1487 **Figure 2-source data 2:** Wild isolates loadings of principal components (PCs) for the PCs that
1488 explain up to 90% of the total variance in the trait data. All traits correspond to those in 1000 µM
1489 arsenic after correcting for strain differences in control conditions

Column	Description
Trait	Trait name
PC	Principal component
loading	Loading value

1490

1491 **Figure 2-source data 3:** Wild isolate trait correlations where each row corresponds to the
1492 Pearson correlation coefficient for two traits. All traits correspond to those in 1000 µM arsenic
1493 after correcting for strain differences in control conditions

Column	Description
trait_b	Trait name
trait_a	Trait name
trait_cor	Pearson correlation coefficient

1494

1495   **Figure 2-source data 4:** GWA mapping results of PC 1 trait in the presence of 1000 μM arsenic
1496   trioxide. Mapping was performed using the EMMA algorithm.

Column	Description
marker	Marker name - (chromosome_position)
chr	Chromosome on which the marker is located
pos	Marker position (bp)
log10p	-log10 of the p-value for a given marker
trait	Trait name
BF	-log10 of the Bonferroni-corrected p-value
aboveBF	1/0 - 1 if marker is above the -log10 of Bonferroni-corrected p-value
strain	Strain name
value	Strain trait value
allele	Strain genotype at QTL peak position
var.exp	Variance explained by QTL, Spearman's rank correlation coefficient squared
startPOS	Start position (bp) of QTL region of interest
endPOS	End position (bp) of QTL region of interest
peak_id	If multiple QTL were identified, this corresponds to the unique QTL ID
interval_size	Size, in bp, of QTL region of interest

1497
1498   **Figure 2-source data 5:** Genotype matrix used for genome-wide mapping. Each row
1499   corresponds to a single-nucleotide variant.

Column	Description
CHROM	Chromosome of variant
POS	Position of variant
REF	Reference allele of variant
ALT	Alternate allele of variant
Strain1	Genotype of strain 1 (-1 = REF, 1 = ALT)
Strain2	Genotype of strain 2 (-1 = REF, 1 = ALT)

...StrainN	Genotype of strain N (-1 = REF, 1 = ALT)

1500

1501 **Figure 2-source data 6:** Genomic heritability estimates from wild isolate phenotype data.
1502 Estimates were calculated by fitting a linear mixed effect model with the equation $y = X\beta +$
1503 $Zu + \epsilon$. Two models were fitted, one using only additive random effects and one using an
1504 additive and epistatic random effects. The Hadamard product of the additive relatedness matrix
1505 was used to calculate the epistatic relatedness matrix.

Column	Description
Broad_H2_add_model	Broad-sense heritability estimate using the realized kinship matrix (K), where $K = ZZ'/2\Sigma p_i(1-p_i)$. Where p_i corresponds to allele frequency.
narrow	Narrow-sense heritability estimate using the realized kinship matrix (K), where $K = ZZ'/2\Sigma p_i(1-p_i)$. Where p_i corresponds to allele frequency.
Broad_H2_add_epp_model	Broad-sense heritability estimate using the realized kinship matrix (K), where $K = ZZ'/2\Sigma p_i(1-p_i)$. Where p_i corresponds to allele frequency.
trait	Trait name

1506

1507 **Figure 2-source data 7:** All QTL detected by GWA mapping

Column	Description
Trait	Trait used for mapping
Chrom	Chromosome on which the marker is located
Start_interval	Start position of QTL
Peak_Position	Peak position of QTL
End_interval	End position of QTL

1508

1509 **Figure 2-source data 8:** Fine mapping of QTL identified from mapping the first principal
1510 component of wild-isolate phenotype data. Each row corresponds to a variant within the QTL
1511 region of interest identified from the genome-wide scan.

Column	Description
MARKER	Marker name - (chromosome_position)
CHROM	Chromosome on which the marker is located
POS	Marker position (bp)
REF	Reference allele

ALT	Alternate allele
MAF_variant	Minor allele frequency of variant
GENE_NAME	WormBase gene name
WBGeneID	WormBase gene ID
WBFeature_TYPE	WormBase feature type
WBFeature_ID	WormBase feature ID
TRANSCRIPT_BIOTYPE	Transcript biotype
VARIANT_IMPACT	SnpEff predicted effect of variant
NUCLEOTIDE_CHANGE	Nucleotide change
AMINO_ACID_CHANGE	Amino acid change that results from variant
STRAND	Strand that gene is located on
TRANSCRIPTION_START_POS	Start position (bp) of transcript
TRANSCRIPTION_END_POS	End position (bp) of transcript
PEAK_MARKER	Most significant marker identified from genome-wide scan
PEAK_MAF	Minor allele frequency of the peak marker
TRAIT	Trait name
QTL_INTERVAL_START	Start position (bp) of QTL region of interest
QTL_INTERVAL_END	End position (bp) of QTL region of interest
VARIANT_LD_WITH_PEAK_MARKER	LD (r^2) of variant with peak marker
VARIANT_LOG10p	-log10 of p-value for a given variant
STRAIN	Strain
STRAIN_GENOTYPE	Strain genotype
Phenotype_Value	Strain trait value

1512
1513 **Figure 4-source data 1:** Metabolite measurements of L1 animals exposed to 100 µM arsenic
1514 trioxide or control conditions

Column	Description
Sample	Strain name
C15iso	C15iso abundance
C15n	C15 straight-chain abundance
C17iso	C17iso abundance
C17n	C17 straight-chain abundance

1515

1516 **Figure 4-source data 2:** Processed metabolite measurements of L1 animals exposed to 100
1517 μM arsenic trioxide or control conditions. Only data for CB4856 and ECA590 were used from
1518 this data set.

Column	Description
strain	Strain name
replicate	Replicate name
concentration	Concentration of arsenic trioxide (100μM or Mock)
compound	Name of metabolite or branched/straight ratio
value	Abundance or ratio of indicated metabolites

1519

1520 **Figure 4-source data 3:** Processed metabolite measurements of N2 and ECA581 L1 animals
1521 exposed to 100 μM arsenic trioxide or control conditions.
1522 Strain,Condition,Replicate,C15_branched,C15n,15_ratio,C17iso,C17n,17_ratio

Column	Description
Strain	Strain name
Condition	Condition tested (100 μM arsenic or Water)
Replicate	Replicate name
C15_branched	C15 branched chain fatty acid abundance
C15n	C15 straight-chain acid abundance
15_ratio	Ratio of C15_branched/C15n
C17iso	C17 branched chain fatty acid abundance

Column	Description
C17n	C17 straight-chain acid abundance
17_ratio	Ratio of C17_branched/C17n

1523

1524 **Figure 4-source data 4:** Processed metabolite measurements of CB4856, N2, ECA590, and
1525 ECA581 young adult animals in control conditions.
1526 Strain,C20n:5,C15iso,C17iso,C18,C18:1,C14,C15_C20n5,C15_C18

Column	Description
Strain	Strain name
C20n:5	C20n:5 abundance
C15iso	C15iso abundance
C17iso	C17iso abundance
C15n	C15 straight-chain acid abundance
C18	C18 straight-chain acid abundance
C18:1	C18:1 straight-chain acid abundance
C15_C20n5	Ratio of C15iso to C20n5 abundance
C15_C18	Ratio of C15iso to C18 abundance

1527

1528 **Figure 4-source data 5:** C15ISO rescue experiment phenotypes of parental and CRISPR allele
1529 swap strains in the presence of 1000 μM arsenic trioxide after correcting for strain differences in
1530 control conditions.

Column	Description
Strain	Strain name
Condition	Condition - all correspond to 1000 μM arsenictrioxide
u_id	Unique strain replicate id
Trait	Trait name
Value	Trait value

1531

1532 **Figure 4-source data 6:** C15ISO rescue experiment trait correlations, where each row
1533 corresponds to the Pearson correlation coefficient for two traits. All traits correspond to those in
1534 1000 μM arsenic after correcting for strain differences in control conditions

Column	Description
trait_b	Trait name
trait_a	Trait name
trait_cor	Pearson correlation coefficient

1535

1536 **Figure 4-source data 7:** C15ISO rescue experiment trait loadings of principal components
1537 (PCs) for the PCs that explain up to 90% of the total variance in the trait data. All traits
1538 correspond to those in 1000 μM arsenic after correcting for strain differences in control
1539 conditions

Column	Description
Trait	Trait name
PC	Principal component
loading	Loading value

1540

1541 **Figure 5-source data 1:** Results from human cell editing experiment.

Column	Description
Guides	CRISPR guide and repair oligos used for swap
Edit	Amino acid edit that the guides generate
Well	Experimental well of replicate
Replicate	Replicate letter
Primers	Primers used to genotype the edit
Arsenic_Concentration	Arsenic concentration in well
Guide1_wt	WT read counts
Guide1_edit	Edited read counts
Guide1_total	Total read counts

1542

1543 **Figure 5-source data 2:** Fisher's exact test *p*-values for human cell editing experiment

Column	Description

p_val	*p*-value from Fisher's exact test of read data
Edit	Amino acid edit
Conc	Arsenic trioxide concentration
Rep	Replicate

1544

1545 **Figure 5-source data 3:** Metabolite measurements for human cell line experiments.

Column	Description
Sample	Treatment condition of cells, mock or arsenic (2.5 µM)
Replicate	Replicate name
C15iso	Abundance of C15iso
C15n	Abundance of C15SC
C15_ratio	Ratio or C15iso/C15SC
C17iso	Abundance of C17iso
C17n	Abundance of C17SC
C17_ratio	Ratio or C17iso/C17SC

1546

1547 **Figure 2-source data 9:** Tajima's D calculation of GWA QTL

Column	Description
snps	SNP index of first SNP in window
position	Genomic coordinate of window start position
window	Window Index
Td	Tajima's D calculation
swindow	Genomic coordinate of window start position
ewindow	Genomic coordinate of window end position
midwindow	Genomic coordinate middle position of window

1548

1549 **Figure 2-source data 10:** Strain isolation locations

Column	Description

strain	Strain name
GT	Allele status of DBT-1(C78S) (REF = C, ALT = S)
long	Longitude coordinate of isolation location
lat	Latitude coordinate of isolation location

1550