Haplotype-phased common marmoset embryonic stem cells for genome editing using CRISPR/Cas9

Bo Zhou[1,2], Steve S. Ho[1,2], Louis C. Leung[1,3], Thomas R. Ward[1,2], Marcus Ho[1,2], Melanie J. Plastini[1,2], Scott C. Vermilyea[4,5], Marina E. Emborg[4,5,6], Thaddeus G. Golos[4,7], Philippe Mourrain[1,3], Dimitri Perrin[8], Karen J. Parker[1,*], Alexander E. Urban[1,2,9,*]

[1]Department of Psychiatry and Behavioral Sciences, Stanford University School of Medicine, Stanford, CA 94305, USA
[2]Department of Genetics, Stanford University School of Medicine, Stanford, California 94305, USA
[3]Stanford Center for Sleep Sciences and Medicine, Stanford University School of Medicine, Stanford, CA 94305, USA
[4]Wisconsin National Primate Research Center, University of Wisconsin-Madison, Madison, WI 53715, USA
[5]Neuroscience Training Program, University of Wisconsin-Madison, Madison, WI 53705, USA
[6]Department of Medical Physics, University of Wisconsin-Madison, Madison, WI 53705, USA
[7]Departments of Comparative Biosciences and Obstetrics and Gynecology, University of Wisconsin-Madison, Madison, WI 53715, USA
[8]Science and Engineering Faculty, Queensland University of Technology, Brisbane, QLD 4001, Australia
[9]Program on Genetics of Brain Function, Stanford Center for Genomics and Personalized Medicine, Tasha and John Morgridge Faculty Scholar, Stanford Child Health Research Institute, Stanford University, Stanford, CA, USA
[*]co-corresponding authors

Corresponding authors:

Alexander E. Urban, Ph.D
3165 Porter Drive, Room 2180
Palo Alto, CA 94304-1213
USA
aeurban@stanford.edu

Karen J. Parker, Ph.D
1201 Welch Road
MSLS, Room P-113
Stanford, CA 94305-5485
USA
kjparker@stanford.edu

Running title: Marmoset stem cells for CRISPR/Cas9 targeting

Keywords: *Callithrix jacchus*, transgenics, embryonic stem cells, haplotype phasing, CRISPR/Cas9, model organisms, marmoset, induced neurons (iNs)

ABSTRACT

Due to anatomical and physiological similarities to humans, the common marmoset (*Callithrix jacchus*) is an ideal organism for the study human diseases.   Researchers are currently leveraging genome-editing technologies such as CRISPR/Cas9 to genetically engineer marmosets for the *in vivo* biomedical modeling of human neuropsychiatric and neurodegenerative diseases**.** The genome characterization of these cell lines greatly reinforces these transgenic efforts. It also provides the genomic contexts required for the accurate interpretation of functional genomics data. We performed haplotype-resolved whole-genome characterization for marmoset ESC line cj367 from the Wisconsin National Primate Research Center. This is the first haplotype-resolved analysis of a marmoset genome and the first whole-genome characterization of any marmoset ESC line. We identified and phased single-nucleotide variants (SNVs) and Indels across the genome. By leveraging this haplotype information, we then compiled a list of cj367 ESC allele-specific CRISPR targeting sites. Furthermore, we demonstrated successful Cas9 Endonuclease Dead (dCas9) expression and targeted localization in cj367 as well as sustained pluripotency after dCas9 transfection by teratoma assay. Lastly, we show that these ESCs can be directly induced into functional neurons in a rapid, single-step process. Our study provides a valuable set of genomic resources for primate transgenics in this post-genome era.

Introduction

Biomedical research has been propelled by the ability to genetically modify

conventional laboratory animals such as mice, fish, flies, and worms that have

traditionally served as model organisms. These species, however, are evolutionarily

distant to, and physiologically distinct from, humans, and are ill-suited for modeling

complex brain functions and neuropsychiatric illness (Nelson and Winslow 2009; Phillips

et al. 2014; Camus et al. 2015). Although there is a critical need for developing

sophisticated animal models with greater homology to human disease (Capitanio and

Emborg 2008; Emborg 2017; Parker et al. 2018), until recently, transgenic non-human

primate (NHP) research endeavors were a prohibitively inefficient and expensive

undertaking (Grow et al. 2016). Recent breakthroughs in NHP genome-editing

technologies, however, have advanced the possibility of gaining mechanistic insights

into the genetic basis of human brain development and function.

The common marmoset (*Callithrix jacchus*) is an ideal model organism by which

to advance the NHP engineering objective. Common marmosets are small New World

monkeys and thrive in captivity under standard laboratory settings. Like humans,

common marmosets are highly social, raise young biparentally, and demonstrate

complex cognitive abilities (Schultz-Darken et al. 2016; Ausderau et al. 2017). Their

metabolic, immunological, endocrinological, and anatomical characteristics are likewise

quite similar to those of humans (Sasaki 2015). Importantly, marmosets and humans

share conserved brain structures, such as enlarged cerebral neocortices (Wise 2008;

Izpisua-Belmonte et al. 2015), as well as complex neural circuitry (Okano et al. 2016;

Okano et al. 2012), rendering marmosets a well-suited model organism for the

development of *in vivo* biomedical modeling for human neurological, psychiatric, and

neurodegenerative diseases. Finally, marmosets also have inherent advantages over

other NHPs, such as macaques, due to their small size, high reproductive efficiency,

year-round ovarian cycles, ease of handling, and absence of serious zoonotic viral

infections which can be fatal to humans (Mansfield 2003).

Several marmoset embryonic stem cell (ESC) lines and induced pluripotent stem cell

(iPSC) lines with the capabilities of differentiating into specified cell lineages are now

available (Thomson et al. 1996; Sasaki et al. 2005; Tomioka et al. 2010; Vermilyea et al.

2017). The first transgenic primate showing successful germline transmission of

transgenes was accomplished using the marmoset in which lentiviruses expressing

enhanced green fluorescent protein (EGFP) were injected into preimplantation embryos

(Sasaki et al. 2009). A similar technology was also used to generate transgenic

marmosets expressing genetically encoded calcium indicators ((Sasaki et al. 2009; Park

et al. 2016; Tomioka et al. 2017). Shortly after the reporting of the first genetically

modified marmosets, the marmoset genome was sequenced and assembled using

various strategies including Sanger sequencing at 6× coverage (Marmoset Genome

Sequencing and Analysis Consortium 2014; Sato et al. 2015). Sequencing of the

marmoset transcriptome across various tissues has also been undertaken in an effort to

provide high-quality gene annotations for its recent genome assembly (Maudhoo et al.

2014). Furthermore, gene-editing technologies based on homologous recombination

and programmable nucleases, such as zinc-finger nucleases (ZFNs) and transcription

activator-like effector nucleases (TALENs), have been successfully applied in marmoset

ESCs and embryos (Shiozawa et al. 2011; Sato et al. 2016).

Advances in CRISPR/Cas9 technology have enabled greater possibilities for NHP

transgenic research including the introduction of copy number variations (CNVs) in

animal models. The ability to create transgenic NHP animals using CRISPR/Cas9 was

successfully demonstrated in cynomolgus monkeys in which multiple genes were

altered in a single step (Niu et al. 2014). Efforts to engineer marmoset cells using

CRISPR/Cas9 are currently underway in multiple labs around the world using the

already-established marmoset ESC and iPSC lines (Vermilyea et al. 2018), and

advances are expected in the coming years (Izpisua-Belmonte et al. 2015; Sato and

Sasaki 2018).

Genomic editing strategies stand to benefit from characterizing the genome of these

cell lines. Whole genome characterization can assist the design of CRISPR gRNAs to

gain sequence specificity and to decrease off target effects. Whole genome

characterization also provides genomic contexts for accurate interpretation of functional

genomics studies, such as measuring genome-wide expression levels and methylation

states. Haplotype phasing across the genome allows for the identification of CRISPR

sites suitable for allele-specific targeting. Whole-genome characterization in the context

of copy-number variation (CNV) analysis also allows for the identification of large CNVs

not evident from standard karyotyping analysis. This is especially important for studies

that utilize *in vitro* differentiation of neurons, as they are known to harbor mosaic ploidy

changes during development via a yet to be understood mechanism (McConnell et al.

2013; Cai et al. 2014; Knouse et al. 2016). Lastly, whole-genome characterization

provides a reference genetic background for the transgenic animals or other cell lines

derived from the ESCs.

Here, we performed whole-genome characterization in a haplotype-resolved manner for marmoset ESC line cj367 from the Wisconsin National Primate Research Center (Vermilyea et al. 2017). To the best of our knowledge, this is the first haplotype-resolved analysis of a marmoset genome and the first whole-genome characterization of any marmoset ESC line. Specifically, we identified single-nucleotide variants (SNVs) and Indels by performing deep-coverage whole-genome sequencing. We then phased these SNV and Indel haplotypes by performing linked-read sequencing at 128.5× physical coverage. Leveraging these phased haplotypes, we then compiled a list of cj367 ESC allele-specific CRISPR targeting sites. Furthermore, we demonstrated successful Cas9 Endonuclease Dead (dCas9) expression and targeted localization in cj367 as well as sustained pluripotency after dCas9 transfection by teratoma assay. We also performed independent confirmation of cj367 ESC pluripotency and also show that these ESCs can be directly induced into functional neurons in a rapid, single-step process. Our datasets provide a valuable set of resources for primate transgenic research.

Results

Characteristics of marmoset ESC line cj367 and direct induction to neurons

Characteristics of the cj367 ESCs were similar to that of human iPSCs, growing in dense, circular colonies adhered to the plate (Figure 1). Following the thawing and first passaging of the ESCs after arrival from the Wisconsin National Primate Research Center, removal of differentiated cells was only needed once. The ESCs required a daily change of media and to be passaged every 4 to 5 days (~50-80% confluency). ESC colonies have clearly defined outlines, identical optical properties, and clear nuclei

(Figure 1a). Extra care is needed when dissociating the cells from the plate because they readily detach with little to no agitation after adding a cell dissociation agent such as Accutase. Instead of aspirating the Accutase before adding PBS to fully dissociate the cells, DMEM/F12 media was added directly to the Accutase to wash the cells and then spun down to remove the Accutase/media mixture. The cj367 ESCs were also confirmed to be pluripotent by immunostaining for pluripotency markers NANOG, SSEA4, and OCT3/4 (Figure 1d-f) as was initially demonstrated elsewhere (Vermilyea et al. 2017).

We differentiated the ESCs into immature cortical glutaminergic neurons using a two-week protocol originally designed and optimized for use in human iPSC lines (Zhang et al. 2013). To test the effectiveness of this protocol, the ESCs were plated at different densities based on total cell count (100k, 150k, 200k, 250k cells). Each well of cells received the same amount of the three lentiviral vectors described in (Zhang et al. 2013), and the protocol was followed for the next nine days. Cell density did not alter the results of the differentiation; however, the ESCs behaved differently from human iPSCs in two major ways. First, compared to human iPSCs, the cells that survived the differentiation process grouped entirely around the edge of the well, and only a few cells were observed in the middle. Second, the marmoset ESCs required 3 to 4 more days to reach complete differentiation compared to human iPSCs. The protocol was stopped short at day 9/10 for the marmoset ESCs, i.e. the human day-seven-equivalent (Zhang et al. 2013), and GFP fluorescence was used as a marker to visualize the differentiation process (Figure 1b, c).

Identification of SNVs and Indels

From deep-coverage (~60×) short-insert Illumina WGS of the cj367 genome, we

identified SNVs and indels using GATK Haplotypecaller (McKenna et al. 2010). In

standard GATK Best Practices workflow (DePristo et al. 2011) for human genomes,

high-confidence SNVs and indels are filtered through a statistical learning approach

using training datasets of variants derived from population-scale data from the 1000

Genomes Project (The 1000 Genomes Project Consortium et al. 2010) or the

International HapMap Project (International HapMap Consortium 2005). Since similar

datasets are not available for marmosets, to identify high-confidence SNVs and indels,

we adopted our own statistical approach. Briefly, we analyzed the distributions of the

following parameters (QD, FS, MQ, MQRankSum, ReadPosRankSum) in the raw

outputs from GATK Haplotypecaller and determined a series of appropriate filtering

thresholds for SNVs and indels separately (see Materials and Methods).

Through this approach, we identified a total of ~3.75M SNVs (2.23M

heterozygous, 1.52M homozygous) and 0.82M indels (0.43M heterozygous, 0.38M

homozygous) (Table 1, Figure 2a, EVA accession PRJEB27676); 33% of SNVs and 35%

of indels are intragenic, and 0.73% of SNVs and 0.41% of indels are within protein

coding exons (Table 1). Furthermore, we also identified long genomic stretches

exhibiting continuous homozygosity by using a Hidden Markov Model (HMM) adopted

from Adey and colleagues (Adey et al. 2013). In cj367, continuous stretches of

chromosome 1, 2, 4, 6, 7, 9, 12, 13, 14, 15, and 22 show long continuous stretches of

homozygosity that likely resulted from inbreeding (Figure 2a, Table S1).

Haplotype Phasing

We then haplotype-phased the identified SNVs and indels in cj367 by performing

10X Genomics Chromium linked-read library preparation and sequencing (Zheng et al.

2016; Marks et al. 2018). Post sequencing analysis showed that approximately 1.26 ng

or 381 genomic equivalents of high molecular weight (HMW) genomic DNA fragments

(average size 36 kb, 66.2% > 20kb, 14.3% >100kb) were partitioned into approximately

1.27 million droplets and uniquely barcoded (16 bp) (Table 1). The linked-read library

was sequenced (2×151 bp) to 30.4× genome coverage with half of the reads coming

from HMW DNA molecules with at least 30 linked-reads (N50 Linked-Reads per

Molecule) (Table 1). We estimate the actual physical coverage (C_F) to be 128.5×.

Coverage of the mean insert by sequencing (C_R) is 8,520 bp (284bp × 30 linked-reads)

or 23.7% of the average input DNA fragment size, thus the overall sequencing coverage

$C = C_R \times C_F = 30.4\times$. In 3115 haplotype blocks (Table 1, EVA accession

PRJEB27676), 2.16M (96.9%) of heterozygous SNVs and 0.79M (83.7%) of indels were

phased. The longest phased haplotype block is 23.4 Mbp (N50 = 2.49Mbp) (Figure 2b,

Table 1, EVA accession PRJEB27676). Haplotype block sizes vary widely across

chromosomes (Figure S1, Figure 2a), and poorly phased regions correspond to regions

exhibiting continuous homozygosity (Table S1, Figure 2a, EVA accession

PRJEB27676).

Cj367 genome-specific and allele-specific CRISPR targets

We identified 44,590 targets in cj367 genome suitable for allele-specific CRISPR

targeting (Table S2). Phased variant sequences (including reverse complement) that

differ by >1 bp between the alleles were extracted to identify all possible CRISPR

targets by pattern searching for [G, C, or A]N_{20}GG (see Materials and Methods). Only

conserved high-quality targets were retained by using a selection method previously

described and validated (Sunagawa et al. 2016). We took the high-quality target

filtering process further by taking the gRNA function and structure into account. Targets

with multiple exact matches, extreme GC fractions, and those with TTTT sequence

(which might disrupt the secondary structure of gRNA) were removed. Furthermore, we

used the Vienna RNA-fold package (Lorenz et al. 2011) to identify gRNA secondary

structure and eliminated targets for which the stem loop structure for Cas9 recognition is

not able to form (Nishimasu et al. 2014). Finally, we calculated the off-target risk score

using the tool as described for this purpose (Ran et al. 2013). A very strict off-target

threshold score was chosen in which candidates with a score below 75 were rejected to

ensure that all targets are as reliable and as specific as possible.

CRISPR/Cas9 in marmoset ESC

In order to test whether CRISPR/Cas9 technology would be suitable to create

transgenic marmoset models, we tested whether sgRNA-guided CRISPR entry was

possible in cj367. After transfecting cj367 marmoset ESCs with plasmids carrying

dCas9-GFP (Chen et al. 2013) and using sgRNA sequences that target telomeres, we

successfully located dCas9-GFP fluorescence from telomeres within the nuclei of live

marmoset ESCs (three nuclei shown; Fig. 3a-c), demonstrating that the Cas9 nuclease

can indeed enter the nuclease and possibly edit the genome in a targeted fashion.

Teratoma assay demonstrate that pluripotency of ESC remains unchanged after dCas9-

GFP transfection (Figure 4).

Discussion

One of the many advantages of using the common marmoset as a model to

study human disease is that its physiology, brain structure, and aspects of behavior

more closely resemble that of humans (Okano et al. 2012; Kishi et al. 2014). The

creation of pluripotent marmoset SC lines provides a platform on which to study

molecular and cellular patterns typical of the species. Traditionally, marmoset ESC

lines have been derived from the inner cell mass of blastocysts, and all of the surviving

lines had been female (Thomson et al. 1996; Sasaki et al. 2005; Debowski et al. 2016).

Recently, three novel marmoset ESC lines have been derived from the natural morula

stage of preimplantation embryos, and one novel male ESC line has been derived from

the expanded blastocyst stage (Debowski et al. 2016). Marmoset iPSCs from fetal liver

cells and adult fibroblasts have also been reported (Tomioka et al. 2010; Farnsworth et

al. 2013; Qiu et al. 2015; Vermilyea et al. 2017). The capacity of marmoset ESC and

iPSC lines to differentiate into cardiomyocytes and hematopoietic cells by induction has

been demonstrated (Sasaki et al. 2005; Tomioka et al. 2010). They can also be

directed to neural cell differentiation using the stromal cell-derived inducing activity

method and to dopaminergic neurons using a dual-SMAD-inhibition induction method

(Sasaki et al. 2005; Tomioka et al. 2010; Vermilyea et al. 2017). Here, we

independently validate the pluripotency of the cj367 cell line (Figure d-f) and show that

ESC line cj367 can be directly induced to neurons in a rapid, single step using a

protocol developed for human pluripotent stem cells (Zhang et al. 2013) (Figure 1b, c).

In addition to revealing these biological characteristics of cj367 that are especially

applicable for neuroscience research, our results allow researchers to perform genome-

editing in cj367 using CRISPR/Cas9 in a "personalized" and allele-specific manner. Our

haplotype-resolved whole-genome characterization of cj367 also provides a reference

for all future functional genomic and epigenomics studies, such as RNA-seq, DNA

methylation analysis, or ChIP-seq, carried out using cj367 or derivative cell

lines/animals. Taken together, our data provide a valuable resource for the primate

transgenic research community in this post-genome era.

The recent successes in using CRISPR-Cas9 to disrupt specific genes such as

Ppar-γ, *Rag1*, and *Dax1* in cynomolgus (Niu et al. 2014; Kang et al. 2015) and *DMD* in

rhesus monkeys (Chen et al. 2015) forecast that the use of CRISPR-Cas9 to genetically

edit marmoset cells or embryos is within reach (Kishi et al. 2014; Izpisua-Belmonte et al.

2015; Kropp et al. 2017; Sato and Sasaki 2018). Gene-editing technologies that

leverage programmable nucleases, such as ZFNs and TALENs have already been

demonstrated in marmoset ESCs (Shiozawa et al. 2011; Sato et al. 2016). Here, we

provide further evidence on this exciting possibility by demonstrating the successful

sgRNA-guided CRISPR entry and expected dCas9-GFP localization in marmoset

ESCs, which suggests that the Cas9 nuclease can indeed enter the nucleus and edit

the genome in a sequence-directed manner. Furthermore, this not only demonstrates

the relative ease of porting existing gene-editing methodologies but also the exciting

future feasibility of creating genetic models of human disorders that provide new

insights only possible in NHP models (Capitanio and Emborg 2008; Zahs and Ashe

2010).

With the imminent application of CRISPR-Cas9 for marmosets on the horizon, it

is important to have genome characterizations of the NHP ESC and iPSC lines that are

being distributed to various research labs across the world from primate cell repositories.

It is now widely recognized that sequence variants can confound the intended on-target

and off-target sites of the gRNAs especially when appropriate controls are lacking

(Pinello et al. 2016). The CRISPR system relies on Watson and Crick base pairing to

mediate genomic editing, thus it is not surprising that genetic sequence variation can

widely affect its efficiency. By taking advantage of the extensive documentation of

human genetic variation, efforts have been made to elucidate the impact of genetic

variation on CRISPR targeting (Canver et al. 2018). Importantly, it has also been

recognized that this effect on CRISPR-targeting efficiency due to genetic sequence

variation can be leveraged to make cell line-specific or allele-specific CRISPR edits for

various studies (Zhou et al. 2018). The need for such resources to facilitate the use of

CRISPR-Cas9 in primate research has already been voiced in the literature (Luo et al.

2016). Here, we performed the first whole-genome characterization for an established

NHP ESC line and the first haplotype-phasing for a marmoset genome. We also

determined the CRISPR-targeting sites suitable for allele-specific editing for cj367

(Table S2). Our haplotype-resolved whole-genome analysis as well as a list of

CRISPR-targeting sites will not only be useful for cj367 in particular but also for future

cell lines and animal models derived from the same genome as cj367.

Possessing the unique property among mammals of routinely producing dizygotic

twins that exchange hematopoietic stem cells *in utero*, all marmoset animals are

naturally chimeric (Benirschke et al. 1962). While such chimerism will affect analyses of

DNA samples obtained from developed animals, it does not affect our whole-genome of

ESCs since the cj367 ESC line was derived before the developmental stage where the

differentiation and exchange of hematopoietic stem cells can occur.

Another important aspect of whole-genome characterization is the analyses of

structural and copy-number variation. The current quality of the marmoset reference

genome assembly does not allow for confident mapping of structural variants

(Marmoset Genome Sequencing and Analysis Consortium 2014; Sato et al. 2015).

Improvements in this regard are greatly needed. This also applies to other NHP

genome assemblies as well (Luo et al. 2016).

Our whole-genome analysis of marmoset ESC line cj367 also allows for the

accurate and confident interpretation of functional genomics datasets. Recently, RNA-

seq experiments have been carried out in marmoset ESCs that show that overall

expression patterns differ between cell lines derived from different animals under the

same culturing conditions (Debowski et al. 2016), which could be a result of different

genetic backgrounds. Furthermore, the expression patterns of various marmoset ESC

lines more closely resemble primed pluripotency states rather than naïve states even

though these cells passed all criteria for expression of established pluripotent markers

(Debowski et al. 2016). This suggests that the culturing methods established for human

pluripotent cells are not directly applicable to marmoset cells. Importantly, these results

also suggest that functional genomic and epigenomics studies such as RNA-seq,

ATAC-seq, and whole-genome bisulfite sequencing could be adopted as a tool to

screen for and identify marmoset ESC lines that may exhibit true naïve pluripotency

states.

Materials and Methods

Cell Culture

The frozen marmoset ESC starter cultures were removed from the liquid nitrogen tank (-

80 °C) and placed in a 37 °C water bath until thawed. Cells were transferred to a 15 mL

centrifuge tube and 10 mL DMEM/F12 (Thermo Scientific, SH30023.01) was added

drop-wise in order to minimize bubble formation. The mixture was centrifuged at 1000

RPM for 3 min. The supernatant was aspirated and the process was repeated allowing

the cells to be washed once more with another 10 mL DMEM/F12. Cells were re-

suspended in 1 mL of medium, which was added to a single well of Matrigel

hESC-qualified Matrix (BD Biosciences, 356234)-coated BioLite 6-well clear multi-dish

(Thermo Scientific, 130184) and incubated at 37 °C and 5% CO_2. The medium in each

well was changed daily and the cells were passaged every 4 to 5 days. For passaging,

cells were washed with DPBS (Life Technologies, 14190-144), dissociated with

Accutase (Stem Cell Technologies, 7920), and added to fresh medium on a new plate

and/or added to freezing medium for storage.

Media ingredients:

• E8 Medium (Thermo Scientific, A15169-01)

• E8 Supplement (Thermo Scientific, A15171-01)

• Glutamax (Life Technologies, 35050-061)

• Lipid Concentrate (Life Technologies, 11905-031)

• Nodal (R&D Systems, 3218-ND)

• Glutathione (Sigma, G4251)

Freezing medium: 20% DMSO (Thermo Fischer, 20688) solution (DMSO +

DMEM/F12).

Differentiating marmoset ESCs into iNs

When the marmoset ESCs reached confluence, the cells were dissociated using Accutase and plated at 250,000, 200,000, 150,000, and 100,000 cells per well. One-day post-passage, a single well was transfected with the 3 viral vectors as described in the two-week iN protocol (Zhang et al. 2013). The vectors were acquired from the Genome Virus and Vector Core (GVVC) at Stanford Neuroscience Institute (neuroscience.stanford.edu). The protocol was stopped prior to plating on mouse glia in order to achieve immature cortical glutamatergic neurons. Following transfection, cells were subjected to puromycin and doxycyclin selection and then monitored for a minimum of 7 days. For human iPSCs, full differentiation is usually completed at day 7. The marmoset ESCs required 10 to 11 days (the day of passage is defined as "day 0") to achieve full differentiation.

CRISPR-Cas9 Transfection

Marmoset cj367 embryonic stem cells were seeded onto a 35mm glass bottom microwell dish (MatTek) that was coated with 0.083 µg of BD Matrigel (BD Biosciences) per well. While the cells were maintained for approximately 48 hours to recover and return to their typical growth rate, they were visually scanned under a light microscope and manually dissected to remove any patches of spontaneously differentiated cells. At 70% confluence on the 35mm dish, the cells were transfected simultaneously with Addgene plasmids pSLQ1658-dCas9-EGFP and pSLQ1651-sgTelomere(F+E). We followed Invitrogen's protocol for the Lipofectamine LTX transfection reagent (Thermo Fisher Scientific) using 5µL with 1µg of DNA from each plasmid for one 35mm dish. The lipid-DNA complexes were adjusted to a final volume of 100µL using OptiMEM (Thermo Fisher Scientific) and added to the marmoset cells for six hours. Following six

hours of incubation, medium was changed to its original E8 formula. Five hours before

imaging, cells were incubated with a photostable DMEM (MEMO EMD Millipore)

supplemented with ROCK inhibitor to reduce photobleaching effects of the E8 media.

Finally, an additional medium change using MEMO was done immediately before

imaging.

Nuclear imaging

Marmoset cell nuclei were imaged with an OMX BLAZE 3D-Structured Illumination

Microscope (Applied Precision, GE). Cells were transferred to a modified stage adaptor

held at 37^{o}C and 5% CO_2. 1µm z-stacks (0.125µm intervals) were taken at 100x

magnification (U-PLANAPO 100X SIM objective N.A. 1.42) with a 488nm laser using

DeltaVision OMX acquisition software (version 3.70.9622.0, GE) and deconvolved with

SoftWoRx 7.0.0 software (Deltavision) before maximum projections were made for the

figures (Fiji/ImageJ).

Teratoma Generation and Histopathology

H9 (2.5million/5ul matrigel mixed with 25ul of PBS) and marmoset ES cells

(3.5million/5ul matrigel mixed with 30ul of PBS) were injected subcutaneously or into

the kidney capsule in 8-week-old male NSG mice (Jackson). Seven weeks after

injection, the mice were euthanized and the teratomas were harvested. All animal

studies were approved by Stanford University IACUC guidelines. For histological

analysis, slides were stained with hematoxylin and eosin (H&E).

Illumina WGS

Marmoset ESC (cj367) genomic DNA was extracted using the Qiagen DNeasy Blood &

Tissue Kit (Cat.D No. 69504) and quantified using the Qubit dsDNA HS Assay Kit

(Invitrogen, Waltham, MA, USA). DNA purity was verified (OD260/280 > 1.8;

OD260/230 > 1.5) using NanoDrop (Thermo Scientific, Waltham, MA, USA). Afterwards,

DNA was sheared to 200 bp to 850 bp (average size = 400 bp) using the Covaris S2

instrument (Covaris, Woburn, MA, USA). DNA was then concentrated to a volume of 50

µL using the DNA Clean & Concentrator kit (Cat. No. D4013) from Zymo Research

(Irvine, CA, USA). One short-insert WGS library was constructed using the Kapa Hyper

Prep Kit (KR0961) from Kapa Biosystems (Woburn, MA, USA) with 150 ng of sheared

DNA as input following standard manufacturer's protocol with 10 cycles of PCR.

Library-size selection (420 bp to 1 kb) was carried out on the BluePippin instrument

using the R2 marker on a 1.5% dye-free cassette from Sage Science (Beverly, MA,

USA). WGS library-size selection was then verified using the 2100 Bioanalyzer DNA

1000 Kit (Agilent Technologies, Santa Clara, CA, USA). The WGS library was then

sequenced (2×150 bp) on two lanes of the Illumina HiSeq 4000 to achieve >600 million

pass-filter paired-end reads.

Determining SNVs and Indels

Paired-end reads from the two lanes of Illumina HiSeq 4000 were combined and aligned

to the *Callithrix jacchus* genome (GenBank accession GCA_000004665.1) using BWA-

MEM version 0.7.5 (Li and Durbin 2009) followed by marking of duplicates using Picard

tools (version 1.129) (http://broadinstitute.github.io/picard/). SNVs and Indels were

called using by GATK Haplotypecaller (version 3.7) (McKenna et al. 2010) and hard

filtered for quality (QD > 2.0, FS < 60.0, MQ > 40.0, MQRankSum > -12.5,

ReadPosRankSum > -8.0).

Identifying genomic regions exhibiting continuous homozygosity

A Hidden Markov Model (HMM) was used to identify genomic regions exhibiting

continuous homozygosity. The HMM is designed with two states: homogzyous and

heterozygous. We used the hard-filtered SNVs we derived from GATK HaplotypeCaller,

and filtered again for GQ > 30 and QUAL > 50. The genome was split into 25 kb bins;

heterozygous and homozygous SNVs were tallied for each bin, and bins with <25 SNVs

were removed. A bin was classified as heterozygous if >50% of the SNVs within the bin

are heterozygous, otherwise it was classified as homozygous. This classification was

used as the HMM emission sequence. The HMM was initialized with the same initiation

and transition probabilities (Prob=10E-8) (Adey et al., 2013), the Viterbi algorithm was

used to estimate a best path, and adjacent homozygous intervals were merged.

10X Genomics linked-read library and sequencing

Marmoset ESC (cj367) genomic DNA was extracted using the MagAttract HMW DNA

Kit from Qiagen (Cat. No. 67563) (Hilden, Germany). The extracted genomic DNA was

verified to be of HMW (average size >30 kb) using using field-inversion gel

electrophoresis on the Pippin Pulse System (Sage Science, Beverly, MA, USA). DNA

was then diluted to 1 ng/µl to be used as input for the 10x Genomics (Pleasanton, CA,

USA) Chromium system (Zheng et al. 2016; Marks et al. 2018) in which HMW DNA

fragments are partitioned into >1 million droplets in emulsion, uniquely barcoded (16 bp)

within each droplet, and subjected to random priming and isothermal amplification

following standard manufacturer's protocol. The barcoded DNA molecules were then

released from the droplets and converted to a linked-read library in which each library

molecule has a 16 bp "HMW fragment barcode". The final linked-read library (8 cycles

of PCR amplification) was quantified using the Kapa Library Quantification Kit (Cat. No.

KK4824) from Kapa Biosystems (Woburn, MA, USA) and diluted to 4 nM. Sequencing

(2×151 bp) was performed on the Illumina NextSeq 500 using the NextSeq 500/550

High Output v2 kit (Cat. No. FC-404-2004) to achieve >30× genomic coverage.

Haplotype phasing using linked-reads

Read-pairs i.e. linked-reads, that come from the same HMW DNA fragment carry the

same "HMW fragment barcode" on Read 1. Virtual long-reads that are representative of

the sequences of the original HMW genomic DNA fragments can be generated by

clustering of linked-reads with identical barcodes. This process allows for the haplotype

phasing of heterozygous SNVs and Indels. Paired-end linked-reads (median insert size

368 bp, duplication rate 5.39%, Q30 Read1 75.6%, Q30 Read2 68.4%) were aligned to

Callithrix jacchus genome (GenBank accession GCA_000004665.1) (alignment rate

87.3%, mean coverage 30.4x, zero coverage 0.938%) and analyzed using the Long

Ranger Software (version 2.1.3) from 10x Genomics (Zheng et al. 2016; Marks et al.

2018) (Pleasanton, CA, USA). Genome indexing was performed prior to alignment

using the Long Ranger *mkref* module. Since Long Ranger *mkref* requires less than 500

contigs, we concatenated contigs in GCA_000004665.1 to meet this requirement.

Reference gaps and repeat regions, downloaded from UCSC Genome Browser

(Karolchik et al. 2004; Kuhn et al. 2013), were excluded from the analysis. Phasing was

performed by specifying the set of pre-determined and filtered heterozygous SNVs and

Indels from using GATK (see above) and formatted using *mkvcf* from Long Ranger

(version 2.1.5). Phasing analysis was performed using the Long Ranger *wgs* module.

Allele-specific CRISPR targets

To identify allele-specific CRISPR targets, we started by extracting variants that

satisfy the following properties from Dataset S2 (phased variants):

1. They passed quality control (VCF field 'Filter' is equal to "PASS")

2. They are phased (VCF field 'GT' uses "|" as separator rather than "/")

3. The alleles are heterozygous (e.g. "0|1" or "1|0", but not "1|1")

4. The difference between the alleles is > 1 bp (to ensure target specificity)

For the 704,571 variants that satisfy these four properties, we extracted the two

haplotype sequences (maximum length: 595). We only worked with the sequences that

were present in the phased genotype. Extracted sequences were tagged them

according to their haplotype, for instance:

• If the sequence in the 'Ref' field was "GTA", the 'Alt' field sequence was "TA",

and phasing was "0|1" i.e. Haplotype_1|Haplotype_2", the sequence containing

"GTA" was tagged "1" for Haplotype 1 and the sequence containing "TA" was

tagged "2" for Haplotype 2.

• If the 'Ref' field was sequence "GTA", the 'Alt' field was "TA,GCTA", phasing was

"1|2", the sequence containing "TA" is tagged "1", the sequence containing

"GCTA" was tagged "2" (and the sequence containing "GTA" was not used)

A regular expression, RegEx, was used to extract all potential CRISPR targets from

these sequences (i.e. all sequences that matched a [G, C, or A]N_{20}GG pattern and

those for which the reverse-complement matched this pattern). This yielded 720,239

candidates, which were then filtered to retain only high-quality targets. The process is

adapted from a selection method previously described and validated (Sunagawa et al.

2016) and has already been used for more than 20 genes (Tatsuki et al. 2016). A high-

quality candidate gRNA needs to target a unique site. All the candidates that have

multiple exact matches in hg19 (irrespective of location) were identified using Bowtie2

(Langmead and Salzberg 2012) and removed. We also removed targets with extreme

GC content (>80% or <20%), and targets that contain TTTT, which tends to break the

gRNA's secondary structure. We also used the Vienna RNA-fold package (Lorenz et al.

2011) to compute the gRNA's secondary structure. We eliminated all candidates for

which the stem loop structure cannot fold correctly for Cas9 recognition (Nishimasu et al.

2014), except if the folding energy was above -18 (indicating that the "incorrectly-folded"

structure is unstable). Finally, we evaluated the off-target risk score using our own

implementation of the Zhang tool (Ran et al. 2013). To ensure that all targets are as

reliable and specific as possible, we used a very strict threshold and rejected

candidates with an off-target risk score less than 75. Candidates that satisfy all these

requirements are considered high quality. For each candidate, we report the location of

the variant (chromosome and position), the haplotype ("1" or "2" from the "Phased

Genotype" column i.e. "Haplotype_1 | Haplotype_2"), the gRNA target sequence, its

position relative to the start of the variant, its orientation, its off-target score, and the

genomic element targeted (gene or enhancer). Note that the position relative to the start

of the variant is for the 5'-most end of the target respective to the genome: if the target

is 5'-3', it is its 5' end; if a target was extracted on the reverse complement, it is its 3'

end.

Data access

All sequencing data are available via NCBI SRA accession SRP141443. Variant data is

available via European Variation Archive accession PRJEB27676.

Acknowledgements

This research was supported by a Tashia and John Morgridge Faculty Scholar Award

from the Stanford Child Health Research Institute (AEU), a the Stanford Neurosciences

Institute Seed Grant (AEU & KJP), the Stanford Child Health Research Institute Grant

(AEU & KJP), and the Stanford Medicine Faculty Innovation Program (AEU). BZ was

funded by NIH grant T32 HL110952. LCL was funded by the Stanford School of

Medicine Dean's Fellowship and NIH grants NS062798, DK090065, and

MH099647. We are thankful for the support of the Stanford CSIF core facility – the

imaging with the DeltaVision OMX system described was supported, in part, by award

1S10OD01227601 from the National Center for Research Resources (NCRR). This

research was partially supported by NIH grants R24OD019803 to TGG and MEE, and

P51OD011106 to the Wisconsin National Primate Research Center. The sequencing

data was generated on an Illumina HiSeq 4000 that was purchased with funds from NIH

award S10OD018220. Lastly, we thank Arineh Khechaduri for performing marmoset

ESC genomic DNA extraction. We also thank Dr. Jeewon S. Kim and Dr. Vittorio

Sebastiano at the Transgenic, Knockout and Tumor Model Center of the Stanford

Cancer Institute for teratoma analyses.

Disclosure Declaration

The authors of this manuscript declare no conflicts of interest.

Author Contributions

B.Z., L.C.L., T.R.W., M.H., and M.J.P. performed experiments. B.Z., S.S.H., and D.P.

performed genomic data analysis. S.C.V., M.E.E., T.G.G., P.M., K.J.P., and A.E.U. provided

materials and reagents. K.J.P. and A.E.U. jointly supervised the study. B.Z. wrote the

manuscript with suggestions from all co-authors.

References

Adey A, Burton JN, Kitzman JO, Hiatt JB, Lewis AP, Martin BK, Qiu R, Lee C, Shendure J. 2013. The haplotype-resolved genome and epigenome of the aneuploid HeLa cancer cell line. *Nature* **500**: 207–211. http://www.nature.com/doifinder/10.1038/nature12064.

Ausderau KK, Dammann C, McManus K, Schneider M, Emborg ME, Schultz-Darken N. 2017. Cross-species comparison of behavioral neurodevelopmental milestones in the common marmoset monkey and human child. *Dev Psychobiol* **59**: 807–821. http://www.ncbi.nlm.nih.gov/pubmed/28763098.

Benirschke K, Anderson JM, Brownhill LE. 1962. Marrow Chimerism in Marmosets. *Science* **138**: 513–5. http://www.ncbi.nlm.nih.gov/pubmed/17753948.

Cai X, Evrony GD, Lehmann HS, Elhosary PC, Mehta BK, Poduri A, Walsh CA. 2014. Single-cell, genome-wide sequencing identifies clonal somatic copy-number variation in the human brain. *Cell Rep* **8**: 1280–9. http://www.ncbi.nlm.nih.gov/pubmed/25159146.

Camus S, Ko WKD, Pioli E, Bezard E. 2015. Why bother using non-human primate models of cognitive disorders in translational research? *Neurobiol Learn Mem* **124**: 123–9. http://www.ncbi.nlm.nih.gov/pubmed/26135120.

Canver MC, Joung JK, Pinello L. 2018. Impact of Genetic Variation on CRISPR-Cas Targeting. *Cris J* **1**: 159–170. http://www.liebertpub.com/doi/10.1089/crispr.2017.0016.

Capitanio JP, Emborg ME. 2008. Contributions of non-human primates to neuroscience research. *Lancet* **371**: 1126–35. http://www.ncbi.nlm.nih.gov/pubmed/18374844.

Chen B, Gilbert LA, Cimini BA, Schnitzbauer J, Zhang W, Li G-W, Park J, Blackburn EH, Weissman JS, Qi LS, et al. 2013. Dynamic imaging of genomic loci in living human cells by an optimized CRISPR/Cas system. *Cell* **155**: 1479–91. http://www.ncbi.nlm.nih.gov/pubmed/24360272.

Chen Y, Zheng Y, Kang Y, Yang W, Niu Y, Guo X, Tu Z, Si C, Wang H, Xing R, et al. 2015. Functional disruption of the dystrophin gene in rhesus monkey using CRISPR/Cas9. *Hum Mol Genet* **24**: 3764–74. http://www.ncbi.nlm.nih.gov/pubmed/25859012.

Debowski K, Drummer C, Lentes J, Cors M, Dressel R, Lingner T, Salinas-Riester G, Fuchs S, Sasaki E, Behr R. 2016. The transcriptomes of novel marmoset monkey embryonic stem cell lines reflect distinct genomic features. *Sci Rep* **6**: 29122. http://www.ncbi.nlm.nih.gov/pubmed/27385131.

DePristo M a, Banks E, Poplin R, Garimella K V, Maguire JR, Hartl C, Philippakis A a, del Angel G, Rivas M a, Hanna M, et al. 2011. A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat Genet* **43**: 491–8.

Emborg ME. 2017. Nonhuman Primate Models of Neurodegenerative Disorders. *ILAR J* **58**: 190–201. http://www.ncbi.nlm.nih.gov/pubmed/28985333.

Farnsworth SL, Qiu Z, Mishra A, Hornsby PJ. 2013. Directed neural differentiation of induced pluripotent stem cells from non-human primates. *Exp Biol Med (Maywood)* **238**: 276–84. http://www.ncbi.nlm.nih.gov/pubmed/23598973.

Grow DA, McCarrey JR, Navara CS. 2016. Advantages of nonhuman primates as preclinical models for evaluating stem cell-based therapies for Parkinson's disease. *Stem Cell Res* **17**: 352–366.

http://linkinghub.elsevier.com/retrieve/pii/S187350611630112X.

International HapMap Consortium. 2005. A haplotype map of the human genome. *Nature* **437**: 1299–320. http://www.ncbi.nlm.nih.gov/pubmed/16255080.

Izpisua-Belmonte JC, Callaway EM, Caddick SJ, Churchland P, Feng G, Homanics GE, Lee K-F, Leopold DA, Miller CT, Mitchell JF, et al. 2015. Brains, genes, and primates. *Neuron* **86**: 617–31. http://www.ncbi.nlm.nih.gov/pubmed/25950631.

Kang Y, Zheng B, Shen B, Chen Y, Wang L, Wang J, Niu Y, Cui Y, Zhou J, Wang H, et al. 2015. CRISPR/Cas9-mediated Dax1 knockout in the monkey recapitulates human AHC-HH. *Hum Mol Genet* **24**: 7255–7264. https://academic.oup.com/hmg/article-lookup/doi/10.1093/hmg/ddv425.

Karolchik D, Hinrichs AS, Furey TS, Roskin KM, Sugnet CW, Haussler D, Kent WJ. 2004. The UCSC Table Browser data retrieval tool. *Nucleic Acids Res* **32**: D493-6. http://www.ncbi.nlm.nih.gov/pubmed/14681465.

Kishi N, Sato K, Sasaki E, Okano H. 2014. Common marmoset as a new model animal for neuroscience research and genome editing technology. *Dev Growth Differ* **56**: 53–62. http://www.ncbi.nlm.nih.gov/pubmed/24387631.

Knouse KA, Wu J, Amon A. 2016. Assessment of megabase-scale somatic copy number variation using single-cell sequencing. *Genome Res* **26**: 376–84. http://www.ncbi.nlm.nih.gov/pubmed/26772196.

Kropp J, Di Marzo A, Golos T. 2017. Assisted reproductive technologies in the common marmoset: an integral species for developing nonhuman primate models of human diseases†. *Biol Reprod* **96**: 277–287. https://academic.oup.com/biolreprod/article/96/2/277/2948764.

Kuhn RM, Haussler D, Kent WJ. 2013. The UCSC genome browser and associated

tools. *Brief Bioinform* **14**: 144–161. https://academic.oup.com/bib/article-

lookup/doi/10.1093/bib/bbs038.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat*

Methods **9**: 357–9. http://www.ncbi.nlm.nih.gov/pubmed/22388286.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics* **25**: 1754–60.

http://www.ncbi.nlm.nih.gov/pubmed/19451168.

Lorenz R, Bernhart SH, Höner Zu Siederdissen C, Tafer H, Flamm C, Stadler PF,

Hofacker IL. 2011. ViennaRNA Package 2.0. *Algorithms Mol Biol* **6**: 26.

http://www.ncbi.nlm.nih.gov/pubmed/22115189.

Luo X, Li M, Su B. 2016. Application of the genome editing tool CRISPR/Cas9 in non-

human primates. *Zool Res* **37**: 214–9.

http://www.ncbi.nlm.nih.gov/pubmed/27469252.

Mansfield K. 2003. Marmoset models commonly used in biomedical research. *Comp*

Med **53**: 383–92. http://www.ncbi.nlm.nih.gov/pubmed/14524414.

Marks P, Garcia S, Barrio AM, Belhocine K, Bernate J, Bharadwaj R, Bjornson K,

Catalanotti C, Delaney J, Fehr A, et al. 2018. Resolving the Full Spectrum of

Human Genome Variation using Linked-Reads. *bioRxiv* 230946.

https://www.biorxiv.org/content/early/2018/01/09/230946.

Marmoset Genome Sequencing and Analysis Consortium. 2014. The common

marmoset genome provides insight into primate biology and evolution. *Nat Genet*

46: 850–7. https://www.nature.com/articles/ng.3042.

Maudhoo MD, Ren D, Gradnigo JS, Gibbs RM, Lubker AC, Moriyama EN, French JA,

Norgren RB. 2014. De novo assembly of the common marmoset transcriptome

from NextGen mRNA sequences. *Gigascience* **3**: 14.

https://academic.oup.com/gigascience/article-lookup/doi/10.1186/2047-217X-3-14.

McConnell MJ, Lindberg MR, Brennand KJ, Piper JC, Voet T, Cowing-Zitron C,

Shumilina S, Lasken RS, Vermeesch JR, Hall IM, et al. 2013. Mosaic copy number

variation in human neurons. *Science* **342**: 632–7.

http://www.ncbi.nlm.nih.gov/pubmed/24179226.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K,

Altshuler D, Gabriel S, Daly M, et al. 2010. The Genome Analysis Toolkit: a

MapReduce framework for analyzing next-generation DNA sequencing data.

Genome Res **20**: 1297–303. http://www.ncbi.nlm.nih.gov/pubmed/20644199.

Nelson EE, Winslow JT. 2009. Non-human primates: model animals for developmental

psychopathology. *Neuropsychopharmacology* **34**: 90–105.

http://www.ncbi.nlm.nih.gov/pubmed/18800061.

Nishimasu H, Ran FA, Hsu PD, Konermann S, Shehata SI, Dohmae N, Ishitani R,

Zhang F, Nureki O. 2014. Crystal structure of Cas9 in complex with guide RNA and

target DNA. *Cell* **156**: 935–49. http://www.ncbi.nlm.nih.gov/pubmed/24529477.

Niu Y, Shen B, Cui Y, Chen Y, Wang J, Wang L, Kang Y, Zhao X, Si W, Li W, et al.

2014. Generation of gene-modified cynomolgus monkey via Cas9/RNA-mediated

gene targeting in one-cell embryos. *Cell* **156**: 836–43.

http://www.ncbi.nlm.nih.gov/pubmed/24486104.

Okano H, Hikishima K, Iriki A, Sasaki E. 2012. The common marmoset as a novel

animal model system for biomedical and neuroscience research applications. *Semin Fetal Neonatal Med* **17**: 336–40. http://www.ncbi.nlm.nih.gov/pubmed/22871417.

Park JE, Zhang XF, Choi S-H, Okahara J, Sasaki E, Silva AC. 2016. Generation of transgenic marmosets expressing genetically encoded calcium indicators. *Sci Rep* **6**: 34931. http://www.ncbi.nlm.nih.gov/pubmed/27725685.

Parker KJ, Garner JP, Oztan O, Tarara ER, Li J, Sclafani V, Del Rosso LA, Chun K, Berquist SW, Chez MG, et al. 2018. Arginine vasopressin in cerebrospinal fluid is a marker of sociality in nonhuman primates. *Sci Transl Med* **10**. http://www.ncbi.nlm.nih.gov/pubmed/29720452.

Phillips KA, Bales KL, Capitanio JP, Conley A, Czoty PW, 't Hart BA, Hopkins WD, Hu S-L, Miller LA, Nader MA, et al. 2014. Why primate models matter. *Am J Primatol* **76**: 801–27. http://www.ncbi.nlm.nih.gov/pubmed/24723482.

Pinello L, Canver MC, Hoban MD, Orkin SH, Kohn DB, Bauer DE, Yuan G-C. 2016. Analyzing CRISPR genome-editing experiments with CRISPResso. *Nat Biotechnol* **34**: 695–697. http://www.nature.com/doifinder/10.1038/nbt.3583.

Qiu Z, Mishra A, Li M, Farnsworth SL, Guerra B, Lanford RE, Hornsby PJ. 2015. Marmoset induced pluripotent stem cells: Robust neural differentiation following pretreatment with dimethyl sulfoxide. *Stem Cell Res* **15**: 141–50. http://www.ncbi.nlm.nih.gov/pubmed/26070112.

Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F. 2013. Genome engineering using the CRISPR-Cas9 system. *Nat Protoc* **8**: 2281–2308. http://www.nature.com/doifinder/10.1038/nprot.2013.143.

Sasaki E. 2015. Prospects for genetically modified non-human primate models,

including the common marmoset. *Neurosci Res* **93**: 110–5.

http://www.ncbi.nlm.nih.gov/pubmed/25683291.

Sasaki E, Hanazawa K, Kurita R, Akatsuka A, Yoshizaki T, Ishii H, Tanioka Y, Ohnishi Y,

Suemizu H, Sugawara A, et al. 2005. Establishment of novel embryonic stem cell

lines derived from the common marmoset (Callithrix jacchus). *Stem Cells* **23**: 1304–

13. http://www.ncbi.nlm.nih.gov/pubmed/16109758.

Sasaki E, Suemizu H, Shimada A, Hanazawa K, Oiwa R, Kamioka M, Tomioka I,

Sotomaru Y, Hirakawa R, Eto T, et al. 2009. Generation of transgenic non-human

primates with germline transmission. *Nature* **459**: 523–7.

http://www.ncbi.nlm.nih.gov/pubmed/19478777.

Sato K, Kuroki Y, Kumita W, Fujiyama A, Toyoda A, Kawai J, Iriki A, Sasaki E, Okano H,

Sakakibara Y. 2015. Resequencing of the common marmoset genome improves

genome assemblies and gene-coding sequence analysis. *Sci Rep* **5**: 16894.

http://www.readcube.com/articles/10.1038/srep16894 (Accessed April 22, 2016).

Sato K, Oiwa R, Kumita W, Henry R, Sakuma T, Ito R, Nozu R, Inoue T, Katano I, Sato

K, et al. 2016. Generation of a Nonhuman Primate Model of Severe Combined

Immunodeficiency Using Highly Efficient Genome Editing. *Cell Stem Cell* **19**: 127–

38. http://linkinghub.elsevier.com/retrieve/pii/S1934590916301539.

Sato K, Sasaki E. 2018. Genetic engineering in nonhuman primates for human disease

modeling. *J Hum Genet* **63**: 125–131.

http://www.ncbi.nlm.nih.gov/pubmed/29203824.

Schultz-Darken N, Braun KM, Emborg ME. 2016. Neurobehavioral development of

common marmoset monkeys. *Dev Psychobiol* **58**: 141–58.

http://www.ncbi.nlm.nih.gov/pubmed/26502294.

Shiozawa S, Kawai K, Okada Y, Tomioka I, Maeda T, Kanda A, Shinohara H, Suemizu H, James Okano H, Sotomaru Y, et al. 2011. Gene targeting and subsequent site-specific transgenesis at the β-actin (ACTB) locus in common marmoset embryonic stem cells. *Stem Cells Dev* **20**: 1587–99.

http://www.ncbi.nlm.nih.gov/pubmed/21126169.

Sunagawa GA, Sumiyama K, Ukai-Tadenuma M, Perrin D, Fujishima H, Ukai H, Nishimura O, Shi S, Ohno R-I, Narumi R, et al. 2016. Mammalian Reverse Genetics without Crossing Reveals Nr3a as a Short-Sleeper Gene. *Cell Rep* **14**: 662–677. http://www.ncbi.nlm.nih.gov/pubmed/26774482.

Tatsuki F, Sunagawa GA, Shi S, Susaki EA, Yukinaga H, Perrin D, Sumiyama K, Ukai-Tadenuma M, Fujishima H, Ohno R, et al. 2016. Involvement of Ca(2+)-Dependent Hyperpolarization in Sleep Duration in Mammals. *Neuron* **90**: 70–85.

http://www.ncbi.nlm.nih.gov/pubmed/26996081.

The 1000 Genomes Project Consortium, Abecasis GR, Altshuler D, Auton A, Brooks LD, Durbin RM, Gibbs RA, Hurles ME, McVean GA. 2010. A map of human genome variation from population-scale sequencing. *Nature* **467**: 1061–73.

http://www.nature.com/doifinder/10.1038/nature09534.

Thomson JA, Kalishman J, Golos TG, Durning M, Harris CP, Hearn JP. 1996. Pluripotent cell lines derived from common marmoset (Callithrix jacchus) blastocysts. *Biol Reprod* **55**: 254–9. http://www.ncbi.nlm.nih.gov/pubmed/8828827.

Tomioka I, Ishibashi H, Minakawa EN, Motohashi HH, Takayama O, Saito Y, Popiel HA,

Puentes S, Owari K, Nakatani T, et al. 2017. Transgenic Monkey Model of the

Polyglutamine Diseases Recapitulating Progressive Neurological Symptoms.

eneuro **4**: ENEURO.0250-16.2017.

http://eneuro.sfn.org/lookup/doi/10.1523/ENEURO.0250-16.2017.

Tomioka I, Maeda T, Shimada H, Kawai K, Okada Y, Igarashi H, Oiwa R, Iwasaki T,

Aoki M, Kimura T, et al. 2010. Generating induced pluripotent stem cells from

common marmoset (Callithrix jacchus) fetal liver cells using defined factors,

including Lin28. *Genes to Cells* **15**: 959–969. http://doi.wiley.com/10.1111/j.1365-

2443.2010.01437.x.

Vermilyea SC, Babinski A, Guthrie S, Golos TG, Emborg ME. 2018. LRRK2 Genomic

Editing in Common Marmoset Stem Cells. *Cell Transplant* **27**: 682–722.

https://doi.org/10.1177/0963689718765742.

Vermilyea SC, Guthrie S, Meyer M, Smuga-Otto K, Braun K, Howden S, Thomson JA,

Zhang S-C, Emborg ME, Golos TG. 2017. Induced Pluripotent Stem Cell-Derived

Dopaminergic Neurons from Adult Common Marmoset Fibroblasts. *Stem Cells Dev*

26: 1225–1235. http://www.ncbi.nlm.nih.gov/pubmed/28635509.

Wise SP. 2008. Forward frontal fields: phylogeny and fundamental function. *Trends*

Neurosci **31**: 599–608.

http://linkinghub.elsevier.com/retrieve/pii/S0166223608002075.

Zahs KR, Ashe KH. 2010. "Too much good news" - are Alzheimer mouse models trying

to tell us how to prevent, not cure, Alzheimer's disease? *Trends Neurosci* **33**: 381–

9. http://www.ncbi.nlm.nih.gov/pubmed/20542579.

Zhang Y, Pak C, Han Y, Ahlenius H, Zhang Z, Chanda S, Marro S, Patzke C, Acuna C,

Covy J, et al. 2013. Rapid single-step induction of functional neurons from human pluripotent stem cells. *Neuron* **78**: 785–98. http://www.ncbi.nlm.nih.gov/pubmed/23764284.

Zheng GXY, Lau BT, Schnall-Levin M, Jarosz M, Bell JM, Hindson CM, Kyriazopoulou-Panagiotopoulou S, Masquelier DA, Merrill L, Terry JM, et al. 2016. Haplotyping germline and cancer genomes with high-throughput linked-read sequencing. *Nat Biotechnol* **34**: 303–11. http://www.ncbi.nlm.nih.gov/pubmed/26829319 (Accessed July 14, 2016).

Zhou B, Ho SS, Zhu X, Zhang X, Spies N, Byeon S, Arthur JG, Pattni R, Ben-Efraim N, Haney MS, et al. 2018. Comprehensive, integrated and phased whole-genome analysis of the primary ENCODE cell line K562. *bioRxiv*. http://biorxiv.org/content/early/2018/01/22/192344.abstract.

FIGURE AND TABLE LEGENDS

Figure 1. (a) cj367 ESC colony at 20× magnification. Days 5 (b) and 9 (c) respectively of cj367 directly induced into immature cortical glutamatergic neurons; white bar = 400 microns. Cells are marked by EGFP. Immunostaining of pluripotency markers OCT3/4 (d), NANOG (e), and SSEA4 (f) in cj367; white bar = 200 microns.

Figure 2. (a) Circos visualization of cj367 genome variants with the following tracks: 1. phased haplotype blocks (demarcated with 4 colors for clearer visualization); 2. SNV density in 1 Mb windows; 3. Indel density in 1 Mb windows; 4. zygosity (heterozygous or homozygous > 50%) in 1 Mb windows; regions with continuous homozygosity. (b) Violin plot of size distribution of haplotype blocks in cj367.

Figure 3. (a-c) Three examples of marmoset ESCs transfected with dCas9-GFP that are targeted telomeric sequences revealing telomeres as fluorescent puncta (white arrowheads). Images are maximum projections of the nuclear volume and the dashed outlines indicate the nuclear boundaries. Scale bar, 5 μm.

Figure 4. Histological analysis of teratomas derived from marmoset ESCs after dCas9-GFP transfection. Scale bar is 50 μm. Mamorset lines are characterized by pluripotency as demonstrated by their ability to form glandular epithelium (endoderm in a); cartilage (mesoderm in b), and neural rosettes-like structures (ectoderm in c).

Figure S1. Violin plot of size distribution of haplotype blocks in cj367 by chromosomes

Table 1. Summary of cj367 SNVs and Indels

Table S1. cj367 genomic regions exhibiting continuous homozygosity.

Table S2. List of CRISPR sites suitable for allele-specific targeting in cj367.

(a) cj367 ESC colony   (b) iN day 5   (c) iN day 9

(d) OCT3/4   (e) NANOG   (f) SSEA4

bar = 400 um (b, c); bar = 200 um (d-f).

a

Track 1
Phase Blocks

Colors alternate, indicating consecutive phase blocks

Track 2
SNP Density

0   4700

Track 3
Indel Density

0   750

Track 4
Zygosity

Heterozygous
Homozygous
Continuous
Homozygosity

cj367

b

Size Distribution for Marmoset Phased Haplotype Blocks

Length (bp)

10,000,000

100,000

1,000

Marmoset

All Marmoset Phase Blocks