Fast and flexible linear mixed models for genome-wide genetics

Daniel E Runcie[1]* and Lorin Crawford[2]

1 Department of Plant Sciences, University of California Davis, Davis, CA, USA
2 Department of Biostatistics, Brown University, Providence, RI, USA

* deruncie@ucdavis.edu

Abstract

Linear mixed effect models are powerful tools used to account for population structure in genome-wide association studies (GWASs) and estimate the genetic architecture of complex traits. However, fully-specified models are computationally demanding and common simplifications often lead to reduced power or biased inference. We describe `Grid-LMM` (https://github.com/deruncie/GridLMM), an extendable algorithm for repeatedly fitting complex linear models that account for multiple sources of heterogeneity, such as additive and non-additive genetic variance, spatial heterogeneity, and genotype-environment interactions. `Grid-LMM` can compute approximate (yet highly accurate) frequentist test statistics or Bayesian posterior summaries at a genome-wide scale in a fraction of the time compared to existing general-purpose methods. We apply `Grid-LMM` to two types of quantitative genetic analyses. The first is focused on accounting for spatial variability and non-additive genetic variance while scanning for QTL; and the second aims to identify gene expression traits affected by non-additive genetic variation. In both cases, modeling multiple sources of heterogeneity leads to new discoveries.

Author summary

The goal of quantitative genetics is to characterize the relationship between genetic variation and variation in quantitative traits such as height, productivity, or disease susceptibility. A statistical method known as the linear mixed effect model has been critical to the development of quantitative genetics. First applied to animal breeding, this model now forms the basis of a wide-range of modern genomic analyses including genome-wide associations, polygenic modeling, and genomic prediction. The same model is also widely used in ecology, evolutionary genetics, social sciences, and many other fields. Mixed models are frequently multi-faceted, which is necessary for accurately modeling data that is generated from complex experimental designs. However, most genomic applications use only the simplest form of linear mixed methods because the computational demands for model fitting can be too great. We develop a flexible approach for fitting linear mixed models to genome scale data that greatly reduces their computational burden and provides flexibility for users to choose the best statistical paradigm for their data analysis. We demonstrate improved accuracy for genetic association tests, increased power to discover causal genetic variants, and the ability to provide accurate summaries of model uncertainty using both simulated and real data examples.

Introduction

Population stratification, genetic relatedness, ascertainment, and other sources of heterogeneity lead to spurious signals and reduced power in genetic association studies [1–5]. When not properly taken into account, non-additive genetic effects and environmental variation can also bias estimates of heritability, polygenic adaptation, and genetic values in breeding programs [5–8]. Both issues are caused by departures from a key assumption underlying linear models that observations are independent. Non-independent samples lead to a form of pseudo-replication, effectively reducing the true sample size. Linear mixed effect models (LMMs) are widely used to account for non-independent samples in quantitative genetics [9]. The flexibility and interpretability of LMMs make them a dominant statistical tool in much of biological research [9–18].

Random effect terms are used in LMMs to account for specific correlations among observations. Fitting an LMM requires estimating the importance of each random effect, called its variance component. General-purpose tools for this are too slow to be practical for genome-scale datasets with thousands of observations and millions of genetic markers [19]. This lack of scalability is caused primarily by two factors: (i) closed-form solutions of maximum-likelihood (ML or REML) or posterior estimates of the variance components are not available and numerical optimization routines require repeatedly evaluating the likelihood function many times, and (ii) each evaluation of the likelihood requires inverting the covariance matrix of random effects, an operation that scales cubically with the number of observations. Repeating this whole process millions of times quickly becomes infeasible.

To this end, several specialized approaches have been developed to improve the speed of LMMs, including the use of sparse matrix operations [20, 21], spectral decomposition of the random effect covariance matrix [22–26], and Monte Carlo REML [27]. These algorithms are particularly useful when the same random effect structure is used many times. For example, in genome-wide association studies (GWAS), each marker is tested with the same LMM. Similarly, in population-level transcriptomics, eQTLs or variance components are estimated for each of tens-of-thousands of genes expression traits. Fast and exact algorithms for fitting LMMs are limited to the case of only a single (full-rank) random effect term, besides the residuals [22–24]. Recently, approximate learning algorithms have been developed for the scalable extension to multiple random effects [28, 29], but few of these ensure guarantees in terms of estimation accuracy. One strategy applicable to studies with multiple random effects is to estimate the variance components only once, in a model without additional marker effects, and then test each marker either using a score test [30] (which does not produce an effect size estimate), or with a conditional F-test assuming the variance component estimates are fixed [31–34]. Given the "known" variance components, closed-form solutions of all other parameters of an LMM can be found using a rotated version of the simple linear model. Unfortunately, both approximations suffer from reduced power when marker effects are large, intractable posterior inference in a Bayesian analysis, and the inability to be applied to parallel analyses over many traits (like gene expression). Table 1 summarizes these different methods, details their respective computational complexities, and provides relevant references.

`Grid-LMM` takes a different approach to fitting LMMs: rather than directly optimizing the variance components separately for each test, we define a grid spanning all valid values of the variance components and fit simple linear models at each grid location. Each evaluation involves a single Cholesky decomposition of the random effect covariance matrix, which is then reused to calculate closed-form ML solutions or Bayesian posterior summaries (under a specific conjugate prior; see S1 Supplementary Methods) for all separate tests. This results in dramatic time-savings in typical GWAS settings (see S1 Fig). After repeating these calculations across the whole grid, we select

Table 1. Performance and limitations of reference models for linear mixed model GWAS. The time complexity of each algorithm is approximate, assuming a model with only a single marker effect and no other fixed effects. Here, l is used to index full-rank random effects; σ_l^2 are variance component parameters; n is the number of observations; and p is number of markers to test. Denote $t_1 \ldots t_7$ to represent the number of iterations needed for convergence, which is expected to vary among methods (particularly for the iterations of grid search in `Grid-LMM-fast`), and may vary across markers. The terms g and g_i are grid sizes for the `Grid-LMM` methods (i.e. the number of grid vertices that must be evaluated). Lastly, p_i is the number of markers that need to be tested in iteration of $i \in \{1 \ldots t_7\}$ of the `Grid-LMM-fast` method. The rate limiting terms in common GWAS applications (where $p \gg n$) are in bold. "Method Type" describes the estimation of σ_l^2. "Exact" means effectively exact, up to machine precision and subject to possible convergence of algorithms to local maxima. "Null" means estimation of parameters under the null model with no marker effects. References list additional methods that are approximately equivalent to the given model classes.

Tool	Max σ_l^2	Method Type	Approx. Time Complexity	References
BOLT-LMM	1	exact	$\mathcal{O}(\boldsymbol{t_1 pn})$	[27]
GEMMA	1	exact	$\mathcal{O}(n^3 + \boldsymbol{pn^2} + pt_2 n)$	[23, 24]
EMMAX	1	null	$\mathcal{O}(n^3 + t_3 n + \boldsymbol{pn^2})$	[31, 32]
LDAK	>1	exact	$\mathcal{O}(\boldsymbol{pt_4 n^3})$	[7, 21, 35, 36]
pylmm	>1	null	$\mathcal{O}(t_5 n^3 + \boldsymbol{pn^2})$	[7, 34, 36, 37]
Grid-LMM	$\sim$ 1-8	grid	$\mathcal{O}(\boldsymbol{g}(n^3 + \boldsymbol{pn^2}))$	Present Work
Grid-LMM-fast	$\sim$ 1-8	grid-sampling	$\mathcal{O}(t_6 n^3 + \boldsymbol{pn^2} + \sum_{i=1}^{t_7} \boldsymbol{g_i}(n^3 + \boldsymbol{p_i n^2}))$	Present Work

the highest ML (or REML) score for each marker to compute approximate likelihood ratio and Wald tests [32, 38], or analogously derive posterior distributions and Bayes factors by summing appropriate statistics across the grid. The `Grid-LMM` approach relies on a re-parameterization of the typical LMM framework from individual variance components σ_l^2 to variance component proportions $h_l^2 = \sigma_l^2/\sigma^2$, where σ^2 without the subscript denotes the total sum of all variance components (including the residual). Since the variance components must be non-negative, their proportions are restricted to the unit interval $[0, 1]$ and sum to 1, forming a simplex. Therefore, a finite grid can span all valid parameter values. While the size of the grid increases rapidly with the number of random effects, for a small number of random effects ($\sim$1-8) and a moderate grid resolution, the size of the grid remains tiny relative to the number of models in a typical GWAS. As we show below, highly accurate test statistics are achieved even with a coarse grid in most reasonable situations, and further improvements to efficiency are possible by using heuristics to adaptively sample the grid or reduce the number of grid locations computed for the majority of tests. This strategy of conditioning on variance components over a grid and then combining solutions can be applied to many other tools in quantitative genetics including set tests for rare variants [39, 40], whole-genome regression models such as LASSO and elastic net [41, 42], and QTL mapping in controlled crosses [43].

In the following sections, we demonstrate the accuracy and advantages of the `Grid-LMM` approach using a simulation study and two real genome-wide quantitative genetics examples. The first is for GWAS, where tens-to-hundreds of thousands of markers are individually tested for associations with a single phenotype. The second is for gene expression, where thousands of traits are each tested for non-additive genetic variance. In both cases, the same random effect structure is used for each model. While approximate, the test-statistics and posterior quantities calculated by `Grid-LMM` are accurate and improve power relative to other approximation methods — all while maintaining dramatically reduced computational burdens than the direct approach. Full derivations of the model and useful heuristics are described in detail in the Methods.

Results

As a first case-study, we used `Grid-LMM` to perform two types of genome-wide association studies (GWAS) that benefit from modeling multiple random effects: (1) the study of gene-environment interactions, and (2) detecting associations for phenotypes driven by non-additive genetic variation or spatial variation. In both cases, there are multiple sources of covariation among observations that can inflate test statistics and bias estimates of heritability if not appropriately accounted for by using mixed models.

As an example of a GWAS with gene-environment interactions, we analyzed data on flowering times for *Arabidopsis thaliana* [44]. First, we benchmarked results from standard LMM methodologies to confirm that `Grid-LMM` association tests are accurate. We ran `Grid-LMM` on both a fine-grained grid with a small step size of 0.01 h^2-units, and a larger step size of 0.1 h^2-units, to test for associations between 216,130 single nucleotide polymorphisms (SNPs) and flowering times of 194 accessions measured at 10C (i.e. in a constant environment). We compared the Wald-test p-values computed by both `Grid-LMM` models to p-values computed using the exact method `GEMMA` [24], and the approximate method `EMMAX` [32]. Each method was applied as an LMM with only the additive relationship matrix as its single random effect. Grid-LMM p-values computed using the finer grid size (i.e. 0.01 h^2-units) were almost identical to those of `GEMMA`, and even p-values computed with the larger step size (i.e. 0.1 h^2-units) were more accurate than those resulting from `EMMAX`. There were particularly noticeable differences in performance for markers with larger scores, which were strongly underestimated by `EMMAX` since its approximations of h^2 are made strictly under the null model (Fig 1a). This pattern held true across all of the available 107 *Arabidopsis thaliana* phenotypes (see S2 Fig). However, despite these results, we do not advocate `Grid-LMM` in this setting; GEMMA (and similar methods) provides an exact test and is more computationally efficient. The real advantage of `Grid-LMM` is its ability to effectively model two (or more) random effects.

To demonstrate this advantage, we tested for gene-environment (G×E) interaction effects on flowering time. Here, we combined flowering time data from two conditions: constant 10C (as described above) and in the field. We limited our analysis to the 175 accessions that were grown under both conditions, yielding a total of $n = 350$ observations. When observations come from different environments, we might expect phenotypes to cluster for at least two reasons: (i) the sharing of alleles with constant effects across environments due to relatedness and population structure, commonly modeled with the additive genomic relationship matrix (**A**) as a random effect, and (ii) the sharing of alleles with effects that differ among environments (or are specific to one environment). Previous work has shown that, when testing for G×E effects with GWAS, modeling the second source of covariance by using a second random effect can prevent spurious signals and increase power [34]. The same result is replicated in this setting using simulations (see S3 Fig and S4 Fig).

Here, we calculated G×E p-values for each genetic marker using `Grid-LMM` (grid step size = 0.01 h^2 units) on the full dataset using both the **A** and the G×E relationship matrices as two random effects. These are compared to p-values from (i) an LMM which ignores the G×E covariance and only considers genetic similarity, and (ii) a model similar to `pylmm` [34] which does consider both random effects but estimates variances components from the null model (this is referred to as `null-LMM-G×E` below). For each model, we included an additional random effect to account for the repetition of accessions which could induce additional covariance among observations. In this dataset, a simpler approach to testing for G×E was also available: test for associations between markers and the *plasticity* (i.e. the difference in flowering time between the field and 10C) of each accession, which requires only a single random effect and can be fit with `GEMMA`. This simple approach is only possible because every genotype was measured in

both environments. Nonetheless, it is expected to produce identical tests to the full three-random-effect model and, therefore, serves as a viable benchmark for the `Grid-LMM` results. We used the plasticity GWAS approach as a baseline to compare the three models that use the raw data directly (Fig 1b). As expected, ignoring the G×E covariance leads to greatly inflated tests for the majority of markers. `Grid-LMM` replicated `GEMMA`'s plasticity p-values almost exactly when run with three random effects; alternatively, a portion of the `null-LMM-G×E`'s tests were deflated, implying lower power. The full analysis using `Grid-LMM` with three random effects took 26 minutes. Fitting the same model for all 216,130 markers directly using an exact method (e.g. `LDAK` [7]) would take approximately 6 hours (about 14× longer). Note that `LDAK` is not designed for re-estimating variance components for each SNP in multiple-random-effect models and so, to conduct time comparisons, we simply ran the program multiple times. This requires `LDAK` to re-load the covariance matrices for each marker. However, by controlling the maximum number of iterations in the `LDAK` optimization engine, we estimate that a maximum $\approx 33\%$ of the running time for these models is due to data import, with the remainder being due to the numerical calculations.

Fig 1. Comparisons of GWAS results for *Arabidopsis* flowering time. (a) Results for days-to-flower (DTF) at 10C. Compared are `EMMAX` and `Grid-LMM` with grid sizes 0.1 and 0.01, respectively. The exact method `GEMMA` is treated as a baseline. Each method was applied to the same LMM with only the additive relationship matrix as a random effect, and p-values were computed using the Wald test. (**b**) GWAS for the gene-environment (G×E) interactions effects on DTF between the constant 10C and field conditions. `GEMMA` results are given for the plasticity of each accession (i.e. the difference DTF_{Field} - DTF_{10C}), and fit with a single random effect of the additive relationship matrix. The other three methods consider the full data with two observations per accession. Namely, `Grid-LMM-A` fit a "standard" GWAS with a grid step size 0.01, but with two random effects — the additive relationship matrix and an *iid* line effect. The other two models, `null-LMM-G×E` and `Grid-LMM-G×E`, fit three random effects — the additive relationship matrix, an *iid* line effect, and a G×E-additive relationship matrix representing the background covariation in gene-environment interactions among accessions.

Even when all individuals are measured in the same environment, the typical additive relationship matrix may not account for all sources of covariation. In particular, spatial variation and the sharing of non-additive alleles may also induce covariance, lead to reduced power, and/or result in inflated tests in GWAS [45]. As an illustrative example, we tested for genetic association between 10,075 bi-allelic autosomal markers and body mass among 1,814 heterogenous stock mice from the Wellcome Trust Centre

for Human Genetics (WTCHG) [46]. We first computed additive and pairwise-epistatic ₁₅₇
relationship matrices, as well as a spatial-environmental covariance matrix based on the ₁₅₈
523 different cages used in the experiment. Next, we compared p-values derived from an ₁₅₉
LMM considering only the additive relationship matrix (as in a typical GWAS) to those ₁₆₀
calculated by `Grid-LMM` using all three relationship matrices (Fig 2a-d). Using the ₁₆₁
three-random effect model, we identified associations on two chromosomes (numbers 4 ₁₆₂
and 11) that were not apparent when just using the additive relationship matrix as a ₁₆₃
random effect. Both loci have been previously identified as size-associated QTL (see ₁₆₄
Table S1 Table) [47,48]. Genomic control values for the two models were both close to ₁₆₅
one (A-only = 0.975, A+E+Cage = 0.974) (Fig 2b,d). The three-random effect model ₁₆₆
took 8.5 minutes to fit using `Grid-LMM`, while a full analysis on all 10,346 markers ₁₆₇
would have taken approximately 10 hours with `LDAK` (more than 100× longer, of which ₁₆₈
we estimate a maximum of ≈ 10% is spent on reading in data). Extrapolating to a ₁₆₉
consortium sized genome-wide analysis with 1 million markers would take ≈ 14 hours ₁₇₀
using `Grid-LMM`, as opposed to 40 days using `LDAK`. We see larger performance gains in ₁₇₁
the mouse dataset compared to the *Arabidopsis* dataset because of the larger sample ₁₇₂
size (n =1,814 vs. 350). `LDAK` (and other exact general-purpose REML methods) is ₁₇₃
dominated by matrix inversions which scale with n^3, while `Grid-LMM` is dominated by ₁₇₄
matrix-vector multiplications which scale with n^2 (again see Table 1). The performance ₁₇₅
advantage of `Grid-LMM` will increase even more for datasets with more individuals. ₁₇₆

**Fig 2. GWAS results for the body weight trait of the Wellcome Trust
heterogenous stock mice**. `Grid-LMM` was run using a grid size of 0.01 h^2-units for
each model. (**a-b**) Wald test p-values from a model with only the additive relationship
matrix (**A**) as a random effect, organized by chromosome and compared to a uniform
distribution via a QQ-plot. (**c-d**) Wald test p-values from a model with three random
effects: the additive, epistatic (**E**), and spatial (**Cage**) relationship matrices. (**e**)
Approximate Bayes factors calculated from the three random effect model, assuming a
standard normal N(0, 1) prior for the marker effects and a uniform prior on the grid of
variance component proportions.

To demonstrate the flexibility of `Grid-LMM` for GWAS, we also ran an analysis on ₁₇₇
the mice body weight trait using Bayesian inference and calculated Bayes Factors by ₁₇₈
comparing the models for each marker to the null model assuming no genetic effects. ₁₇₉
Here, we used a uniform prior over the grid of variance component proportions and ₁₈₀
assumed a standard normal prior for the marker effect sizes. In this setting, the Bayes ₁₈₁
Factors were highly correlated with the frequentist p-values — they also highlight the ₁₈₂

same associations on chromosomes 4 and 11 (Fig 2e). In general, Bayes Factors provide additional flexibility for incorporating prior information on individual markers or even combining results among multiple studies [49,50]. The full Bayesian analysis for `Grid-LMM` took 15.5 minutes, just 7 minutes longer than the frequentist analysis, making it practical for genome-wide studies.

As a third case-study, we used `Grid-LMM` to estimate the additive and pairwise-epistatic variance components for 20,178 gene expression traits measured on 681 *Arabidopsis* accessions from the 1001 Genomes Project [51]. Using a grid-size of 0.05 h^2 units, we estimated the magnitude of each variance component by REML (Fig 3a). The whole analysis took ≈ 6 minutes. Finding REML solutions for the same two-random effect model, on each of the traits separately, using the exact method `LDAK` took ≈ 90 minutes (of which $\approx 20\%$ was due to data import). Grid-LMM variance component estimates replicated those of `LDAK` accurately, but with less precision due to the coarse grid size (see S5 Fig). Notably, for many genes, the proportion of variance in expression attributed to additive variance dropped considerably when the epistatic variance was also modeled (see S6 Fig). Therefore, including multiple random effects can have significant impact on downstream conclusions for many traits.

Even with this relatively large sample size for a population-level gene expression dataset, considerable uncertainty remains in the estimated variance components. The point-estimates calculated by REML do not capture this uncertainty and estimating confidence intervals for the variance components using REML is difficult. The full `Grid-LMM` approach can be used to calculate posterior distributions for each variance component with little additional cost — note that MCMC sampling is not needed because a reasonably-sized grid can span all valid values of each variance component proportion (see Methods). Using this approach, we identify 8,585 genes with a posterior probability of non-zero epistatic variance greater than 90%, and 28 more genes with a posterior mean epistatic variance component greater than 80%. For two example genes, we show that the fitted posterior distributions are similar to those estimated via MCMC using `rstan` [52,53] (see Fig 3b-c). The `rstan` analyses took ≈ 20 hours per gene to generate an effective sample size of $\approx 200 - 400$ for the variance component parameters. Therefore, posterior inference by MCMC for all 20,178 genes would take about 50 years of computational time.

Now that we have demonstrated in real data examples that `Grid-LMM` is accurate, fast, and expands the range of genetic analyses that are computationally feasible, we turn to the question of scalability. Specifically, we assess whether `Grid-LMM` is sufficiently accurate for the much larger sample sizes commonly used in modern human GWAS's. There are no conceptual hurdles to implementing `Grid-LMM` for studies with tens-to-hundreds of thousands of samples and the improvement in time, relative to a direct mixed modeling approach, should increase dramatically (see S1 Figa). Unfortunately, total computational time and memory requirements will grow significantly as well (see S1 Figb). For example, storing a single Cholesky decomposition of the random effect covariance matrix for 100,000 samples would require approximately 80 Gb RAM and would take approximately two days to compute. This contrasts with `BOLT-LMM` which can run a GWAS analysis of this size in less than an hour, while using less than 10 Gb RAM [27]. However, `BOLT-LMM` is restricted to a single random effect and uses a "null-LMM" approach for estimating variance component parameters as part of a two-step analysis.

To test if `Grid-LMM`'s accuracy changes with larger sample sizes, we artificially increased and decreased the sample size of the WTCHG mouse dataset by a factor of 5, simulated phenotypic data based on a randomly selected marker and a random effect with an additive relationship matrix. We then compared the marker's p-values calculated with the exact mixed model (`Exact-LMM`) to those calculated using `Grid-LMM`

Fig 3. Variance component estimates for *Arabidopsis* gene exprssion traits.
`Grid-LMM` was fit using a grid size of 0.05 h^2-units for each of 20,178 genes. (**a**) REML
estimates of the variance components proportions for the additive ($\mathbf{K}_A$) and
pairwise-epistatic ($\mathbf{K}_E$) genetic variances. Estimates are jittered for clarity. (**b-c**)
Posterior distributions of variance component proportions for two example genes,
highlighted with red dots in (**a**). The area of each point is proportional to the posterior
mass at that combination of the two variance components. A uniform prior over the
grid was assumed, and the intercept was assigned a Gaussian prior with infinite
variance. The grey contours show the posterior density as estimated by `rstan` using
half-Student-t(3,0,10) priors on each standard deviation parameter. In each plot, the
red dot shows the REML estimates. The red cross is the posterior mean as estimated by
`Grid-LMM`. The blue cross shows the posterior mean as estimated by `rstan`.

with two resolutions: 0.1 and 0.01 h^2 units. As a baseline, we also calculated p-values ²³⁵
with the two-step method that estimates variance components under the null ²³⁶
(`null-LMM`). See Methods for details on the simulations. We used a single random effect ²³⁷
for this experiment so that we could utilize the "diagonalization" trick [23, 24] to solve ²³⁸
for the variance components rapidly at large sample sizes. However, we expect the ²³⁹
results to be similar for other models. For the `Grid-LMM` tests, we assumed that the ²⁴⁰
nearest grid vertices were exactly centered around the variance component estimate ²⁴¹
from the exact mixed model. This represented a "worse-case scenario" for `Grid-LMM`. ²⁴²

As a function of the variance contributed by the tested marker, the mean relative ²⁴³
difference in p-values among the four methods was approximately constant across the ²⁴⁴
three sample sizes (Fig 4a). There were large differences when the marker effect was ²⁴⁵
large, diminishing to no difference for small effects. `Grid-LMM(0.01)` was barely ²⁴⁶
distinguishable from the `Exact-LMM` across all sample sizes and marker effect sizes. ²⁴⁷
Mean (-$\log_{10}$) p-values from `Grid-LMM(0.1)` were similar to `null-LMM` for small effect ²⁴⁸
sizes, but closer to `Exact-LMM` for large effect sizes. This is expected because most ²⁴⁹
randomly selected markers are correlated with the dominant eigenvectors of the additive ²⁵⁰
relationship matrix; hence, large effect markers will affect the variance attributed to the ²⁵¹
random effect, but small effect markers will not. ²⁵²

While the relative change in (-$\log_{10}$) p-values is approximately constant, the range of ²⁵³
effect sizes where the approximate methods have a negative impact on power changes ²⁵⁴
across sample sizes. Assuming a genome-wide significance threshold of -$\log_{10}(p) = 8$ in ²⁵⁵
this dataset, even the `null-LMM` method will consistently declare any marker with an ²⁵⁶
effect size $> \approx 0.02\%$ of the total variance as significant for sample sizes $\approx 10,000$. If we ²⁵⁷
focus specifically on the range of effect sizes where the difference among methods may ²⁵⁸
have an impact on power, the rankings of the methods do change with sample size. At ²⁵⁹
the smallest sample size (i.e. $n = 362$), the mean -$\log_{10}(p)$-values of `Grid-LMM(0.1)` ²⁶⁰
cross the significance threshold at a smaller effect size than `null-LMM`. At the medium ²⁶¹
and large sample sizes ($n = 1814$ and $n = 9070$), the mean -$\log_{10}(p)$-values for ²⁶²

Fig 4. `Grid-LMM` is accurate but less important at larger sample sizes. We used a simulation study to assess the accuracy of the `GridLMM` approximations for GWAS at larger sample sizes. Data were simulated for three sample sizes (362, 1,814 and 9,070) samples, based off the WTCHG mice genotype data. Each simulation was parameterized such that a single randomly chosen marker explained a defined percentage of the total variance $(0 - 0.15\%)$ and 10% of the residual variation was explained by the additive relationship matrix between lines. Each simulation was run 1000 times with different causal markers. The $-\log_{10}$ (p)-values were calculated with 4 methods: (i) `Exact-LMM`, an exact LMM implementation, (ii) `null-LMM`, a two-step method with the variance component percentage estimated under the null model, and (iii)-(iv) `Grid-LMM` with grid sizes of 0.1 and 0.01 h^2 units, respectively. For the `Grid-LMM` tests, we assumed that the grid was exactly centered around the variance component estimate from the exact mixed model — a "worse-case scenario" for `Grid-LMM`. (**a**) curves were fit using the `geom_smooth` function from the `ggplot2` package [54]. The horizontal line is at $-\log_{10}(p) = 8$, a typical genome-wide significance level. (**b**) The same curves as in (**a**), but zoomed in to the approximate range of marker effects with power $\ll 1$ for each sample size. (**c**) Absolute differences in $-\log_{10}(p)$ between each approximate method and `Exact-LMM` for the first 300 simulations (i.e. different causal markers) at each effect size. Note that the y-axes are different for each sample size in (**a**), and the x-axes are different for each sample size in (**b**).

`Grid-LMM(0.1)` cross the significance threshold at a larger effect size than `null-LMM` (Fig 4b) — implying lower power. However, the results of the finer `Grid-LMM(0.01)` model remain nearly indistinguishable from those of `Exact-LMM`, irregardless of the number of samples. Fig 4c shows the distribution of changes in $-\log_{10}(p)$-values for each approximate method relative to the `Exact-LMM` for individual tests. Note that when using the `Grid-LMM-fast` heuristic, `Grid-LMM` will never perform worse than `null-LMM` because we peg the grid to the variance component estimate under the null model. We repeated these tests for datasets with different variance components and sample sizes up to $n = 90,700$. These results were qualitatively similar; although, the difference between `Exact-LMM` and `null-LMM` decreased when h^2 increased.

Discussion ₂₇₃

Grid-LMM addresses a central obstacle to the practical use of linear mixed models: the ₂₇₄
computational time needed to find optimal solutions for variance components. Our key ₂₇₅
observation is that for many common quantitative genetics analyses, optimizing ₂₇₆
variance component parameters to high precision is not necessary. When sample sizes ₂₇₇
are large, statistical power will be sufficient to detect associations even under the ₂₇₈
approximate `null-LMM` methods such as `EMMAX` [32], `pylmm` [34], or `BOLT-LMM` [27]. ₂₇₉
However when sample sizes are more limited, as in the examples we have shown here, ₂₈₀
the closer approximations achieved by `Grid-LMM` can increase power without greatly ₂₈₁
increasing computational requirements. From a Bayesian perspective, the posterior ₂₈₂
distribution of variance components tends to be broad even with large sample sizes, and ₂₈₃
coarse approximations can be sufficient given the uncertainty in their true values [55]. ₂₈₄
In GWAS applications, magnitudes of variance components are of less interest than the ₂₈₅
accuracy of test statistics for the (fixed) SNP effects, and we show that these are ₂₈₆
sufficiently accurate even with approximate variance component proportions. ₂₈₇

The advantage to relaxing the need for perfect variance component solutions is a ₂₈₈
vast reduction in both computational time and algorithmic complexity. This reduces the ₂₈₉
time required for a typical GWAS sized dataset with two-or-more random effects from ₂₉₀
days to hours, and provides a framework for applying LMMs to even more powerful ₂₉₁
statistical tools [56–58]. We optimize variance component estimation with a grid search, ₂₉₂
the simplest type of optimization algorithms. At each grid vertex, after conditioning on ₂₉₃
(relative) variance component values, the LMM simplifies to a simple linear model; ₂₉₄
therefore, general purpose solutions to normal linear models are available. This means ₂₉₅
that the simple LMMs we explored in this paper can easily be generalized to more ₂₉₆
complex methods used in other GWAS-type applications that currently cannot easily be ₂₉₇
extended to experiments with heterogeneous samples (e.g. set-tests and multi-marker ₂₉₈
regressions). ₂₉₉

We demonstrated `Grid-LMM` using three examples that are broadly representative of ₃₀₀
many experimental settings in quantitative genetics. The first was a study of ₃₀₁
gene-environment interactions, while the second and third focused on the partitioning of ₃₀₂
genetic variance among additive and non-additive components. Recent studies have ₃₀₃
shown how neglecting gene-environment covariance when estimating heritability [8] or ₃₀₄
testing genetic associations [34] can lead to biased estimates. There is little reason to ₃₀₅
expect these results to be limited to controlled environmental settings, as we have ₃₀₆
studied here. Previous work has found that incorporating spatial covariance through a ₃₀₇
Gaussian radial basis function improved estimates of trait heritability within a sample ₃₀₈
of individuals from Uganda [8]. Spatial variability also exists in agricultural field trials, ₃₀₉
and two-step approaches are frequently used where spatial variation is removed first and ₃₁₀
then genetic associations are tested on the residuals. This two-step procedure may be ₃₁₁
underpowered when true effect sizes are large. Similarly, epistatic and other non-linear ₃₁₂
genetic variation are known to be large for many traits [59–61] and accounting for this ₃₁₃
variation may improve our ability to detect both the main effects of markers (as we ₃₁₄
demonstrated above), as well as possibly interacting loci [17]. ₃₁₅

The slight differences in posterior means between the `Grid-LMM` and MCMC-based ₃₁₆
posterior estimates in Fig 3b-c are due to differences in the priors. MCMC-based LMM ₃₁₇
implementations classically use inverse-Gamma priors for variance components because ₃₁₈
of conjugacy [20]. However, others have recommended uniform or half-t-family priors for ₃₁₉
the standard-deviation parameters of hierarchical models [62], which are easily ₃₂₀
implemented in `Stan` [52]. We used a half-Student-t(3,0,10) distribution for each ₃₂₁
variance component in our `rstan` model to produce Fig 3b-c. This is easy to ₃₂₂
approximate in `Grid-LMM`; relative prior weights can simply be applied to each ₃₂₃
grid-vertex, resulting in much closer agreement of posterior summaries between the two ₃₂₄

methods (see S7 Fig). As we show in S8 Fig, supposedly "uniformative" versions of ³²⁵ both the inverse-Gamma and half-Cauchy-type priors are actually highly informative for ³²⁶ variance component *proportions*. In our experience, it is more natural to elicit priors on ³²⁷ variance component proportions than variance components themselves, particularly ³²⁸ when the phenotypes are on very different scales, because these can be interpreted as ³²⁹ the relative importance of the various factors. This is an additional advantage of the ³³⁰ LMM parameterization that we utilize in `Grid-LMM`. ³³¹

The `Grid-LMM` approach does have limitations. First, the size of the grid spanning ³³² the variance components increases nearly exponentially with the number of random ³³³ effects. Since each grid vertex requires a separate Cholesky decomposition of the ³³⁴ observation-level covariance matrix, a grid search quickly becomes prohibitively ³³⁵ expensive with more than ∼6-8 variance components. This is a general problem for ³³⁶ mixed model algorithms, and it may be possible to adapt efficient derivative-based ³³⁷ algorithms to the grid space. Our fast-heuristic search method converges on the correct ³³⁸ answer in most cases we have examined; however, likelihood surfaces of linear mixed ³³⁹ models are not always convex and this algorithm may converge onto a local maximum ³⁴⁰ in such cases. We note that most general-purpose algorithms for LMMs with multiple ³⁴¹ random effects are also sensitive to this issue. Second, for REML or posterior inference ³⁴² of variance component proportions, `Grid-LMM` estimates are accurate but not precise; ³⁴³ specifically, they are limited by the resolution of the grid. We show that this has little ³⁴⁴ impact on hypothesis testing for fixed effects, for example in GWAS settings. However, ³⁴⁵ boundaries of posterior intervals in particular may not be reliable. Nevertheless, ³⁴⁶ summaries like the posterior mean or estimates of the joint posterior density are highly ³⁴⁷ accurate (e.g. Fig 3). Third, the `Grid-LMM` approach is limited to Gaussian linear mixed ³⁴⁸ models. Generalized linear mixed model algorithms rely on iteratively re-weighting the ³⁴⁹ observations, a function that changes the covariance matrix in a way that cannot be ³⁵⁰ discretized. Finally, we have not explored LMMs with correlated random effects, ³⁵¹ although these are commonly used in quantitative genetics. Since correlation ³⁵² parameters are restricted to the interval $(-1, 1)$, discretizing correlations in parallel ³⁵³ with the variance component proportions may be feasible and is an avenue that is worth ³⁵⁴ future study. ³⁵⁵

Methods ³⁵⁶

Linear Mixed Models ³⁵⁷

We consider the following parameterization of the standard linear mixed model:

$$
\begin{aligned}
\mathbf{y} &= \mathbf{W}\boldsymbol{\alpha} + \mathbf{X}\boldsymbol{\beta} + \sum_{l=1}^{L} \mathbf{Z}_l \boldsymbol{u}_l + \mathbf{e}, \\
\boldsymbol{u}_l &\sim \mathrm{N}(\mathbf{0}, \sigma^2 h_l^2 \mathbf{K}_l), \\
\mathbf{e} &\sim \mathrm{N}(\mathbf{0}, \sigma^2 h_e^2 \mathbf{I}),
\end{aligned}
\tag{1}
$$

where n is the number of observations, L is the number of random effect terms (not ³⁵⁸ including the residuals), $\mathbf{y}$ is an $n \times 1$ vector of quantitative traits, and $\mathbf{W}$ and $\mathbf{X}$ are ³⁵⁹ $n \times c$ and $n \times p$ design matrices for covariates and marker effects, respectively, with $\boldsymbol{\alpha}$ ³⁶⁰ and $\boldsymbol{\beta}$ corresponding $c \times 1$ and $p \times 1$ vectors of coefficients. Similarly, $\mathbf{Z}_l$ are $n \times r_l$ ³⁶¹ design matrices with corresponding random effects $\boldsymbol{u}_l$, which are normally distributed ³⁶² around zero and have covariance matrices proportional to the known positive ³⁶³ semi-definite $r_l \times r_l$ matrices $\mathbf{K}_l$. Lastly, $\mathbf{e}$ is a $n \times 1$ vector of uncorrelated normally ³⁶⁴ distributed errors, and $\mathrm{N}(\bullet, \bullet)$ denotes the multivariate normal distribution. The ³⁶⁵ common variance of all random effects are denoted by σ^2, and the vector ³⁶⁶

$\mathbf{h}^2 = (h_1^2, \ldots, h_L^2, h_e^2)$ represents the proportion of variance attributed to each random
effect term or the residual error. Elements of $\mathbf{h}^2$ are all non-negative and sum to one,
forming an L-dimensional simplex.

In GWAS applications, we assume that $\mathbf{W}$ is constant across markers and the value of $\boldsymbol{\alpha}$ is not of central interest; meanwhile, $\mathbf{X}$ varies for each test and we aim to perform statistical inference on a subset of $\boldsymbol{\beta}$. In heritability estimation applications, we focus on inferring the vector $\mathbf{h}^2$. In both cases, the vectors $\boldsymbol{u}_l$ and $\mathbf{e}$ are nuisance parameters and we can integrate them out resulting in the following equivalent model:

$$\mathbf{y} \sim \mathrm{N}(\mathbf{W}\boldsymbol{\alpha} + \mathbf{X}\boldsymbol{\beta}, \sigma^2 \mathbf{V}),$$
$$\mathbf{V} = \sum_{l=1}^{L} h_l^2 \mathbf{Z}_l \mathbf{K}_l \mathbf{Z}_l^\mathsf{T} + h_e^2 \mathbf{I}. \tag{2}$$

If the matrix $\mathbf{V}$ is full-rank (which is guaranteed if $h_e^2 > 0$), we can use the inverse of the Cholesky decomposition $\mathbf{V} = \mathbf{L}\mathbf{L}^\mathsf{T}$ to transform Equation 2 to the following:

$$\mathbf{y}^* \sim \mathrm{N}(\mathbf{W}^*\boldsymbol{\alpha} + \mathbf{X}^*\boldsymbol{\beta}, \sigma^2 \mathbf{I}) \tag{3}$$

where $\mathbf{y}^* = \mathbf{L}^{-1}\mathbf{y}$, $\mathbf{W}^* = \mathbf{L}^{-1}\mathbf{W}$ and $\mathbf{X}^* = \mathbf{L}^{-1}\mathbf{X}$. Equation 3 is a simple linear model for $\mathbf{y}^*$, with the likelihood function:

$$l_F(\mathbf{y}^*; \boldsymbol{\alpha}, \boldsymbol{\beta}, \sigma^2 \,|\, \mathbf{h}^2) =$$
$$\frac{1}{2}\left[-n\log(2\pi\sigma^2) - \frac{1}{\sigma^2}(\mathbf{y}^* - \mathbf{W}^*\boldsymbol{\alpha} - \mathbf{X}^*\boldsymbol{\beta})^\mathsf{T}(\mathbf{y}^* - \mathbf{W}^*\boldsymbol{\alpha} - \mathbf{X}^*\boldsymbol{\beta})\right],$$

where efficient methods for inference of $[\boldsymbol{\alpha}, \boldsymbol{\beta}]$ and σ^2 are well known. We derive maximum-likelihood and restricted-maximum likelihood solutions for these parameters here, as well as posterior distributions under the conditional normal-inverse-gamma prior below. The log-likelihood and restricted-likelihood functions (respectively) for Equation 2 are:

$$l_F(\mathbf{y}; \boldsymbol{\alpha}, \boldsymbol{\beta}, \sigma^2, \mathbf{h}^2) = l_F(\mathbf{y}^*; \boldsymbol{\alpha}, \boldsymbol{\beta}, \sigma^2 \,|\, \mathbf{h}^2) - \log|\mathbf{L}| \tag{4}$$

and

$$l_R(\mathbf{y}; \boldsymbol{\alpha}, \boldsymbol{\beta}, \sigma^2, \mathbf{h}^2) =$$
$$l_F(\mathbf{y}; \boldsymbol{\alpha}, \boldsymbol{\beta}, \sigma^2, \mathbf{h}^2) + \frac{1}{2}\left[(c+p)\log(2\pi\sigma^2) + \log|\widetilde{\mathbf{X}}^\mathsf{T}\widetilde{\mathbf{X}}| - \log|\widetilde{\mathbf{X}}^{*\mathsf{T}}\widetilde{\mathbf{X}}^*|\right] \tag{5}$$

which are simple (and computationally inexpensive) updates to the likelihood function of Equation 3. Here, we denote $|\bullet|$ as the matrix determinant and let $\widetilde{\mathbf{X}} = [\mathbf{W}; \mathbf{X}]$ and $\widetilde{\mathbf{X}}^* = [\mathbf{W}^*; \mathbf{X}^*]$, respectively. For ML and REML applications, we can calculate the profile likelihoods $l_F(\mathbf{y}; \mathbf{h}^2)$ and $l_R(\mathbf{y}; \mathbf{h}^2)$ as functions of the profile likelihood of the rotated data, which is formed by maximizing $l_F(\mathbf{y}^*; \boldsymbol{\alpha}, \boldsymbol{\beta}, \sigma^2 \,|\, \mathbf{h}^2)$ with respect to $\boldsymbol{\alpha}$, $\boldsymbol{\beta}$, and σ^2. Namely:

$$l_F(\mathbf{y}; \mathbf{h}^2) = \frac{n}{2}\left[\log\left(\frac{n}{2\pi}\right) - 1 - \log(\mathrm{RSS}_{\mathbf{y}^*})\right] - \log|\mathbf{L}| \tag{6}$$

where $\mathrm{RSS}_{\mathbf{y}^*} = \mathbf{y}^{*\mathsf{T}}[\mathbf{I} - \widetilde{\mathbf{P}}]\mathbf{y}^*$ is the residual sum of squares, and
$\widetilde{\mathbf{P}} = \widetilde{\mathbf{X}}^*(\widetilde{\mathbf{X}}^{*\mathsf{T}}\widetilde{\mathbf{X}}^*)^{-1}\widetilde{\mathbf{X}}^{*\mathsf{T}}$ is a projection (hat) matrix.

Statistical inference by parallel grid search

We now outline the `Grid-LMM` approach for calculating approximate Wald-test statistics, and then show extensions for computing Bayesian posterior distributions of variance components and marker specific Bayes Factors.

A Wald test for the null hypothesis $\mathbf{M}\boldsymbol{\theta} = \mathbf{0}$ for $\boldsymbol{\theta} = [\boldsymbol{\alpha}^{\intercal}, \boldsymbol{\beta}^{\intercal}]^{\intercal}$ and an arbitrary $q \times (c + p)$ matrix $\mathbf{M}$ uses the general F-statistic:

$$F_{\text{Wald}} = \frac{\widehat{\boldsymbol{\theta}}^{\intercal}\mathbf{M}^{\intercal}(\mathbf{M}(\widetilde{\mathbf{X}}^{\intercal}\mathbf{V}^{-1}\widetilde{\mathbf{X}})^{-1}\mathbf{M}^{\intercal})^{-1}\mathbf{M}\widehat{\boldsymbol{\theta}}}{q} \tag{7}$$

with q and $(n - c - p)$ degrees of freedom, where $\widehat{\boldsymbol{\theta}}$ is the estimate of $\boldsymbol{\theta}$ using the REML estimate of $\mathbf{V}$ [63].

To calculate genome-wide Wald statistics, we must estimate $\widehat{\mathbf{h}}^2$ for each of the p markers tested. There are no closed-form ML (or REML) solutions for $\mathbf{h}^2$; therefore, iterative algorithms are required. A naive approach involves repeatedly inverting $\mathbf{V}$, which scales cubically in the number of observations. Since the estimates for $\widehat{\mathbf{h}}^2$ differ for each test, the total computational complexity of in a GWAS setting is $\mathcal{O}(tpn^3)$ assuming an average of $\approx t$ inversions per run of the optimization algorithm (generally $\approx 3 - 100$, increasing with the number of random effect parameters). The fast algorithm used in `GEMMA` [24] and `FaST-LMM` [23] reduces this to $\mathcal{O}(n^3 + pn^2 + ptc^2n)$ by utilizing the eigenvalue decomposition of the similarity matrix $\mathbf{K}$. However, this only works for a single random effect. We are unaware of any exact algorithm with lower computational complexity for models with multiple random effects (and full-rank covariance matrices).

With `Grid-LMM`, rather than optimizing $\mathbf{h}^2$ separately for each marker, we instead define a grid of candidate values for $\mathbf{h}^2$ and calculate the restricted profile-likelihood $l_R(\mathbf{h}^2 \,|\, \mathbf{y}^*)$ at every grid vertex for each marker. At each grid vertex, we must invert $\mathbf{V}$ once, but we can re-use this calculation for every marker. This has a computational complexity of approximately $\mathcal{O}(g(n^3 + pn^2))$ for a grid with g vertices. For all analyses reported in the main text, we use a rectangular grid with a resolution of 0.1 or 0.01 h^2-units, with the restriction that all $h_l^2 \geq 0$ and $\sum_{l=1}^{L} h_l^2 < 1$. As described below, we either peg this grid to the origin (i.e. $h_0^2 = 0$, $\forall l$), or to the REML estimate $\widehat{\mathbf{h}}_0^2$ derived from the null model with no marker effects. This grid search generates a vector of g profiled Restricted Likelihood scores for each marker. We select the values of $\widehat{\mathbf{h}}^2$ that correspond to the highest such score for each marker and use Equation 7 to calculate the approximate Wald statistics.

To calculate approximate likelihood ratio test statistics, we use the `Grid-LMM` approach to calculate the full profile-likelihoods for models under both the null and alternative hypothesis.

For Bayesian inference, rather than working with the profile likelihoods, we instead use a conditional normal-inverse-gamma prior $([\boldsymbol{\alpha}^{\intercal}, \boldsymbol{\beta}^{\intercal}]^{\intercal}, \sigma^2) \sim \text{NIG}(\mathbf{0}, \boldsymbol{\Psi}, a_0, b_0)$, and then integrate over the prior to calculate the marginal likelihood of the data given $\mathbf{h}^2$. This results in the following:

$$p(\mathbf{y} \,|\, \mathbf{h}^2) = \frac{b_0^{a_0}}{(2\pi)^{n/2 + (c+p)/2}|\mathbf{V}|^{1/2}|\boldsymbol{\Psi}|^{1/2}\Gamma(a)} \times \frac{(2\pi)^{(p+c)/2}|\boldsymbol{\Psi}^*|^{1/2}\Gamma(a^*)}{(b^*)^{a^*}}, \tag{8}$$

where $\Gamma(\bullet)$ is the gamma function, $\boldsymbol{\Psi}^* = (\boldsymbol{\Psi}^{-1} + \widetilde{\mathbf{X}}^{\intercal}\mathbf{V}^{-1}\widetilde{\mathbf{X}})^{-1}$, $a^* = a_0 + n/2$, and $b^* = b_0 + \text{RSS}_{\mathbf{y}^*, \boldsymbol{\Psi}}/2$ with $\text{RSS}_{\mathbf{y}^*, \boldsymbol{\Psi}}$ having the same form as $\text{RSS}_{\mathbf{y}^*}$ in Equation 6 — except with $\widetilde{\mathbf{P}} = \widetilde{\mathbf{X}}^*\boldsymbol{\Psi}^*\widetilde{\mathbf{X}}^{*\intercal}\widetilde{\mathbf{X}}^*\boldsymbol{\Psi}^*\widetilde{\mathbf{X}}^{*\intercal}$. See S1 Supplementary Methods for more detail on the specific derivations. We calculate the marginal likelihood $p(\mathbf{y} \,|\, \mathbf{h}^2)$ at each vertex

of the grid as described above. Assuming a discrete-valued prior $p(\mathbf{h}^2)$, we can then compute the posterior distribution of $\mathbf{h}^2$ as:

$$p(\mathbf{h}^2 \mid \mathbf{y}) = \frac{|\mathbf{V}_{\mathbf{h}^2}|^{-1/2}|\mathbf{\Psi}_{\mathbf{h}^2}^*|^{1/2}(b_{\mathbf{h}^2}^*)^{-a^*}p(\mathbf{h}^2)}{\sum_{\mathbf{h}^2}|\mathbf{V}_{\mathbf{h}^2}|^{-1/2}|\mathbf{\Psi}_{\mathbf{h}^2}^*|^{1/2}(b_{\mathbf{h}^2}^*)^{-a^*}p(\mathbf{h}^2)} \qquad (9)$$

where, for clarity, parameters that are a function of $\mathbf{h}^2$ are denoted with a subscript. 404
Continuous target densities $\pi(\mathbf{h}^2)$ can be approximated as $p(\mathbf{h}^2)$ by assigning each grid 405
vertex a probability equal to the integral of $\pi(\mathbf{h}^2)$ over the L-dimensional rectangle 406
centered at the corresponding value $\mathbf{h}^2$. We assume a uniform prior over our grid for all 407
analyses presented in the main text. Bayes factors are computed by comparing models 408
under the alternative and null hypotheses as the ratios in Equation 9. All analytical 409
calculations — including the summation in Equation 9 — can be performed on the 410
log-scale to prevent numerical underflows. Terms common to both models drop out of 411
the ratio; therefore, limiting improper priors on $\boldsymbol{\alpha}$ and σ^2 can be used, which results in 412
scale-independence [49]. 413

Accelerated grid searches 414

The full grid search described above is dramatically faster than the naive algorithm for
mixed-model GWAS analyses, as long as the vertices of the grid is less than the number
of genetic markers (i.e. $g < p$) and can easily be parallelized across multiple computers.
However, g grows rapidly as the grid resolution and number of random effects increases:

$$g = \sum_{k=1}^{L} \binom{m}{k}\binom{L-1}{k-1}, \qquad (10)$$

for a grid with m divisions per h_l^2 and, therefore, can still be slow. If we make two 415
assumptions which are commonly true, we can develop heuristics for both the 416
ML/REML and Bayesian algorithms that prevent the need to evaluate every grid vertex: 417

- The vast majority of markers have little association with the phenotype. This is a 418
  common hypothesis in GWAS settings [64–67], and for our purposes, we merely 419
  require that the percentage of variance explained individually by most markers is 420
  smaller than the difference between grid vertices $\approx 1/m$. 421

- The likelihood and/or posterior function is convex. This is not always true, since 422
  both the likelihood and posterior functions can have > 1 maximum. However, the 423
  conditions that cause these events are rare [68], and most exact LMM algorithms 424
  are also susceptible to converging to local maxima. 425

To search for the ML or REML solutions, we first find $\widehat{\mathbf{h}}_0^2$ under the null model with no 426
marker effects. We calculate the profile (restricted)-likelihood scores for each test at $\widehat{\mathbf{h}}_0^2$, 427
and then form a grid centered at this value by adding or subtracting $1/m$ to each h_l^2 in 428
all combinations. For two random effects, this grid will be a ring around $\widehat{\mathbf{h}}_0^2$ with $g_1 \leq 8$ 429
vertices (depending on if $\widehat{\mathbf{h}}_0^2$ is within $1/m$ of a boundary of the simplex). We calculate 430
the scores for each test at each vertex of the grid, and then compare the maximum 431
scores to the scores at $\widehat{\mathbf{h}}_0^2$. For every test, when no greater value is found, we choose $\widehat{\mathbf{h}}_0^2$ 432
as the maximum and skip the corresponding marker in all future calculations. For the 433
remaining p_2 markers, we select the set $\{\widehat{\mathbf{h}}_0^2, \mathbf{h}_1^2, \ldots, \mathbf{h}_j^2\}$ of grid vertices that maximized 434
the scores for 1+ tests, and form a new grid of size g_2 around all of them, dropping 435
vertices already tested. This procedure is repeated t_7 times until the new grid no-longer 436
increases scores for any test and we are confident that all (approximate) maximums 437

have been found. This accelerated search has total complexity $\mathcal{O}(t_6 n^3 + pn^2 + \sum_{i=1}^{t_7} g_i(n^3 + p_i n^2))$, with t_6 the number of iterations needed to optimize variance components under the null model, and $p_1 = p$ (see Table 1).

Analogously, to accelerate evaluations of posterior distributions, we evaluate $p(\mathbf{y} \,|\, \mathbf{h}^2)$ over an initial grid of resolution $1/m_1$ with discrete prior $p_{m_1}(\mathbf{h}^2)$ and estimate the posterior distribution as in Equation 9. Summary statistics, such as the posterior mean and variance, will be more accurate if the posterior mass is distributed across multiple vertices of the grid. Therefore, we identify the set $\mathcal{H} = \{\mathbf{h}_l^2\}$ of vertices that together explain 99% of the posterior mass. If the size of this set is below a threshold (say $|\mathcal{H}| = 10$), we then form a new grid with double the resolution $m_2 = 2m_1$ and a new prior $p_{m_2}(\mathbf{h}^2)$. Vertices that overlap between the grids can be filled in directly. We then begin filling in the new grid by evaluating vertices within $1/m_2$ distance from any h_l^2 corresponding to the vertices in $\mathcal{H}$. After each iteration, we re-calculate the set $\mathcal{H}$, and continue evaluating the neighboring vertices as long as $\mathcal{H}$ continues to grow. If the size of $\mathcal{H}$ remains smaller than the threshold after whole grid is evaluated, or no new vertices are added to $\mathcal{H}$ at the end of an iteration, we double the resolution again: $m_{i+1} = 2m_i$ and repeat the grid-filling steps again. We note that this procedure is only appropriate if the posterior is convex and, therefore, is limited to the case of uniform priors on $\mathbf{h}^2$. A similar procedure was proposed for Bayesian inference in Gaussian process models in the `GPstuff` toolbox, although it is not optimized for parallel inference in GWAS [55].

To accelerate evaluations of GWAS Bayes Factors, we combine the two previously described algorithms. In particular, we define a grid centered on $\widehat{\mathbf{h}}_0^2$ with a resolution $1/m$ that is sufficiently fine such that we expect each posterior to be distributed across multiple vertices. We then calculate $p(\mathbf{y} \,|\, \mathbf{h}^2)$ for each test, starting from $\widehat{\mathbf{h}}_0^2$ and moving out in concentric rings on the grid. After each iteration, if the new ring contributes $< 0.01\%$ to the total posterior mass (among evaluated vertices) for that test, we assume the posterior is well characterized and stop evaluating $p(\mathbf{y} \,|\, \mathbf{h}^2)$ for that marker. As for the ML and REML solutions above, markers with little to no association with the phenotype will lead to posteriors of $\mathbf{h}^2$ that are concentrated close to $\widehat{\mathbf{h}}_0^2$; hence, only markers with large effects will shift $p(\mathbf{h}^2 \,|\, \mathbf{y})$ strongly to new regions of the grid.

Unless otherwise specified, we used accelerated grid searches for all `Grid-LMM` analyses presented here with grid step sizes of 0.01 h_l^2 units.

GWAS datasets

Genotype and phenotype data on 107 *Arabidopsis thaliana* traits and 216,130 genetic markers were downloaded from `https://github.com/Gregor-Mendel-Institute/atpolydb/wiki`. We follow practices suggested by the original authors of these data and log-transformed a subset of the phenotypes prior to analyses (except for traits that had values less-than or equal to zero) [44]. For the analysis of gene-environment interactions in flowering time, we (1) extracted the trait identifiers "7_FT10" (i.e. growth chamber at 10C) and "57_FT Field" (i.e. field setting), (2) selected data from the 175 accessions that were measured in both environments, and (3) concatenated the two datasets into a single vector of 350 observations. The two traits were individually standardized to have mean zero and standard deviation one prior to analysis. We used the `sommer` package in R to calculate an additive relationship matrix from all 216,130 markers [69], and then created a G×E kinship matrix as $\mathbf{DZKZ^{\intercal}D}$ where $\mathbf{K}$ is the 175×175 additive genomic relationship matrix, $\mathbf{Z}$ is a 350×175 incidence matrix linking observations to accessions, and $\mathbf{D}$ is a 350×350 diagonal matrix with elements equal to -1 or 1 corresponding to observations measured under "7_FT10" and "57_FT Field", respectively. Plasticities for each accession were calculated as the difference between "57_FT Field" and "7_FT10". We

ran `GEMMA` (version 0.97) with the "-lmm1" option for Wald tests using the **K** matrix described above. We emulated `EMMAX` and `pylmm` functionality by estimating variance components using a null model with no marker effects, and either a single **K** matrix (for single-trait analyses) or 3 random effects (for G×E analysis). Wald test statistics were computed for each marker using our GridLMM `R` function with a grid consisting of only a single vertex. For G×E analyses, we fit a model with a main effect of the environment, a main effect for each marker of interest, and an interaction between the marker and the environment — and then calculated the Wald-F statistic only for the interaction effect.

The heterogeneous stock of mice data from the Wellcome Trust Centre for Human Genetics (`http://mtweb.cs.ucl.ac.uk/mus/www/mouse/index.shtml`) consists of 1,814 individuals from 85 families, all descending from eight inbred progenitor strains [46]. We used the marker and phenotype data provided in the `BGLR R` package [70] from which we extracted the "EndNormalBW" trait and information on the gender and cage of each mouse. Gender was used as a covariate in all analyses, and cage assignment was treated as a random effect in the three-random-effect models. Additive and epistatic kinship matrices were calculated using `sommer`. Wald test statistics and Bayes Factors were calculated using the heuristic (accelerated) grid search of `Grid-LMM`. Bayes Factors were calculated assuming flat priors on the intercept and residual variance term, a standard normal prior $N(0, 1)$ on the marker effects — similar to the previously proposed D_2 prior [49] — as well as a uniform prior over the grid of variance component proportions.

Computational timings for GWAS analyses are reported for a MacBookPro14,3 with a 2.9Ghz Intel Core i7 processor and using only a single CPU core. Further speedups are possible by parallelizing the grid search. For accelerated grid searches, REML estimates of variance components under the null model (i.e. the starting points of the grid search) were calculated with `LDAK` and are included in the computational time estimates.

Gene expression dataset

Gene expression data on 24,175 genes from 728 *Arabidopsis* accessions was downloaded from `http://signal.salk.edu/1001.php` and subsetted to genes with average counts ≥ 10. A genomic relationship matrix ($\mathbf{K}_A$) for 1,135 accessions was downloaded from `http://1001genomes.org/data/GMI-MPI/releases/v3.1/SNP_matrix_imputed_hdf5/`. Both sets of data were subsetted to an overlapping set of 665 accessions. $\mathbf{K}_A$ was then centered by projecting out the intercept and normalized to have mean diagonal elements equal to one. The pairwise-epistasis genomic relationship matrix $\mathbf{K}_E$ was calculated with element-wise multiplication as $\mathbf{K}_A \odot \mathbf{K}_A$, and then also normalized to have mean diagonal elements equal to one. The gene expression matrix was normalized and variance-stabilized using the `varianceStabilizingTransformation` function of the `DEseq2` package [71]. Grid-LMM REML estimates were compared to exact REML estimates from the `LDAK` program with variance components constrained to be non-negative. Grid-LMM posterior distributions estimates were compared to those estimated using `Stan` [52] with the `rstan R` package [53]. To speed computation, we diagonalized the $\mathbf{K}_A$ covariance matrix by calculating the singular value decomposition $\mathbf{K}_A = \mathbf{U}\mathbf{S}\mathbf{U}^\intercal$, and pre-multiplying both sides of the LMM by $\mathbf{U}^\intercal$. We used a half-Student-t prior distribution with 3 degrees of freedom and a scale of 10 for the three standard deviation parameters. We ran four MCMC chains each of length 10,000, with the first 5,000 as warmup and a thinning rate of 20. Because of poor mixing, this resulted in an "n_{eff}" of approximately 200-400 per variance component parameter for each trait.

Power simulations

We compared the power of one and two-random effect models for detecting G×E
markers based on the Arabidopsis flowering time data. We calculated additive genetic
and G×E relationship matrices as described above using the actual genotypes. We then
simulated phenotypes by summing together a G×E effect of a particular marker,
(co)variance from the two random effects, and normally distributed error. For each
combination of marker effect sizes ($\{0, 0.025, 0.05, 0.1, 0.15, 0.2\}\%$ of the total
phenotypic variance), and random effect proportions ($h_l^2 \in \{0, 0.4, 0.8\}$ for each random
effect), we ran 10,000 simulations with different randomly selected markers from the
Arabidopsis genotype data. We then fit five models to each dataset and calculated
Wald-tests for marker G×E effects using each. The five methods consisted of three
two-random effect approaches including: (1) an exact two-random effect model fit with
`LDAK`, (2) the approximate two-random effect model fit with `Grid-LMM` with a grid size
of 0.1 h^2-units, and (3) the approximate two-random effect model `pylmm` that conditions
on variance components estimated under the null model [34]. We also consider two
one-random effect models that could be fit with `GEMMA`: (4) a model that only included
the additive genetic relationships and ignored the G×E covariance, and (5) a model
that only included the G×E covariance and ignored the additive genetic relationships.

For each method and for each simulation scenario, we first calculated a genomic
control inflation factor [1] for the GxE marker tests as the ratio between the median
value of the the F-statistics returned by each model and the median value of a $F_{1,312}$
distribution since $n = 316$ and each model included $p = 4$ predictors (overall intercept,
environmental effect, the main effect of the marker, and the G× E interaction between
the environment and the marker). We then "corrected" all F-statistics by dividing by
the appropriate inflation factor, and calculated statistical power for each GWAS method
as the proportion of corrected Wald test p-values exceeding the genome-wide significance
level at the conventional Bonferroni corrected threshold $P = 2 \times 10^{-7}$ (see S4 Fig).

The results show that `Grid-LMM` maintains nearly identical power to the exact
method in each simulation, all while also maintaining well-calibrated p-values under the
null-distribution. The approximate method based on `pylmm` has uniformly lower power,
but maintains an accurate null-distribution. The one-random effect methods show
opposite results: when including only the additive relationship matrix, the
null-distribution of $-\log_{10}(p)$-values is greatly inflated when G×E-variation is large. On
the other hand, when only G×E variance is modeled and yet the additive genetic
variance is large, $-\log_{10}(p)$-values are greatly deflated. These results all confirm our
expectations that modeling covariance accurately, but not necessarily precisely, is
important for accurate association mapping in GWAS.

Sample size simulations

We developed a simulation strategy to evaluate the effect of sample size on the accuracy
of `Grid-LMM` in a reasonable amount of time without the confounding issue of changes
in population structure across different populations. Here, we focused on the case of a
mixed model with a single random effect (e.g. the additive relationship matrix $\mathbf{K}_A$)
because the REML solutions can be found very rapidly after rotating by the
eigenvectors of $\mathbf{K}_A$ [23, 24]. In particular, the restricted maximum likelihood solutions
$\{\widehat{\boldsymbol{\beta}}, \widehat{h}^2\}$ and Wald F-test for $H_0 : \boldsymbol{\beta} = \mathbf{0}$ are identical for the normal model:

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{g} + \mathbf{e}, \; \mathbf{g} \sim \mathrm{N}(0, \sigma^2 h^2 \mathbf{K}_A), \; \mathbf{e} \sim \mathrm{N}(0, \sigma^2(1 - h^2)\mathbf{I}), \qquad (11)$$

and the rotated model:

$$\mathbf{y}^* = \mathbf{X}^*\boldsymbol{\beta} + \mathbf{g}^* + \mathbf{e}^*, \ \mathbf{g}^* \sim \mathrm{N}(0, \sigma^2 h^2 \mathbf{D}_A), \ \mathbf{e}^* \sim \mathrm{N}(0, \sigma^2(1-h^2)\mathbf{I}), \qquad (12)$$

where $\mathbf{K}_A = \mathbf{U}\mathbf{D}_A\mathbf{U}$ is a singular value decomposition of $\mathbf{K}_A$, and $\mathbf{y}^* = \mathbf{U}^\mathsf{T}\mathbf{y}$, $\mathbf{X}^* = \mathbf{U}^\mathsf{T}\mathbf{X}$, $\mathbf{g}^* = \mathbf{U}^\mathsf{T}\mathbf{g}$, and $\mathbf{e}^* = \mathbf{U}^\mathsf{T}\mathbf{e}$. In model 12, both random effects have diagonal covariance matrices, so $\mathbf{V} = h^2\mathbf{D}_A + (1-h^2)\mathbf{I}$ is also diagonal and can be trivially inverted. We can find $\widehat{h}^2$ efficiently for any sample size n as a bounded one-dimensional optimization problem where all operations have $\mathcal{O}(n)$ once $\mathbf{U}$ and $\mathbf{D}_A$ are known. However, computing $\mathbf{K}_A = \mathbf{U}\mathbf{D}_A\mathbf{U}$ is $\mathcal{O}(n^3)$ and, thus, prohibitive for large samples. To circumvent this, we simulated directly from model 12 by creating the eigenvalues $\mathbf{D}_A$ and the candidate markers $\mathbf{X}^*$ directly.

We started with the genotypes of the 1,814 WTCHG heterogeneous stock mice described above. We calculated $\mathbf{K}_A$ as before and then calculated $\mathbf{U}$ and $\mathbf{D}_A$ by singular value decomposition. Next, we defined $\mathbf{X}^* = \mathbf{U}^\mathsf{T}\mathbf{X}$ for 1,000 randomly selected autosomal markers from the original dataset. This initial procedure made up our "medium" sized population. To create a "large" population, we simply repeated each diagonal element of $\mathbf{D}_A$ (i.e. the eigenvalues of $\mathbf{K}_A$) five times, and correspondingly repeated the n values of $\mathbf{X}^*$ five times to create $\mathbf{D}_{A_l}$ and $\mathbf{X}_l^*$ with sample size 9,070. While not completely realistic, this created a population similar to what we would expect if the heterogeneous stock population was created five separate times from five independent sets of progenitors; therefore, it has a similar level of structure relative to its size as the original $n = 1,814$ population. Lastly, we created a "small" population of 1/5 the original sample size by defining $\mathbf{D}_{A_s}$ with the first 362 diagonal elements of $\mathbf{D}_A$, and correspondingly the first 362 rows of $\mathbf{X}^*$. This small dataset had approximately the same amount of genetic structure as the original dataset (and so more structure relative to its size), but it represents a reasonable approximation to a highly structured small dataset. As part of this exercise, we also created an "extra-large" dataset of size $n = 90,700$. However, those results were similar to the "large" dataset, and so we do not present them here. The advantage of this simulation strategy is that we can create simulated phenotypes with defined genetic and background effects without having to repeatedly calculate large singular value decompositions.

With each set of 1,000 markers $\mathbf{X}^*$ and random effect covariance matrix $\mathbf{D}_A$, we created phenotype vectors $\mathbf{y}^*$ such that the variance contributed by $\mathbf{X}^*\boldsymbol{\beta}$ was a defined fraction between 0 and 0.15 in 16 steps, and then created residual vectors with $h^2 = 0.1$ by sampling from a multivariate normal distribution with covariance $h^2\mathbf{D}_A + (1-h^2)\mathbf{I}$. We also experimented with $h^2 = 0.5$ and $h^2 = 0.9$ but results were similar except that the differences among the methods were even smaller. For each simulated phenotype vector, we calculated Wald-test p-values using four methods: (1) `Exact-LMM` with $\widehat{h}^2$ estimated by REML using a full model including the causal marker, (2) `null-LMM` with $\widehat{h}_0^2$ estimated by REML using a null model *without* the causal marker, (3) `Grid-LMM(0.1)` with $\widehat{h}_{g.1}^2$ selected as $\widehat{h}^2 \pm 0.05$ with the greater REML score, and (4) `Grid-LMM(0.1)` with $\widehat{h}_{g.01}^2$ selected as $\widehat{h}^2 \pm 0.005$ with the greater REML score.

Software Availability

Software for computing the `Grid-LMM` is carried out in R code, which is freely available at https://github.com/deruncie/GridLMM. Scripts for running the analyses reported in the manuscript are available at https://github.com/deruncie/GridLMM_scripts.

Author Contributions Statement ₆₂₆

D.E.R. and L.C. conceived the experiment(s), D.E.R. conducted the experiment(s), 627
D.E.R. and L.C. analyzed the results. Both authors wrote and revised the manuscript. 628

Competing Financial Interests ₆₂₉

The authors have declared that no competing interests exist. 630

Supporting information ₆₃₁

S1 Fig. Approximate time and memory requirements of `Grid-LMM` as a 632
function of sample size. (**a**) Computational times for the most costly steps of typical 633
mixed model fitting algorithms: inverting an $n \times n$ covariance matrix (generally using a 634
Cholesky decomposition), and multiplying the inverse matrix by an n-vector (i.e. a 635
vector of marker genotypes). The red curve shows the time required for a Cholesky 636
decomposition using the base `R` function `chol` as a function of n. The green curve shows 637
the time required for a Cholesky matrix-by-marker matrix multiplication with 1×10^5 638
markers, as a function of n. The blue curve is the sum of the Cholesky decomposition 639
and matrix-vector multiplication operations for a single grid-cell in `Grid-LMM` with 640
1×10^5 markers. The green and blue curves are barely distinguishable across most of 641
the range because the Cholesky decomposition is generally not limiting. The purple 642
curve would be the expected time for a separate Cholesky decomposition and 643
matrix-vector multiplication for *each marker* in a GWAS with 1×10^5 markers (i.e. the 644
cost of a typical `exact-LMM` method such as `LDAK` for a single iteration). Both the 645
`Grid-LMM` and LDAK times are *per-iteration*. `Grid-LMM` requires this time at each grid 646
cell. LDAK requires multiple iterations for the REML optimization separately for each 647
marker. Generally, `Grid-LMM` will evaluate more grid cells than `LDAK` requires iterations 648
per test. However this will not cause a reversal in the relative time requirements unless 649
a very large grid is used. (**b**) Memory requirements for storing an $n \times n$ Cholesky 650
matrix as a function of sample size. In both panels, the curves were extrapolated based 651
on tests with n between 256 and 4096 (actual times shown with points). All timings 652
were estimated using base `R` functions. 653

S2 Fig. Accuracy of the log-transformed p-values across the 107 654
Arabidopsis phenotypes [44]. GWASs were run for each phenotype using 216,130 655
markers and up to 199 accessions, with a single random effect controlling for additive 656
genetic relationships among lines. For each phenotype (represented as a single point in 657
the plots), we compared the exact Wald-test $-\log_{10}(p)$ calculated by `GEMMA` to p-values 658
calculated by the approximate methods `EMMAX`, and `Grid-LMM` using either the naive 659
approach with a complete grid of size 0.1 h^2-units, or the fast heuristic algorithm 660
`Grid-LMM-fast` with a fine grid size of 0.01-h^2 units. `Grid-LMM` p-values were always at 661
least as accurate (as measured by root mean-squared-error, RMSE) as those calculated 662
by `EMMAX`. Specifically, p-values calculated with a fine grid-size of 0.01 (using the fast 663
algorithm) were nearly indistinguishable from those of `GEMMA`, except in the rare cases 664
where the REML surface was not unimodal. This was generally restricted to a small 665
subset of rare markers with small-moderate effect sizes, and only occurred for a few 666
traits. In these cases, the complete — but more coarse — grid search of `Grid-LMM` with 667
step sizes of 0.1 was more accurate. 668

S3 Fig. Genomic control inflation factors for simulated data. Simulated datasets were created based on the Atwell genotype data and the G×E analysis. We randomly selected 10,000 markers, generated simulated data with different proportions of additive (G) and gene-environment interaction (G× E) variation for each marker, and calculated Wald F-statistics for an interaction between the marker and the environment. Bars show an estimate of genomic control inflation factors [1] for each of the following five methods. `exact-LMM-G+GxE` is an exact LMM algorithm fit with `LDAK`. This model included both random effects and the marker effect. At a genome-wide scale, it is very slow, with computational complexity $\mathcal{O}(ptn^3)$. `exact-LMM-G` and `exact-LMM-GxE` are exact LMM algorithms, similar to `GEMMA`, which included only one random effect and the marker effect. `null-LMM` is an approximate method similar to `pylmm` that conditions on variance components estimated under a null model with no marker effect. It was run with both random effects. *Grid-LMM* was run with a grid size of 0.1 h^2-units and included both random effects and the marker effect. The λ values were calculated as the ratio between the median value of the the F-statistics returned by each model and the median value of a $F_{1,316-4}$ distribution. The horizontal line shows the expected value $\lambda = 1$ under the true model.

S4 Fig. Power analysis for simulated data. Bars show the genome-wide power for randomly selected SNPs in the Atwell genotype data under simulations with different proportions of additive (G) and gene-environment interaction (G×E) variation with different marker effect sizes. Simulations were generated as described in S3 Fig, and included only a single marker with zero main effect and G×E effects scaled to a defined percentage of the phenotypic variation. The remaining phenotypic variation was simulated from a multivariate normal distribution constructed by appropriately weighting the additive relationship matrix, the G×E covariance matrix, and the uncorrelated residual variation. Each simulation was run separately for 10,000 randomly selected markers. Wald F-statistics from each method were normalized by dividing by the genomic control inflation factor computed for Figure , and then *p-values* were calculated and compared to the Bonferroni corrected threshold $P = 2 \times 10^{-7}$ to determine significance.

S5 Fig. Comparison of REML estimates between `Grid-LMM` and `LDAK` for 20,843 *Arabidopsis* genes. (**a**) REML estimates for the additive genetic variance (variance component for $\mathbf{K}_A$). (**b**) REML estimates for the epistatic genetic variance (variance component for $\mathbf{K}_E$).

S6 Fig. Comparison of REML estimates for the additive genetic variance between models with and without an additional pairwise-epistasis random effect for 20,843 *Arabidopsis* genes. Both models were fit using `Grid-LMM` with a grid size of 0.05 h^2 units. Point positions are jittered for clarity.

S7 Fig. Posterior distributions of variance components for two genes under the half-Student-t(3,0,10) prior. Panels (**b**) and (**c**) of Figure 3 in the main text are repeated, except the half-Student-t(3,0,10) prior on the standard deviation of the variance components of $\mathbf{K}_A$ and $\mathbf{K}_E$ for the random effects was applied to each grid vertex. The prior was approximated by simulating 1×10^4 independent draws for σ_A, σ_E and σ_e, converting these to prior draws for h_A^2 and h_E^2, and then measuring the proportion of draws closest to each grid vertex.

S8 Fig. Inverse-Gamma and half-Student-t priors are informative for variance component proportions. We compare the implied prior distributions on

variance component proportions for three classes of priors in a two-random effect model (e.g. $\mathbf{K}_A$ and $\mathbf{K}_E$ as random effects plus uncorrelated random error). (**a**)-(**b**) Each standard deviation parameter was assigned a half-Student-t prior with 3 degrees of freedom and scale parameter of 10. (**c**)-(**d**) Each variance parameter was assigned an inverse-Gamma prior with shape parameter 2 and scale parameter 1. (**e**)-(**g**) A uniform prior was applied to the 2-dimensional simplex of $[h_A^2, h_E^2, h_e^2]$. This is the default prior in GridLMM and equivalent to all analyses reported in the main text. (**a**)-(**c**)-(**e**) 2D-density plots for the two variance component proportions. Lighter blue denotes higher prior density. (**b**)-(**d**)-(**f**) Marginal densities for the $\mathbf{K}_A$ variance component proportion under each prior. The half-Student-t prior implies high probability that only one variance component is important. The inverse-Gamma prior implies high probability that all variance component proportions are non-zero.

S1 Table   Markers associated with heterogenous stock mice body weight only when accounting for non-additive genetic covariance. Markers with $-\log_{10}(p) > 3$ on chromosomes 4 and 11 are shown. Position information for each marker is derived from [72]. Other studies that identified the same marker associations are listed. WTC: Wellcome Trust Center Heterogeneous Stock Mice. LG,SM Advanced Intercross: F_9 and F_{10} generation mice from of an intercross between LG/J and SM/J strains selected for differences in body size.

S1 Supplementary Methods   Derivation of Bayesian posterior and Bayes Factors.

Acknowledgements

D.E.R was supported by the United States Department of Agriculture (USDA) National Institute of Food and Agriculture (NIFA), Hatch project 1010469. L.C. would like to acknowledge the support of Institutional Development Award Number P20GM109035 from the National Institute of General Medical Sciences of the National Institutes of Health, which funds COBRE Center for Computational Biology of Human Disease. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of any of the funders or supporters.

References

1. Devlin B, Roeder K. Genomic control for association studies. Biometrics. 1999;55:997–1004.

2. Astle W, Balding DJ. Population Structure and Cryptic Relatedness in Genetic Association Studies. Statistical Science. 2009;24(4):451–471.

3. Price AL, Zaitlen NA, Reich D, Patterson N. New approaches to population stratification in genome-wide association studies. Nature reviews Genetics. 2010;11(7):459–463. doi:10.1038/nrg2813.

4. Vilhjálmsson BJ, Nordborg M. The nature of confounding in genome-wide association studies. Nature Reviews Genetics. 2012;14(1):1–2.

5. Berg JJ, Harpak A, Sinnott-Armstrong N, Joergensen AM, Mostafavi H, Field Y, et al. Reduced signal for polygenic adaptation of height in UK Biobank. bioRxiv. 2018;doi:10.1101/354951.

6. Howard R, Carriquiry AL, Beavis WD. Parametric and Nonparametric Statistical Methods for Genomic Selection of Traits with Additive and Epistatic Genetic Architectures. G3: Genes|Genomes|Genetics. 2014;4(6):1027–1046. doi:10.1534/g3.114.010298.

7. Speed D, Balding DJ. MultiBLUP: Improved SNP-based prediction for complex traits. Genome Research. 2014;24(9):gr.169375.113–1557.

8. Heckerman D, Gurdasani D, Kadie C, Pomilla C, Carstensen T, Martin H, et al. Linear mixed model for heritability estimation that explicitly addresses environmental variation. Proceedings of the National Academy of Sciences USA. 2016;113(27):7377–7382.

9. Lynch M, Walsh B. Genetics and Analysis of Quantitative Traits. Sinauer; 1998.

10. Lynch M. Methods for the Analysis of Comparative Data in Evolutionary Biology. Evolution; international journal of organic evolution. 1991;45(5):1065–1080.

11. Wang L, Zhang B, Wolfinger RD, Chen X. An Integrated Approach for the Analysis of Biological Pathways using Mixed Models. PLoS Genet. 2008;4(7):e1000115–.

12. Wilson AJ, Réale D, Clements MN, Morrissey MM, Postma E, Walling CA, et al. An ecologist's guide to the animal model. Journal Of Animal Ecology. 2010;79(1):13–26.

13. Wang L, Jia P, Wolfinger RD, Chen X, Grayson BL, Aune TM, et al. An efficient hierarchical generalized linear mixed model for pathway analysis of genome-wide association studies. Bioinformatics. 2011;27(5):686–692.

14. Korte A, Vilhjálmsson BJ, Segura V, Platt A, Long Q, Nordborg M. A mixed-model approach for genome-wide association studies of correlated traits in structured populations. Nature genetics. 2012;44(9):1066–1071. doi:10.1038/ng.2376.

15. Weissbrod O, Lippert C, Geiger D, Heckerman D. Accurate liability estimation improves power in ascertained case-control studies. Nature Methods. 2015;12:332 EP –.

16. Dutta D, Scott L, Boehnke M, Lee S. Multi-SKAT: General framework to test multiple phenotype associations of rare variants. bioRxiv. 2017;.

17. Crawford L, Zeng P, Mukherjee S, Zhou X. Detecting epistasis with the marginal epistasis test in genetic mapping studies of quantitative traits. PLoS Genet. 2017;13(7):e1006869. doi:10.1371/journal.pgen.1006869.

18. Moore R, Casale FP, Bonder MJ, Horta D, Franke L, Barroso I, et al. A linear mixed model approach to study multivariate gene-environment interactions. bioRxiv. 2018; p. 270611.

19. Sudlow C, Gallacher J, Allen N, Beral V, Burton P, Danesh J, et al. UK Biobank: An Open Access Resource for Identifying the Causes of a Wide Range of Complex Diseases of Middle and Old Age. PLOS Medicine. 2015;12(3):e1001779.

20. Hadfield JD. MCMC Methods for Multi-Response Generalized Linear Mixed Models: The MCMCglmm R Package. Journal of Statistical Software. 2010;33(2):1–22.

21. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using lme4. Journal of Statistical Software. 2015;67(1):1–48. doi:10.18637/jss.v067.i01.

22. Mishchenko K, Neytcheva M. New Algorithms for Evaluating the Log-Likelihood Function Derivatives in the AI-REML Method. Communications in Statistics - Simulation and Computation. 2009;38(6):1348–1364.

23. Lippert C, Listgarten J, Liu Y, Kadie CM, Davidson RI, Heckerman D. FaST linear mixed models for genome-wide association studies. Nature methods. 2011;8(10):833–835.

24. Zhou X, Stephens M. Genome-wide efficient mixed-model analysis for association studies. Nature Genetics. 2012;44(7):821–824.

25. Hannah MV, Casale FP, Stegle O, Birney E. LiMMBo: a simple, scalable approach for linear mixed models in high-dimensional genetic association studies. bioRxiv. 2018;.

26. Kadie CM, Heckerman D. Ludicrous Speed Linear Mixed Models for Genome-Wide Association Studies. bioRxiv. 2018;.

27. Loh PR, Tucker G, Bulik-Sullivan BK, Vilhjálmsson BJ, Finucane HK, Salem RM, et al. Efficient Bayesian mixed-model analysis increases association power in large cohorts. Nature Genetics. 2015;47(3):284–290.

28. Zhou X. A unified framework for variance component estimation with summary statistics in genome-wide association studies. The Annals of Applied Statistics. 2017;11(4):2027–2051.

29. Tan Z, Roche K, Zhou X, Mukherjee S. Scalable Algorithms for Learning High-Dimensional Linear Mixed Models; 2018.

30. Zhou H, Blangero J, Dyer TD, Chan KhK, Lange K, Sobel EM. Fast Genome-Wide QTL Association Mapping on Pedigree and Population Data. Genet Epidemiol. 2016;41(3):174–186.

31. Zhang Z, Ersoz E, Lai CQ, Todhunter RJ, Tiwari HK, Gore MA, et al. Mixed linear model approach adapted for genome-wide association studies. Nature Genetics. 2010;42(4):355–360.

32. Kang HM, Sul JH, Service SK, Zaitlen NA, Kong Sy, Freimer NB, et al. Variance component model to account for sample structure in genome-wide association studies. Nature Genetics. 2010;42(4):348–354.

33. Rakitsch B, Lippert C, Stegle O, Borgwardt K. A Lasso multi-marker mixed model for association mapping with population structure correction. Bioinformatics. 2013;29(2):206–214.

34. Sul JH, Bilow M, Yang WY, Kostem E, Furlotte N, He D, et al. Accounting for Population Structure in Gene-by-Environment Interactions in Genome-Wide Association Studies Using Mixed Models. PLOS Genetics. 2016;12(3):e1005849.

35. Gilmour AR, Agriculture N. ASREML reference manual / A R Gilmour ... [et al.]. NSW Agriculture [Orange]; 1999.

36. Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: A Tool for Genome-wide Complex Trait Analysis. American journal of human genetics. 2011;88(1):76–82.

37. Wang M, Roux F, Bartoli C, Huard-Chauveau C, Meyer C, Lee H, et al. Two-way mixed-effects methods for joint association analysis using both host and pathogen genomes. Proceedings of the National Academy of Sciences USA. 2018;8:201710980.

38. Yu J, Pressoir G, Briggs WH, Bi IV, Yamasaki M, Doebley JF, et al. A unified mixed-model method for association mapping that accounts for multiple levels of relatedness. Nature Genetics. 2006;38(2):203–208.

39. Schifano ED, Epstein MP, Bielak LF, Jhun MA, Kardia SLR, Peyser PA, et al. SNP Set Association Analysis for Familial Data. Genet Epidemiol. 2012;66(5):797–810.

40. Listgarten J, Lippert C, Kang EY, Xiang J, Kadie CM, Heckerman D. A powerful and efficient set test for genetic markers that handles confounders. Bioinformatics. 2013;29(12):1526–1533.

41. Friedman J, Hastie T, Tibshirani R. Regularization Paths for Generalized Linear Models via Coordinate Descent. Journal of Statistical Software. 2010;33(1):1–22.

42. Schelldorfer J, Bühlmann P, van de Geer S. Estimation for High-Dimensional Linear Mixed-Effects Models Using ℓ1-Penalization. Scandinavian Journal of Statistics. 2011;38(2):197–214.

43. Gilmour AR. Mixed model regression mapping for QTL detection in experimental crosses. Computational Statistics & Data Analysis. 2007;51(8):3749–3764.

44. Atwell S, Huang YS, Vilhjálmsson BJ, Willems G, Horton M, Li Y, et al. Genome-wide association study of 107 phenotypes in Arabidopsis thaliana inbred lines. Nature. 2010;465(7298):627–631.

45. Vitezica ZG, Varona L, Legarra A. On the Additive and Dominant Variance and Covariance of Individuals Within the Genomic Selection Scope. Genetics. 2013;195(4):1223–1230. doi:10.1534/genetics.113.155176.

46. Valdar W, Solberg LC, Gauguier D, Burnett S, Klenerman P, Cookson WO, et al. Genome-wide genetic association of complex traits in heterogeneous stock mice. Nature Genetics. 2006;38(8):879–887.

47. Norgard EA, Jarvis JP, Roseman CC, Maxwell TJ, Kenney-Hunt JP, Samocha KE, et al. Replication of long-bone length QTL in the F9-F10 LG,SM advanced intercross. Mammalian genome : official journal of the International Mammalian Genome Society. 2009;20(4):224–235.

48. Liu J, Wang K, Ma S, Huang J. Accounting for linkage disequilibrium in genome-wide association studies: A penalized regression method. Statistics and its interface. 2013;6(1):99–115.

49. Servin B, Stephens M. Imputation-Based Analysis of Association Studies: Candidate Regions and Quantitative Traits. PLOS Genetics. 2007;3(7):e114.

50. Wakefield J. Bayes factors for genome-wide association studies: Comparison with P-values. Genet Epidemiol. 2009;33(1):79–86.

51. Kawakatsu T, Huang SSC, Jupe F, Sasaki E, Schmitz RJ, Urich MA, et al. Epigenomic Diversity in a Global Collection of Arabidopsis thaliana Accessions. - PubMed - NCBI. Cell. 2016;166(2):492–505.

52. Carpenter B, Gelman A, Hoffman M, Lee D, Goodrich B, Betancourt M, et al. Stan: A Probabilistic Programming Language. Journal of Statistical Software, Articles. 2017;76(1):1–32. doi:10.18637/jss.v076.i01.

53. Stan Development Team. RStan: the R interface to Stan; 2018. Available from: `http://mc-stan.org/`.

54. Wickham H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York; 2016. Available from: `http://ggplot2.org`.

55. Vanhatalo J, Pietiläinen V, Vehtari A. Approximate inference for disease mapping with sparse Gaussian processes. Statistics in Medicine. 2010;29(15):1580–1607.

56. Runcie D, Mukherjee S. Dissecting High-Dimensional Phenotypes with Bayesian Sparse Factor Analysis of Genetic Covariance Matrices. Genetics. 2013;194(3):753–767.

57. Lea AJ, Tung J, Zhou X. A Flexible, Efficient Binomial Mixed Model for Identifying Differential DNA Methylation in Bisulfite Sequencing Data. PLOS Genetics. 2015;11(11):e1005650.

58. Sun S, Hood M, Scott L, Peng Q, Mukherjee S, Tung J, et al. Differential expression analysis for RNAseq using Poisson mixed models. Nucleic Acids Research. 2017;45(11):e106–e106.

59. Phillips PC. Epistasis—the essential role of gene interactions in the structure and evolution of genetic systems. Nat Rev Genet. 2008;9(11):855–867. doi:10.1038/nrg2452.

60. Mackay TFC. Epistasis and quantitative traits: using model organisms to study gene-gene interactions. Nat Rev Genet. 2014;15(1):22–33.

61. Kerwin RE, Feusier J, Muok A, Lin C, Larson B, Copeland D, et al. Epistasis x environment interactions among Arabidopsis thaliana glucosinolate genes impact complex traits and fitness in the field. New Phytol. 2017;215(3):1249–1263. doi:10.1111/nph.14646.

62. Gelman A. Prior distributions for variance parameters in hierarchical models . Bayesian Analysis. 2006;1(3):515–533.

63. Kang HM, Zaitlen NA, Wade CM, Kirby A, Heckerman D, Daly MJ, et al. Efficient control of population structure in model organism association mapping. Genetics. 2008;178(3):1709–1723.

64. Zhou X, Carbonetto P, Stephens M. Polygenic Modeling with Bayesian Sparse Linear Mixed Models. PLOS Genetics. 2013;9(2):e1003264–.

65. Loh PR, Tucker G, Bulik-Sullivan BK, Vilhjálmsson BJ, Finucane HK, Salem RM, et al. Efficient Bayesian mixed model analysis increases association power in large cohorts. Nature genetics. 2015;47(3):284–290. doi:10.1038/ng.3190.

66. Moser G, Lee SH, Hayes BJ, Goddard ME, Wray NR, Visscher PM. Simultaneous Discovery, Estimation and Prediction Analysis of Complex Traits Using a Bayesian Mixture Model. PLoS Genetics. 2015;11(4):e1004969. doi:10.1371/journal.pgen.1004969.

67. Tang Z, Shen Y, Zhang X, Yi N. The Spike-and-Slab Lasso Generalized Linear Models for Prediction and Associated Genes Detection. Genetics. 2017;205(1):77.

68. Guiard V. About the Multimodality of the Likelihood Function when Estimating the Variance Components in a One-Way Classification by Means of the ML or REML Method. In: Proceedings of the International Conference on Linear Statistical Inference LINSTAT '93. Dordrecht: Springer, Dordrecht; 1994. p. 139–146.

69. Covarrubias-Pazaran G. Genome assisted prediction of quantitative traits using the R package sommer. PLoS ONE. 2016;11:1–15.

70. de los Campos G, Rodriguez PP. BGLR: Bayesian Generalized Linear Regression; 2016. Available from: `https://CRAN.R-project.org/package=BGLR`.

71. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biology. 2014;15:550. doi:10.1186/s13059-014-0550-8.

72. Shifman S, Bell JT, Copley RR, Taylor MS, Williams RW, Mott R, et al. A High-Resolution Single Nucleotide Polymorphism Genetic Map of the Mouse Genome. PLoS Biology. 2006;4(12):e395.