The molecular signatures of compatible and incompatible pollination

Authors: Chie Kodera[1*], Jérémy Just[1], Martine Da Rocha[2], Antoine Larrieu[1,3], Lucie Riglet[1], Jonathan Legrand[1,4], Frédérique Rozier[1], Thierry Gaude[1], Isabelle Fobis-Loisy[1*,]

[1] Laboratoire Reproduction et Développement des Plantes, Univ Lyon, ENS de Lyon, UCB Lyon 1, CNRS, INRA, Lyon, France

[2]INRA, Université Côte d'Azur, INRA, CNRS, ISA 400 route des Chappes BP 167, 06903 Sophia Antipolis Cedex, France

[3]Centre for Plant Sciences, Faculty of Biological Sciences, University of Leeds, Leeds, UK

[4] Laboratoire Reproduction et Développement des Plantes, Univ Lyon, ENS de Lyon, UCB Lyon 1, CNRS, INRA, Inria, F-69342, Lyon, France

[*]chie.kodera@ens-lyon.fr, isabelle.fobis-loisy@ens-lyon.fr

Abstract:

Fertilization in flowering plants depends on the early contact and acceptance of pollen grains by the receptive papilla cells of the stigma. To identify the associated molecular pathways, we developed a transcriptomic analysis based on single nucleotide polymorphisms (SNPs) present in two *Arabidopsis thaliana* accessions, one used as female and the other as male. We succeeded in distinguishing 80 % of transcripts according to their parental origins and drew up a catalog of genes whose expression is modified after pollen-stigma interaction. Global analysis of our data reveals pattern-triggered immunity (PTI). From this analysis, we predicted the activation of the Mitogen-activated Protein Kinase 3 on the female side after compatible pollination, which we confirmed through expression and mutant analysis. Altogether this works provides the molecular signatures of compatible and incompatible pollination, highlights the active status of incompatible stigmas, and unravels a new MPK3-dependent cell wall feature associated with stigma-pollen interaction.

Introduction:

In flowering plants, the early interaction between the tip of the female organ (stigma) and the male gametophyte (pollen grain) act as a checkpoint for fertilization. This first step of the female-male interaction includes recognition by the female tissues of the male partner. In the Brassicaceae, sophisticated mechanisms have evolved that allow the papilla cells of the stigma to reject self pollen while accepting non-self pollen. These self/non-self recognition mechanisms are known as self-incompatibility and promote genetic variability within species. Following compatible pollination, the dry pollen grain starts to hydrate on the stigma papilla, germinates a tube that penetrates the wall of stigmatic cells and grows down to convey the male gamete towards the ovules for fertilization[1,2]. By contrast, when a pollen grain is recognized as incompatible, it fails to hydrate properly and does, not or only poorly, germinate. This rejection mechanism is initiated by a ligand-receptor interaction and is genetically controlled by a single polymorphic locus called the S-locus[3]. The S-locus Cysteine Rich protein (SCR)/ S-locus protein 11 (SP11) located on the pollen surface interacts with its cognate S-locus Receptor Kinase (SRK) localized to the plasma membrane of the papilla cells[4,5]. This interaction leads to the phosphorylation of SRK that triggers the downstream cascade leading to pollen rejection[1,5,6]. Cellular events triggered in the stigma papillae by these two pathways, compatible and incompatible, start to be more clearly defined. Compatible pollination induces actin network orientation, calcium export and polarized secretion towards the pollen grain[7–11]. Incompatible pollen leads to inhibition of actin polymerization and vesicular trafficking accompanied by a strong calcium influx within the stigmatic cell[7,12,13]. Stigmatic calcium fluxes were reported to involve the autoinhibited Ca^{2+}-ATPase13 (ACA13) for pollen acceptance[8] and a glutamate receptor-like channel (GLR) for pollen rejection[12]. In addition, the stigmatic EXO70A1 protein was identified as a factor required for polarized secretion during compatible pollination, which is negatively regulated in incompatible reaction[13,14].

Whereas these results help us understand specific downstream pathways, they are dependent on strong a priori. To get a global picture of the early fertilization events with no a priori, transcriptome approaches were also conducted. The main goal was to draw up catalogues of genes modulated during pollination so as to unravel the stigmatic response to compatible or incompatible pollen[8,15–18]. However, the main drawback of these approaches was the difficulty in accurately distinguishing between pollen and stigma derived transcripts. To address that issue, translatome analysis[19] has recently been applied to identify sex-specific genes expressed during

pollination, but this strategy needs great amounts of tissues from the specific transgenic line and

fine techniques of biochemistry.

Here, inspired by previous RNA-seq analysis[20], we developed a new experimental procedure

coupled with a bioinformatic analysis of sequencing data to unravel the comprehensive molecular

dynamics that occurs both in the stigma and pollen grain following compatible and incompatible

pollinations. We took advantage of the SNPs existing between two distinct *Arabidopsis thaliana*

accessions, one used as female (Col-0) and the other as male (C24), to differentiate male and

female transcripts based on a new statistical methodology, ASE-TIGAR[21] which can take all

sequenced reads in account even those without SNPs while previous study used only reads with

SNPs[20]. Our analysis allowed the identification of 80 % of mRNAs according to their parental

origin and revealed transcriptional changes occurring specifically in either the stigma or pollen

grain/tube. Gene Ontology and signaling pathway prediction of up-regulated genes showed that

pattern-triggered immunity (PTI) pathways, including induction of the Mitogen-activated Protein

Kinase 3 (MPK3), were activated on the female side after compatible pollination. Mutant analysis

then confirmed that MPK3 is implicated in the growth of pollen tubes in papilla cells.

Results:

Experimental setup to isolate transcripts from compatible and incompatible pollination in *A. thaliana*

To compare self-compatible and self-incompatible pollination, we needed to have the same

genetic background, albeit with different stigmatic responses to pollen. *Arabidopsis thaliana* is

highly self-compatible, derived from an obligate outbreeding ancestor by mutations that disrupt

the self-incompatible system. To obtain a self-incompatible *Arabidopsis thaliana* background, we

transformed *A. thaliana* with a functional *SRK-SCR* gene pair isolated from its close self-

incompatible relative *A. lyrata*[22,23]. To analyze both compatible and incompatible reactions and

take advantage of nucleotide polymorphism between *A. thaliana* accessions, we generated two

transgenic lines that restored incompatible response: one in Col-0 background expressing the

SRK gene from the *A. lyrata* *S*14 haplotype (Col-0/*SRK14*) and the other in C24 background

expressing the *SCR* gene from the same *S*-haplotype (C24/*SCR14*) (Fig. 1a, Supplementary Fig.

1a). To express the SRK protein in stigmatic cell, we used the *SLR1* promoter that displays a high

stigma-specific activity in *Brassicaceae*[24,25]. Expression of the *SCR* gene was controlled by its

own promoter. We selected one transgenic line for Col-0/*SRK14* and one C24/*SCR14*. Both lines

were self-fertile.

Next, we tested the fertilization response in the crosses. As expected, stigmas of Col-0/*SRK14*

pollinated with wild-type C24 pollen showed a clear compatible reaction with hydration of pollen

grains and more than 55 pollen tubes elongation on papilla cells (Fig. 1a upper part,

Supplementary Fig. 1bd). By contrast, when Col-0/*SRK14* stigmas were pollinated with

C24/*SCR14* pollen, a strong incompatible reaction was observed as deduced from poor pollen

hydration and the absence of pollen tube germination (Fig. 1a lower part, Supplementary Fig.

1cd).

Building on these experimental validations, we performed a time-course experiment focusing on

two time-points of the interaction to identify genes whose expression is modified following

pollen-stigma interaction. We selected an early stage (10 minutes after pollen deposition), which

corresponds to the start of pollen grain hydration, and a later stage (one hour after pollination)

when pollen tubes reach the bottom of the stigma. We sequenced mRNAs extracted from

pollinated stigmas at 10, and 60 min after compatible (Col-0/*SRK14* x C24) pollination (C10,

C60, respectively) and for incompatible (Col-0/*SRK14* x C24/*SCR14*) pollination, we sequenced

only mRNAs extracted at 10 and 60 min after pollination (I10, I60, respectively) (Fig. 1a). We

also extracted mRNA right after pollination, 0 min (Col-0/SRK14 x C24), as a control for both

compatible and incompatible pollination (C0).

SNP-based transcriptome analysis using variants between Col-0 and

C24

To distinguish the parental origin of the transcripts, we developed a method based on the

detection of small genomic variations, including SNPs and small insertions and deletions (indels)

between Col-0 and C24 accessions. The pipeline includes three main steps (Fig. 1b,

Supplementary Fig. 2). In the first step, variations between Col-0/*SRK14* and C24 genomes were

identified by whole-genome sequencing of the two strains with the depth of 9.3 X for Col-

0/*SRK14* and 14.2 X for C24 (Supplementary Table 1). After aligning clean reads against the

reference genome (TAIR10, accession Col-0), SNPs and short indels between the mapped reads

and the reference genome sequence were identified. Reads from Col0/*SRK14* and C24 covered

95.8% and 90.6% of TAIR10 genome sequence, respectively. When compared to the reference,

the line Col-0/*SRK1*4 had 2,032 variants, while C24/*SCR14* had 732,767 variants, respectively. In

a second step, we introduced previously identified variants in the sequence of the TAIR10

template to produce two new reference genome sequences, one for Col-0/*SRK14* and one for C24

(Fig. 1b, Supplementary Fig.2). At the end of this process, between the new genomes of Col-

0/*SRK14* and C24, we got 616,781 SNPs and 446,999 indels (Supplementary Table 2). 24 % and

31 % of genomic polymorphism were found in UTRs and CDSs for Col-0/SRK14 and C24,

respectively, and led to sequence variations in predicted mRNAs (Supplementary Table 2). We

then annotated these two new genome sequences by projecting gene models from TAIR10. We

predicted 39,205 gene models in Col-0/*SRK14* and 39,206 in C24. We then extracted predicted

mRNA sequences for each gene model from these annotated Col-0/*SRK14* and C24 genome

sequences to produce maternal and paternal reference transcripts. We obtained a list of 39,204

predicted gene models that were shared between maternal and paternal reference. The number of

predicted mRNAs with at least one SNP between Col-0/*SRK14* and C24 was 31,271 among the

total common predicted mRNAs (39,204) (Fig. 1c). This result allowed us to distinguish the

origin of about 80 % (31,271/39,204 = 79.8%) of mRNAs with SNP-based analysis. The last step

included RNA read mapping and the estimation of sex-specific isoform abundances using ASE-

TIGAR. Total length of raw reads from each condition was more than 6,400 Mb (Supplementary

Table 3). ASE-TIGAR uses a Bayesian approach to estimate allele-specific expression[21]. This

approach allowed us to use all sequenced reads even those without SNPs. After getting the

estimated read counts from ASE-TIGAR, we used DESeq2[26] to normalize counts, for female and

male transcripts, respectively (Fig. 1b, Supplementary Fig. 2 Supplementary Table. 4).

From this large dataset, we first determined the contribution of each tissue (stigma vs pollen) in

mixed samples harvested immediately after compatible pollination (C0). We found that among

the 47.0 million RNA reads, 69 % were estimated as derived from Col-0/*SRK14* (stigma), and

31 % were from C24 (pollen) (Fig. 1d) although 15 % of reads were assigned to the genes

without using SNPs. This proportion is stable over time (0, 10 min, 60 min) and independent of

the pollination type (compatible, incompatible), thus suggesting that these proportions are

reflecting the relative abundance of stigmatic cells compared with pollen grains collected by our

manipulation.

Then, we analyzed the relative abundance of each transcript between stigma and pollen. To do so,

transcript abundance was computed by ASE-TIGAR and was expressed as Fragments Per Kilo

base of exon per Million reads mapped (FPKM), which accounts for sequencing depth and gene

length. We then normalized FPKM (nFPKM) by dividing FPKM of each transcript by the ratio of

the transcript counts from stigma [nFPKM(stigma)] or pollen [nFPKM(pollen)] at C0 (Fig. 1d,

Supplementary Table 4). After computing nFPKM, we plotted them as a hexbin[27] plot to see the

distribution of gene expression levels. Genes without SNPs did not show any particular pattern

(Fig. 1e, bottom line). Gene expression levels were widely distributed and most of highly

expressed genes were specific of stigma or pollen, suggesting the existence of distinct transcript

signatures between stigma and pollen (Fig. 1e).

Post-validation of the SNP-based analysis

To have a global view of expressed genes in stigma and pollen, we built three classes of genes

from calculated nFPKM: expressed genes, sex-preferentially and sex-specifically expressed

genes (see methods for precise criteria of gene selection) (Supplementary Table 5). Briefly, we

defined genes that showed nFPKM >1 as expressed genes, those that were expressed at least a

ten-fold higher in stigma than in pollen as stigma-preferentially expressed genes, and those that

were at least one hundred-fold higher expressed in stigma than in pollen as stigma-specific

expressed genes (and vice versa for pollen preferentially and specifically expressed genes) (Fig.

2a). To evaluate the accuracy of our SNP-based analysis, we first focused on sex-specific genes

and analyzed the top 20 highly expressed genes (Supplementary Table 5) with the ThaleMine

database (https://apps.araport.org/thalemine/begin.do). Heat maps of gene expression levels

based on Cheng et al., 2016 were derived (Supplementary Fig. 3). Interestingly, stigma specific

and highly expressed genes were clearly excluded from pollen even though many of them were

also expressed in various tissues. By contrast, most pollen specific and highly expressed genes

showed expression restricted to pollen and stage 12-inflorescences, which contain mature pollens

(Supplementary Fig.3) Moreover, within the list of stigma-specific genes, we found *AtS1*

(*AT3G12000.1*), which has been described as specifically expressed in *Arabidopsis* stigmas[28,29],

and in the list of pollen-specific genes, we found *CPK34* (*AT5G19360.*1), whose derived protein

is involved in pollen tube regulation[30]. Finally, to confirm the specific expression of these genes

in stigma or pollen, we carried out RT-PCR and sequence the cDNAs of SNP-containing regions

using C0 sample as template. From the top 20 highly expressed genes, we selected genes whose

location of SNPs permitted designing of primers. We found that, as expected, mRNAs of genes in

the stigma-specific genes had SNPs only from Col-0/*SRK14* (stigma), whereas mRNAs of genes

in the pollen-specific genes had SNPs only from C24 (pollen) SNPs (Fig. 2b).

To assess the global consistency of our data sets with already published studies, we further

analyzed our SNP-based data by correlating them tissue-specific transcripts reported from

microarray experiments[31–33]. Stigma associated transcripts from our analysis showed poor

correlation with male transcriptomes from mature pollen or growing pollen tube, whereas the

highest correlation was observed with transcriptome from unpollinated stigmas. Conversely, our

pollen associated transcriptome showed a high correlation with male transcriptomes and only

poorly correlated with female transcriptome or transcriptomes of vegetative tissues (Fig. 2c).

Correlations between another recent SNP-based analysis, which identified pistil- and pollen tube-

specific transcripts 8 hour after pollination[20] and the tissue/organ-specific transcripts showed

similar trends to our stigma- and pollen-transcripts (Fig. 2c).

To further characterize the pollen vs. stigma associated transcripts, we looked for Gene Ontology

(GO) term enrichment at C0 within the list of sex-preferentially expressed genes (FDR < 0.05).

Although GO terms may be somewhat subjective or not fully consolidated by functional tests,

with the current annotation, we observed a clear difference between the two sets of transcripts

(Fig. 2d). From GO term enrichment of the top 1000 highly expressed genes among stigma or

pollen -preferentially expressed genes (Supplementary Table 5, Supplementary Table 6), our

analysis revealed high enrichment of several GO terms associated with metabolism in stigmas

(such as "photosynthesis", "mitochondrial-", and "-metabolic process") suggesting an active

metabolic state of stigmatic cells (Fig. 2d left). Conversely, GO terms on the pollen side were

specific to pollen functions such as "pollen tube growth", "pollen sperm cell differentiation" and

"cell tip growth" (Fig. 2d right).

Osaka et al.[34] previously reported the transcriptome of unpollinated stigmas using laser

microdissection of *Arabidopsis* stigmatic cells. Among the top 100 expressed genes in this latter

analysis, 44 were common with the top 100 expressed genes in stigma of our analysis at C0[34].

The exocyst complex, including the EXO70A1 protein, has been shown to play a role in pollen

acceptance, probably through its function in the secretion of factors required for pollen hydration

and pollen tube growth[13,14]. We found EXO70A1 to be stably expressed in stigmas over time

(nFPKM(stigma) = 27.1 at C0, with no expression change following pollination). ACA13 which

was shown to be involved in calcium fluxes was up-regulated after both compatible (FC=3.2 at

C60) and incompatible (FC=2.4 at I60) pollinations; similar up-regulation after compatible

pollination was also reported by Iwano et al., 2014.

Altogether, these results validate our SNP-based workflow, which allows identification of

female- and male- derived transcripts from a combination of tissues following pollination.

Gene expression dynamics triggered after pollination

To examine the transcriptomic response of pollen and stigma following compatible or

incompatible pollination, we first performed a principal component analysis (PCA) [35] using the

gene expression levels of each biological replicate in all conditions (Fig. 3a). Data from stigma-

and pollen-transcripts were treated separately (Fig. 3a left and right, respectively). The total

explained variance of the first two principal components (PC1 and PC2) is around 39% (28 +11)

for stigma and 35% (24 + 11) for pollen. These relatively low percentages are explained by the

fact we took all genes and not only those that are specific for compatible or incompatible

reaction. It also explain the fact that the different conditions (C0, C10, C60, I10 & I60) were not

always well separated along those axis.

Nevertheless, comparing the PCA of stigma and of pollen transcripts suggested different

dynamics in each tissue. On the stigma side, the PCA showed that both compatible and

incompatible pollinations triggered a molecular response. Along PC1 axis, capturing almost 30%

of the explained variance, samples are temporally sorted from 0 min to 60 minutes, thus

suggesting that PC1 can be interpreted as a time-axis. For the second axis, explaining 14% of the

observed variability, compatible samples are located in the lower part when incompatible samples

are located in the upper part, close to those of the starting point (C0). This seems to indicate that this axis is oriented by the compatible/incompatible effect. Meanwhile, on the pollen side, all the conditions except C60 were clustered together. Again PC1 captures the temporality of the compatible response, however we cannot conclude that PC2 is related to the compatible response effect since early compatible and incompatible clusters overlap.

The clear response pattern of pollen at 1 hour is coherent with the massive changes displayed by compatible pollens, which hydrated and germinated a pollen tube growing in the stigmatic tissue (Fig. 3a).

After performing differentially expressed gene (DEG) analysis, we found more up regulated genes than down regulated genes, and more DEGs in stigma than in pollen (1841 up in stigma vs. 595 in pollen, 513 down in stigma vs. 113 down in pollen) (Supplementary Table 7).). To get a clearer picture on the number of genes whose expressions was modified during the course of pollination, we used a Venn diagram representation (Fig. 1b) only with up regulated genes (padj < 0.1, FC > 2) to study. The Venn diagram of up-regulated genes in stigma (Fig. 3b left) showed a similar dynamics to PCA, with a moderate alteration of gene expression in incompatible reactions, with only 45 genes at I10 and 344 genes at I60. By contrast, following compatible pollination, a massive and progressive change of gene expression was detected with 414 up-regulated genes at C10 and 1038 at C60 (Fig. 3b left). On the pollen side (Fig. 3b right), very few genes had their expression modified, except in condition C60, where 551 genes were up-regulated (6 common in C10 and C60, 533 exclusively in C60) (Supplementary Table 8). Again this correlates perfectly with the PCA. Although the pollen is completely changing its physiology during compatible pollination, the stigma does not remain passive and undergoes a massive molecular reprogramming.

PTI pathways induced after compatible pollination in stigma

The rapid and global molecular changes in stigma after compatible pollination motivated us to investigate the molecular events occurring at this stage. The 944 genes (158 + 163 + 623) induced only after compatible reaction (Fig. 3b left) showed dramatically different GO terms and enrichment from those preferentially expressed at C0 in the stigma (Fig. 2d left, Fig. 3c). There were 126 terms with high confidence (FDR < 0.05) and 85 terms with FDR < 0.005

(Supplementary Table 9). In particular, we found many GO terms including "response" and

"signalling", thus suggesting the prompt activation of signalling pathways in response to the

pollination event. We also applied the enrichment analysis for 108 genes (104 + 4) that were

induced only after incompatible pollination in stigma. But it was not informative when compared

with compatible reaction as only 9 terms showed a FDR < 0.05 with the minimum FDR = 0.006.

To predict induced pathways after pollination, we then mapped the 944 up-regulated genes after

compatible pollination to KEGG pathways[36]. We found many metabolism related pathways,

signaling pathways including hormone-signaling (with 20 genes), plant-pathogen interaction

(with 22 genes) and MAPK (with 11 genes) pathways. Interestingly, two plant-pathogen

interaction pathways known as pattern-triggered immunity (PTI) induced by bacterial flg22 and

EF-Tu[37] were clearly distinguishable with 5 up-regulated genes in compatible situation (Fig. 4,

green). On the other hand, among the 108 (4+104) exclusively up-regulated genes after

incompatible pollination, we could not distinguish clear induction of PTI pathways as only one

gene, *EFR*, was up-regulated (Fig. 4, gray).

In the compatible pollination, *SERK4*, *BAK1*, *EFR* and *FLS2* - all genes known to encode

receptors of bacterial components - were expressed in stigma at C0 and *SERK4* was up regulated

at C60 (FC = 2.3). Although *MKK6* and transcription factors *WRKY22* and *WRKY25* were not

expressed at C0, the first two were already induced at C10 and induction was maintained at C60,

while *WRKY25* was induced only at C60 (all FCs were between 2 and 2.8). *MPK4* and *MPK6*

were highly expressed from C0 to C60 compared with other members of the pathway but did not

exhibit significant expression change. Interestingly, even though *MPK3*, *WRKY33* and *WRKY53*

did not have SNPs, their high induction following compatible (FC = 6.3, 10.0, 11.0 respectively)

compared with incompatible (FC = 1.5, 2.8, 5.2) reaction suggests that this PTI pathway is a

hallmark of compatible pollination (Fig. 5, Supplementary Table. 10). Altogether, our results

show that PTI pathway is rapidly induced in stigma only after compatible pollination.

Stigma-pollen interaction activates and requires MPK3

The predicted mRNA of MPK3 did not have SNPs between Col-0/*SRK14* and C24, preventing to

accurately assign the increased expression to the stigma. To check whether activation of MPK3-

associated pathways occurs in stigma, as predicted from our analysis, we derived the list of genes

known to be co-expressed with MPK3 from ATTED-II (http://atted.jp/)[38]. Genes in the list were

ordered based on Pearson's correlation coefficient with MPK3 expression (from high to low)

(Supplementary Table 11). After removing the genes without SNPs, we checked the expression

of the top 50 genes using our SNP-based analysis in stigma (Supplementary Fig. 4, upper plot)

and pollen (Supplementary Fig. 4, lower plot). At C0, most of the genes were moderately

expressed in stigma but not in pollen. After compatible pollination, they were induced only in

stigma, supporting our hypothesis of a stigma-specific MPK3 pathway induction. Subsequently,

we took advantage of the MPK3-knockout line *mpk3-DG*[39–41] to monitor the expression and

activation of MPK3 protein *in vivo*.

Immunoblotting was performed on protein extracts of pollinated stigmas from the following

combinations of crosses: Col-0 x *mpk3* (i.e., MPK3 protein only present in stigma), *mpk3* x Col-0

(i.e., MPK3 only in pollen), Col-0 x Col-0 (MPK3 in both), and *mpk3* x *mpk3* as a negative

control. Pollinated stigmas were harvested immediately or 60 min after pollination,

corresponding to C0 and C60, respectively. The absence of a specific band for MPK3 from the

mpk3 x *mpk3* sample confirmed the complete knockout of the mutant (Fig. 6a). MPK3 protein

was detected in both stigma (Col-0 x *mpk3*) and pollen (*mpk3* x Col-0) but with a higher

abundance in the stigma (Fig. 6ab). MPK3 was slightly accumulated 60 min after pollination in

stigma and in pollen (FC = 1.6 and 2.4, respectively). Subsequently, we checked whether MPK3

was activated through phosphorylation using an anti-Phospho MPK antibody. At C0, the band

corresponding to phosphorylated MPK3 was very weak, but its intensity clearly increased after

compatible pollination in stigma but not in pollen (FC = 5.2 and 0.9 respectively) (Fig. 6cd).

Altogether these results demonstrate that MPK3 is both transcriptionally and post-translationally

activated in the stigma during compatible pollination.

To elucidate whether the loss of MPK3 function might have any effect on pollen-stigma

interaction, we pollinated *mpk3* stigmas with Col-0 pollen and observed the pollinated stigmas 60

min after pollination by scanning electron microscopy. Col-0 x Col-0 pollination was used as a

control. Before pollination, there were no apparent differences between stigmas of *mpk3* and Col-

0 (Supplementary Fig. 5). Hydration of pollen grains and elongation of pollen tubes occurred

normally on *mpk3* stigmas compared to the control. However, when we measured the width of

the contact sites between the pollen tube and stigma (white arrows on Fig. 7a), even if the

distributions of the values were partly overlapping we found that contact sites were significantly

wider on *mpk3* stigmas compared with Col-0 (Fig. 7b). This demonstrates that MPK3 contributes

to the stigma-pollen interaction and structure.

328   **Discussion:**

To decipher the mechanisms that control reproduction in flowering plants, we extracted the

global molecular signatures associated with both pollen and stigmatic cells following their

interaction during compatible or incompatible pollination. Although transcriptomes of pollinated

stigmas have been recently published[17,18], these analyses did not try to distinguish between pollen

and stigma transcripts. In the present work, we provide a comprehensive transcriptomic dataset

on stigma-pollen interactions using a novel SNP-based analysis. We took advantage of a recently

developed statistical tool called ASE-TIGAR, which is based on a Bayesian approach to estimate

allele-specific expression in diploid cells[21]. Nariai et al. showed an accurate estimation of gene

expression from RNA-seq with ASE-TIGAR and identified some autosomal genes as allele-

specific genes in a human reference lymphoblastoid cell line. We applied this new tool to

discriminate transcripts from female-male mixed tissues during the first steps of the fertilization

process. Our experimental design and bioinformatic pipeline allowed us to unveil the

transcriptomic response of the pollen/pollen tube and that of the stigma following compatible or

incompatible pollination at two early (10 min and 60 min) time points of the male-female

interaction (Fig. 1). It is worth noting that another SNP-based pipeline has recently been

published[20], identifying transcripts derived from the pistil and pollen tubes collected 8h after

compatible pollination. Although we cannot compare the two analyses directly, as pollination

time points were different and only compatibility was studied in Leydon et al., both analyses

succeeded in distinguishing male and female transcripts as presented in Fig. 2c. However, while

Leydon et al., 2017 used 12 % of reads that had SNPs among the total sequenced reads, in our

analysis we took all the RNA-seq reads into account, including those without SNPs[20,21]. This

difference in the approaches has likely contributed to making our analysis more exhaustive, as

exemplified by the remarkable segregation we found between male transcripts and those derived

from female and vegetative tissues (Fig.2 c). The sex preferentially- or specific-expressed gene

lists we drew up contain many of the already identified female or male specific genes and

reported GO term enrichment[8,13,14] (Fig.2). Together with the correlation analysis (Fig. 2c) and

experimental evaluations (Fig. 2b) we can conclude that our female- / male-transcripts globally

represent stigma-/pollen-transcripts. Transcriptomic dynamics shown by PCA and Venn

diagrams were not substantial in terms of variance and number of DEGs (Fig. 3ab). These results

are in line with those of Matsuda et al.[17] who reported only a few percent of difference between

transcripts before and 60 min after pollination for both compatible and incompatible reaction.

While our data are rather consistent with other transcriptomic analyses of pollinated stigmas[8,17,34]

(Fig. 2, 3), the clear distinction we made between male and female transcripts also reveals

surprises. The Venn diagram suggested a rapid transcriptomic response in the stigma for

compatible (414 up-regulated genes at C10) but not for incompatible pollination (45 up-regulated

genes at I10) (Fig. 3b left). Since almost no specific transcriptional induction was detected at I10

(only 4 up-regulated genes), this suggests that key molecules required for the incompatibility

reaction are already present in the stigma (Fig.3b left, I10up). This may be consistent with the

observation that the pollen rejection response in *A. thaliana* lines exhibiting a restored

incompatibility system is extremely rapid and occurs within minutes[12]. Astonishingly, while no

cellular changes were detectable by SEM on stigmas one hour after incompatible pollination (Fig.

1a), we identified more than one hundred up-regulated genes in stigmas and only 6 in pollen at

I60 (Fig. 3b). This demonstrates that the stigma undergoes massive molecular changes following

incompatible pollination, i.e. that incompatibility is actively maintained through a global

physiological modification of the stigmatic cells. The stigma up-regulated genes can be

assimilated to the stigmatic response to self-incompatible pollen. As incompatibility reaction is

maintained for several days, at least as long as SRK is properly expressed in the stigma, we may

propose that among the stigma up-regulated genes there are key factors required for the

maintenance of SI. Since components of the signaling pathway downstream of the SRK-

SP11/SCR receptor-ligand complex are still not fully unveiled, our data provide a key resource to

identify new genes involved in the incompatibility response and its maintenance over time.

Our results revealed a molecular signature of compatible pollination in stigma through the

activation of stress response genes and PTI pathways (Figs. 3, 4, 5). The similarity between the

stigma-pollen interaction and plant-pathogen interaction has been discussed and has drawn

researchers' attention for a long time[10,42–44]. Several transcriptomic analyses of pollinated

stigmas/pistils reported enrichment in transcripts linked to stress or defense response in female

tissues[18,20,33,44,45]. Tung et al. and Leydon et al. suggested that such defense-related genes would

have function in pollination. Moreover, the incompatible response has been considered to share

common mechanism with plant-pathogen interaction as it is a response to block the invader[18,44].

However, in our analysis, GO terms associated with defense responses did not clearly appear

after incompatible pollination but rather were mainly represented in compatible response (Fig.

3c).

Recently, Zhang et al.[18] performed a time course transcriptome analysis of compatible and

incompatible reaction in *Brassica napus*. They speculated that compatible responses has close

parallels with plant-pathogen interactions, mainly with effector-triggered susceptibility, while

incompatible response rather resembles effector-triggered immunity, and that PTI could be

common to both compatible and incompatible responses. Although we found up-regulated

defense related genes in the incompatible reaction, no clear defense pathways were identified by

KEGG analysis. By contrast, our study clearly indicates that two PTI pathways were induced

during compatible response (Fig.4). Indeed, the SNP-based analysis demonstrated the stigma

specific expression and induction of genes involved in these pathways (Fig. 5). This result was

not completely unexpected since pollen tube growth has many similar cellular responses to

pathogen attacks, for example invasive growth recognition of an external invader, cell wall

digestion, calcium burst[8,46,47]. We suggest that plant responses to pollen and pathogens share

conserved molecular mechanisms at early stages of the interactions but later downstream

branches would lead to different reactions, one inhibiting the pathogen invasion, whereas the

other would promote pollen tube growth in the pistil. Because links between the symbiotic and

pathogen responses have been already underlined[48,49], we can also speculate that the response of

plant cell to symbiotic organisms could share common mechanisms with the response of stigma

to pollen. How these shared modules evolved together to end up to highly specific responses (i.e.

pollination, PTI, symbiosis) is an intriguing question for further research.

MPK3 is one of the key components of PTI pathways[37], and we confirmed both the induction of

genes associated with MPK3 expression and MPK3 activation in compatible pollinated stigmas

(Supplementary Fig. 4, and Fig. 6). An increased pollen-stigma contact site was observed in

mpk3 mutant suggesting that the structural prosperities of the cell wall in the mutant are somehow

impaired (Fig. 7). Although the role of mechanical signals and cell wall reorganization has been

studied in growing pollen tube *in vitro*[50–52], very little is known on component involved in pollen

tube penetration and elongation within the stigma. Important cell wall modifications occur to

allow pollen tube growth through softening and loosening of the cell matrix[47]. How the pollen

tube and stigma papilla contribute to these changes is still unknown even though both male and

female partners are likely involved. The identification of a MPK3-dependent stigma-pollen

interaction structure provides an interesting phenotype feature for such future studies.

MPK3 is involved in sensing cell wall modifications and reorganization during development as

well as following perception of external signals like pathogen attacks and wounding through

receptor-like Wall-Associated Kinases (WAKs)[53,54]. WAKs bind pectin fragments and

oligogalacturonic acids that are generated by pathogen attacks and wounding. In stigmas, two of

the *WAK* family members (*AT2G23450* and *AT1G79670*) were stably expressed (nFPKM(stigma)

= 11.7 and 12.2, respectively at C0 with no expression change following pollination), and another

member, *AT1G21210*, was up-regulated after compatible pollination (FC = 2.3 at C60).

Interestingly, Glycine-Rich Protein3 (GRP3), a possible ligand for WAKs that is known to bind

to WAK1[55] was highly expressed (nFPKM(stigma) = 303.0 at C0 and no expression change

following pollination) in stigmas. In addition, we found that several pectin related enzymes were

up-regulated in stigmas after compatible pollination. Taken together, we propose that cell wall

reorganization that allows pollen tube growth in the stigma is dependent on MPK3-based

pathways (Fig. 8).

Our study provides the molecular signatures of pollen and stigma responses following pollen-

stigma interaction in compatible and incompatible situation. Most remarkably, we reveal that the

acceptance of pollen grains involves unexpected PTI pathways and we confirm the long-time

thought that fertilization and plant pathogen responses share common molecular signaling

components.

Materials and Methods:

A. lyrata SCR14 **and** *SRK14* **gene cloning and plasmid construction**

We used Gateway® vectors[56] (Life Technologies, USA) for expression of transgenes in

Arabidopsis thaliana. The *AlSRK14* genomic sequence spanning the coding region from the start

to the stop codons (3,620 kb) was amplified from genomic DNA of an *Arabidopsis lyrata*

individual containing the *S14*-locus with specific AttB-containing primers

(5'GGGGACAAGTTTGTACAAAAAAGCAGGCTACCATGAGAGGTGTAATACCAAAGT

ACC3' and

5'GGGGACCACTTTGTACAAGAAAGCTGGGTTTACCGAGGTTCCACTTCCGTGGTGG3'

) and subsequently inserted by BP recombination into a pDon207 plasmid. A fragment of 4,081

kb containing the *AlSCR14* gene, 1,844 kb of the 5' upstream region and 817 bp of the 3'

downstream region were amplified from genomic DNA of *Arabidopsis lyrata* individual

containing the *S14*-locus using the AttB-containing primer

(5'GGGGACAAGTTTGTACAAAAAAGCAGGCTCGGGTAGCTCAACCTAGCTAAG3' and

5' ACCACTTTGTACAAGAAAGCTGGGT CATGATCACCAAAGACAAGATCC3'). This

fragment was subsequently inserted by BP recombination into a pDONR-Zeo plasmid. The DNA

fragment containing the *Brassica oleracea SLR1* promoter (1.5 kb upstream of the *SLR1* start

codon[24,25]) was inserted into a the pDONR P4-P1R plasmid[57]. The *SLR1* promoter, the genomic

sequence *AlSRK14* and a 3'mock sequence were inserted in the pK7m34GW destination vectors

by three fragment LR recombination. The genomic sequence *AlSCR14*, a 5'mock sequence and a

3'mock sequence were inserted in the pB7m34G. GenBank accession numbers for pg*AlSCR14*

(promoter and gene of *AlSCR14*) and p*SLR1*-g*AlSRK14* (*SLR1* promoter and *AlSRK14* gene) are

MH680584 and MH680585, respectively.

Plant material and growth condition

Arabidopsis thaliana C24 and Col-0 transgenic plants were generated using *Agrobacterium*

tumefaciens-mediated transformation according to Logemann et al. 2006[58]. *AlSRK14* construct

was introduced in Col-0 (Col-0/*SRK14*). *AlSCR14* construct was introduced in C24 (C24/*SCR14*).

Unique insertion lines homozygotes for the transgene were selected. mpk3-DG[39–41] was kindly

gifted from Marcel Wiermer and Yuelin Zhang. All plants including Col-0 and C24 were grown

in growth chambers under long-day cycles (16h light/8h dark at 21°C/19°C).

Aniline blue staining and observation

Buds at the end of stage 12[59] were emasculated and 18 hours later, stigmas which have reached

the stage 13 or Early 14 (14E), were pollinated with mature pollen grains. After 2 h, pollinated

stigmas were fixed in acetic acid 10%, ethanol 50% and stained with Aniline Blue for

epifluorescence microscopy observation. Germinated pollen grains with pollen tubes within the

stigmas were manually counted. We considered that pollination is incompatible when fewer than

5 pollen tubes were able to overcome the stigmatic barrier[60].

Genomic DNA and mRNA preparation

For genomic DNA extraction, about 2 mL volume young inflorescences of Col-0/SRK14 and

C24 were harvested, respectively. After grinding the material in liquid nitrogen, DNA was

extracted as described previously[61]. For RNA extraction, late stage 12[59] Col-0/SRK14 flowers

were emasculated and 16-20h after emasculation stigmas were pollinated with compatible (C24)

or incompatible pollen (C24/*SCR14*). 0, 10 or 60 minutes after compatible pollination (C0, C10

and C60) or incompatible pollination (I10, I60), 50 stigmas were harvested manually using fine

tweezers, then frozen in liquid nitrogen and stored at − 80 °C until further processing. After

grinding the material in liquid nitrogen, RNA was extracted and purified by using Arcturus®

PicoPure® RNA Isolation Kit (Applied Biosystems / Thermo Fisher Scientific) following the

manufacturer instruction except that we added a DNAse treatment (Qiagen, catalog#79254). Five

biological replicates of RNA at each point were prepared then four replicates were selected for

sequencing based on their quality and quantity.

Whole genome sequencing and variant calling

Library preparation and whole genome sequencing of Col-0/SRK14 and C24 were performed by

HELIXIO (Clermont-Ferrand, France; http://www.helixio.com/) with TruSeq® DNA PCR-free

Library Preparation kit (Illumina) and NextSeq500 platform (Illumina) applying paired-end

sequencing (2×150 bp) (Supplementary Table 1). We then proceeded to the analysis of the

sequencing data using GATK3.5 after quality check by FastQC

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc), trimming and pairing of the

resulting reads using custom Perl scripts. We aligned the reads to the Col-0 reference genome

(TAIR10) with BWA[62], applied GATK[63] base quality score recalibration, indel realignment,

duplicate removal, and performed SNP and INDEL discovery across all samples according to

GATK Best Practices recommendations[64,65]. After checking the depth of coverage of two samples

by using GATK DepthOfCoverage (Supplementary Fig. 6), we performed filtering applying

depth 3 for Col-0/SRK14 and depth 6 for C24 and homozygous for both, then obtained the

complete set of variants for each genome as VCF files.

Production of new reference genomes

Col-0/SRK14 and C24 genome sequences were derived from TAIR10 genome, by introducing

the called variants in the sequence, by using GATK FastaAlternateReferenceMaker. Gene

annotations from TAIR10 genome were transferred onto the two new genome sequences using

RATT[66] in "Strain" mode and seqret from EMBOSS suite[67]. To characterize polymorphism

(SNPs and short indels) between obtained Col-0/SRK14 and C24 genome sequences, inside and

outside predicted genes, we aligned chromosome sequences using LAST (v. 938)[68], after training

LAST on chromosomes 1.

RNA sequencing and expression analysis

Library preparation and RNA sequencing were performed by HELIXIO with TruSeq® Standard

mRNA sample Preparation kit (Illumina) and NextSeq500 platform (Illumina) applying paired-

end sequencing (2x75 bp). We then proceeded to the analysis of the sequencing data using ASE-

TIGAR after quality check by FastQC, trimming and pairing of the resulting reads using custom

Perl scripts. First, we derived mRNA reference sequences from new reference genomes and

merged them in one FASTA file, then mapped the clean reads on the mRNA reference with

Bowtie2[69]. We then run ASE-TIGAR with SAM files produced by mapping and the mRNA

reference[21] (http://nagasakilab.csml.org/ase-tigar/). Raw sequencing data is available at NCBI

database (https://www.ncbi.nlm.nih.gov/sra/SRP154565) under SRA accession: SRP154565.

Comparison with published datasets

To check the consistency of our RNA-seq data with previously published microarray data[31–33],

for each pair of condition, we computed a Pearson correlation coefficient (r) over all genes with

SNPs from our variant analysis, using mean of estimated read counts from three or four

biological replicates for RNA-seq and mean absolute intensity for multiple microarray replicates.

We also computed Pearson correlation coefficients (r) between previously published RNA-seq

data[20] and the microarray data sets.

Differential gene expression analysis and calculation of nFPKM.

Differential expression analysis on the whole transcriptome was performed using DESeq2[26].

After performing principal component analysis (PCA) for all biological replicates at each

condition, we removed one replicate at I10 from female samples and one at I60 from male

samples, as they were isolated from their other replicates. All the analyses, including the

presented PCA, were performed with the sample set after the removal.

Normalized FPKM (nFPKM) was calculated by dividing stigma- or pollen-FPKM by female or

male transcript proportion at each condition (C0, C10, C60, I10, I60).

Criteria to select expressed genes, sex–preferentially or specifically expressed genes

To select expressed genes, sex-preferentially or specifically expressed genes at C0 (Fig. 2a,

Supplementary Table 5), we used nFPKM. We defined stigma-expressed genes consisting

[nFPKM(stigma) >1 $\cap$ with SNPs] and stigma-preferentially expressed genes consisting

[nFPKM(stigma) > 1 $\cap$ nFPKM(stigma) $\geq 10 \times$ nFPKM(pollen)]. Sex-specifically expressed

genes were selected among sex-preferentially expressed genes applying more stringent criteria.

Stigma-specifically expressed genes consist [stigma preferentially expressed genes $\cap$

nFPKM(pollen) ≤ 1 $\cap$ nFPKM(stigma) > 100 $\cap$ nFPKM(stigma) > 100 x nFPKM(pollen)], then

sorted by nFPKM(stigma) from the largest to the smallest. Vice-versa for pollen (preferentially /

specifically) expressed genes

GO term and enrichment analysis and pathway analysis

GO and gene enrichment analyses were performed using Gene Ontology Consortium website

(http://www.geneontology.org/) and PANTHER13.1 software[70,71]. We used "GO biological

process complete" for the classification. GO terms with False Discovery Rate (FDR) < 0.05 were

considered as significantly enriched.

Pathway analysis was performed using and KEGG PATHWAY Database using KEGG Mapper

software (http://www.kegg.jp/kegg/tool/map_pathway2.html)[36].

RT-PCR and sequencing

For experimental evaluation of the SNP-based analysis with reverse transcription polymerase

chain reaction (RT-PCR) and sequencing, we selected genes among the top-20 specifically

expressed genes that have SNPs at suitable positions allowing discrimination of parental origin

(stigma vs pollen) (Fig. 2b). RNA at C0 was purified as described above. cDNA was generated

using SuperScript® VILO™ cDNA Synthesis Kit (Invitrogen). Primers were designed by

Primer3web (http://bioinfo.ut.ee/primer3/) at the flanking region of SNP including sites. PCR was

performed with GoTaq (Promega). PCR products were purified using PCR clean-up Gel

extraction kit (Macherey-Nagel) and then sequenced.

Protein extraction and western blotting

566 Proteins from around 50 pollinated stigmas at C10 and C60 were extracted in SDG buffer (Tris-

567 HCl 62.5 mM pH 6.8, Glycerol 10%, DTT 2%, SDS 2.5 %). Proteins were separated on a 10 %

568 SDS-PAGE gel and immunodetected with Anti-AtMPK3 (Sigma, M8318), anti-phospho p44/p42

569 MAPK (Cell Signaling, #9101S), or anti-alpha-Tubulin (Sigma, T5168). Band intensities were

570 normalized to tubulin signal using ImageJ software.

Environmental Scanning Electron Microscopy (SEM).

572 Late stage 12 stigmas were pollinated with mature Col-0 pollen and 60 min after pollination,

573 stigmas were observed under Hirox SEM (SH-3000 table-top) at -20 °C with an accelerating

574 voltage of 10 kV. The width of each pollen tube–stigma contact site was measured using ImageJ

575 sofware. We applied a Student test to the measurements.

References:

1. Doucet, J., Lee, H. K. & Goring, D. R. Pollen Acceptance or Rejection : A Tale of Two Pathways. *Trends*

Plant Sci. **21,** 1058–1067 (2016).

2. Mizuta, Y. & Higashiyama, T. Chemical signaling for pollen tube guidance at a glance. *J Cell Sci* **131,**

jcs208447 (2018).

3. Bateman, A. J. Self-incompatibility systems in angiosperms. *Heredity* **9,** 52–68 (1955).

4. Kachroo, a, Schopfer, C. R., Nasrallah, M. E. & Nasrallah, J. B. Allele-specific receptor-ligand

interactions in Brassica self-incompatibility. *Science* **293,** 1824–1826 (2001).

5. Takayama, S. *et al.* Direct ligand-receptor complex interaction controls Brassica self-incompatibility.

Nature **413,** 534–538 (2001).

6. Cabrillac, D., Cock, J. M., Dumas, C. & Gaude, T. The *S*-locus receptor kinase is inhibited by

thioredoxins and activated by pollen coat proteins. *Nature* **410,** 220–223 (2001).

7. Iwano, M. *et al.* Actin dynamics in papilla cells of Brassica rapa during self- and cross-pollination. *Plant*

Physiol. **144,** 72–81 (2007).

8. Iwano, M. *et al.* A pollen coat-inducible autoinhibited Ca2+-ATPase expressed in stigmatic papilla cells

is required for compatible pollination in the Brassicaceae. *Plant Cell* **26,** 636–49 (2014).

9. Elleman, C. J. & Dickinson, H. G. Pollen-stigma interactions in Brassica. IV. Structural reorganization in

the pollen grains during hydration. *J. Cell Sci.* **80,** 141–157 (1986).

10.   Elleman, C. J. & Dickinson, H. G. Identification of pollen components regulating pollination-

specific responses in the stigmatic papillae of Brassica oleracea. *New Phytol.* **133,** 197–205 (1996).

11.   Safavian, D. & Goring, D. R. Secretory Activity Is Rapidly Induced in Stigmatic Papillae by

Compatible Pollen, but Inhibited for Self-Incompatible Pollen in the Brassicaceae. *PLOS ONE* **8,** e84286

(2013).

12.   Iwano, M. *et al.* Calcium signalling mediates self-incompatibility response in the Brassicaceae.

Nat. Plants **1,** 1–8 (2015).

13.   Safavian, D. & Goring, D. R. Secretory activity is rapidly induced in stigmatic papillae by

compatible pollen, but inhibited for self-incompatible pollen in the brassicaceae. *PLoS ONE* **8,** (2013).

14.   Indriolo, E., Safavian, D. & Goring, D. R. The ARC1 E3 Ligase Promotes Two Different Self-Pollen

Avoidance Traits in Arabidopsis. *Plant Cell* **26,** 1525–1543 (2014).

15.   Swanson, R., Edlund, A. F. & Preuss, D. Species Specificity in Pollen-Pistil Interactions. *Annu. Rev.*

Genet. **38,** 793–818 (2004).

16.   Sankaranarayanan, S., Jamshed, M. & Samuel, M. A. Proteomics Approaches Advance Our

Understanding of Plant Self-Incompatibility Response. *J. Proteome Res.* **12,** 4717–4726 (2013).

17.   Matsuda, T. *et al.* Transcriptional Characteristics and Differences in Arabidopsis Stigmatic Papilla

Cells Pre- and Post-Pollination. *Plant Cell Physiol.* **56,** 663–673 (2015).

18.   Zhang, T. *et al.* Time-Course Transcriptome Analysis of Compatible and Incompatible Pollen-

Stigma Interactions in Brassica napus L. *Front. Plant Sci.* **8,** 682 (2017).

19.   Lin, S.-Y. *et al.* Profiling of translatomes of in vivo-grown pollen tubes reveals genes with roles in

micropylar guidance during pollination in Arabidopsis. *Plant Cell* **26,** 602–18 (2014).

20.   Leydon, A. R. *et al.* The molecular dialog between flowering plant reproductive partners defined

by SNP-informed RNA-Sequencing. *Plant Cell* **29,** 984–1006 (2017).

21.   Nariai, N., Kojima, K., Mimori, T., Kawai, Y. & Nagasaki, M. A Bayesian approach for estimating

allele-specific expression from RNA-Seq data with diploid genomes. *BMC Genomics* **17,** 2 (2016).

22.   Kusaba, M. *et al.* Self-incompatibility in the genus Arabidopsis: characterization of the S locus in

the outcrossing A. lyrata and its autogamous relative A. thaliana. *Plant Cell* **13,** 627–643 (2001).

23.   Nasrallah, M. E., Liu, P. & Nasrallah, J. B. Generation of Self-Incompatible Arabidopsis thaliana by

Transfer of Two S Locus Genes from A. lyrata. *Science* **297,** 247–249 (2002).

24.   Hackett, R. M., Cadwallader, G. & Franklin, F. Functional Analysis of a Brassica oleracea SLR1

Gene Promoter. *Plant Physiol.* **112,** 1601–1607 (1996).

25.   Fobis-Loisy, I., Chambrier, P. & Gaude, T. Genetic transformation of Arabidopsis lyrata: Specific

expression of the green fluorescent protein (GFP) in pistil tissues. *Plant Cell Rep.* **26,** 745–753 (2007).

26. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-

seq data with DESeq2. *Genome Biol.* **15,** 550 (2014).

27. Carr, D., Lewin-Koh, ported by N., Maechler, M. & Sarkar, contains copies of lattice functions

written by D. *hexbin: Hexagonal Binning Routines*. (2018).

28. Dwyer, K. G., Lalonde, B. A., Nasrallah, J. B. & Nasrallah, M. E. Structure and expression of AtS1,

an Arabidopsis thaliana gene homologous to the S-locus related genes of Brassica. *Mol. Gen. Genet.*

MGG **231,** 442–448 (1992).

29. Dwyer, K. G. *et al.* A superfamily of S locus-related sequences in Arabidopsis: diverse structures

and expression patterns. *Plant Cell* **6,** 1829–1843 (1994).

30. Myers, C. *et al.* Calcium-dependent protein kinases regulate polarized tip growth in pollen tubes.

Plant J. **59,** 528–539 (2009).

31. Honys, D. & Twell, D. Transcriptome analysis of haploid male gametophyte development in

Arabidopsis. *Genome Biol.* **5,** R85.1–R.85.13 (2004).

32. Qin, Y. *et al.* Penetration of the stigma and style elicits a novel transcriptome in pollen tubes,

pointing to genes critical for growth in a pistil. *PLoS Genet.* **5,** (2009).

33. Boavida, L. C., Borges, F., Becker, J. D. & Feijo, J. A. Whole Genome Analysis of Gene Expression

Reveals Coordinated Activation of Signaling and Metabolic Pathways during Pollen-Pistil Interactions

in Arabidopsis. *Plant Physiol.* **155,** 2066–2080 (2011).

34. Osaka, M. *et al.* Cell type-specific transcriptome of brassicaceae stigmatic papilla cells from a

combination of laser microdissection and RNA sequencing. *Plant Cell Physiol.* **54,** 1894–1906 (2013).

35. Ringnér, M. What is principal component analysis? *Nat. Biotechnol.* **26,** 303–304 (2008).

36. Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y. & Morishima, K. KEGG: new perspectives on

genomes, pathways, diseases and drugs. *Nucleic Acids Res.* **45,** D353–D361 (2017).

37.   Bigeard, J., Colcombet, J. & Hirt, H. Signaling mechanisms in pattern-triggered immunity (PTI).

Mol. Plant **8,** 521–539 (2015).

38.   Obayashi, T., Aoki, Y., Tadaka, S., Kagaya, Y. & Kinoshita, K. ATTED-II in 2018: A Plant

Coexpression Database Based on Investigation of the Statistical Property of the Mutual Rank Index.

Plant Cell Physiol. **59,** e3–e3 (2018).

39.   Li, X., Lassner, M. & Zhang, Y. Deleteagene: A fast neutron deletion mutagenesis-based gene

knockout system for plants. *Comp. Funct. Genomics* **3,** 158–160 (2002).

40.   Miles, G. P., Samuel, M. A., Zhang, Y. & Ellis, B. E. RNA interference-based (RNAi) suppression of

AtMPK6, an Arabidopsis mitogen-activated protein kinase, results in hypersensitivity to ozone and

misregulation of AtMPK3. *Environ. Pollut.* **138,** 230–237 (2005).

41.   Genenncher, B. *et al.* Nucleoporin-Regulated MAP Kinase Signaling in Immunity to a

Necrotrophic Fungal Pathogen. *Plant Physiol.* **172,** 1293–1305 (2016).

42.   Dickinson, H. Simply a social disease? *Nature* **367,** 517–518 (1994).

43.   Nasrallah, J. B. Recognition and rejection of self in plant self-incompatibility: Comparisons to

animal histocompatibility. *Trends Immunol.* **26,** 412–418 (2005).

44.   Mondragon, M., John-Arputharaj, A., Pallmann, M. & Dresselhaus, T. Similarities between

reproductive and immune pistil transcriptomes of Arabidopsis species. *Plant Physiol.* pp.00390.2017

(2017). doi:10.1104/pp.17.00390

45.   Tung, C. *et al.* Genome-Wide Identification of Genes Expressed in Arabidopsis Pistils Specifically

along the Path of Pollen Tube Growth 1 [w]. *Plant Physiol.* **138,** 1–13 (2005).

46.   Sanati Nezhad, A. & Geitmann, A. The cellular mechanics of an invasive lifestyle. *J. Exp. Bot.* **64,**

4709–4728 (2013).

47.   Marsollier, A.-C. & Ingram, G. Getting physical: invasive growth events during plant

development. *Curr. Opin. Plant Biol.* **46,** 8–17 (2018).

48.   Velázquez, E. *et al.* The Coexistence of Symbiosis and Pathogenicity-Determining Genes in

Rhizobium rhizogenes Strains Enables Them to Induce Nodules and Tumors or Hairy Roots in Plants.

Mol. Plant. Microbe Interact. **18,** 1325–1332 (2005).

49.   Chen, T. *et al.* Interplay of pathogen-induced defense responses and symbiotic establishment in

Medicago truncatula. *Front. Microbiol.* **8,** 1–13 (2017).

50.   Hamilton, E. S. *et al.* Mechanosensitive channel MSL8 regulates osmotic forces during pollen

hydration and germination. *Science* **350,** 438–441 (2015).

51.   Mecchia, M. A. *et al.* RALF4/19 peptides interact with LRX proteins to control pollen tube growth

in Arabidopsis. *Science* **358,** 1600–1603 (2017).

52.   Ge, Z. *et al.* Arabidopsis pollen tube integrity and sperm release are regulated by RALF-mediated

signaling. *Science* **358,** 1596–1600 (2017).

53.   Kohorn, B. D. & Kohorn, S. L. The cell wall-associated kinases, WAKs, as pectin receptors. *Front.*

Plant Sci. **3,** (2012).

54.   Kohorn, B. D. The state of cell wall pectin monitored by wall associated kinases: A model. *Plant*

Signal. Behav. **10,** e1035854 (2015).

55.   Park, A. R. *et al.* Interaction of the Arabidopsis Receptor Protein Kinase Wak1 with a Glycine-rich

Protein, AtGRP-3. *J. Biol. Chem.* **276,** 26688–26693 (2001).

56.   Karimi, M., Inzé, D. & Depicker, A. GATEWAY^{TM} vectors for Agrobacterium-mediated plant

transformation. *Trends Plant Sci.* **7,** 193–195 (2002).

57.   Durand, E. *et al.* Dominance hierarchy arising from the evolution of a complex small RNA

regulatory network. *Science* **346,** 1200–1205 (2014).

58.   Logemann, E., Birkenbihl, R. P., Ülker, B. & Somssich, I. E. An improved method for preparing

Agrobacterium cells that simplifies the Arabidopsis transformation protocol. *Plant Methods* **2,** 16

(2006).

59.   Smyth, D. R., Bowman, J. L. & Meyerowitz, E. M. Early flower development in Arabidopsis. *Plant*

Cell **2,** 755–767 (1990).

60.   Kitashiba, H., Liu, P., Nishio, T., Nasrallah, J. B. & Nasrallah, M. E. Functional test of Brassica self-

incompatibility modifiers in Arabidopsis thaliana. *Proc. Natl. Acad. Sci.* **108,** 18173–18178 (2011).

61.   Verger, S., Chabout, S., Gineau, E. & Mouille, G. Cell adhesion in plants is under the control of

putative O-fucosyltransferases. *Development* **143,** 2536–2540 (2016).

62.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler transform.

Bioinformatics **25,** 1754–1760 (2009).

63.   McKenna, A. *et al.* The Genome Analysis Toolkit: A MapReduce framework for analyzing next-

generation DNA sequencing data. *Genome Res.* **20,** 1297–1303 (2010).

64.   DePristo, M. A. *et al.* A framework for variation discovery and genotyping using next-generation

DNA sequencing data. *Nat. Genet.* **43,** 491–498 (2011).

65.   Auwera, G. A. V. der *et al.* From FastQ Data to High-Confidence Variant Calls: The Genome

Analysis Toolkit Best Practices Pipeline. *Curr. Protoc. Bioinforma.* **43,** 11.10.1–11.10.33 (2013).

66.   Otto, T. D., Dillon, G. P., Degrave, W. S. & Berriman, M. RATT: Rapid Annotation Transfer Tool.

Nucleic Acids Res. **39,** e57–e57 (2011).

67.   Rice, P., Longden, I. & Bleasby, A. EMBOSS: The European Molecular Biology Open Software

Suite. *Trends Genet.* **16,** 276–277 (2000).

68.   Hamada, M., Ono, Y., Asai, K., Frith, M. C. & Hancock, J. Training alignment parameters for

arbitrary sequencers with LAST-TRAIN. *Bioinformatics* **33,** 926–928 (2017).

69.   Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient alignment of

short DNA sequences to the human genome. *Genome Biol.* **10,** R25 (2009).

70.   Mi, H., Muruganujan, A. & Thomas, P. D. PANTHER in 2013: modeling the evolution of gene

function, and other gene attributes, in the context of phylogenetic trees. *Nucleic Acids Res.* **41,** D377–

D386 (2013).

71.   Thomas, P. D. *et al.* PANTHER: A Library of Protein Families and Subfamilies Indexed by Function.

Genome Res. **13,** 2129–2141 (2003).

Acknowledgments: We thank Yvon Jaillais, Olivier Hamant, Gwyneth Ingram for critical

reading of the manuscript, SiCE team members for fruitful discussions and comments, Fabrice

Besnard for critical reading of the manuscript and for specific advice and script sharing relative to

SNP calling, Naoki Nariai for the help to perform analysis with ASE-TIGAR, Alexander R.

Leydon for the information for SNP-based RNAseq analysis, Marcel Wiermer and Yuelin Zhang

for *mpk*3-DG line, Agnès Attard for the discussion of the bioinformatics data analysis. We

gratefully acknowledge support from the PSMN (Pôle Scientifique de Modélisation Numérique)

of the ENS de Lyon for the computing resources. This work was supported by Grant ANR-14-

CE11-0021.

Author Contributions: CK, JJ, MDR developed the data analysis pipeline and performed the

bioinformatics analyses. AL, JL supported the bioinformatics analyses. AL helped with

interpretation of the transcriptomic data. CK is responsible for all experiments and analysis

performed in the study except described below. FR produced and characterized Col-0/*SRK14* and

C24/*SCR14* lines. CK and LR performed the image acquisition by scanning electron microscope.

CK, LR, FR, IFL harvested plant materials and extracted RNA. CK, TG, IFL designed the study

and CK, JJ, TG, IFL wrote the manuscript. All the authors contributed to the discussion.

Figure legends:

Fig. 1 | SNP-based RNA-seq analysis. a, Time course of sample collection. Flowers of Col-
0/*SRK14* were emasculated 16h - 20h before pollination with compatible (C24) or incompatible
(C24/*SCR14*) pollen grains, respectively. Stigmas were harvested just before the stigma
(dashed line) 0, 10, 60 min after compatible (C0, C10, C60) or incompatible (I10, I60) pollination

for RNA extraction. Typical scanning electron microscope images for compatible (Col-0 /*SRK14*
x C24) and incompatible (Col-0 /*SRK14* x C24/*SCR14*) reaction observed 60 min after
pollination. Scale bar = 20 μm. **b**, Three steps of a SNP-based data analysis pipeline. **c**, Number
of predicted mRNAs derived from the reference sequence produced at the second step in b.
Around 80 % of coding regions have SNPs between Col-0/SRK14 and C24. **d**, Number and ratio
of estimated read counts for Col-0/*SRK14* and C24 at C0 computed by ASE-TIGAR at the third
step in b. Reads estimated to the stigma or pollen include reads assigned to the genes without
using SNPs. **e**, Hexibin plot of nFPKM for Col-0/*SRK14* and C24 at C0. nFPKM(stigma) of genes
without SNPs were plotted outside.

Fig. 2 | Validation of the SNP-based analysis. a, Number of stigma or pollen (preferentially /
specifically) -expressed genes at C0 selected by nFPKM (Supplementary Table 5). **b**, RT-PCR
and sequence analysis of stigma or pollen specifically-expressed genes at C0. The genes are
selected among the 20 specifically-expressed genes and the rank of the genes according to the
expression level are presented. RNA was extracted from pollinated stigmas at C0 then we
analysed their SNP-information by RT-PCR and sequencing. SNP number corresponds to the
number of SNPs present in the sequenced regions. **c**, Heat map of Pearson's correlation
coefficient between the stigma/pollen transcripts from the SNP-based analysis and the
previously published transcripts. **d**, Top ten gene enrichment categories of stigma or pollen
preferentially-expressed genes at C0 according to the GO term on biological processes.
Enrichment analysis was performed with the one thousand top expressed genes in stigma (left)
or pollen (right) among the list, respectively. Only top 10 first enrichment categories are shown.
Selection criteria for genes analysed in **b** (sex-specific), and **d** (sex-preferential), are described
in the manuscript.

Fig. 3 | Gene expression dynamics after pollination. a, PCA of stigma (left) and pollen (right)
transcripts. Biological replicates of all the conditions were analysed. **b**, Venn diagrams showing
the number of up-regulated (FC > 2) genes in stigma (left) and pollen (right), after compatible
(green and violet) or incompatible (grey) pollinations. **c**, Gene enrichment categories of up-
regulated genes after compatible pollination in stigma according to GO term on biological
processes. Only genes specifically induced after compatible reaction (b, left; 158+163+623)
were analysed. Only top 20 enrichment categories are shown.

Fig. 4 | Induced genes in PTI pathways after pollination in stigma.
KEGG pathway mapping applied for the genes exclusively up-regulated after compatible
pollination (Fig.3 b, left; 158+163+623) and incompatible pollination (Fig.3 b, right; 4+0+104),
respectively. Green colored genes were identified from the compatible gene set and grey were
from the incompatible set. Up-regulated gene in our analysis assigned to *BAK1*/*BKK1* was
SERK4, and to *MKK1/2* was *MKK6* (see Fig. 5 and Supplementary Table 10).

Fig. 5 | Induced pathways in stigma after compatible pollination. Genes in green boxes with
solid line were with SNPs and up-regulated (FC > 2) at 10 min and 60 min (light green), or only
at 60 min (green), genes in orange boxes with dashed line were without SNPs and up-regulated
(FC > 2) at 10 min and 60 min (orange), or only at 60 min (dark orange). Genes written in bold
were expressed at C0; nFPKM(stigma) > 1. The pathway map from KEGG database was
modified with the information from Bigeard et al. 2015.

Fig.6 | Activation of a MPK3-dependent pathway after compatible pollination.
Immunoblotting assays after compatible pollination with *mpk3* mutant line. Pollination between
Col-0 x Col-0, *mpk3* x *mpk3*, Col-0 x *mpk3*, and *mpk3* x Col-0 were performed. Protein was

extracted from pollinated stigmas at C0 and C60 then analysed. Anti-MPK3 antibody was used
for a, and anti-phospho-p44/42 MPK antibody was used for c. The shown western blots are
representative of three independent biological replicates. All of them showed similar results. b.,
d. Semi-quantification of the band intensity (b for a, d for c). Green, violet and grey bars indicate
signals from stigma side, from pollen side, and from both sides, respectively. Error bars indicate
standard error of the mean, ** indicates non-specific signal, *** indicates non-specific signal or
signals from other MPK proteins.
806
Fig.7 | *mpk3* stigma changes the width of pollen tube contact sites. a, Pollen tube contact
sites on Col-0 (left) or *mpk3* (right) papilla cells observed by SEM 60 min after pollination. One
stigma papilla is artificially colorized in green and one pollen grain and its tube in violet,
respectively. Scale bar = 10 µm. Double-arrows indicate the width of pollen tube contact site
analysed in b. **b**, Width of the pollen tube contact site for each pollination type presented in a.
Observations were performed three different days; n is the total number of contact sites
measured on 16 Col-0 and 25 *mpk3* stigmas. Orange points in the boxes are averages of each
day.*** indicates p-value < 0.01.
815
816 **Fig.8 | Stigma-pollen interaction involves MPK3-dependent pathway.** Schematic model for
817 stigma-pollen interaction. i, Signals at the contact site between pollen tube and papilla cell. ii,
818 Signal transduction involving MPK3 pathways. iii, Gene expression change. iv, Protein
819 activations without transcriptomic change v, Stigma cell wall reorganization. vi, Effect on pollen
820 tube by stigma surface property.
821

822 Additional information:

Supplementary Fig. 1

Supplementary Fig. 2

Supplementary Fig. 3

Supplementary Fig. 4

Supplementary Fig. 5

Supplementary Fig. 6

Supplementary Table. 1

Supplementary Table. 2

Supplementary Table. 3

Supplementary Table. 4

Supplementary Table. 5

Supplementary Table. 6

Supplementary Table. 7

Supplementary Table. 8

Supplementary Table. 9

Supplementary Table. 10

Supplementary Table. 11

Fig. 1 | SNP-based RNA-seq analysis. a, Time course of sample collection. Flowers of Col-0/*SRK14* were emasculated 16h - 20h before pollination with compatible (C24) or incompatible (C24/*SCR14*) pollen grains, respectively. Stigmas were harvested just before the stigma (dashed line) 0, 10, 60 min after compatible (C0, C10, C60) or incompatible (I10, I60) pollination for RNA extraction. Typical scanning electron microscope images for compatible (Col-0 /*SRK14* x C24) and incompatible (Col-0 /*SRK14* x C24/*SCR14*) reaction observed 60 min after pollination. Scale bar = 20 μm. **b**, Three steps of a SNP-based data analysis pipeline. **c**, Number of predicted mRNAs derived from the reference sequence produced at the second step in b. Around 80 % of coding regions have SNPs between Col-0/SRK14 and C24. **d**, Number and ratio of estimated read counts for Col-0/*SRK14* and C24 at C0 computed by ASE-TIGAR at the third step in b. Reads estimated to the stigma or pollen include reads assigned to the genes without using SNPs. **e**, Hexibin plot of nFPKM for Col-0/*SRK14* and C24 at C0. nFPKM(stigma) of genes without SNPs were plotted outside.

a

Gene expression in stigma and pollen

	Stigma	Pollen
Expressed	16964	6857
Preferentially-expressed	11094	1111
Specifically-expressed	587	59

b

Stigma specific genes	Rank	SNP number	Col-0/*SRK14* reference	C24 reference	Sequence after RT-PCR
AT4G38770.1	2	2	T/C	C/A	T/C
AT3G12000.1	3	2	G/T	A/C	G/T
AT5G24780.1	5	8	T/G/T/G/A/A/G/G	C/A/C/A/C/G/A/A	T/G/T/G/A/A/G/G
AT2G34810.1	7	6	T/T/C/C/C/A	C/C/A/A/G/G	T/T/C/C/C/A
AT5G19800.1	8	1	T	G	T

Pollen specific genes	Rank	SNP number	Col-0/*SRK14* reference	C24 reference	Sequence after RT-PCR
AT5G19360.1	1	1	C	T	T
AT3G54240.1	2	1	G	T	T
AT3G60570.1	3	3	G/A/C	A/T/T	A/T/T
AT3G56600.2	7	1	T	C	C
AT1G60410.1	19	10	T/G/C/-/-/A/G/T/T/G	C/-/-/T/G/T/T/C/C/A	C/-/-/T/G/T/T/C/C/A

c

d

Fig. 2 | Validation of the SNP-based analysis. **a**, Number of stigma or pollen (preferentially / specifically) -expressed genes at C0 selected by nFPKM (Supplementary Table 5). **b**, RT-PCR and sequence analysis of stigma or pollen specifically-expressed genes at C0. The genes are selected among the 20 specifically-expressed genes and the rank of the genes according to the expression level are presented. RNA was extracted from pollinated stigmas at C0 then we analysed their SNP-information by RT-PCR and sequencing. SNP number corresponds to the number of SNPs present in the sequenced regions. **c**, Heat map of Pearson's correlation coefficient between the stigma/pollen transcripts from the SNP-based analysis and the previously published transcripts. **d**, Top ten gene enrichment categories of stigma or pollen preferentially-expressed genes at C0 according to the GO term on biological processes. Enrichment analysis was performed with the one thousand top expressed genes in stigma (left) or pollen (right) among the list, respectively. Only top 10 first enrichment categories are shown. Selection criteria for genes analysed in **b** (sex-specific), and **d** (sex-preferential), are described in the manuscript.

Fig. 3 | Gene expression dynamics after pollination. a, PCA of stigma (left) and pollen (right) transcripts. Biological replicates of all the conditions were analysed. **b**, Venn diagrams showing the number of up-regulated (FC > 2) genes in stigma (left) and pollen (right), after compatible (green and violet) or incompatible (grey) pollinations. **c**, Gene enrichment categories of up-regulated genes after compatible pollination in stigma according to GO term on biological processes. Only genes specifically induced after compatible reaction (b, left; 158+163+623) were analysed. Only top 20 enrichment categories are shown.

| | Up-regulated after compatible pollination |
| | Up-regulated after incompatible pollination |

Fig. 4 | Induced genes in PTI pathways after pollination in stigma.
KEGG pathway mapping applied for the genes exclusively up-regulated after compatible pollination (Fig.3 b, left; 158+163+623) and incompatible pollination (Fig.3 b, right; 4+0+104), respectively. Green colored genes were identified from the compatible gene set and grey were from the incompatible set. Up-regulated gene in our analysis assigned to *BAK1*/*BKK1* was *SERK4*, and to *MKK1/2* was *MKK6* (see Fig. 5 and Supplementary Table 10).

Fig. 5 | Induced pathways in stigma after compatible pollination. Genes in green boxes with solid line were with SNPs and up-regulated (FC > 2) at 10 min and 60 min (light green), or only at 60 min (green), genes in orange boxes with dashed line were without SNPs and up-regulated (FC > 2) at 10 min and 60 min (orange), or only at 60 min (dark orange). Genes written in bold were expressed at C0; nFPKM(stigma) > 1. The pathway map from KEGG database was modified with the information from Bigeard et al. 2015.

Fig.6 | Activation of a MPK3-dependent pathway after compatible pollination. Immunoblotting assays after compatible pollination with *mpk3* mutant line. Pollination between Col-0 x Col-0, *mpk3* x *mpk3*, Col-0 x *mpk3*, and *mpk3* x Col-0 were performed. Protein was extracted from pollinated stigmas at C0 and C60 then analysed. Anti-MPK3 antibody was used for a, and anti-phospho-p44/42 MPK antibody was used for c. The shown western blots are representative of three independent biological replicates. All of them showed similar results. b., d. Semi-quantification of the band intensity (b for a, d for c). Green, violet and grey bars indicate signals from stigma side, from pollen side, and from both sides, respectively. Error bars indicate standard error of the mean, ** indicates non-specific signal, *** indicates non-specific signal or signals from other MPK proteins.

Fig.7 | *mpk3* stigma changes the width of pollen tube contact sites. a, Pollen tube contact sites on Col-0 (left) or *mpk3* (right) papilla cells observed by SEM 60 min after pollination. One stigma papilla is artificially colorized in green and one pollen grain and its tube in violet, respectively. Scale bar = 10 µm. Double-arrows indicate the width of pollen tube contact site analysed in b. **b**, Width of the pollen tube contact site for each pollination type presented in a. Observations were performed three different days; n is the total number of contact sites measured on 16 Col-0 and 25 *mpk3* stigmas. Orange points in the boxes are averages of each day.*** indicates p-value < 0.01.

Fig.8 | Stigma-pollen interaction involves MPK3-dependent pathway. Schematic model for stigma-pollen interaction. i, Signals at the contact site between pollen tube and papilla cell. ii, Signal transduction involving MPK3 pathways. iii, Gene expression change. iv, Protein activations without transcriptomic change v, Stigma cell wall reorganization. vi, Effect on pollen tube by stigma surface property.