Title: A Metagenomic Comparison of Tracheal Aspirate and Mini-Bronchial

Alveolar Lavage for Assessment of Respiratory Microbiota

Running Title: Sequencing of Tracheal Aspirate and Mini-BAL Specimens

Authors: Katrina L. Kalantar[1]; Farzad Moazed[2]; Stephanie C. Christenson[2]; Jenny

Wilson[3]; Thomas Deiss[2]; Annika Belzer[2]; Kathryn Vessell[2], Alejandra Jauregui[2],

Samaneh Boulourchi[2]; Joseph L. DeRisi[1,4], Carolyn S. Calfee[2], Charles Langelier[5]

Affiliations:

[1] Department of Biochemistry and Biophysics, University of California, San Francisco,

San Francisco, CA, USA

[2] Division of Pulmonary, Critical Care, Sleep and Allergy, Department of Medicine,

University of California, San Francisco, CA, USA

[9] Department of Emergency Medicine, Stanford University, Stanford, California

[4] Chan Zuckerberg Biohub, San Francisco, CA, USA

[5] Division of Infectious Diseases, Department of Medicine, University of California, San

Francisco, CA, USA

Corresponding author:

Charles Langelier, MD, PhD

Division of Infectious Diseases

Department of Medicine

1700 4th St, Room 403

San Francisco CA, 94158

chaz.langelier@ucsf.edu

Funding:

NHLBI K12HL119997 (Langelier C), NHLBI K23HL123778 (Christensen SC), NHLBI

R01 HL110969 (Calfee CS), NHLBI R01 HL140026 (Calfee CS), Chan Zuckerberg

Biohub (DeRisi JL).

Keywords: next generation sequencing, lower respiratory tract infection, pneumonia,

tracheal aspirate, mini-bronchial alveolar lavage

Abstract (220 / 250 words):

Accurate and informative microbiologic testing is essential for guiding diagnosis and management of pneumonia in critically ill patients. Sampling of tracheal aspirate (TA) is less invasive compared to mini-bronchoalveolar lavage (mBAL) and is now recommended as a frontline diagnostic approach in mechanically ventilated patients, despite the historical belief that TA was suboptimal due to contamination from oral microbes. Advancements in metagenomic next generation sequencing (mNGS) now permit assessment of airway microbiota without a need for culture, and as such provide an opportunity to examine differences between mBAL and TA at a resolution previously unachievable. Here, we engaged shotgun mNGS to quantitatively assess the airway microbiome in matched mBAL and TA specimens from a prospective cohort of critically ill adults. We observed moderate differences between sample types across all patients (Pearson correlation of 0.72, 95% CI: 0.68 – 0.76), however we found significant compositional similarity in patients with bacterial pneumonia, whose microbial communities were characterized by a dominant pathogen (Pearson correlation of 0.92, 95% CI: 0.88 – 0.95). In addition, we found that both mBAL and TA were similar in terms of microbial burden, abundance of oropharyngeal taxa, and microbial diversity. Our findings suggest that TA sampling provides a similar assessment of airway microbiota as more invasive testing by mBAL, and that this similarity is most significant in the setting of bacterial pneumonia.

Importance (150 / 150 words):

Less invasive tracheal aspirate (TA) sampling has historically been considered inferior to mini-bronchial alveolar lavage for both diagnostic microbiology and for investigations of the lung microbiome. Recent studies demonstrating that sample type does not affect patient outcomes has led to shift in clinical practice guidelines, which now recommend TA. Despite the implications of this shift for both patient care and lung microbiome research, differences in microbial composition between fluid types had not been assessed using metagenomic methods. To address this gap, we employed metagenomic sequencing to characterize the microbial composition of matched mBAL and TA specimens. While modest differences in taxonomic composition were observed, mBAL and TA were highly similar with respect to microbial burden and diversity, and in patients with bacterial pneumonia. Our findings provide metagenomic evidence that TA is an acceptable replacement for more invasive mBAL sampling, which has implications both for clinical microbiology and lung microbiome research.

Observation (997 / 1200 words):

Pneumonia causes more deaths each year in the United States than any other type of infectious disease (1). The ability to accurately detect etiologic pathogens and distinguish them from background commensal microbiota is essential for guiding optimal antimicrobial treatment. In patients requiring mechanical ventilation, less invasive tracheal aspirate (TA) sampling has historically been considered inferior to specimen collection by mini-bronchoalveolar lavage/telescoping catheter (mBAL) due to the potential for oropharyngeal microbiota contamination (2, 3). This idea has been challenged, however, by studies demonstrating a lack of clinically significant differences between sample types, (2–5)   and a greater acceptance of TA sampling is now reflected in recent updates to clinical practice guidelines (6). Despite the broad potential implications of this shift in diagnostic sampling approach, relatively little information exists regarding microbial composition differences between mBAL and TA specimens and the potential implications of such differences for both clinical diagnostic testing and airway microbiome studies.

To address this gap in knowledge, we used shotgun metagenomic next generation sequencing (mNGS) to evaluate the microbial compositions of matched mBAL and TA specimens from 52 adults who were admitted to the intensive care unit with acute respiratory failure and enrolled under University of California San Francisco protocol 10-02701 as detailed in (**Table S1**) and in supplemental materials . Two-physician adjudication based on retrospective medical record review (blinded to mNGS results) and the United States Centers for Disease Control surveillance case definition of pneumonia was used to identify 15 subjects with culture-confirmed bacterial

pneumonia (PNA-pos) (8). Adjudication also identified 12 patients with a clear

alternative non-infectious etiology of acute respiratory failure (PNA-neg), and 25

patients with acute respiratory illnesses of indeterminate etiology (PNA-ind), which

included patients with negative bacterial cultures but suspected pneumonia based on

clinical criteria alone (**Table S2**). Patients with PCR-confirmed viral etiologies were also

included in the PNA-ind group because occult bacterial co-infection could not be

excluded.

Excess mBAL and TA specimens collected on the same day, within 72 hours of

intubation, underwent DNA extraction and sequencing library preparation according to

previously described methods (8). Following paired-end Illumina sequencing, we

employed a custom bioinformatics pipeline (8) to detect and profile airway microbes

(**Figure S1**) . We sequenced no-template water control samples and restricted analyses

to taxa present at > 1% of the microbial population by abundance, as previously

described (2, 9). No microbe was universally present in every sample, suggesting that

systematic contamination across TA or mBAL sampling methods was unlikely.

We began by evaluating the correlation between genus abundance in mBAL and

TA specimens and identified a Pearson correlation of 0.72 (95% CI: 0.68 – 0.76),

suggesting moderate differences (**Figure 1A**). Next, we examined microbial burden

(total genus alignments per million reads sequenced) and found no significant

differences between TA and mBAL specimens (Median = 43.30 (IQR: 6.69 – 327.93)

versus 26.89 (5.66 - 167.63), p = 0.66 by Wilcoxon rank sum). Oropharyngeal

microbiota have historically been suspected to compromise TA specimens, and thus we

next evaluated for differences in the abundance of common oropharyngeal microbiota

(2). Surprisingly, we found no statistically significant differences between mBAL and TA

specimens with respect to *Prevotella, Veillonella, Streptococcus, Fusobacterium,*

Rothia, or *Neisseria* abundance (**Table 1B**). We also identified no significant inter-

specimen type differences in microbial β-diversity based on Bray-Curtis dissimilarity

index (p = .31 by PERMANOVA).

We reasoned that differences in microbial composition between mBAL and TA

specimens would be most clinically significant if they impacted diagnostic accuracy in

patients with pneumonia, and thus assessed taxonomic similarity between the PNA-pos

and PNA-neg groups. We found significantly greater similarity in PNA-pos subjects as

compared to the PNA-neg subjects (Pearson correlation of 0.92 (0.88 – 0.95) versus

0.39 (0.28 – 0.49), p < 1.00 x 10^{-3}), suggesting that pathogen dominance of the lung

microbiome during infection may drive compositional similarity (9, 10) (**Table 1A, Figure**

1B). For both sample types, a culture-confirmed pathogen was the most abundant

microbe detected by mNGS in 14 (93%) of PNA-pos subjects, and the second most

abundant in the remaining subject (**Table S3**). mNGS of mBAL specimens detected all

23 culture-identified microbes, while mNGS of TA samples identified 22. The discrepant

microbe was from a polymicrobial culture and was detected by mNGS in the TA

specimen, but present at < 1% relative abundance and therefore indistinguishable from

background using our bioinformatic approach (**Table S3**).

Reduced alpha diversity of the human respiratory microbiome has been

described as an ecological marker of infection(9, 10), and thus we next asked whether

Shannon's Diversity Index (SDI) differed by specimen type. We found that SDI differed

significantly between PNA-pos and PNA-neg patients within mBAL samples, and a

similar, albeit less significant difference, was also observed within TA samples

(Wilcoxon rank sum p = 5.2×10^{-6} and 4.7×10^{-2}, respectively) (**Figure S2**). Community

richness (total number of different genera identified in each sample) was significantly

decreased in PNA-pos compared to PNA-neg subjects when assessed by either

specimen type, although the difference was again more significant for mBAL than for TA

(Wilcox rank sum p = 1.2×10^{-3} versus 6.5×10^{-2}, **Table 1**).

Advances in genome sequencing have revealed that the lung, previously

considered sterile, supports diverse microbial communities that play a role in both

health and disease (2). Using shotgun mNGS to compare mBAL and TA samples, we

found that while some differences exist in microbial composition between these two

sample types, the dissimilarities become insignificant in the setting of a pathogen-

dominated infection. As such, our data indicate that from a metagenomic perspective,

TA sampling is an effective surrogate for more invasive testing by mBAL, a conclusion

consistent with findings of prior clinical studies (4, 5) and the Clinical Practice

Guidelines from the Infectious Diseases Society of America and the American Thoracic

Society (6). These results may help inform both culture-independent clinical

microbiologic testing, and research on the lung microbiome.

Data availability: Data are available publicly via BioProject Accession ID

PRJNA445982. Analyses of primary sequencing data including pre-processed data files

and an R markdown file with documentation are available on GitHub:

(https://github.com/katrinakalantar/mBALvTA_comparison).

References:

1. Hansen V, Oren E, Dennis LK, Brown HE. 2016. Infectious disease mortality trends in the united states, 1980-2014. JAMA 316:2149–2151.

2. Dickson RP, Erb-Downward JR, Freeman CM, McCloskey L, Falkowski NR, Huffnagle GB, Curtis JL. 2017. Bacterial Topography of the Healthy Human Lower Respiratory Tract. mBio 8.

3. Boersma WG, Erjavec Z, van der Werf TS, de Vries-Hosper HG, Gouw ASH, Manson WL. 2007. Bronchoscopic diagnosis of pulmonary infiltrates in granulocytopenic patients with hematologic malignancies: BAL versus PSB and PBAL. Respir Med 101:317–325.

4. Canadian Critical Care Trials Group. 2006. A randomized trial of diagnostic techniques for ventilator-associated pneumonia. N Engl J Med 355:2619–2630.

5. Berton DC, Kalil AC, Teixeira PJZ. 2014. Quantitative versus qualitative cultures of respiratory secretions for clinical outcomes in patients with ventilator-associated pneumonia. Cochrane Database Syst Rev CD006482.

6. Kalil AC, Metersky ML, Klompas M, Muscedere J, Sweeney DA, Palmer LB, Napolitano LM, O'Grady NP, Bartlett JG, Carratalà J, El Solh AA, Ewig S, Fey PD, File J Thomas M, Restrepo MI, Roberts JA, Waterer GW, Cruse P, Knight SL, Brozek JL. 2016. Management of Adults With Hospital-acquired and Ventilator-associated Pneumonia: 2016 Clinical Practice Guidelines by the Infectious

Diseases Society of America and the American Thoracic Society. Clin Infect Dis

63:e61–e111.

8.   United States Centers for Disease Control and Prevention. 2017. CDC/NHSN

Surveillance Definitions for Specific Types of Infections.

9.   Langelier C, Zinter MS, Kalantar K, Yanik GA, Christenson S, O'Donovan B, White

C, Wilson M, Sapru A, Dvorak CC, Miller S, Chiu CY, DeRisi JL. 2017.

Metagenomic Sequencing Detects Respiratory Pathogens in Hematopoietic

Cellular Transplant Patients. Am J Respir Crit Care Med.

10. Abreu NA, Nagalingam NA, Song Y, Roediger FC, Pletcher SD, Goldberg AN,

Lynch SV. 2012. Sinus microbiome diversity depletion and Corynebacterium

tuberculostearicum enrichment mediates rhinosinusitis. Sci Transl Med

4:151ra124.

Figure Legends and Tables

Figure 1. A. Pearson correlation demonstrates taxonomic similarity by genus in mBAL

versus TA specimens (r = 0.72 overall, (95% CI: 0.68 – 0.76)). **B.** Pearson correlation

was stronger among microbes from PNA-pos subjects (red plus, r = 0.92 (95% CI: 0.88

– 0.95)) than PNA-neg subjects (blue plus, r = 0.39 (95% CI: 0.28 – 0.49)).

Table 1. A) Microbial community metrics in tracheal aspirate (TA) and mini-bronchial

alveolar lavage (mBAL) samples, including differences between subjects with

pneumonia (PNA-pos) and those with respiratory failure due to etiologies other than

respiratory infection (PNA-neg). **B)** Abundance differences (reads per million reads

mapped, rpM) for common oropharyngeal microbes by genus rpm between TA and

mBAL samples calculated using Wilcoxon Rank Sum.

A.	mBAL		TA		
Metric	Median (IQR)	PNA-pos vs. PNA-neg	Median (IQR)	PNA-pos vs. PNA-neg	mBAL vs. TA
Richness	6.00 (4.00-9.00)	$p = 6.5 \times 10^{-2}$	8.00 (4.00-13.50)	$p = 1.2 \times 10^{-3}$	$p = 5.0 \times 10^{-2}$
Shannon Diversity	1.05 (0.71-1.55)	$p = 4.7 \times 10^{-2}$	1.45 (0.74-2.05)	$p = 5.2 \times 10^{-6}$	$p = 6.0 \times 10^{-2}$

B.	mBAL	TA	
Genus	**mean rpM (IQR)**	**mean rpM (IQR)**	mBAL vs. TA
Prevotella	0.07 (0.00 - 0.06)	0.02 (0.00 - 0.03)	p = 0.15
Veillonella	0.03 (0.00 - 0.02)	0.03 (0.00 - 0.02)	p = 0.99
Streptococcus	0.20 (0.00 - 0.33)	0.18 (0.00 - 0.34)	p = 0.88
Fusobacterium	0.02 (0.00 – 0.00)	0.01 (0.00 - 0.00)	p = 0.53
Rothia	0.03 (0.00 - 0.03)	0.01 (0.00 - 0.00)	p = 0.31
Neisseria	0.04 (0.00 - 0.02)	0.03 (0.00 - 0.02)	p = 0.72