Experimental demonstration that screening can enable the

environmental recruitment of a defensive microbiome

Tabitha Innocent[1]*[†], Neil Holmes[2]*, Mahmoud Al Bassam[2][††], Morten Schiøtt[1], István

Scheuring[3,4], Barrie Wilkinson[5], Matthew I Hutchings[2][†], Jacobus J Boomsma[1][†], and

Douglas W Yu[2,6,7][†]

[1] Centre for Social Evolution, Department of Biology, University of Copenhagen, 2100

Copenhagen Ø, Denmark

[2] School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich,

Norfolk NR4 7TJ United Kingdom

[3] MTA-ELTE Theoretical Biology and Evolutionary Ecology Research Group, Pázmány Péter

sétány, 1/C, 1117, Budapest, Hungary

[4] MTA Centre for Ecological Research, Evolutionary Systems Research Group, Klebelsberg K. u.

3, 8237, Tihany, Hungary

[5] Department of Molecular Microbiology, John Innes Centre, Norwich Research Park, Norwich,

Norfolk NR4 7UH United Kingdom

[6] State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology,

Chinese Academy of Sciences, Kunming Yunnan, 650223 China

[7] Center for Excellence in Animal Evolution and Genetics, Chinese Academy of Sciences,

Kunming Yunnan, 650223 China

* Joint first authors

[†] Correspondence authors: tabitha.innocent@bio.ku.dk, m.hutchings@uea.ac.uk,

jjboomsma@bio.ku.dk, dougwyu@mac.com

[††] Current address: Division of Host-Microbe Systems & Therapeutics, Department of Pediatrics,

University of California, San Diego, La Jolla, California 92093, USA

Keywords: Attini, game theory, microbiome, mutualism, Actinobacteria, partner choice, leaf-cutting

ants, *Pseudonocardia*, horizontal transmission, symbiosis

Article type: Letters.

Word count: Abstract = 148; Main text = 4767; Box 1 = 244

Number of references: 51

Number of Figures, Tables, Boxes: 3, 1 and 1 respectively

Data accessibility statement: The R scripts and data for analyses will be archived at

github.com/dougwyu/screening_Innocent_et_al.

Author declaration: DY, IS, JJB, MH conceived the research, and DY, MH, JJB, TI, NH, MAB, BW

and IS designed the experiments. TI, MAB and MS isolated strains. TI and NH carried out the

experiments. DY performed the statistical analysis. DY, JJB, TI and NH wrote the manuscript, with

comments from all other authors.

Abstract

Many animals and plants recruit beneficial microbes from the environment, enhancing

their defence against pathogens. However, we have only a limited understanding of

the assembly mechanisms involved. A game-theoretical concept from economics,

screening, potentially explains how a host can selectively recruit antibiotic-producing

microbes from the environment, by fomenting and biasing competition among

potential symbionts in such a way that the likely winners are mutualists. The cuticular

microbiomes of *Acromyrmex* leaf-cutting ants inspired one of the first applications of

screening theory to mutualisms, and here we use inoculation experiments to test the

efficacy of screening *in vitro.* Using agar infused with antibacterial metabolites from

the ants' vertically transmitted *Pseudonocardia* symbionts, we show that secondary

antibiotic-producing bacteria have higher growth rates than do non-producer strains

and are more likely to win in direct competition. Our results demonstrate how game-

theoretical concepts from economics can provide powerful insight into host-

microbiome coevolution.

Introduction

Many, perhaps most, animal and plant species recruit multiple strains of beneficial

microbes from the environment, with one of the most common benefits being defence

against pathogens (Kaltenpoth 2009; Barke *et al.* 2011; Seipke *et al.* 2012; Antwis *et*

al. 2015; Loudon *et al.* 2016; Duarte *et al.* 2018; Engl *et al.* 2018). It is widely

expected that diverse microbiomes provide a more reliable defence against pathogens

than do single-species microbiomes, by allowing the equivalent of multi-drug therapy

(reviews in Scheuring & Yu 2012; Seipke *et al.* 2012; Antwis & Harrison 2018;

Duarte *et al.* 2018; Engl *et al.* 2018). However, despite an abundance of recent

studies, two recent reviews still conclude that "our ability to make predictions about

these dynamic, highly complex communities is limited" (Hoye & Fenton 2018), and

that "integration between theory and experiments is a crucial 'missing link' in current

microbial ecology" (Widder *et al.* 2016).

The Problem of Hidden Information. – We have previously shown that a game-

theoretical approach provides a powerful shortcut to modelling the coevolution of

host-microbiome assemblies. Specifically, screening theory from economics (Archetti

et al. 2011a; b; Scheuring & Yu 2012) conceptualises symbioses as a Principal-Agent

game (Edwards *et al.* 2006; Weyl *et al.* 2010), in which a Principal (host) tries to

recruit one or more suitable Agents (symbionts) out of all possible Agents (e.g. all

microbes in the environment), some or most of which are unsuitable. The challenge is

that the Principal does not know which Agents are suitable – that is, *the Agent's*

characteristics are hidden. This is the Problem of Hidden Information. The equivalent

statement in evolutionary biology is that in horizontally acquired symbioses, a host is

under selection to recruit mutualistic symbionts out of a species pool that includes

both commensals and parasites, but the host is faced with the Partner Choice problem

of not being able to recognise which species are mutualistic (Bull & Rice 1991).

Principal-Agent theory provides two solutions to the Problem of Hidden Information:

signalling and *screening* (Archetti *et al.* 2011a; b). Honest signalling uses the display

of costly phenotypes to reveal the quality of potential partners (Spence 1973; Grafen

1990), but it is difficult to envision a cost that can signal cooperativeness *per se* (but

see Archetti et al. 2016). Also, even if the host is able to discern symbiont qualities,

the host might be unable to actively choose amongst the symbionts. Both problems

apply to the recruitment of defensive symbionts where the host cannot use *signalling*

because the host has no mechanism for actively choosing amongst symbiont lineages,

cannot identify their characteristics before establishment, or both (Scheuring & Yu

2012).

Here, we present a set of *in vitro* experiments to demonstrate that ant hosts can

instead use *screening* to selectively take up antibiotic-producing microbes from the

pool of bacteria in the environment. First, we recap how screening works.

Screening. – Imagine a host faced with multiple potential symbionts differing in their

benefit to the host. For simplicity, we assume two types, mutualistic and parasitic,

where the latter includes commensals because they can impose opportunity costs on

hosts (Yu 2001). It is possible for the host to selectively 'screen-in' mutualistic

symbionts, provided that the host evolves a 'demanding environment', which imposes

a cost on colonising symbionts that is easier to endure if they are mutualists. The

benefit of enduring the cost is a host-provided reward that is high enough for the

mutualist to enjoy a net benefit (see Box 1 for a human example).

Screening succeeds if the nature and size of the cost and benefit evolve so that only

the mutualist reaps a sufficiently large net benefit for natural selection to reward

colonisation of the host over any alternatives, including free living. Multiple

screening mechanisms appear to exist in nature (Archetti *et al.* 2011a; b; see also

Ranger *et al.* 2018). In particular, we have proposed that hosts evolve to foment

competition among potential symbionts in such a way that the winners have a high

probability of being mutualists, which we call *competition-based screening*. A good

example is given by Heil (2013), who studied *Acacia* ant-plants, which can be

colonised by ant species that either protect their hostplants (mutualists) or not

(parasites). *Acacia* species that provide high amounts of food and housing to incipient

ant colonies promote colony growth and worker activity, and the colony that wins the

hostplant is the one whose workers kill off the other incipient colonies. Having

numerous, aggressive workers is also the defining characteristic of a mutualistic

colony, since the workers attack herbivores. Thus, the demanding environment is

interference competition amongst ants, fueled by the hostplant, and the ant species

that are best able to endure this cost are the aggressive ones, which is correlated with

their mutualistic service to the plant. The benefit of enduring the demanding

environment is the high levels of plant-provided reward. This benefit, minus the cost

of the risk of colony death due to fighting, is presumably greater than the alternative

fitness from living in other microhabitats. What makes this an especially satisfying

example is that there also exist *Acacia* species that provide low amounts of food and

housing, and these species are regularly colonised by parasitic, non-defending ant

colonies, whereas high-reward *Acacia* species are rarely colonised by parasitic ants.

Box 1. A human example of a successful screening mechanism

Any automobile-breakdown insurance company is faced with a *hidden information*

problem. Potential clients differ in the probability that their cars will need rescue.

Owners of poor-quality cars are more willing to purchase insurance but will impose

higher costs on insurers with their more frequent callouts. Owners of high-quality cars

are less willing to pay for insurance but would be more profitable to insurers. The

challenge for insurers is to find a way to charge a higher price to owners of poor-

quality cars and a lower price to owners of high-quality cars, without needing to

inspect the cars.

One solution can be viewed at www.rac.co.uk/breakdown-cover (accessed 30 May

2018). Adding the 'At home cover' rescue option costs an additional £5/month, nearly

doubling the cost of the cheapest cover at £5.50/month. Poor-quality cars run a

nontrivial risk of failing to start in the morning. For high-quality cars, this risk is

negligible. In screening terms, the absence of 'At Home' recovery is *a burden that*

owners of high-quality cars are better able to endure. If priced and designed

correctly, the two types of owners will voluntarily sign up for the two different

coverage levels.

A similar design challenge applies to costly, honest signalling, in which the signal has

to be costly in a way that reveals a specific hidden quality. For instance, an expensive

car is good at signalling wealth and poor at signalling fidelity or niceness.

Attine defensive microbiomes. – We have argued that competition-based screening is

consistent with several biological details of the defensive cuticular microbiomes of

attine fungus-growing ants (Scheuring & Yu 2012). These microbiomes produce

antibiotics that can control specialised crop diseases, in particular mycopathogens of

the genus *Escovopsis* (Currie *et al.* 2006), and may also provide brood hygienic

protection (Mattoso *et al.* 2012). They are dominated by Actinobacteria, which are

renowned as antibiotic producers (Kaltenpoth 2009; Barka *et al.* 2016; Worsley *et al.*

2018), but only species of the actinobacterial genus *Pseudonocardia* are documented

to be vertically transmitted and sufficiently well-studied to qualify as a co-evolving

symbiont of attine ants (Cafaro *et al.* 2011; Andersen *et al.* 2013). Other detected

actinobacterial strains (a less restrictive term used in preference to 'species' in

microbiology) might also be resident but are horizontally acquired, or alternatively

might be transient and provide no benefits. Although the relevance of these two

transmission routes is not in doubt, their relative importance has been extensively

debated without consensus (Sen *et al.* 2009; Cafaro *et al.* 2011; Mueller 2012).

Scheuring and Yu's (2012) model tried to resolve this debate by pointing out that the

Pseudonocardia biofilm coats the worker surface in antibacterial compounds,

rendering this a demanding environment that we expect antibiotic-resistant strains to

be best able to endure. This is because antibiotic-producing bacteria must also have

antibiotic resistance, or they would commit suicide when producing antibacterial

compounds. Thus, we expect the quality that allows potential symbionts to endure this

demanding environment (resistance to antibacterials) to be strongly correlated with

the quality that makes them mutualistic (antibiotic production). Attine biology

supports the assumption that the demanding environment is paired with high rewards,

because the genus-level lineages that host actinobacterial biofilms appear to feed them

with subcuticular exocrine glands (Currie *et al.* 2006) – although the secretions of

these glands have not yet been characterised.

Screening can potentially explain two other aspects of attine ant biology. Firstly, in

the crown-group genus *Acromyrmex*, newly hatched workers are inoculated with

Pseudonocardia by their sister workers (Poulsen *et al.* 2003b; Andersen *et al.* 2013;

Marsh *et al.* 2014), and the monoculture biofilm subsequently blooms over almost the

entire worker surface, then retreats to the ant's ventral chest plates after ca. 25 days,

when the workers start to forage outside the nest (Poulsen *et al.* 2003a; Andersen *et*

al. 2013). At this time it becomes possible to detect the presence of other bacteria on

the ant's surface, many of which are also Actinobacteria (Poulsen *et al.* 2005;

Andersen *et al.* 2013).

Secondly, Andersen *et al.* (2013) have documented that *Acromyrmex* colonies hosting

one of the native *Pseudonocardia* species (*P. octospinosus*, also known as 'Ps1') are

colonised by a more diverse secondary cuticular microbiome than are *Acromyrmex*

colonies hosting the other species (*P. echinatior*, 'Ps2') (species described in Holmes

et al. 2016). For simplicity, we refer to these as Ps1 and Ps2. Screening theory

predicts that Ps2 imposes the more demanding environment for secondary invaders,

by making stronger or more antibacterials. This would reduce the likelihood of

successful colonisation by competitors, but may make the mutualist more costly for

the host that provides the resources for its growth. Genomic analysis has confirmed

that Ps1 and Ps2 have different sets of secondary metabolite clusters and thus likely

produce different spectra of antimicrobial compounds (Holmes *et al.* 2016).

Tests of screening. – Experimental tests of mechanisms that could promote selective

recruitment are missing from the microbiome literature (Widder *et al.* 2016; Hoye &

Fenton 2018). We address this gap with an *in vitro* model emulating the dynamics on

the cuticle of *Acromyrmex* ants. Using *Pseudonocardia* cultures, we created two types

of environment on agar media: *demanding* (where growth media are infused with

secondary metabolites from *Pseudonocardia*), and *undemanding* (where symbiont

metabolites were absent). We then introduced a selection of different bacterial strains

representing two classes of potential colonisers from the environment – a set of

antibiotic-producing Actinobacteria (*producers*) and a set of bacteria that do not

produce antibiotics (*non-producers*) – and we tested the prediction that *producers* are

better invaders (by measuring growth rates in the demanding environment) than are

non-producers. We also staged direct-competition experiments between *producers*

and *non-producers* on both *demanding* and *undemanding* media.

We show that antibiotic-*producers* grow relatively more quickly on *demanding*

(toxic) media despite growing relatively more slowly on *undemanding* (non-toxic)

media, as predicted by screening theory. This result also parsimoniously explains the

outcome of the direct-competition experiment, in which antibiotic-*producers* are

superior competitors on toxic media, while *non-producers* are superior competitors on

non-toxic media. Finally, we also confirm the assumption that antibiotic-*producers*

are more resistant to antibiotics and the prediction that *Pseudonocardia* Ps2 strains

are stronger growth-suppressors of other bacteria.

Materials & Methods

We obtained:

(**1**) a set of *Pseudonocardia* Ps1 and Ps2 strains to grow on agar plates, after which

we flipped the agar to expose a surface that was open for colonisation but infused

with *Pseudonocardia* metabolites. We and others have shown that *Pseudonocardia*

releases antibacterials and antifungals when grown in culture (Barke *et al.* 2010;

Holmes *et al.* 2016; Van Arnam *et al.* 2016);

(**2**) a set of 'environmental' *antibiotic-producing* bacterial strains, defined as not

having an evolutionary history of growing on the cuticle of *Acromyrmex* ants;

(**3**) a set of 'environmental' *non-antibiotic-producing* bacterial strains, defined as not

having an evolutionary history of growing on the cuticle of *Acromyrmex* ants; and

(**4**) a set of bacterial strains that reside in the cuticular microbiomes of *Acromyrmex*

workers while being *non-antibiotic-producing*, which in theory should preclude their

survival in cuticular biofims. We asked why these bacteria do not competitively

exclude the resident antibiotic-*producers*, since they are presumably resistant to

Pseudonocardia metabolites but do not pay the cost of producing antibiotics. We use

the term 'resident' only to indicate that we isolated these strains from *Acromyrmex*

cuticles; we do not imply that these strains are true native residents like

Pseudonocardia.

For bacterial growth media, we used Soya Flour + Mannitol (SFM) agar (20g soya

flour, 20g mannitol, 20g agar in 1L tap water) and Lysogeny Broth (LB) Lennox (10g

tryptone, 5g yeast extract, 5g NaCl in 1L deionised water).

Collection and isolation of bacterial strains

(**1**) Nineteen Ps1 and Ps2 strains were isolated from the cuticles of individual workers

of the leaf-cutting ant *Acromyrmex echinatior* (Hymenoptera, Formicidae, Attini).

The 19 colonies of these workers were collected in Soberania National Park, Panama,

between 2001 and 2014, and reared in climate-controlled rooms at the University of

Copenhagen at ca. 25 ºC and 70% relative humidity. The propleural chest plates of

individual ants – where the white *Pseudonocardia* biofilms grows most conspicuously

(Poulsen *et al.* 2003b; Currie *et al.* 2006) – were scraped with a sterile needle under a

stereomicroscope and immediately streak-inoculated across Lennox (LB) media under

sterile conditions. The plates were then incubated at 30 °C until visible growth of

distinctive white 'cotton ball' colonies of *Pseudonocardia* were clearly apparent.

These individual colonies were removed under sterile (LAF bench) conditions and

inoculated onto fresh plates, a process that was repeated until clean *Pseudonocardia*

cultures were isolated. The *Pseudonocardia* isolates were then grown on Mannitol

Soya Flour (SFM) media (optimal for actinobacterial growth and spore production),

from which spore stock solutions were prepared in 20% glycerol and kept at -20 °C

until use. Each isolate was genotyped to Ps1 or Ps2 (Supplementary Information S1).

Five strains from each of these two species have previously been genome-sequenced,

formally described, and functionally characterised as different *Pseudonocardia*

species by Holmes et al. (2016): *P. octospinosus* (Ps1) and *P. echinatior* (Ps2).

(**2**) The 10 environmental antibiotic-producing strains (all *Streptomyces*) were taken

from general collections in the Hutchings lab (Tables 1, S2).

(**3**) The 10 environmental non-antibiotic-producing strains were obtained from the

Hutchings lab (2 strains) and from the ESKAPE suite (8 strains) – a set with varying

origins (human skin, soil, etc.) associated with hospital-acquired infections and used

to test antibiotic resistance or efficacy in clinical/research settings (Tables 1, S2).

(**4**) The 10 *Acromyrmex*-resident, non-antibiotic-producing strains were isolated from

A. echinatior and *A. octospinosus* ants by culturing from cuticular washes or from the

Pseudonocardia isolations above. These strains were 16S-genotyped and where

possible assigned to genus level by comparison to the NCBI 16S RefSeq database

(Supplementary Information S1, Tables 1, S2).

275   **Table 1.** The bacterial strains used in our experiments: the 19 *Pseudonocardia* strains isolated from

276   *Acromyrmex echinatior* (columns 1 and 2); the 10 environmental *antibiotic-producer* strains (column 3);

277   the 10 environmental *antibiotic-non-producer* strains (column 4); and the 10 *Acromyrmex-resident,*

278   *antibiotic- non-producer* strains (column 5). Details for all strains not isolated from *Acromyrmex* in Table

279   S2.

8 *Pseudonocardia* Ps1 strains (colony of origin)	11 *Pseudonocardia* Ps2 strains (colony of origin)	10 environmental antibiotic-producing strains (all *Streptomyces*)	10 environmental non-producer strains	10 *Acromyrmex*-resident non-producer strains
Ae356*	Ae406*	S1. *Streptomyces* M1146	St1. *Escherichia coli*	Sr1. *Ochrobactrum sp.*
Ae263*	Ae160	S2. *S. lividans*	St2. *Lysobacter antibioticus*	Sr2. *Erwinia* sp.
Ae322	Ae505*	S3. *S. coelicolor*	St3. *Bacillus subtilis*	Sr3. *Acinetobacter* sp.
Ae150a*	Ae717*	S4. *S. scabies*	St4. *Pseudomonas putida*	Sr4. *Sphingobacterium sp.*
Ae168*	Ae703	S5. *S. venezuelae*	St5. *Erwinia caratova*	Sr5. *Acinetobacter* sp.
Ae707-CP-A2*	Ae702	S6. *S.* Ae150A-B1	St6. *Enterobacter aerogenes*	Sr6. *Luteibacter sp.*
Ae712†	Ae707	S7. *S.* Ae356-S1	St7. *Acinetobacter baylyi*	Sr7. *Flavobacterium sp.*
Ae280†	Ae331*	S8. *S.* KY5	St8. *Staphylococcus epidermidis*	Sr8. *Brevundimonas sp.*
-	Ae706*	S9. *S.* S4	St9. *Micrococcus luteus*	Sr9. *Acinetobacter sp.*
-	Ae704	S10. *S.* Amy1	St10. *Serratia* KY15	Sr10. *Brachybacterium sp.*
-	Ae715	-	-	-

* *Pseudonocardia* strains that have been genome-sequenced (Holmes *et al.* 2016)
† *Pseudonocardia* strains that were only used in the growth-rate experiment with *Acromyrmex*-resident *non-producers*.

Individual growth-rate experiments

We measured bacterial colony growth rates by growing all *producer* and *non-*

producer strains on antibiotic-infused and control media.

To create the antibiotic-infused media, we grew lawns of the 19 *Pseudonocardia*

isolates (Table 1), plating 30 µl on 90 mm SFM agar plates. The control plates were

inoculated with glycerol only (20%; identical to that used for making solutions from

Pseudonocardia strains). We incubated these plates at 30 °C for 6 weeks, which

ultimately produced lawns from 17 strains that could be included in the experiments

(6 Ps1 and 11 Ps2 strains). Once each plate was fully (not necessarily uniformly)

covered, we flipped the agar to reveal a surface open for colonisation.

The sets of 10 environmental antibiotic-*producers* (all *Streptomyces* spp.) and of 10

environmental *non-producers* (Table 1) were inoculated onto the different media

types. Each agar plate was inoculated with 10 evenly spaced colonies, with 3

replicates, generating 2 invader types x 17 Ps-media-types x 3 replicates = 102 Ps1/2-

infused plates and 2 x 3 = 6 control plates, i.e. a total of 1020 treatment inoculations

and 60 control inoculations. Each strain inoculation used 5 μl of solution (at

approximately 1 x 10^6 cells per ml in 20% glycerol), spotted at evenly spaced

positions and without coming into direct contact. All plates were incubated at 30 °C

for five days, after which photographs were taken under sterile conditions.

Images were processed in Fiji software (Schindelin *et al.* 2012; Rueden *et al.* 2017),

creating binary negatives (black & white) so automated tools could identify discrete

areas of growth (regions in black) and measure growth areas for each invading strain;

in the few cases where binary image resolution was insufficient, outlines were added

manually before area calculation. 48 *producer*-inoculated and 57 *non-producer*-

inoculated treatment measurements were excluded because plate condition had

deteriorated to become unscorable or they were contaminated, leaving a final sample

size of 1020-48-57= 915 treatment inoculations and 60 controls.

The second growth-rate experiment compared the 10 *Acromyrmex*-resident, *non-*

producer strains with 9 of the environmental *producer* strains. All 19 *Pseudonocardia*

strains grew sufficiently to be included in this experiment, and each plate was again

inoculated with 10 or 9 evenly spaced colonies. Starting sample sizes were therefore 2

invader types x 19 Ps-media-types x 3 replicates = 114 Ps1/2-infused plates and 2 x 3

= 6 control plates, giving a total of 1083 treatment inoculations and 57 controls. 50

producer and 20 *non-producer* treatment measurements were excluded for the same

reasons as above, giving a final sample of 1083-50-20=1013 treatment inoculations

and 57 controls for scoring as described above.

Pairwise competition experiment

Building on the above experiment, where *producer* strains had relatively higher

growth rates on *Pseudonocardia*-infused media, we tested whether this growth

advantage (plus any direct interference interactions) could result in competitive

exclusion of *non-producers*. We co-inoculated pairs of environmental *producers* and

non-producers on both *Pseudonocardia*-infused media and control media (prepared as

in the individual growth-rate experiment), and we measured the outcome of

competition as a win, loss, or draw. We predicted that the *producer* strains would be

more likely to win when competing on *Pseudonocardia*-infused media.

To keep the number of tests manageable, we used only two combinations of

Pseudonocardia and *Streptomyces*: *Pseudonocardia* Ps1 (strain Ae707) +

Streptomyces S8 and *Pseudonocardia* Ps2 (strain Ae717) + *Streptomyces* S2 (Table

1). These combinations are representative of the results from the growth-rate

experiment: the two *Streptomyces* strains grew more slowly than most of the *non-*

producers on control media and either near the median growth rate of (S8) or faster

than (S2) the *non-producers* on *Pseudonocardia*-infused media.

We competed *Streptomyces* S8 and S2 against the 10 environmental *non-producer*

strains (20 pairings), using spore titrations of S8 and S2 consisting of 10^6 spores per

ml for each strain in 20% glycerol. Non-producers were grown overnight in 10 ml

LB-Lennox, subcultured with a 1:100 dilution into a fresh 10 ml LB-Lennox, grown

at 37 °C for 3-4 hours, after which OD600 was measured, assuming that OD600 = 1

represented 8×10^9 cells. Similar dilutions of 10^6 cells per ml were made for each

non-producer strain in 20% glycerol, after which *producer* and *non-producer*

dilutions were mixed 1:1. Each pair of *producer* and *non-producer* was co-inoculated

as a mixture of 20 µl (10^4 spore-cells of each) on the designated *Pseudonocardia*-

infused media (set up as above) with 5 replicates per pairing. We used 150 plates for

the S8 experiment (including 100 control plates; 10 replicates per pairing) and 100

plates for the S2 experiment (including 50 control plates; 5 replicates per pairing).

Plates were incubated at 30 °C for 5 days and then photographed, after which images

were scored with respect to the *producer* as: win (dominant growth), draw (both

strains growing with no clear dominant), or loss (little or no visible growth), with

reference to images of each strain grown alone on control media to minimise observer

bias. One plate's outcome was too ambiguous to score and was omitted. All plates

were independently scored by two observers, one using only photos, which produced

data sets giving the same statistical results. We report the results from the observer

who scored from direct observation.

Antibiotic-resistance profiles

The key assumption of screening theory is that antibiotic-*producers* are better at

resisting antibiotics, as measured by growth rates in the presence of antibiotics,

because this correlation is what allows *producer* strains to better endure the

demanding environment produced by *Pseudonocardia*. We tested this assumption by

growing the 10 environmental *producer* strains and the 10 environmental *non-*

producer strains (Table 1, S2) in the presence of 8 different antibiotics, representing a

range of chemical classes and modes of action. Antibiotics were added to 1 ml of LB-

Lennox medium in a 24-well microtitre plate at 6 different concentrations. The

relative concentration range was the same for each antibiotic, although actual

concentrations reflected activity (Table S3). *Producers* and *non-producers* were

inoculated onto plates and incubated at 30 °C for 7 days, then photographed (Table

S3). Lowest Effective Concentration (LEC, lowest concentration with inhibitory

effect) and Minimum Inhibitory Concentration (MIC, lowest concentration with no

growth) scores were assigned on a Likert scale of 1–6, where 1 was no resistance and

6 was resistance above the concentrations tested (adapted from generalised MIC

methods; reviewed by Balouiri *et al.* 2016).

Statistical analyses

Initial models revealed correlated residuals by *Pseudonocardia* strain media and by

inoculant strain, so we used mixed-effects models to incorporate strain identities as

random intercepts, and we nested plate replicate within the *Pseudonocardia* strains.

Statistical significance was determined by term deletion, and final-model residuals

were approximately homoscedastic and uncorrelated within random factor levels.

Analyses were carried out in *R* 3.4.4 (R Core Team 2017) with *lme4* 1.1-17 (Bates *et*

al. 2015), *tidyverse* 1.2.1 (Wickham 2017), and *RColorBrewer* 1.1-2

(www.colorbrewer.org, accessed 4 Mar 2018). R scripts and data are available in

Supplementary Information (S4) and at

github.com/dougwyu/screening_Innocent_et_al.

Results

Individual growth-rate experiments. – As predicted by the screening model, *non-*

producers grew more quickly on undemanding control media and *producers* had

higher growth rates on the demanding *Pseudonocardia*-infused media, producing a

highly significant statistical interaction effect (Fig. 1A). There was also a significant

main effect of *Pseudonocardia* genotype, with both *non-producers* and *producers*

exhibiting a lower growth rate on Ps2-infused media, relative to Ps1-infused media,

consistent with Andersen et al.'s (2013) observation that *Acromyrmex* colonies

hosting Ps2-dominated cuticular microbiomes were less prone to secondary invasion

by other bacteria.

The resident *non-producers* isolated from cuticular microbiomes had significantly

slower growth rates overall, even on control media without antibiotics (Fig. 1B). This

suggests that they are unable to outcompete *producer* strains on the cuticle of

Acromyrmex ants and raises the question why these *non-producer* species can persist

at all.

A. Growth rates of environmental *non-producers* vs. *producers*

B. Growth rates of *Acromyrmex* resident *non-producers* vs. *producers*

Figure 1. Individual growth-rate experiments. Bacterial colony sizes after 5 days at 30 ºC,

with the boxplots indicating medians ± one quartile. The white section shows growth rates

on control media, and the grey section shows growth rates on the *Pseudonocardia*-infused

media. Coral boxes represent *non-producer* strains, and blue boxes represent *producer*

strains. For analysis, a linear mixed-effects model, including *Pseudonocardia* strain

(`Ps.strain`, 17 groups) and inoculated bacterial species (`Inv.strain`, 20 groups) as

random intercepts, was used to test for interaction and main effects of growth media

(`Control` vs. `Ps1.infused` vs. `Ps2.infused`) and antibiotic production

(`Non.producers` vs. `Streptomyces`): (*lme4::lmer*(Growth.score ~ Media *

Antibiotic.production + (1|Ps.strain/Plate) + (1|Inv.strain)). **A.**

Environmental strains. There was a highly significant interaction effect, such that

environmental non-producer strains grow more rapidly on control media, while

Streptomyces grow more rapidly on *Pseudonocardia*-infused media (n = 975, χ^2 = 45.86, df =

2, p < 0.0001). There was also a highly significant main effect of *Pseudonocardia* genotype,

with growth being lower on Ps2-infused media (n = 915, χ^2 = 24.55, df = 1, p < 0.0001,

control-media data omitted for this analysis). One non-producer strain (*Staphylococcus*

epidermidis) grew more rapidly than all other strains (open triangle points), demonstrating

the need to control for correlated residuals by strain. The y-axis has been truncated at 2.5

for clarity (the full figure is in Supplementary Information). **B.** *Acromyrmex*-resident, *non-*

producer strains. There was no interaction effect (χ^2 = 2.64, df = 2, p = 0.27), but both main

effects were highly significant. The *non-producers* grow more slowly on all media types (χ^2 =

20.96, df = 1, p < 0.0001), and bacterial growth was generally slower on Ps2-infused media

(χ^2 = 21.43, df = 1, p < 0.0001, control-media data omitted).

Pairwise competition experiment. – The two *producer* strains were much more likely

to win in direct competition against the *non-producer* strains when grown on the toxic

(*Pseudonocardia*-infused) media (Fig. 2), which demonstrates that growth rates in

isolation (Fig. 1A) are sufficient predictors of success in competitive interactions,

although other interactions, such as antibiotic production by the *Streptomyces,* likely

also contribute. These results support the screening model prediction that a

demanding environment (facilitated by a host) can determine the profile of successful

colonisers of the cuticular microbiome.

Figure 2. Pairwise competition experiment, scoring whether or not *producers* prevail. **A**.

Representative images of agar plates at five days post-inoculation showing examples of the

three competitive outcomes: Win (S2 vs. St3 on Ps2 media), Draw (S2 vs. St3 on control

media), and Loss (S8 vs. St3 on control media; strain details in Table 1). **B**. Barcharts of

competitive outcomes for the two *Streptomyces producer* strains (S8 and S2; Table 1), each

pairwise co-inoculated with ten non-antibiotic-producer strains on *Pseudonocardia*-infused

media and on control media. For analysis, draw outcomes were omitted, and a general

linear mixed-effects model, including non-antibiotic-producer strain as a random intercept

(10 groups), was used to test for an effect of the `medium` term (`Control` vs. `Ps1/2-`

`infused`) on competitive outcome (`Win` vs. `Loss`) ((*lme4::glmer*(outcome ~

medium + (1|non.producer.strain), family = binomial)). Significance was

estimated using term deletion. In both experiments, *producer* strains (*Streptomyces*) were

more likely to be competitively superior when grown on the *Pseudonocardia*-infused media

(Left: n = 129, χ^2 = 103.6, df = 1, p < 0.0001; Right: n = 94, χ^2 = 87.9, df = 1, p < 0.0001).

The screening model assumes that *producer* strains have greater resistance to

antibiotics than do *non-producer* strains. We confirmed this assumption for this

experiment because both the Lowest Effective Concentration (LEC, lowest

concentration with inhibitory effect) and the Minimum Inhibitory Concentration

(MIC, lowest concentration with no growth) were greater for the *producer* strains

(Fig. 3). We also observed that the *resident non-producers* had high levels of

antibiotic resistance (Fig. 3), as expected (Fig. 1B).

Figure 3. Antibiotic resistance profiles for *producer*, *non-producer*, and *resident non-producer* strains (Table 1). Boxplots indicate medians (notches) ± one quartile. For analysis, we calculated each strain's mean growth score across the eight tested antibiotics (reducing from n = 155 to n = 20; details in S3). *Producers* showed higher levels of resistance than did *non-producers* for both measurements: Wilcoxon two-sided test (`wilcox.test`), W = 94.5, p = 0.0017 for LEC (**A**) and W = 80, p = 0.0253 for MIC (**B**), after correction for two tests (*p.adjust*(`method="fdr"`)). *Producers* and *Resident Non-producers* showed no difference in resistance levels (p = 0.44 and 0.25).

Discussion

We used *in vitro* experiments to test key predictions and assumptions of a screening model (Scheuring & Yu 2012) that was designed to explain how leaf-cutting ants could use a vertically transmitted symbiont, *Pseudonocardia*, to selectively recruit

additional antibiotic-producing bacterial strains from the large pool of potential

bacterial symbionts present in the environment.

We showed that *non-producer* invasive strains grow more slowly than do antibiotic-

producer strains on *Pseudonocardia*-infused media (Fig. 1A). Since the control plates

showed that *non-producers* have higher intrinsic growth rates, we attribute the

restricted growth on *Pseudonocardia*-infused plates to inhibition by *Pseudonocardia*

metabolites, consistent with the large number of antibacterials documented as

encoded in the genomes of both Ps1 and Ps2 (Holmes *et al.* 2016). We also showed

that Ps2 is a stronger suppressor of bacterial growth in general, consistent with

Andersen et al.'s (2013) observation that Ps2-hosting *Acromyrmex* colonies assemble

lower-diversity (and more Ps2-dominated) cuticular biofilms on foraging workers.

Finally, we confirmed that our *producer* strains (all genus *Streptomyces*) had higher

resistance to antibiotics than did our *non-producer* strains (Fig. 3).

Our pairwise competition experiment confirmed that growing on *Pseudonocardia*-

infused media renders *producer* strains competitively superior (i.e. they usually win)

or at least resistant to exclusion (they more often draw than lose, especially on the

less-toxic Ps1-infused media) (Fig. 2). This *in vitro* experiment mimics the conditions

we expect to occur on the cuticle of *Acromyrmex* workers after their full-body

Pseudonocardia biofilm (obtained shortly after hatching) has retreated to the

laterocervical chest plates (and possibly into the subcuticular glands), when the

workers begin to forage outside the nest, where they are exposed to many bacterial

colonists (Poulsen *et al.* 2005; Andersen *et al.* 2013; Marsh *et al.* 2014; Andersen *et*

al. 2015).

Taken together, these results show how a single, vertically transmitted, antibiotic-

producing symbiont such as *Pseudonocardia* can be used by the ant host to create a

demanding environment on the ant surface that favours the recruitment of a more

diverse set of antibiotic-producers. In short, vertical transmission of one mutualist

enables the successful horizontal acquisition of other mutualists. Vertical transmission

and horizontal symbiont acquisition are therefore not mutually exclusive

explanations, consistent with a mixture of both not being uncommon in other

symbioses (Ebert 2013; Bourguignon *et al.* 2018; da Costa & Poulsen 2018; Ivens *et*

al. 2018). Under some circumstances, mutualistic stability through screening may be

able to do without a vertically transmitted symbiont; Duarte et al. (2018) have argued

that a host might be able to create a suitably demanding environment by evolving the

right set of chemical exudates.

We note that in a colonial social insect, one needs to be precise about symbiont

transmission and symbiont inheritance. Since virgin *Acromyrmex* queens are

inoculated upon hatching by nurse workers that maintain monocultures of either Ps1

or Ps2, the secondary acquisition of other *producer* symbionts by foraging workers is

highly unlikely to disrupt verticality in the inheritance of native *Pseudonocardia*

symbionts down the generations.

Acromyrmex-*resident non-producers*. – We also ran a growth-rate experiment with a

set of *resident non-producer* strains that had been isolated from *Acromyrmex* ant

cuticles (Fig. 1B) and found that they generally had low intrinsic growth rates,

making it unlikely that they can competitively exclude *producers*. It seems surprising

that we found these *non-producer* strains at all, but we note that the fact that we could

isolate them does not imply that they are abundant. Resident *non-producer* taxa are

only found on some (not all) field-sampled *Acromyrmex* workers (Andersen *et al.*

2013; Innocent et al., in prep) and generally below 1% prevalence when measured by

amplicon read number. They may only be transient environmental contaminants,

since many were isolated via whole-ant washes, or strains persisting in non-

equilibrium dynamics. In simulation models (Boza, G., Worsley, S., Yu, D.,

Scheuring, I., unpubl. data), we have found that if bacterial competition is limited to

nearest neighbours and if antibiotics slowly diffuse across the ant surface, even non-

antibiotic-resistant strains can sometimes persist for long periods of time.

It remains an open question why three different *Acromyrmex* species in Panama all

host the same two *Pseudonocardia* strains, Ps1 and Ps2 (Andersen *et al.* 2013). The

genomes of Ps1 and Ps2 strains each encode a single distinct nystatin-like antifungal,

plus variable and partially segregated sets of secondary metabolite biosynthetic gene

clusters with predicted antibacterial functions (Holmes *et al.* 2016). These two

Pseudonocardia species are almost never found in the same colony even though they

are encountered at approximately equal rates in field colonies (Poulsen *et al.* 2005).

The difference between Ps1 and Ps2 in microbiome invasibility (Fig. 1A) may reflect

a tradeoff between stronger endogenous production of antimicrobials early in life

against the recruitment of a more diverse set of secondary antibiotic producers later in

life.

Comparable studies to ours using other species of fungus-growing ants will be needed

to test the generalisability of screening to other attine ant lineages that maintain

actinobacterial biofilms, including species that appear to have cuticular microbiota

with abundant actinomycete genera other than *Pseudonocardia* (Sen *et al.* 2009;

Andersen *et al.* 2013; Meirelles *et al.* 2014). Our results suggest that cuticular

biofilms can be adequately simulated on agar plates, providing a tractable

experimental model with which to test microbiome assembly mechanisms. Such

studies might also shed light on why *Atta* and some other genus-level branches of the

attine fungus-growing ants have secondarily lost their cuticular biofilms, a

phenomenon that remains poorly understood. The screening requirement to pair a

demanding environment with a high reward imposes nontrivial costs on hosts, and in

some cases, this cost might have tipped the cost-benefit ratio in favour of chemical

pest control over biological control (Fernandez-Marin *et al.* 2009; Fernandez-Marin *et*

al. 2015).

Toward a general game theory of hosted microbiomes

Hosted microbiomes are coevolutionary metacommunities in which most of the

species interactions are difficult to observe. There are numerous calls to use

ecological theory to understand microbiomes (Widder *et al.* 2016; Hoye & Fenton

2018), but we propose that a game theory approach may be more promising for

producing testable predictions. Screening (Archetti *et al.* 2011a) and password

signalling (Archetti *et al.* 2016) are candidate solutions to the Problem of Hidden

Information. When hosts evolve to use screening, the emergence of demanding

environments can be thought of as an exercise in applied ecology, where the host

evolves a mechanism to foment and bias competition by modifying resource and

toxicity levels to encourage cooperative microbial communities (see Foster *et al.*

2017; Duarte *et al.* 2018 for similar perspectives).

Acknowledgments

The work was supported by the European Research Council (ERC Advanced grant 323085 to

JJB), a Marie Curie Individual European Fellowship (IEF grant 627949 to TI), NERC grants

NE/J01074X/1 to MH and DY and NE/M015033/1 and NE/M014657/1 to MH, DY and BW. D.

W. Yu was supported by the National Natural Science Foundation of China (41661144002,

31670536, 31400470, 31500305), the Key Research Program of Frontier Sciences, CAS (

QYZDY-SSW-SMC024), the Bureau of International Cooperation project (GJHZ1754), the

Ministry of Science and Technology of China (2012FY110800), the University of East Anglia, the

State Key Laboratory of Genetic Resources and Evolution at the Kunming Institute of Zoology,

and the University of Chinese Academy of Sciences. IS is supported by National Research,

Development and Innovation Office of Hungary (GINOP 2.3.2.-15-2016-0057). We thank the

Smithsonian Tropical Research Institute for logistical help and facilities for all work in Gamboa,

Panama, and the Autoridad Nacional del Ambiente y el Mar for permission to sample and export

ants. We thank Ryan Seipke and Gergely Boza for input in developmental stages of the work;

Elaine Patrick and Sylvia Matthiesen for lab and logistics support; Michael Poulsen, Saria Otani

and Panos Sapountzis for useful technical advice and discussion. The authors declare no conflicts

of interest.

References

583   Andersen, S.B., Hansen, L.H., Sapountzis, P., Sorensen, S.J. & Boomsma, J.J. (2013).
584   Specificity and stability of the *Acromyrmex-Pseudonocardia* symbiosis. *Molecular*
585   *Ecology*, 22, 4307-4321.
586
587   Andersen, S.B., Yek, S.H., Nash, D.R. & Boomsma, J.J. (2015). Interaction
588   specificity between leaf-cutting ants and vertically transmitted *Pseudonocardia*
589   bacteria. *BMC Evolutionary Biology*, 15:27.
590
591   Antwis, R.E. & Harrison, X.A. (2018). Probiotic consortia are not uniformly effective
592   against different amphibian chytrid pathogen isolates. *Molecular Ecology*, 27, 577-
593   589.
594
595   Antwis, R.E., Preziosi, R.F., Harrison, X.A. & Garner, T.W.J. (2015). Amphibian
596   symbiotic bacteria do not show a universal ability to inhibit growth of the global
597   panzootic lineage of *Batrachochytrium dendrobatidis*. *Applied and Environmental*
598   *Microbiology*, 81, 3706-3711.
599
600   Archetti, M., Miller, J.B. & Yu, D.W. (2016). Evolution of passwords for cost-free,
601   honest signalling between symbionts and hosts. *bioRxiv*.
602
603   Archetti, M., Scheuring, I., Hoffman, M., Frederickson, M.E., Pierce, N.E. & Yu,
604   D.W. (2011a). Economic game theory for mutualism and cooperation. *Ecology*
605   *Letters*, 14, 1300-1312.
606

607 Archetti, M., Ubeda, F., Fudenberg, D., Green, J., Pierce, N.E. & Yu, D.W. (2011b).
608 Let the right one in: a microeconomic approach to partner choice in mutualisms.
609 *American Naturalist*, 177, 75-85.
610
611 Balouiri, M., Sadiki, M. & Ibnsouda, S.K. (2016). Methods for in vitro evaluating
612 antimicrobial activity: A review. *Journal of Pharmaceutical Analysis*, 6, 71-79.
613
614 Barka, E.A., Vatsa, P., Sanchez, L., Gaveau-Vaillant, N., Jacquard, C., Klenk, H.-P. *et*
615 *al.* (2016). Taxonomy, physiology, and natural products of Actinobacteria.
616 *Microbiology and Molecular Biology Reviews*, 80, 1-43.
617
618 Barke, J., Seipke, R.F., Grüschow, S., Heavens, D., Drou, N., Bibb, M.J. *et al.* (2010).
619 A mixed community of actinomycetes produce multiple antibiotics for the fungus
620 farming ant *Acromyrmex octospinosus*. *BMC Biology*, 8:109.
621
622 Barke, J., Seipke, R.F., Yu, D.W. & Hutchings, M.I. (2011). A mutualistic
623 microbiome. *Communicative & Integrative Biology*, 4, 41-43.
624
625 Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015). Fitting linear mixed-effects
626 models using lme4. *Journal Of Statistical Software* 1-48.
627
628 Bourguignon, T., Lo, N., Dietrich, C., Šobotník, J., Sidek, S., Roisin, Y. *et al.* (2018).
629 Rampant host switching shaped the termite gut microbiome. *Current Biology*, 28,
630 649-654.
631
632 Bull, J.J. & Rice, W.R. (1991). Distinguishing mechanisms for the evolution of co-
633 operation. *Journal of Theoretical Biology*, 149, 63-74.
634
635 Cafaro, M.J., Poulsen, M., Little, A.E.F., Price, S.L., Gerardo, N.M., Wong, B. *et al.*
636 (2011). Specificity in the symbiotic association between fungus-growing ants and
637 protective *Pseudonocardia* bacteria. *Proceedings of the Royal Society B-Biological*
638 *Sciences*, 278, 1814-1822.
639
640 Currie, C.R., Poulsen, M., Mendenhall, J., Boomsma, J.J. & Billen, J. (2006).
641 Coevolved crypts and exocrine glands support mutualistic bacteria in fungus-growing
642 ants. *Science*, 311, 81-83.
643
644 da Costa, R.R. & Poulsen, M. (2018). Mixed-Mode Transmission Shapes Termite Gut
645 Community Assemblies. *Trends in Microbiology*, 7, 557-559.
646
647 Duarte, A., Welch, M., Swannack, C., Wagner, J. & Kilner, R.M. (2018). Strategies
648 for managing rival bacterial communities: Lessons from burying beetles. *Journal of*
649 *Animal Ecology*, 87, 414-427.
650
651 Ebert, D. (2013). The epidemiology and evolution of symbionts with mixed-mode
652 transmission. *Annual Review of Ecology, Evolution, and Systematics*, 44, 623-643.
653
654 Edwards, D.P., Hassall, M., Sutherland, W.J. & Yu, D.W. (2006). Selection for
655 protection in an ant–plant mutualism: host sanctions, host modularity, and the

656   principal–agent game. *Proceedings of the Royal Society B: Biological Sciences*, 273,
657   595-602.
658
659   Engl, T., Kroiss, J., Kai, M., Nechitaylo, T.Y., Svatoš, A. & Kaltenpoth, M. (2018).
660   Evolutionary stability of antibiotic protection in a defensive symbiosis. *Proceedings*
661   *of the National Academy of Sciences*, 115, E2020-E2029.
662
663   Fernandez-Marin, H., Nash, D.R., Higginbotham, S., Estrada, C., van Zweden, J.S.,
664   d'Ettorre, P. *et al.* (2015). Functional role of phenylacetic acid from metapleural gland
665   secretions in controlling fungal pathogens in evolutionarily derived leaf-cutting ants.
666   *Proceedings of the Royal Society B-Biological Sciences*, 282: 20150212.
667
668   Fernandez-Marin, H., Zimmerman, J.K., Nash, D.R., Boomsma, J.J. & Wcislo, W.T.
669   (2009). Reduced biological control and enhanced chemical pest management in the
670   evolution of fungus farming in ants. *Proceedings of the Royal Society B-Biological*
671   *Sciences*, 276, 2263-2269.
672
673   Foster, K.R., Chluter, J.S., Oyte, K.Z.C. & Rakoff-Nahoum, S. (2017). The evolution
674   of the host microbiome as an ecosystem on a leash. *Nature*, 548, 43-51.
675
676   Grafen, A. (1990). Biological signals as handicaps. *Journal of Theoretical Biology*,
677   144, 517-546.
678
679   Heil, M. (2013). Let the best one stay: screening of ant defenders by Acacia host
680   plants functions independently of partner choice or host sanctions. *Journal of*
681   *Ecology*, 101, 684-688.
682
683   Holmes, N.A., Innocent, T.M., Heine, D., Al Bassam, M., Worsley, S.F., Trottmann,
684   F. *et al.* (2016). Genome analysis of two *Pseudonocardia* phylotypes associated with
685   *Acromyrmex* leafcutter ants reveals their biosynthetic potential. *Frontiers in*
686   *Microbiology*, 7:2073.
687
688   Hoye, B.J. & Fenton, A. (2018). Animal host–microbe interactions. *Journal of Animal*
689   *Ecology*, 87, 315-319.
690
691   Ivens, A.B.F., Gadau, A., Kiers, E.T. & Kronauer, D.J.C. (2018). Can social
692   partnerships influence the microbiome? Insights from ant farmers and their
693   trophobiont mutualists. *Molecular Ecology*, 27, 1898-1914.
694
695   Kaltenpoth, M. (2009). Actinobacteria as mutualists: general healthcare for insects?
696   *Trends in Microbiology*, 17, 529-535.
697
698   Loudon, A.H., Venkataraman, A., Van Treuren, W., Woodhams, D.C., Parfrey, L.W.,
699   McKenzie, V.J. *et al.* (2016). Vertebrate hosts as islands: Dynamics of selection,
700   immigration, loss, persistence, and potential function of bacteria on salamander skin.
701   *Frontiers in Microbiology*, 7:333.
702
703   Marsh, S.E., Poulsen, M., Pinto-Tomas, A. & Currie, C.R. (2014). Interaction
704   between workers during a short time window is required for bacterial symbiont
705   transmission in *Acromyrmex* leaf-cutting ants. *Plos One*, 9, e103269.

Mattoso, T.C., Moreira, D.D.O. & Samuels, R.I. (2012). Symbiotic bacteria on the cuticle of the leaf-cutting ant *Acromyrmex subterraneus subterraneus* protect workers from attack by entomopathogenic fungi. *Biology Letters*, 8, 461-464.

Meirelles, L.A., Mendes, T.D., Solomon, S.E., Bueno, O.C., Pagnocca, F.C. & Rodrigues, A. (2014). Broad *Escovopsis* inhibition activity of *Pseudonocardia* associated with *Trachymyrmex* ants. *Env Microbiol Rep*, 6, 339-345.

Mueller, U.G. (2012). Symbiont recruitment versus ant-symbiont co-evolution in the attine ant–microbe symbiosis. *Current Opinion in Microbiology*, 15, 269-277.

Poulsen, M., Bot, A.N.M. & Boomsma, J.J. (2003a). The effect of metapleural gland secretion on the growth of a mutualistic bacterium on the cuticle of leaf-cutting ants. *Naturwissenschaften*, 90, 406-409.

Poulsen, M., Bot, A.N.M., Currie, C.R., Nielsen, M.G. & Boomsma, J.J. (2003b). Within-colony transmission and the cost of a mutualistic bacterium in the leaf-cutting ant *Acromyrmex octospinosus*. *Functional Ecology*, 17, 260-269.

Poulsen, M., Cafaro, M., Boomsma, J.J. & Currie, C.R. (2005). Specificity of the mutualistic association between actinomycete bacteria and two sympatric species of *Acromyrmex* leaf-cutting ants. *Molecular Ecology*, 14, 3597-3604.

Ranger, C.M., Biedermann, P.H.W., Phuntumart, V., Beligala, G.U., Ghosh, S., Palmquist, D.E. *et al.* (2018). Symbiont selection via alcohol benefits fungus farming by ambrosia beetles. *Proceedings of the National Academy of Sciences*, 4447-4452.

Rueden, C.T., Schindelin, J., Hiner, M.C., DeZonia, B.E., Walter, A.E., Arena, E.T. *et al.* (2017). ImageJ2: ImageJ for the next generation of scientific image data. *BMC Bioinformatics*, 18, 529.

Scheuring, I. & Yu, D.W. (2012). How to assemble a beneficial microbiome in three easy steps. *Ecology Letters*, 15, 1300-1307.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T. *et al.* (2012). Fiji: an open-source platform for biological-image analysis. *Nature Methods*, 9, 676.

Seipke, R.F., Kaltenpoth, M. & Hutchings, M.I. (2012). *Streptomyces* as symbionts: an emerging and widespread theme? *FEMS Microbiology Reviews*, 36, 862-876.

Sen, R., Ishak, H.D., Estrada, D., Dowd, S.E., Hong, E.K. & Mueller, U.G. (2009). Generalized antifungal activity and 454-screening of *Pseudonocardia* and *Amycolatopsis* bacteria in nests of fungus-growing ants. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 17805-17810.

Spence, M. (1973). Job market signaling *Quarterly Journal of Economics*, 87, 355–374.

756 Van Arnam, E.B., Ruzzini, A.C., Sit, C.S., Horn, H., Pinto-Tomás, A.A., Currie, C.R.
757 *et al.* (2016). Selvamicin, an atypical antifungal polyene from two alternative
758 genomic contexts. *Proceedings of the National Academy of Sciences*, 113, 12940-
759 12945.
760
761 Weyl, E.G., Frederickson, M.E., Yu, D.W. & Pierce, N.E. (2010). Economic contract
762 theory tests models of mutualism. *Proceedings of the National Academy of Sciences*,
763 107, 15712-15716.
764
765 Wickham, H., and G. Grolemund (2017). *R for data science*. O'Reilly Media.
766
767 Widder, S., Allen, R.J., Pfeiffer, T., Curtis, T.P., Wiuf, C., Sloan, W.T. *et al.* (2016).
768 Challenges in microbial ecology: building predictive understanding of community
769 function and dynamics. *The ISME Journal*, 10, 2557-2568.
770
771 Worsley, S.F., Innocent, T.M., Heine, D., Murrell, J.C., Yu, D.W., Wilkinson, B. *et*
772 *al.* (2018). Symbiotic partnerships and their chemical interactions in the leafcutter
773 ants (Hymenoptera: Formicidae). *Myrmecological News*, 27, 59-74.
774
775 Yu, D.W. (2001). Parasites of mutualisms. *Biological Journal of the Linnean Society*,
776 72, 529-546.
777
778