Improving the diagnostic yield of exome-sequencing, by predicting gene-phenotype associations using large-scale gene expression analysis

Patrick Deelen,[1,2,4] Sipko van Dam,[1,4] Johanna C. Herkert,[1,5] Juha M. Karjalainen,[1,5] Harm Brugge,[1,5] Kristin M. Abbott,[1] Cleo C. van Diemen,[1] Paul A. van der Zwaag,[1] Erica H. Gerkes,[1] Pytrik Folkertsma,[1] Tessa Gillett,[1] K. Joeri van der Velde,[1,2] Roan Kanninga,[1,2] Peter C. van den Akker,[1] Sabrina Z. Jan,[1] Edgar T. Hoorntje,[1,3] Wouter P. te Rijdt,[1,3] Yvonne J. Vos,[1] Jan D.H. Jongbloed,[1] Conny M.A. van Ravenswaaij-Arts,[1] Richard Sinke,[1] Birgit Sikkema-Raddatz,[1] Wilhelmina S. Kerstjens-Frederikse,[1] Morris A. Swertz,[1,2] Lude Franke[1]

[1] University of Groningen, University Medical Center Groningen, Department of Genetics, Groningen, 9700 VB, the Netherlands

[2] University of Groningen, University Medical Center Groningen, Genomics Coordination Center, Groningen, 9700 VB, the Netherlands

[3] Netherlands Heart Institute, Utrecht, the Netherlands

[4] These authors contributed equally to this work
[5] These authors contributed equally to this work

Corresponding author:
Lude Franke
E-mail: Lude@ludesign.nl

Abstract

Clinical interpretation of exome and genome sequencing data remains challenging and time consuming, with many variants with unknown effects found in genes with unknown functions. Automated prioritization of these variants can improve the speed of current diagnostics and identify previously unknown disease genes. Here, we used 31,499 RNA-seq samples to predict the phenotypic consequences of variants in genes. We developed GeneNetwork Assisted Diagnostic Optimization (GADO), a tool that uses these predictions in combination with a patient's phenotype, denoted using HPO terms, to prioritize identified variants and ease interpretation. GADO is unique because it does not rely on existing

knowledge of a gene and can therefore prioritize variants missed by tools that rely on

existing annotations or pathway membership. In a validation trial on patients with a known

genetic diagnosis, GADO prioritized the causative gene within the top 3 for 41% of the

cases. Applying GADO to a cohort of 38 patients without genetic diagnosis, yielded new

candidate genes for seven cases. Our results highlight the added value of GADO

(www.genenetwork.nl) for increasing diagnostic yield and for implicating previously

unknown disease-causing genes.

40   Introduction

With the increasing use of whole-exome sequencing (WES) and whole-genome sequencing

(WGS) to diagnose patients with a suspected genetic disorder, diagnostic yield is steadily

increasing [1]. Although our knowledge of the genetic basis of Mendelian diseases has

improved considerably, the underlying cause remains elusive for a substantial proportion of

cases. The diagnostic yield of genome sequencing varies from 8% to 70% depending on the

patient's phenotype and the extent of genetic testing [2]. Sequencing all ~20,000 protein-

coding genes by WES and entire genomes by WGS usually increases sensitivity but

decreases specificity: it results in off-target noise and reveals many variants of uncertain

clinical significance. In a study by Yang *et al.*, proband-only WES identified approximately

875 variants in each patient, even after removing low quality variants [3].

One strategy to manage the list of genetic variants is to perform trio analysis of samples

from the proband and both of his or her biological parents to ascertain, for instance,

whether a variant has *de novo* status [4]. Another strategy is to limit the analyses to a gene

panel of Online Mendelian Inheritance in Men (OMIM) disease-annotated genes [5] or genes

known to be directly related to the patient's phenotype. However, determining the actual

disease-causing variant requires further variant filtering based on information about its

predicted functional consequence, population frequency data, conservation, disease-specific

databases (such as the Human Gene Mutation Database [6]), literature, and segregation

analysis [7].

Several tools have been developed that aid in variant filtering and prioritization [8,9].

Annotation tools, such as VEP [10] and GAVIN [9], offer additional functionality that allows

variants to be filtered according to their population frequency and variant class. Other tools

use phenotype descriptions to rank potential candidates genes [11]. The phenotypes are

typically described in a structured manner, e.g. using Human Phenotype Ontology (HPO)

terms [12]. AMELIE (Automatic Mendelian Literature Evaluation), for example, prioritizes

candidate genes by their likelihood of causing the patient's phenotype based on automated
literature analysis [13]. However, this focus on what is known may inadvertently filter out
variants in potential novel disease genes. Alternatively, the causative gene defect could be
missed if a patient's phenotype differs from the features previously reported to be
associated to a disease gene. Tools like Exomiser can identify novel human disease genes,
as it prioritizes variants based on semantic phenotypic similarity between a patient's
phenotype described by HPO terms and HPO-annotated diseases, Mammalian Phenotype
Ontology (MPO)-annotated mouse and Zebrafish Phenotype Ontology (ZPO)-annotated fish
models associated with each exomic candidate and/or its neighbors in an interaction
network [14]. However, most available algorithms are based on existing knowledge on human
disease genes, their orthologues in animal models, or well-described biological pathways
(for a detailed review see [11]).

To overcome this, we hypothesized that co-regulation of expression data could be used to
prioritize variants, including those in less well studied genes. We assumed that if a gene or
a gene set is known to cause a specific disease or disease symptom, these genes will often
have similar molecular functions or be involved in the same biological process or pathway.
We reasoned that variants in genes with yet unknown function that are involved in the same
biological pathway or co-regulated with known disease genes likely result in the same
phenotype. In order to identify groups of genes with a related biological function, we used
an expansive compendium of 31,499 RNA-sequencing (RNA-seq) gene expression samples
to predict functions for genes with high accuracy.

We then developed a user-friendly tool that can prioritize variants in known *and* unknown
genes based on our functional predictions, which we designated GeneNetwork Assisted
Diagnostic Optimization (GADO). GADO ranks variants based on gene co-regulation in
publicly available expression data of a wide range of tissues and cell types using HPO terms
to describe a patient's phenotype. To validate our prioritization method, we tested how well
our method predicts disease-causing genes based on features described for each of the
genes in the OMIM database. We then used exome sequencing data of patients with a
known genetic diagnosis to benchmark GADO. Finally, we applied our methodology to
previously inconclusive WES data and identified several genes that contain variants that
likely explain the phenotype of the respective patients. Thus, we show that our methodology
is successful in identifying variants in novel, potentially relevant genes explaining the
patient's phenotype.

Results

Gene prioritization using GADO

We have developed GADO to perform gene prioritizations using the phenotypes observed in patients denoted as HPO terms [15]. In combination with a list of candidate genes (i.e. genes harboring rare and possibly damaging variants), this results in a ranked list of genes with the most likely candidate genes on top (**Figure 1**a). The gene prioritizations are based on the predicted involvement of the candidate genes for the specified set of HPO terms. These predictions are made by analyzing public RNA-seq data from 31,499 samples (**Figure 1**b), resulting in a gene prediction score for each HPO term. These predictions are solely based on co-regulation of genes annotated to a certain HPO term with other genes. This makes it possible to also prioritize genes that currently lack any biological annotation.

Figure 1: Schematic overview of GADO. *(a) Per patient, GADO requires a set of phenotypic features and a list of candidate genes (i.e. genes harboring rare alleles that are predicted to be pathogenic) as input. It then ascertains whether genes have been predicted to cause these features, and which ones are present in the set of candidate genes that has been provided as input. The predicted HPO phenotypes are based on the co-regulation of genes with sets of genes that are already known to be associated with that phenotype. (b) Overview of how disease symptoms are predicted using gene expression data from 31,499 human RNA-seq samples. A principal component analysis on the co-expression matrix results in the identification of 1,588 significant principal components. For each HPO term we investigate every component: per component we test whether there is a significant difference between eigenvector coefficients of genes known to cause a specific phenotype and a background set of genes. This results in a matrix that indicates which principal components are informative for every HPO term. By correlating this matrix to the eigenvector coefficients of every individual gene, it is possible to infer the likely HPO disease phenotype term that would be the result of a pathogenic variant in that gene.*

Public RNA-seq data acquisition and quality control

To predict functions of genes and HPO term associations, we downloaded all human RNA-seq samples publicly available in the European Nucleotide Archive (accessed June 30, 2016) (supplementary table 1) [16]. We quantified gene-expression using Kallisto [17] and removed samples for which a limited number of reads are mapped. We used a principal component analysis (PCA) on the correlation matrix to remove low quality samples and samples that were annotated as RNA-seq but turned out to be DNA-seq. In the end, we included 31,499

samples and quantified gene expression levels for 56,435 genes (of which 22,375 are

protein-coding).

Although these samples are generated in many different laboratories, we previously

observed that, after having corrected for technical biases, it is possible to integrate these

samples into a single expression dataset [18]. We validated that this is also true for our new

dataset by visualizing the data using t-Distributed Stochastic Neighbor Embedding (t-SNE).

We labeled the samples based on cell-type or tissue and we observed that samples cluster

together based on cell-type or tissue origin (**Figure 2**a). Technical biases, such as whether

single-end or paired-end sequencing had been used, did not lead to erroneous clusters,

which suggests that this heterogeneous dataset can be used to ascertain co-regulation

between genes and can thus serve as the basis for predicting the functions of genes.

Prediction of gene HPO associations and gene functions

To predict HPO term associations and putative gene functions using co-regulation (**Figure**

1b), we used a method that we had previously developed and applied to public expression

microarrays [19]. Since these microarrays only cover a subset of the protein-coding genes (n

= 14,510), we decided to use public RNA-seq data instead. This allows for more accurate

quantification of lower expressed genes and the expression quantification of many more

genes, including a large number of non-protein-coding genes. [20].

We applied this prediction methodology [19] to the HPO gene sets and also to Reactome [21],

KEGG pathways [22], Gene Ontology (GO) molecular function, GO biological process and GO

cellular component [23] gene sets. For 5,088 of the 8,657 gene sets (59%) with at least 10

genes annotated, the gene function predictions had significant predictive power (see

materials and methods). For the 8,657 gene sets with at least 10 genes annotated, the

median predictive power, denoted as Area Under the Curve (AUC), ranged between 0.73

(HPO) to 0.87 (Reactome) (**Figure 2**b).

A

B

Database	Number of gene sets	Gene sets ≥ 10 genes	Gene sets with significant predictive power	Median AUC
Reactome	2,143	1,388	1,150	0.87
GO molecular function	4,070	726	398	0.82
GO biological process	11,753	2,576	1,115	0.82
GO cellular component	1,609	500	370	0.84
KEGG	186	186	168	0.84
HPO	7,920	3,281	1,887	0.73

 Figure 2: A compendium of gene expression profiles that can be used for gene function
prediction (a) 31,499 RNA-seq samples derived from many different studies show coherent clustering
after correcting for technical biases. Generally, samples originating from the same tissue, cell-type or
cell-line cluster together. The two axes denote the first t-SNE components. (b) Gene co-expression
information of 31,499 samples is used to predict gene functions. We show the prediction accuracy for
gene sets from different databases. AUC, Area Under the Curve, GO, Gene Ontology, HPO, Human
Phenotype Ontology.

Prioritization of known disease genes using the annotated HPO terms

Once we had calculated the prediction scores of HPO disease phenotypes, we leveraged these scores to prioritize genes found by sequencing the DNA of a patient. For each individual HPO term–gene combination, we calculated a prediction z-score that can be used to rank genes. In practice, however, patients often present with not one feature but a combination of multiple features. Therefore, we combined the z-scores for each HPO term [24] to generate an overall z-score that explains the full spectrum of features in a patient. GADO uses these combined z-scores to prioritize the candidate genes: the higher the combined z-score for a gene, the more likely it explains the patient's phenotype.

Because many HPO terms have fewer than 10 genes annotated, and since we were unable to make significant predictions for some HPO terms, certain HPO terms are not suitable to use for gene prioritization. We solved this problem by taking advantage of the way HPO terms are structured. Each term has at least one parent HPO term that describes a more generic phenotype and thus has also more genes assigned to it. Therefore, if an HPO term cannot be used, GADO will make suggestions for suitable parental terms (supplementary figure 1).

To benchmark our prioritization method, we used the OMIM database [5]. We tested how well our method was able to retrospectively rank disease-causing genes listed in OMIM based on the annotated symptoms of these diseases. We took each OMIM disease gene (n = 3,382) and used the associated disease features (15 per gene on average) as input for GADO. What we found was that for 49% of the diseases GADO ranks the causative gene in the top 5% (**Figure 3**a, b). Moreover, we observed a statistically significant difference between the performance of GADO on true gene-phenotype combinations and its performance using a random permutation of gene-phenotype combinations (p-value = 2.16×10^{-532}).

Gene predictability scores explains performance differences between genes

For some combinations of genes and HPO terms listed in OMIM, GADO could not establish

the gene-phenotype combination (**Figure 3**). For example, variants in *SLC6A3* are known to

cause infantile Parkinsonism-dystonia (MIM 613135) [25–27], but GADO was unable predict the

annotated HPO terms related to the Parkinsonism-dystonia for this gene. This may,

however, be due to very low expression levels of *SLC6A3* in most tissues except specific

brain regions [28].

To better understand why we can't predict HPO terms for all genes, we used the Reactome,

GO and KEGG prediction scores. Jointly these databases comprise thousands of gene sets.

Since these databases describe such a wide range of biology, we assumed that if a gene

does not show any prediction signal for any gene set in these databases, gene co-

expression is probably not informative for this gene. To quantify this, we calculated, per
gene, the average skewness of the z-score distribution of the Reactome, GO and KEGG gene
sets. From this we were able to derive a 'gene predictability score' for every gene that is
independent of whether this gene is already known to play a role in any a disease or
pathway (**Figure 3**c, d, e). We then ascertained whether these 'gene predictability scores'
are correlated with the prediction z-score of the OMIM diseases, and found a strong
correlation (Pearson r = 0.54, p-value = 1.14×10^{-332}) between the gene predictability
scores and GADO's ability to identify a known disease gene (**Figure 3**c).

To investigate why some genes have a high 'gene predictability score' but low prediction
performance, we scored a set of genes known to cause cardiomyopathy (CM) for the
amount of literature evidence that these genes cause CM. We found several genes for which
the prediction score for the CM phenotype is lower than expected based on the gene
predictability scores (supplementary figure 2a). Pathogenic variants in the *TTR* gene
implicated in hereditary amyloidosis (MIM 105210) [29], for instance, cause accumulation of
the transthyretin protein in different organ systems, including the heart, resulting in CM.
However, this gene is primarily expressed in the liver. Therefore, its disease mechanism is
different from other mechanisms resulting in CM, as many inherited CMs are caused by
deleterious variants in genes highly expressed in the heart and directly affecting the
function of the cardiac sarcomere. Therefore, the phenotypic function prediction for this
gene may be worse than we would expect based on the predictability score. We performed a
similar analysis using the HPO term 'dilated cardiomyopathy' and observed a low prediction
performance for the *TMPO* gene, despite a high gene predictability score (supplementary
figure 2b). Previously, this gene was reported to be related to dilated cardiomyopathy
(DCM) and listed as such by OMIM. However, recent reclassification of the reported variants
using the ExAC data revealed that the reported variant was far too common to be causative
for DCM [30].

Benchmarking GADO using solved cases with realistic phenotyping
Although *in silico* benchmarking demonstrated the potential of GADO, it used all annotated
HPO terms for a disease. In practice, however, patients may only present with a limited
number of the annotated features. To perform a validation that was a more realistic
reflection of clinical practice, we used exome sequencing data of 83 patients with a known
genetic diagnosis. We used their phenotypic features as listed in their medical records prior
to the genetic diagnosis (supplementary table 2). On average, per patient, GADO yielded 56
possible disease-causing genes with variants that are rare and predicted to be deleterious.

In 41% of the patients the actual causative gene was ranked in the top 3 and in 50% of the

cases it was in the top 5 (mean rank 10) (**Figure 4**a).

Clustering of HPO terms

In addition to ranking potentially causative genes based on a patient's phenotype, we

observed that GADO can be used to cluster HPO terms based on the genes that are predicted

to be associated to these HPO terms. This can help identify pairs of symptoms that often occur

together, as well as symptoms that rarely co-occur, and we actually observed this for a patient

suspected of having two different diseases. This patient is diagnosed with a glycogen storage

disease, GSD type Ib, caused by compound heterozygous variants in *SLC37A4* (MIM 602671)

and DCM that is probably caused by a truncating variant in *TTN* (MIM 188840). Clustering of

the assigned HPO terms placed the phenotypic features related to GSD type Ib ('leukopenia'

(HP:0001882) and 'inflammation of the large intestine' (HP:0002037)) together, while

Cardiomyopathy (HP:0001638) was only weakly correlated to these specific features (**Figure**

4b).

Figure 4: Performance of GeneNetwork on solved cases *(a) Rank of the known causative gene*
among the candidate disease causing variants. (b) Our cohort contained a case with two distinct
conditions, and clustering showed the HPO terms of the same disease are closest to each other. Note,
the HPO term "Inflammation of the large intestine" did not yield a significant prediction profile and
therefore the parent terms "Abnormality of the large intestine", "Increased inflammatory response"
and "Functional abnormality of the gastrointestinal tract" where used for this case.

Reanalysis of previously unsolved cases

To assess GADO's ability to discover new disease genes, we applied it to data from 38 patients who are suspected to have a Mendelian disease but who have not had a genetic diagnosis. All patients had undergone prior genetic testing (WES with analysis of a gene panel according to their phenotype, supplementary table 3). On average three genes had a z-score ≥ 5 (which we used as an arbitrary cut-off and that correspond to a p-value of 5.7 X 10^{-7}) and were further assessed. In seven cases, we identified variants in genes not associated to a disease in OMIM or other databases, but for which we could find literature or for which we gained functional evidence implicating their disease relevance (**Table 1**). For example, we identified two cases with DCM with rare compound heterozygous variants in the *OBSCN* gene (MIM 608616) that are predicted to be damaging. In literature, inherited variant(s) in *OBSCN*, encoding obscurin, are associated with hypertrophic CM [31] and DCM [32]. Furthermore, obscurin is a known interaction partner of titin (TTN), a well-known DCM-related protein [31]. Another example came from a patient with ichthyotic peeling skin syndrome, which is caused by a damaging variant in *FLG2 (*MIM 616284). We recently published this case where we prioritized this gene using an alpha version of GADO [33].

HPO terms used	Number of genes with candidate variant	Number of genes with z ≥ 5	Candidate gene	Variants	CADD scores	GnomAD minor allele frequency	Supporting papers	Expression in relevant tissue
HP:0001644	247	5	*OBSCN*	NM_001098623.2: c.[15037C>T]; [20963delC]	24.8 25.2	8.0×10^{-5} 1.7×10^{-3}	[31, 32]	Yes
HP:0001644	226	3	*OBSCN*	NM_001098623.2: c.[5545C>T]; [22384+3_22384 +21del]	14.7 7.8	3.2×10^{-4} 0	[31, 32]	Yes
HP:0008066 HP:0008064	359	3	*FLG2*	NM_001014342.2: c.[632C>G]; [632C>G]	35.0 35.0	1.1×10^{-5} 1.1×10^{-5}	[34]	Yes
HP:0001263 HP:0001249 HP:0000717 HP:0000708 HP:0002167 HP:0002360 HP:0000664	206	12	*INO80*	NM_017553.2: c. [898C>T]	34	0	[35, 36]	Yes
HP:0001644	346*	2	MB	NM_00203377.1: c.[214G>A]	22.4	3.6×10^{-5}	[37]	Yes
HP:0001644	126*	1	*SYNPO2L***	NM_001114133.2: c.[473G>A]	24.1	5.4×10^{-4}	[38]	Yes
HP:0001638	336	4	*NRAP***	NM_001261463.1: c.[4648C>T]	20.4	8.7×10^{-4}	[39]	Yes

*Table 1: unsolved cases with new candidate genes. Out of the 38 unsolved patients investigated, we identified candidate genes in seven patients. For these genes we have found literature that indicates these genes fit the phenotype of these patients or for which we gained functional evidence implicating their disease relevance. *These variants where pre-filtered for family segregation. **The variants in these genes do not fully explain the phenotype but are likely contributing to the phenotype.*

www.genenetwork.nl

All analyses described in this paper can be performed using our online toolbox at

www.genenetwork.nl. Users can perform gene prioritizations using GADO by providing a set

of HPO terms and a list of candidate genes (**Figure 5**a). Per gene, it is also possible to

download all prediction scores for the HPO terms and pathways. Our co-regulation scores

between genes can be used for clustering. Furthermore, the predicted pathway and HPO

293   annotations of genes can be used to perform function enrichment analysis (**Figure 5**b). We

294   also support automated queries to our database.

A

B

Figure 5: www.genenetwork.nl *(a) Prioritization results of one of our previously solved cases. This*
patient was diagnosed with Kleefstra syndrome. The patient only showed a few of the phenotypic
features associated with Kleefstra syndrome and additionally had a neoplasm of the pituitary (which is
not associated with Kleefstra syndrome). Despite this limited overlap in phenotypic features, GADO
was able to rank the causative gene (EHMT1) second. Here, we also show the value of the HPO
clustering heatmap, the two terms related to the neoplasm cluster separately from the intellectual
disability and the facial abnormalities that are associated to Kleefstra syndrome. (b) Clustering of a set
of genes allowing function / HPO enrichment of all genes or specific enrichment of automatically
defined sub clusters. Here we loaded all known DCM genes and OBSCN, and we focus on a sub-cluster
of genes containing OBSCN (highlighted by the arrow). We see that it is strongly co-regulated with
many of the known DCM genes. Pathway enrichment of this sub-cluster reveals that these genes are
most strongly enriched for the muscle contraction Reactome pathway. DCM, Dilated Cardiomyopathy.

Discussion

Prioritizing genes from WES or WGS data remains challenging. To meet this challenge, we

developed GADO, a novel tool to prioritize genes based on the phenotypic features of a

patient. Since the classification of variants is labor-intensive, prioritization of the most likely

candidate variants saves time in the diagnostic process.

Importantly, GADO can also aid in the discovery of currently unknown disease genes. The

main advantage of our methodology is that it does not rely on any prior knowledge about

disease-gene annotations. Instead, we used predicted gene functions based on co-

expression networks extracted from a large compendium of publicly available RNA-seq

samples. RNA-seq has previously shown to be very helpful to accurately quantify expression

levels of lowly expressed genes and non-coding genes [18]. To evaluate our diagnostic

algorithm, we developed a testing scenario based on simulated patients presenting with all

clinical features listed in OMIM for a certain disease or syndrome. This validation test

showed that for 49% of the diseases the causative gene ranks in the top 5%. We also

investigated the OMIM "provisional" category of genes for which there is limited evidence.

Both the OMIM disease-gene annotation and the provisional annotations perform

significantly better than a random permutation. While we do find a small but significant

difference in prediction performance between the provisionally annotated genes and the

more established disease associated genes, we conclude, based on our findings, that these

provisional OMIM annotations are generally of similar reliability to the other OMIM disease

annotations.

Benchmarking on sequence data of patients with a known genetic diagnosis revealed that

GADO returned the real causative variant within the top 3 results for 41% of the samples,

indicating the potential power of GADO for a large number of diseases. Finally, in seven

patients, GADO was able to identify potential novel disease genes that are strong candidates

based on literature or functional evidence. For other cases we have identified genes with a

strong prediction score harboring variants that might explain the phenotype. However, since
very little is known about these genes it is not yet possible to draw firm conclusions.
Hopefully this will become possible in the near future through initiatives like Genematcher
[40].

Potential to discover novel human disease genes
Over the last decade, several computational tools have been developed to prioritize variants
in genes. Some, such as GAVIN, focus on variant filtering and prioritization based on
deleteriousness scores, allele frequency and inheritance model [9]. Other methods measure
the similarity between the clinical manifestations observed in a patient and those
representing each of the diseases in a database or literature. Exomiser is closely related to
GADO as it prioritizes genes based on specified HPO terms and also infers HPO annotation
for unknown genes [14]. The gene prioritization by Exomiser is based on the effects of
orthologs in model organisms and applies a guilt-by-association method using protein-
protein associations provided by STRING [41]. Exomiser performs better than GADO in ranking
known disease-causing genes (supplementary table 4) and is also able to identify potential
new genes in human disease. However, Exomiser has a limitation in that only a subset of
the protein-coding genes has orthologous genes in other species for which a knockout
model also exists. Additionally, the used STRING interactions are biased towards well
studied genes and rely heavily on existing annotations to biological pathways
(supplementary figure 4). There are however, still 3,922 protein-coding genes that are not
currently annotated in any of the databases we used, and there are even more non-coding
genes for which the biological function or role in disease is unknown. Since GADO does not
rely on prior knowledge, it can be used to prioritize variants in both coding *and* non-coding
genes (for which no or limited information is available). GADO thus enables the discovery of
novel human disease genes and can complement existing tools in analyzing the genomic
data of patients who have a broad spectrum of phenotypic abnormalities.

Limitations
The gene predictability score indicates for which genes we can reliably predict phenotypic
associations and for which genes we cannot based on gene co-regulation. This score gives
insight into which genes are expected to perform poorly in our prioritization. We found
strong correlation between these gene predictability scores and the gene prioritization z-
scores. Thus, genes with a high predictability score have more accurate HPO term
predictions. However, since our predictions primarily rely on co-activation patterns that we
identified from RNA-seq data, our method does not perform well for genes where gene-
expression patterns are not informative of their function. This could, for instance, be the

case for proteins relying heavily on post-translation modifications for regulation or genes for
which different transcripts have distinct functions. This last limitation can potentially be
overcome by predicting HPO-isoform associations by using transcript-based expression
quantification.

Insufficient statistical power to obtain accurate predictions may be another explanation for
the low predictability scores of certain genes. This may be true for genes that are poorly
expressed or expressed in only a few of the available RNA-seq samples. The latter issue we
expect to overcome in the near future as the availability of RNA-seq data in public
repositories is rapidly increasing. Initiatives such as Recount enable easy analysis on these
samples [42], allowing us to update our predictions in the future, thereby increasing our
prediction accuracy.

For some genes we are unable to predict annotated disease associations despite having a
high gene predictability scores. Some genes, such as *TTR,* simply act in a manner unique to
a specific phenotype. Other genes, such as *TMPO,* turned out to be false positive disease
associations. These examples show that our gene predictability score has the potential to
flag genes acting in a unique manner as well as genes that might be incorrectly assigned to
a certain disease or phenotype.

We noted that the median prediction performance of HPO terms is lower compared to the
other gene sets databases used in our study, such as Reactome. This may be due to the
fact that phenotypes can arise by disrupting multiple distinct biological pathways. For
instance, DCMs can be caused by variants in sarcomeric protein genes, but also by variants
in calcium/sodium handling genes or by transcription factor genes [43]. As our methodology
makes guilt-by-association predictions based on whether genes are showing similar
expression levels, the fact that multiple separately working processes are related to the
same phenotype can reduce the accuracy of the predictions (although it is often still
possible to use these predictions as the DCM HPO phenotype prediction performance AUC =
0.76).

Complexity

Given that nearly 5% of patients with a Mendelian disease have another genetic disease [44],
it is important to consider that multiple genes might each contribute to specific phenotypic
effects. Clinically, it can be difficult to assess if a patient suffers from two inherited
conditions, which may hinder variant interpretation based on HPO terms. We showed that
GADO can disentangle the phenotypic features of two different diseases manifesting in one
patient by correlating and subsequently clustering the profiles of HPO terms describing the

patient's phenotype. If the HPO terms observed for a patient do not correlate, it is more

likely that they are caused by two different diseases. An early indication that this might be

the case for a specific patient can simplify subsequent analysis because the geneticist or

laboratory specialist performing the variant interpretation can take this in consideration.

GADO also facilitates separate prioritizations on subsets of the phenotypic features.

Conclusion

Connecting variants to disease is a complex multistep process. The early steps are usually

highly automated, but the final most critical interpretations still rely on expert review and

human interpretation. GADO is a novel approach that can aid users in prioritizing genes

using patient-specific HPO terms, thereby speeding-up the diagnostic process. It prioritizes

variants in coding *and* non-coding genes, including genes for which there is no current

knowledge about their function and those that have not been annotated in any ontology

database. This gene prioritization is based on co-regulation of genes identified by analyzing

31,499 publicly available RNA-seq samples. Therefore, in contrast to many other existing

prioritization tools, GADO has the capacity to identify novel genes involved in human

disease. By providing a statistical measure of the significance of the ranked candidate

variants, GADO can provide an indication for which genes its predictions are reliable. GADO

can also detect phenotypes that do not cluster together, which can alert users to the

possible presence of a second genetic disorder and facilitate the diagnostic process in

patients with multiple non-specific phenotypic features. GADO can easily be combined with

any filtering tool to prioritize variants within WES or WGS data and can also be used in gene

panels such as PanelApp [45]. GADO is freely available at www.genenetwork.nl to help guide

the differential diagnostic process in medical genetics.

426   Materials and Methods

Sample acquisition

All RNA-seq data used in this project was acquired from the European Nucleotide Archive

(ENA) database [46]. Of the 67,090 human RNA-seq samples, with at least 500,000 reads,

registered in the ENA on June 30, 2016 (supplementary table 1), 67,019 were successfully

downloaded. For 71 of the registered samples, the files were missing. Sample annotations

were acquired from [18,47] and through manual curation based on study meta-information in

the ENA database (supplementary table 1).

Gene expression quantification

The 67,019 downloaded samples were mapped to transcript annotations from Ensemble

release 83 which uses build GRCh38.p5 of the human genome [48] using Kallisto [17] version

0.42.4, and the number of reads assessed. The number of reads mapped per sample was

obtained from the Kallisto summary file. The following genome files were used:

ftp://ftp.ensembl.org/pub/release-

83/fasta/homo_sapiens/cdna/Homo_sapiens.GRCh38.cdna.all.fa.gz

ftp://ftp.ensembl.org/pub/release-

83/fasta/homo_sapiens/ncrna/Homo_sapiens.GRCh38.ncrna.fa.gz

These files were merged and used to build the Kallisto reference index file. The following

setting, in addition to all default settings, was used: –k 31.

The following Kallisto settings were used mapping all 67,019 samples using default settings

for paired-end data mapping. For single-end data mapping we used the following settings in

addition to the defaults: –l 200 and –s 20 –bias.

After obtaining the transcript counts per sample, these transcript-level counts were summed

to gene-level counts for each sample.

Gene quality control

We quantified 66,233 genes, which were filtered on the criteria described below, after which

56,435 genes remained. Twenty-nine gene names were duplicates/identical. After these

were removed, 66,203 genes remained. Of these, 3,628 genes are not expressed (0 reads

detected among 31,499 samples) and were removed, leaving 62,575 genes. Next, we

detected a number of duplicate genes (100% sequence similarity). Since these genes with

perfect sequence similarity have exactly the same number of reads mapping, we were

concerned they would appear as perfectly co-expressed genes in our analysis. Most of these

genes are either incorrectly mapped genes in the genome build or duplicates of their

biological counterpart. Due to their high sequence similarity they are indistinguishable to the

mapping tool (potentially introducing false correlations). To avoid potential biases resulting

in deceptively high co-expression values, we decided to remove this bias prior to our

analysis. 5,471 of these were not located on chromosomes (but on scaffolds), and were

removed, leaving 57,104 genes. Another 665 genes had identical transcripts: different IDs,

but 100% identical sequences (e.g. ENST00000442165 and ENST00000446969).

An additional four genes had no expression in any of the remaining samples after removing

outlier/poor-quality samples, as described below, and were also removed prior to the PCA

analysis. The 56,435 genes that remained were used for our analyses (supplementary figure

5).

RNA-seq sample quality control

We excluded all samples in which less than 70% of the reads successfully mapped to the

genome, as reported by Kallisto, resulting in 36,761 samples.

Principal component analysis to identify outlier samples

To identify outlier samples, we conducted a principal component analysis (PCA) along the

following steps. First, all estimated counts were log2 transformed. Second, the data was

quantile normalized. Third, the covariance over the samples was calculated. Fourth, genes

without variance were removed from the dataset. Fifth, a PCA was conducted on the

covariance matrix. An arbitrary cut-off on PC 1 was selected at 0.0049 (supplementary

figure 6), leaving us with 32,142 samples.

Removal of non-Illumina samples

Since only a small number of samples that passed quality control (147 samples, <0.5% of

the total number of samples) were not sequenced on Illumina machines, we removed these

to avoid potential biases as a result of these different sequencing tools. This left 31,995

samples in our dataset.

Removing duplicate samples

A number of samples had identical values for all genes. Upon inspection, some of these

samples appeared to be have been used by multiple studies and uploaded to the ENA

database multiple times. To remove duplicate samples, we identified all samples with a

correlation >0.9999, randomly selected one of them to include and removed the other.

After this step, 31,499 samples remained.

Removal of technical biases

To identify potential technical biases in our data, we calculated the correlation between the

PC-scores for each PC and the following potential confounders: read length, paired/single

end, total reads in the dataset and percentage mapping reads (supplementary figure 7). We

found that all these factors significantly correlated to our sample PC scores for multiple PCs

(p-value < 0.01), indicating that these technical factors would affect the co-expression

detected in the dataset, if not removed. We decided not to correct for GC content per gene

as this may also have biological meaning [49]. For a manual of the covariate removal pipeline

we refer to: https://github.com/molgenis/systemsgenetics/tree/master/eqtl-mapping-

pipeline. To remove covariates, we used the "adjustcovariates" option.

PCA

After correcting our dataset for technical biases, we conducted the following steps on the

matrix. First, we calculated the correlation over the genes. Second, we conducted a PCA

over the correlation matrix over the genes. Third, we calculated PC scores for each sample

for all PCs.

Inspection of gene PC eigencoefficients

To investigate if any technical biases were present for the different gene types (coding,

miRNA, pseudogene, etc.), we plotted the gene eigencoefficients for the first 10 PCs and

colored the genes by biotype (supplementary figure 8) and detected an outlier cluster on

PC8 and PC9, which were further investigated (supplementary figure 9).

Inspection of sample PC scores

To better understand the origin of the outlier genes in eigenvector coefficients of PC 8 and

PC 9, we investigated the PC scores of the samples for these PCs. Additionally, we created a

plot for each of the sample PC scores of the first 10 PCs (supplementary figure 10). We

observed that there is a clear biological explanation for these outliers, and therefore we

decided to retain these signals in the data (supplementary figure 11).

Gene co-regulation analysis

After the quality control steps described above, we conducted a co-regulation analysis using

the 31,499 sample by 56,435 gene matrix. The co-regulation analysis was performed using

the PC eigencoefficients of the genes for each of the reliable PCs obtained from our gene-co-

expression matrix. To determine which PCs are reliable, Cronbach's alpha [50] was calculated

for each PC (based on PCA of the gene-correlation matrix). Those PCs with a Cronbach's

Alpha ≥ 0.7 were considered reliable, and is a commonly used cutoff [51]. In total, 1,588 PCs

have a Cronbach's Alpha ≥ 0.7. Additionally, we calculated the variance explained by each

of these PCs and found the first 1588 PCs explain 66 percent of the variance

(supplementary figure 12). By including signals from only these PCs, we aimed to remove

signals that are not reliable from our analysis. This method was previously shown to

perform better than using the correlation matrix directly [19]. The co-regulation scores were

calculated by calculating the correlation between the eigencoefficients of each gene pair.

Prior to this step the eigencoefficients were standard normalized per gene, after which the

eigencoefficients per PC were standard normalized. The logic to this step is to let the signal

a gene has for each PC weigh equally when determining the correlation between 2 genes.

Here we presumed each PC represents some biological process and those genes that are co-

expressed in multiple processes should be reported as strongly co-expressed. This is

illustrated and further explained in [19]. The p-values of co-regulated genes can be queried via
the website.

Data visualization of sample PC scores using a t-SNE plot

To identify clusters for each cell type and tissue type, we used the sample PC scores, which
indicate how strong the signal of each sample is for each PC in the data. Here, each PC is a
gene expression signature for the complete set of genes. To visualize how the samples
cluster in a two dimensional figure, we constructed a t-SNE plot [52] based on these sample
PC-scores using the Rtsne library [53] (version 0.13). The t-SNE was run with a perplexity of
50, and we ran 10,000 iterations on our sample PC score matrix. We found that single
clusters were visible for many cell- and tissue-types (**Figure 2**a). Most of these clusters
contain samples from different studies, which suggests that these clusters are not merely a
representation of study-specific biases. The fact that studies with multiple cell/tissue types
show multiple clusters further supports the suggestion that the clusters are not driven by
non-biological inter-study differences.

Gene function and HPO association predictions

Next, we used the PC eigenvector coefficients calculated in the previous steps to predict
functions for genes and to predict which phenotypes they are most likely to play a role in
(also described in [19]). For each of the 1,588 reliable PCs, we determined the extent to which
each PC captures the activity of a biological module (defined as a group of genes annotated
to a term, e.g. a GO function term or HPO phenotype).

To do this, the following steps were taken. First, for each PC, a student's T-test was
conducted between the eigencoefficients of the genes annotated to a particular term and a
group of genes serving as a background. This background consisted of all genes annotated
to any term in a specific database, except for those annotated to the term for which the T-
test was conducted. Genes that were not annotated to any term in a database were
excluded from this background, as these genes have not yet been annotated to any
biological functions/terms (because they have not been studied yet). Second, the resulting
p-values were transformed into a z-score, which are indicating to which extend each PC
represents a biological function/term. This was repeated for each of the 1,588 significant
PCs, resulting in a z-score for each PC-term combination. Higher absolute z-values between
a term and a PC indicate that the signal for that PC is more strongly related to that term.
We applied this methodology to the gene sets described by terms in the following
databases: Reactome and KEGG pathways, Gene Ontology (GO) molecular function, GO
biological process and GO cellular component terms and finally to HPO terms. We excluded

terms for which fewer than 10 genes are annotated because predictions for smaller groups

of genes are less accurate and might be misleading. Predictions were made for 8,657 gene

sets in total. For each term, we calculated how well each PC captured the signal of the

genes that are annotated to that term. Third and last, to predict which genes are correlated

to a particular HPO term, we correlated the 1,588 z-scores for that term (as calculated

above) with the 1,588 eigenvector coefficients of a gene. These correlations were

transformed into z-scores, which we refer to as prioritization scores. This can be done for

any gene-to-HPO term combination. However, when a gene is already explicitly annotated

to the term and we wish to predict whether that gene is predicted to be involved in that

term, there is a small circular bias as the z-scores for this term were partly calculated based

on this gene. To remove this bias in these circumstances, the 1,588 z-scores for a gene set

were first re-calculated while assuming these gene is not involved in that term, after which

the prediction for this gene was made.

Validation of the GO, HPO and Reactome term predictions

To determine the accuracy of our GO, HPO and Reactome term predictions, we calculated

how well we could predict genes that are part of a term. To do so, we used the prioritization

z-scores that the genes had for a particular term. For each term, we calculated an Area

Under the Curve (AUC), using a Mann-Whitney U test, on the prioritization scores of the

genes that are part of the term versus those that are not part of the term. These AUCs

indicate how accurate the predictions were, with an AUC of 1 indicating perfect predictions

and an AUC of 0.5 indicating no predictive power. The average AUC for each category was

calculated based on all terms with at least 10 genes annotated and for which the p-value

was less than 0.05 (Bonferroni corrected for the number of pathways for the category

tested) (**Figure 2**b).

GADO predictions

To identify potential causative variants in patients, we used HPO term annotations

describing the patient's features. The gene prediction z-scores for an HPO term were used

to rank the genes. If a patient's phenotype was described by more than one HPO term, a

meta-analysis was conducted. In this case a weighted z-score was calculated by adding the

HPO z-scores for all the patient's HPO terms and then dividing by the square root of the

number of HPO terms. In this calculation, we used only those HPO terms, which have

significant predictive power (based on whether genes annotated to this term have

significantly absolute higher z-scores than those not annotated to the term as calculated in

the section "Gene function and HPO association predictions"). If the predictions for a

patient's HPO term were not significant, the parent/umbrella HPO term(s) was used. (The

online GADO tool supplies the user with a list of parent terms from which the user can then
manually select which terms should be used in the analysis (supplementary figure 1)). If
this parent term also did not have significant predictive power, the parent's parent term was
used (thus moving up the HPO tree until a parent term is found which has significant
predictive power). If an HPO term has multiple parents, predictions were made using each
parent and the results are reported separately. The genes with the highest z-scores are
most relevant for the patient according to GADO's predictions. This analysis can be
conducted at: https://www.genenetwork.nl/gado.

Validation of disease-gene predictions
To benchmark our method we used the OMIM morbid map [5] downloaded on March 26,
2018, containing all disease-gene-phenotype entries. From this list, we extracted the
disease-gene associations, excluding non-disease and susceptibility entries. We extracted
the provisional disease-gene associations separately. For each disease in OMIM, we used
GADO to determine the rank of the causative gene among all genes in the OMIM morbid
map. For this we used all phenotypes annotated to the OMIM disease. If any of the HPO
terms did not have significant predictive power, the parent term(s) was used.

To determine if these distributions were significantly different from what we expect by
chance, we permuted the data. We replaced the existing gene-OMIM annotation but
assigned every gene to a new disease (keeping the phenotypic features for a disease
together), assuring that the randomly selected gene was not already annotated to any of
the phenotypes of the original gene.

Cohort of previously solved cases
To test if GADO could help prioritize genes that contain the causative variant, we used 83
samples of patients who were previously genetically diagnosed through whole exome
analysis or gene panel analysis. These samples encompass a wide variety of different
Mendelian disorders (supplementary table 2). To assess which genes harbor potentially
causative variants, we first called and annotated the variants from the exome sequencing
files.

Variant calling
We used the available WES or WGS data from patients with and without genetic diagnosis.
These samples were genotyped using a relatively standard BWA and GATK pipeline. For a
detailed description of the genotype pipeline see: https://molgenis.gitbooks.io/ngs_dna/
(version 3.4.0). For the WGS samples, we confined our analysis to the exome.

Variant annotation

We used GAVIN to annotate our variants to obtain a list of candidate variants. GAVIN

prioritizes genes based on, among other factors, minor allele frequency and gene-

recalibrated CADD scores (for details see [9]). For 11 of the previously solved cases, GAVIN

did not flag the causative variant as a candidate. To be able to include these samples in our

GADO benchmark, we added the causative genes for these cases manually to the candidate

list.

GADO ranking

The phenotypic features of a patient were translated into HPO terms, which were used as

input to GADO for ranking all genes based on how likely they are to cause that set of

features. If any of the HPO terms did not have significant predictive power, the parent

term(s) was used. From the resulting list of ranked genes, the known disease genes

harboring a potentially causative variant were selected. Next, we determined the rank of the

gene with the known causative variant among the selected genes. If a patient harbored

multiple causative variants in different genes, in case of di-genic inheritance or two

inherited conditions, the median rank of these genes was reported (supplementary table 2).

Benchmark comparison with Exomiser

To evaluate GADO's performance, we compared GADO with Exomiser [54] (version 10.1.0,

with exomiser-phenotype-1802 and exomiser-genome-hg19-1805 files from

https://data.monarchinitiative.org/exomiser/data/). Both GADO and Exomiser were given

each patient candidate gene list along with their respective set of phenotypes as input.

Default settings were used. We used the gene rankings based on

"EXOMISER_GENE_COMBINED_SCORE" and identified the rank of the causative gene

(supplementary table 4). In case of a tie, the average rank of the ties was reported. If a

patient harbored multiple causative variants, the median rank of the genes harboring the

causative variants was reported. To ensure a fair comparison, we used GADO on the set of

genes reported by Exomiser (supplementary table 4).

Unsolved cases cohorts

In addition to the patients with a known genetic diagnosis, we tested 38 unsolved cases

(supplementary table 3). These are patients with mainly cardiomyopathies or developmental

delay. All patients were previously investigated using exome sequencing, by analyzing a

gene panel appropriate for their phenotype. To allow discovery of potential novel disease

genes, we used GADO to score all genes with candidate variants. For genes with a

prediction z-score ≥ 5, a literature search for supporting evidence was performed to assess

whether these genes are likely candidate genes.

GADO web-tool

To make the gene-co-regulation-based HPO predictions publicly available a website was

constructed: www.genenetwork.nl

On this website the user can conduct the following analyses:

1. Predict putative functions of genes. This can be achieved by querying a gene, for which

gene-network will then predict the function based on the functional enrichment of its co-

regulation partners. Enrichment for GO, Reactome, KEGG and HPO phenotypes can be

retrieved.

2. Prioritize potential causative disease genes for patients: Based on HPO terms or a group

of genes annotated to a patient, the GADO tool will rank all genes based on how likely they

are to be related to the patient's phenotype. These can be further filtered for genes of

interest, by providing a list of genes known to harbor likely causative variants.

Gene network visualization

Edges are drawn between two genes/nodes based on a z-score cutoff. The cutoff at which a

line/edge between two genes should be drawn can be manually altered with the bar in the

top right corner. The network is drawn based on a force directed layout and clusters are

assigned using affinity propagation [55]

HPO, Reactome, KEGG and GO enrichment calculations

On the network page it is possible to retrieve which HPO, Reactome, KEGG and GO

categories are enriched among the visualized genes. It is also possible to retrieve this for a

sub-selection of these genes. The enrichment is calculated based on the z-scores of each of

these genes for each category. For each category/term, a Mann-Whitney U test is conducted

between the z-scores of the genes in the network versus the z-scores of genes that are not

part of the visualized network. The pathways with the most significant p-values are then

ranked highest.

It is also possible to identify which other genes are strongly co-regulated with those

visualized in the network. This is done similarly to how the correlation between a gene and

a pathway is calculated, as described above in "Gene function and HPO association

predictions". First, the z-scores for each PC of the genes visualized in the network is

calculated. After the z-scores of this group of genes have been calculated for each4

pathway, the correlation of the PC coefficients for each gene not in the network with these
z-scores is calculated. The genes with the most significant correlation are ranked highest.

Gene predictability scores

To explain why for some genes we cannot predict known HPO annotation, we have
established a gene predictability score. We have calculated this gene predictability using the
prioritization z-scores based on Reactome, GO and KEGG. For each gene and for each
database we calculated the skewness in the distribution of the prioritization z-scores of the
gene sets. We used the average skewness as the gene predictability score.

710 Description of Supplemental Data

Supplementary figure 1. Selection of parent HPO term if GADO does not have significant
predictive power for query term

Supplementary figure 2. Comparison of GADO performance with the level of evidence for
each cardiomyopathy-related gene

Supplementary figure 3. Comparison between GADO and Exomiser rankings

Supplementary figure 4. Correcting for biases in co-expression networks

Supplementary figure 5. Histogram of the gene types included in our analyses

Supplementary figure 6. PCA plot of 36,761 samples

Supplementary figure 7. Investigation of principal components capturing technical biases

Supplementary figure 8. Visualization of PC1 to PC 10 of PCA over gene correlation matrix

Supplementary figure 9. Outlier genes in PC 8 and PC 9 of PCA over gene correlation matrix

Supplementary figure 10. PC sample scores to distinguish different tissues

Supplementary figure 11. Outlier samples in PC sample scores of PC 8 and PC 9

Supplementary figure 12. Variance explained by first 1588 PCs

Supplementary table 1. A list of samples annotated in the European Nucleotide Archive June
30, 2016

Supplementary table 2. A list of 83 diagnosed patients with Mendelian disorders and
corresponding predictions with GADO

Supplementary table 3. A list of 38 undiagnosed patients with suspected Mendelian
disorders

Supplementary table 4. A comparison between GADO and Exomiser predictions using a list
of 83 diagnosed patients with Mendelian disorders

Declaration of Interests

734 The authors declare no competing interests.

Acknowledgments

We are grateful for the participation of the patients and their parents in this study. We
thank Kate Mc Intyre for editing the manuscript and Marieke Bijlsma, Gerben van der Vries,
Sido Haakma and Pieter Neerincx for support with the computational analyses. This work
was carried out on the Groningen Center for Information Technology (Strikwerda, W. Albers,
R. Teeninga, H. Gankema and H. Wind) and Target storage (E. Valentyn and R. Williams).
Target is supported by Samenwerkingsverband Noord Nederland, the European Fund for
Regional Development, the Dutch Ministry of Economic Affairs, Pieken in de Delta and the
provinces of Groningen and Drenthe. Wouter P. te Rijdt is supported by Young Talent
Program (CVON PREDICT) grant 2017T001 from the Dutch Heart Foundation. Netherlands
Heart Institute, Utrecht, the Netherlands. This work is supported by a grant from the
European Research Counsil (ERC Starting Grant agreement number 637640 ImmRisk) to
Lude Franke and a VIDI grant (917.14.374) from the Netherlands Organisation for Scientific
Research (NWO) to Lude Franke. This work was supported by BBMRI-NL, a research
infrastructure financed by the Dutch government (NWO 184.021.007).

Web Resources

Gene Network, www.genenetwork.nl
GADO, https://www.genenetwork.nl/gado
European Nucleotide Archive, https://www.ebi.ac.uk/ena
Ensembl, https://www.ensembl.org
OMIM, https://www.omim.org
Genotyping pipeline, https://molgenis.gitbooks.io/ngs_dna/
Covariate removal pipeline, https://github.com/molgenis/systemsgenetics/tree/master/eqtl-
mapping-pipeline

References

1. Brown, T.L., and Meloche, T.M. (2016). Exome sequencing a review of new strategies for rare genomic disease research. Genomics *108*, 109–114.

2. Wright, C.F., FitzPatrick, D.R., and Firth, H. V. (2018). Paediatric genomics: diagnosing rare disease in children. Nat. Rev. Genet. *19*, 253–268.

3. Yang, Y., Muzny, D.M., Xia, F., Niu, Z., Person, R., Ding, Y., Ward, P., Braxton, A., Wang, M., Buhay, C., et al. (2014). Molecular findings among patients referred for clinical whole-exome sequencing. JAMA *312*, 1870–1879.

4. Wright, C.F., Fitzgerald, T.W., Jones, W.D., Clayton, S., McRae, J.F., van Kogelenberg, M., King, D.A., Ambridge, K., Barrett, D.M., Bayzetinova, T., et al. (2015). Genetic diagnosis of developmental disorders in the DDD study: a scalable analysis of genome-wide research data. Lancet (London, England) *385*, 1305–1314.

5. McKusick-Nathans Institute of Genetic Medicine, and Johns Hopkins University Online Mendelian Inheritance in Man, OMIM , https://omim.org/.

6. Stenson, P.D., Mort, M., Ball, E. V., Evans, K., Hayden, M., Heywood, S., Hussain, M., Phillips, A.D., and Cooper, D.N. (2017). The Human Gene Mutation Database: towards a comprehensive repository of inherited mutation data for medical research, genetic diagnosis and next-generation sequencing studies. Hum. Genet. *136*, 665–677.

7. Lek, M., Karczewski, K.J., Minikel, E. V., Samocha, K.E., Banks, E., Fennell, T., O'Donnell-Luria, A.H., Ware, J.S., Hill, A.J., Cummings, B.B., et al. (2016). Analysis of protein-coding genetic variation in 60,706 humans. Nature *536*, 285–291.

8. Eilbeck, K., Quinlan, A., and Yandell, M. (2017). Settling the score: variant prioritization and Mendelian disease. Nat. Rev. Genet. *18*, 599–612.

9. van der Velde, K.J., de Boer, E.N., van Diemen, C.C., Sikkema-Raddatz, B., Abbott, K.M., Knopperts, A., Franke, L., Sijmons, R.H., de Koning, T.J., Wijmenga, C., et al. (2017). GAVIN: Gene-Aware Variant INterpretation for medical sequencing. Genome Biol. *18*, 6.

10. McLaren, W., Gil, L., Hunt, S.E., Riat, H.S., Ritchie, G.R.S., Thormann, A., Flicek, P., and Cunningham, F. (2016). The Ensembl Variant Effect Predictor. Genome Biol. *17*, 122.

11. Smedley, D., and Robinson, P.N. (2015). Phenotype-driven strategies for exome prioritization of human Mendelian disease genes. Genome Med. *7*, 81.

12. Robinson, P.N., Köhler, S., Bauer, S., Seelow, D., Horn, D., and Mundlos, S. (2008).
The Human Phenotype Ontology: A Tool for Annotating and Analyzing Human Hereditary
Disease. Am. J. Hum. Genet. *83*, 610–615.

13. Birgmeier, J., Haeussler, M., Deisseroth, C.A., Jagadeesh, K.A., Ratner, A.J., Guturu, H.,
Wenger, A.M., Stenson, P.D., Cooper, D.N., Re, C., et al. (2017). AMELIE accelerates
Mendelian patient diagnosis directly from the primary literature. BioRxiv 171322.

14. Bone, W.P., Washington, N.L., Buske, O.J., Adams, D.R., Davis, J., Draper, D., Flynn,
E.D., Girdea, M., Godfrey, R., Golas, G., et al. (2016). Computational evaluation of exome
sequence data using human and model organism phenotypes improves diagnostic efficiency.
Genet. Med. *18*, 608–617.

15. Köhler, S., Vasilevsky, N.A., Engelstad, M., Foster, E., McMurry, J., Aym?, S., Baynam,
G., Bello, S.M., Boerkoel, C.F., Boycott, K.M., et al. (2017). The Human Phenotype Ontology
in 2017. Nucleic Acids Res. *45*, D865–D876.

16. Leinonen, R., Akhtar, R., Birney, E., Bower, L., Cerdeno-Tárraga, A., Cheng, Y., Cleland,
I., Faruque, N., Goodgame, N., Gibson, R., et al. (2011). The European Nucleotide Archive.
Nucleic Acids Res. *39*, D28-31.

17. Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic
RNA-seq quantification. Nat. Biotechnol. *34*, 525–527.

18. Deelen, P., Zhernakova, D. V, de Haan, M., van der Sijde, M., Bonder, M.J., Karjalainen,
J., van der Velde, K.J., Abbott, K.M., Fu, J., Wijmenga, C., et al. (2015). Calling genotypes
from public RNA-sequencing data enables identification of genetic variants that affect gene-
expression levels. Genome Med. *7*, 30.

19. Fehrmann, R.S.N., Karjalainen, J.M., Krajewska, M., Westra, H., Maloney, D., Simeonov,
A., Pers, T.H., Hirschhorn, J.N., Jansen, R.C., Schultes, E.A., et al. (2015). Gene expression
analysis identifies global gene dosage sensitivity in cancer. Nat. Genet. *47*, 115–125.

20. Zhao, S., Fung-Leung, W.-P., Bittner, A., Ngo, K., and Liu, X. (2014). Comparison of
RNA-Seq and Microarray in Transcriptome Profiling of Activated T Cells. PLoS One *9*,
e78644.

21. Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P., Haw,
R., Jassal, B., Korninger, F., May, B., et al. (2018). The Reactome Pathway Knowledgebase.
Nucleic Acids Res. *46*, D649–D655.

22. Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y., and Morishima, K. (2017). KEGG:
new perspectives on genomes, pathways, diseases and drugs. Nucleic Acids Res. *45*, D353–
D361.

23. The Gene Ontology Consortium (2017). Expansion of the Gene Ontology knowledgebase
and resources. Nucleic Acids Res. *45*, D331–D338.

24. Zaykin, D. V (2011). Optimally weighted Z-test is a powerful method for combining
probabilities in meta-analysis. J. Evol. Biol. *24*, 1836–1841.

25. Kurian, M.A., Zhen, J., Cheng, S.-Y., Li, Y., Mordekar, S.R., Jardine, P., Morgan, N. V.,
Meyer, E., Tee, L., Pasha, S., et al. (2009). Homozygous loss-of-function mutations in the
gene encoding the dopamine transporter are associated with infantile parkinsonism-
dystonia. J. Clin. Invest. *119*, 1595–1603.

26. Puffenberger, E.G., Jinks, R.N., Sougnez, C., Cibulskis, K., Willert, R.A., Achilly, N.P.,
Cassidy, R.P., Fiorentini, C.J., Heiken, K.F., Lawrence, J.J., et al. (2012). Genetic Mapping
and Exome Sequencing Identify Variants Associated with Five Novel Diseases. PLoS One *7*,
e28936.

27. Kurian, M.A., Li, Y., Zhen, J., Meyer, E., Hai, N., Christen, H.-J., Hoffmann, G.F.,
Jardine, P., von Moers, A., Mordekar, S.R., et al. (2011). Clinical and molecular
characterisation of hereditary dopamine transporter deficiency syndrome: an observational
cohort and experimental study. Lancet Neurol. *10*, 54–62.

28. The Gtex Consortium (2013). The Genotype-Tissue Expression (GTEx) project. Nat.
Genet. *45*, 580–585.

29. Benson, M.D. (1991). Inherited amyloidosis. J. Med. Genet. *28*, 73–78.

30. Nouhravesh, N., Ahlberg, G., Ghouse, J., Andreasen, C., Svendsen, J.H., Haunsø, S.,
Bundgaard, H., Weeke, P.E., and Olesen, M.S. (2016). Analyses of more than 60,000
exomes questions the role of numerous genes previously associated with dilated
cardiomyopathy. Mol. Genet. Genomic Med. *4*, 617–623.

31. Arimura, T., Matsumoto, Y., Okazaki, O., Hayashi, T., Takahashi, M., Inagaki, N.,
Hinohara, K., Ashizawa, N., Yano, K., and Kimura, A. (2007). Structural analysis of obscurin
gene in hypertrophic cardiomyopathy. Biochem. Biophys. Res. Commun. *362*, 281–287.

32. Marston, S., Montgiraud, C., Munster, A.B., Copeland, O., Choi, O., dos Remedios, C.,
Messer, A.E., Ehler, E., and Knöll, R. (2015). OBSCN Mutations Associated with Dilated

Cardiomyopathy and Haploinsufficiency. PLoS One *10*, e0138568.

33. Bolling, M.C., Jan, S.Z., Pasmooij, A.M.G., Lemmink, H.H., Franke, L.H., Yenamandra,

V.K., Sinke, R.J., van den Akker, P.C., and Jonkman, M.F. (2018). Generalized Ichthyotic

Peeling Skin Syndrome due to FLG2 Mutations. J. Invest. Dermatol.

34. Alfares, A., Al-Khenaizan, S., and Al Mutairi, F. (2017). Peeling skin syndrome

associated with novel variant in *FLG2* gene. Am. J. Med. Genet. Part A *173*, 3201–3204.

35. Alazami, A.M., Patel, N., Shamseldin, H.E., Anazi, S., Al-Dosari, M.S., Alzahrani, F.,

Hijazi, H., Alshammari, M., Aldahmesh, M.A., Salih, M.A., et al. (2015). Accelerating novel

candidate gene discovery in neurogenetic disorders via whole-exome sequencing of

prescreened multiplex consanguineous families. Cell Rep. *10*, 148–161.

36. Runge, J.S., Raab, J.R., and Magnuson, T. (2018). Identification of Two Distinct Classes

of the Human INO80 Complex Genome-Wide. G3 (Bethesda). *8*, 1095–1102.

37. Meeson, A.P., Radford, N., Shelton, J.M., Mammen, P.P., DiMaio, J.M., Hutcheson, K.,

Kong, Y., Elterman, J., Williams, R.S., and Garry, D.J. (2001). Adaptive mechanisms that

preserve cardiac function in mice without myoglobin. Circ. Res. *88*, 713–720.

38. van der Harst, P., van Setten, J., Verweij, N., Vogler, G., Franke, L., Maurano, M.T.,

Wang, X., Mateo Leach, I., Eijgelsheim, M., Sotoodehnia, N., et al. (2016). 52 Genetic Loci

Influencing Myocardial Mass. J. Am. Coll. Cardiol. *68*, 1435–1448.

39. Truszkowska, G.T., Bilińska, Z.T., Muchowicz, A., Pollak, A., Biernacka, A., Kozar-

Kamińska, K., Stawiński, P., Gasperowicz, P., Kosińska, J., Zieliński, T., et al. (2017).

Homozygous truncating mutation in NRAP gene identified by whole exome sequencing in a

patient with dilated cardiomyopathy. Sci. Rep. *7*, 3362.

40. Sobreira, N., Schiettecatte, F., Valle, D., and Hamosh, A. (2015). GeneMatcher: A

Matching Tool for Connecting Investigators with an Interest in the Same Gene. Hum. Mutat.

36, 928–930.

41. Szklarczyk, D., Morris, J.H., Cook, H., Kuhn, M., Wyder, S., Simonovic, M., Santos, A.,

Doncheva, N.T., Roth, A., Bork, P., et al. (2017). The STRING database in 2017: quality-

controlled protein–protein association networks, made broadly accessible. Nucleic Acids Res.

45, D362–D368.

42. Collado-Torres, L., Nellore, A., and Jaffe, A.E. (2017). recount workflow: Accessing over

70,000 human RNA-seq samples with Bioconductor. F1000Research *6*, 1558.

43. Posafalvi, A., Herkert, J.C., Sinke, R.J., van den Berg, M.P., Mogensen, J., Jongbloed,

J.D.H., and van Tintelen, J.P. (2013). Clinical utility gene card for: dilated cardiomyopathy

(CMD). Eur. J. Hum. Genet. *21*,.

44. Posey, J.E., Harel, T., Liu, P., Rosenfeld, J.A., James, R.A., Coban Akdemir, Z.H.,

Walkiewicz, M., Bi, W., Xiao, R., Ding, Y., et al. (2017). Resolution of Disease Phenotypes

Resulting from Multilocus Genomic Variation. N. Engl. J. Med. *376*, 21–31.

45. Genomics England PanelApp , https://panelapp.genomicsengland.co.uk.

46. Silvester, N., Alako, B., Amid, C., Cerdeño-Tárraga, A., Cleland, I., Gibson, R.,

Goodgame, N., ten Hoopen, P., Kay, S., Leinonen, R., et al. (2015). Content discovery and

retrieval services at the European Nucleotide Archive. Nucleic Acids Res. *43*, D23–D29.

47. Lachmann, A., Torre, D., Keenan, A.B., Jagodnik, K.M., Lee, H.J., Wang, L., Silverstein,

M.C., and Ma 'ayan, A. Massive mining of publicly available RNA-seq data from human and

mouse.

48. Cunningham, F., Amode, M.R., Barrell, D., Beal, K., Billis, K., Brent, S., Carvalho-Silva,

D., Clapham, P., Coates, G., Fitzgerald, S., et al. (2015). Ensembl 2015. Nucleic Acids Res.

43, D662–D669.

49. Vinogradov, A.E. (2003). DNA helix: the importance of being GC-rich. Nucleic Acids Res.

31, 1838–1844.

50. Cronbach, L.J. (1951). Coefficient alpha and the internal structure of tests.

Psychometrika *16*, 297–334.

51. Bresciani, M.J., Oakleaf, M., Kolkhorst, F., Nebeker, C., Barlow, J., Duncan, K., and

Hickmott, J. (2009). Examining Design and Inter-Rater Reliability of a Rubric Measuring

Research Quality across Multiple Disciplines - Practical Assessment, Research &

Evaluation. *14*,.

52. Van Der Maaten, L., and Hinton, G. (2008). Visualizing Data using t-SNE. J. Mach.

Learn. Res. *9*, 2579–2605.

53. Krijthe, J.H. (2015). T-Distributed Stochastic Neighbor Embedding using Barnes-Hut , .

54. Smedley, D., Jacobsen, J.O.B., Jäger, M., Köhler, S., Holtgrewe, M., Schubach, M.,

Siragusa, E., Zemojtel, T., Buske, O.J., Washington, N.L., et al. (2015). Next-generation

diagnostics and disease-gene discovery with the Exomiser. Nat. Protoc. *10*, 2004–2015.

55. Frey, B.J., and Dueck, D. (2007). Clustering by passing messages between data points.

Science *315*, 972–976.