Bed nucleus of the stria terminalis regulates fear to ambiguous threat signals

Travis D. Goode[1], Reed L. Ressler[1], Gillian M. Acca[1], Olivia W. Miles[1], Stephen Maren[1,2]

[1]*Department of Psychological and Brain Sciences and Institute for Neuroscience, Texas A&M University, College Station, TX, 77843-3474, USA*

[2]*Corresponding author*

Email:	maren@tamu.edu
Phone:	979-458-7960
Address:	Stephen Maren
	Department of Psychological and Brain Sciences
	Texas A&M University
	301 Old Main Dr.
	College Station, TX 77843-3474

ORCiD

Travis D. Goode:	https://orcid.org/0000-0003-1432-8894
Reed L. Ressler:	https://orcid.org/0000-0003-0514-8269
Olivia W. Miles:	https://orcid.org/0000-0001-9422-8855
Stephen Maren:	https://orcid.org/0000-0002-9342-7411

ABSTRACT

Anxiety is characterized by a fear of potential future threats. The bed nucleus of the stria terminalis (BNST) has been implicated in anxiety, but its role in processing ambiguous threats is unclear. Here we examined whether the BNST mediates conditioned freezing to conditioned stimuli (CSs) that poorly predict the onset of aversive unconditioned stimuli (USs) in rats. Reversible inactivation of the BNST selectively reduced freezing to CSs that poorly predicted US onset (e.g., a backward CS that followed the US), but did not affect freezing to forward CSs even when they predicted USs of variable intensity. Importantly, backward (but not forward) CSs selectively increased Fos in the ventral BNST and in BNST-projecting neurons in the infralimbic region of the medial prefrontal cortex (mPFC), but not in the hippocampus or amygdala. These data reveal that BNST circuits process ambiguous threat signals central to the etiology and expression of anxiety.

IMPACT STATEMENT

The bed nucleus of the stria terminalis (BNST) is required for the expression of defensive behavior to uncertain threats, a function that is central to pathological anxiety.

INTRODUCTION

Excessive apprehension about potential future threats, including financial loss, illness, or death, is a defining feature of many anxiety disorders, including generalized anxiety disorder (GAD) (Behar et al., 2009). Anxiety and trauma-related disorders are widespread and costly (Blanco et al., 2011; Comer et al., 2011; Kinley et al., 2011; Salas-Wright et al., 2014; Stein et al., 2017; Wittchen, 2002), and remain difficult to treat (Colvonen et al., 2017; Costello et al., 2014; Iza et al., 2013; Sinnema et al., 2015). Understanding the neural circuits underlying anxiety is important for refining behavioral and pharmacotherapeutic treatments (Berridge, 2018; Deslauriers et al., 2017; Fanselow and Pennington, 2018, 2017; Graham et al., 2014; LeDoux and Daw, 2018; Nees et al., 2015; Pine and LeDoux, 2017; Tye, 2018). Several recent studies have demonstrated changes in the function of the bed nucleus of the stria terminalis (BNST) in individuals with post-traumatic stress disorder (PTSD) and anxiety disorders (Andreescu et al., 2015; Brinkmann et al., 2018; L Brinkmann et al., 2017; Leonie Brinkmann et al., 2017; Buff et al., 2017; Münsterkötter et al., 2015; Rabellino et al., 2018; Straube et al., 2007; Yassa et al., 2012). However, the conditions that recruit the BNST to aversive learning and memory processes believed to underlie anxiety disorders are still not understood (Avery et al., 2016, 2014; Ch'ng et al., 2018; Daniel and Rainnie, 2016; Davis et al., 2010; Fox and Shackman, 2017; Goode and Maren, 2017; Gungor and Paré, 2016; Lebow and Chen, 2016; Perusini and Fanselow, 2015; Shackman and Fox, 2016; Walker et al., 2009; Walker and Davis, 2008).

Early work on this question revealed that BNST lesions in rats impair defensive behaviors evoked by unconditioned threats (Gewirtz et al., 1998). For example, BNST inactivation results in a loss of unconditioned defensive responding to the presence of predator odors (Breitfeld et al., 2015; Fendt et al., 2003; Xu et al., 2012). Additionally, unconditioned increases in the acoustic

startle reflex produced by either intracranial administration of corticotropin-releasing factor (CRF)

or exposure to bright light require the BNST, whereas fear-potentiated startle to punctate

conditioned stimuli (CSs) do not (Walker et al., 2009). However, the involvement of the BNST in

defensive responding is not restricted to unconditioned threat: BNST lesions produce deficits in

both freezing and corticosterone release elicited by contextual, but not auditory, CSs after

Pavlovian fear conditioning in rats (LeDoux et al., 1988; Sullivan et al., 2004). Interestingly,

however, it is not stimulus modality that differentiates the effects of BNST lesions on tone and

context freezing. BNST lesions impair freezing responses to long-duration (e.g., 10-min) auditory

conditioned stimuli (CSs) (Waddell et al., 2006), and spare freezing to short-duration (1-min)

contextual stimuli (Hammack et al., 2015). Based on this work, it has been proposed that the BNST

is required to organize behavioral and hormonal responses to sustained threats (whether

conditioned or unconditioned) (Hammack et al., 2015, 2009; Takahashi, 2014; Waddell et al.,

2008, 2006). Of course, the duration of the behavioral responses in these situations is confounded

with the duration of the eliciting stimulus. Thus, it has been argued that the BNST mediates

sustained defensive *responses* independent of the specific features and durations of the stimuli that

evoke them (Davis, 2006; Davis et al., 2010; Walker et al., 2009; Walker and Davis, 2008).

Of course, another factor that differentiates discrete tones from contexts, or short- from

long-duration stimuli (whether tones or contexts), is the temporal information the stimuli provide

about when an aversive event will occur (Goode and Maren, 2017). For example, long-duration

CSs or contexts provide relatively poor information about *when* a future US will occur (i.e., the

US will occur at some time in the future), whereas short-duration CSs or contexts provide more

certainty about the temporal imminence of US onset (i.e., the US will occur soon). We and others

have proposed that the BNST may have a critical role in processing temporally ambiguous threats,

particularly in mediating defensive responses to threatening stimuli that poorly predict when an

aversive event will occur (Goode and Maren, 2017; Lange et al., 2017; Luyck et al., 2018a).

Consistent with this view, punctate auditory CSs that are followed by shock at unpredictable

latencies yield freezing responses that are sensitive to BNST manipulations [(Lange et al., 2017);

also, see (Daldrup et al., 2016)]. This suggests that a crucial parameter that determines the role for

the BNST in defensive behavior is neither the duration nor modality of the threat (nor the duration

of the elicited defensive response), but rather the information a signal provides about when an

aversive event will occur.

Here we directly examined this possibility by investigating the role of the BNST in

conditioned freezing responses elicited by procedures that equated both the duration and modality

of the threat CSs, as well as the total number of unconditioned stimulus (US) presentations, but

differed according to the timing of the aversive US in relation to the CS. Specifically, we arranged

a brief (10-sec) auditory CS to either precede (forward conditioning, FW) or follow (backward

conditioning, BW) a footshock US in rats. Although extensively-trained BW CSs become

conditioned inhibitors that dampen responding to other first-order excitatory cues (Andreatta et

al., 2012; Ayres et al., 1976; Christianson et al., 2011; Gerber et al., 2014; Moscovitch and

LoLordo, 1968; Siegel and Domjan, 1971), minimally-trained BW CSs elicit excitatory

conditioned responses that transfer across contexts (Ayres et al., 1987; Barnet and Miller, 1996;

Bevins and Ayres, 1992; Chang et al., 2003; Connor et al., 2017; Heth, 1976; Mahoney and Ayres,

1976; Prével et al., 2018, 2016; Rescorla, 1968). After conditioning, we examined the effect of

pharmacological inactivation of the BNST on freezing to FW or BW CSs. We hypothesized that

pharmacological inactivation of the BNST would selectively disrupt fear expression to the BW CS

and that the presentation of the BW CS would recruit BNST neurons (as assessed by Fos) in greater

numbers than the FW CS. Furthermore, we anticipated that the backward CS would increase the

activity of BNST-projecting neurons in afferent brain regions implicated in anxiety states.

Consistent with our hypothesis, we found that BNST inactivation reduced freezing to a BW, but

not FW, CS. This effect was related to the ambiguity of the BW CS in predicting US onset, because

the same outcome was observed when USs occurred randomly with respect to the CS. We also

observed that BW CSs selectively increased Fos expression in the BNST and BNST-projecting

medial prefrontal cortical (mPFC) neurons. These data reveal that BNST circuits process the

expression of defensive behavior in the presence of ambiguous threat signals.

RESULTS

Reversible inactivation of the BNST attenuates fear to backward, not forward, CSs. To examine

the role of the BNST in threat uncertainty, we reversibly inactivated the BNST during retrieval of

fear to either a forward- ("FW"; imminent threat) or backward-trained ("BW"; ambiguous threat)

CS. A schematic of the behavioral design is shown in Figure 1A. Representative cannula tracts

and histological placements are presented in Figure 1—Figure Supplement 1 and Figure 1—Figure

Supplement 2 (respectively). Freezing behavior during the conditioning session is shown in Figure

1B. A main effect of conditioning trial was observed (repeated measures: $F_{6,288} = 37.87$, $p <$

0.0001). No other significant main effects or interactions were observed for any of the conditioning

or drug assignments across the conditioning trials (F's < 0.70, p's > 0.60).

One day after conditioning, the animals were infused with NBQX, an AMPA receptor

antagonist, to reversibly inactivate the BNST; saline ("VEH") infusions served as a control.

Immediately after the infusions, the rats were placed in a novel context and received twelve

presentations of the CS [some BW-trained rats received no CS exposure at the test, "NoCS(Test)"].

As shown in (Figure 1B), inactivation of the BNST produced a selective decrease in conditioned

freezing to the BW CS. Analysis of freezing behavior across the entire session (including the

baseline) revealed a main effect of trial (repeated measures: $F_{6,264} = 8.06$, $p < 0.0001$), a significant

main effect of CS exposure ($F_{2,44} = 25.38$, $p < 0.0001$), and a significant CS × drug interaction

($F_{2,44} = 3.66$, $p < 0.05$). There were no other main effects or interactions (F's < 1.6, p's > 0.05).

Fisher's PLSD indicated that rats in the FW condition exhibited significantly more freezing during

the retrieval test than rats in all other groups (p's < 0.05). Importantly, BW-VEH rats exhibited

significantly higher levels of freezing than BW-NBQX rats ($p < 0.005$); BW-NBQX rats did not

significantly differ from the NoCS(Test) groups.

Figure 1. *Reversible inactivation of the BNST attenuates conditioned fear expression to a backward, but not forward, CS.* **(A)** Behavioral schematic. **(B)** Freezing behavior at conditioning and retrieval testing. For conditioning, the left panel depicts mean percentage freezing during the 5-min baseline (BL) and across each conditioning block (each 138-sec block is comprised of two trials; conditioning trials consist of freezing during the 10-sec CS followed by the 58-sec interstimulus interval). For retrieval testing, the center panel shows mean percentage freezing at the 5-min baseline (BL) and across each test block (each 140-sec block is comprised of two trials; trials consist of freezing during the 10-sec CS followed by the 60-sec interstimulus interval). The right panel shows mean percentage freezing during the first half of the test (trials 1-6; corresponding to 420 sec of behavior). All data are represented as means ± s.e.m [FW-VEH (*n* = 5); FW-NBQX (*n* = 4); BW-VEH (*n* = 13); BW-NBQX (*n* = 12); NoCS(Test)-VEH (*n* = 8); NoCS(Test)-NBQX (*n* = 8)]; * = $p < 0.05$.

Given that freezing to the BW CS in the VEH-treated animals was maximal in the first half

of the test, a separate factorial ANOVA was performed on the average percentage of freezing

during trials 1-6 (Figure 1B). For this period, there was a main effect of CS treatment ($F_{2,44} =$

18.61, $p < 0.0001$) as well as a CS treatment × drug interaction ($F_{2,44} = 3.81$, $p < 0.05$); there was

no main effect of drug across all groups ($F < 3.00$, $p > 0.05$). Fisher's PLSD revealed significant

differences in NBQX- and VEH-treated animals in the BW group ($p < 0.0005$). Additionally, rats

in the NoCS(Test) groups differed from all others (p's < 0.0005) except those in the BW-NBQX

group, indicating selective reduction of CS-elicited freezing in the BW-NBQX animals. It has been

suggested that the BNST mediates sustained but not acute fear responses (Davis, 2006; Davis et

al., 2010; Walker et al., 2009; Walker and Davis, 2008). However, it is notable that in this

experiment BNST inactivation selectively attenuated conditioned freezing to the backward CS,

which produced less sustained freezing during the retrieval test compared to the FW CS.

116   The different effects of BNST inactivation on freezing to FW and BW CSs may have been

related to differences in memory strength between the two CSs (reflected in overall levels of

freezing to each CS across the retrieval test). Indeed, to generate conditioned freezing to a

backward CS we used more conditioning trials than typical in our laboratory. Of course, this may

have resulted in moderate overtraining of the FW CS, which might have rendered the FW response

less sensitive to disruption. To examine this possibility, we also examined the effects of BNST

inactivation of FW and BW conditioning after five conditioning trials (Figure 1—Figure

Supplement 3). The behavioral design for the five-trial experiment is shown in Figure 1—Figure

Supplement 3A; cannula placements for this experiment are shown in Figure 1—Figure

Supplement 4.

126   During the 5-trial conditioning procedure (Figure 1—Figure Supplement 3B), animals

exhibited reliable increases in freezing across the session (repeated measures: $F_{5,115}$ = 12.154, $p <$

0.0001; there were no main effects of conditioning procedure or drug treatment and no interaction

of these variables: F's < 2.3, p's > 0.15). After infusions of NBQX or VEH, animals were tested

to the CS in a familiar context that was distinct from the conditioning context. As shown in Figure

1—Figure Supplement 3B, BNST inactivation did not affect conditioned freezing to either the FW

or BW CS. An ANOVA revealed a main effect of trial (repeated measures [including baseline

freezing]: $F_{5,105}$ = 22.118, p < 0.0001), a main effect of conditioning procedure ($F_{1,21}$ = 15.930, p

< 0.001), and a trial × conditioning procedure interaction (repeated measures: $F_{5,105} = 12.346$, $p <$

0.0001). A separate ANOVA performed on trials 1-5 (excluding baseline; Figure 1—Figure

Supplement 3B) revealed a significant main effect of conditioning procedure ($F_{1,21} = 21.585$, $p =$

0.0001; there was no main effect of drug and no interaction: F's < 0.15, p's > 0.7). Thus, the failure

of BNST inactivation to impair conditioned freezing to the FW CS in the 12-trial procedure does

not appear to be due to a ceiling effect; freezing to the FW CS in the 5-trial procedure was not

maximal and remained insensitive to BNST inactivation. That said, the absence of an effect of

BNST inactivation on freezing to the BW CS in this experiment may be due to a floor effect insofar

as the 5-trial procedure did not yield conditional significant freezing to the CS (at least freezing

that exceeded the pre-CS baseline). Together, these data indicate that the BNST is required for the

expression of conditioned freezing to an excitatory BW, but not FW, CS.

One index that might reveal differences in the ability of the CS to predict the US is the US-

evoked response itself. Footshock USs elicit an unconditioned response (UR) that includes

vocalization, autonomic adjustments, and burst of locomotor activity (Bali and Jaggi, 2015;

Fanselow, 1994). To better understand the mechanisms of forward and backward conditioning on

behavior, we examined shock-evoked activity bursts (Kunwar et al., 2015; Zelikowsky et al., 2018)

during the conditioning session in a separate cohort of animals (Figure 1—Figure Supplement 5).

Fear conditioning resulted in robust freezing in both groups of animals (Figure 1—Figure

Supplement 5A). An ANOVA revealed a significant main effect of trial (repeated measures: $F_{6,180}$

$= 67.02$, $p < 0.0001$) with no differences between levels of freezing in FW or BW animals (no

other main effects and no interactions: F's < 1.80, p's > 0.15). In contrast, shock-induced activity

differed in FW and BW animals (Figure 1—Figure Supplement 5C). All animals exhibited a

reliable decrease in shock-induced activity across the conditioning session ($F_{5,150} = 9.59$, $p <$

0.0001) and the rate of decline in activity was similar in the two groups (no trial × group

interaction: $F < 1.70$, $p > 0.09$), but the overall level of shock-induced activity was significantly

lower in the FW animals (main effect of group: $F_{1,30} = 9.59$, $p < 0.05$). For comparison, two-tailed

unpaired t-test of mean levels of activity during the 5-min baseline revealed no significant

difference between FW and BW rats (Figure 1—Figure Supplement 5B; $t < 0.30$; $p > 0.75$). Hence,

USs that were not signaled by a forward CS evoked greater activity bursts than those that were,

which may suggest greater regulation of the shock-induced activity when the US is preceded by a

temporally predictive cue.

Inactivation of the BNST disrupts freezing to a discrete CS associated with random USs. The

differential effect of BNST inactivation on freezing to a FW or BW CS suggests that the BNST

regulates defensive behavior to stimuli that poorly signal US onset. However, the BW CS does not

completely lack temporal information about US onset; the BW conditioning procedure may have

resulted in forward trace conditioning to the CS, which always preceded the next US by 60-sec

(after trial 1). To address this possibility, a separate cohort of rats was implanted with cannulas in

the BNST and submitted to one of three conditioning procedures (Figure 2). In the first group,

animals were conditioned to a backward CS (identical to 12-trial experiment above; "BW"). In the

second group, animals received CS presentations followed by randomized onset of the US

("RANDOM"). In the third control group, animals were conditioned using the RANDOM

procedure but in the absence of any CS ["NoCS(Cond)"]. A summary of the behavioral design is

shown in Figure 2A. Cannula placements for these animals are shown in Fig. 2—Figure

Supplement 1.

The percentage of freezing during fear conditioning for the RANDOM, BW, and

NoCS(Cond) procedures is shown in Figure 2B. For conditioning, an ANOVA revealed a main

effect of trial (repeated measures: $F_{6,204} = 85.209$, $p < 0.0001$), insofar as all groups reliably

increased their freezing across the conditioning trials. There was no main effect of conditioning

procedure ($F < 3.10$, $p > 0.05$) and no main effect or interactions with regard to future drug or

vehicle assignments (F's < 1.00, p's > 0.30).

Figure 2. *BNST inactivation attenuates freezing to a discrete CS paired with random onset of shock.* **(A)** Behavioral schematic. **(B)** Freezing behavior during conditioning and retrieval testing. For conditioning, the left panel depicts mean percentage freezing during the 5-min baseline (BL) and across each conditioning block (each block is comprised of two trials; trials consist of freezing during the 10-sec CS followed by the 58-sec interval for the BW animals, whereas blocks for the other groups include equivalent time periods which include the CS [if present] and inter-CS intervals [excluding the duration of the shock]). For retrieval testing, the center panel shows mean percentage freezing at the 5-min baseline (BL) and across each test block (each 140-sec block is comprised of two trials; trials consist of freezing during the 10-sec CS followed by the 60-sec interval). The right panel shows mean percentage freezing across the first two test trials (after BL; corresponding to 140 sec of behavior). All data are represented as means ± s.e.m [RANDOM-VEH (*n* = 6); RANDOM-MUS (*n* = 9); BW-VEH (*n* = 5); BW-MUS (*n* = 7); NoCS(Cond)-VEH (*n* = 7); NoCS(Cond)-VEH (*n* = 6)].

As shown in Figure 2B, inactivation of the BNST produced deficits in freezing during

retrieval testing in the BW and RANDOM groups, but did not affect freezing in the NoCS(Cond)

control. This was confirmed in an ANOVA that revealed a significant drug × trial interaction

(repeated measures: $F_{6,204} = 7.91$, $p < 0.0001$). A significant main effect of trial (repeated measures:

$F_{6,204} = 7.80$, $p < 0.0001$) indicated significant changes in freezing across the entire test amongst

all groups. No other significant main effects or interactions were detected for this test (F's < 1.80,

p's > 0.05). Similar to the aforementioned experiments, freezing was maximal in the early test

trials, so a separate ANOVA was performed on mean responding during the first two trials of

testing (Figure 2B). These analyses revealed a significant main effect of drug ($F_{1,34} = 6.217$, $p <$

0.05), indicating a disruption of freezing in drug-treated animals; no other significant main effects

or interactions were detected for these trials (F's < 3.00, p's > 0.07). Excluding NoCS(Cond) animals, a main effect of drug treatment was also identified for these trials ($F_{1,23}$ = 28.115, p < 0.0001). Additionally, we ran an ANOVA solely on baseline freezing. MUS-treated animals exhibited a trending, but nonsignificant ($F < 4.1, p > 0.05$), decrease in freezing during the baseline [no other main effect or interact was detected: (F's < 1.00, p's > 0.30)]. While baseline freezing may impact CS-responding, subtracting baseline responding from CS-elicited freezing (of the first two trials for RANDOM and BW animals) did not eliminate a main effect a drug-treatment ($F_{1,23}$ = 5.877, $p < 0.05$). Thus, these data replicate the effects of BNST inactivation on BW CS-elicited freezing as seen in our first experiment and extend these results to show that the BNST mediates freezing to CSs that signal uncertain latency of shock onset.

Temporary inactivation of the BNST does not eliminate fear to a forward CS that is paired with a US of variable intensity. To determine whether the BNST is involved in other conditioning procedures imbued with outcome (but not temporal) uncertainty, we examined whether freezing to a FW CS that is paired with a US of variable intensity is also BNST-dependent. In this case, rats received forward fear conditioning with either a fixed ("FIXED") or variable ("VARIABLE") US intensity (Figure 3). A schematic of the behavioral design is shown in Figure 3A; cannula placements are illustrated in Figure 3—Figure Supplement 1. During conditioning (Figure 3B), a repeated measures ANOVA revealed a main effect of trial (repeated measures: $F_{6,168}$ = 68.15, p < 0.0001), with no main effect of drug or conditioning procedure, and no interactions (F's < 1.50, p's > 0.20).

Twenty-four hours after conditioning and immediately before a test to the CS in a novel context (Figure 3B), the animals were infused with either muscimol ("MUS") to reversibly inactivate the BNST; saline ("VEH") infusions served as a control. As shown in Figure 3B, BNST

inactivation did not affect freezing to a FW CS paired with either a fixed or variable US. During

retrieval testing, there was a main effect of trial (repeated measures: $F_{6,156} = 24.31$, $p < 0.0001$),

however no other main effects or interactions were detected (F's < 2.70, p's > 0.15). Although

average baseline freezing was highest in VARIABLE-VEH animals, an ANOVA of baseline

freezing did not reveal any main effects or interactions (F's < 3.60, p's > 0.05). To equate for

differences in pre-CS freezing in the experimental groups, baseline freezing was subtracted from

CS-elicited freezing (Figure 3C). This did not reveal an effect of BNST inactivation on freezing;

there were no significant main effects or interactions in the ANOVA (F's < 0.15, p's > 0.70).

Although BNST inactivation trended towards reducing generalized contextual freezing in this

experiment, it did not eliminate CS-elicited freezing to the temporally predictable cue, regardless

of the variable magnitude of the US.

Figure 3. *Temporary inactivation of the BNST does not prevent conditioned fear expression to a forward CS paired with a US of fixed or variable intensity.* **(A)** Behavioral schematic. **(B)** Freezing behavior during conditioning and retrieval testing. For conditioning, the left panel depicts mean percentage freezing during the 5-min baseline (BL) and across each conditioning block (each 138-sec block is comprised of two trials; trials consist of freezing during the 10-sec CS followed by the 58-sec post-shock interval). For retrieval testing, the center panel shows mean percentage freezing at the 5-min baseline (BL) and across each test block (each 140-sec block is comprised of two trials; trials consist of freezing during the 10-sec CS followed by the 60-sec interval). The right panel shows mean percentage freezing across all test trials (after BL; corresponding to 840 sec of behavior). **(C)** Freezing percentages across all twelve trials, with BL levels of freezing subtracted from these values. All data are represented as means ± s.e.m [FIXED-VEH ($n = 7$); FIXED-MUS ($n = 8$); VARIABLE-VEH ($n = 7$); VARIABLE-MUS ($n = 8$)].

Backward CSs selectively increase Fos expression in the ventral BNST. To further examine the

role of the BNST in the expression of conditioned fear, we quantified Fos expression in multiple

subregions of the BNST following the presentation of either a FW or BW CS during a shock-free

retrieval test (Figure 4). The behavioral design is summarized in Figure 4A. Four experimental

groups were compared: rats conditioned and tested to a forward CS ("FW"), rats conditioned and

tested to a backward CS ("BW"), rats conditioned to a forward or backward CS but not receiving

a CS at test ["NoCS(Test)"], and animals that were conditioned but not tested ("NoTest").

Conditioning (Figure 4B) was similar to previous experiments (main effect of trial; repeated

measures: $F_{6,258} = 77.346$, $p < 0.0001$; no other main effects or interactions: F's < 1.30, p's > 0.30).

As shown in Figure 4B, the groups differed in their levels of conditioned freezing during the

retrieval test. ANOVA revealed a main effect of trial (repeated measures: $F_{6,216} = 15.54$, $p <$

0.0001), conditioning procedure ($F_{2,36} = 11.42$, $p < 0.0001$), and a procedure × trial interaction

($F_{6,216} = 3.65$, $p < 0.0001$) (includes baseline). Post-hoc analyses revealed that FW ($p < 0.0001$)

and BW ($p < 0.005$) rats exhibited significantly higher levels of freezing behavior than NoCS(Test)

rats. Similarly, an ANOVA of average freezing across the test trials (without baseline) revealed a

main effect of group ($F_{2,36} = 13.76$, $p < 0.0001$). Post-hoc comparisons revealed significant

differences between FW vs. BW ($p < 0.05$) and FW vs. NoCS(Test) rats ($p < 0.0001$), as well as

BW vs. NoCS(Test) animals ($p < 0.005$).

Ninety minutes after the retrieval test, the animals were sacrificed for Fos

immunohistochemistry. Fos-positive nuclei were counted in three BNST subregions (Figure 5A):

"ovBNST" (Fos counts targeting the oval nucleus of the BNST), "dBNST" (counts in an area

containing the dorsal region of the anteromedial BNST), and "vBNST" [Fos counts in a region

containing the BNST's anterolateral, fusiform, and anteromedial (ventral) nuclei; refer to

(Swanson, 2003)]. As shown in Figure 5B, the average number of Fos-positive nuclei for each of

these regions differed in rats undergoing FW or BW conditioning in the ovBNST and vBNST, but

not the dBNST. In the vBNST, an ANOVA revealed a main effect of group ($F_{3,43} = 11.41$, $p <$

0.0001). Fisher's PLSD revealed that BW rats exhibited significantly higher levels of Fos

compared to FW ($p < 0.0005$), NoCS(Test) ($p < 0.001$), and NoTest ($p < 0.0001$) rats. Additionally,

NoCS(Test) rats exhibited significantly greater Fos expression than NoTest rats ($p < 0.05$). In the

ovBNST, an ANOVA revealed a main effect of group ($F_{3,43} = 3.26$, $p < 0.05$). Post-hoc analyses

revealed that NoTest rats exhibited significantly higher Fos levels compared to FW ($p < 0.05$) and

No-CS ($p < 0.01$) rats. In dBNST, there was no effect of conditioning procedure on Fos expression

(factorial ANOVA: $F < 2.60$, $p > 0.07$). Together, these data indicate that presentation of a BW

CS increases Fos expression in the vBNST. Moreover, exposure to the temporally predictive FW

CS (which elicited the highest levels of fear) was associated with low levels of Fos in the BNST.

Figure 4. *CS-evoked freezing in rats utilized for Fos analyses.* **(A)** Behavioral schematic. **(B)** Freezing during conditioning and retrieval testing. For conditioning, the left panel depicts mean percentage freezing during the 5-min baseline (BL) and across each conditioning block (each block is comprised of two trials; conditioning trials consist of freezing during the 10-sec CS followed by the 58-sec interstimulus interval). For retrieval testing, the center panel shows mean percentage freezing at the 5-min baseline (BL) and across each test block (each block is comprised of two trials; trials consist of freezing during the 10-sec CS followed by the 60-sec interval). The right panel shows mean percentage freezing after BL. Animals were sacrificed for Fos analyses 90 min after trial 1. All data are represented as means ± s.e.m [FW ($n = 8$); BW ($n = 14$); NoCS(Test) ($n = 17$); NoTest ($n = 8$)]; * = $p < 0.05$.

Backward CSs selectively increase Fos expression in mPFC afferents of the BNST. The BNST

receives input from many areas involved in the regulation of fear (Fox and Shackman, 2017; Goode

and Maren, 2017; Lebow and Chen, 2016). Therefore, we used a functional tracing procedure to

quantify Fos expression in neurons targeting the BNST. Specifically, rats were injected with a

retrograde tracer (CTb-488) into the BNST (prior to conditioning) so that we could quantify Fos

expression in BNST-projecting neurons in its major limbic and cortical afferents. We quantified

Fos in the infralimbic (IL) and prelimbic (PL) regions of the medial prefrontal cortex (mPFC), the

basolateral amygdala (BLA), and the ventral hippocampus (HPC). The behavioral data for these

animals are shown in Figure 4.

Figure 5. *Fos expression in the BNST following exposure to a temporally predictable or uncertain CS.* **(A)** Schematic depicting regions counted within the BNST (left panel). The right panel shows example of Fos expression in the ventral BNST (vBNST). **(B)** Mean Fos-positive cells per 0.1 mm^2 for each of the quantified regions. All data are represented as means ± s.e.m [FW ($n = 8$); BW ($n = 14$); NoCS(Test) ($n = 17$); NoTest ($n = 8$)]; * = $p < 0.05$.

A representative image of CTb infusion into the BNST is shown in Figure 6A and an

illustration of the largest and smallest CTb spread of injection included in the analyses is shown

in Figure 6B. Approximate microinjection sites for CTb for all animals are shown in Figure 6—

Figure Supplement 1. Representative Fos and CTb co-labeling is shown in Figure 6C. As shown

in Figure 6D, there were reliable differences in CTb labeling among the BNST afferents we

quantified. An ANOVA revealed a significant main effect of region ($F_{3,116} = 42.34$, $p < 0.0001$).

Post-hoc comparisons indicated that the number of CTb-positive cells in the HPC were

significantly higher than that in all other regions ($p < 0.0001$, per comparison); IL and BLA

exhibited significantly greater CTb counts compared to PL ($p < 0.0001$, per comparison). CTb

counts in each region did not differ by behavioral condition (F's < 0.5, p's > 0.75) and were

collapsed for Figure 6D. These data indicate extensive connectivity of the PFC, BLA, and HPC

with the BNST.

Figure 6. *Functional tracing in afferents targeting the BNST.* **(A)** Coronal section (10×) showing representative fluorescence of a CTb infusion (green) into the BNST (dotted outline; "ac" = anterior commissure, "lv" = lateral ventricle). Black "X" denotes approximate lowest point of the infusion (as an example of how injection sites are documented in Figure 6—Figure Supplement 1). **(B)** Coronal schematic (-0.26 mm from bregma) showing the approximate largest (green) and smallest (dark green) areas of CTb spread in the BNST for animals included in the analyses; the black dotted outline represents the extent of spread of CTb in the BNST in the image shown in panel **A**. **(C)** Example CTb-positive (green) and Fos-positive (red; nuclei) cells in a coronal section (40 μm) of the IL; open white squares denote double-labeled cells. **(D)** Mean number of BNST-targeting CTb-positive cells (per 0.1 mm^2) for each of the quantified regions (shows FW, BW, and NoTest animals corresponding to Figure 8). All data are represented as means ± s.e.m (for each region, n = 30); * = $p < 0.05$.

As shown in Figure 7A, the number of Fos-positive nuclei in the IL, PL, HPC, and BLA

differed among the behavioral groups. Factorial ANOVA of Fos counts in IL revealed a main

effect of group ($F_{2,27}$ = 8.55, $p < 0.005$). Post-hoc comparisons for IL revealed significant

differences between BW vs. NoTest rats ($p < 0.0005$) and FW vs. NoTest rats ($p < 0.05$). In PL, a

main effect of group was also identified ($F_{2,27}$ = 6.79, $p < 0.005$). Post-hoc analyses indicated that

BW rats exhibited significantly more Fos in PL as compared to NoTest animals ($p < 0.001$). For

BLA, a main effect of group was detected (factorial ANOVA: $F_{2,27}$ = 8.10, $p < 0.005$). Post-hoc

analyses revealed that FW and BW rats (which did not significantly differ) exhibited significantly

more Fos in BLA as compared to No Test rats ($p < 0.01$, FW vs. NoTest; $p < 0.001$, BW vs.

NoTest). In HPC, an ANOVA revealed a significant main effect of group ($F_{2,27}$ = 4.28, $p < 0.05$).

Post-hoc comparisons indicated that FW and BW rats (again, which did not significantly differ)

had significantly more Fos expression in HPC as compared to NoTest animals ($p < 0.05$ for FW

vs. NoTest and BW vs. NoTest). These data indicate that conditioned freezing to forward or

backward CSs increased the number of Fos-positive neurons in the PFC, BLA, and HPC.

To quantify the fraction of BNST-projecting neurons within the PFC, BLA, and HPC that

were activated by a FW or BW CS, we calculated a ratio of Fos-positive to CTb-positive nuclei in

each afferent region ("Fos-CTb%"; Figure 7B). Interestingly, within the IL, animals in the BW

group exhibited the highest fraction of double-labeled cells. An ANOVA revealed a significant

main effect of group ($F_{2,27} = 14.22$, $p < 0.0001$). Post-hoc comparisons revealed significant

differences between BW and FW rats ($p < 0.005$) as well as BW and NoTest animals ($p < 0.0001$).

In contrast, there were no differences between FW and BW rats in any of the other regions

quantified. In PL, there was not a significant main effect of group ($F < 1.90$, $p > 0.17$) and no

group effects were observed in the BLA ($F < 2.0$, $p > 0.15$). In the HPC, rats in the FW and BW

groups exhibited greater numbers of double-labeled cells than those in the NoTest group ($F_{2,27} =$

7.82, $p < 0.01$). Additionally, we examined whether freezing significantly correlates with any of

the aforementioned Fos measures; gross measures of freezing percentages (trials 1-12) for FW and

BW rats did not significantly correlate with Fos levels across the multiple regions (data not shown

in figures), suggesting that the presence or absence of the fear CS was more predictive of activity

in these circuits than freezing performance *per se*. Nonetheless, these data indicate that conditioned

freezing (in both FW and BW rats) is associated with increased activity in BNST-targeting cells

of HPC, and that BW CSs selectively increase Fos in BNST-projecting neurons in IL. Hence, IL

projections to the BNST may regulate the effects of BW CSs on BNST Fos expression and

freezing.

Figure 7. *Fos expression in BNST-targeting cells of the prefrontal cortex, amygdala, and hippocampus following exposure to a forward or backward CS.* **(A)** Mean number of Fos-positive cells (per 0.1 mm^2) for each of the quantified regions. **(B)** Mean percentage (per 0.1 mm^2) of Fos-positive and CTb-positive cells divided by the total number of CTb-positive cells for each region. All data are represented as means ± s.e.m [FW ($n = 8$); BW ($n = 14$); NoTest ($n = 8$)]; * = $p < 0.05$.

DISCUSSION

In the present study, we demonstrate that the BNST is critical for processing ambiguous threats, specifically discrete auditory CSs that are poor predictors of when aversive footshocks will occur. Reversible inactivation of the BNST inactivation disrupted CS-elicited freezing to a backward CS, but did not prevent freezing to a CS that reliably signaled the imminent onset of shock, even when that CS signaled an unpredictable intensity of the US. Moreover, pharmacological blockade of the BNST attenuated fear to both a backward CS and a CS that was trained with random US onset. This data suggests that uncertain timing of shock is a critical factor in recruiting the BNST to conditioned fear. Moreover, BNST inactivation did not affect freezing to a forward CS that was

paired with a US that varied in intensity, suggesting that contributions of the BNST to conditioned

fear may depend on the type of uncertainty imbued by the CS. Interestingly, freezing to the

backward CS was less sustained than that to the forward CS, suggesting that neither long-duration

stimuli nor sustained freezing responses are required to recruit the BNST. Finally, although both

forward and backward CSs increased Fos expression in the hippocampus, BNST, amygdala, and

mPFC, only backward CSs selectively increased Fos expression in both the ventral BNST and

BNST-projecting neurons in the infralimbic cortex. Collectively, these data reveal that the BNST

and its mPFC afferents may mediate defensive responding to threats that signal uncertain or remote

onset of shock.

Unpredictable threats come in many forms [including uncertain timing, magnitude or type,

or probability of threat: (Bennett et al., 2018; Daldrup et al., 2015; Davies and Craske, 2015;

McHugh et al., 2015; McNally et al., 2011; Schroijen et al., 2016; Seidenbecher et al., 2016)];

these subtypes of uncertainty may have dissociable and cumulative contributions to anxiety. In

particular, uncertainty about the timing of the onset of aversive events has been linked to measures

of anxiety in humans and other animals (Amadi et al., 2017; Bennett et al., 2018; Shankman et al.,

2011). Interestingly, ambiguous threats (of varying types and degrees) are associated with

increases or changes in regional cerebral blood flow and functional connectivity in the primate

BNST (Alvarez et al., 2015, 2011; Brinkmann et al., 2018; L Brinkmann et al., 2017; Leonie

Brinkmann et al., 2017; Buff et al., 2017; Choi et al., 2014, 2012; Coaster et al., 2011; Fox et al.,

2018; Grupe et al., 2013; Herrmann et al., 2016; Kalin et al., 2008, 2005, Klumpers et al., 2017,

2015; McMenamin et al., 2014; Meyer et al., 2018; Mobbs et al., 2010; Münsterkötter et al., 2015;

Naaz et al., 2019; Pedersen et al., 2018, 2017; Schlund et al., 2013; Sladky et al., 2018; Somerville

et al., 2010; Torrisi et al., 2018). In turn, BNST lesions or circuit inactivations reduce defensive

responding to contextual cues that poorly signal the imminence of the aversive US [(Asok et al.,

2018; Davis and Walker, 2014; Goode et al., 2015; Hammack et al., 2015; Luyten et al., 2011;

Poulos et al., 2010; Sullivan et al., 2004; Waddell et al., 2006; Zimmerman and Maren, 2011); but,

see (Luyck et al., 2018b)]. Consistent with the possibility that the BNST mediates conditioned fear

to temporally uncertain threats, we observed significant and selective decreases in the expression

of defensive freezing when we inactivated the BNST in the presence of BW CSs or CSs trained

with randomized USs. Conversely, cues that predicted imminent shock were not affected by BNST

inactivation. Additionally, we observed significant differences in shock reactivity during forward

and backward conditioning, insofar as rats exhibited more robust shock-elicited activity bursts than

during forward conditioning. These data may suggest that conditioning in the presence of

predictive cues may reduce the magnitude of the unconditioned response, in a way that is not

present for the backward CS.

A critical finding in the present study is that the BNST mediates conditioned freezing to

ambiguous short-duration discrete CSs. This contrasts with prior work suggesting that the BNST

only mediates freezing behavior to long-duration cues or contexts (Hammack et al., 2015; Waddell

et al., 2006). However, recent data in other paradigms suggest a role for the BNST in the processing

of responses to relatively brief threat stimuli (Brinkmann et al., 2018; Choi et al., 2014; Haufler et

al., 2013; Kinnison et al., 2012; Kiyokawa et al., 2015; Klumpers et al., 2017; Luyck et al., 2018b,

2018a; Naaz et al., 2019). Together, these data suggest that stimulus duration is not necessarily a

determinant of BNST involvement. Similarly, it does not appear that response duration is a

determinant of BNST in threat reactions. In the present work, conditioned freezing to the backward

CS was generally shorter-lived than that to the forward CS, but was more sensitive to BNST

inactivation than freezing to the forward CS. One might conclude from this observation that the

BNST mediates weak (but not strong) freezing responses. However, we and others have shown

that BNST lesions or inactivation reduce the expression of freezing even when those levels are

high (Goode et al., 2015; Hammack et al., 2015). Hence, the BNST mediates fear to cues that

signal ambiguous or remote threat, independent of the duration of the antecedent stimuli or

behavioral responses associated with the threat (Goode and Maren, 2017).

Interestingly, it has been suggested that fear to a backward CS relies on associations

between the CS and the conditioning context (Chang et al., 2003), which more strongly predicts

the US. That is, extinction of fear to the conditioning context reduces fear to a BW, but not FW,

CS. Thus, it is possible that the BNST mediates fear expression to a backward CS by retrieving a

memory of the context-US association. Indeed, considerable work indicates that BNST is involved

in the expression of contextual fear (Davis and Walker, 2014; Goode et al., 2015; Hammack et al.,

2015; Luyten et al., 2011; Sullivan et al., 2004; Waddell et al., 2006; Zimmerman and Maren,

2011). Increases in immediate early gene expression, metabolic activity, and changes in

electrophysiological firing patterns have been observed in the BNST in response to direct exposure

to conditioned contextual cues (Campeau et al., 1997; Daldrup et al., 2016; Jennings et al., 2013;

Lemos et al., 2010; Luyten et al., 2012; Marcinkiewcz et al., 2016). If the backward CS functions

similarly to a conditioned context, then one might expect Fos activity to be elevated in the BNST

of animals exposed to the backward CS. Indeed, we observed significantly more Fos expression

in the BNST in response to the backward compared to the forward CS. The backward CS-elicited

Fos expression was specific to vBNST, insofar as backward and forward rats did not differ in

overall levels of Fos in dBNST or ovBNST. Similarly, others have shown elevated Fos expression,

in regions in or near vBNST after stress or conditioned fear-related behavior [(Besnard et al., 2019;

Sterrenburg et al., 2012; Verma et al., 2018); also, see (Radley and Johnson, 2018; Radley and

Sawchenko, 2011)].

Of course, another possibility is that the backward conditioning procedure yielded forward

trace conditioning after the first trial (Burman et al., 2014; Marchand et al., 2004; Raybuck and

Lattal, 2014; Tipps et al., 2014). Although there has been considerable work on the neural

mechanisms of trace conditioning (Raybuck and Lattal, 2014), a role for the BNST in this form of

learning has not yet been established. Because trace conditioning also degrades the temporal

imminence of the US following the CS, it might be expected to require the BNST. Consistent with

this possibility, we found that inactivation of the BNST reduced fear in the presence of a cue was

conditioned with randomized onset of shock after the CS. In either event, the BNST appears to be

recruited by stimuli, whether cues or contexts, that signal remote or ambiguous onset of shock.

The amygdala, PFC, and hippocampus have strong connections with the BNST (Canteras

and Swanson, 1992; deCampo and Fudge, 2013; Dong et al., 2001; Glangetas et al., 2017; Johnson

et al., 2018; McDonald et al., 1999; Reichard et al., 2017; Reynolds and Zahm, 2005; Torrisi et

al., 2015; Vertes, 2004; Weller and Smith, 1982; Wood et al., 2018). In line with previous work,

the expression of freezing (whether elicited by a forward or, in our case, backward CS) increased

the number of Fos-positive neurons in these regions (Herry et al., 2008; Jin and Maren, 2015;

Knapska and Maren, 2009; Lemos et al., 2010; Wang et al., 2016). In addition, we found that

forward and backward CSs increased Fos expression in BNST-targeting cells of the HPC (with

similar, but trending, effects within the BLA). HPC and BLA projections to the BNST are thought

to regulate hypothalamic-pituitary-adrenal (HPA) axis activity (Crestani et al., 2013; Forray and

Gysling, 2004; Zhu et al., 2001), and these afferents may be engaged in high-fear states to mobilize

corticosterone release, for example [also, see (Kim et al., 2013)]. However, unlike the HPC and

BLA, we observed a greater number of Fos-positive cells in the IL in response to a backward CS

relative to forward CS. This suggests a unique role for BNST-projecting neurons of the IL [which

appear to be glutamatergic: (Crowley et al., 2016; Glangetas et al., 2017)] in processing excitatory

backwards CSs. There is growing evidence for a regulatory role for prefrontal connections to the

BNST in stress and anxiety-like behavior (Fox et al., 2010; Glangetas et al., 2017, 2013; Johnson

et al., 2018; Kinnison et al., 2012; Motzkin et al., 2015; Naaz et al., 2019; Radley et al., 2009), the

present data adds to this work by suggesting that the IL may be involved in generating freezing

and/or stress responding to the backward CS. This would stand in contrast to considerable data

indicating that the IL is involved in the inhibition of freezing behavior after extinction, for example

(Milad and Quirk, 2002; Quirk and Mueller, 2008). However, the fact that the backward CS elicits

less fear (or more extinguishable fear) could suggest this circuit is active, in part, to gate or regulate

fear expression. Indeed, there is evidence for a role of the infralimbic cortex in regulating

discrimination of conditioned excitors and inhibitors (Sangha et al., 2014). An alternative view is

that the IL is involved in processing contextual information conveyed by the hippocampus

(Corcoran and Quirk, 2007; Marek et al., 2018; Zelikowsky et al., 2013) during threat processing.

To the extent that that backward CSs elicit fear via CS-(context-US) associations (Chang et al.,

2003), it is possible that the IL is involved in retrieving CS-context associations.

In conclusion, we demonstrate a novel role for the BNST and its circuits in processing

discrete cues, including excitatory backward CSs and randomized CSs. Specifically, BNST

inactivation impaired conditioned freezing to a backward, but not forward, auditory CS that

differed only in their temporal relationship to a footshock US. This reveals that neither stimulus

modality nor duration (or response duration) are the critical parameters driving BNST involvement

in defensive behavior. We further show that backward CSs increase the number of Fos-positive

neurons in the vBNST and in BNST-projecting neurons in the infralimbic cortex. This suggests a

novel role for mPFC projections to the BNST in processing ambiguous threats, possibly of the

aversive contexts to which backward CSs are associated. These findings may help in our

understanding of the broader contributions of the BNST to motivated behaviors that depend on

uncertainty, such as during BNST-dependent triggers of fear and drug relapse (Aston-Jones and

Harris, 2004; Avery et al., 2016; Goode et al., 2018; Goode and Maren, 2018; Harris and Winder,

2018; Miles et al., 2018; Silberman and Winder, 2013; Stamatakis et al., 2014). In total, these

experiments provide new insight into the antecedents for BNST-dependent defensive behavior and

highlight behavioral procedures to explore these processes.

MATERIALS AND METHODS

Subjects. All experiments used adult (200-240 g upon arrival; $n = 285$, before exclusions) male

Long-Evans rats (Envigo; Indianapolis, IN). Rats were housed in a climate-controlled vivarium

and kept on a fixed light/dark cycle (lights on starting at 7:00 AM and off at 9:00 PM; experiments

took place during the light phase of the cycle). Rats were individually housed in clear plastic cages

(with bedding consisting of wood shavings; changed weekly) on a rotating cage rack. Group

assignments for behavioral testing was randomized for cage position on the racks. Animals had

access to standard rodent chow and water *ad libitum*. Animals were handled by the experimenter(s)

(~30 sec/day) for five consecutive days prior to the start of any surgeries or behavior. All

procedures were in accordance with the US National Institutes of Health (NIH) Guide for the Care

and Use of Laboratory Animals and were approved by the Texas A&M University Institutional

Animal Care and Use Committee.

Apparatuses. All behavioral testing occurred within distinct rooms in the laboratory. Each

behavioral room housed eight identical rodent conditioning chambers (30 cm × 24 cm × 21 cm;

MED Associates, Inc.). Each chamber was housed in a larger, external sound-attenuating cabinet.

Rear walls, ceilings, and the front doors of the testing chambers were made of Plexiglas, while

their sidewalls were made of aluminum. Grid floors of the chambers were comprised of nineteen

stainless steel bars (4 mm in diameter), and spaced 1.5 cm apart (center to center). The grid floors

were attached to an electric shock source and a solid-state grid scrambler for delivery of the US

(MED Associates, Inc.). A speaker attached to each chamber was used to deliver the auditory CS.

As needed for each context, the chambers were equipped with 15 W house lights, and small fans

were embedded in the cabinets (providing background noise of ~70 dB). An aluminum pan was

inserted beneath the grid floor to collect animal waste. A small camera was attached to the top of

the cabinet for video monitoring of behavior.

Measurements of freezing were performed using an automated and unbiased system

(Maren, 1998). Specifically, each behavioral testing chamber rested on a load-cell platform that

was sensitive to cage displacement due to the animal's movements. During behavioral testing,

load-cell activity values (ranging from -10 to +10 V) were collected and digitized at 5 Hz using

Threshold Activity Software (MED Associates, Inc.). Offline conversions of the load-cell activity

values were performed to generate absolute values ranging from 0 to 100; lower values indicate

minimal cage displacement, which coincided with freezing behaviors in the chambers.

Accordingly, freezing bouts were defined as absolute values of $\leq$10 for 1 s or more. The percentage

of freezing behavior during the pre-CS baseline, the CS, and the inter-trial intervals was computed

for each behavioral session. Shock reactivity was analyzed by directly reporting the absolute

values generated by the Threshold Activity Software (i.e., larger values indicated more movement

in the cage).

Unique contexts (A and B) were used for conditioning retrieval testing. Chamber

assignments were unique to each context and group assignments were counterbalanced across test

chambers when possible. For each experiment, context A was assigned to one of the behavioral

rooms, and B the other. For context A, the test chamber was wiped down with an acetic acid

solution (3%) and a small amount was poured in the pans beneath the grid floors. The cage lights

were turned on, while the chamber fans were turned off. The cabinet doors were closed. The

behavioral room was illuminated with dim red light. Animals were transported to and from the

chambers using white plastic transport boxes. For context B, an ammonium hydroxide solution

(1%) was used to wipe down and scent the chambers, black Plexiglas panels were placed over the

grid floors, the cage lights were turned off, the chamber fans were turned on, and the cabinet doors

remained open. The behavioral room was illuminated with white light (red room lights were turned

off). Rats were transported to and from the context using black plastic transport boxes that included

a layer of clean bedding.

Surgeries. For animals receiving intracranial microinfusions into the BNST, rats were transported

to the surgical suite and deeply anesthetized using isoflurane (5% for induction, 1-2% for

maintenance) (Acca et al., 2017; Goode et al., 2015; Nagaya et al., 2015; Zimmerman and Maren,

2011). Rats were then secured in a stereotaxic frame (Kopf Instruments). Hair on top of the

rodent's head was shaved, povidone-iodine was applied, and a small incision was made in the scalp

to expose the skull. Bregma and lambda were aligned on a horizontal plane, and small holes were

drilled in the skull for the placement of anchoring screws and bilateral stainless-steel guide

cannulas (26 gauge, 8 mm from the bottom of their plastic pedestals; Small Parts). The cannulas

were inserted into the BNST at the following coordinates: -0.15 mm posterior to bregma, ±2.65

mm lateral to the midline, and -5.85 mm dorsal to dura (guide cannulas were angled at 10° with

their needles directed at the midline). Dental cement was used to secure the cannulas to the screws

and stainless steel obturators (33 gauge, 9 mm; Small Parts) were inserted into the guide cannulas.

Animals were given at least one week to recover prior to the onset of behavioral training.

For rats injected with cholera toxin subunit B (CTb) conjugated with Alexa Fluor-488

(CTb-488; ThermoFisher Scientific) in the BNST, the surgical procedures were identical to those

described above. A single small hole was drilled into the skull to allow for the insertion of a glass

micropipette. Injection tips were backfilled with mineral oil and secured in the injector; CTb-488

was then drawn up into the injector immediately before use. Rats received unilateral CTb-488

infusions into either the left or right hemisphere (group assignments were randomized for sites of

CTb-488 infusion): -0.15 mm posterior to bregma, ±2.65 mm lateral to the midline, and -6.50 mm

dorsal to dura (the pipette was angled at 10° with the tip directed at the midline). CTb-488 (5.0

mg/μl; total volume of 0.25 μl) was microinfused into the brain using a Nanoject II auto-nanoliter

injector (Drummond Scientific Co.) secured to the arms of the stereotaxic frame. For the infusion

process, 50 nl (25 nl/s) of CTb-488 was infused once per min for 5 min to achieve 0.25 μl of the

total infusion (the injection needle was left in the brain for 5 additional minutes to allow for

diffusion of CTb-488). Following the infusion procedures, the incision was closed with sutures

and the animals were returned to their homecages where they recovered for ~10 days.

Intracranial infusions. Prior to behavioral testing, the animals were acclimated to the process of

intracranial infusions. On two occasions, the animals were transported from the vivarium to the

infusion room in 5-gallon buckets containing a layer of bedding; the experimenter(s) removed the

obturators and replaced them with clean ones. On the day of infusions, animals (in predetermined

squads of four to eight rats; representing all drug/behavioral groups as possible) were transported

to the laboratory, obturators were removed, and injectors were inserted into the guides. Stainless

steel injectors (33 gauge, Small Parts; 9-mm extending 1 mm beyond the end of the guide cannula)

were connected to polyethylene tubing (PE-20; Braintree Scientific); the other end of the tubing

was connected to gastight 10 μl syringes (Hamilton, Co.). The syringes were mounted to an

infusion pump (KD Scientific, Inc.). For the experiments shown in Figure 1 and Figure 1—Figure

Supplement 3, the α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptor

antagonist 2,3-dihydroxy-6-nitro-7-sulfamoyl-benzo[f]quinoxaline-2,3-dione (NBQX) was used

to reversibly inactivate the BNST (Adami et al., 2017; Davis and Walker, 2014; Goode et al., 2015;

Zimmerman and Maren, 2011). NBQX disodium salt hydrate (Sigma Life Sciences) was dissolved

in saline to a concentration of 10.0 μg/μl ("NBQX"); physiological saline served as the vehicle

("VEH" for all experiments). For the experiments shown in Figure 2 and 3, the γ-aminobutyric

acid (GABA)$_A$ receptor agonist muscimol was used to reversibly inactivate the BNST (Bangasser

et al., 2005; Breitfeld et al., 2015; Buffalari and See, 2011; Fendt et al., 2003; Goode et al., 2015;

Markham et al., 2009; Pina et al., 2015; Sajdyk et al., 2008; Xu et al., 2012). Muscimol (Sigma-

Aldrich) was dissolved in physiological saline to a concentration of 0.1 μg/μl ("MUS" for Figure

3 and 5); physiological saline served as the vehicle ("VEH"). For the representative image showing

drug spread in the BNST (Figure 1—Figure Supplement 1), muscimol TMR-X conjugate (Thermo

Fisher Scientific) was dissolved in physiological saline to a concentration of 0.1 μg/μl and used

for infusions. For all of the aforementioned experiments, drug or vehicle was drawn into the

injectors (immediately prior to the infusions) and a total volume of 0.275 μl/hemisphere of drug

or vehicle was infused at a rate of 0.275 μl/min; injectors were left in the cannulas for 1 min

following the infusions to allow for diffusion. Once the injectors were removed, clean obturators

were inserted into the guides.

Behavioral procedures and exclusions. Overviews of each behavioral experiment are provided

in the figures. The conditioned stimulus (CS) for all experiments was an auditory tone (80 dB, 2

kHz, 10 sec), which was paired with the unconditioned stimulus (US, footshock; 1.0 mA, 2 sec).

For the variable shock intensity experiment (Figure 4), US intensity varied as described.

FW/BW/NoCS(Test) BNST inactivation (twelve training trials). In a 3 × 2 design, animals

(*n* = 64, prior to exclusions) were randomly assigned to receive a CS at test [either a forward

("FW")- or backward ("BW")-trained CS] or no CS at test [animals were trained to a BW CS;

"No-CS"], and NBQX ("NBQX") or vehicle infusions ("VEH"). Of these rats, seven were

excluded due to off-target cannulas, three additional rats were excluded due to illness, and four

more were excluded due to a technical error during infusions that resulted in no drug or vehicle to

be infused (fourteen total exclusions). This resulted in the following (final) group numbers (shown

in data/figures): FW-VEH ($n = 5$); FW-NBQX ($n = 4$); BW-VEH ($n = 13$); BW-NBQX ($n = 12$);

No-CS-VEH ($n = 8$); No-CS-NBQX ($n = 8$). For conditioning, animals (in squads of eight rats)

were transported from the vivarium to context A; FW and BW procedures were run in alternating

squads of animals. Drug and vehicle assignments were counterbalanced for position in the

chambers. For FW conditioning, were placed in the context for 5 min before the delivery of twelve

CS-US trials (CS offset immediately preceded US onset; 70-sec intertrial interval). After the final

conditioning trial, the animals remained in the chamber for 1 min before being returned to their

homecages (the entire conditioning session consisted of 19 min total; for both FW and BW

conditioning). For BW conditioning, all aspects of conditioning were identical to the FW training

except the order of the CS and US was reversed. That is, after a 5-min baseline period, the animals

received 12 US-CS trials (the onset of the CS occurred immediately after the offset of the US; 70-

sec intertrial intervals). After conditioning, rats were returned to their homecages.

Twenty-four hours after conditioning, animals (in squads of four) were infused with NBQX

or VEH into the BNST immediately before being placed in context B. After 5 min of acclimation

to the context, FW and BW animals (intermixed in each squad) received twelve presentations of

the CS in the absence of the US (70-sec intertrial interval). The rats remained in the chambers for

1 min following the final CS (19 min session, in total). For rats not receiving the CS at test, the

animals remained in the test context for 19 min without presentation of the CS or US. We alternated

running squads that received the CS and those that did not after the infusions. Following the test,

animals were returned to their homecages.

FW/BW BNST inactivation (five training trials). In a 2 × 2 design, animals (*n* = 28, prior

to exclusions) were randomly assigned to receive forward ("FW") or backward ("BW")

conditioning using five training trials, and NBQX ("NBQX") or vehicle infusions ("VEH"). Of

these rats, three were excluded due to off-target cannulas. This resulted in the following (final)

group numbers (shown in data/figures): FW-NBQX (*n* = 6); FW-VEH (*n* = 6); BW-NBQX (*n* =

6); BW-VEH (*n* = 7). At the start of behavior, rats (in squads of seven) were transported from the

vivarium and placed in context A. We alternated squads that received forward or backward

conditioning. For forward conditioning, rats were given 3 min of acclimation to the context prior

to the onset of five CS-US trials. Rats remained in the chamber for 1 min following the final trial

(530 sec total conditioning session). For backward conditioning, rats were placed in context a for

a 3-min baseline (530 sec session total), after which they received five US-CS trials. Rats were

returned to their homecages after conditioning. Twenty-four hours after conditioning, the animals

were placed in context B in the absence of the CS or US for 530 sec to acclimate them to the test

context. After this acclimation session, rats were again returned to their homecages.

Forty-eight hours after conditioning, the animals received NBQX or VEH infusions

(identical to above) immediately prior to retrieval testing. FW and BW animals were intermixed

in each squad. Immediately after the infusions, animals were placed in context B for a retrieval

test to assess freezing to the CS. The test (530 sec in total) consisted of a 3-min baseline period

followed by five CS-alone presentations (70-s intertrial interval). Rats were returned to their

homecages following the test session.

RANDOM/BW/NoCS(Cond) BNST inactivation. In a 3 × 2 design, rats (*n* = 70, prior to

exclusions) were randomly assigned to receive backward conditioning ("BW"; identical to the 12-

trial BW conditioning described above), randomized CS/US trials in which the duration of the

intervals between CS offset and US onset vary on each trial ("RANDOM"), or US-only contextual

conditioning [shock intervals matched the RANDOM group; "NoCS(Cond)"]. Of these groups,

animals were randomly assigned to receive muscimol ("MUS") or vehicle ("VEH") infusions

immediately prior to retrieval. For this experiment, 30 animals were excluded due to off-target

cannulas. This resulted in the following (final) group numbers (shown in data/figures): BW-VEH

($n = 5$); BW-MUS ($n = 7$); RANDOM-VEH ($n = 6$); RANDOM-MUS ($n = 9$); NoCS(Cond)-VEH

($n = 7$); NoCS(Cond)-MUS ($n = 6$).

For conditioning, animals (in squads of five to eight rats; counterbalanced for the type of

conditioning) were transported to context A and underwent BW, RANDOM, or NoCS(Cond)

training. Drug and vehicle assignments were counterbalanced for the conditioning procedure and

chamber placements for each squad. For BW animals, the conditioning parameters matched those

used for the 12-trial experiment described above. For RANDOM animals, and after the 5-min

baseline, CS presentations matched the 60 s interstimulus intervals used for the 12-trial FW

paradigms (noted above); US onset occurred at the following intervals after offset of each CS at

each trial: 26 s, 52 s, 10 s, 40 s, 16 s, 52 s, 14 s, 50 s, 18 s, 36 s, 48 s, 6 s (the entire session last 19

min). For NoCS(Cond) animals, rats experienced the same protocol as the RANDOM group, but

no auditory CS occurred. Rats were returned to their homecages following conditioning.

Twenty-four hours later, animals (in squads of four to six) were transported to the

laboratory to receive pre-retrieval infusions of MUS or VEH (counterbalanced in each squad).

BW, RANDOM, and NoCS(Cond) animals were intermixed in each squad. Immediately after the

intracranial infusions, and after a 5-min baseline in context B, each squad of rats received twelve

presentations of the CS in the absence of the US. Each CS presentation was separated by 70 sec

intertrial intervals, with the entire test lasting 19 min. Rats were returned to their homecages

following the test session.

FIXED/VARIABLE BNST inactivation. In a 2 × 2 design, rats (n = 31, prior to exclusions)

were randomly assigned to receive forward conditioning with consistent ("FIXED") or variable

magnitudes of the US ("VARIABLE"); muscimol ("MUS") or vehicle ("VEH") infusions were

made prior to retrieval testing. For FIXED animals, US intensity was 1 mA at each trial. For

VARIABLE animals, shock intensity (in mA) varied across trials in this order (mean = 1 mA):

0.5, 1.8, 0.4, 1.6, 1.4, 0.3, 0.5, 1.8, 0.4, 1.6, 1.4, 0.3. In this experiment, one rat was excluded due

to off-target cannulas. This resulted in the following (final) group numbers (shown in data/figures):

FIXED-VEH (n = 7); FIXED-MUS (n = 8); VARIABLE-VEH (n = 7); VARIABLE-MUS (n =

8).

For conditioning, animals (in squads of seven to eight) were transported to context A

(squads alternated between FIXED and VARIABLE paradigms). The rats were given 5 min to

acclimate to the context before the onset of 12 CS-US pairings (70-sec intertrial intervals). Rats

remained in the chambers for 1 min following the final CS, with the entire conditioning session

lasting 19 min. Rats were returned to their homecages following conditioning.

Twenty-four hours later, rats (in squads of seven to eight) received intracranial infusions

of MUS or VEH into the BNST immediately before being placed in context B for retrieval testing.

Rats were given a 5-min baseline period before the onset of twelve CS-alone presentations (70-

sec intertrial intervals). Rats remained in the chambers for 1 min following the final CS, with the

entire test session lasting 19 min. Rats were returned to their homecages following the test.

FW/BW Fos-CTb. Animals (n = 60, before exclusions) were randomly assigned to receive

a forward ("FW")- or backward ("BW")-trained CS at testing, or no CS retrieval at test

["NoCS(Test)"]. Note that the No-CS group consists of animals that were trained to either a FW

or BW CS. Additionally, a group of BW-trained animals remained in their homecages ("NoTest")

during the final test and were sacrificed alongside the other groups. Twelve rats were excluded for

either excessive or off-target infusion of CTb-488 outside the borders of the BNST or an absence

of CTb-488 labeling in the ventral BNST. One additional rat was excluded due to a technical issue

that resulted in the loss of tissue at the level of the prefrontal cortex. This resulted in the following

(final) group numbers (shown in data/figures): FW ($n = 8$); BW ($n = 14$); No-CS [$n = 17$ (BW-

trained: $n = 9$; FW-trained: $n = 8$)]; NoTest ($n = 8$). For behavioral training, rats (in squads of six

to eight; with all groups intermixed) were transported to the laboratory and placed in context B to

acclimate for 5 min (no CS or US) and then returned to their homecages. Later that day, the animals

were transported to context A where they received twelve trials of either FW or BW fear

conditioning, which was identical to the other aforementioned procedures. We alternated FW and

BW conditioning squads. After conditioning, the rats were returned to their homecages. Twenty-

four hours after conditioning, all rats were exposed to context B for 20 min in the absence of the

CS or US to acclimate them to the test context.

Forty-eight hours later, FW, BW, and No-CS rats were transported to context B to receive

a retrieval test to the CS or were merely placed in the context (No-CS condition). Squads alternated

between CS and No-CS test. For rats undergoing CS retrieval, FW and BW animals (intermixed

in each squad) received CS trials as described previously. Rats were perfused 90 min following

the first CS of the test, in groups of three to four. For No-CS rats (with FW- and BW-trained

animals intermixed), animals were exposed to context B in the absence of the CS or US. No-CS

rats were perfused 95 min after being placed in the test context, in groups of three to four. NoTest

rats were perfused alongside groups of FW, BW, and No-CS rats. Rats were returned to their

homecages after testing and prior to the perfusions.

FW/BW intra-shock reactivity. Rats ($n = 32$, no exclusions) were randomly assigned to

receive forward ("FW") or backward ("BW") conditioning. No rats were excluded from this

experiment (no infusions occurred; $n = 16$, per group); only data from the conditioning session are

shown. Animals (in squads of eight) were transported to context A for either FW or BW

conditioning. Parameters for FW and BW conditioning were identical to the procedures for the

previous 12-trial experiments. We alternated FW and BW squads. Rats were returned to their

homecages following training.

Histological procedures. At the conclusion of behavioral testing, cannula-implanted animals were

overdosed with sodium pentobarbital (Fatal Plus; 100 mg/ml, 0.5 ml, i.p.). Transcardial perfusions

were then performed using chilled physiological saline followed by 10% formalin. Brains were

extracted and stored in 10% formalin for 24 hr at 4° C; brains were transferred to a 30% sucrose-

formalin solution for three or more days (at 4° C) before sectioning. Brains were flash frozen with

crushed dry ice and coronal sections (40 μm) containing the BNST were collected using a cryostat

(Leica Microsystems) at -20° C. The tissue was wet-mounted to gelatin-subbed microscope slides

and stained with 0.25% thionin prior to adding glass coverslips secured with mounting medium

(Permount, Sigma). To further examine the spread of drug, a subset of animals was infused

(identical to the aforementioned infusion parameters) with fluorescent muscimol (1.0 μg/μl;

EverFluor TMR-X conjugate, Setareh Biotech) before being sacrificed (these animals were not

perfused) and having brains dissected (40 μm; brains were stored in 30% sucrose solution at 4° C

until sectioning). These tissues were wet-mounted to slides and aqueous mounting medium

(Fluoromount; Sigma-Aldrich) was used to secure glass coverslips.

For CTb-injected animals, post-behavior perfusions mirrored the aforementioned

procedures. For sectioning, coronal sections (which included regions of the prefrontal cortex,

BNST, basolateral amygdala, and ventral hippocampus) were collected into well plates containing

phosphate-buffered saline (1× PBS) and stored in the dark at 4° C until immunohistochemistry

could be performed. For localization of the CTb-injection site, separate sections at the level of the

BNST were wet-mounted and coverslipped using Fluoromount mounting medium.

Immunohistochemistry. Immunohistochemistry for Fos was performed on free-floating brain

tissue similar to prior reports (Jin and Maren, 2015; Marek et al., 2018; Orsini et al., 2011; Wang

et al., 2016). For sections containing the BNST, Fos was stained using the following procedures

[all steps were performed at room temperature (~20° C) on a shaker, unless stated otherwise; rinses

were brief (~20 sec)]. The tissue was first rinsed in 1× tris-buffered saline (TBS; 7.4 pH), and then

incubated in 0.3% H_2O_2 (in TBS) for 15 min, followed by rinses (×3) in TBS. Slices were

transferred to primary antibody [rabbit anti-c-fos, 1:10,000 in 1× TBS containing Tween 20

(TBST); Millipore] and incubated overnight. Sections were then rinsed (×3) in TBS before

incubating in secondary antibody for 1 hr (biotinylated goat anti-rabbit, 1:1,000 in TBST; Jackson

Immunoresearch). Sections were rinsed (×3) again in TBS. The slices were transferred to wells

containing avidin biotin complex (ABC, 1:1,000 in TBST; Vector Labs) for 45 min. The tissues

were again rinsed (×3) in TBS. Tissue was transferred to wells containing 3,3′ diaminobenzidine

[(DAB) 5% stock, 1:200], nickel ammonium sulfate (5% stock, 1:10), and 30% H_2O_2 (1:2,000) in

TBS for 10 min to generate purple/black nuclear chromophore products. After another rinse (×3)

in TBS, the tissue was subsequently wet-mounted to slides and secured with coverslips using

Permount mounting medium.

For sections containing the prefrontal cortex, amygdala, and hippocampus (Fos-CTb

experiment), Fos was stained using the following procedures [unless stated otherwise, each step

occurred at room temperature (~20° C) on a shaker (and away from excess light)]. First, the tissue

was rinsed (10 min; ×2) in 1× TBS, followed by a 10 min wash in 1× TBST. The tissue was

incubated in 10% normal donkey serum (NDS; in TBST) for 1 hr. The slices were then rinsed (5

min; ×2) in TBST. Sections were transferred to primary antibody [goat anti-c-fos, 1:2,000 in 3%

NDS (in TBST); Santa Cruz Biotechnology] and incubated on a rotator in the dark for 72 hr at 4°

C. The tissue was rinsed (10 min; ×3) in TBST before incubating in secondary antibody

[biotinylated donkey anti-goat, 1:200 in 3% NDS (in TBST); Santa Cruz Biotechnology] for 2 hr.

Slices were then rinsed (10 min; ×3) in TBST. Sections were transferred to wells containing

streptavidin (Alexa-Fluor 594-conjugate, 1:500 in 3% NDS (in TBST); Thermo Fisher Scientific]

for 1 hr. Tissue was rinsed (10 min; ×3) in TBS before being wet-mounted to slides and secured

with coverslips using Fluoromount mounting medium.

Image analyses. All imaging and counting procedures (for all regions) were performed with the

experimenter(s) blind to the group assignments of the animals. For thionin-stained coronal tissue,

photomicrographs of cannula in the BNST were generated (10× magnification) using a Leica

microscope (MZFLIII) and Leica Firecam software. For animals infused with fluorescent

muscimol into the BNST, infusion sites were imaged (10× magnification) using a Zeiss

microscope and Axio Imager 2 software (Zen Pro 2012). For CTb-injected animals, BNST images

were generated (10× magnification) using the same Zeiss microscope and software.

For Fos analyses in BNST, brightfield images of BNST [in BNST regions ranging from

approximately -0.00 to -0.50 mm posterior to bregma of the skull, and from both left and right

hemispheres (randomized for site of CTb-injection)] were generated (10× magnification) using a

Zeiss microscope and Axio Imager 2 software (Zen Pro 2012). ImageJ software (Schneider et al.,

2012) was used to count cells. Counts were confined to the following areas of interest: (1)

"ovBNST" [an area of 0.217 mm × 0.558 mm (oval); targeting the oval nucleus of the BNST], (2)

"dBNST" [0.372 mm^2 (circle); targeting the dorsal and medial subregions of the anterior BNST,

including the anteromedial area of the BNST (dorsal to the anterior commissure)], and (3)

"vBNST" [0.434 mm^2 (circle); targeting the ventral regions (ventral to the anterior commissure)

of the anterior BNST, which includes the ventral portion of the anteromedial area, the anterolateral

area, and the fusiform nucleus of the BNST] (Swanson, 2003). For each of these regions, three to

six images were quantified and averaged for each animal (Fos levels were standardized to 0.1

mm^2).

For Fos-CTb analyses in the prefrontal cortex, amygdala, and hippocampus, images (10×

magnification) of Fos and CTb expression were generated using a Zeiss microscope and Axio

Imager 2 software (Zen Pro 2012). Cell counts were analyzed using ImageJ software. Images were

generated and analyzed only for the hemisphere injected with CTb. Counts were confined to the

following areas of interest: (1) "IL" [an area of 0.805 mm × 0.217 mm (rectangle); targeting deep

and superficial layers of the infralimbic cortex (approximately +2.0 to +3.2 mm anterior to bregma

of the skull)], (2) "PL" [1.115 mm × 0.217 mm (rectangle); targeting deep and superficial layers

of the prelimbic cortex (approximately +2.0 to +3.2 mm anterior to bregma of the skull)], (3)

"BLA" [0.434 mm^2 (circle); targeting the basolateral amygdala (approximately -1.8 to -3.5 mm

posterior to bregma of the skull)], and (4) "HPC" [0.496 mm × 0.217 mm (rectangle); targeting

the ventral subiculum but may include some CA1 cells of the ventral hippocampus (approximately

within -4.7 to -7.0 mm posterior to bregma of the skull)] (Swanson, 2003). For each of these

regions, three to six images were quantified and averaged for each animal (all counts and

percentages were normalized to 0.1 mm^2).

Statistics. All data were submitted to repeated or factorial analysis of variance (ANOVA) or two-

tailed *t*-tests as described for each experiment. Fisher's protected least significant difference

(PLSD) test was used for *post hoc* comparisons of group means following a significant omnibus *F*

ratio in the ANOVA (α was set at 0.05). No statistical methods were used to predetermine group

sizes (group sizes were selected based on prior work and what is common for the field). Data

distributions were assumed to be normal, but these were not formally tested. Unless stated

otherwise, all data are represented as means ± s.e.m.

ACKNOWLEDGEMENTS

The authors thank Carolyn Evemy, Kaitlyn French, and Sohmee Kim for technical assistance.

Supported by grants from the National Institutes of Health (R01MH065961 and R01MH117852

to S.M. and F31MH107113 to T.D.G.), as well as a McKnight Foundation Memory and Cognitive

Disorders Award and a Brain & Behavior Research Foundation NARSAD Distinguished

Investigator Grant to S.M.

ADDITIONAL INFORMATION

Competing interests

The authors declare no competing interests.

Author contributions

T.D.G. and S.M. designed the research; T.D.G., R.R., G.M.A., and O.W.M. performed the

research; T.D.G., R.R., G.M.A., O.W.M., and S.M. analyzed data; T.D.G. and S.M. wrote the

paper.

Figure 1—Figure Supplement 1
Representative bilateral cannula placements in the BNST. Photomicrograph (10×) of a thionin-stained coronal section depicting representative cannula tracts in the BNST (top panel). Fluorescent image (gold filter) of a coronal section (10×) showing spread of drug in BNST (BNST outlined in dotted line) (bottom panel).

Figure 1—Figure Supplement 2
Bilateral cannula placements for BNST microinfusions. Schematic depicting cannula placements for Figure 1. Symbols (split by each group) correspond to injector tips (approximate borders of BNST are shown by red dotted outline; approximate borders of BNST are shown by red dotted outline).

Figure 1—Figure Supplement 3

Effects of BNST inactivation on freezing to a forward vs. backward CS trained with five trials. **(A)** Behavioral schematic. **(B)** Freezing behavior during conditioning and retrieval testing. For conditioning, the left panel depicts mean percentage freezing during the 3-min baseline (BL) and across each conditioning block (each 138-sec block is comprised of two trials; conditioning trials consist of freezing during the 10-sec CS followed by the 58-sec interstimulus interval). For retrieval testing, the center panel shows mean percentage freezing at the 3-min baseline (BL) and across each test trial (each trial consists of freezing during the 10-sec CS followed by the 60-sec interstimulus interval). The right panel shows mean percentage freezing after baseline (trials 1-5; corresponding to 350 sec of behavior). All data are represented as means ± s.e.m [FW-VEH ($n = 6$); FW-NBQX ($n = 6$); BW-VEH ($n = 7$)]; BW-NBQX ($n = 6$)]; * = $p < 0.05$.

Figure 1—Figure Supplement 4

Bilateral cannula placements for BNST microinfusions. Schematic depicting cannula placements for Figure 1—Figure Supplement 3. Symbols (split by each group) correspond to injector tips (approximate borders of BNST are shown by red dotted outline).

Figure 1—Figure Supplement 5. *Shock-induced activity during conditioning to a forward vs. backward CS.* **(A)** Mean percentage of freezing at baseline (BL; 5-min) and across conditioning blocks (each 138-sec block is comprised of two trials; trials consist of freezing during the 10-sec CS followed by the 58-sec interstimulus interval). **(B)** Mean activity values across the 5-min BL (no shock present). **(C)** The left panel shows mean shock-induced activity (during shocks) at conditioning [averaged into 4-sec (two-shock) blocks]. The right panel shows mean shock-induced activity across all trials. All data are represented as means ± s.e.m [FW (*n* = 16); BW (*n* = 16)]; * = *p* < 0.05.

Figure 2—Figure Supplement 1
Bilateral cannula placements for BNST microinfusions. Schematic depicting cannula placements for Figure 2. Symbols (split by each group) correspond to injector tips (approximate borders of BNST are shown by red dotted outline).

Figure 3—Figure Supplement 1
Bilateral cannula placements for BNST microinfusions. Schematic depicting cannula placements for Figure 3. Symbols (split by each group) correspond to injector tips (approximate borders of BNST are shown by red dotted outline).

Figure 6—Figure Supplement 1
CTb injection sites in BNST. Approximation of the most ventral and centermost sites of unilateral microinjection of CTb (green X's) for all animals shown in Figure 6 and 7 (red outline approximates borders of BNST).

References

Acca, G.M., Mathew, A.S., Jin, J., Maren, S., Nagaya, N., 2017. Allopregnanolone induces state-dependent fear via the bed nucleus of the stria terminalis. Horm. Behav. 89, 137–144. doi:10.1016/j.yhbeh.2017.01.002

Adami, M.B., Barretto-de-Souza, L., Duarte, J.O., Almeida, J., Crestani, C.C., 2017. Both N-methyl-D-aspartate and non-N-methyl-D-aspartate glutamate receptors in the bed nucleus of the stria terminalis modulate the cardiovascular responses to acute restraint stress in rats. J. Psychopharmacol. (Oxford) 31, 674–681. doi:10.1177/0269881117691468

Alvarez, R.P., Chen, G., Bodurka, J., Kaplan, R., Grillon, C., 2011. Phasic and sustained fear in humans elicits distinct patterns of brain activity. Neuroimage 55, 389–400. doi:10.1016/j.neuroimage.2010.11.057

Alvarez, R.P., Kirlic, N., Misaki, M., Bodurka, J., Rhudy, J.L., Paulus, M.P., Drevets, W.C., 2015. Increased anterior insula activity in anxious individuals is linked to diminished perceived control. Transl. Psychiatry 5, e591. doi:10.1038/tp.2015.84

Amadi, U., Lim, S.H., Liu, E., Baratta, M.V., Goosens, K.A., 2017. Hippocampal processing of ambiguity enhances fear memory. Psychol. Sci. 28, 143–161. doi:10.1177/0956797616674055

Andreatta, M., Fendt, M., Mühlberger, A., Wieser, M.J., Imobersteg, S., Yarali, A., Gerber, B., Pauli, P., 2012. Onset and offset of aversive events establish distinct memories requiring fear and reward networks. Learn. Mem. 19, 518–526. doi:10.1101/lm.026864.112

Andreescu, C., Sheu, L.K., Tudorascu, D., Gross, J.J., Walker, S., Banihashemi, L., Aizenstein, H., 2015. Emotion reactivity and regulation in late-life generalized anxiety disorder: functional connectivity at baseline and post-treatment. Am. J. Geriatr. Psychiatry 23, 200–214. doi:10.1016/j.jagp.2014.05.003

Asok, A., Draper, A., Hoffman, A.F., Schulkin, J., Lupica, C.R., Rosen, J.B., 2018. Optogenetic silencing of a corticotropin-releasing factor pathway from the central amygdala to the bed nucleus of the stria terminalis disrupts sustained fear. Mol. Psychiatry 23, 914–922. doi:10.1038/mp.2017.79

Aston-Jones, G., Harris, G.C., 2004. Brain substrates for increased drug seeking during protracted withdrawal. Neuropharmacology 47 Suppl 1, 167–179. doi:10.1016/j.neuropharm.2004.06.020

Avery, S.N., Clauss, J.A., Blackford, J.U., 2016. The human BNST: functional role in anxiety and addiction. Neuropsychopharmacology 41, 126–141. doi:10.1038/npp.2015.185

Avery, S.N., Clauss, J.A., Winder, D.G., Woodward, N., Heckers, S., Blackford, J.U., 2014. BNST neurocircuitry in humans. Neuroimage 91, 311–323. doi:10.1016/j.neuroimage.2014.01.017

Ayres, J.J.., Mahoney, W.J., Proulx, D.T., Benedict, J.O., 1976. Backward conditioning as an extinction procedure. Learn Motiv 7, 368–381. doi:10.1016/0023-9690(76)90043-6

Ayres, J.J.B., Haddad, C., Albert, M., 1987. One-trial excitatory backward conditioning as assessed by conditioned suppression of licking in rats: Concurrent observations of lick suppression and defensive behaviors. Anim Learn Behav 15, 212–217. doi:10.3758/BF03204964

Bali, A., Jaggi, A.S., 2015. Electric foot shock stress: a useful tool in neuropsychiatric studies. Rev Neurosci 26, 655–677. doi:10.1515/revneuro-2015-0015

Bangasser, D.A., Santollo, J., Shors, T.J., 2005. The bed nucleus of the stria terminalis is critically involved in enhancing associative learning after stressful experience. Behav. Neurosci. 119, 1459–1466. doi:10.1037/0735-7044.119.6.1459

Barnet, R.C., Miller, R.R., 1996. Second-order excitation mediated by a backward conditioned inhibitor. Journal of Experimental Psychology: Animal Behavior Processes 22, 279–296. doi:10.1037/0097-7403.22.3.279

Behar, E., DiMarco, I.D., Hekler, E.B., Mohlman, J., Staples, A.M., 2009. Current theoretical models of generalized anxiety disorder (GAD): conceptual review and treatment implications. J. Anxiety Disord. 23, 1011–1023. doi:10.1016/j.janxdis.2009.07.006

Bennett, K.P., Dickmann, J.S., Larson, C.L., 2018. If or when? Uncertainty's role in anxious anticipation. Psychophysiology 55, e13066. doi:10.1111/psyp.13066

Berridge, K.C., 2018. Evolving concepts of emotion and motivation. Front. Psychol. 9, 1647. doi:10.3389/fpsyg.2018.01647

Besnard, A., Gao, Y., TaeWoo Kim, M., Twarkowski, H., Reed, A.K., Langberg, T., Feng, W., Xu, X., Saur, D., Zweifel, L.S., Davison, I., Sahay, A., 2019. Dorsolateral septum somatostatin interneurons gate mobility to calibrate context-specific behavioral fear responses. Nat. Neurosci. doi:10.1038/s41593-018-0330-y

Bevins, R.A., Ayres, J.J., 1992. One-trial backward excitatory fear conditioning transfers across contexts. Behav. Res. Ther. 30, 551–554. doi:10.1016/0005-7967(92)90041-E

Blanco, C., Xu, Y., Schneier, F.R., Okuda, M., Liu, S.-M., Heimberg, R.G., 2011. Predictors of persistence of social anxiety disorder: a national study. J. Psychiatr. Res. 45, 1557–1563. doi:10.1016/j.jpsychires.2011.08.004

Breitfeld, T., Bruning, J.E.A., Inagaki, H., Takeuchi, Y., Kiyokawa, Y., Fendt, M., 2015. Temporary inactivation of the anterior part of the bed nucleus of the stria terminalis blocks alarm pheromone-induced defensive behavior in rats. Front. Neurosci. 9, 321. doi:10.3389/fnins.2015.00321

Brinkmann, L., Buff, C., Feldker, K., Neumeister, P., Heitmann, C.Y., Hofmann, D., Bruchmann, M., Herrmann, M.J., Straube, T., 2018. Inter-individual differences in trait anxiety shape the functional connectivity between the bed nucleus of the stria terminalis and the amygdala during brief threat processing. Neuroimage 166, 110–116. doi:10.1016/j.neuroimage.2017.10.054

Brinkmann, L, Buff, C., Feldker, K., Tupak, S.V., Becker, M.P.I., Herrmann, M.J., Straube, T., 2017. Distinct phasic and sustained brain responses and connectivity of amygdala and bed nucleus of the stria terminalis during threat anticipation in panic disorder. Psychol. Med. 47, 2675–2688. doi:10.1017/S0033291717001192

Brinkmann, Leonie, Buff, C., Neumeister, P., Tupak, S.V., Becker, M.P.I., Herrmann, M.J., Straube, T., 2017. Dissociation between amygdala and bed nucleus of the stria terminalis during threat anticipation in female post-traumatic stress disorder patients. Hum. Brain Mapp. 38, 2190–2205. doi:10.1002/hbm.23513

Buff, C., Brinkmann, L., Bruchmann, M., Becker, M.P.I., Tupak, S., Herrmann, M.J., Straube, T., 2017. Activity alterations in the bed nucleus of the stria terminalis and amygdala during threat anticipation in generalized anxiety disorder. Soc. Cogn. Affect. Neurosci. 12, 1766–1774. doi:10.1093/scan/nsx103

Buffalari, D.M., See, R.E., 2011. Inactivation of the bed nucleus of the stria terminalis in an animal model of relapse: effects on conditioned cue-induced reinstatement and its enhancement by yohimbine. Psychopharmacology 213, 19–27. doi:10.1007/s00213-010-2008-3

Burman, M.A., Simmons, C.A., Hughes, M., Lei, L., 2014. Developing and validating trace fear conditioning protocols in C57BL/6 mice. J. Neurosci. Methods 222, 111–117. doi:10.1016/j.jneumeth.2013.11.005

Campeau, S., Falls, W.A., Cullinan, W.E., Helmreich, D.L., Davis, M., Watson, S.J., 1997.

Elicitation and reduction of fear: behavioural and neuroendocrine indices and brain induction of the immediate-early gene c-fos. Neuroscience 78, 1087–1104. doi:10.1016/S0306-4522(96)00632-X

Canteras, N.S., Swanson, L.W., 1992. Projections of the ventral subiculum to the amygdala, septum, and hypothalamus: a PHAL anterograde tract-tracing study in the rat. J. Comp. Neurol. 324, 180–194. doi:10.1002/cne.903240204

Chang, R.C., Blaisdell, A.P., Miller, R.R., 2003. Backward conditioning: mediation by the context. J Exp Psychol Anim Behav Process 29, 171–183. doi:10.1037/0097-7403.29.3.171

Ch'ng, S., Fu, J., Brown, R.M., McDougall, S.J., Lawrence, A.J., 2018. The intersection of stress and reward: BNST modulation of aversive and appetitive states. Prog. Neuropsychopharmacol. Biol. Psychiatry 87, 108–125. doi:10.1016/j.pnpbp.2018.01.005

Choi, J.M., Padmala, S., Pessoa, L., 2012. Impact of state anxiety on the interaction between threat monitoring and cognition. Neuroimage 59, 1912–1923. doi:10.1016/j.neuroimage.2011.08.102

Choi, J.M., Padmala, S., Spechler, P., Pessoa, L., 2014. Pervasive competition between threat and reward in the brain. Soc. Cogn. Affect. Neurosci. 9, 737–750. doi:10.1093/scan/nst053

Christianson, J.P., Jennings, J.H., Ragole, T., Flyer, J.G.N., Benison, A.M., Barth, D.S., Watkins, L.R., Maier, S.F., 2011. Safety signals mitigate the consequences of uncontrollable stress via a circuit involving the sensory insular cortex and bed nucleus of the stria terminalis. Biol. Psychiatry 70, 458–464. doi:10.1016/j.biopsych.2011.04.004

Coaster, M., Rogers, B.P., Jones, O.D., Viscusi, W.K., Merkle, K.L., Zald, D.H., Gore, J.C., 2011. Variables influencing the neural correlates of perceived risk of physical harm. Cogn Affect Behav Neurosci 11, 494–507. doi:10.3758/s13415-011-0047-9

Colvonen, P.J., Glassman, L.H., Crocker, L.D., Buttner, M.M., Orff, H., Schiehser, D.M., Norman, S.B., Afari, N., 2017. Pretreatment biomarkers predicting PTSD psychotherapy outcomes: A systematic review. Neurosci. Biobehav. Rev. 75, 140–156. doi:10.1016/j.neubiorev.2017.01.027

Comer, J.S., Blanco, C., Hasin, D.S., Liu, S.-M., Grant, B.F., Turner, J.B., Olfson, M., 2011. Health-related quality of life across the anxiety disorders: results from the national epidemiologic survey on alcohol and related conditions (NESARC). J. Clin. Psychiatry 72, 43–50. doi:10.4088/JCP.09m05094blu

Connor, D.A., Kutlu, M.G., Gould, T.J., 2017. Nicotine disrupts safety learning by enhancing fear associated with a safety cue via the dorsal hippocampus. J. Psychopharmacol. (Oxford) 31, 934–944. doi:10.1177/0269881117695861

Corcoran, K.A., Quirk, G.J., 2007. Activity in prelimbic cortex is necessary for the expression of learned, but not innate, fears. J. Neurosci. 27, 840–844. doi:10.1523/JNEUROSCI.5327-06.2007

Costello, E.J., He, J., Sampson, N.A., Kessler, R.C., Merikangas, K.R., 2014. Services for adolescents with psychiatric disorders: 12-month data from the National Comorbidity Survey-Adolescent. Psychiatr. Serv. 65, 359–366. doi:10.1176/appi.ps.201100518

Crestani, C.C., Alves, F.H., Gomes, F.V., Resstel, L.B., Correa, F.M., Herman, J.P., 2013. Mechanisms in the bed nucleus of the stria terminalis involved in control of autonomic and neuroendocrine functions: a review. Curr Neuropharmacol 11, 141–159. doi:10.2174/1570159X11311020002

Crowley, N.A., Bloodgood, D.W., Hardaway, J.A., Kendra, A.M., McCall, J.G., Al-Hasani, R., McCall, N.M., Yu, W., Schools, Z.L., Krashes, M.J., Lowell, B.B., Whistler, J.L.,

Bruchas, M.R., Kash, T.L., 2016. Dynorphin controls the gain of an amygdalar anxiety circuit. Cell Rep. 14, 2774–2783. doi:10.1016/j.celrep.2016.02.069

Daldrup, T., Lesting, J., Meuth, P., Seidenbecher, T., Pape, H.-C., 2016. Neuronal correlates of sustained fear in the anterolateral part of the bed nucleus of stria terminalis. Neurobiol. Learn. Mem. 131, 137–146. doi:10.1016/j.nlm.2016.03.020

Daldrup, T., Remmes, J., Lesting, J., Gaburro, S., Fendt, M., Meuth, P., Kloke, V., Pape, H.C., Seidenbecher, T., 2015. Expression of freezing and fear-potentiated startle during sustained fear in mice. Genes Brain Behav. 14, 281–291. doi:10.1111/gbb.12211

Daniel, S.E., Rainnie, D.G., 2016. Stress modulation of opposing circuits in the bed nucleus of the stria terminalis. Neuropsychopharmacology 41, 103–125. doi:10.1038/npp.2015.178

Davies, C.D., Craske, M.G., 2015. Psychophysiological responses to unpredictable threat: effects of cue and temporal unpredictability. Emotion 15, 195–200. doi:10.1037/emo0000038

Davis, M., 2006. Neural systems involved in fear and anxiety measured with fear-potentiated startle. Am. Psychol. 61, 741–756. doi:10.1037/0003-066X.61.8.741

Davis, M., Walker, D.L., 2014. Role of bed nucleus of the stria terminalis and amygdala AMPA receptors in the development and expression of context conditioning and sensitization of startle by prior shock. Brain Struct. Funct. 219, 1969–1982. doi:10.1007/s00429-013-0616-5

Davis, M., Walker, D.L., Miles, L., Grillon, C., 2010. Phasic vs sustained fear in rats and humans: role of the extended amygdala in fear vs anxiety. Neuropsychopharmacology 35, 105–135. doi:10.1038/npp.2009.109

deCampo, D.M., Fudge, J.L., 2013. Amygdala projections to the lateral bed nucleus of the stria terminalis in the macaque: comparison with ventral striatal afferents. J. Comp. Neurol. 521, 3191–3216. doi:10.1002/cne.23340

Deslauriers, J., Toth, M., Der-Avakian, A., Risbrough, V.B., 2017. Current status of animal models of posttraumatic stress disorder: behavioral and biological phenotypes, and future challenges in improving translation. Biol. Psychiatry. doi:10.1016/j.biopsych.2017.11.019

Dong, H.W., Petrovich, G.D., Swanson, L.W., 2001. Topography of projections from amygdala to bed nuclei of the stria terminalis. Brain Res Brain Res Rev 38, 192–246. doi:10.1016/S0165-0173(01)00079-0

Fanselow, M.S., 1994. Neural organization of the defensive behavior system responsible for fear. Psychon. Bull. Rev. 1, 429–438. doi:10.3758/BF03210947

Fanselow, M.S., Pennington, Z.T., 2017. The Danger of LeDoux and Pine's Two-System Framework for Fear. Am. J. Psychiatry 174, 1120–1121. doi:10.1176/appi.ajp.2017.17070818

Fanselow, M.S., Pennington, Z.T., 2018. A return to the psychiatric dark ages with a two-system framework for fear. Behav. Res. Ther. 100, 24–29. doi:10.1016/j.brat.2017.10.012

Fendt, M., Endres, T., Apfelbach, R., 2003. Temporary inactivation of the bed nucleus of the stria terminalis but not of the amygdala blocks freezing induced by trimethylthiazoline, a component of fox feces. J. Neurosci. 23, 23–28.

Forray, M.I., Gysling, K., 2004. Role of noradrenergic projections to the bed nucleus of the stria terminalis in the regulation of the hypothalamic-pituitary-adrenal axis. Brain Res Brain Res Rev 47, 145–160. doi:10.1016/j.brainresrev.2004.07.011

Fox, A.S., Oler, J.A., Birn, R.M., Shackman, A.J., Alexander, A.L., Kalin, N.H., 2018. Functional Connectivity within the Primate Extended Amygdala Is Heritable and Associated with Early-Life Anxious Temperament. J. Neurosci. 38, 7611–7621. doi:10.1523/JNEUROSCI.0102-18.2018

Fox, A.S., Shackman, A.J., 2017. The central extended amygdala in fear and anxiety: Closing the gap between mechanistic and neuroimaging research. Neurosci. Lett. doi:10.1016/j.neulet.2017.11.056

Fox, A.S., Shelton, S.E., Oakes, T.R., Converse, A.K., Davidson, R.J., Kalin, N.H., 2010. Orbitofrontal cortex lesions alter anxiety-related activity in the primate bed nucleus of stria terminalis. J. Neurosci. 30, 7023–7027. doi:10.1523/JNEUROSCI.5952-09.2010

Gerber, B., Yarali, A., Diegelmann, S., Wotjak, C.T., Pauli, P., Fendt, M., 2014. Pain-relief learning in flies, rats, and man: basic research and applied perspectives. Learn. Mem. 21, 232–252. doi:10.1101/lm.032995.113

Gewirtz, J.C., McNish, K.A., Davis, M., 1998. Lesions of the bed nucleus of the stria terminalis block sensitization of the acoustic startle reflex produced by repeated stress, but not fear-potentiated startle. Prog. Neuropsychopharmacol. Biol. Psychiatry 22, 625–648.

Glangetas, C., Girard, D., Groc, L., Marsicano, G., Chaouloff, F., Georges, F., 2013. Stress switches cannabinoid type-1 (CB1) receptor-dependent plasticity from LTD to LTP in the bed nucleus of the stria terminalis. J. Neurosci. 33, 19657–19663. doi:10.1523/JNEUROSCI.3175-13.2013

Glangetas, C., Massi, L., Fois, G.R., Jalabert, M., Girard, D., Diana, M., Yonehara, K., Roska, B., Xu, C., Lüthi, A., Caille, S., Georges, F., 2017. NMDA-receptor-dependent plasticity in the bed nucleus of the stria terminalis triggers long-term anxiolysis. Nat. Commun. 8, 14456. doi:10.1038/ncomms14456

Goode, T.D., Jin, J., Maren, S., 2018. Neural circuits for fear relapse, in: Neurobiology of Abnormal Emotion and Motivated Behaviors. Academic Press.

Goode, T.D., Kim, J.J., Maren, S., 2015. Reversible inactivation of the bed nucleus of the stria terminalis prevents reinstatement but not renewal of extinguished fear. Eneuro 2. doi:10.1523/ENEURO.0037-15.2015

Goode, T.D., Maren, S., 2017. Role of the bed nucleus of the stria terminalis in aversive learning and memory. Learn. Mem. 24, 480–491. doi:10.1101/lm.044206.116

Goode, T.D., Maren, S., 2018. Common neurocircuitry mediating drug and fear relapse in preclinical models. Psychopharmacology. doi:10.1007/s00213-018-5024-3

Graham, B.M., Callaghan, B.L., Richardson, R., 2014. Bridging the gap: Lessons we have learnt from the merging of psychology and psychiatry for the optimisation of treatments for emotional disorders. Behav. Res. Ther. 62, 3–16. doi:10.1016/j.brat.2014.07.012

Grupe, D.W., Oathes, D.J., Nitschke, J.B., 2013. Dissecting the anticipation of aversion reveals dissociable neural networks. Cereb. Cortex 23, 1874–1883. doi:10.1093/cercor/bhs175

Gungor, N.Z., Paré, D., 2016. Functional heterogeneity in the bed nucleus of the stria terminalis. J. Neurosci. 36, 8038–8049. doi:10.1523/JNEUROSCI.0856-16.2016

Hammack, S.E., Guo, J.-D., Hazra, R., Dabrowska, J., Myers, K.M., Rainnie, D.G., 2009. The response of neurons in the bed nucleus of the stria terminalis to serotonin: implications for anxiety. Prog. Neuropsychopharmacol. Biol. Psychiatry 33, 1309–1320. doi:10.1016/j.pnpbp.2009.05.013

Hammack, S.E., Todd, T.P., Kocho-Schellenberg, M., Bouton, M.E., 2015. Role of the bed nucleus of the stria terminalis in the acquisition of contextual fear at long or short context-shock intervals. Behav. Neurosci. 129, 673–678. doi:10.1037/bne0000088

Harris, N.A., Winder, D.G., 2018. Synaptic Plasticity in the Bed Nucleus of the Stria Terminalis: Underlying Mechanisms and Potential Ramifications for Reinstatement of Drug- and Alcohol-Seeking Behaviors. ACS Chem. Neurosci. 9, 2173–2187. doi:10.1021/acschemneuro.8b00169

Haufler, D., Nagy, F.Z., Pare, D., 2013. Neuronal correlates of fear conditioning in the bed

nucleus of the stria terminalis. Learn. Mem. 20, 633–641. doi:10.1101/lm.031799.113

Herrmann, M.J., Boehme, S., Becker, M.P.I., Tupak, S.V., Guhn, A., Schmidt, B., Brinkmann, L., Straube, T., 2016. Phasic and sustained brain responses in the amygdala and the bed nucleus of the stria terminalis during threat anticipation. Hum. Brain Mapp. 37, 1091–1102. doi:10.1002/hbm.23088

Herry, C., Ciocchi, S., Senn, V., Demmou, L., Müller, C., Lüthi, A., 2008. Switching on and off fear by distinct neuronal circuits. Nature 454, 600–606. doi:10.1038/nature07166

Heth, C.D., 1976. Simultaneous and backward fear conditioning as a function of number of CS-UCS pairings. Journal of Experimental Psychology: Animal Behavior Processes 2, 117–129. doi:10.1037/0097-7403.2.2.117

Iza, M., Olfson, M., Vermes, D., Hoffer, M., Wang, S., Blanco, C., 2013. Probability and predictors of first treatment contact for anxiety disorders in the United States: analysis of data from the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC). J. Clin. Psychiatry 74, 1093–1100. doi:10.4088/JCP.13m08361

Jennings, J.H., Sparta, D.R., Stamatakis, A.M., Ung, R.L., Pleil, K.E., Kash, T.L., Stuber, G.D., 2013. Distinct extended amygdala circuits for divergent motivational states. Nature 496, 224–228. doi:10.1038/nature12041

Jin, J., Maren, S., 2015. Fear renewal preferentially activates ventral hippocampal neurons projecting to both amygdala and prefrontal cortex in rats. Sci. Rep. 5, 8388. doi:10.1038/srep08388

Johnson, S.B., Emmons, E.B., Lingg, R.T., Anderson, R.M., Romig-Martin, S.A., LaLumiere, R.T., Narayanan, N.S., Viau, V., Radley, J.J., 2018. Prefrontal-bed nucleus circuit modulation of a passive coping response set. J. Neurosci. doi:10.1523/JNEUROSCI.1421-18.2018

Kalin, N.H., Shelton, S.E., Fox, A.S., Oakes, T.R., Davidson, R.J., 2005. Brain regions associated with the expression and contextual regulation of anxiety in primates. Biol. Psychiatry 58, 796–804. doi:10.1016/j.biopsych.2005.05.021

Kalin, N.H., Shelton, S.E., Fox, A.S., Rogers, J., Oakes, T.R., Davidson, R.J., 2008. The serotonin transporter genotype is associated with intermediate brain phenotypes that depend on the context of eliciting stressor. Mol. Psychiatry 13, 1021–1027. doi:10.1038/mp.2008.37

Kim, S.-Y., Adhikari, A., Lee, S.Y., Marshel, J.H., Kim, C.K., Mallory, C.S., Lo, M., Pak, S., Mattis, J., Lim, B.K., Malenka, R.C., Warden, M.R., Neve, R., Tye, K.M., Deisseroth, K., 2013. Diverging neural pathways assemble a behavioural state from separable features in anxiety. Nature 496, 219–223. doi:10.1038/nature12018

Kinley, D.J., Walker, J.R., Enns, M.W., Sareen, J., 2011. Panic attacks as a risk for later psychopathology: results from a nationally representative survey. Depress. Anxiety 28, 412–419. doi:10.1002/da.20809

Kinnison, J., Padmala, S., Choi, J.-M., Pessoa, L., 2012. Network analysis reveals increased integration during emotional and motivational processing. J. Neurosci. 32, 8361–8372. doi:10.1523/JNEUROSCI.0821-12.2012

Kiyokawa, Y., Mikami, K., Mikamura, Y., Ishii, A., Takeuchi, Y., Mori, Y., 2015. The 3-second auditory conditioned stimulus is a more effective stressor than the 20-second auditory conditioned stimulus in male rats. Neuroscience 299, 79–87. doi:10.1016/j.neuroscience.2015.04.055

Klumpers, F., Kroes, M.C., Heitland, I., Everaerd, D., Akkermans, S.E.A., Oosting, R.S., van Wingen, G., Franke, B., Kenemans, J.L., Fernández, G., Baas, J.M.P., 2015. Dorsomedial prefrontal cortex mediates the impact of serotonin transporter linked polymorphic region

genotype on anticipatory threat reactions. Biol. Psychiatry 78, 582–589. doi:10.1016/j.biopsych.2014.07.034

Klumpers, F., Kroes, M.C.W., Baas, J.M.P., Fernández, G., 2017. How human amygdala and bed nucleus of the stria terminalis may drive distinct defensive responses. J. Neurosci. 37, 9645–9656. doi:10.1523/JNEUROSCI.3830-16.2017

Knapska, E., Maren, S., 2009. Reciprocal patterns of c-Fos expression in the medial prefrontal cortex and amygdala after extinction and renewal of conditioned fear. Learn. Mem. 16, 486–493. doi:10.1101/lm.1463909

Kunwar, P.S., Zelikowsky, M., Remedios, R., Cai, H., Yilmaz, M., Meister, M., Anderson, D.J., 2015. Ventromedial hypothalamic neurons control a defensive emotion state. Elife 4. doi:10.7554/eLife.06633

Lange, M.D., Daldrup, T., Remmers, F., Szkudlarek, H.J., Lesting, J., Guggenhuber, S., Ruehle, S., Jüngling, K., Seidenbecher, T., Lutz, B., Pape, H.C., 2017. Cannabinoid CB1 receptors in distinct circuits of the extended amygdala determine fear responsiveness to unpredictable threat. Mol. Psychiatry 22, 1422–1430. doi:10.1038/mp.2016.156

Lebow, M.A., Chen, A., 2016. Overshadowed by the amygdala: the bed nucleus of the stria terminalis emerges as key to psychiatric disorders. Mol. Psychiatry 21, 450–463. doi:10.1038/mp.2016.1

LeDoux, J., Daw, N.D., 2018. Surviving threats: neural circuit and computational implications of a new taxonomy of defensive behaviour. Nat. Rev. Neurosci. 19, 269–282. doi:10.1038/nrn.2018.22

LeDoux, J.E., Iwata, J., Cicchetti, P., Reis, D.J., 1988. Different projections of the central amygdaloid nucleus mediate autonomic and behavioral correlates of conditioned fear. J. Neurosci. 8, 2517–2529.

Lemos, J.I., Resstel, L.B., Guimarães, F.S., 2010. Involvement of the prelimbic prefrontal cortex on cannabidiol-induced attenuation of contextual conditioned fear in rats. Behav. Brain Res. 207, 105–111. doi:10.1016/j.bbr.2009.09.045

Luyck, K., Goode, T.D., Lee Masson, H., Luyten, L., 2018a. Distinct Activity Patterns of the Human Bed Nucleus of the Stria Terminalis and Amygdala during Fear Learning. Neuropsychol Rev. doi:10.1007/s11065-018-9383-7

Luyck, K., Nuttin, B., Luyten, L., 2018b. Electrolytic post-training lesions of the bed nucleus of the stria terminalis block startle potentiation in a cued fear conditioning procedure. Brain Struct. Funct. 223, 1839–1848. doi:10.1007/s00429-017-1591-z

Luyten, L., Casteels, C., Vansteenwegen, D., van Kuyck, K., Koole, M., Van Laere, K., Nuttin, B., 2012. Micro-positron emission tomography imaging of rat brain metabolism during expression of contextual conditioning. J. Neurosci. 32, 254–263. doi:10.1523/JNEUROSCI.3701-11.2012

Luyten, L., van Kuyck, K., Vansteenwegen, D., Nuttin, B., 2011. Electrolytic lesions of the bed nucleus of the stria terminalis disrupt freezing and startle potentiation in a conditioned context. Behav. Brain Res. 222, 357–362. doi:10.1016/j.bbr.2011.03.066

Mahoney, W.J., Ayres, J.J.B., 1976. One-trial simultaneous and backward fear conditioning as reflected in conditioned suppression of licking in rats. Anim Learn Behav 4, 357–362. doi:10.3758/BF03214421

Marchand, A.R., Luck, D., Di Scala, G., 2004. Trace fear conditioning: a role for context? Arch Ital Biol 142, 251–263.

Marcinkiewcz, C.A., Mazzone, C.M., D'Agostino, G., Halladay, L.R., Hardaway, J.A., DiBerto, J.F., Navarro, M., Burnham, N., Cristiano, C., Dorrier, C.E., Tipton, G.J., Ramakrishnan, C., Kozicz, T., Deisseroth, K., Thiele, T.E., McElligott, Z.A., Holmes, A., Heisler, L.K.,

Kash, T.L., 2016. Serotonin engages an anxiety and fear-promoting circuit in the extended amygdala. Nature 537, 97–101. doi:10.1038/nature19318

Marek, R., Jin, J., Goode, T.D., Giustino, T.F., Wang, Q., Acca, G.M., Holehonnur, R., Ploski, J.E., Fitzgerald, P.J., Lynagh, T., Lynch, J.W., Maren, S., Sah, P., 2018. Hippocampus-driven feed-forward inhibition of the prefrontal cortex mediates relapse of extinguished fear. Nat. Neurosci. 21, 384–392. doi:10.1038/s41593-018-0073-9

Maren, S., 1998. Overtraining does not mitigate contextual fear conditioning deficits produced by neurotoxic lesions of the basolateral amygdala. J. Neurosci. 18, 3088–3097.

Markham, C.M., Norvelle, A., Huhman, K.L., 2009. Role of the bed nucleus of the stria terminalis in the acquisition and expression of conditioned defeat in Syrian hamsters. Behav. Brain Res. 198, 69–73. doi:10.1016/j.bbr.2008.10.022

McDonald, A.J., Shammah-Lagnado, S.J., Shi, C., Davis, M., 1999. Cortical afferents to the extended amygdala. Ann. N. Y. Acad. Sci. 877, 309–338. doi:10.1111/j.1749-6632.1999.tb09275.x

McHugh, S.B., Barkus, C., Lima, J., Glover, L.R., Sharp, T., Bannerman, D.M., 2015. SERT and uncertainty: serotonin transporter expression influences information processing biases for ambiguous aversive cues in mice. Genes Brain Behav. 14, 330–336. doi:10.1111/gbb.12215

McMenamin, B.W., Langeslag, S.J.E., Sirbu, M., Padmala, S., Pessoa, L., 2014. Network organization unfolds over time during periods of anxious anticipation. J. Neurosci. 34, 11261–11273. doi:10.1523/JNEUROSCI.1579-14.2014

McNally, G.P., Johansen, J.P., Blair, H.T., 2011. Placing prediction into the fear circuit. Trends Neurosci. 34, 283–292. doi:10.1016/j.tins.2011.03.005

Meyer, C., Padmala, S., Pessoa, L., 2018. Dynamic Threat Processing. J. Cogn. Neurosci. 1–21. doi:10.1162/jocn_a_01363

Milad, M.R., Quirk, G.J., 2002. Neurons in medial prefrontal cortex signal memory for fear extinction. Nature 420, 70–74. doi:10.1038/nature01138

Miles, O.W., May, V., Hammack, S.E., 2018. Pituitary Adenylate Cyclase-Activating Peptide (PACAP) Signaling and the Dark Side of Addiction. J. Mol. Neurosci. doi:10.1007/s12031-018-1147-6

Mobbs, D., Yu, R., Rowe, J.B., Eich, H., FeldmanHall, O., Dalgleish, T., 2010. Neural activity associated with monitoring the oscillating threat value of a tarantula. Proc. Natl. Acad. Sci. USA 107, 20582–20586. doi:10.1073/pnas.1009076107

Moscovitch, A., LoLordo, V.M., 1968. Role of safety in the Pavlovian backward fear conditioning procedure. J Comp Physiol Psychol 66, 673–678. doi:10.1037/h0026548

Motzkin, J.C., Philippi, C.L., Oler, J.A., Kalin, N.H., Baskaya, M.K., Koenigs, M., 2015. Ventromedial prefrontal cortex damage alters resting blood flow to the bed nucleus of stria terminalis. Cortex 64, 281–288. doi:10.1016/j.cortex.2014.11.013

Münsterkötter, A.L., Notzon, S., Redlich, R., Grotegerd, D., Dohm, K., Arolt, V., Kugel, H., Zwanzger, P., Dannlowski, U., 2015. Spider or no spider? neural correlates of sustained and phasic fear in spider phobia. Depress. Anxiety 32, 656–663. doi:10.1002/da.22382

Naaz, F., Knight, L.K., Depue, B.E., 2019. Explicit and ambiguous threat processing: functionally dissociable roles of the amygdala and bed nucleus of the stria terminalis. J. Cogn. Neurosci. 1–17. doi:10.1162/jocn_a_01369

Nagaya, N., Acca, G.M., Maren, S., 2015. Allopregnanolone in the bed nucleus of the stria terminalis modulates contextual fear in rats. Front. Behav. Neurosci. 9, 205. doi:10.3389/fnbeh.2015.00205

Nees, F., Heinrich, A., Flor, H., 2015. A mechanism-oriented approach to psychopathology: The

role of Pavlovian conditioning. Int. J. Psychophysiol. 98, 351–364. doi:10.1016/j.ijpsycho.2015.05.005

Orsini, C.A., Kim, J.H., Knapska, E., Maren, S., 2011. Hippocampal and prefrontal projections to the basal amygdala mediate contextual regulation of fear after extinction. J. Neurosci. 31, 17269–17277. doi:10.1523/JNEUROSCI.4095-11.2011

Pedersen, W.S., Balderston, N.L., Miskovich, T.A., Belleau, E.L., Helmstetter, F.J., Larson, C.L., 2017. The effects of stimulus novelty and negativity on BOLD activity in the amygdala, hippocampus, and bed nucleus of the stria terminalis. Soc. Cogn. Affect. Neurosci. 12, 748–757. doi:10.1093/scan/nsw178

Pedersen, W.S., Muftuler, L.T., Larson, C.L., 2018. Conservatism and the neural circuitry of threat: economic conservatism predicts greater amygdala-BNST connectivity during periods of threat vs safety. Soc. Cogn. Affect. Neurosci. 13, 43–51. doi:10.1093/scan/nsx133

Perusini, J.N., Fanselow, M.S., 2015. Neurobehavioral perspectives on the distinction between fear and anxiety. Learn. Mem. 22, 417–425. doi:10.1101/lm.039180.115

Pina, M.M., Young, E.A., Ryabinin, A.E., Cunningham, C.L., 2015. The bed nucleus of the stria terminalis regulates ethanol-seeking behavior in mice. Neuropharmacology 99, 627–638. doi:10.1016/j.neuropharm.2015.08.033

Pine, D.S., LeDoux, J.E., 2017. Elevating the role of subjective experience in the clinic: response to fanselow and pennington. Am. J. Psychiatry 174, 1121–1122. doi:10.1176/appi.ajp.2017.17070818r

Poulos, A.M., Ponnusamy, R., Dong, H.-W., Fanselow, M.S., 2010. Compensation in the neural circuitry of fear conditioning awakens learning circuits in the bed nuclei of the stria terminalis. Proc. Natl. Acad. Sci. USA 107, 14881–14886. doi:10.1073/pnas.1005754107

Prével, A., Rivière, V., Darcheville, J.-C., Urcelay, G.P., 2016. Conditioned reinforcement and backward association. Learn Motiv 56, 38–47. doi:10.1016/j.lmot.2016.09.004

Prével, A., Rivière, V., Darcheville, J.-C., Urcelay, G.P., Miller, R.R., 2018. Excitatory second-order conditioning using a backward first-order conditioned stimulus: A challenge for prediction error reduction. Q J Exp Psychol (Colchester) 1747021818793376. doi:10.1177/1747021818793376

Quirk, G.J., Mueller, D., 2008. Neural mechanisms of extinction learning and retrieval. Neuropsychopharmacology 33, 56–72. doi:10.1038/sj.npp.1301555

Rabellino, D., Densmore, M., Harricharan, S., Jean, T., McKinnon, M.C., Lanius, R.A., 2018. Resting-state functional connectivity of the bed nucleus of the stria terminalis in post-traumatic stress disorder and its dissociative subtype. Hum. Brain Mapp. 39, 1367–1379. doi:10.1002/hbm.23925

Radley, J.J., Gosselink, K.L., Sawchenko, P.E., 2009. A discrete GABAergic relay mediates medial prefrontal cortical inhibition of the neuroendocrine stress response. J. Neurosci. 29, 7330–7340. doi:10.1523/JNEUROSCI.5924-08.2009

Radley, J.J., Johnson, S.B., 2018. Anteroventral bed nuclei of the stria terminalis neurocircuitry: Towards an integration of HPA axis modulation with coping behaviors - Curt Richter Award Paper 2017. Psychoneuroendocrinology 89, 239–249. doi:10.1016/j.psyneuen.2017.12.005

Radley, J.J., Sawchenko, P.E., 2011. A common substrate for prefrontal and hippocampal inhibition of the neuroendocrine stress response. J. Neurosci. 31, 9683–9695. doi:10.1523/JNEUROSCI.6040-10.2011

Raybuck, J.D., Lattal, K.M., 2014. Bridging the interval: theory and neurobiology of trace conditioning. Behav. Processes 101, 103–111. doi:10.1016/j.beproc.2013.08.016

Reichard, R.A., Subramanian, S., Desta, M.T., Sura, T., Becker, M.L., Ghobadi, C.W., Parsley, K.P., Zahm, D.S., 2017. Abundant collateralization of temporal lobe projections to the accumbens, bed nucleus of stria terminalis, central amygdala and lateral septum. Brain Struct. Funct. 222, 1971–1988. doi:10.1007/s00429-016-1321-y

Rescorla, R.A., 1968. Probability of shock in the presence and absence of CS in fear conditioning. J Comp Physiol Psychol 66, 1–5. doi:10.1037/h0025984

Reynolds, S.M., Zahm, D.S., 2005. Specificity in the projections of prefrontal and insular cortex to ventral striatopallidum and the extended amygdala. J. Neurosci. 25, 11757–11767. doi:10.1523/JNEUROSCI.3432-05.2005

Sajdyk, T., Johnson, P., Fitz, S., Shekhar, A., 2008. Chronic inhibition of GABA synthesis in the bed nucleus of the stria terminalis elicits anxiety-like behavior. J. Psychopharmacol. (Oxford) 22, 633–641. doi:10.1177/0269881107082902

Salas-Wright, C.P., Kagotho, N., Vaughn, M.G., 2014. Mood, anxiety, and personality disorders among first and second-generation immigrants to the United States. Psychiatry Res. 220, 1028–1036. doi:10.1016/j.psychres.2014.08.045

Sangha, S., Robinson, P.D., Greba, Q., Davies, D.A., Howland, J.G., 2014. Alterations in reward, fear and safety cue discrimination after inactivation of the rat prelimbic and infralimbic cortices. Neuropsychopharmacology 39, 2405–2413. doi:10.1038/npp.2014.89

Schlund, M.W., Hudgins, C.D., Magee, S., Dymond, S., 2013. Neuroimaging the temporal dynamics of human avoidance to sustained threat. Behav. Brain Res. 257, 148–155. doi:10.1016/j.bbr.2013.09.042

Schneider, C.A., Rasband, W.S., Eliceiri, K.W., 2012. NIH Image to ImageJ: 25 years of image analysis. Nat. Methods 9, 671–675.

Schroijen, M., Fantoni, S., Rivera, C., Vervliet, B., Schruers, K., van den Bergh, O., van Diest, I., 2016. Defensive activation to (un)predictable interoceptive threat: The NPU respiratory threat test (NPUr). Psychophysiology 53, 905–913. doi:10.1111/psyp.12621

Seidenbecher, T., Remmes, J., Daldrup, T., Lesting, J., Pape, H.-C., 2016. Distinct state anxiety after predictable and unpredictable fear training in mice. Behav. Brain Res. 304, 20–23. doi:10.1016/j.bbr.2016.02.009

Shackman, A.J., Fox, A.S., 2016. Contributions of the central extended amygdala to fear and anxiety. J. Neurosci. 36, 8050–8063. doi:10.1523/JNEUROSCI.0982-16.2016

Shankman, S.A., Robison-Andrew, E.J., Nelson, B.D., Altman, S.E., Campbell, M.L., 2011. Effects of predictability of shock timing and intensity on aversive responses. Int. J. Psychophysiol. 80, 112–118. doi:10.1016/j.ijpsycho.2011.02.008

Siegel, S., Domjan, M., 1971. Backward conditoning as an inhibitory procedure. Learn Motiv 2, 1–11. doi:10.1016/0023-9690(71)90043-9

Silberman, Y., Winder, D.G., 2013. Emerging role for corticotropin releasing factor signaling in the bed nucleus of the stria terminalis at the intersection of stress and reward. Front. Psychiatry 4, 42. doi:10.3389/fpsyt.2013.00042

Sinnema, H., Majo, M.C., Volker, D., Hoogendoorn, A., Terluin, B., Wensing, M., van Balkom, A., 2015. Effectiveness of a tailored implementation programme to improve recognition, diagnosis and treatment of anxiety and depression in general practice: a cluster randomised controlled trial. Implement. Sci. 10, 33. doi:10.1186/s13012-015-0210-8

Sladky, R., Geissberger, N., Pfabigan, D.M., Kraus, C., Tik, M., Woletz, M., Paul, K., Vanicek, T., Auer, B., Kranz, G.S., Lamm, C., Lanzenberger, R., Windischberger, C., 2018. Unsmoothed functional MRI of the human amygdala and bed nucleus of the stria terminalis during processing of emotional faces. Neuroimage 168, 383–391.

doi:10.1016/j.neuroimage.2016.12.024

Somerville, L.H., Whalen, P.J., Kelley, W.M., 2010. Human bed nucleus of the stria terminalis indexes hypervigilant threat monitoring. Biol. Psychiatry 68, 416–424. doi:10.1016/j.biopsych.2010.04.002

Stamatakis, A.M., Sparta, D.R., Jennings, J.H., McElligott, Z.A., Decot, H., Stuber, G.D., 2014. Amygdala and bed nucleus of the stria terminalis circuitry: Implications for addiction-related behaviors. Neuropharmacology 76 Pt B, 320–328. doi:10.1016/j.neuropharm.2013.05.046

Stein, D.J., Scott, K.M., de Jonge, P., Kessler, R.C., 2017. Epidemiology of anxiety disorders: from surveys to nosology and back. Dialogues Clin Neurosci 19, 127–136.

Sterrenburg, L., Gaszner, B., Boerrigter, J., Santbergen, L., Bramini, M., Roubos, E.W., Peeters, B.W.M.M., Kozicz, T., 2012. Sex-dependent and differential responses to acute restraint stress of corticotropin-releasing factor-producing neurons in the rat paraventricular nucleus, central amygdala, and bed nucleus of the stria terminalis. J. Neurosci. Res. 90, 179–192. doi:10.1002/jnr.22737

Straube, T., Mentzel, H.-J., Miltner, W.H.R., 2007. Waiting for spiders: brain activation during anticipatory anxiety in spider phobics. Neuroimage 37, 1427–1436. doi:10.1016/j.neuroimage.2007.06.023

Sullivan, G.M., Apergis, J., Bush, D.E.A., Johnson, L.R., Hou, M., Ledoux, J.E., 2004. Lesions in the bed nucleus of the stria terminalis disrupt corticosterone and freezing responses elicited by a contextual but not by a specific cue-conditioned fear stimulus. Neuroscience 128, 7–14. doi:10.1016/j.neuroscience.2004.06.015

Swanson, L., 2003. Brain Maps, Third Edition: Structure of the Rat Brain (Vol 3), 3rd ed. Academic Press.

Takahashi, L.K., 2014. Olfactory systems and neural circuits that modulate predator odor fear. Front. Behav. Neurosci. 8, 72. doi:10.3389/fnbeh.2014.00072

Tipps, M.E., Raybuck, J.D., Buck, K.J., Lattal, K.M., 2014. Delay and trace fear conditioning in C57BL/6 and DBA/2 mice: issues of measurement and performance. Learn. Mem. 21, 380–393. doi:10.1101/lm.035261.114

Torrisi, S., Gorka, A.X., Gonzalez-Castillo, J., O'Connell, K., Balderston, N., Grillon, C., Ernst, M., 2018. Extended amygdala connectivity changes during sustained shock anticipation. Transl. Psychiatry 8, 33. doi:10.1038/s41398-017-0074-6

Torrisi, S., O'Connell, K., Davis, A., Reynolds, R., Balderston, N., Fudge, J.L., Grillon, C., Ernst, M., 2015. Resting state connectivity of the bed nucleus of the stria terminalis at ultra-high field. Hum. Brain Mapp. 36, 4076–4088. doi:10.1002/hbm.22899

Tye, K.M., 2018. Neural circuit motifs in valence processing. Neuron 100, 436–452. doi:10.1016/j.neuron.2018.10.001

Verma, D., Tasan, R., Sperk, G., Pape, H.-C., 2018. Neuropeptide Y2 receptors in anteroventral BNST control remote fear memory depending on extinction training. Neurobiol. Learn. Mem. 149, 144–153. doi:10.1016/j.nlm.2018.01.011

Vertes, R.P., 2004. Differential projections of the infralimbic and prelimbic cortex in the rat. Synapse 51, 32–58. doi:10.1002/syn.10279

Waddell, J., Bouton, M.E., Falls, W.A., 2008. Central CRF receptor antagonist a-helical CRF9-41 blocks reinstatement of extinguished fear: the role of the bed nucleus of the stria terminalis. Behav. Neurosci. 122, 1061–1069. doi:10.1037/a0013136

Waddell, J., Morris, R.W., Bouton, M.E., 2006. Effects of bed nucleus of the stria terminalis lesions on conditioned anxiety: aversive conditioning with long-duration conditional stimuli and reinstatement of extinguished fear. Behav. Neurosci. 120, 324–336.

    doi:10.1037/0735-7044.120.2.324

Walker, D.L., Davis, M., 2008. Role of the extended amygdala in short-duration versus sustained fear: a tribute to Dr. Lennart Heimer. Brain Struct. Funct. 213, 29–42. doi:10.1007/s00429-008-0183-3

Walker, D.L., Miles, L.A., Davis, M., 2009. Selective participation of the bed nucleus of the stria terminalis and CRF in sustained anxiety-like versus phasic fear-like responses. Prog. Neuropsychopharmacol. Biol. Psychiatry 33, 1291–1308. doi:10.1016/j.pnpbp.2009.06.022

Wang, Q., Jin, J., Maren, S., 2016. Renewal of extinguished fear activates ventral hippocampal neurons projecting to the prelimbic and infralimbic cortices in rats. Neurobiol. Learn. Mem. 134 Pt A, 38–43. doi:10.1016/j.nlm.2016.04.002

Weller, K.L., Smith, D.A., 1982. Afferent connections to the bed nucleus of the stria terminalis. Brain Res. 232, 255–270. doi:10.1016/0006-8993(82)90272-4

Wittchen, H.-U., 2002. Generalized anxiety disorder: prevalence, burden, and cost to society. Depress. Anxiety 16, 162–171. doi:10.1002/da.10065

Wood, M., Adil, O., Wallace, T., Fourman, S., Wilson, S.P., Herman, J.P., Myers, B., 2018. Infralimbic prefrontal cortex structural and functional connectivity with the limbic forebrain: a combined viral genetic and optogenetic analysis. Brain Struct. Funct. doi:10.1007/s00429-018-1762-6

Xu, H.Y., Liu, Y.J., Xu, M.Y., Zhang, Y.H., Zhang, J.X., Wu, Y.J., 2012. Inactivation of the bed nucleus of the stria terminalis suppresses the innate fear responses of rats induced by the odor of cat urine. Neuroscience 221, 21–27. doi:10.1016/j.neuroscience.2012.06.056

Yassa, M.A., Hazlett, R.L., Stark, C.E.L., Hoehn-Saric, R., 2012. Functional MRI of the amygdala and bed nucleus of the stria terminalis during conditions of uncertainty in generalized anxiety disorder. J. Psychiatr. Res. 46, 1045–1052. doi:10.1016/j.jpsychires.2012.04.013

Zelikowsky, M., Bissiere, S., Hast, T.A., Bennett, R.Z., Abdipranoto, A., Vissel, B., Fanselow, M.S., 2013. Prefrontal microcircuit underlies contextual learning after hippocampal loss. Proc. Natl. Acad. Sci. USA 110, 9938–9943. doi:10.1073/pnas.1301691110

Zelikowsky, M., Hui, M., Karigo, T., Choe, A., Yang, B., Blanco, M.R., Beadle, K., Gradinaru, V., Deverman, B.E., Anderson, D.J., 2018. The neuropeptide tac2 controls a distributed brain state induced by chronic social isolation stress. Cell 173, 1265–1279.e19. doi:10.1016/j.cell.2018.03.037

Zhu, W., Umegaki, H., Suzuki, Y., Miura, H., Iguchi, A., 2001. Involvement of the bed nucleus of the stria terminalis in hippocampal cholinergic system-mediated activation of the hypothalamo--pituitary--adrenocortical axis in rats. Brain Res. 916, 101–106.

Zimmerman, J.M., Maren, S., 2011. The bed nucleus of the stria terminalis is required for the expression of contextual but not auditory freezing in rats with basolateral amygdala lesions. Neurobiol. Learn. Mem. 95, 199–205. doi:10.1016/j.nlm.2010.11.002