Title: The *Drosophila* chromodomain protein Kismet activates steroid hormone receptor

transcription to govern axon pruning and memory *in vivo*

Authors: Nina K. Latcheva[1,2], Jennifer M. Viveiros[1], and Daniel R. Marenda[1,2,3*]

Addresses: [1]Department of Biology, Drexel University, Philadelphia, PA, and [2]Program

in Molecular and Cellular Biology and Genetics, Drexel University College of Medicine,

Philadelphia, PA; [3]Department of Neurobiology and Anatomy, Drexel University College

of Medicine, Philadelphia, PA

*Correspondence: Daniel R. Marenda, Department of Biology, Drexel University, 3141

Chestnut St., Philadelphia, PA 19104.

e-mail: daniel.marenda@drexel.edu

Short Title: Epigenetic control of axon pruning

Key Words: *Drosophila*, *kismet*, axon pruning, Ecdysone Receptor, CHARGE Syndrome,

CHD7.

Abstract

Axon pruning is critical for the proper synapse elimination required for sculpting precise neural circuits. Though axon pruning has been described in the literature for decades, relatively little is known about the molecular and cellular mechanisms that govern axon pruning *in vivo*. Here, we show that the epigenetic reader Kismet (Kis) binds to cis-regulatory elements of the steroid hormone receptor *Ecdysone Receptor* (*EcR*) gene in *Drosophila* neurons. Kis is required to activate *EcR* transcription at these elements and promote H3K36 di- and tri-methylation. We show that exogenous EcR can rescue axon pruning and memory defects associated with loss of Kis, and that the histone deacetylase inhibitor SAHA can rescue these phenotypes. EcR protein abundance is a cell-autonomous, rate-limiting step required to initiate axon pruning in *Drosophila,* and our data suggests that this step is under epigenetic control.

Running Title

Kismet activates EcR to promote axon pruning *in vivo*

Highlights (bullet points up to 4)

- The chromodomain reader Kismet activates transcription of the steroid hormone receptor *EcR-B1* in the *Drosophila* to initiate developmental axon pruning.

- Kismet promotes H3K36 di- and tri-methylation at the *EcR* locus and upstream cis-regulatory sites.

- Axon pruning and memory defects associated with loss of Kismet are significantly rescued by general HDAC inhibition.

Introduction

The elimination and refinement of synaptic connections is an integral part of normal development in vertebrates and invertebrates alike. Early in the developing nervous system periods of progressive growth result in an overelaboration of synaptic connections onto a target. Inappropriate synapses then need to be eliminated to establish functional organization of the neuronal circuitry [1]. The pruning of these exuberant connections can occur on a small scale, as with dendritic remodeling, or on a large scale such as with axon retraction and degeneration, with each type occurring through distinct molecular mechanisms [2]. Precise control of axon pruning is critical for proper nervous system function, as defects in pruning have been well documented to lead to developmental brain and psychiatric disorders [1]. Despite its vital role, relatively little is known about the mechanisms that govern axon pruning *in vivo*. What is well known about this process is that it requires tight regulation of gene expression in order to execute the necessary signaling pathways in a temporal and tissue specific manner [3,4]. Epigenetic regulation is key to orchestrating precise gene expression programs for many tightly controlled processes in the body. However, its involvement in axon pruning is still unclear.

Holometabolous insects provide an attractive model for studying axon pruning as their nervous system undergoes extensive reorganization during metamorphosis [5,6]. In *Drosophila melanogaster* the larval neuronal circuitry is eliminated to make way for adult specific circuitry that governs adult behavior. The most notable changes occur in the learning and memory processing center of the fly brain known as the mushroom bodies (MBs) [7-10]. MB neurons undergo a selective, stereotypical pruning during metamorphosis, which is initiated by the steroid hormone 20-hydroxyecdysone (ecdysone) [3,11,12]. Ecdysone

binds to the Ecdysone Receptor (EcR), and translocates to the nucleus to initiate

transcription of a subset of regulatory target genes required for axon pruning. MB neurons

that undergo remodeling express a particular isoform of the EcR, EcR-B1, which has been

shown to be a rate-limiting and cell-autonomous step required for developmental axon

pruning [11].

Our lab previously identified the chromodomain protein Kismet (Kis) as necessary for

proper developmental axon pruning in the *Drosophila* MB neurons [13]. Kis is the

Drosophila ortholog of the mammalian chromatin ATPase CHD7, a chromatin "reader"

that is thought to play a role in chromatin remodeling by binding to methylated histone

tails [14]. In humans, heterozygous mutations in *CHD7* cause CHARGE syndrome [15], an

autosomal dominant neurodevelopmental disorder.

Here, we investigated the role of Kis in the developmental axon pruning of the

Drosophila MB neurons. We determined that the loss of Kis in the MBs results in pruning

defects during metamorphosis, which persist into adulthood and are due to a decrease in

transcription of *EcR-B1*. We show that endogenous Kis is enriched at cis-regulatory

enhancer sites as well as the transcriptional start site (TSS) of the *EcR* locus. Further, loss

of Kis leads to a decrease in the histone marks H3K36me2 and H3K36me3, which have

been associated with actively transcribed genes. Adult flies with Kis specifically decreased

in the MB neurons display a loss of immediate recall memory, which is rescued by

exogenous expression of EcR-B1. Finally, we show that pharmacological intervention via

the general histone deacetylase (HDAC) inhibitor suberoylanilide hydroxamic acid

(SAHA) can rescue the decrease in *EcR-B1* mRNA, axon pruning, and memory defects

associated with decreased Kis in MB neurons. Taken together, these data show that Kis-

mediated regulation of *EcR-B1* is required for proper developmental axon pruning *in vivo*

by mediating the epigenetic marks H3K36me2 and H3K36me3. These novel findings

suggest that the rate−limiting step required to initiate axon pruning (*EcR-B1* expression) is

under epigenetic control.

Results

Kismet is required for MB pruning

To characterize the pruning defects previously observed in *kis* mutant MB neurons,

we utilized the mosaic analysis with a repressible cell marker (MARCM) system to

generate homozygous mutant neuroblast clones tagged with a membrane bound GFP using

the *201Y*-Gal-4 driver [13,16-18]. We measured dorsal, medial, and total surface area of the

MB lobes in pupal brains 18-22 hours after puparium formation (APF). At this timepoint,

the MB lobes are mostly pruned away in control animals (Fig. 1A, 1E). In the null mutant

kis^{LM27}, MB clones had significantly larger medial and total lobe surface areas compared

to control MB clones (Figs. 1A, 1B, 1E). MB clones expressing the wild-type Kis-L protein

in the *kis^{LM27}* mutant background showed a significant reduction of the medial and total

lobe areas back to control levels (Figs. 1D, 1E). Overexpression of Kis-L alone did not

have an effect on lobe surface area (Figs. 1C, 1E). In addition to MARCM analysis, we

utilized RNA interference (RNAi) mediated knockdown of Kis with two separate

previously validated RNAi constructs to verify the pruning defects [13]. Pan-neural

knockdown of Kis using the *elav*-Gal-4 driver showed a significant increase in the medial

and total lobe areas compared to outcross controls (Supplemental Figure 1). Similar to the

MARCM analysis, pan-neural expression of Kis-L in the knockdown genetic backgrounds

was able to significantly rescue the medial and total surface area levels (Supplemental

Figure 1). Taken together, these data show that Kis is required for the developmental axon

pruning of MB neurons during metamorphosis.

Kismet binds EcR and activates EcR transcription

Expression of the steroid hormone receptor EcR-B1 is the first step in the

developmental axon pruning of the MB neurons, and loss of EcR-B1 function produces

defects in pruning similar to what we observe in decreased Kis function [11]. Therefore, we

sought to determine if Kis affects levels of EcR-B1. RNAi knockdown of Kis in MB

neurons *in vivo* showed a significant decrease in EcR-B1 immunofluorescence compared

to control MBs (Figs. 1F-1K, Fig. 1R). In support of this, mRNA levels of EcR-B1 were

significantly reduced in pupae with decreased Kis as well (Fig. 1S). Finally, we wanted to

determine if addition of exogenous Kis-L in Kis knockdown MBs could rescue the loss of

EcR-B1 expression. We found that replacement of Kis-L protein did successfully rescue

both decreased EcR-B1 mRNA and protein expression in these neurons (Figs. 1L-1S).

Taken together, these data suggest that Kis is required to promote EcR-B1 expression

within MB neurons.

Given that Kis is an epigenetic chromatin reader, we theorized that it may be

affecting EcR-B1 mRNA and protein levels by promoting transcription. To test this, we

performed chromatin immunoprecipitation (ChIP) followed by qPCR to analyze Kis

occupancy at previously published cis-regulatory elements of the *EcR* locus as well as the

EcR TSS (Fig. 2A) [19]. We utilized a Kis-eGFP protein trap animal previously described to

express a eGFP tagged version of the Kis protein [20] [21]. Importantly, this animal shows

normal localization and function of Kis, and we have shown that our RNAi reagents can

successfully knockdown the eGFP tagged Kis protein and *kis* mRNA (Supplemental Figure

2) [20]. We utilized the *forkhead* (*fkh*) TSS as a positive control, as it was previously reported

to be bound by endogenous Kis [22]. Additionally, we used the *dynamin* homolog *shibire*

(*shi*) as a negative control, as we had previously shown via microarray analysis that loss of

Kis did not have any significant effect on *shi* expression [20]. We verified that Kis was not

enriched at the *shi* promoter region (Fig. 2A), and that knockdown of Kis did not have any

significant effect on Kis abundance at the *shi* promoter (Fig. 2A). In contrast, wild-type

control brains showed enrichment of Kis at the two previously reported *EcR* cis-regulatory

elements (*EcR.1* and *EcR.2*) as well as the *EcR* and *fkh* TSS (Fig. 2A). Finally, we observed

a significant decrease in Kis enrichment at the *fkh* TSS, *EcR.1*, *EcR.2*, and *EcR* TSS sites

when Kis was knocked down, confirming specificity of Kis binding at these sites (Fig. 2A).

These results suggest that Kis binds to known cis-regulatory sites and the TSS of the *EcR*

locus in *Drosophila* larval brains.

To determine if Kis can control transcriptional output from these sites, we utilized

an EcR transcriptional reporter (*EcR.1*-Gal-4, described in [23,24]). This reporter drives the

expression of the exogenous Gal-4 protein from the *EcR.1* cis-regulatory element that is

partially responsible for the expression of *EcR-B1* in MB neurons [19] . Using this reporter,

we determined the levels of Gal-4 mRNA and protein in wild-type control MBs and Kis

knockdown MBs. We observed a significant decrease in both Gal-4 mRNA and protein

levels from this reporter in Kis knockdown animals compared to controls (Figs. 2C-G).

Taken together, these data suggest that Kis protein can bind to *EcR* cis-regulatory elements

and the *EcR* TSS to promote transcriptional activation and increase EcR-B1 expression in

the *Drosophila* nervous system.

Kismet promotes H3K36me2 and H3K36me3

165   One mechanism by which chromatin remodeling proteins promote transcriptional

activation is by mobilizing nucleosomes and allowing transcriptional machinery access to

enhancer sites and promoters of target genes [25]. Given Kis's homology to CHD7 and its

conserved ATPase domain, we hypothesized that Kis may be removing nucleosomes from

the *EcR* locus to allow access to the transcriptional machinery. To test this possibility, we

immunoprecipitated total histone 3 (H3) protein at the *EcR* locus in control and Kis

knockdown brains. However, we found no significant difference between these conditions

at each of the cis-regulatory loci we analyzed (Fig. 2B). To verify this, we performed an

MNase protection assay. Consistent with our total H3 analysis, we saw no change in the

digestion of chromatin upon Kis knockdown (Supplemental Figure 3). Taken together,

these data suggest that Kis is not affecting the movement of nucleosomes as a mechanism

to promote transcription at the *EcR* locus.

177   Another way chromatin readers can affect gene expression is by altering the histone

modifications present at relevant genomic loci [25], and Kis has been shown to affect histone

modifications previously [22]. We began by analyzing H3K4 methylation states at *EcR* loci

in control and Kis knockdown animals, as it is the type of methylation most commonly

associated with actively transcribed genes. ChIP-qPCR analysis showed no change in all

types of H3K4 methylation levels upon Kis knockdown at these loci (Supplemental Figs.

4A-C). We next analyzed H3K27 trimethylation, as this modification is often associated

with gene repression. We did not observe any significant change in H3K27 trimethylation

upon Kis knockdown at any of the loci examined (Supplemental Fig. 4D). Taken together,

these data suggest that alteration of H3K4 methylation or H3K27 trimethylation is not part

of the mechanism by which Kis controls *EcR* transcription.

188   Previous studies demonstrated a global decrease in H3K36 di- and tri-methylation

upon Kis loss in *Drosophila* salivary gland polytene chromosomes [26]. In *Drosophila*, this

type of modification is usually associated with actively transcribed genes [26-28]. We

therefore sought to determine if the same effect on H3K36 methylation was present in the

Drosophila nervous system upon pan-neural knockdown of Kis protein. We observed that

H3K36me2 was significantly decreased at all the *EcR* cis-regulatory sites, as well as at the

fkh positive control, in Kis knockdown brains compared to controls (Fig. 2H). Additionally,

H3K36me3 was also significantly decreased at the *EcR* and *fkh* TSS (Fig. 2I). Importantly,

this decrease in H3K36 methylation was not global, as there was no significant change with

either H3K36 di- or tri-methylation at the *shi* promoter which is not bound by Kis (Figs.

2A, 2H-I). Also, given that H3K36me2 is the substrate for the tri-methylated form, it is

likely that the decrease in H3K36me3 is due to the decrease in H3K36me2. Combined,

these results suggest Kis affects transcription at the *EcR* locus by promoting the active

H3K36 di- and tri-methylation histone marks.

EcR-B1 rescues axon pruning defects

204   EcR-B1 is well documented to be a key player in initiating the axon pruning

cascade in MB neurons [4,11]. Given that we have shown Kis acts to promote transcription of

EcR-B1 in MB neurons, we sought to determine if we could rescue the pruning defects we

observe is Kis mutants by expression of exogenous EcR-B1. Utilizing the MARCM

system, we expressed exogenous EcR-B1 protein within *kis^{LM27}* mutant MB clones (Fig.

3). We observed that exogenous expression of EcR-B1 significantly reduced the abnormal

pruning observed in *kis^{LM27}* mutant MB clones (Figs. 3B, 3D, 3E). We also observed a

significant reduction in pruning defects when exogenous EcR-B1 was pan-neurally co-

expressed with Kis RNAi constructs (Supplemental Fig. 5). Interestingly, EcR-B1

overexpression alone, as well as EcR-B1 expression in Kis loss of function backgrounds,

produced smaller surface areas in the dorsal and total lobes compared to outcross controls

(Figs. 3A, 3C, 3E-3F). This is consistent with Kis functioning upstream of EcR-B1 in the

pruning process, as well as with EcR-B1 being the rate-limiting factor for pruning. Taken

together, these data suggest that Kis mediates axon pruning in MB neurons by

transcriptionally activating *EcR-B1*, thereby controlling EcR-B1 protein levels.

Pruning defects observed during metamorphosis may simply reflect a delay in

normal pruning. Therefore, to determine that homozygous MARCM *kis^{LM27}* mutant clones

indeed have defective developmental pruning, as opposed to having delayed pruning, we

sought to determine if the unpruned axons persisted into adulthood. We generated

MARCM clones with homozygous *kis^{LM27}* MBs as previously described and aged the

adults for 5 days after eclosion. We then immunostained the adult brains with anti-FASII,

a transmembrane cell adhesion protein, which is differentially expressed in the separate

populations of MB neurons [29,30]. FASII expression is lowest in the early born gamma

neurons and highest in the late born alpha/beta neurons; therefore, the appearance of GFP

labeled MARCM axons in the dorsal lobe that are weakly or unstained for FASII would

constitute aberrant unpruned axons that persisted into adulthood [29]. Compared to control

MBs, kis^{LM27} MARCM clones had significantly more weakly-stained and/or unstained

FASII GFP-positive axons outside the dorsal lobe bundle, indicating that pruning is in fact

prevented and not delayed in this mutant (Figs. 3F-3K, 3R). Given that exogenous

expression of EcR-B1 in the kis^{LM27} MARCM background rescued the pruning defects at

the pupal stage, we tested whether the rescue effects of exogenous EcR-B1 provided at the

pupal stage also persisted into adulthood. We observed that expression of EcR-B1 in kis^{LM27}

mutant MARCM clones showed significantly fewer GFP-positive axons outside the dorsal

lobe in adult MB axons, while expression of EcR-B1 alone had no effect (Figs. 3L-3Q,

3R). Collectively, these data suggest that the defective pruning observed during

metamorphosis in kis loss-of-function MB neurons persists into adulthood.

EcR-B1 rescues memory defects

Previous studies from our lab showed that reduction of Kis levels in MB neurons

produced significant defects in immediate recall memory [13]. Since loss of Kis leads to

pruning defects that persist into adulthood, and exogenous EcR-B1 expression is able to

rescue the defects, we were interested in determining if exogenous expression of EcR-B1

could also rescue the memory defect associated with loss of Kis function as well. We

utilized the conditioned courtship suppression assay which takes advantage of the innate

courting behaviors carried out by male *Drosophila* in response to multimodal signals

transduced by females [31-33]. Wild-type males will decrease their rate of courting during

training when exposed to an unresponsive female, and continue to court at lower rates even

with subsequent receptive females for an average of 1-3 hours after exposure [31-33]. We

utilized RNAi to knockdown Kis in MB neurons using the *OK107*-Gal4 driver.

Importantly, expression of the Gal4 alone, or Gal4-mediated expression of exogenous EcR-

B1 did not produce any learning (Fig. 4A) or memory (Fig. 4B) defects. We observed that

males with decreased Kis protein displayed intact learning, as evident by the significant

decrease in courtship from the initial to final stages of exposure to an unresponsive female

(Fig. 4C). Similarly, males with both decreased Kis and exogenously expressed EcR-B1

also had intact learning (Fig. 4C). However, males with decreased Kis showed abnormal

memory (Fig. 4D), as these males had rates of courtship that were not significantly different

than sham males, which did not receive exposure to an unreceptive female. In contrast,

males with both decreased Kis and exogenously expressed EcR-B1 showed significantly

reduced courtship in trained males compared to sham males, indicative of intact immediate

recall memory (Fig. 4D). Taken together, these data suggest that the memory defects

associated with loss of Kis in the MB neurons are due to decreased EcR-B1 levels.

SAHA rescues Kis defects

Recently, our lab showed that pharmacological inhibition of HDACs can rescue

multiple defects associated with loss of Kis at the neuromuscular junction [34]. HDAC

inhibition (HDACi) did not significantly affect *kis* mRNA levels [34], so we hypothesized

that rescue may be due to effect on target genes. To further examine this hypothesis, we

tested whether HDACi treatment could rescue the decreased EcR-B1 levels, pruning, and

memory defects observed in Kis knockdown. We observed that treatment of Kis

knockdown animals with SAHA significantly increased *EcR-B1* mRNA levels compared

to dimethyl sulfoxide (DMSO) treated controls (Fig. 5S). SAHA treatment alone had no

significant impact on axon pruning in either pupal or adult MBs (Figs. 5E, 5R). However,

we observed that SAHA treatment significantly decreased the number of unpruned axons

in both pupal and adult brains in Kis loss-of-function MB neurons (Figs. 5E, 5R). Finally,

SAHA treatment itself did not have an adverse effect on learning or memory in controls

(Figs. 6A-6B) and was able to significantly rescue the immediate recall defect in Kis

knockdown animals compared to DMSO treatment alone (Figs. 6C-6D). Taken together,

these results show that HDACi treatment significantly rescues multiple defects associated

with Kis loss-of-function.

Discussion

Axon pruning and elimination are critical steps to establishing and refining neural

circuitry. However, relatively little is known about the molecular details that control

developmental axon pruning. Here we begin to unravel the epigenetic mechanisms

essential for initiating developmental axon pruning *in vivo*. We show that the chromatin

reader Kis activates transcription of *EcR-B1* in the *Drosophila* MBs and promotes

methylation of H3K36 at cis-regulatory sites and the TSS of *EcR-B1*. We show that proper

regulation of EcR-B1 by Kis is required for developmental pruning and immediate recall

memory in adults. Finally, we show that the general HDACi SAHA can increase *EcR-B1*

levels in animals with decreased Kis and rescue the pruning and memory defects. This is

the first study demonstrating that the essential rate-limiting step in developmental axon

pruning, EcR-B1 expression, is under epigenetic control. Few studies have hinted at an

epigenetic mechanism of EcR activation. The *Drosophila* Set2 K36 histone methyl

transferase (HMT) was shown to genetically interact with the EcR signaling pathway and

positively regulate expression of EcR target genes [27]. However, as dSet2 is required for

general transcriptional elongation, this may be a non-specific effect of dSet2 function.

Nonetheless, this implicates H3K36 methylation as an important factor in EcR expression.

While Kis does not appear to affect expression of dSet2, Ash1 or dMes-4 (data not shown),

it might be acting to recruit the HMTs to relevant target genes. To this effect, Kis was

previously shown to increase the global association of Ash1 on polytene chromosomes

[26,35]. While other studies with Kis have implied a role for the chromatin reader in

transcriptional elongation [36], we demonstrate a selectivity for Kis binding and Kis-

mediated elevation of H3K36 methylation as no significant changes were observed at the

shi promoter in Kis knockdown animals. Further work needs to be done to tease apart the

relationship between Kis, H3K36 methylation, and any HMTs responsible for this

modification in EcR regulation.

Much work has been done to understand the crosstalk between different histone

modifications and their cumulative outcome on transcription. For example, H3K36me2

was reported to specifically increase H4K16 acetylation, a modification most well-known

for its role in decondensation of chromatin structure [37,38]. Additionally, this type of

acetylation   directly   influences   transcription   by   positively   regulating   chromatin

accessibility to non-histone proteins [39]. It is therefore plausible that Kis might be affecting

transcriptional output by promoting H4K16 acetylation via H3K36me2 crosstalk. This may

explain why HDAC inhibition with SAHA, seemingly unrelated to histone methylation, is

able to restore *EcR-B1* mRNA levels due to loss of Kis and rescue pruning and memory

defects. Furthermore, SAHA treatment has been previously shown to increase H4K16ac

and ultimately increase transcriptional output in cancer cells [40]. Further work is essential

to building a complete understanding of the integration of extrinsic and intrinsic cues that

control transcriptional regulation. This is fundamental to unraveling the mechanisms that

underlie the refinement of neural circuits.

Figure 1

Figure 1. Kis is required for axon pruning and EcR expression. (A-D) Representative
images of MARCM generated MB clones expressing membrane bound GFP using the
201Y-Gal-4 driver 18-22 hours APF. (E) Quantification of dorsal, medial, and total MB
lobe surface areas in MARCM animals (n = 8-12). (F-Q) Representative *elav*-Gal-4 driven
control, knockdown, overexpression, and rescue pupal brains expressing membrane bound
GFP or stained with α-EcR-B1, as indicated. (R) Quantification of fluorescent intensity of
EcR-B1 in *elav*-Gal-4 driven pupae (n = 9-12). (S) *EcR-B1* mRNA levels in pupal brains
of *elav*-Gal-4 driven animals analyzed by RT-qPCR (n = 3-4). Scale is 10 μm in (A) and
20 μm in (F).

Figure 2

Figure 2. Kis binds to EcR loci promoting EcR-B1 transcription and H3K36 methylation. (A) ChIP-qPCR analysis using chromatin isolated from 3rd instar larval brains. Kis enrichment at the *EcR* TSS and enhancer sites, the *fkh* TSS, and the *shi* promoter site (n = 4-5). (B) Total H3 as a percentage of the input DNA was determined by qPCR at the previously noted genomic loci (n = 6). (C-E) Representative images of control and knockdown *EcR.1*-Gal-4 driven pupal brains stained with α-Gal-4. (F) Quantification of fluorescent intensity of α-Gal-4 (n = 10). (G) *Gal-4* mRNA levels of *EcR.1*-Gal-4 driven animals analyzed by RT-qPCR (n = 3). (H, I) ChIP-qPCR analysis of

371 H3K36me2 and H3K36me3 abundance relative to total H3 at the above mentioned
372 genomic loci, respectively (n = 3). Scale is 20 μm in (C).
373
374

Figure 3.

Figure 3. Exogenous EcR-B1 rescues defective axon pruning associated with loss of
Kis. (A-D) Representative images of MARCM generated MB clones expressing
membrane bound GFP using the *201Y*-Gal-4 driver 18-22 hours APF. (E) Quantification
of dorsal, medial, and total MB lobe surface areas in MARCM animals (n = 10-12). (F-
Q) Representative adult MARCM generated MB clones expressing membrane bound
GFP or stained with α-FasII, as indicated. (R) Quantification of aberrant axons in MBs (n
= 10). Scale is 10 μm in (A) and 20 μm in (F).

Figure 4

Figure 4. Ectopic EcR-B1 expression rescues memory defects rescued in Kis
knockdown MBs. (A, C) Change in courtship index of *OK107*-Gal-4 driven males
during the initial (light grey) and final (dark grey) 10 minutes was assessed (n = 20-25).
(B, D) Immediate recall of trained *OK107*-Gal-4 driven males was assessed and
compared to genetically identical sham trained males (n = 20-25).
392

Figure 5.

Figure 5. HDAC inhibition rescues defective axon pruning and EcR-B1 expression
in loss of Kis animals. (A-D) Representative images of DMSO or SAHA treated
MARCM generated MB clones expressing GFP using the *201Y*-Gal-4 driver 18-22 hours
APF. (E) Quantification of dorsal, medial, and total MB lobe surface areas in DMSO or
SAHA treated MARCM animals (n = 10-12). (F-Q) Representative DMSO or SAHA
treated adult MARCM generated MB clones expressing membrane bound GFP or stained
with α-FasII, as indicated. (R) Quantification of aberrant axons in MBs of flies treated
with DMSO or SAHA (n = 10-12). (S) *EcR-B1* mRNA levels in brains of DMSO or
SAHA treated *elav*-Gal-4 driven pupae analyzed by RT-qPCR (n = 3-4). Scale is 10 μm
in (A) and 20 μm in (F).

Figure 6.

Figure 6. SAHA treatment rescues immediate recall defects rescued by in Kis
knockdown animals. (A, C) Change in courtship index of DMSO or SAHA treated
OK107-Gal-4 driven males during the initial (light grey) and final (dark grey) 10 minutes
was assessed (n = 20-25). (B, D) Immediate recall of DMSO or SAHA treated trained
OK107-Gal-4 driven males assessed and compared to genetically identical and treated
sham trained males (n = 20-25).
422
423

431   **Supplemental Figure 1**

433   **Supplemental Figure 1. Kis is required for axon pruning in Kis knockdown pupae.**
434   Quantification of dorsal, medial, and total MB lobe surface areas in *elav*-Gal-4, mCD8-
435   GFP driven pupal brains stained with α-EcR-B1 (n = 10).
436

Supplemental Figure 2.

Supplemental Figure 2. Kis-eGFP is knocked down by Kis.RNAi.a. (A-F)
Representative images of Kis-eGFP and *elav*-Gal-4 driven knockdown 3rd instar larval
brains stained with DAPI. (G) Quantification of GFP fluorescence intensity compared to
that of DAPI (n = 9-12). (H) *Kis* mRNA levels in 3rd instar larval brains of *elav*-Gal-4
driven control and knockdown analyzed by RT-qPCR (n = 4).

Supplemental Figure 3.

Supplemental Figure 3. Kis does not alter DNA protection at *EcR* loci. Quantification
of MNase protection assay followed by qPCR at the *EcR* TSS, EcR enhancer sites, and
the *fkh* TSS (n = 3).
458

   Supplemental Figure 4.

Supplemental Figure 4. Kis does not alter H3K4 or H3K27 methylation at *EcR* loci.
(A-D) qPCR analysis H3K4me1, H3K4me2, H3K4me3, H3K27me3 abundance at the
EcR TSS and enhancer sites, the *fkh* TSS, and the *shi* promoter site, respectively (n = 2-
4).

 Supplemental Figure 5.

485 Supplemental Figure 5. **Exogenous EcR-B1 rescues defective axon pruning in Kis**
486 **knockdown pupae.** Quantification of dorsal, medial, and total MB lobe surface areas in
487 e*lav*-Gal-4, mCD8-GFP driven pupae (n = 10).
488

Methods

Drosophila stocks and genetics

Unless otherwise noted, all crosses were carried out at 25 °C in a 12:12 light:dark cycle at 60% humidity on standard cornmeal-molasses-agar medium. BL numbers refer to Bloomington Stock Center (BSC) stock numbers (http://flystocks.bio.indiana.edu/bloomhome.htm). VDRC numbers refer to the Vienna Drosophila Resource Center stock numbers (http://stockcenter.vdrc.at/control/main). To drive the expression of transgenes in Drosophila, the Gal-4/UAS bipartite system was used as previously described [41]. All Gal-4 stocks (*elav*-Gal-4 (BL #458), *elav*-Gal-4,mCD8-GFP (BL #5146), (*OK107*-Gal-4 (BL #44407)), w^{1118} (BL #5905), UAS:EcRB1 (BL #6469), and Canton S (wild-type) were obtained from Bloomington Stock Center. UAS:Kis.RNAi.a and UAS:Kis.RNAi.b (VDRC #10762 and #46685, respectively) came from the VDRC and were previously described [13]. The *kis*^{LM27} allele was generated by EMS mutagenesis, as previously described [42]. UAS:Kis-L, Kis-eGFP, *201Y*-Gal-4 MARCM stocks were gifts from J. Tamkun, A. Spralding, and L. Luo, respectively [20,21]. MARCM mosaic analysis clones were generated as previously described [16,43]. The EcR transcriptional reporter (*EcR.1*-Gal-4) was obtained as described [23,24].

Pharmacological treatment media was prepared as described [34]. Treated fly media was made using dried instant food (Nutri-Fly Instant, Genesee Scientific) with water containing 1.6% of 10% w/v tegosept (methyl p-hydroxybenzoate in 95% ethanol) and 0.1% of DMSO vehicle or 10μM SAHA. *Drosophila* were raised on drug containing food for their entire lifespan.

Immunohistochemistry

Immunohistochemical staining was carried out as previously described [20].

Dissections were performed on Sylgard-coated plates in phosphate buffer and fixed in 2%

paraformaldehyde. Primary antibodies obtained from the Iowa Developmental Hybridoma

Bank included α-EcR-B1 (1:200), α-Gal-4 (1:200), and α-FasII (1:200). Fluorescently

conjugated goat α-rabbit or goat α-mouse secondary antibodies (1:100, Jackson

Immunoresearch Labs). Brains were mounted in Vectashield with DAPI (Vector

Laboratories, H-1000) and images were obtained using an Olympus Fluoview 1000 laser

scanning confocal microscope.

Quantitative RT-PCR

Pupal (approximately 18 hours APF) heads or third instar larval brains were

dissected in ice-cold phosphate buffer and immediately transferred to RNA Later (Abion)

and stored in −80 °C. Isolation of total RNA was done using phenol:chloroform extraction

followed by alcohol precipitation for purification. RNA was stored in DEPC water at −80

°C. An adapted version of iTaq^{TM} Universal SYBR® Green One-Step protocol (Bio-Rad)

was utilized and samples were run on Bio-Rad C1000 Thermal Cycler CFX96 Real-Time

system. Primers were made to *Kismet, EcR-B1, Gal-4* mRNAs (IDT). ΔC(t) values were

calculated by subtracting the C(t) value of each primer set from C(t) value of *RP49*

housekeeping control. Fold change in expression was calculated from ΔΔC(t) values. Each

experiment was performed in triplicate with at least three biological replicates.

Chromatin Immunoprecipitation

Brains from third instar larvae were isolated in ice-cold phosphate buffer. Brains

were transferred to 1X phosphate buffered saline (PBS, Hyclone) and stored at −80 °C. A

modified version of truChIP Tissue Chromatin Shearing Kit with SDS Shearing Buffer

protocol (Covaris) was used to shear the DNA. Heads were washed twice with 1X PBS

and then fixed in Buffer A with 1% methanol-free formaldehyde for 5 minutes at room

temperature. Fixing was stopped with Quenching Buffer E followed by incubation for 5

minutes at room temperature. Tissue was pelleted by centrifugation at 4 °C for 5 minutes.

Supernatant was removed, and tissue was washed twice with cold 1X PBS. Wash buffer

(WB) was removed and tissue was homogenized for 2-3 minutes in 500µL Lysis Buffer

(LB) B. Volume was increased to 1mL with LB B followed by incubation on rocker at 4

°C for 20 minutes with 3 second vortex every 10 minutes. Lysed tissue was pelleted and

resuspened in WB C. Tissue was washed on rocker for 10 minutes at 4°C at which time it

was pelleted, followed by an additional washing with WB C without incubation. Pelleted

lysed and washed tissue, largely consisting of nuclei, was resuspended in Covaris SDS

Shearing Buffer D. The aggregate of nuclei was incubated with Buffer D for 10 minutes

with occasional vortex prior to transfer to a TC 12X12 tube for shearing. Shearing followed

the S- and E-Series Shearing recommendations for 15 minutes. Aliquots were stored at −

80 °C.

Sheared DNA was confirmed to be within a target range of 100-600 bp fragments.

Chromatin was immunoprecipitated using Magna ChIP™ HiSens kit (Millipore).

Antibodies against modifications H3K27me3 (rabbit, ab195477), H3K4me1 (rabbit,

ab8895), H3K4me2 (rabbit, ab7766), H3K4me3 (rabbit, ab8580), H3K36me2 (rabbit,

ab9049), H3K36me3 (rabbit, ab9050) were used and compared to Histone H3 antibody

(rabbit, ab1791). To examine Kismet abundance, α-GFP (rabbit, ab290) was used and α-

IgG (rabbit, ab171870) was used as a background control. After elution, samples were

incubated with RNaseA (10mg/mL, ThermoScientific) at 37 °C for 30 minutes followed

by an incubation with proteinase K (10mg/mL, Millipore) at 57 °C overnight and then

inactivate at 75 °C for 15 minutes the next day. Isolated DNA was purified via QIAquick®

PCR Purification Kit (Qiagen) and stored at –20 °C.

MNase Protection Assay

Brains from third instar larvae were dissected in ice-cold phosphate buffer,

transferred to 1X PBS, and stored at –80 °C. An MNase protection assay was performed

using an adapted protocol from [44,45]. Tissue was homogenized in 500μL of crosslinking

buffer (60mM KCl, 15mM NaCl, 4mM MgCl2,15mM HEPES pH 7.6, 0.5mM DTT, 0.5%

Triton X-100, protease inhibitor (100X), 2% formaldehyde) and incubated at room

temperature for 15 minutes. Crosslinking was quenched with 50μL of 2.5M of glycine and

incubated at room temperature for 5 minutes. Samples were washed twice in crosslinking

buffer and twice in D1 buffer (25% glycerol, 5mM Mg Acetate, 50mM Tris pH 8.0, 0.1mM

EDTA, 5mM DTT) and resuspended in 1mL of MNase buffer (60mM KCl, 15mM NaCl,

15mM Tris pH 7.4, 0.5mM DTT, 0.25M sucrose, 1.0mM CaCl2). 10units MNase (70196Y,

Affymetrix) was added to sample tubes and incubated at 37 °C for 30 minutes. Reaction

was quenched with EDTA (final 12.5mM) and SDS (final 0.5%). Samples were

equilibrated with NaCl (final 140mM) and incubated with RNaseA (10mg/mL) at 37 °C

and then overnight with proteinase K (10mg/mL) at 65 °C and then 15 minutes at 75 °C

the next day. DNA was purified via QIAquick PCR Purification Kit and stored at –20 °C.

Q-PCR

Purified DNA was used to prep PCR reaction mixes according to DyNAmo Flash

SYBR Green qPCR Kit (ThermoFisher Scientific). Samples were run using Bio-Rad

C1000 Thermal Cycler CFX96 Real-Time system. Primers were made to the TSS of *EcR-*

B1, a 16kb upstream enhancer of *EcR-B1* (*EcR.1*), and a 7.4kb upstream enhancer of *EcR-*

B1 (*EcR.2*) (IDT). Additionally, positive and negative control primers were made to the

TSS of *fkh* and the *shi* promoter site, respectively (IDT). For control and RNAi knockdown

analysis values were adjusted for input at each primer set and $\Delta\Delta C(t)$ values were

calculated by subtracting the adjusted $\Delta C(t)$ value of each primer set from the

corresponding $\Delta C(t)$ IgG control. Fold change in expression was calculated from $\Delta\Delta C(t)$

values. Each experiment was performed in triplicate with at least three biological

replicates.

Behavioral testing

To evaluate learning and memory, the canonical fly courtship behavior was used as

a readout in an associative conditioning assay described by Siegel and Hall [32]. Virgin male

flies (0 to 6 hours following eclosion) were collected in individual food vials and aged 5

days. Similarly, virgin female wild-type flies were collected, transferred to collective food

vials, and aged 5 days. 24 hours before assessment, virgin wild-type females were mated

individually using wild-type males. These flies were subsequently separated from virgin

females. This behavioral test was executed in a separate room kept at 25°C and 50%

humidity, recorded using a Sony DCR-SR47 Handycam with Carl Zeiss optics, and

illuminated from below using a constant 115V white light transilluminator. Genotypes of

each male were blinded on the day of the assay and all fly transfers were performed without

anesthesia. Aged male flies were transferred to mating chambers (Aktogen) each

containing a portioned-off mated female fly. Flies were allowed to acclimate for 2 minutes

before the assay. Training was recorded and commenced for 60 minutes. After, the male

fly was transferred to a clean mating chamber containing a portioned-off virgin female fly.

After an additional 2-minute acclimation period, the divider was removed, and immediate

recall was recorded for 10 minutes. Shams experienced the same manipulations however

these aged males were not exposed to any fly during the training portion. Digital video

analysis of the time spent courting was performed using iMovie software (Apple).

Courtship indices were calculated by total time observed performing courtship behaviors

divided by total time assayed.

<u>Statistical analysis</u>

All statistical analyses were performed using GraphPad Prism (v. 7.03).

Significance was determined at the 95% confidence interval. Unpaired student's t-test was

used for all experiments, except pupal pruning analysis (utilized one-way ANOVA test)

and the learning portion of the associative conditioning assay (utilized paired student's t-

test). Statistical significance in figures is represented by * = $p < 0.05$, ** = $p < 0.001$, ***

= $p < 0.0001$. Error bars represent the standard error of the mean (SEM).

633
634
635

Acknowledgements

We would like to thank the members of the Marenda lab for their helpful feedback during the course of this project, the Bloomington Drosophila Stock Center, the Vienna Drosophila Stock Center, and the Developmental Studies Hybridoma Bank for fly stocks, antibodies, and reagents; Drexel CIC for imaging analysis and assistance. We would like to thank J. Tamkun, L. Luo, and A. Spralding for stocks. This work was supported by grants from the CHARGE syndrome Foundation (to D.R.M.) and the National Science Foundation IOS 1256114 (to D.R.M.). Some of this material is based upon work supported by (while serving at) the National Science Foundation. Any opinion, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

References

1   Tau, G. Z. & Peterson, B. S. Normal development of brain circuits.
Neuropsychopharmacology **35**, 147-168, doi:10.1038/npp.2009.115 (2010).
2   Low, L. K. & Cheng, H. J. Axon pruning: an essential step underlying the
developmental plasticity of neuronal connections. *Philos Trans R Soc Lond B Biol*
Sci **361**, 1531-1544, doi:463621T5G6131557 [pii]
10.1098/rstb.2006.1883 (2006).
3   Awasaki, T., Huang, Y., O'Connor, M. B. & Lee, T. Glia instruct developmental
neuronal remodeling through TGF-beta signaling. *Nat Neurosci* **14**, 821-823,
doi:10.1038/nn.2833 (2011).
4   Yu, F. & Schuldiner, O. Axon and dendrite pruning in Drosophila. *Curr Opin*
Neurobiol **27**, 192-198, doi:10.1016/j.conb.2014.04.005 (2014).
5   Truman, J. W. Metamorphosis of the central nervous system of Drosophila. *J*
Neurobiol **21**, 1072-1084, doi:10.1002/neu.480210711 (1990).
6   Levine, R. B., Morton, D. B. & Restifo, L. L. Remodeling of the insect nervous
system. *Curr Opin Neurobiol* **5**, 28-35 (1995).
7   Heisenberg, M., Borst, A., Wagner, S. & Byers, D. Drosophila mushroom body
mutants are deficient in olfactory learning. *J Neurogenet* **2**, 1-30 (1985).
8   Connolly, J. B. *et al.* Associative learning disrupted by impaired Gs signaling in
Drosophila mushroom bodies. *Science* **274**, 2104-2107 (1996).
9   Ferveur, J. F., Stortkuhl, K. F., Stocker, R. F. & Greenspan, R. J. Genetic
feminization of brain structures and changed sexual orientation in male Drosophila.
Science **267**, 902-905 (1995).
10   McBride, S. M. *et al.* Mushroom body ablation impairs short-term memory and
long-term memory of courtship conditioning in Drosophila melanogaster. *Neuron*
24, 967-977, doi:S0896-6273(00)81043-0 [pii] (1999).
11   Lee, T., Marticke, S., Sung, C., Robinow, S. & Luo, L. Cell-autonomous
requirement of the USP/EcR-B ecdysone receptor for mushroom body neuronal
remodeling in Drosophila. *Neuron* **28**, 807-818, doi:S0896-6273(00)00155-0 [pii]
(2000).
12   Zheng, X. *et al.* TGF-beta signaling activates steroid hormone receptor expression
during neuronal remodeling in the Drosophila brain. *Cell* **112**, 303-315,
doi:S0092867403000722 [pii] (2003).
13   Melicharek, D. J., Ramirez, L. C., Singh, S., Thompson, R. & Marenda, D. R.
Kismet/CHD7 regulates axon morphology, memory and locomotion in a
Drosophila model of CHARGE syndrome. *Hum Mol Genet* **19**, 4253-4264,
doi:ddq348 [pii]
10.1093/hmg/ddq348 (2010).
14   Layman, W. S., Hurd, E. A. & Martin, D. M. Chromodomain proteins in
development: lessons from CHARGE syndrome. *Clin Genet* **78**, 11-20,
doi:CGE1446 [pii]
10.1111/j.1399-0004.2010.01446.x (2010).
15   Vissers, L. E. *et al.* Mutations in a new member of the chromodomain gene family
cause CHARGE syndrome. *Nat. Genet.* **36**, 955-957, doi:10.1038/ng1407
ng1407 [pii] (2004).

16   Lee, T. & Luo, L. Mosaic analysis with a repressible cell marker for studies of gene
function in neuronal morphogenesis. *Neuron* **22**, 451-461 (1999).
17   Schuldiner, O. *et al.* piggyBac-based mosaic screen identifies a postmitotic
function for cohesin in regulating developmental axon pruning. *Dev Cell* **14**, 227-
238, doi:S1534-5807(07)00419-4 [pii]
10.1016/j.devcel.2007.11.001 (2008).
18   Yang, M. Y., Armstrong, J. D., Vilinsky, I., Strausfeld, N. J. & Kaiser, K.
Subdivision of the Drosophila mushroom bodies by enhancer-trap expression
patterns. *Neuron* **15**, 45-54, doi:0896-6273(95)90063-2 [pii] (1995).
19   Boulanger, A. *et al.* ftz-f1 and Hr39 opposing roles on EcR expression during
Drosophila mushroom body neuron remodeling. *Nat Neurosci* **14**, 37-44,
doi:nn.2700 [pii]
10.1038/nn.2700 (2011).
20   Ghosh, R. *et al.* Kismet positively regulates glutamate receptor localization and
synaptic transmission at the Drosophila neuromuscular junction. *PLoS One* **9**,
e113494, doi:10.1371/journal.pone.0113494 (2014).
21   Buszczak, M. *et al.* The carnegie protein trap library: a versatile tool for Drosophila
developmental   studies.   *Genetics*   **175**,   1505-1531,
doi:10.1534/genetics.106.065961 (2007).
22   Srinivasan, S., Dorighi, K. M. & Tamkun, J. W. Drosophila Kismet regulates
histone H3 lysine 27 methylation and early elongation by RNA polymerase II. *PLoS*
Genet. **4**, e1000217, doi:10.1371/journal.pgen.1000217 (2008).
23   Pfeiffer, B. D. *et al.* Tools for neuroanatomy and neurogenetics in Drosophila.
Proceedings of the National Academy of Sciences of the United States of America
105, 9715-9720 (2008).
24   Pfeiffer, B. D. *et al.*   (2011).
25   Clapier, C. R., Iwasa, J., Cairns, B. R. & Peterson, C. L. Mechanisms of action and
regulation of ATP-dependent chromatin-remodelling complexes. *Nat Rev Mol Cell*
Biol **18**, 407-422, doi:10.1038/nrm.2017.26 (2017).
26   Dorighi, K. M. & Tamkun, J. W. The trithorax group proteins Kismet and ASH1
promote H3K36 dimethylation to counteract Polycomb group repression in
Drosophila. *Development* **140**, 4182-4192, doi:10.1242/dev.095786 (2013).
27   Stabell, M., Larsson, J., Aalen, R. B. & Lambertsson, A. Drosophila dSet2
functions in H3-K36 methylation and is required for development. *Biochem*
Biophys Res Commun **359**, 784-789, doi:10.1016/j.bbrc.2007.05.189 (2007).
28   Wagner, E. J. & Carpenter, P. B. Understanding the language of Lys36 methylation
at histone H3. *Nat Rev Mol Cell Biol* **13**, 115-126, doi:10.1038/nrm3274 (2012).
29   Bornstein, B. *et al.* Developmental Axon Pruning Requires Destabilization of Cell
Adhesion   by   JNK   Signaling.   *Neuron*   **88**,   926-940,
doi:10.1016/j.neuron.2015.10.023 (2015).
30   Stewart, B. A. & McLean, J. R. Population density regulates Drosophila synaptic
morphology in a Fasciclin-II-dependent manner. *J Neurobiol* **61**, 392-399,
doi:10.1002/neu.20096 (2004).
31   McBride, S. M. *et al.* Pharmacological rescue of synaptic plasticity, courtship
behavior, and mushroom body defects in a Drosophila model of fragile X
syndrome. *Neuron* **45**, 753-764, doi:S0896-6273(05)00076-0 [pii]

10.1016/j.neuron.2005.01.038 (2005).

32   Siegel, R. W. & Hall, J. C. Conditioned responses in courtship behavior of normal
and mutant Drosophila. *Proc Natl Acad Sci U S A* **76**, 3430-3434 (1979).

33   Siwicki, K. K. *et al.* The role of cuticular pheromones in courtship conditioning of
Drosophila males. *Learn Mem* **12**, 636-645, doi:lm.85605 [pii]

10.1101/lm.85605 (2005).

34   Latcheva, N. K. *et al.* Epigenetic crosstalk: Pharmacological inhibition of HDACs
can rescue defective synaptic morphology and neurotransmission phenotypes
associated with loss of the chromatin reader Kismet. *Mol Cell Neurosci* **87**, 77-85,
doi:10.1016/j.mcn.2017.11.007 (2018).

35   Srinivasan, S., Dorighi, K. M. & Tamkun, J. W. Drosophila Kismet regulates
histone H3 lysine 27 methylation and early elongation by RNA polymerase II. *PLoS*
Genet **4**, e1000217, doi:10.1371/journal.pgen.1000217 (2008).

36   Srinivasan, S. *et al.* The Drosophila trithorax group protein Kismet facilitates an
early step in transcriptional elongation by RNA Polymerase II. *Development* **132**,
1623-1635, doi:10.1242/dev.01713 (2005).

37   Bell, O. *et al.* Localized H3K36 methylation states define histone H4K16
acetylation during transcriptional elongation in Drosophila. *EMBO J* **26**, 4974-
4984, doi:10.1038/sj.emboj.7601926 (2007).

38   Shogren-Knaak, M. *et al.* Histone H4-K16 acetylation controls chromatin structure
and protein interactions. *Science* **311**, 844-847, doi:10.1126/science.1124000
(2006).

39   Zhang, R., Erler, J. & Langowski, J. Histone Acetylation Regulates Chromatin
Accessibility: Role of H4K16 in Inter-nucleosome Interaction. *Biophys J* **112**, 450-
459, doi:10.1016/j.bpj.2016.11.015 (2017).

40   Barbetti, V. *et al.* Time- and residue-specific differences in histone acetylation
induced by VPA and SAHA in AML1/ETO-positive leukemia cells. *Epigenetics* **8**,
210-219, doi:10.4161/epi.23538 (2013).

41   Brand, A. H. & Perrimon, N. Targeted gene expression as a means of altering cell
fates and generating dominant phenotypes. *Development* **118**, 401-415 (1993).

42   Melicharek, D. *et al.* Identification of Novel Regulators of Atonal Expression in the
Developing Drosophila Retina. *Genetics* **180**, 2095–2110,
doi:10.1534/genetics.108.093302 (2008).

43   Wu, J. S. & Luo, L. A protocol for mosaic analysis with a repressible cell marker
(MARCM) in Drosophila. *Nature protocols* **1**, 2583-2589 (2006).

44   Berson, A. *et al.* TDP-43 Promotes Neurodegeneration by Impairing Chromatin
Remodeling. *Curr Biol* **27**, 3579-3590 e3576, doi:10.1016/j.cub.2017.10.024
(2017).

45   Chereji, R. V. *et al.* Genome-wide profiling of nucleosome sensitivity and
chromatin accessibility in Drosophila melanogaster. *Nucleic Acids Res* **44**, 1036-
1051, doi:10.1093/nar/gkv978 (2016).