Temporal scaling of ageing as an adaptive strategy of *Escherichia coli*

Yifan Yang[1,2,*], Ana L. Santos[1], Luping Xu[1,2,3], Chantal Lotton[1], François Taddei[1,2] and Ariel B. Lindner[1,2,*]

[1] INSERM U1001, Faculté de Médecine, Université Paris Descartes, Paris, France

[2] Center for Research and Interdisciplinarity, Paris, France

[3] Current address: Center for Nano and Micro Mechanics, School of Aerospace Engineering, Tsinghua University, Beijing, China

[*] Correspondence: yifan.yang@inserm.fr and ariel.lindner@inserm.fr

Abstract

Natural selection has long been hypothesised to shape ageing patterns, but whether and how ageing contributes to life-history evolution remains elusive. The complexity of various ageing-associated molecular mechanisms and their inherent stochasticity hinder reductionist approaches to the understanding of functional senescence, *i.e.* reduced fecundity and increased mortality. Recent bio-demographic work demonstrated that high-precision statistics of life-history traits such as mortality rates could be used phenomenologically to understand the ageing process. We adopted this approach to study cellular senescence in growth-arrested *E. coli* cells, where damages to functional macromolecules are no longer diluted by fast *de novo* biosynthesis. We acquired high-quality longitudinal physiological and life history data of large environmentally controlled clonal *E. coli* populations at single-cell resolution, using custom-designed microfluidic devices coupled to time-lapse microscopy. We show that *E. coli* lifespan distributions follow the Gompertz law of mortality, a century-old actuarial observation of human populations, despite developmental, cellular and genetic differences between bacteria and metazoan organisms. Measuring the shape of the hazard functions allowed us to disentangle quantitatively the demographic effects of ageing, which accumulate with time, from age-independent genetic longevity-modulating interventions. A

pathway controlling cellular maintenance, the general stress response, not only promotes

longevity but also temporally scales the whole distribution by reducing ageing rate. We

further show that *E. coli*, constrained by the amount of total biosynthesis, adapt to their

natural feast-or-famine lifestyle by modulating the amount of maintenance investment,

rendering ageing rate a highly evolvable life-history trait.

Background

The biology of ageing and senescence is centred on the duality of individual frailty

and population resiliency. Throughout the course of normal metabolism, components of

living systems such as cells, lipids, proteins and DNA inevitably suffer from wear-and-tear

such as free-radical damages. This constant and collective decay eventually leads to the loss

of vital functions and collapse of individuals. Understanding the way that system failure

emerges out of distributed microscopic damages could reveal how functional components are

organised into self-maintaining individuals in the first place[1,2]. Yet for some organisms in the

tree of life, ageing does not lead to increased mortality and declined fertility[3]. Organisms

possess the abilities to repair or replace most of the damages to their components,

exemplified by the "immortal germ line"[4]. Evolutionary biologists attribute the apparent

senescence of metazoan somas to the inadequate investment in cellular maintenance as an

adaptive strategy to maximise lifetime reproductive success, due to the trade-offs between the

survival and reproduction of the young on one hand, and maintenance for the benefit of the

old on the other[5,6]. Presumably, natural selection has to operate through the molecular

"levers" of damage accumulation and/or repair to achieve such life-history optimisation.

Escherichia coli, a single-cell prokaryote with short lifespans, has historically served

as a model organism that resolved many fundamental questions in biology. *E.coli* cells, as

their metazoan counterparts, suffer from damages to their components, which lead to cellular

senescence[7,8]. In exponential growth, these damages are quickly diluted by *de novo*

biosynthesis, and the effects of cellular senescence mitigated by rapid and robust

reproduction[9]. Yet, the natural life cycle of *E.coli* entails a much wider range of physiological

conditions than exponential growth. Most of bacterial cells spend much of their lives in

resource-limited growth arrested conditions, where *de novo* biosynthesis rates are slower[10]

and cells undergo senescence due to the accumulation of molecular damages such as protein

misfolding and oxidation (Fig. 1)[11]. Despite the lack of fixed separation of germ and soma

cells, the ability to survive the wear and tear of cellular components during growth arrests

contribute to bacterial overall fitness as much as the ability for exponential growth. We,

therefore, adopted a bio-demographic approach[1,2] to understand how modulation of molecular

damage repair could shape ageing dynamics in growth-arrested *E.coli*.

64 Custom-designed microfluidic devices allow longitudinal tracking of vitality at

65 single-cell level

We first set out to acquire high quality longitudinal physiological and life-history data

at single-cell resolution of clonal populations of *E.coli*. For decades, limitations of traditional

cell culture methods have frustrated quantitative physiologists' attempts to understand

bacterial maintenance[12,13], due to both media-cell interactions and cellular interaction such as

cross-feeding and cannibalism. To be able to measure individual *E.coli* cell lifespan in

constant and homogenous environmental settings, we designed a novel microfluidic device

with cell-dimension chambers (Fig. 2). Cells are trapped in an array of single-cell-wide

dead-ended wells, with openings to a main flow channel. Constant flow of fresh media in this

main channel provides necessary nutrients and eliminates metabolic waste, cell debris and

intercellular crosstalk so that the environmental conditions are maintained constant over time.

Cells with appropriately expressed fluorescent markers are imaged bottom-up and appear as

fluorescent spots (Fig. 2b). This experimental system allows easy tracking of cohorts of a

large number of individual bacterial cells for prolonged periods (up to 7 days).

In order to observe cell mortality, we included in the carbon-source-free medium a

red-fluorescent, DNA-binding, bacterial viability dye, Propidium Iodide (P.I.), which

penetrates the cells only when cellular membrane potentials are disrupted. P.I. staining has

been established as an effective proxy of cellular death and correlates well with cell viability

assessed by proliferating potential[14]. We used P.I. at a concentration 4-fold lower than the

previous concentration that had no effects on *E.coli* viability and growth[14]. Automatic time-

lapse fluorescence microscopy and fixed geometry of our devices allowed longitudinal

quantification of P.I. signal for every single cell in the population. We defined the half point

between peak P.I. signal and background fluorescence as a threshold to establish the time-of-

death. At the end of the experiments, 70% ethanol was injected into the device in order to

account for every cell in the cohort, and establish a lower bound for the time-of-death of

surviving cells. These cells were censored at the time that their P.I. signals crossed their

respective thresholds (see Methods).

Despite being genetically clonal and environmentally controlled, cells did not share

the same time or manner of death, as measured by the P.I. time-series. The transition from

life to death of a representative cohort (N= 4744) could be visualised by the boundary

between dark to light in Fig. 2d. Because cells were sorted vertically according to estimated

lifespans, the shape of this boundary represents the survival function of the population.

Considering ageing and death as a stochastic process of system reliability reduction[1,2], the

observed time-of-death distribution could be viewed as the population's first-passage time

distribution. In addition, we observed individual differences in death trajectories. Short-lived

cells tended to have very sharp P.I. increases that associated with abrupt losses of membrane

integrity; while those that die late tended to suffer a type of "slow death" characterised by a

gradual P.I. increase over the course of 10-15 hours.

***E. coli* lifespan distributions follow the Gompertz law**

For both demographers and reliability engineers, the age-associated increase in death

probability, also known as the hazard rate $h(t)$, is considered to be a hallmark of ageing[1-3].

The detailed population statistics derived from our experiments are particularly useful in

estimating the shape of the hazard functions over the whole lifespan. We find that *E.coli*

lifespan distributions have, as their main feature, regimes with exponentially increasing

hazard rates, *i.e.* the Gompertz law of mortality *$h(t) \sim h(t_0) \, e^{b(t-t_0)}$*, where *$h(t_0)$* is the initial

hazard rate and *b* is the Gompertz ageing rate[15,16]. We estimated *$h(t)$* directly with binomial

error by binning mortality events within discrete time intervals (τ_i, τ_{i+1}), *i.e.* $\tilde{h}_i \sim d_i/Y_i/\Delta\tau_i$,

where Y_i is the number of individuals at risk at time τ_i and d_i is the number of cell deaths

within (τ_i, τ_{i+1}) (see Fig. 3). For wildtype cells, this exponential regime spans from about 13

hours to 93 hours, corresponding to approximately 90% of all cell deaths (N_{total}=4744, see

Fig. 3b, top x-axis), and ranges at least 100-fold changes in hazard rates ($2*10^{-3}$ h^{-1} to $2*10^{-1}$

h^{-1}). Within this exponential regime, the doubling time of hazard rate is 9.4±0.5 hours. In

comparison, the doubling time of human mortality hazards is about 8 years[15].

Hazard rates not only define lifespan distributions but can also be thought of as

surrogates for system vulnerability, whose dynamics through time reflect the physiological

consequence of ageing. In this light, the Gompertz law can be interpreted as a dynamic

equation governing the ageing process: *$dh(t)/dt = bh(t)$*, where *b* is the Gompertz ageing rate.

The integral version, *$h(t) = bH(t) + h(t_0)$*, where $H(t) = \int_0^t h(t^{'})dt^{'}$, can be directly observed in

the phase plane of the cumulative hazards *$H(t)$* without any free parameters (Fig. 3b). The

integral equation is used because *H(t)* can be estimated independent of binning, by using the

Nelson-Aalen (N-A) estimator $\tilde{H}_{NA}(\tau_i)$[17]. The fact that the theoretical trajectory, including

the aforementioned linear regime, lies within the confidence intervals (CI) of almost every

state coordinate ($\tilde{H}_{NA}(\tau_i)$, $\tilde{h}_i$) lends strong statistical credence to the Gompertz law for

describing our data. In addition, the observed trajectory in the phase plane links the

instantaneous hazard rate *h(t)* at any given time *t*, with the total proportion of population that

have died before time *t* (because *H(t) = -ln[S(t)]*, where *S(t)* is the survival function).

The hazard rate observed deviates from the exponential regime at the beginning (t<15

hours, 3% cell deaths) and the end (t>93 hours, 7% cell deaths) of the total lifespan. These

observations are reminiscent of similar deviations from Gompertz law in human mortality

data[18,19]. In our case, the additional mortality at the early age might result from harvesting

and transferring exponentially growing cells from batch culture directly to growth-arresting

conditions inside our microfluidic chip, in a way analogous to infant mortality in human

mortality data, which can be modelled by an age-independent component (λ in the Gompertz-

Makeham model $h_{gm}(t) = \lambda + \frac{b}{\beta}e^{bt}$, where β controls the age at which the Gompertz regime

overtakes the Makeham term λ).

Late-life hazard decelerations are common phenomena for both model organisms and

human populations[18,20]. It is thought that these hazard decelerations often do not reflect

ageing decelerations for individuals, but result from changes in composition of heterogeneous

populations. The more fragile individuals are more likely to die and be removed from the

cohort. When hazard rates are high late in life, significant portions of the populations are

removed so that the observed hazard rates of the surviving populations would be smaller than

the original cohorts on average. We found that an extension to the Gompertz law named

Gamma-Gompertz, originally used to model old-age human model data[21], could also

satisfactorily model late life decelerations in our data, for both wildtype and mutant strains

(see below). It could be understood as accounting for the compositional changes by an

additional parameter, *s*, controlling the level of frailty heterogeneity:

$h_{gg}(t) = \dfrac{b}{\beta} e^{bt} \cdot s[S_{gg}(t)]^{1/s}$, where $S_{gg}(t)$ is the survival function of Gamma-Gompertz, and

other parameters are as before. Despite being genetically and environmentally non-

distinguishable, we think it is reasonable to assume frailty heterogeneity exist in our single-

cell populations due to the stochastic nature of cellular biochemistry, explaining the

suitability of the Gamma-Gompertz for our data.

We thus used the Gamma-Gompertz-Makeham (GGM) extension model with 4

parameters to fully model our data across the whole lifespan, with parameters estimated by

likelihood maximisation:

$$h_{ggm}(t) = \lambda + \dfrac{bs}{1 + (\beta - 1)e^{-bt}}$$

Goodness-of-fit was evaluated using a one-sample Kolmogorov-Smirnov test for

right-censored survival time data. The appropriateness of the model terms was evaluated by

comparing Akaike Information Criterion (AIC) of the candidate models (see methods).

165   The general stress response of *E. coli* modulates the ageing rate

Evolutionary theories of ageing predict the existence of ageing-modulating

mechanisms and their activation by nutrient limitation[22]. Several nutrient-sensing pathways in

both bacteria and metazoans have been shown to control stress resistance and reduce

mortality[23-25]. Yet their roles in delaying ageing-related damage accumulations are often

controversial. It is often difficult and requires large experimental cohorts to disentangle their

effects on ageing from age-independent components of longevity[26]. Having established a

method to measure ageing rates using high-quality mortality statistics, we harnessed this

system to shed light on potential ageing-modulating mechanisms of *E.coli* that might be

activated in response to nutrient deprivation.

Many bacteria species regulate the level of cellular maintenance through a genetic

pathway called general stress response (GSR), controlled by the master transcriptional

regulator *rpoS* that is activated by nutrient-deprivation among other signals[27] (Fig. 4a). To

assess the role of *rpoS*-controlled cellular maintenance in the ageing of growth-arrested

E.coli, we measured the lifespan distributions of two GSR mutants, $\Delta rpoS$, and $\Delta rssB$ with

that of the wildtype strain. $\Delta rpoS$ is the null mutant and $\Delta rssB$ displays an elevated GSR due

to increased RpoS stability[28]. We observed that higher GSR promotes longevity in the

microfluidic experiments, whereas the absence of GSR results in shortened longevity (Fig. 4b

and Fig. 5a). Our large sample sizes allowed us to directly measure the hazard dynamics (Fig.

4c) of each strain, which could disentangle GSR's effects on ageing rate, as opposed to age-

independent components of longevity.

Significantly, we found that increased GSR reduces the rate of ageing (Fig. 4d and

Fig. 5c). Given how well the GGM model fitted for all 3 strains, the impact of genotypes on

ageing parameters such as ageing rates could be extracted using generalised linear models

(GLM). With enhanced GSR, $\Delta rssB$ cells double their mortality risk every 14.1 hours, with a

95% CI ranging from 13.1 to 15.3 hours, compared to 9.4 hours for the wildtype (CI 8.9 - 9.9

hours) and 7.3 hours for the null strain $\Delta rpoS$ (CI 6.8 to 7.7 hours). Variability of ageing rate

measurements was assessed using 3 independent experimental cohorts for each strain (Figure

S1). We visualised non-parametrically the overall experimental variations (Fig. 5a and b).

Parametric differences in ageing rates among experimental repeats were tested using GLM

models and AIC. We confirmed that experimental repeats shared similar ageing rates (H_0:

Same aging rate for experimental repeats. d.f. H_0 - H_1 = -2; ΔAIC_{rssB}=-2.56, ΔAIC_{wt}=-2.26,

ΔAIC_{rpoS}=-2.30; N_{wt}=6867, N_{rssB}=6969, N_{rpoS}=4793).

The systematic increase in vulnerability of *E.coli* in our experiments is likely driven

by the catabolism of pre-existing macromolecules and dissimilation of biomass[29,30], which is

necessary to provide energy to express housekeeping genes and maintain physiological

homeostasis[10,31]. Our finding that GSR modulates ageing rate suggests that optimising this

maintenance energy requirement is likely one of the physiological functions of the RpoS

regulon (see Discussion).

Evolutionary trade-offs mediated by the general stress response

Is slower ageing an adaptive life-history trait? This was one of original question

raised by Sir Peter Medawar[32] that motivated much of later ageing research. Optimal life-

history theory suggested that trade-offs and constraints among fitness components shape

metazoan ageing rates[6]. The possibility of modulating ageing rate through GSR offers us the

opportunity to test these ideas in a fast-evolving organism as *E. coli*.

The relatively well-understood GSR pathway provides a clear molecular mechanism

for a trade-off between growth and maintenance. The master regulator *rpoS* encodes the RNA

polymerase (RNAP) sigma subunit σ^S, which competes with the other sigma factors

including the vegetative σ^D to recruit the core RNAP and direct the transcription and

translation machinery towards the RpoS regulon. By titrating protein synthesis activity away

from metabolic and ribosomal genes controlled by σ^D, RpoS activity inhibits growth and

nutrient assimilation[33]. We measured quantitatively the growth impact of modulating GSR

levels, and modelled its effect using a simple course-grained model of proteome sectors[34].

We found that the proportion of protein synthesis devoted to the RpoS regulon linearly

increases the timescale of growth (see top axis in Fig. 6a).

To assess the life-history optimality of different ageing rates, we integrated

numerically the experimentally derived growth rates and mortality rates into fitness, defined

as the long-term population growth rates. In contrast to metazoans, for whom fertility and

mortality schedules are connected through fixed age structures, our model of *E. coli* life-

history consisted of alternating environmental episodes of varying durations, in which *E. coli*

populations either grow ("feast"), or decline ("famine") (Fig. 6a). These feast-or-famine

cycles were parameterised by two transition rates, controlling respectively the average

lengths of "feast" and "famine" episodes. In order to identify selective pressure for ageing

rates, we directly compared the fitness of the three strains with different GSR phenotypes,

each representing a different strategic position in the growth-maintenance trade-off. We

identified the environmental regimes selecting for faster or slower ageing (Fig. 6b). The

boundaries between these regimes were characterised along two axes: lifestyle ratio, defined

as the ratio of time spent in "famine" versus "feast"; and "famine" mean residence time, or in

other words, average length of "famine" episodes.

There are two necessary conditions for selecting slower ageing strategies represented

by $\Delta rssB$. First, populations have to spend much more time in "famine" rather than "feast", so

that over the long term, population decline rather than grow. Secondly but no less important,

given the same lifestyle ratio, famines should consist of longer episodes rather than short but

more frequent ones. This condition is necessary due to the exponential mortality dynamics

described by Gompertz law: investing in cellular maintenance only becomes beneficial at old

age, when the exponentially increasing benefits of slower ageing eventually overcome the

more immediate cost on growth. It is the typical timescale of "famine" that provides the

selective pressure for ageing rates.

Despite representing the complex regulations of GSR with two mutants, we can now

understand the ecological role of GSR activation and its adaptive consequences. Activated by

declining nutrient availability, GSR directs resources towards internal maintenance to wait

out the adverse conditions, whose lengths determine the optimal activation level. Previous

observations from experimental evolution of *E. coli* support our predictions. In continuous

cultures of *E. coli* where the populations do not pass through prolonged growth-arresting

bottlenecks, mutations that attenuate or knock out RpoS activity are among the first to

arise[35,36]. In contrast to the isolated populations under constant environmental conditions in

our chip experiments, *E. coli* populations in nature influence their environments, and also

interact with each other. These interactions may give rise to frequency-dependent selection

and evolutionary game dynamics between slower and faster ageing strategies, as is observed

in experimental evolution[37,38].

Discussion

We obtained high quality single-cell demographic data, using a novel microfluidic

device, to demonstrate that ageing of growth-arrested *E. coli* follows the Gompertz law of

mortality. Moreover, bacterial general stress response could temporally rescale the lifespan

distribution by modulating the ageing rate. We further articulated in a demographic model the

trade-offs and selective pressure driving the evolution of ageing rate.

In our work, two different conceptual perspectives of ageing are integrated and

applied to one of the most iconic model organism, *E. coli*. One perspective, held by

biochemists and physicists, sees ageing as the stochastic and inevitable erosion of organismal

order and biochemical redundancy, created by self-reproducing and self-maintaining

networks during growth and development. The other perspective, from evolutionary biology,

views ageing as a component of the organismal life-history strategy, optimised and fine-tuned

by natural selection. These two approaches constitute the proximate and ultimate causes of

ageing respectively, with the former providing the constraints and the "lever" for the latter. In

bacteria, we indeed observed the stochastic process of ageing and mortality. Lifespan vary

significantly among genetically identical individuals in constant, homogeneous environment.

However, at the population level, ageing is characterised qualitatively by the Gompertz law

of mortality. In response to environmental conditions, the lifespan distribution is modulated

by regulatory and selective forces through temporal scaling, while its general, exponential,

shape is preserved. These new empirical observations provided an integrated perspectives to

bacterial ageing, and by extension, potentially to the breath of the tree of life.

Although the Gompertz law has been shown to characterise ageing of many metazoan

organisms, the physiological correlates of its parameters are not well understood. To rescale

the lifespan distribution without changing the general shape of hazard dynamics, GSR has to

orchestrate a coordinated response to manage various molecular damages that the organism

encounters[1]. Indeed, RpoS regulates hundreds of genes conferring resistance to both internal

and external stresses such as oxidative, thermal, acid, alkaline, osmotic, and UV[39]. For this

reason, the ageing rate in our case reflects the general level of macromolecular catabolism

and may relate to the maintenance energy of bacteria[13]. We hypothesise that the energetic

costs of homeostasis and molecular damage repair are paid by the loss of biomass, leading to

the gradual increase in the probability of death. It is the rate of energy dissipation, or the rate

of living, that correlates with the ageing rate in our system (see Fig. 7).

Despite the simplicity of bacteria and its vast differences from animals, we find that

concepts in evolutionary theories of ageing, such as antagonistic pleiotropy[5], still apply.

RpoS mediates trade-offs between growth and maintenance, between assimilation of nutrients

and dissimilation of biomass. Sigma competition, a well-understood mechanism in the

physiology of bacterial stress response, provides a molecular basis for such trade-offs often

only hypothesised in metazoan organisms. Activated by nutritional deprivation and shown to

decrease the ageing rate, GSR has immediate analogies in calorie-restriction-induced

longevity of metazoan model organisms. Whereas in many studies on animal models, calorie-

restriction-related pathways have only been shown to promote longevity, our demographic

data allow us to show that GSR indeed changes the rate of ageing. It can be thought of as not

only providing resistance against stress, but also insurance against prolonged growth arrests,

or in other words, protection against the progression of time.

In the future, the experimental approach and statistical framework described here

could combine with classical genetic methods to examine further ways hazard dynamics

could be perturbed, and to reveal dynamic features of the complex network shaping cellular

senescence under various environmental constraints. Future modelling efforts should clarify

how adaptive processes could optimise redundancy of living systems, in trade-offs with other

traits, to produce the Gompertz law.

Acknowledgements

This study was supported by the Axa Research Fund Chair on Longevity. We thank

X. Song for contributions on image analysis, D. Misevic and other member of INSERM

U1001 for their advice and discussions.

Author contributions

Conceived and designed project: Y.Y., L.X., F.T., A.B.L.; Conceived and designed

experiments: Y.Y., A.B.L.; performed experiments: Y.Y. A.S. C.L.; Data analysis: Y.Y.,

Wrote the paper: Y.Y. with contribution from A.B.L. .

Figure 1. The feast-and-famine life cycle of *E. coli*. Single-cell organisms such as *E. coli*

undergo developmental transitions not in response to fixed developmental programs but

opportunistically in response to environmental changes. Because long-term growth rates in a

stable ecosystem remain close to zero, life-history traits in "feast" and "famine" conditions

should contribute rather equally to overall fitness. In "feast" conditions, the traits under

strong selection are resource assimilation and reproduction, while in "famine" conditions are

maintenance and survival. Senescence has been shown to occur in "famine" conditions[11].

325
Figure 2. Coupling microfluidic chip and time-lapse microscopy to measure *E. coli* lifespan

distribution at single-cell resolution. **a**, 3D model of microfluidic devices used to trap and

isolate large number of single cells. The blue-gray cube in the upper half of the picture

represents the main flow channel where fresh, carbon-source-free media are supplied. The

array of light-blue rods represents *E.coli* single-cells trapped in a 2D-array of cell-sized dead-

end chambers. The arrows represent the media flow that maintain the environmental

homogeneity and removes debris. **b**, Fluorescence microscopy image of the microfluidic

device loaded with *E.coli* cells. Each fluorescent dot corresponds to a single cell trapped in

the 2D array of dead-end wells. Z-axis focus is adjusted so that dead-ends of the micro-wells

are imaged. The cyan pseudo-colour represents constitutively expressed fluorescent protein

fluorescence signal; red pseudo-colour - Propidium Iodide (P.I.) fluorescent signal described

in the text. **c**, Sample time-lapse P.I. images of mortality events, from early (top) to late

(bottom) deaths. **d**, Heat map of P.I. signal time-series for a population of single cells. The

colour values of each row correspond to the time-series of one cell, ranked according to

lifespan, top-to-bottom, from the shortest living to the longest living cell. The boundary

between dark and bright red indicates the transition from life to death and defines the lifespan

distribution. Dark strips after cell deaths indicate decaying DNA or empty wells after cell

debris have been washed away. Data shown in **d** have been smoothed according to the

procedure described in supplementary material to remove minor signal fluctuations coming

from microscopy focusing. The same procedure is used before time-of-death is estimated.

Figure 3. Non-parametric estimation indicates that wildtype *E. coli* lifespan distribution can

be characterised by the Gompertz law. Shown are non-parametric estimations of the hazard

functions and cumulative hazard functions of the same populations in one particular

microfluidic experiment. **a**, Binomial estimators of the hazard rates, shown in log scale. Cell

deaths are binned into discrete time intervals, which are marked by the x-axis error bars. The

time bins are chosen so that the Nelson-Aalen confidence intervals (CI) in b do not overlap

with each other. Error bars on the y-axis are the 95% CI for hazard rate estimated based on

binomial distribution. Yellow shading covers regions that deviate from the exponential

hazard regime for the shortest and longest living individuals. **b**, Cumulative hazard dynamics

in the phase plane. Data, binning intervals and y-axis are the same as those in **a**.

Instantaneous hazard rates plotted against the cumulative hazards (bottom x-axis), or

equivalently the negative logarithm of survivorship (top x-axis). Nelson-Aalen estimators are

used for the cumulative hazard function. The horizontal error bars are the 95% CI of the

Nelson-Aalen estimator at the centre of the time bins. Dashed lines are maximum likelihood

parametric estimations using the Gamma-Gompertz-Makeham model. The inset provides a

zoom-out view of the whole data range, while the main figure zooms in on the first 63% of

cell deaths.

Figure 4. The general stress response of *E. coli* modulates ageing rate. The lifespan

distribution for the wildtype (*wt*), Δ*rpoS* (lacking general stress response) and Δ*rssB*

(overexpressing the general stress response) strains are measured multiple times by

independent microfluidic experiments. **a**, A scheme representing relevant regulatory features

of the general stress response, and in particular the functions of the genes *rpoS* and *rssB*. **b**,

Experimental and GGM model survivorship. Representing the experimental survivorship,

colour bands are the 95% CI of the Kaplan-Meier (K-M) estimators. Coloured dashed lines

are Gamma-Gompertz-Makeham (GGM) models whose parameters are estimated from

maximum-likelihood (ML) methods. **c**, Hazard rates estimated using only cell deaths within

discrete time intervals (Error bar markers), and GGM hazard models estimated from the

whole dataset using ML methods. Similar to Fig. 3a, Vertical error bars are binomial 0.95

confidence intervals and horizontal error bars are the binning time intervals. Data from **b** and

c are from the same representative experimental cohorts for each strain. **d**, Ageing rates for

each strain, estimated by ML-fitted GGM models to 3 independent experiments. Error bars

represent 95% CI. The ageing rates are: $b_{rpoS} = 0.095 \pm 0.006$ h^{-1}, $b_{rssB} = 0.049 \pm 0.004$ h^{-1} and

$b_{wt} = 0.074 \pm 0.004$ h^{-1}.

Figure 5. Experimental repeatability and variances. For each of wildtype (*wt*), Δ*rpoS* and

Δ*rssB* knockout strains, data from 3 independent experimental replicates are shown. **a**,

Survivorship (Kaplan-Meier estimator, K-M) and cumulative hazards (Nelson-Aalen

estimator, N-A) of sub-populations from different imaging positions. Thick opaque lines

correspond to the data shown in Fig. 4b and c. Thin semi-transparent lines are the K-M and

N-A estimators of sub-populations from different imaging positions, which are used to

constitute the population represented by the opaque lines. Each sub-population is from one

imaging position that corresponds to a roughly 125μm-sided square-patch on the microfluidic

chip, and consists of 150-600 cells. Imaging positions with less than 50 cells are not shown.

b, Hierarchical clustering of the 3 independent experimental replicates of each strain.

Standard agglomerative clustering algorithm is applied using complete linkage and the 2-

sample modified Kolmogorov-Smirnov statistic sup$\{|Y(t)|\}$ (See methods) as a distance

metric. **c**, Ageing rates 3 independent experimental replicates of each strain. Ageing rates are

estimated similarly as those in Fig. 4d, but independently for each experimental replicate

without pooling them together using GLM. Error bars represent 95% CI.

Figure 6. Trade-offs between growth and maintenance mediated by the *rpoS* pathway, and its

fitness consequences. **a**, Scheme of the ecological processes (dashed arrows) and regulatory

relationships (solid arrows) involved in the trade-offs mediated by *rpoS*. Environments are

structured by alternating episodes of "feast" and "famine". Fitness of Δ*rpoS* (red), *wild-type*

(blue) and Δ*rssB* (green) as functions of time spent in each environmental episode (age) are

plotted in the top and bottom axes. Fitness is defined as the logarithmic change of population

sizes. **b**, Fitness comparison of the 3 strains across a range of environmental conditions to

identify regimes favouring faster and slower ageing strategies. The colour-coded regions

identify environmental conditions under which one strain dominates over the other two.

Absolute fitness of the 3 strains are converted to RGB tuples using softmax normalisation.

For each strain and each pair of environmental parameters, fitness is calculated by averaging

the population growth/decline rates over 5000 episodes of both "feast" and "famine". Episode

lengths are independently drawn from exponential distributions, parameterised by the two

ecological rates, k_1 and k_2, as shown in **a**. The fitness functions, population gain for "feast"

and survivorship for "famine" (shown in the axes in **a**), are used to determine population size

changes in each episode. See methods for details.

Figure 7. Gompertz law of mortality informs and constraints physiological models of ageing

and mortality in growth-arrested *E.coli* cells. Depicted are the abstract states and pathways

cells pass through during ageing and mortality. Acute damages to essential cellular

components, when unrepaired, often lead to cell death, quantified by *h(t)*. Cells could also

triage and repair these damages to return themselves back to physiological homeostasis. The

energy cost of this maintenance process, *J(t)*, termed maintenance energy[13], need to be repaid

by either an external energy source or internal storage in the form of existing biomass. If we

assume it is the same set of acute molecular damages that kill the cells or lead to ageing, then

J(t) could relate to *h(t)* proportionally, *J(t) = R h(t)* (eq1), where *R* is the "friction" coefficient

controlled by the relative proportions of the two fates of damaged cells: repair vs death. For

carbon-starved cells in our experiments, this flux of energy dissipation decreases the capacity

for molecular repair in the future, *U(t)*, leading to cellular senescence. Put it another way,

dU(t)/dt = -J(t) (eq2). Through these assumptions, the hazard dynamics we described in this

work could be recast as the dynamics of maintenance capacity. Our experimental results

place non-trivial constraints on how the amount of maintenance capacity regulates the rate at

which cells are damaged molecularly in the first place. If we only consider the Gompertz

regime, *i.e. dh(t)/dt = bh(t)* (eq3), combining with (eq1) (eq2) and differentiating on both

sides, we have $d^2U/dt^2 + b\,dU/dt = 0$ (eq3) and *dU(t)/dt = -R h(0)* , where *b* is the Gompertz

ageing rate and R balance out the choice of t0 which should not alter the dynamics of (eq3).

This framework could also be extended to growing populations by considering the equation

$d^2U/dt^2 + b\,dU/dt + c(U) = 0$, where c(*U*) is zero in low levels of *U* such as in growth arrest

conditions, but theoretically could be larger than zero, pushing the physiology away from

senescence but toward rejuvenation.

References

Stroustrup, N. *et al.* The temporal scaling of Caenorhabditis elegans ageing. *Nature* **530**, 103-+, doi:10.1038/nature16550 (2016).

Gavrilov, L. A. & Gavrilova, N. S. The reliability theory of aging and longevity. *J Theor Biol* **213**, 527-545, doi:10.1006/jtbi.2001.2430 (2001).

Jones, O. R. *et al.* Diversity of ageing across the tree of life. *Nature* **505**, 169-173, doi:10.1038/nature12789 (2014).

Kirkwood, T. B. & Holliday, R. The evolution of ageing and longevity. *Proc R Soc Lond B Biol Sci* **205**, 531-546 (1979).

Williams, G. C. Pleiotropy, Natural-Selection, and the Evolution of Senescence. *Evolution* **11**, 398-411, doi:Doi 10.2307/2406060 (1957).

Partridge, L. & Barton, N. H. Optimality, Mutation and the Evolution of Aging. *Nature* **362**, 305-311, doi:DOI 10.1038/362305a0 (1993).

Stewart, E. J., Madden, R., Paul, G. & Taddei, F. Aging and death in an organism that reproduces by morphologically symmetric division. *PLoS Biol* **3**, e45, doi:10.1371/journal.pbio.0030045 (2005).

Lindner, A. B., Madden, R., Demarez, A., Stewart, E. J. & Taddei, F. Asymmetric segregation of protein aggregates is associated with cellular aging and rejuvenation. *Proc Natl Acad Sci U S A* **105**, 3076-3081, doi:10.1073/pnas.0708931105 (2008).

Wang, P. *et al.* Robust growth of Escherichia coli. *Curr Biol* **20**, 1099-1103, doi:10.1016/j.cub.2010.04.045 (2010).

Gefen, O., Fridman, O., Ronin, I. & Balaban, N. Q. Direct observation of single stationary-phase bacteria reveals a surprisingly long period of constant protein production activity. *Proc Natl Acad Sci U S A* **111**, 556-561, doi:10.1073/pnas.1314114111 (2014).

Nystrom, T. Aging in bacteria. *Curr Opin Microbiol* **5**, 596-601 (2002).

Monod, J. *Recherches sur la croissance des cultures bactériennes*. (Hermann, 1942).

Pirt, S. J. The maintenance energy of bacteria in growing cultures. *Proc R Soc Lond B Biol Sci* **163**, 224-231 (1965).

Ericsson, M., Hanstorp, D., Hagberg, P., Enger, J. & Nystrom, T. Sorting out bacterial viability with optical tweezers. *J Bacteriol* **182**, 5551-5555 (2000).

Gompertz, B. On the Nature of the Function Expressive of the Law of Human Mortality, and on a New Mode of Determining the Value of Life Contingencies. *Philosophical Transactions of the Royal Society of London* **115**, 513-583 (1825).

Kirkwood, T. B. Deciphering death: a commentary on Gompertz (1825) 'On the nature of the function expressive of the law of human mortality, and on a new mode of determining the value of life contingencies'. *Philos Trans R Soc Lond B Biol Sci* **370**, doi:10.1098/rstb.2014.0379 (2015).

Aalen, O., Borgan, O. & Gjessing, H. *Survival and Event History Analysis: A Process Point of View (Statistics for Biology and Health)*. (Springer, 2008).

Barbi, E., Lagona, F., Marsili, M., Vaupel, J. W. & Wachter, K. W. The plateau of human mortality: Demography of longevity pioneers. *Science* **360**, 1459-1461, doi:10.1126/science.aat3119 (2018).

Makeham, W. M. On the Law of Mortality and the Construction of Annuity Tables. *The Assurance Magazine, and Journal of the Institute of Actuaries* **8**, 301-310 (1860).

Vaupel, J. W. *et al.* Biodemographic trajectories of longevity. *Science* **280**, 855-860 (1998).

Yashin, A. I., Vaupel, J. W. & Iachine, I. A. A duality in aging: the equivalence of mortality models based on radically different concepts. *Mech Ageing Dev* **74**, 1-14 (1994).

Kirkwood, T. B. & Austad, S. N. Why do we age? *Nature* **408**, 233-238, doi:10.1038/35041682 (2000).

Fontaine, F., Stewart, E. J., Lindner, A. B. & Taddei, F. Mutations in two global regulators lower individual mortality in Escherichia coli. *Mol Microbiol* **67**, 2-14, doi:10.1111/j.1365-2958.2007.05988.x (2008).

Fontana, L., Partridge, L. & Longo, V. D. Extending healthy life span--from yeast to humans. *Science* **328**, 321-326, doi:10.1126/science.1172539 (2010).

Kenyon, C., Chang, J., Gensch, E., Rudner, A. & Tabtiang, R. A C. elegans mutant that lives twice as long as wild type. *Nature* **366**, 461-464, doi:10.1038/366461a0 (1993).

Pedro de Magalhaes, J. *et al.* A Reassessment of Genes Modulating Aging in Mice Using Demographic Measurements of the Rate of Aging. *Genetics* **208**, 1617-1630, doi:10.1534/genetics.118.300821 (2018).

Battesti, A., Majdalani, N. & Gottesman, S. The RpoS-mediated general stress response in Escherichia coli. *Annu Rev Microbiol* **65**, 189-213, doi:10.1146/annurev-micro-090110-102946 (2011).

28 Peterson, C. N., Levchenko, I., Rabinowitz, J. D., Baker, T. A. & Silhavy, T. J. RpoS proteolysis is
controlled directly by ATP levels in Escherichia coli. *Genes Dev* **26**, 548-553,
doi:10.1101/gad.183517.111 (2012).
29 Kuroda, A. *et al.* Role of inorganic polyphosphate in promoting ribosomal protein degradation by the
Lon protease in E. coli. *Science* **293**, 705-708, doi:10.1126/science.1061315 (2001).
30 Lever, M. A. *et al.* Life under extreme energy limitation: a synthesis of laboratory- and field-based
investigations. *FEMS Microbiol Rev* **39**, 688-728, doi:10.1093/femsre/fuv020 (2015).
31 Radzikowski, J. L. *et al.* Bacterial persistence is an active sigmaS stress response to metabolic flux
limitation. *Mol Syst Biol* **12**, 882, doi:10.15252/msb.20166998 (2016).
32 Medawar, P. B. *An Unsolved Problem of Biology: An Inaugural Lecture Delivered at University*
College, London, 6 December, 1951. (H.K. Lewis and Company, 1952).
33 Mauri, M. & Klumpp, S. A model for sigma factor competition in bacterial cells. *PLoS Comput Biol*
10, e1003845, doi:10.1371/journal.pcbi.1003845 (2014).
34 Scott, M., Gunderson, C. W., Mateescu, E. M., Zhang, Z. & Hwa, T. Interdependence of cell growth
and gene expression: origins and consequences. *Science* **330**, 1099-1102, doi:10.1126/science.1192588
(2010).
35 Maharjan, R., Seeto, S., Notley-McRobb, L. & Ferenci, T. Clonal adaptive radiation in a constant
environment. *Science* **313**, 514-517, doi:10.1126/science.1129865 (2006).
36 Spira, B., de Almeida Toledo, R., Maharjan, R. P. & Ferenci, T. The uncertain consequences of
transferring bacterial strains between laboratories - rpoS instability as an example. *BMC Microbiol* **11**,
248, doi:10.1186/1471-2180-11-248 (2011).
37 Hol, F. J. *et al.* Spatial structure facilitates cooperation in a social dilemma: empirical evidence from a
bacterial community. *PLoS One* **8**, e77042, doi:10.1371/journal.pone.0077042 (2013).
38 Vulic, M. & Kolter, R. Evolutionary cheating in Escherichia coli stationary phase cultures. *Genetics*
158, 519-526 (2001).
39 Hengge, R. in *Bacterial Stress Responses, Second Edition*   (American Society of Microbiology,
2011).
529

Methods

Experimental methods

Microfluidic chip fabrication

Our Polydimethylsiloxane (PDMS) lab-on-a-chip system consists of two layers,

containing the flow channel and the array of cell-sized chambers. The two layers are

fabricated separately using soft-lithography technology, and then bonded together to produce

the microfluidic device (Fig. S2)

To fabricate the negative master for the piece containing the flow channel, SU8 3050

photoresist (Microchem, MA, USA) was patterned on a silicon wafer using photolithography.

SU8 3050 was spin-coated on a silicon wafer at 4000 rpm for 30 seconds, baked at 95 °C for

15 minutes, and then subjected to UV exposure (25 s, 10m W/cm2). After post-exposure

baking (95 °C for 5 minutes), the master was developed using SU8 developer (Microchem,

USA), rinsed with isopropanol and dried with filtered nitrogen.

The master for the layer with cell-sized chambers was fabricated using reactive ion

etching (R.I.E.) technology on a silicon wafer. The mask was patterned on a silicon wafer

using photoresist AZ5214 (Microchem, USA), and a 100-nm layer of nickel was sputtered on

to the substrate. A lift-off procedure was applied to remove the photoresist layer yielding the

metal mask for the R.I.E. process. By adjusting the R.I.E. parameters (SF_6 = 4 sccm, CHF_3 =

16 sccm, pressure = 10 mTorr and power = 30 W), we managed to achieve a large array of

micro-pillars with high aspect ratio (diameter = 1.2 μm and height = 6 μm).

To form the device, PDMS mixtures (RTV615, Momentive Performance Materials

Inc., Waterford, NY) were poured (flow channel) and spin-coated (array) onto the masters to

a thickness of 5 mm (main channel) and 80 μm (array) respectively. Heat curing initially

formed solid PDMS layers with patterned surfaces. After drilling inlets and outlets through

the flow channel layer, and mounting the array layers onto cover glasses, the two layers were

then bonded together using oxygen plasma (90s, 1000 mTorr). Finally, the assemblies were

cured at 80 °C overnight to produce the integrated microfluidic chips. On the day of use, the

wetted surfaces of the PDMS chip were first activated by 90s exposure to oxygen plasma

(90s, 250 mTorr) immediately followed by infusion of 20% (v/v) polyethylene glycol

(PEG400) solution to prevent bacterial adhesion and biofilm formation.

Media preparation

All equipment used for media preparation, sterilisation and infusion were made of

non-leaching materials (glass, Polytetrafluoroethylene or similar perfluoropolymer material)

to avoid contamination with trace level carbon sources from leachable plastic additives (see

supplementary material for details). Media were filter sterilised (0.2 μm) to avoid volatile

organic contamination during autoclaving, and glassware was sterilised by dry heat. Carbon-

free minimum media mentioned below refer to those prepared in this fashion.

Strain information

All lifespan distributions described in the main text are measured for strains of the

Keio *E. coli* BW25113 strain ("widltype") single-gene knockout collection[40]. For the

knockout strains, the presence and location of genomic inserts were verified by kanamycin

resistance and PCR amplification. The general stress response phenotypes of Δ*rpoS* and

Δ*rssB* were verified using the catalase test. In addition, in developing the microfluidic device

and validation of our method, we used an MG1655 derived *E. coli* strain with a

chromosomally integrated CFP under P_{2rrnB} constitutive promoter[41] (Fig. 2b).

Cell culture and loading

Single isolated colonies of the bacterial strains *E. coli* wildtype, Δ*rpoS* and Δ*rssB*

were grown overnight in minimal medium (1 × M9 salts, 2 mM $MgSO_4$ and 0.1 mM $CaCl_2$)

supplemented with 20% w/v glucose (final concentration 0.4%). The following day the

overnight cultures were diluted 200-fold in 50 mL of fresh medium in 250 mL Erlenmeyer

flasks and grown to early exponential phase (OD_{600} = 0.2). This growth phase was chosen to

guarantee that the variation of birth time among cells was less than one cell cycle and thus to

minimise the uncertainty of lifespan measurement. Cells were concentrated by centrifugation

(4,000 rpm x 15 min, 37 °C) and washed by 3 cycles of gentle re-suspension with carbon-free

minimal medium and centrifugation prior to injection into the microfluidic channels. Cells

were then trapped into the dead-ended wells by centrifugation at 2,000 rpm for 15 min at 37

°C with a surface density up to 6.25×10^4 cells/mm^2. The main channels were then thoroughly

washed with carbon-free minimal medium.

Experimental setup and microscopy

A constant flow of carbon-free M9 minimal medium at 20µl per hour was provided to

the micro-channels using a high-precision syringe pump (Harvard Apparatus PHD 2000

Programmable) and Hamilton GC-grade glass/PTFE syringes (Gastight 1000 Series). PTFE

tubing was used to connect the syringes to the microfluidic chip. The medium was

supplemented with 1.5% (v/v) polyethylene glycol (PEG400) to prevent unspecific adherence

of cells to the channels and 5 µg/mL propidium iodide, as a fluorescent indicator of cell

viability, was added. Cell viability was monitored using temperature-controlled (37 °C)

automatic time-lapse microscopy (Zeiss AX10, 63x oil-immersion objective, controlled with

MetaMorph® software). Focus was maintained by a Z-scanning maximum-contrast

procedure using phase-contrast illumination. Focus was re-adjusted before each imaging

cycle for every position and maintained within a Z-range of 0.2µm around the maximum-

contrast Z-position. Phase-contrast and fluorescence images (P.I. signal ex. 546nm/12nm,

em. 605nm/75nm) were acquired for every stage position once every hour for up to 150

hours.

Growth phenotypes

Growth phenotypes of aforementioned *E.coli* strains on selected carbon sources in

minimal media were measured in the 96-well format using a TECAN Spark® microplate

reader. To induce the appropriate metabolic enzymes before growth curves can be measured,

strains were first grown for 24 hours in minimal medium (M9, as in those used for

microfluidic experiments) supplemented with the assayed carbon sources. The optimal

densities of these cultures were determined, and diluted into fresh media identical to those

used for the overnight cultures. The dilution ratios were chosen so that all experimental cell

cultures have an initial optimal density of $OD_{600} = 0.002$. For each experimental well on the

microplates, 50µl mineral oil was added to 100µl cell culture to prevent evaporation during

the experiments. The microplates were then maintained at 37°C, and shaken constantly in

double-orbital motion at 150rpm by the plate reader. Microplates with flat-bottomed wells

were used to maximise agitation. OD_{600} readings were taken every 10min. The growth

phenotypes used in Fig. 6 are based on minimum media culture supplemented with 60mM

acetate as the carbon source.

Statistical and computational methods

Image analysis

The cells in our experiments were trapped in an evenly spaced 2D-grid. The

fluorescence signals of every cell throughout their whole lifespans could be extracted at fixed

positions, once images in the time-lapse stack were properly registered. A simple registration

procedure might misidentify one cell for another because the cells are vertically imaged and

look very similar to each other if only local features are considered. To register the images

based on global features, such as the presence/absence of cells at individual grid positions,

we devised a two-pass, coarse-to-fine registration strategy. Cells were first identified and

segmented within the images using the Point-Spread Function[42]. These segmented binary

image stacks containing global information were registered in a coarse pass using least-

square minimisation. The obtained 2D-translations were applied to the original images. A

second fine registration pass was executed on these pre-registered original images using the

Pyramid Approach[43]. After registration, the salient positions of cells were detected on the Z-

projected images of the whole time-lapse stacks, and fluorescence time-series were extracted

from these positions.

Regularised estimation of P.I. signal

To determine the true P.I. signal of each individual cell and to remove the noisy

effects of focusing fluctuations, we designed and implemented a correction algorithm to the

raw fluorescence intensity time-series (Fig. S3). Our three-step algorithm estimated the

focusing noises of each imaging position, and deduced them from the raw time-series to

arrive at the true P.I. signals of each cell. The first step of our algorithm took advantage of the

fact that focusing fluctuations should change synchronously the intensities of all fluorescent

objects within an imaging position, while the true P.I. signal from the cells should move

independently of each other. By averaging the fold changes in intensities over all cells within

a given time-lapse image stack, the focusing noise was enhanced while cell-specific signals

were spread over hundreds of independent time-series. In the second step, the averaged fold

changes was decomposed into focusing noises and population-wide P.I. trends. This was

possible because noise from focusing should have very quick fluctuations (small

autocorrelation time on the order of imaging cycles) yet no long-term trends (focusing was

maintained with a narrow Z-range of 0.2μm). We applied a Total Variation Regularisation

algorithm[44] to effectively de-noise the average fold changes to produce the population-wide

long term trends. In the last step, the population-wide P.I. trends were combined with the

cell-specific signals to recover the true P.I. signals used to determine the times of death. See

Fig. S3 for the details and the effects of the algorithm.

Survival analysis

Populations within the same microfluidic channels were followed at multiple imaging

positions, distributed evenly throughout the flow channel. We tested the statistical

consistency and homogeneity of lifespan distributions among subpopulations at different

imaging positions within the same channel (see Fig. 5a and Fig. S4 for examples of such

subpopulations). The P.I. time-series data and lifespan distributions from each imaging

position were visualised in the style of Fig. 2d (Fig. S4a). Statistically, we tested for non-

parametric differences between subpopulation lifespan distributions using a two-sample

Kolmogorov-Smirnov statistic $\sup\{|Y(t)|\}$, *i.e.* the supremum of empirical distribution

distances $|Y(t)|$ normalised to account for censorship[45] (Fig. S4b).

After passing the consistency test, the subpopulations from the same channel were

merged to form the experimental cohorts. The lifespan data of each cohort was fitted

parametrically using the family of lifespan distributions described in the main text. Maximum

likelihood (ML) estimators of the model parameters and their confidence intervals were

obtained using the R package 'flexsurv'. We tested the 2-, 3- or 4-parameter hazard models

mentioned in the main text, with the 4-parameter Gamma-Gompertz-Makeham (GGM)

model being the most general, and chose the best among these candidate models according to

the Akaike Information Criterion (AIC) at their maximum-likelihood parameters.

Goodness-of-fit of the ML models were tested using the one-sample version of

$\sup\{|Y(t)|\}$. The best models and their fitting residues $Y(t)$ were plotted in supplementary

material. For comparison and visual inspection, we also plotted alongside the Kaplan-Meier

estimators for survivorship, Nelson-Aalen estimators for cumulative hazards and their

respective 95% C.I..

Experimental design and experimental variation analysis

Three independent experimental cohorts were tracked on separate dates for each

strain. Two statistical methodologies were used to assess the level of experimental variability,

and test for significant differences in lifespan distribution and its parameters among the

strains. Overall variations in lifespan distribution were analysed using hierarchical clustering.

The two-sample Kolmogorov-Smirnov statistic $\sup\{|Y(t)|\}$[1,45] mentioned in the previous

section was used as a distance measure in complete linkage clustering. Preferential

aggregation of experimental cohorts of the same genotype within the lower branches of the

clustering tree was taken as evidence that lifespan distributions of the strains were

significantly different.

Because the GGM model adequately described our data, we also analysed variability

of its parameters and the sources of said variability. Generalised Linear Models (GLMs) with

GGM as the probability distribution were built to examine the explanatory power of

categorical covariates such as experimental cohorts (*exp*) or strain genotypes (*strain*).

Specifically, the null hypothesis H_0 that cohorts with the same genotype have the same

ageing rate b were tested against the alternative hypothesis H_1 that experiment cohorts all

have significantly different ageing rates regardless of their genotypes. Since H_0 is a special

case of H_1, both ΔAIC and likelihood ratio test were used and H_1 was rejected. See

supplementary material to see more details on GLM model selection and hypothesis testing.

Fitness models

Fitness is defined over a time period (t_1, t_2) as $f(s, t_1, t_2) = \Delta \ln(\textit{population size})/(t_2 - t_1)$,

where s denotes one of the 3 strains. We calculate fitness based on time spent in "feast" or

"famine" episodes. These fitness functions were extrapolated from experimental data. For

"famine" episodes, changes in logarithmic population size were simply the negatives of

cumulative hazards $H(t_1)-H(t_2)$, so that $f_{\text{famine}}(s, t_1, t_2) = [H_s(t_1) - H_s(t_2)]/(t_2 - t_1)$. Extrapolations

from experimental data were done using the fitted GGM models. For "feast" episodes,

$f_{feast}(s, t_1, t_2) = \dfrac{1}{t_2 - t_1} \int_{t_1}^{t_2} \mu_s(t)dt$. Specific growth rates $\mu_s(t)$ and maximum growth rates μ_s^{max}

were determined from the experimental growth curves smoothed by cubic B-Splines. For

"feast" episodes longer than the time at which μ_s^{max} was reached, we assumed exponential

growth at μ_s^{max} .

Overall fitness f(s, k_1, k_2) of each strain at a given set of environmental parameters k_1

and k_2, is calculated by averaging the episodic fitness f_{famine} and f_{feast} over 5000 episodes of

both "feast" $\tau_{e,i}$ and "famine" $\tau_{m,i}$, where i = 1, ... , 5000. Episode lengths $\tau_{e,i}$ and $\tau_{m,i}$ are

independently drawn from exponential distributions parameterised by the two ecological

rates, k_1 and k_2, as shown in Fig. 6a. In summary, the formula for over fitness is f(s, k_1, k_2) =

$\dfrac{\int_0^{\tau_{e,i}} \mu_s(t)dt - \Delta_0^{\tau_{m,i}} H_s(t)}{\sum\limits_i \tau_{e,i} + \sum\limits_i \tau_{m,i}}$.

For visual comparison in the form of the colourmap in Fig. 6b, for each set of

environmental parameters (k_1, k_2), the fitness of the 3 strains are combined into a 3-tuple and

normalised using the softmax function. If we simply denote average fitness of strain s as f_s,

the formula for the elements of 3-tuple is $\dfrac{e^{Tf_s}}{\sum\limits_s e^{Tf_s}}$, where T is the comparison timespan and is

chosen to be 200 hours in Fig. 6b.

Programming codes and data availability

The image analysis procedure described above was implemented in Java as an ImageJ

plugin. Fitness modelling, plotting and timeseries analyses including noise correction and

time-of-death determination were implemented in Python. We relied on R package 'flexsurv'

for the core algorithm of survival analysis and GLM, and Python package 'rpy2' was used for

interoperability between the Python and R codes. All source codes and data are made

available through GitHub.

40   Baba, T. *et al.* Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio
collection. *Mol Syst Biol* **2**, 2006 0008, doi:10.1038/msb4100050 (2006).
41   Giraud, A. *et al.* Dissecting the genetic components of adaptation of Escherichia coli to the mouse gut.
PLoS Genet **4**, e2, doi:10.1371/journal.pgen.0040002 (2008).
42   Yang, Y., Song, X. & Lindner, A. B. in *Methods in Microbiology* Vol. 43 (eds Colin Harwood &
Grant J. Jensen) 49-68 (Academic Press, 2016).
43   Thevenaz, P., Ruttimann, U. E. & Unser, M. A pyramid approach to subpixel registration based on
intensity. *IEEE Trans Image Process* **7**, 27-41, doi:10.1109/83.650848 (1998).
44   Chartrand, R. Numerical Differentiation of Noisy, Nonsmooth Data. *ISRN Applied Mathematics* **2011**,
11, doi:10.5402/2011/164564 (2011).
45   Fleming, T. R., O'Fallon, J. R., O'Brien, P. C. & Harrington, D. P. Modified Kolmogorov-Smirnov
Test Procedures with Application to Arbitrarily Right-Censored Data. *Biometrics* **36**, 607-625,
doi:10.2307/2556114 (1980).

Available resources

C P N

"Feast"

Reproduction

Colonisation

Death

Dispersal

"Famine"

Senescence

Dissipated resources

C P N

Physiological processes

Stochastic events

a

b

c

Time

d

Cell #

1000

2000

3000

4000

0   20   40   60   80   100   120

Time (h)

a

b

a

"Feast"

Available resources

Population gain

Time (h)

Anabolism

Biomass

Colonisation rate, k_2

RpoD ⊣ RpoS

Dispersal rate, k_1

Biomass

Catabolism

Time (h)

Survivorship

Dissipated resources

"Famine"

b

Δ*rpoS*   *wt*   Δ*rssB*

Lifestyle ratio, k_1/k_2

"Famine" mean residence time, $1/k_2$ (h)

Maintenance capacity

U

$U - J\Delta t$

Physiological homeostasis

Molecular damage

Molecular repair

Maintenance energy flux J

Hazard rate h

Death

Free radicals damaging protein, membrane & DNA

Physiological processes

Molecular processes