Temporal scaling of ageing as an adaptive strategy of *Escherichia coli*

Yifan Yang[1,2,*], Ana L. Santos[1], Luping Xu[1,2,3], Chantal Lotton[1], François Taddei[1,2] and Ariel B. Lindner[1,2,*]

[1] INSERM U1001, Faculté de Médecine, Université Paris Descartes, Paris, France

[2] Center for Research and Interdisciplinarity, Paris, France

[3] Current address: Center for Nano and Micro Mechanics, School of Aerospace Engineering, Tsinghua University, Beijing, China

[*] Correspondence: yifan.yang@inserm.fr and ariel.lindner@inserm.fr

Abstract

Natural selection has long been hypothesised to shape ageing patterns, but whether and how ageing contributes to life-history evolution remains elusive. The complexity of various ageing-associated molecular mechanisms and their inherent stochasticity hinder reductionist approaches to the understanding of functional senescence, *i.e.* reduced fecundity and increased mortality. Recent bio-demographic work demonstrated that high-precision statistics of life-history traits such as mortality rates could be used phenomenologically to understand the ageing process. We adopted this approach to study cellular senescence in growth-arrested *E. coli* cells, where damages to functional macromolecules are no longer diluted by fast *de novo* biosynthesis. We acquired high-quality longitudinal physiological and life history data of large environmentally controlled clonal *E. coli* populations at single-cell resolution, using custom-designed microfluidic devices coupled to time-lapse microscopy. We show that *E. coli* lifespan distributions follow the Gompertz law of mortality, a century-old actuarial observation of human populations, despite developmental, cellular and genetic differences between bacteria and metazoan organisms. Measuring the shape of the hazard functions allowed us to disentangle quantitatively the demographic effects of ageing, which accumulate with time, from age-independent genetic longevity-modulating interventions. A

pathway controlling cellular maintenance, the general stress response, not only promotes

longevity but also temporally scales the whole distribution by reducing ageing rate. We

further show that *E. coli*, constrained by the amount of total biosynthesis, adapt to their

natural feast-or-famine lifestyle by modulating the amount of maintenance investment,

rendering ageing rate a highly evolvable life-history trait.

Background

The biology of ageing and senescence is centred on the duality of individual frailty

and population resiliency. Throughout the course of normal metabolism, components of

living systems such as cells, lipids, proteins and DNA inevitably suffer from wear-and-tear

such as free-radical damages. This constant and collective decay eventually leads to the loss

of vital functions and collapse of individuals. Understanding the way that system failure

emerges out of distributed microscopic damages could reveal how functional components are

organised into self-maintaining individuals in the first place[1,2]. Yet for some organisms in the

tree of life, ageing does not lead to increased mortality and declined fertility[3]. Organisms

possess the abilities to repair or replace most of the damages to their components,

exemplified by the "immortal germ line"[4]. Evolutionary biologists attribute the apparent

senescence of metazoan somas to the inadequate investment in cellular maintenance as an

adaptive strategy to maximise lifetime reproductive success, due to the trade-offs between the

survival and reproduction of the young on one hand, and maintenance for the benefit of the

old on the other[5,6]. Presumably, natural selection has to operate through the molecular

"levers" of damage accumulation and/or repair to achieve such life-history optimisation.

Escherichia coli, a single-cell prokaryote with short lifespans, has historically served

as a model organism that resolved many fundamental questions in biology. *E.coli* cells, as

their metazoan counterparts, suffer from damages to their components, which lead to cellular

senescence[7,8]. In exponential growth, these damages are quickly diluted by *de novo* biosynthesis, and the effects of cellular senescence mitigated by rapid and robust reproduction[9]. Yet, the natural life cycle of *E.coli* entails a much wider range of physiological conditions than exponential growth. Most of bacterial cells spend much of their lives in resource-limited growth arrested conditions, where *de novo* biosynthesis rates are slower[10] and cells undergo senescence due to the accumulation of molecular damages such as protein misfolding and oxidation (Fig. 1)[11]. Despite the lack of fixed separation of germ and soma cells, the ability to survive the wear and tear of cellular components during growth arrests contribute to bacterial overall fitness as much as the ability for exponential growth. We, therefore, adopted a bio-demographic approach[1,2] to understand how modulation of molecular damage repair could shape ageing dynamics in growth-arrested *E.coli*.

Custom-designed microfluidic devices allow longitudinal tracking of vitality at single-cell level

We first set out to acquire high quality longitudinal physiological and life-history data at single-cell resolution of clonal populations of *E.coli*. For decades, limitations of traditional cell culture methods have frustrated quantitative physiologists' attempts to understand bacterial maintenance[12,13], due to both media-cell interactions and cellular interaction such as cross-feeding and cannibalism. To be able to measure individual *E.coli* cell lifespan in constant and homogenous environmental settings, we designed a novel microfluidic device with cell-dimension chambers (Fig. 2). Cells are trapped in an array of single-cell-wide dead-ended wells, with openings to a main flow channel. Constant flow of fresh media in this main channel provides necessary nutrients and eliminates metabolic waste, cell debris and intercellular crosstalk so that the environmental conditions are maintained constant over time. Cells with appropriately expressed fluorescent markers are imaged bottom-up and appear as

fluorescent spots (Fig. 2b). This experimental system allows easy tracking of cohorts of a large number of individual bacterial cells for prolonged periods (up to 7 days).

In order to observe cell mortality, we included in the carbon-source-free medium a red-fluorescent, DNA-binding, bacterial viability dye, Propidium Iodide (P.I.), which penetrates the cells only when cellular membrane potentials are disrupted. P.I. staining has been established as an effective proxy of cellular death and correlates well with cell viability assessed by proliferating potential[14]. We used P.I. at a concentration 4-fold lower than the previous concentration that had no effects on *E.coli* viability and growth[14]. Automatic time-lapse fluorescence microscopy and fixed geometry of our devices allowed longitudinal quantification of P.I. signal for every single cell in the population. We defined the half point between peak P.I. signal and background fluorescence as a threshold to establish the time-of-death. At the end of the experiments, 70% ethanol was injected into the device in order to account for every cell in the cohort, and establish a lower bound for the time-of-death of surviving cells. These cells were censored at the time that their P.I. signals crossed their respective thresholds (see Methods).

Despite being genetically clonal and environmentally controlled, cells did not share the same time or manner of death, as measured by the P.I. time-series. The transition from life to death of a representative cohort (N= 4744) could be visualised by the boundary between dark to light in Fig. 2d. Because cells were sorted vertically according to estimated lifespans, the shape of this boundary represents the survival function of the population. Considering ageing and death as a stochastic process of system reliability reduction[1,2], the observed time-of-death distribution could be viewed as the population's first-passage time distribution. In addition, we observed individual differences in death trajectories. Short-lived cells tended to have very sharp P.I. increases that associated with abrupt losses of membrane

integrity; while those that die late tended to suffer a type of "slow death" characterised by a gradual P.I. increase over the course of 10-15 hours.

E. coli lifespan distributions follow the Gompertz law

For both demographers and reliability engineers, the age-associated increase in death probability, also known as the hazard rate h(t), is considered to be a hallmark of ageing[1-3]. The detailed population statistics derived from our experiments are particularly useful in estimating the shape of the hazard functions over the whole lifespan. We find that *E.coli* lifespan distributions have, as their main feature, regimes with exponentially increasing hazard rates, *i.e.* the Gompertz law of mortality $h(t) \sim h(t_0) e^{b(t-t_0)}$, where $h(t_0)$ is the initial hazard rate and b is the Gompertz ageing rate[15,16]. We estimated $h(t)$ directly with binomial error by binning mortality events within discrete time intervals (τ_i, τ_{i+1}), *i.e.* $\tilde{h}_i \sim d_i/Y_i/\Delta\tau_i$, where Y_i is the number of individuals at risk at time τ_i and d_i is the number of cell deaths within (τ_i, τ_{i+1}) (see Fig. 3). For wildtype cells, this exponential regime spans from about 13 hours to 93 hours, corresponding to approximately 90% of all cell deaths (N_{total}=4744, see Fig. 3b, top x-axis), and ranges at least 100-fold changes in hazard rates ($2*10^{-3}$ h^{-1} to $2*10^{-1}$ h^{-1}). Within this exponential regime, the doubling time of hazard rate is 9.4±0.5 hours. In comparison, the doubling time of human mortality hazards is about 8 years[15].

Hazard rates not only define lifespan distributions but can also be thought of as surrogates for system vulnerability, whose dynamics through time reflect the physiological consequence of ageing. In this light, the Gompertz law can be interpreted as a dynamic equation governing the ageing process: $dh(t)/dt = bh(t)$, where b is the Gompertz ageing rate. The integral version, $h(t) = bH(t)+h(t_0)$, where $H(t) = \int_0^t h(t')dt'$, can be directly observed in the phase plane of the cumulative hazards $H(t)$ without any free parameters (Fig. 3b). The

integral equation is used because *H(t)* can be estimated independent of binning, by using the

Nelson-Aalen (N-A) estimator $\tilde{H}_{NA}(\tau_i)$[17]. The fact that the theoretical trajectory, including

the aforementioned linear regime, lies within the confidence intervals (CI) of almost every

state coordinate ($\tilde{H}_{NA}(\tau_i)$, $\tilde{h}_i$) lends strong statistical credence to the Gompertz law for

describing our data. In addition, the observed trajectory in the phase plane links the

instantaneous hazard rate *h(t)* at any given time *t*, with the total proportion of population that

have died before time *t* (because *H(t) = -ln[S(t)]*, where *S(t)* is the survival function).

The hazard rate observed deviates from the exponential regime at the beginning (t<15

hours, 3% cell deaths) and the end (t>93 hours, 7% cell deaths) of the total lifespan. These

observations are reminiscent of similar deviations from Gompertz law in human mortality

data[18,19]. In our case, the additional mortality at the early age might result from harvesting

and transferring exponentially growing cells from batch culture directly to growth-arresting

conditions inside our microfluidic chip, in a way analogous to infant mortality in human

mortality data, which can be modelled by an age-independent component (λ in the Gompertz-

Makeham model $h_{gm}(t) = \lambda + \frac{b}{\beta}e^{bt}$, where β controls the age at which the Gompertz regime

overtakes the Makeham term λ).

Late-life hazard decelerations are common phenomena for both model organisms and

human populations[18,20]. It is thought that these hazard decelerations often do not reflect

ageing decelerations for individuals, but result from changes in composition of heterogeneous

populations. The more fragile individuals are more likely to die and be removed from the

cohort. When hazard rates are high late in life, significant portions of the populations are

removed so that the observed hazard rates of the surviving populations would be smaller than

the original cohorts on average. We found that an extension to the Gompertz law named

Gamma-Gompertz, originally used to model old-age human model data[21], could also

satisfactorily model late life decelerations in our data, for both wildtype and mutant strains (see below). It could be understood as accounting for the compositional changes by an additional parameter, *s*, controlling the level of frailty heterogeneity:

$$h_{gg}(t) = \frac{b}{\beta} e^{bt} \cdot s[S_{gg}(t)]^{1/s}$$, where $S_{gg}(t)$ is the survival function of Gamma-Gompertz, and other parameters are as before. Despite being genetically and environmentally non-distinguishable, we think it is reasonable to assume frailty heterogeneity exist in our single-cell populations due to the stochastic nature of cellular biochemistry, explaining the suitability of the Gamma-Gompertz for our data.

We thus used the Gamma-Gompertz-Makeham (GGM) extension model with 4 parameters to fully model our data across the whole lifespan, with parameters estimated by likelihood maximisation:

$$h_{ggm}(t) = \lambda + \frac{bs}{1 + (\beta - 1)e^{-bt}}$$

Goodness-of-fit was evaluated using a one-sample Kolmogorov-Smirnov test for right-censored survival time data. The appropriateness of the model terms was evaluated by comparing Akaike Information Criterion (AIC) of the candidate models (see methods).

The general stress response of *E. coli* modulates the ageing rate

Evolutionary theories of ageing predict the existence of ageing-modulating mechanisms and their activation by nutrient limitation[22]. Several nutrient-sensing pathways in both bacteria and metazoans have been shown to control stress resistance and reduce mortality[23-25]. Yet their roles in delaying ageing-related damage accumulations are often controversial. It is often difficult and requires large experimental cohorts to disentangle their effects on ageing from age-independent components of longevity[26]. Having established a method to measure ageing rates using high-quality mortality statistics, we harnessed this

system to shed light on potential ageing-modulating mechanisms of *E.coli* that might be activated in response to nutrient deprivation.

Many bacteria species regulate the level of cellular maintenance through a genetic pathway called general stress response (GSR), controlled by the master transcriptional regulator *rpoS* that is activated by nutrient-deprivation among other signals[27] (Fig. 4a). To assess the role of *rpoS*-controlled cellular maintenance in the ageing of growth-arrested *E.coli*, we measured the lifespan distributions of two GSR mutants, $\Delta rpoS$, and $\Delta rssB$ with that of the wildtype strain. $\Delta rpoS$ is the null mutant and $\Delta rssB$ displays an elevated GSR due to increased RpoS stability[28]. We observed that higher GSR promotes longevity in the microfluidic experiments, whereas the absence of GSR results in shortened longevity (Fig. 4b and Fig. 5a). Our large sample sizes allowed us to directly measure the hazard dynamics (Fig. 4c) of each strain, which could disentangle GSR's effects on ageing rate, as opposed to age-independent components of longevity.

Significantly, we found that increased GSR reduces the rate of ageing (Fig. 4d and Fig. 5c). Given how well the GGM model fitted for all 3 strains, the impact of genotypes on ageing parameters such as ageing rates could be extracted using generalised linear models (GLM). With enhanced GSR, $\Delta rssB$ cells double their mortality risk every 14.1 hours, with a 95% CI ranging from 13.1 to 15.3 hours, compared to 9.4 hours for the wildtype (CI 8.9 - 9.9 hours) and 7.3 hours for the null strain $\Delta rpoS$ (CI 6.8 to 7.7 hours). Variability of ageing rate measurements was assessed using 3 independent experimental cohorts for each strain (Figure S1). We visualised non-parametrically the overall experimental variations (Fig. 5a and b). Parametric differences in ageing rates among experimental repeats were tested using GLM models and AIC. We confirmed that experimental repeats shared similar ageing rates (H_0: Same aging rate for experimental repeats. d.f. $H_0 - H_1 = -2$; $\Delta AIC_{rssB}=-2.56$, $\Delta AIC_{wt}=-2.26$, $\Delta AIC_{rpoS}=-2.30$; $N_{wt}=6867$, $N_{rssB}=6969$, $N_{rpoS}=4793$).

The systematic increase in vulnerability of *E.coli* in our experiments is likely driven by the catabolism of pre-existing macromolecules and dissimilation of biomass[29,30], which is necessary to provide energy to express housekeeping genes and maintain physiological homeostasis[10,31]. Our finding that GSR modulates ageing rate suggests that optimising this maintenance energy requirement is likely one of the physiological functions of the RpoS regulon (see Discussion).

Evolutionary trade-offs mediated by the general stress response

Is slower ageing an adaptive life-history trait? This was one of original question raised by Sir Peter Medawar[32] that motivated much of later ageing research. Optimal life-history theory suggested that trade-offs and constraints among fitness components shape metazoan ageing rates[6]. The possibility of modulating ageing rate through GSR offers us the opportunity to test these ideas in a fast-evolving organism as *E. coli*.

The relatively well-understood GSR pathway provides a clear molecular mechanism for a trade-off between growth and maintenance. The master regulator *rpoS* encodes the RNA polymerase (RNAP) sigma subunit σ^S, which competes with the other sigma factors including the vegetative σ^D to recruit the core RNAP and direct the transcription and translation machinery towards the RpoS regulon. By titrating protein synthesis activity away from metabolic and ribosomal genes controlled by σ^D, RpoS activity inhibits growth and nutrient assimilation[33]. We measured quantitatively the growth impact of modulating GSR levels, and modelled its effect using a simple course-grained model of proteome sectors[34]. We found that the proportion of protein synthesis devoted to the RpoS regulon linearly increases the timescale of growth (see top axis in Fig. 6a).

To assess the life-history optimality of different ageing rates, we integrated numerically the experimentally derived growth rates and mortality rates into fitness, defined

as the long-term population growth rates. In contrast to metazoans, for whom fertility and mortality schedules are connected through fixed age structures, our model of *E. coli* life-history consisted of alternating environmental episodes of varying durations, in which *E. coli* populations either grow ("feast"), or decline ("famine") (Fig. 6a). These feast-or-famine cycles were parameterised by two transition rates, controlling respectively the average lengths of "feast" and "famine" episodes. In order to identify selective pressure for ageing rates, we directly compared the fitness of the three strains with different GSR phenotypes, each representing a different strategic position in the growth-maintenance trade-off. We identified the environmental regimes selecting for faster or slower ageing (Fig. 6b). The boundaries between these regimes were characterised along two axes: lifestyle ratio, defined as the ratio of time spent in "famine" versus "feast"; and "famine" mean residence time, or in other words, average length of "famine" episodes.

There are two necessary conditions for selecting slower ageing strategies represented by Δ*rssB*. First, populations have to spend much more time in "famine" rather than "feast", so that over the long term, population decline rather than grow. Secondly but no less important, given the same lifestyle ratio, famines should consist of longer episodes rather than short but more frequent ones. This condition is necessary due to the exponential mortality dynamics described by Gompertz law: investing in cellular maintenance only becomes beneficial at old age, when the exponentially increasing benefits of slower ageing eventually overcome the more immediate cost on growth. It is the typical timescale of "famine" that provides the selective pressure for ageing rates.

Despite representing the complex regulations of GSR with two mutants, we can now understand the ecological role of GSR activation and its adaptive consequences. Activated by declining nutrient availability, GSR directs resources towards internal maintenance to wait out the adverse conditions, whose lengths determine the optimal activation level. Previous

observations from experimental evolution of *E. coli* support our predictions. In continuous cultures of *E. coli* where the populations do not pass through prolonged growth-arresting bottlenecks, mutations that attenuate or knock out RpoS activity are among the first to arise[35,36]. In contrast to the isolated populations under constant environmental conditions in our chip experiments, *E. coli* populations in nature influence their environments, and also interact with each other. These interactions may give rise to frequency-dependent selection and evolutionary game dynamics between slower and faster ageing strategies, as is observed in experimental evolution[37,38].

Discussion

We obtained high quality single-cell demographic data, using a novel microfluidic device, to demonstrate that ageing of growth-arrested *E. coli* follows the Gompertz law of mortality. Moreover, bacterial general stress response could temporally rescale the lifespan distribution by modulating the ageing rate. We further articulated in a demographic model the trade-offs and selective pressure driving the evolution of ageing rate.

In our work, two different conceptual perspectives of ageing are integrated and applied to one of the most iconic model organism, *E. coli*. One perspective, held by biochemists and physicists, sees ageing as the stochastic and inevitable erosion of organismal order and biochemical redundancy, created by self-reproducing and self-maintaining networks during growth and development. The other perspective, from evolutionary biology, views ageing as a component of the organismal life-history strategy, optimised and fine-tuned by natural selection. These two approaches constitute the proximate and ultimate causes of ageing respectively, with the former providing the constraints and the "lever" for the latter. In bacteria, we indeed observed the stochastic process of ageing and mortality. Lifespan vary significantly among genetically identical individuals in constant, homogeneous environment.

However, at the population level, ageing is characterised qualitatively by the Gompertz law of mortality. In response to environmental conditions, the lifespan distribution is modulated by regulatory and selective forces through temporal scaling, while its general, exponential, shape is preserved. These new empirical observations provided an integrated perspectives to bacterial ageing, and by extension, potentially to the breath of the tree of life.

Although the Gompertz law has been shown to characterise ageing of many metazoan organisms, the physiological correlates of its parameters are not well understood. To rescale the lifespan distribution without changing the general shape of hazard dynamics, GSR has to orchestrate a coordinated response to manage various molecular damages that the organism encounters[1]. Indeed, RpoS regulates hundreds of genes conferring resistance to both internal and external stresses such as oxidative, thermal, acid, alkaline, osmotic, and UV[39]. For this reason, the ageing rate in our case reflects the general level of macromolecular catabolism and may relate to the maintenance energy of bacteria[13]. We hypothesise that the energetic costs of homeostasis and molecular damage repair are paid by the loss of biomass, leading to the gradual increase in the probability of death. It is the rate of energy dissipation, or the rate of living, that correlates with the ageing rate in our system (see Fig. 7).

Despite the simplicity of bacteria and its vast differences from animals, we find that concepts in evolutionary theories of ageing, such as antagonistic pleiotropy[5], still apply. RpoS mediates trade-offs between growth and maintenance, between assimilation of nutrients and dissimilation of biomass. Sigma competition, a well-understood mechanism in the physiology of bacterial stress response, provides a molecular basis for such trade-offs often only hypothesised in metazoan organisms. Activated by nutritional deprivation and shown to decrease the ageing rate, GSR has immediate analogies in calorie-restriction-induced longevity of metazoan model organisms. Whereas in many studies on animal models, calorie-restriction-related pathways have only been shown to promote longevity, our demographic

data allow us to show that GSR indeed changes the rate of ageing. It can be thought of as not only providing resistance against stress, but also insurance against prolonged growth arrests, or in other words, protection against the progression of time.

In the future, the experimental approach and statistical framework described here could combine with classical genetic methods to examine further ways hazard dynamics could be perturbed, and to reveal dynamic features of the complex network shaping cellular senescence under various environmental constraints. Future modelling efforts should clarify how adaptive processes could optimise redundancy of living systems, in trade-offs with other traits, to produce the Gompertz law.

Acknowledgements

This study was supported by the Axa Research Fund Chair on Longevity. We thank X. Song for contributions on image analysis, D. Misevic and other member of INSERM U1001 for their advice and discussions.

Author contributions

Conceived and designed project: Y.Y., L.X., F.T., A.B.L.; Conceived and designed experiments: Y.Y., A.B.L.; performed experiments: Y.Y. A.S. C.L.; Data analysis: Y.Y., Wrote the paper: Y.Y. with contribution from A.B.L. .

Figure 1. The feast-and-famine life cycle of *E. coli*. Single-cell organisms such as *E. coli* undergo developmental transitions not in response to fixed developmental programs but opportunistically in response to environmental changes. Because long-term growth rates in a stable ecosystem remain close to zero, life-history traits in "feast" and "famine" conditions should contribute rather equally to overall fitness. In "feast" conditions, the traits under strong selection are resource assimilation and reproduction, while in "famine" conditions are maintenance and survival. Senescence has been shown to occur in "famine" conditions[11].

Figure 2. Coupling microfluidic chip and time-lapse microscopy to measure *E. coli* lifespan

distribution at single-cell resolution. **a**, 3D model of microfluidic devices used to trap and

isolate large number of single cells. The blue-gray cube in the upper half of the picture

represents the main flow channel where fresh, carbon-source-free media are supplied. The

array of light-blue rods represents *E.coli* single-cells trapped in a 2D-array of cell-sized dead-

end chambers. The arrows represent the media flow that maintain the environmental

homogeneity and removes debris. **b**, Fluorescence microscopy image of the microfluidic

device loaded with *E.coli* cells. Each fluorescent dot corresponds to a single cell trapped in

the 2D array of dead-end wells. Z-axis focus is adjusted so that dead-ends of the micro-wells

are imaged. The cyan pseudo-colour represents constitutively expressed fluorescent protein

fluorescence signal; red pseudo-colour - Propidium Iodide (P.I.) fluorescent signal described

in the text. **c**, Sample time-lapse P.I. images of mortality events, from early (top) to late

(bottom) deaths. **d**, Heat map of P.I. signal time-series for a population of single cells. The

colour values of each row correspond to the time-series of one cell, ranked according to lifespan, top-to-bottom, from the shortest living to the longest living cell. The boundary between dark and bright red indicates the transition from life to death and defines the lifespan distribution. Dark strips after cell deaths indicate decaying DNA or empty wells after cell debris have been washed away. Data shown in **d** have been smoothed according to the procedure described in supplementary material to remove minor signal fluctuations coming from microscopy focusing. The same procedure is used before time-of-death is estimated.

Figure 3. Non-parametric estimation indicates that wildtype *E. coli* lifespan distribution can be characterised by the Gompertz law. Shown are non-parametric estimations of the hazard functions and cumulative hazard functions of the same populations in one particular microfluidic experiment. **a**, Binomial estimators of the hazard rates, shown in log scale. Cell deaths are binned into discrete time intervals, which are marked by the x-axis error bars. The time bins are chosen so that the Nelson-Aalen confidence intervals (CI) in b do not overlap with each other. Error bars on the y-axis are the 95% CI for hazard rate estimated based on binomial distribution. Yellow shading covers regions that deviate from the exponential hazard regime for the shortest and longest living individuals. **b**, Cumulative hazard dynamics in the phase plane. Data, binning intervals and y-axis are the same as those in **a**. Instantaneous hazard rates plotted against the cumulative hazards (bottom x-axis), or equivalently the negative logarithm of survivorship (top x-axis). Nelson-Aalen estimators are

used for the cumulative hazard function. The horizontal error bars are the 95% CI of the

Nelson-Aalen estimator at the centre of the time bins. Dashed lines are maximum likelihood

parametric estimations using the Gamma-Gompertz-Makeham model. The inset provides a

zoom-out view of the whole data range, while the main figure zooms in on the first 63% of

cell deaths.

Figure 4. The general stress response of *E. coli* modulates ageing rate. The lifespan distribution for the wildtype (*wt*), Δ*rpoS* (lacking general stress response) and Δ*rssB* (overexpressing the general stress response) strains are measured multiple times by independent microfluidic experiments. **a**, A scheme representing relevant regulatory features of the general stress response, and in particular the functions of the genes *rpoS* and *rssB*. **b**, Experimental and GGM model survivorship. Representing the experimental survivorship, colour bands are the 95% CI of the Kaplan-Meier (K-M) estimators. Coloured dashed lines are Gamma-Gompertz-Makeham (GGM) models whose parameters are estimated from maximum-likelihood (ML) methods. **c**, Hazard rates estimated using only cell deaths within discrete time intervals (Error bar markers), and GGM hazard models estimated from the whole dataset using ML methods. Similar to Fig. 3a, Vertical error bars are binomial 0.95 confidence intervals and horizontal error bars are the binning time intervals. Data from **b** and **c** are from the same representative experimental cohorts for each strain. **d**, Ageing rates for each strain, estimated by ML-fitted GGM models to 3 independent experiments. Error bars represent 95% CI. The ageing rates are: $b_{rpoS} = 0.095 \pm 0.006$ h^{-1}, $b_{rssB} = 0.049 \pm 0.004$ h^{-1} and $b_{wt} = 0.074 \pm 0.004$ h^{-1}.

Figure 5. Experimental repeatability and variances. For each of wildtype (*wt*), Δ*rpoS* and

Δ*rssB* knockout strains, data from 3 independent experimental replicates are shown. **a**,

Survivorship (Kaplan-Meier estimator, K-M) and cumulative hazards (Nelson-Aalen

estimator, N-A) of sub-populations from different imaging positions. Thick opaque lines

correspond to the data shown in Fig. 4b and c. Thin semi-transparent lines are the K-M and

N-A estimators of sub-populations from different imaging positions, which are used to

constitute the population represented by the opaque lines. Each sub-population is from one

imaging position that corresponds to a roughly 125μm-sided square-patch on the microfluidic

chip, and consists of 150-600 cells. Imaging positions with less than 50 cells are not shown.

b, Hierarchical clustering of the 3 independent experimental replicates of each strain.

Standard agglomerative clustering algorithm is applied using complete linkage and the 2-

sample modified Kolmogorov-Smirnov statistic sup{|Y(t)|} (See methods) as a distance

metric. **c**, Ageing rates 3 independent experimental replicates of each strain. Ageing rates are

estimated similarly as those in Fig. 4d, but independently for each experimental replicate without pooling them together using GLM. Error bars represent 95% CI.

Figure 6. Trade-offs between growth and maintenance mediated by the *rpoS* pathway, and its fitness consequences. **a**, Scheme of the ecological processes (dashed arrows) and regulatory relationships (solid arrows) involved in the trade-offs mediated by *rpoS*. Environments are structured by alternating episodes of "feast" and "famine". Fitness of Δ*rpoS* (red), *wild-type* (blue) and Δ*rssB* (green) as functions of time spent in each environmental episode (age) are plotted in the top and bottom axes. Fitness is defined as the logarithmic change of population sizes. **b**, Fitness comparison of the 3 strains across a range of environmental conditions to identify regimes favouring faster and slower ageing strategies. The colour-coded regions identify environmental conditions under which one strain dominates over the other two.

Absolute fitness of the 3 strains are converted to RGB tuples using softmax normalisation. For each strain and each pair of environmental parameters, fitness is calculated by averaging the population growth/decline rates over 5000 episodes of both "feast" and "famine". Episode lengths are independently drawn from exponential distributions, parameterised by the two ecological rates, k_1 and k_2, as shown in **a**. The fitness functions, population gain for "feast" and survivorship for "famine" (shown in the axes in **a**), are used to determine population size changes in each episode. See methods for details.

Figure 7. Gompertz law of mortality informs and constraints physiological models of ageing and mortality in growth-arrested *E.coli* cells. Depicted are the abstract states and pathways cells pass through during ageing and mortality. Acute damages to essential cellular components, when unrepaired, often lead to cell death, quantified by *h(t)*. Cells could also triage and repair these damages to return themselves back to physiological homeostasis. The energy cost of this maintenance process, *J(t)*, termed maintenance energy[13], need to be repaid by either an external energy source or internal storage in the form of existing biomass. If we assume it is the same set of acute molecular damages that kill the cells or lead to ageing, then *J(t)* could relate to *h(t)* proportionally, *J(t) = R h(t)* (eq1), where *R* is the "friction" coefficient controlled by the relative proportions of the two fates of damaged cells: repair vs death. For carbon-starved cells in our experiments, this flux of energy dissipation decreases the capacity for molecular repair in the future, *U(t)*, leading to cellular senescence. Put it another way, *dU(t)/dt = -J(t)* (eq2). Through these assumptions, the hazard dynamics we described in this work could be recast as the dynamics of maintenance capacity. Our experimental results place non-trivial constraints on how the amount of maintenance capacity regulates the rate at

which cells are damaged molecularly in the first place. If we only consider the Gompertz regime, *i.e. dh(t)/dt = bh(t)* (eq3), combining with (eq1) (eq2) and differentiating on both sides, we have $d^2U/dt^2 + b\ dU/dt = 0$ (eq3) and *dU(t)/dt = -R h(0)* , where *b* is the Gompertz ageing rate and R balance out the choice of t0 which should not alter the dynamics of (eq3). This framework could also be extended to growing populations by considering the equation $d^2U/dt^2 + b\ dU/dt + c(U) = 0$, where c(*U*) is zero in low levels of *U* such as in growth arrest conditions, but theoretically could be larger than zero, pushing the physiology away from senescence but toward rejuvenation.

References

Stroustrup, N. *et al.* The temporal scaling of Caenorhabditis elegans ageing. *Nature* **530**, 103-+, doi:10.1038/nature16550 (2016).

Gavrilov, L. A. & Gavrilova, N. S. The reliability theory of aging and longevity. *J Theor Biol* **213**, 527-545, doi:10.1006/jtbi.2001.2430 (2001).

Jones, O. R. *et al.* Diversity of ageing across the tree of life. *Nature* **505**, 169-173, doi:10.1038/nature12789 (2014).

Kirkwood, T. B. & Holliday, R. The evolution of ageing and longevity. *Proc R Soc Lond B Biol Sci* **205**, 531-546 (1979).

Williams, G. C. Pleiotropy, Natural-Selection, and the Evolution of Senescence. *Evolution* **11**, 398-411, doi:Doi 10.2307/2406060 (1957).

Partridge, L. & Barton, N. H. Optimality, Mutation and the Evolution of Aging. *Nature* **362**, 305-311, doi:DOI 10.1038/362305a0 (1993).

Stewart, E. J., Madden, R., Paul, G. & Taddei, F. Aging and death in an organism that reproduces by morphologically symmetric division. *PLoS Biol* **3**, e45, doi:10.1371/journal.pbio.0030045 (2005).

Lindner, A. B., Madden, R., Demarez, A., Stewart, E. J. & Taddei, F. Asymmetric segregation of protein aggregates is associated with cellular aging and rejuvenation. *Proc Natl Acad Sci U S A* **105**, 3076-3081, doi:10.1073/pnas.0708931105 (2008).

Wang, P. *et al.* Robust growth of Escherichia coli. *Curr Biol* **20**, 1099-1103, doi:10.1016/j.cub.2010.04.045 (2010).

Gefen, O., Fridman, O., Ronin, I. & Balaban, N. Q. Direct observation of single stationary-phase bacteria reveals a surprisingly long period of constant protein production activity. *Proc Natl Acad Sci U S A* **111**, 556-561, doi:10.1073/pnas.1314114111 (2014).

Nystrom, T. Aging in bacteria. *Curr Opin Microbiol* **5**, 596-601 (2002).

Monod, J. *Recherches sur la croissance des cultures bactériennes.* (Hermann, 1942).

Pirt, S. J. The maintenance energy of bacteria in growing cultures. *Proc R Soc Lond B Biol Sci* **163**, 224-231 (1965).

Ericsson, M., Hanstorp, D., Hagberg, P., Enger, J. & Nystrom, T. Sorting out bacterial viability with optical tweezers. *J Bacteriol* **182**, 5551-5555 (2000).

Gompertz, B. On the Nature of the Function Expressive of the Law of Human Mortality, and on a New Mode of Determining the Value of Life Contingencies. *Philosophical Transactions of the Royal Society of London* **115**, 513-583 (1825).

Kirkwood, T. B. Deciphering death: a commentary on Gompertz (1825) 'On the nature of the function expressive of the law of human mortality, and on a new mode of determining the value of life contingencies'. *Philos Trans R Soc Lond B Biol Sci* **370**, doi:10.1098/rstb.2014.0379 (2015).

Aalen, O., Borgan, O. & Gjessing, H. *Survival and Event History Analysis: A Process Point of View (Statistics for Biology and Health).* (Springer, 2008).

Barbi, E., Lagona, F., Marsili, M., Vaupel, J. W. & Wachter, K. W. The plateau of human mortality: Demography of longevity pioneers. *Science* **360**, 1459-1461, doi:10.1126/science.aat3119 (2018).

Makeham, W. M. On the Law of Mortality and the Construction of Annuity Tables. *The Assurance Magazine, and Journal of the Institute of Actuaries* **8**, 301-310 (1860).

Vaupel, J. W. *et al.* Biodemographic trajectories of longevity. *Science* **280**, 855-860 (1998).

Yashin, A. I., Vaupel, J. W. & Iachine, I. A. A duality in aging: the equivalence of mortality models based on radically different concepts. *Mech Ageing Dev* **74**, 1-14 (1994).

Kirkwood, T. B. & Austad, S. N. Why do we age? *Nature* **408**, 233-238, doi:10.1038/35041682 (2000).

Fontaine, F., Stewart, E. J., Lindner, A. B. & Taddei, F. Mutations in two global regulators lower individual mortality in Escherichia coli. *Mol Microbiol* **67**, 2-14, doi:10.1111/j.1365-2958.2007.05988.x (2008).

Fontana, L., Partridge, L. & Longo, V. D. Extending healthy life span--from yeast to humans. *Science* **328**, 321-326, doi:10.1126/science.1172539 (2010).

Kenyon, C., Chang, J., Gensch, E., Rudner, A. & Tabtiang, R. A C. elegans mutant that lives twice as long as wild type. *Nature* **366**, 461-464, doi:10.1038/366461a0 (1993).

Pedro de Magalhaes, J. *et al.* A Reassessment of Genes Modulating Aging in Mice Using Demographic Measurements of the Rate of Aging. *Genetics* **208**, 1617-1630, doi:10.1534/genetics.118.300821 (2018).

Battesti, A., Majdalani, N. & Gottesman, S. The RpoS-mediated general stress response in Escherichia coli. *Annu Rev Microbiol* **65**, 189-213, doi:10.1146/annurev-micro-090110-102946 (2011).

Peterson, C. N., Levchenko, I., Rabinowitz, J. D., Baker, T. A. & Silhavy, T. J. RpoS proteolysis is controlled directly by ATP levels in Escherichia coli. *Genes Dev* **26**, 548-553, doi:10.1101/gad.183517.111 (2012).

Kuroda, A. *et al.* Role of inorganic polyphosphate in promoting ribosomal protein degradation by the Lon protease in E. coli. *Science* **293**, 705-708, doi:10.1126/science.1061315 (2001).

Lever, M. A. *et al.* Life under extreme energy limitation: a synthesis of laboratory- and field-based investigations. *FEMS Microbiol Rev* **39**, 688-728, doi:10.1093/femsre/fuv020 (2015).

Radzikowski, J. L. *et al.* Bacterial persistence is an active sigmaS stress response to metabolic flux limitation. *Mol Syst Biol* **12**, 882, doi:10.15252/msb.20166998 (2016).

Medawar, P. B. *An Unsolved Problem of Biology: An Inaugural Lecture Delivered at University College, London, 6 December, 1951.* (H.K. Lewis and Company, 1952).

Mauri, M. & Klumpp, S. A model for sigma factor competition in bacterial cells. *PLoS Comput Biol* **10**, e1003845, doi:10.1371/journal.pcbi.1003845 (2014).

Scott, M., Gunderson, C. W., Mateescu, E. M., Zhang, Z. & Hwa, T. Interdependence of cell growth and gene expression: origins and consequences. *Science* **330**, 1099-1102, doi:10.1126/science.1192588 (2010).

Maharjan, R., Seeto, S., Notley-McRobb, L. & Ferenci, T. Clonal adaptive radiation in a constant environment. *Science* **313**, 514-517, doi:10.1126/science.1129865 (2006).

Spira, B., de Almeida Toledo, R., Maharjan, R. P. & Ferenci, T. The uncertain consequences of transferring bacterial strains between laboratories - rpoS instability as an example. *BMC Microbiol* **11**, 248, doi:10.1186/1471-2180-11-248 (2011).

Hol, F. J. *et al.* Spatial structure facilitates cooperation in a social dilemma: empirical evidence from a bacterial community. *PLoS One* **8**, e77042, doi:10.1371/journal.pone.0077042 (2013).

Vulic, M. & Kolter, R. Evolutionary cheating in Escherichia coli stationary phase cultures. *Genetics* **158**, 519-526 (2001).

Hengge, R. in *Bacterial Stress Responses, Second Edition*   (American Society of Microbiology, 2011).

Available resources

C P N

"Feast"

Reproduction

Death

Colonisation

Dispersal

"Famine"

Senescence

Dissipated resources

C P N

Physiological processes

Stochastic events

a

b

c

Time

d

Cell #

1000

2000

3000

4000

0   20   40   60   80   100   120

Time (h)

a

b

a

b

c

Maintenance capacity

U

$U - J\Delta t$

Physiological homeostasis

Molecular damage

Molecular repair

Maintenance energy flux J

Hazard rate h

Death

Free radicals damaging protein, membrane & DNA

→ Physiological processes

⇢ Molecular processes