Fungal-bacterial interaction selects for quorum sensing mutants and a metabolic shift

towards the production of natural antifungal compounds

Andrea G. Albarracín Orio[1,2,3], Daniel Petras[3], Romina A. Tobares[2], Alexander Aksenov[3], Mingxun Wang[3],

Florencia Juncosa[1], Pamela Sayago[1], Alejandro Moyano[2], Pieter C. Dorrestein[3], Andrea M. Smania[2] and

Daniel A. Ducasse[1,4]

[1]IRNASUS, Universidad Católica de Córdoba, CONICET, Facultad de Ciencias Agropecuarias. Av. Armada

Argentina 3555 (X5016DHK) Córdoba, Argentina. [2]Universidad Nacional de Córdoba. Facultad de Ciencias

Químicas. Departamento de Química Biológica Ranwel Caputto. Córdoba, Argentina. CONICET.

Universidad Nacional de Córdoba. Centro de Investigaciones en Química Biológica de Córdoba

(CIQUIBIC), Córdoba, Argentina. [3]Collaborative Mass Spectrometry Innovation Center, 9500 Gilman Drive,

University of California, San Diego, CA 92093, USA. [4]Instituto de Patología Vegetal IPAVE-CIAP, Instituto

Nacional de Tecnología Agropecuaria, Camino 60 Cuadras Km 5[1/2] (5019) Córdoba, Argentina.

To whom correspondence may be addressed. Email: asmania@fcq.unc.edu.ar,

a.albarracinorio@conicet.gov.ar.

Abstract

Metabolic exchange may define adaptation to the changes imposed by microbial interactions. We found

that antifungal activity of cell-free supernatants of *Bacillus subtilis* ALBA01 is acquired after interacting with

the soil fungus *Setophoma terrestris* (ST). Here we evaluated different traits of *B. subtilis* ALBA01 before

(pre-ST) and after (post-ST) interacting with the fungus in co-cultures and inquired about the molecular

mechanisms underlying this inter-kingdom interaction dependent antagonistic activity. Apart from the high

fungal inhibition activity, we observed that the ability to form robust biofilms *in vitro* was a prevailing feature

of post-ST and that biofilm formation was positively correlated to biocontrol efficacy. Moreover, nuclear

magnetic resonance and mass spectrometry analysis revealed a differential metabolomics profile in post-

ST variants, including the unexpected absence of surfactin or plipastatin production. Disruption of the *srfAA*

gene implicated in surfactin synthesis resulted in an abolition of the ST-driven antifungal activity and ST

growth inhibition levels similar to those observed for post-ST variants. The post-ST phenotype was stable

after several passages on solid media, indicating that a mutation-based process underlies this phenotypic

adaptation process. In fact, whole genome comparison revealed mutations that converge in genes of the

ComQPXA quorum sensing system in post-ST variants. Altogether, our results suggest a role for a still

unknown quorum sensing regulated pathway, which by classical antimicrobial lipopeptide abrogation

mediates *B. subtilis* ALBA01 adaptive phenotypic variation and that may be responsible for the antagonistic

behavior it exerts on *S. terrestris*.

Introduction

The immense microbial diversity within the soil leads to extensive interspecies interactions (1). Generation

of antagonistic and mutualistic behaviors, mediated by exchange of small diffusible secondary metabolites,

facilitates microbial adaptation to complex communal life. Microorganism co-cultures are inspired by the

natural microbe communities that are omnipresent in nature. It has been shown that microorganism co-

culturing can stimulate silent gene clusters, but the molecular mechanisms that can prompt this

phenomenon remain unknown. It has been reported that co-culturing *Aspergillus nidulans* with soil-dwelling

actinobacteria result in the expression and production of specialized fungal metabolites (2). Moreover,

co-culturing of *Streptomyces lividans* TK23 with *Tsukamurella pulmonis* TP-B0596 induce the production

of a still unidentified red pigment, while *S. endus* S-522 co-cultured with *T. pulmonis* TP-B0596 produces

the novel antibiotic alchivemycin A65 (3). The compounds secreted by microbes in response to species

interactions can frequently antagonize other microorganisms, have roles as intraspecific signaling cues,

alter general microbial development and/or their confinement in particular environmental niches (4, 5).

Bacillus subtilis, responds to distinct signals by differentiating into specialized cell types (6), which are

related   to   its   survival   in   the   environment.   Indeed,   by   responding   to   quorum   sensing

molecules, *B. subtilis* can undergo several developmental processes such as formation of biofilms, within

which a subpopulation of cells differentiates to produce an extracellular matrix that encases the community

(7, 8). As a relevant molecule in *B. subtilis*, surfactin is a lipopetide biosurfactant and antibiotic that displays

hemolytic, antiviral, antimycoplasma and antibacterial activities, for example against the plant pathogen

Pseudomonas syringae (9, 10). It has been reported that production of surfactin is essential for biofilm

formation and general medium/niche colonization capacity in *B. subtilis* (11-13). The production of surfactin

is quorum-sensing dependent and is activated by the end of the exponential growth phase (14).

Transcription of *srfA* operon, which encodes surfactin production genes, is directly activated by the

phosphorylated form of the master regulator ComA, which is part of the ComQXPA quorum-sensing system

(14-16). In this system, the ComX autoinducer, encoded by the *comX* gene and post-translationally cleaved

and prenylated by ComQ, binds to and activates ComP, a membrane-anchored histidine kinase receptor,

which then phosphorylates the response regulator ComA that finally activates transcription of more than

one hundred genes (17, 18). Modifications in the regulation of *srfA* operon directly affect the expression of

the competence regulatory gene *comS*, which resides within the same operon, and thus alter competence,

sporulation and motility regulation pathways (19). In fact, ComS causes the release of the factor ComK that

also activates the transcription of several genes involved in DNA cleavage, transformability, motility,

arginine biosynthesis and transport (20). On the other hand, it has also been reported that ComA activates

the expression of *degQ* gene in undomesticated *B. subtilis* strains, an essential gene for the synthesis of

another non-ribosomal peptide, plipastatin (21, 22). Plipastatin, a member of the fengycin family of

lipopetides, is a powerful antifungal compound with a specific and strong activity against filamentous fungi

(23-26). All these evidences highlight the complex and refined surfactin-mediated connection among

development pathways in *B. subtilis*. It has been reported that cellular differentiation and biofilm formation

in *B. subtilis* can be affected not only by intraspecific but also by interspecific signals and that metabolites

secreted by other soil microbes different from *Bacillus spp.* might alter biofilm development in *B. subtilis*

(27). For example, nystatin, amphotericin, valinomycin and gramicidin can all induce the quorum-sensing

pathway and cause differentiation of the subpopulation of matrix producers (28). The ability to sense small

molecules produced by a range of diverse soil microorganisms suggests a broad mechanism

that *B. subtilis* uses to respond to neighboring communities by altering its development. Thus, it is important

to complement our current knowledge on the interspecies communication strategies of rhizosphere

B. subtilis because systems involved in development programs can regulate the expression of anti-

pathogenic and biocontrol agents like surfactins and biofilm-inducing compounds (29).

We have previously shown that *Bacillus subtilis* ALBA01, a strain isolated from the rhizosphere of onion

plants, is capable of strongly inhibit the growth of *S. terrestris in vitro* (30), one of the most severe pathogens

of onion and the causative agent of its major limiting diseases, the pink root. Interestingly, we found that

BsA01 acquires a strong antagonistic behavior against ST after previous interaction with the fungus in co-

culture. Moreover, we determined that the antifungal inhibition activity is mediated by extracellular diffusible

factors since it is observed in cell-free supernatants of BsA01 previously grown in the presence of the

fungus (post-ST variants) but not in those supernatants obtained from variants grown without any previous

contact with it (pre-ST variants) (30). It is noteworthy that this behavior appeared to be specifically provoked

by the interaction with *S. terrestris* since no antagonistic activity against two other onion pathogens,

Fusarium oxysporum and *F. proliferatum*, was observed (30). In order to gain insights into the underlying

mechanisms responsible for this behavior, we analyzed differentiation processes of the bacterium that lead

to the release of diffusible bacterial inhibitory compounds. In this study, we combined phenotypic

characterization, untargeted LC-MS/MS and NMR-based metabolomics approaches and comparative

genomics to distinguish the differences between the pre and post-ST variants. Our findings suggest that a

mutation-based phenotype variation involving quorum sensing genes arises in BsA01 after co-culturing with

S. terrestris and mediates both a suppression of surfactin and plipastatin production and a metabolomics

profile change that might be the bases of the specific antagonistic behavior and the stability of the

phenomenon.

Results

Bacillus subtilis **post-ST variants undergo phenotypic variation and form robust biofilms**

By co-culturing assays, we found that after 15 days of interaction with the soil fungal pathogen *S. terrestris*

(ST), the strain of *B. subtilis* ALBA01 acquires a strong antagonistic activity mediated by secreted factors,

since fungal growth inhibition is observed in bacterial cell-free supernatants (30) (Figure 1 A and B). These

variants were named as post-ST to distinguish them from those variants grown without previous contact

with the fungus (pre-ST variants) and whose cell-free supernatants do not display antagonistic activity (30)

(Figure 1B). To gain further insight into the *S. terrestris*-driven bacillus behavior, we first performed a

phenotypic characterization of both, the pre-ST and the post-ST variants. As shown in Figure 1C, the post-

ST variants exhibited changes in morphology with an apparent increase in roughness and wrinkly colony

phenotype relative to *B. subtilis* colonies grown without previous contact with the fungus (Figure 1C, panel

4 and 5). A qualitative analysis by electron microscopy of *B. subtilis* after co-culture with *S. terrestris*

revealed a high number of longer cells as well as a reduced number of sporulated cells in comparison to

control pre-ST cells (Figure 1C, panel 6). We then characterized the capacity to form biofilm in both, pre-

ST and post-ST variants. In contrast to general assumption that soil microorganisms might release

compounds that would inhibit rather than stimulate cellular differentiation in *B. subtilis*, we observed that

both pre and post-ST variants were capable of forming floating biofilms (pellicles) characteristically

associated with biofilm formation in LBGM medium (31). However, the pre-ST variants formed a more

featureless biofilm with slightly thinner pellicles that easily disintegrate (Figure 2A). In contrast, the post-ST

variants were able to develop into more wrinkled and thicker pellicles, which is consistent with a robust

biofilm formation (Figure 2A). We also evaluated the ability of the *B. subtilis* variants to form submerged

biofilms on borosilicate surfaces and quantified biofilm formation by crystal violet staining. As shown in

Figure 2B, biofilm formation was significantly increased in post-ST variants. Moreover, post-ST lacked

swarming motility when spotted in semisolid LB medium, while pre-ST colonized the entire Petri dish

showing full swarming motility (Figure 2C). Remarkably, when re-inoculated onto a fresh growth medium

(passaging) the post-ST variants still maintained not only the structured colony morphology but also the

increased capacity of biofilm formation, indicating that *B. subtilis* experiences a genetic stable phenotypic

variation after co-culture with *S. terrestris*.

The interaction between *S. terrestris* and *B. subtilis* ALBA01 results in differential secondary

metabolite profile in post-ST variants

To gain deeper understanding on the differential antifungal activity observed in the cell-free supernatants

of the post-ST variants, we applied an untargeted metabolomics profiling based on Proton Nuclear Magnetic

Resonance (^{1}H-NMR) and multivariate data analysis tools to characterize the difference in secondary

metabolite profile of cell-free supernatants of the BsA01 pre-ST and post-ST variants. Twelve samples of

cell-free bacterial supernatants were analyzed, six of which belonged to the pre-ST variant group and the

other six to the post-ST variant group. Principal Component Analysis (PCA) was applied to the data

obtained from NOESY spectra of each sample. As shown in Figure 3A, samples from the pre-ST group

clustered together and clearly separated from post-ST variants samples. To discard potential biases in

sample separation as a consequence of the sample run order, the samples were analyzed by varying the

initial order and we found the same grouping profile as observed before (Figure S1a). Moreover, by

Orthogonal partial least squares discriminative analysis (OPLS-DA) we could maximize the separation

between both groups (Figure S1b). To visualize the differences between classes, we generated S-line plots

in OPLS-DA models and observed different signals that represent discriminant metabolites between both

BsA01 pre-ST and post-ST groups of samples (Figure 3B), all or some of which might constitute the factors

responsible for the antagonistic antifungal behavior.

We then compared the chemotypes of whole cells of BsA01 pre and post-ST variants to test whether the

metabolome is significantly altered between both variants. We carried out a non-targeted liquid

chromatography tandem mass spectrometry (LC-MS/MS) based metabolomics analysis and observed a

clear separation in the global metabolome between the variants in a Principal Coordinate Analysis (PCoA)

(Figure 4A). According to random forest analysis, lipopetides surfactin and plipastatin appeared to be the

most important variables that determine the difference between pre- and post-ST variants (Figure 4B).

Intriguingly, the post-ST variants showed a marked reduction in the intensity of surfactin ions (m/z 1008.66;

1022.67; 1036.69; 1044.65; 1058.67) and also a decrease in the intensity of peaks with masses indicative

of the plipastatin (Figure 4C and D, respectively), both of which were identified through spectrum library

matching and spectral networking (32, 33). Moreover, the post-ST variants clustered together with a *B.*

subtilis NCBI 3610 surfactin-defective mutant, highlighting the role of surfactin in the separation between

the groups (Figure 4A). Figure S2 shows the molecular network obtained by analysis of the chemical profile

of the BsA01 variants. The pie charts in the network indicate relative abundances of ion features between

the different groups. The molecular networks indicate that not only the known surfactin and plipastatin

derivatives were diminished in post-ST variants but also other related unknown derivatives. In a second

approach, we applied LC-MS/MS based metabolomics to analyze cell-free supernatants of the BsA01 pre

and post-ST variants. In accordance with the results found for the whole bacterial cell analysis, we observed

a clear discrimination according to the origin of the cell-free supernatants with PcoA plots showing that

post-ST samples clustered separated from pre-ST variants (Figure 5A). Moreover, for post-ST cell-free

supernatants we could not detect surfactin (Figure 5B) nor plipastatin (Figure 5C) ions in levels comparable

to those found in the pre-ST variant of *B. subtilis* ALBA01 strain. Consistent with this, we also found that

post-ST variants did not show hemolytic activity (Figure 5D). Taking together, these results indicate that

what separates the post-ST variants most is the absence/decrease of the antimicrobial compounds

surfactin and plipastatin. In this sense, the abrogation of surfactin production in the post-ST variants is even

most interesting since this lipopeptide can be considered a key player in environmental adaptation

processes, quorum-sensing and biofilm formation in *B. subtilis.*

Biofilm is positively correlated with the antifungal activity of cell-free supernatants

So far, our findings suggest that the biofilm formation and the absence of the lipopetide surfactin have a

role in the antagonistic behavior of the post-ST variants. Biofilm formation depends on the synthesis of

extracellular matrix, whose production is regulated by two major operons: the *epsA-epsO* and the *tapA*,

both under the control of the transcriptional repressor and biofilm master regulator SinR (34, 35). On the

other hand, surfactin, which is involved in swarming motility and also acts as an antimicrobial metabolite (9,

36), has been identified as a communication signal that triggers biofilm formation (6, 28). To test if key

genes involved in biofilm formation and surfactin production are related to the antifungal activity acquired

by post-ST variants, we generated two BsA01 mutants: a hyper robust biofilm-forming Δ*sinR* and a

surfactin-defective producer Δ*srfAA*. The activity of the BsA01 mutants against *S. terrestris* was screened

by 15-day co-cultures on PDA dishes before and after which samples of each mutant were recovered and

tested. When cell-free supernatants of Δ*sinR* pre-ST mutant were examined for antifungal activity, although

lower than BsA01 pre-ST variant, it displayed a significantly higher activity than cell-free supernatants

obtained from Δ*sinR* post-ST cells (Figure 6). While this result suggest that there is a positive correlation

between biofilm formation and antagonism, the mere activation of the biofilm pathway cannot completely

explain the ST-driven behavior of BsA01. On the contrary, antifungal activity differences between pre-ST

and post-ST supernatants disappeared when the Δ*srfAA* surfactin-less mutant was analysed (Figure 6).

Moreover, the absence of surfactin *per se* led to the acquisition of an equalled mycelium growth inhibitory

activity with respect to the BsA01 post-ST variants that previously interacted with the fungus (Figure 6).

The inability to produce and release surfactin not only resulted in an enhanced level of antagonism towards

ST but also in a total suppression of the ST-driven phenomenon.

B. subtilis post-ST variant undergoes a mutation-based phenotypic variation

To gain insight into the mechanisms underlying the specific and stable phenotypic variation in *B. subtilis*

ALBA01 directed by the interaction with *S. terrestris*, we decided to sequence the whole genome of three

different post-ST variants (post-ST1, post-ST2 and post-ST3), which were independently derived from the

ALBA01 strain that was taken as a pre-ST ancestor and used as a reference. Reads from each post-ST

variant were aligned against the ALBA01 genome (NCBI Bioproject PRJNA316980) to assess the genetic

changes accumulated in post-ST during the interaction with the fungus. Overall the complete genomes, we

found between one and three different mutations in coding sequence regions for post-ST variants (Table

1). Surprisingly, each one of them carried at least one mutation in genes of the ComQXPA quorum-sensing

system. Among these, we found that post-ST 1 and post-ST 2 variants hold a mutation in the coding

sequence of the *comA* gene, whereas the remaining post-ST 3 variant carried a mutation in the kinase

encoding gene, *comP.* For both post-ST 1 and 2 variants, mutations in *comA* resulted in an insertion of one

A in a homopolymeric tract of six adenines at position 415 of the gene ($+1A_{415}$), causing a frameshift

mutation and the generation of a stop codon in position 157 of the protein sequence. Structurally, ComA

belongs to the NarL family of transcriptional activators with a C-terminal DNA-binding domain that shares

sequence homology with the quorum sensing regulator LuxR of *Vibrio fischeri*, and is a member of the

widespread tetrahelical helix-turn-helix (HTH) subfamily (17). The N-terminal domain, which comprises the

response regulator receiver motif (sites 3 to 121), includes the phosphorylation site D55, and the loop linker

sites with the residue T141. The C-terminal region contains the Helix-Turn-Helix motif (sites 147 to 212)

and within this the DNA binding domain (171 to 190). According to prediction analysis by InterPro (37), the

$+1A_{415}$ mutation may lead to a truncated version of the protein that lost the expected DNA-binding domain.

On the other hand, the mutation in the *comP* gene in post-ST3 variant consisted of a 100 bp-deletion at

position 632 of the gene and is predicted to cause the loss of the complete histidine kinase domain of the

protein. It is important to mention that the presence of these mutations was confirmed by PCR amplification

of *comA* and *comP* followed by direct sequencing. We then aimed to explore the involvement of *com*

mutations in the post-ST phenotype and to inquire to what extent *com* genes could be mutated in BsA01

after interacting with *S. terrestris*. For that, we conducted a whole-genome sequencing of a pool of 15 *B.*

subtilis post-ST variants independently obtained after interaction with the fungus in co-culture (Pool-seq).

To identify allelic variations in the pool, we first mapped the sequence reads to the reference genome (*B.*

subtilis ALBA01, NCBI Bioproject PRJNA316980) and then performed the variant calling (see Material and

Methods). Surprisingly, we identified six different mutations in the *comQXPA* operon within the population

of post-ST variants, revealing a marked convergence of mutation within these genes that might explain the

metabolomics changes observed. Remarkably, we found the previously described *comA* insertional

mutation and also three new mutations in *comA*, two substitutions, T215A and C601T, and one 5

nucleotide-insertion at position 299 (Table 2). The T215A mutation generated a premature stop codon at

position 126 of ComA whereas the C601T and the 5 nucleotide-insertion generated stop codons at position

201 and 78 of the protein, respectively. Moreover, two new mutations in the *comP* gene were observed,

and both of them represented insertions. The GC insertion at position 1517 of the gene generated a

truncated version of ComP with a stop codon at postion 512, and the insertion of AT produced a stop codon

at position 232 of ComP (Table 2). These findings present evidence of a mutation-based phenotypic change

in the post-ST variants and support the hypothesis that the environmental conditions that *B. subtilis* ALBA01

encountered when interacting with *S. terrestris* may force an adaptive process that allows the bacterium to

resist and outperform the fungus and that involved a not yet fully understood pathway controlled by

ComQXPA.

Discussion

The soil-dwelling bacterium *Bacillus subtilis* and closely related species have been extensively used as

biological control agents in agriculture, mainly due to their production of antibiotics with a great variety of

chemical structures (10). In contrast, relatively little is known about the molecular mechanisms underlying

the *B. subtilis* biocontrol behavior. A typical mechanism of competition among microorganism is the

secretion of metabolites that produce diverse effects during intra-, inter-species and even inter-kingdom

interactions. In this sense, *B. subtilis*, secretes numerous secondary metabolites including the antibiotic

compound subtilin, the bacteriocin subtilosin, lipopeptides as surfactins, plipastatins, and others, and the

polyketide/nonribosomal peptide hybrid compound bacillaene (25). Many of these metabolites play dual

roles as antimicrobial compounds and signaling molecules during interactions, participating in development

modulation, biofilm formation and inhibition of virulence factors of competitor individuals (4, 26, 38). In a

previous work, we isolated a strain of *B. subtilis* subsp. *subtilis* from the rhizosphere of onion plants, BsA01,

that is capable of strongly inhibiting the growth of the soil-borne fungal pathogen *S. terrestris in vitro* (30).

Remarkably, the antagonistic behavior of BsA01 is manifested in cell-free supernatants of cells previously

grown in the presence of the fungus (post-ST variants) but not in cells never exposed to the fungus (pre-

ST variants), indicating that the interaction with *S. terrestris* promotes a bacterial transformation that leads

to the secretion of antifungal factors. In this work, we aimed to understand this phenomenon through an

exhaustive characterization of BsA01 post-ST variants that included phenotypic, genetic, genomics and

metabolomics analysis. We present evidences that after 15 days of co-culture with *S. terrestris*, BsA01

exhibits changes in colony morphology and acquires the ability to form robust biofilms. In fact, *B. subtilis*

post-ST variants display a remarkable increase in colony roughness/wrinkles and highly structured biofilm

pellicles in the air-liquid medium interphase. We also observed that post-ST variants show neither hemolytic

activity nor swarming motility indicating an absence of surfactin or the impossibility of producing this

lipopeptide. In agreement with these findings, we observed that both a BsA01 hyper biofilm-forming Δ*sinR*

mutant and a surfactin-deficient Δ*srfAA* mutant display higher antifungal activity when compared to the pre-

ST variant. Is noteworthy that the sole absence of surfactin in the Δ*srfAA* mutant was capable of erasing

the differences in activity that pre and post-ST cell-free supernatants exhibited toward *S. terrestris*.

Furthermore, LC-MS/MS-based metabolomics expose major differences in the metabolomes of pre and

post-ST variants, with post-ST showing a marked reduction or absence of surfactin ions. Interestingly, PcoA

analysis showed that post-ST variants clustered both together and with a *B. subtilis* NCBI 3610 knock-out

surfactin derivative mutant suggesting that key differences between the BsA01 variants might rely on the

synthesis of surfactin. Surfactin is a heptapeptide linked to a fatty acid via a cyclic lactone ring structure

that is capable of inserting itself into cellular membranes where interacts with its lipid moiety leading to the

loss of membrane barrier properties and eventually resulting in its complete disintegration by a detergent-

type mechanism (39). It has been widely reported that surfactin displays a significant antimicrobial activity

(9, 40). However, we found that the BsA01 post-ST variants show enhanced activity against the onion

pathogen *S. terrestris* but the biocontrol properties of the cell-free supernatant seemed to be independent

of surfactin in our study model. Most striking, *B. subtilis* appeared to have suppressed or reduced to a

minimum level the production of this classical antimicrobial lipopeptide after interacting with the fungus.

One could think that the lack of surfactin is a mere consequence of the metabolic change that arise along

with the phenotype variation experienced by BsA01 and that is not directly related to the antagonism *per*

se. However, it has been suggested that surfactins could interfere with the activity of other lipopeptides with

proven antifungal activity such as fengycins/plipastatins and iturins (38, 41). In a study, *B.*
amyloliquefaciens FZB42 mutants selectively repressed in the synthesis of one or two types of lipopeptides
revealed that the co-production of surfactin and fengycin lowered the global antifungal activity (38). Such
an effect of mixed lipopeptides was previously observed against the fungus *Verticillium dahliae*
(42). Moreover, a *Bacillus subtilis* subsp. *krictiensis* surfactin-less mutant showed significantly higher
antifungal activity against *Fusarium oxysporum* than its wild-type strain (41). Thus, we speculate that by
reducing or preventing the biosynthesis of surfactin, BsA01 could avoid some negative impact on fitness
while making an effective use of other inhibitory compounds against *S. terrestris.* However, we also found
that of the main differences between pre and post-ST variants is a strong reduction in plipastatin abundance
in the post-ST variants. Plipastatins are lipo-decapeptides with an internal lactone ring and with a β-hydroxy
fatty acid chain (C14- C18) (43). Lipopeptides of this family are less hemolytic than surfactins but have a
strong antifungal activity (43). Among the *B. subtilis* antibiome, cyclic lipopetides of surfactin and
fengycin/plipastatin families are widely recognized as the main antimicrobials that can be secreted in
biologically relevant amounts (44, 45). So, the striking reduction in the production of these two classical
compounds in post-ST variants raises the question of what the essential antifungal factor that BsA01
produces to fight against *S. terrestris* is.

After repeated passages onto fresh growth media, the post-ST variants still maintained not only their colony
morphology but also their increased ability to form biofilm and the absence of surfactin. This pointed out a
possible participation of biofilm and/or surfactin-related pathways in the phenotype variation that
characterize the BsA01 post-ST variants. Moreover, this observation indicates that *B. subtilis* ALBA01
undergoes a heritable phenotypic variation after interacting with the fungus in co-culture. Comparative
whole genome as well as genomic pool-seq analysis between the BsA01 strain (pre-ST variant) and
independently derived post-ST variants revealed mutations in the coding regions of the *comA* and *comP*
genes, which are part of the quorum sensing system ComQXPA. In response to certain environmental
stresses, the regulator ComA is activated by the kinase ComP, which is required for the transcription of
several genes involved in competence development, antibiotic production, exopolysaccharide and
degradative enzyme production as well as transport and fatty acid metabolism (17, 18). Among those
genes, ComA activates the transcription of the *srfA* operon responsible for surfactin production (15, 46).

Mutations found in *comA* gene in post-ST variants all consist of frameshift and non-sense mutations, which

result in the generation of premature stop codons and render truncated versions of the protein that losses

the predicted DNA-binding domain. It is then expected that this alteration produces the loss-of-function of

ComA and as a consequence the decrease or loss of surfactin-related genes expression. The same could

be speculated for the mutations found in the coding sequence of *comP*, which result in the loss of the

complete histidine kinase domain responsible of activating ComA. These observations highlight the

importance of the ComQXPA system in the evolutionary adaptation that BsA01 undergoes during the

interaction with *S. terrestris in vitro*.

It has been described that *Bacillus subtilis* 916 strain was an effective biocontrol agent against the rice

pathogen *Rhizoctonia solani* and that a mutant version of this strain, artificially obtained by means of low-

energy ion implantation, carries three missense mutations within the *comA* gene that the authors discuss

to be the cause of the lower levels of surfactin produced by the mutant. However, in this work in opposition

to what we observe, the reduction in surfactin production leads to a non-optimal performance of *B. subtilis*

916 against the fungus (47). *Bacillus subtilis* has an average of 4-5% of its genome devoted to antibiotic

synthesis and has the potential to produce more than two dozens of structurally diverse antimicrobial

compounds. More than 10% of its genome is controlled by the sophisticated ComQXPA quorum sensing

system centered on the transcriptional regulator, ComA. It has also been found that numerous genes with

unknown function were also affected by ComA (18). As mentioned before, the abrogation of surfactin in

post-ST variants might be explained by a loss-of-function of mutations in the *comA* gene. But, how could

mutations in *comQXPA* genes explain the reduction in the production of plipastatin also observed in the

post-ST variants? ComQXPA system has also been described to be involved in the transcription of the

pleiotropic regulator DegQ, which controls the production of degradative enzymes and is necessary for

plipastatin production (21, 48, 49). Thus, we hypothesize that the reduction in plipastatin might represent

an indirect effect of the mutations in *comA* and *comP* genes through alteration in *degQ* expression.

Interestingly, co-cultures have also been used to explore the modulation of quorum sensing in the search

for new antibiotics since quorum sensing molecules play effective regulatory roles within bacterial

populations by adapting global growth to the level of nutrient availability (50, 51). However, at present no

quorum sensing response mutants were observed in natural populations of *Bacillus subtilis*. Most bacterial

species are confronted with frequent and often extreme changes in environmental conditions to which they

must rapidly adapt. To this end, they operate diverse regulatory networks being one of the most important

involved in quorum sensing control, which in response to particular environmental signals brings global

changes in gene expression that in turn regulate complex physiological behaviors and confer adaptability

and competiveness (52). As of yet, reports have demonstrated neither a mutational evolution of the ComA

protein mediating a stable phenotypic change nor the role of ComQXPA mutations in positive expression

or proper activity/functioning of extracellular antifungal metabolites that could mediate an adaptive response

in *B. subtilis*.

Taken together, our findings allows us to speculate that the mutation found in the ComQXPA system of *B.*

subtilis post-ST variants may be involved in a regulatory mechanism that leads to the activation of a cryptic

biosynthetic pathway, which specifically responds to extracellular signals involved in the interaction

between the *B. subtilis* strain and *S. terrestris* in co-cultures. Future investigations will be necessary to

expand our knowledge on the identity and the role that the antagonistic metabolites play in this iteraction-

driven evolution process.

Materials and Methods

Materials and methods are described in the SI Appendix, including microorganisms growth conditions, *B.*

subtilis mutants construction, phenotyping characterization analysis, whole genome sequencing and pool-

seq analysis and metabolomics studies based on liquid chromatography – tandem mass spectrometry and

nuclear magnetic resonance spectrometry.

Acknowledgments

The authors are grateful to Dr. Alejandro Moyano for valuable discussion and comments on the manuscript.

We would like to thank Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Agencia

Nacional de Promoción Científica y Tecnológica (ANPCyT). This work was supported by by MINCyT |

Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT), grant PICT 2013-2592 to A.G.A.O and

PICT 2012-2088 to A.M.S.; the US National Science Foundation (NSF) Inspire Track II IOS-1343020;

National Institutes of Health (NIH) Grants GMS10RR029121, 5P41GM103484–07 and the Deutsche

Forschungsgemeinscht (DFG) with Grant PE 2600/1.

Author Contributions

A.G.A.O, A.M.S, D.P., D.A.D and P.C.D. designed the experiments and supervised the study. A.G.A.O,

P.S. and F.S. cultivated the strains. A.G.A.O. performed the interaction studies. A.G.A.O, R.A.T. and A.M.S.

performed DNA isolation and analyzed bioinformatics data of bacterial genomes. A.G.A.O. analyzed NMR

data. A.G.A.O. and D.P. collected the mass spectrometry data. A.G.A.O., D.P. and P.C.D. analyzed the

metabolomics data and performed molecular networking. A.G.A.O. and A.M.S. integrated all experimental

data and performed the final interpretation. A.G.A.O., A.M.S., D.P. and D.A.D. wrote the paper. All authors

discussed, edited and agreed on the final manuscript.

References

1.   Kolter R & Greenberg EP (2006) Microbial sciences: the superficial life of microbes. *Nature*

441(7091):300-302.

2.   Schroeckh V*, et al.* (2009) Intimate bacterial–fungal interaction triggers biosynthesis of archetypal

polyketides in Aspergillus nidulans. *Proceedings of the National Academy of Sciences*

106(34):14558-14563.

3.   Onaka H, Mori Y, Igarashi Y, & Furumai T (2011) Mycolic acid-containing bacteria induce natural-

product biosynthesis in *Streptomyces* species. *Applied and environmental microbiology* 77(2):400-

406.

4.   Straight PD & Kolter R (2009) Interspecies Chemical Communication in Bacterial Development.

Annual Review of Microbiology 63(1):99-118.

5.   Ryan RP & Dow JM (2008) Diffusible signals and interspecies communication in bacteria.

Microbiology 154(7):1845-1858.

6.   López D, Vlamakis H, Losick R, & Kolter R (2009) Paracrine signaling in a bacterium. *Genes &*

Development 23(14):1631-1638.

7.   Branda SS, Chu F, Kearns DB, Losick R, & Kolter R (2006) A major protein component of the

Bacillus subtilis biofilm matrix. *Molecular Microbiology* 59(4):1229-1238.

8.   Vlamakis H, Aguilar C, Losick R, & Kolter R (2008) Control of cell fate by the formation of an

architecturally complex bacterial community. *Genes & Development* 22(7):945-953.

9.   Bais HP, Fall R, & Vivanco JM (2004) Biocontrol of *Bacillus subtilis* against Infection of *Arabidopsis*

Roots by *Pseudomonas syringae* Is Facilitated by Biofilm Formation and Surfactin Production. *Plant*

Physiology 134(1):307-319.

10.   Ongena M & Jacques P (2008) *Bacillus* lipopeptides: versatile weapons for plant disease

biocontrol. *Trends in Microbiology* 16(3):115-125.

11.   B. KD & Richard L (2003) Swarming motility in undomesticated *Bacillus subtilis*. *Molecular*

Microbiology 49(3):581-590.

12.   Kinsinger RF, Shirk MC, & Fall R (2003) Rapid surface motility in *Bacillus subtilis* is dependent on

extracellular surfactin and potassium ion. *Journal of Bacteriology* 185(18):5627-5631.

13.   Julkowska D, Obuchowski M, Holland IB, & Séror SJ (2005) Comparative analysis of the

development of swarming communities of *Bacillus subtilis* 168 and a natural wild type: critical

effects of surfactin and the composition of the medium. *Journal of Bacteriology* 187(1):65-76.

14.   Cosby WM, Vollenbroich D, Lee OH, & Zuber P (1998) Altered srf Expression in *Bacillus subtilis*

resulting from changes in culture pH is dependent on the Spo0K oligopeptide permease and the

ComQX System of Extracellular Control. *Journal of Bacteriology* 180(6):1438-1445.

15.   Roggiani M & Dubnau D (1993) ComA, a phosphorylated response regulator protein of *Bacillus*

subtilis, binds to the promoter region of *srfA*. *Journal of Bacteriology* 175(10):3182-3187.

16.   Nakano MM & Zuber P (1991) The primary role of *comA* in establishment of the competent state

in *Bacillus subtilis* is to activate expression of *srfA*. *Journal of Bacteriology* 173(22):7269-7274.

17.   Wolf D*, et al.* (2016) The quorum-sensing regulator ComA from *Bacillus subtilis* activates

transcription using topologically distinct DNA motifs. *Nucleic Acids Research* 44(5):2160-2172.

18.   Comella N & Grossman AD (2005) Conservation of genes and processes controlled by the quorum

response in bacteria: characterization of genes controlled by the quorum‐sensing transcription

factor ComA in *Bacillus subtilis*. *Molecular Microbiology* 57(4):1159-1174.

19.   Hamoen LW, Eshuis H, Jongbloed J, Venema G, & Sinderen D (1995) A small gene, designated

414   *comS*, located within the coding region of the fourth amino acid-activation domain of *srfA*, is

415   required for competence development in Bacillus subtilis. *Molecular Microbiology* 15(1):55-63.

416   20.   Ogura M*, et al.* (2002) Whole-genome analysis of genes regulated by the *Bacillus subtilis*

417   competence transcription factor ComK. *Journal of Bacteriology* 184(9):2344-2351.

418   21.   Tsuge K, Ano T, Hirai M, Nakamura Y, & Shoda M (1999) The Genes *degQ*, *pps*, and *lpa-8* (*sfp*)

419   are responsible for conversion of *Bacillus subtilis* 168 to plipastatin production. *Antimicrobial Agents*

420   *and Chemotherapy* 43(9):2183-2192.

421   22.   Tsuge K, Matsui K, & Itaya M (2007) Production of the non-ribosomal peptide plipastatin in *Bacillus*

422   *subtilis* regulated by three relevant gene blocks assembled in a single movable DNA segment.

423   *Journal of Biotechnology* 129(4):592-603.

424   23.   Vanittanakom N, Loeffler W, Koch U, & Jung G (1986) Fengycin-a novel antifungal lipopeptide

425   antibiotic produced by *Bacillus subtilis* F-29-3. *The Journal of antibiotics* 39(7):888-901.

426   24.   Falardeau J, Wise C, Novitsky L, & Avis T (2013) Ecological and mechanistic insights into the direct

427   and indirect antimicrobial properties of *Bacillus subtilis* lipopeptides on plant pathogens. *Journal of*

428   *chemical ecology* 39(7):869-878.

429   25.   Cawoy H*, et al.* (2015) Lipopeptides as main ingredients for inhibition of fungal phytopathogens by

430   *Bacillus subtilis*/amyloliquefaciens. *Microbial Biotechnology* 8(2):281-295.

431   26.   Tang Q*, et al.* (2014) Effects of fengycin from *Bacillus subtilis* fmbJ on apoptosis and necrosis in

432   *Rhizopus stolonifer*. *Journal of Microbiology* 52(8):675-680.

433   27.   Shank EA*, et al.* (2011) Interspecies interactions that result in *Bacillus subtilis* forming biofilms are

434   mediated mainly by members of its own genus. *Proceedings of the National Academy of Sciences*

435   108(48):E1236–E1243.

436   28.   López D, Fischbach MA, Chu F, Losick R, & Kolter R (2009) Structurally diverse natural products

437   that cause potassium leakage trigger multicellularity in *Bacillus subtilis*. *Proceedings of the National*

438   *Academy of Sciences of the United States of America* 106(1):280-285.

29.   Zeriouh H, de Vicente A, Pérez-García A, & Romero D (2014) Surfactin triggers biofilm formation

of *Bacillus subtilis* in melon phylloplane and contributes to the biocontrol activity. *Environmental*

Microbiology 16(7):2196-2211.

30.   Albarracín Orio AG, Brücher E, & Ducasse DA (2016) A strain of *Bacillus subtilis* subsp. *subtilis*

shows a specific antagonistic activity against the soil-borne pathogen of onion *Setophoma*

terrestris. European Journal of Plant Pathology 144(1):217-223.

31.   Shemesh M & Chai Y (2013) A Combination of Glycerol and Manganese Promotes Biofilm

Formation in *Bacillus subtilis* via Histidine Kinase KinD Signaling. *Journal of Bacteriology*

195(12):2747-2754.

32.   Watrous J*, et al.* (2012) Mass spectral molecular networking of living microbial colonies.

Proceedings of the National Academy of Sciences 109(26):E1743–E1752.

33.   Wang M*, et al.* (2016) Sharing and community curation of mass spectrometry data with Global

Natural Products Social Molecular Networking. *Nature Biotechnology* 34:828.

34.   Nakano MM*, et al.* (1991) *srfA* is an operon required for surfactin production, competence

development, and efficient sporulation in Bacillus subtilis. *Journal of Bacteriology* 173(5):1770-

1778.

35.   Kearns DB, Chu F, Branda SS, Kolter R, & Losick R (2005) A master regulator for biofilm formation

by *Bacillus subtilis. Molecular Microbiology* 55(3):739-749.

36.   Branda SS, González-Pastor JE, Ben-Yehuda S, Losick R, & Kolter R (2001) Fruiting body

formation by *Bacillus subtilis. Proceedings of the National Academy of Sciences* 98(20):11621-

11626.

37.   Finn RD*, et al.* (2017) InterPro in 2017—beyond protein family and domain annotations. *Nucleic*

Acids Research 45(D1):D190-D199.

38.   Zihalirwa Kulimushi P, Argüelles Arias A, Franzil L, Steels S, & Ongena M (2017) Stimulation of

Fengycin-Type Antifungal Lipopeptides in *Bacillus amyloliquefaciens* in the Presence of the Maize

Fungal Pathogen *Rhizomucor variabilis. Frontiers in Microbiology* 8(850).

39.   Carrillo C, Teruel JA, Aranda FJ, & Ortiz A (2003) Molecular mechanism of membrane
466   permeabilization by the peptide antibiotic surfactin. *Biochimica et Biophysica Acta (BBA) -*
467   *Biomembranes* 1611(1):91-97.

468   40.   Fan H*, et al.* (2017) Biocontrol of Bacterial Fruit Blotch by *Bacillus subtilis* 9407 via Surfactin-
469   Mediated Antibacterial Activity and Colonization. *Frontiers in Microbiology* 8(1973).

470   41.   Kim YT*, et al.* (2017) Organization and characterization of genetic regions in *Bacillus subtilis* subsp.
471   *krictiensis* ATCC55079 associated with the biosynthesis of iturin and surfactin compounds. *PLoS*
472   *ONE* 12(12):e0188179.

473   42.   Liu J, Hagberg I, Novitsky L, Hadj-Moussa H, & Avis TJ (2014) Interaction of antimicrobial cyclic
474   lipopeptides from *Bacillus subtilis* influences their effect on spore germination and membrane
475   permeability in fungal plant pathogens. *Fungal Biology* 118(11):855-861.

476   43.   Steller S*, et al.* (1999) Structural and functional organization of the fengycin synthetase
477   multienzyme system from *Bacillus subtilis* b213 and A1/3. *Chemistry & Biology* 6(1):31-41.

478   44.   Kinsella K, P. Schulthess C, Morris T, & D. Stuart J (2009) *Rapid quantification of Bacillus subtilis*
479   *antibiotics in the rhizosphere* pp 374-379.

480   45.   Debois D*, et al.* (2014) Spatiotemporal monitoring of the antibiome secreted by *Bacillus* biofilms on
481   plant roots using MALDI mass spectrometry imaging. *Analytical chemistry* 86(9):4431-4438.

482   46.   Nakano MM, Xia LA, & Zuber P (1991) Transcription initiation region of the *srfA* operon, which is
483   controlled by the comP-comA signal transduction system in *Bacillus subtilis*. *Journal of Bacteriology*
484   173(17):5487-5493.

485   47.   Wang X*, et al.* (2010) Three non-aspartate amino acid mutations in the ComA Response regulator
486   receiver motif severely decrease surfactin production, competence development and spore
487   formation in *Bacillus subtilis*. *Journal of microbiology and biotechnology* 20(2):301-310.

488   48.   Msadek T, Kunst F, Klier A, & Rapoport G (1991) DegS-DegU and ComP-ComA modulator-effector
489   pairs control expression of the *Bacillus subtilis* pleiotropic regulatory gene *degQ*. *Journal of*
490   *Bacteriology* 173(7):2366-2377.

491   49.   Wang P*, et al.* (2015) DegQ regulates the production of fengycins and biofilm formation of the
492   biocontrol agent *Bacillus subtilis* NCD-2. *Microbiological Research* 178:42-50.

493   50.   Bertrand S*, et al.* (2014) Metabolite induction via microorganism co-culture: A potential way to

494   enhance chemical diversity for drug discovery. *Biotechnology Advances* 32(6):1180-1204.

495   51.   Charusanti P*, et al.* (2012) Exploiting Adaptive Laboratory Evolution of *Streptomyces clavuligerus*

496   for Antibiotic Discovery and Overproduction. *PLoS ONE* 7(3):e33727.

497   52.   Miller MB & Bassler BL (2001) Quorum sensing in bacteria. *Annual Reviews in Microbiology*

498   55(1):165-199.

Figure 1. A specific induced phenotype change was observed in _B. subtilis_ upon the interaction with _S. terrestris_.

After 15 days of co-culture with _S. terrestris_ (ST) _in vitro_, a sample of the _B. subtilis_ A01 (BsA01) colony was taken, stored and named post-ST variant to differentiate it from its original version never exposed to ST in co-culture (pre-ST variant). (A) The antifungal activity of cell-free supernatants (SN) of both variants were assayed according to the workflow shown. (B) A high growth inhibition of _S. terrestris_ was observed in dishes containing only SN of post-ST. The figure shows the fungal growth inhibition at 4 days after inoculation. (C)The interaction with the fungus also leads BsA01 to experience a phenotypic change since post-ST variants develop into a rougher and wrinkled colonies in LB agar compared to the pre-ST variant. Panels 2 and 5 show pictures of individual colonies of a pre-ST and a post-ST variant, respectively, and panels 1 and 4 their respective agar plates showing the overall macroscopic appearance of each variant culture. Electron microscopy of _B. subtilis_ post-ST variants exposed an increment in the number of longer cells and a reduction of the number of sporulated cells (panel 6) in contrast to control cells never exposed to the fungus or pre-ST variants (panel 3) (bar scale, 2 μm). Arrows in panels 3 and 6 indicate sporulated cells.

519
520
521
Figure 2. *B. subtilis* post-ST variants form robust biofilm and lacks swarming motility.
(A) *B. subtilis* post-ST variants were able to develop into pellicles characteristically associated with strong
biofilm formation in liquid medium while, surprisingly, the pre-ST variant formed pellicles with diminished
wrinkles. (B) The quantification of biofilm formation in borosilicate surfaces by using crystal violet staining
also revealed an increase in biofilm formation in *B. subtilis* post-ST variants compared to the pre-ST variant.
(C) Swarming motility was assessed in 0.7% agar LB plates. The *B. subtilis* pre-ST variant occupied the
entire plate showing the expected full motility while *B. subtilis* post-ST variants clearly showed a swarming
deficiency as the colony could not develop and extend beyond the region of inoculation.
530

Figure 3. A metabolomics profile variation in *B. subtilis* ALBA01 is observed after the interaction with *S. terrestris in vitro*.

The high antifungal activity found in cell-free supernatants (SN) of post-ST variants appeared to be associated to changes in the metabolomics profile of released compounds since differences in SN between the pre and post-ST variants could be observed by an untargeted NMR approach. (A) The PCA scores plot shows a clear separation of the SN samples according to their type, pre-ST variant (black) and post-ST variant (grey) samples, in the first two principal components derived from the 1H NMR spectra of *Bacillus* conditioned media (percent variation in the NMR data explained by the model, R2x= 70.4%; percent variation in the NMR data predicted by the model from cross validation, Q2x= 57 %, based on a 2 components model). Quality control (QC) samples are shown as white circles. (B) OPLS-DA S-line plots with pairwise comparison of data from NMR spectra obtained from *Bacillus* variants SN. Colors are associated with the correlation of metabolites characterized from the 1H NMR data for the class of interest with the scale shown on the right-hand side of the plot.

Figure 4. LC-MS/MS based metabolomics exposed the chemical signatures that differentiate the *B. subtilis* pre and post-ST variants.

(A) A strong separation between the pre and post-ST variants could be observed in PcoA plots. (B) Random Forest analysis revealed that surfactin and plipastatin were one of the most important variables that determined this separation. The post-ST variants showed a marked reduction in the intensity of surfactin (C) and plipastatin ions (D).

561
562
563
Figure 5. LC-MS/MS metabolomics analysis of the cell-free supernatants of *B. subtilis* pre and post-
ST variants.
(A) PcoA plots show a clear discrimination according to the origin of the cell-free supernatants, with post-
ST samples clustering separated from pre-ST variants. Surfactin (B) and plipastatin (C) could not be
detected in post-ST cell-free supernatants in levels comparable to those found in the pre-ST variants. (D)
Consistent with the fact that apparently the post-ST variants have lost the ability to produce surfactin, they
did not show the typical hemolytic activity associated to this lipopeptide.

576
577
Figure 6. Biofilm correlates with the antagonism and surfactin is not the metabolite responsible of
the induced antifungal activity.
The activity of cell-free supernatants of BsA01 Δ*sinR* and Δ*srfAA* mutants, pre- and post-contact with *S.*
terrestris, was screened in co-culture as described in the text. Graph columns represent mean values of
mycelium growth. The experiments were independently repeated at least four times. * indicates that the
mycelium growth is significantly different between the strains tested and ns denotes no significant
differences (Tukey's Multiple Comparison Test, *p*-value < 0.0001). The antifungal activity of the cell-free
supernatants of the surfactin deficient Δ*srfAA* mutant pre- and post-contact with the fungus were similar but
lower than those exhibited by the BsA01 post-ST variants and the Δ*sinR* mutant indicating that the inability
to produce surfactin cannot completely abrogate the antifungal activity of *B. subtilis* ALBA01.
588

Figure 7. Schematic representation of the interaction between *S. terrestris* and *B. subtilis* and its influence on the cellular processes implied in the mutation-based phenotypic variation experienced by the bacterium. The phenotypic variation observed in *B. subtilis* after interacting with the fungus manifests an enhanced antifungal activity in cell-free supernatants, biofilm formation and deficient swarming motility. Metabolic exchange and the chemical scenario defined during the confrontation of the two individuals (fungus and bacterium) drive a mutational event and selection of *B. subtilis* cells that carry truncated versions of either the master regulator ComA and/or the kinase ComP (represented by asterisks). The absence of the activated forms of this two key protein components of the ComQXPA quorum-sensing system results in a lack of activation of the expression of ComA-controlled genes among which are the surfactin-encoded operon and *degQ*. Transcription of *degQ* is necessary for the production of the non-ribosomal lipopeptide, plipastatin. Besides, DegQ positively regulates motility-associated genes through modulation of the activity of the pleiotropic factor DegU. Surfactin has been described as a signaling molecule that induces the activation of the SinI-SinR pathway causing extracellular matrix production and biofilm formation. The effect of its abrogation might be overcome by the action of some particular chemical signal, produced during the interaction, which is able to either activate the phosphorilation of Spo0A, and thus matrix gene expression, or act via a still unknown mechanism that finally results in the biofilm formation characteristically observed in *B. subtilis* after interacting with *S. terrestris*. The physicochemical conditions that arose during the interaction may contribute to the adaptation of *B. subtilis* and the improvement of its antagonistic behavior towards *S. terrestris*.

613   **Tables**

614   **Table 1.** Whole genome sequencing comparative analysis of *B. subtilis* post-ST variants revealed mutations

615   in *comQXPA* coding regions

Variant	Scaffold/position	Gene	Nucleotide substitution	Description
Post-ST 1	1/69978	*comA*	Ins A 415	Transcriptional regulatory protein ComA
Post-ST 2	1/69978	*comA*	Ins A 415	Transcriptional regulatory protein ComA
Post-ST 2	2/165291	*yhfV*	ΔA 1089	Methyl-accepting chemotaxis protein
Post-ST 2	3/371952	*spo0A*	G289C	Sporulation two-component response regulator
Post-ST 3	1/71309	*comP*	Δ632-731	Two-component sensor kinase ComP

618   **Table 2.** Mutations found in *comPQXA* operon by whole-genome sequencing of pools of individuals

619   experiment (Pool-seq)

| | | | | 621 |
Scaffold (ALBA01)	Position	Gene	Nucleotide Substitution	622
1	69979	*comA*	C601T	
1	69978	*comA*	Ins A 415	
1	70101	*comA*	Ins CTGGG 299	
1	70185	*comA*	T215A	
1	71264	*comP*	Ins CG 1517	
1	71919	*comP*	Ins AT 863	