Extreme genetic structure and dynamic range evolution in a montane passerine bird: implications for tropical diversification

Carlos Daniel Cadena[1][*], Jorge L. Pérez-Emán[2,3], Andrés M. Cuervo[4,5] Laura N. Céspedes[1], Kevin L. Epperly[6] & John T. Klicka[6]

[1]Laboratorio de Biología Evolutiva de Vertebrados, Departamento de Ciencias Biológicas, Universidad de Los Andes, Bogotá, Colombia

[2]Instituto de Zoología y Ecología Tropical, Universidad Central de Venezuela, Caracas, Venezuela

[3]Colección Ornitológica Phelps, Caracas, Venezuela

[4]Louisiana State University Museum of Natural Science

[5]Instituto de Investigación en Recursos Biológicos Alexander von Humboldt

[6]Department of Biology and Burke Museum of Natural History and Culture, University of Washington, Seattle

[*]Corresponding author. Email: ccadena@uniandes.edu.co

Abstract

18 **Aim**: Employ phylogeographic analyses of a widespread species complex to examine the role of historical and
19 evolutionary processes in the origin and maintenance of high species diversity in the Neotropical montane region.
20
21 **Location**: Neotropical highlands.
22
23 **Taxon**: *Henicorhina* wood-wrens (Aves, Troglodytidae).
24
25 **Methods:** We collected mtDNA sequence data for 288 individuals thoroughly covering the range of the *Henicorhina*
26 *leucophrys* complex from Mexico to Bolivia. Sequences were employed to characterize population structure, infer
27 phylogenetic relationships among populations and their divergence times, examine lineage accumulation through time,
28 and identify presumptive species using coalescent methods. We also explored the origin of elevational and latitudinal
29 replacements involved in spatial changes in species assemblages in the Andes.
30
Results: We found remarkable genetic structure within the complex, which consists of numerous lineages reaching
>12% sequence divergence; most divergent populations occur in areas separated by topographic barriers but several of
them, typically not sister to each other, co-occur with elevational segregation on mountain slopes or replace each other
with latitude along the Andes. Some close relatives occur in areas separated by thousands of kilometers, with more
distant relatives occupying intervening areas. The complex likely originated in the Mexican highlands and expanded
extensively in South America while diverging rapidly at a constant rate into many different lineages which have
persisted for millions of years. Coalescent analyses consistently revealed that the complex may comprise more than 30
species; while we do not suggest these presumptive species should be recognized by taxonomists in the absence of
additional data, *H. leucophrys* is a distant outlier among New World birds in terms of high lineage diversity within a
single recognized species.
41
Main Conclusions: Our study captured wood-wren lineages in the act of building up diversity via divergence and
persistence in allopatry, achievement of secondary sympatry, and coexistence at the landscape scale mediated by
ecological and evolutionary divergence. Although dispersal by wood-wrens is restricted at present and this likely
accounts for strong population structure across topographic barriers, their ranges have been dynamic, managing to
disperse over much of the montane Neotropics. Phases of expansion and contraction of ranges and localized extinctions
of populations likely account for phylogeographic patterns which are precursors to the origin of new species and the
accumulation of diversity in tropical mountains.
49
50 **Key words:** Andes, elevational replacement, range expansion, speciation

Introduction

Mountains in the tropics contribute disproportionately to regional species richness given their area in comparison to lowlands, and are often considered global hotspots of biological diversity and endemism (Stattersfield *et al.*, 1997; Orme *et al.*, 2005; Fjeldså *et al.*, 2012). Tropical mountains exhibit particularly high beta diversity (i.e. species turnover in space) because (1) species assemblages shift along elevational gradients, and (2) related species occupy similar elevations in different mountains or in sectors of a mountain separated by geographic barriers. While knowledge of the biodiversity of mountains has advanced conceptually (Graham *et al.*, 2014; Bertuzzo *et al.*, 2016; Badgley *et al.*, 2017) and empirically (e.g., Patterson *et al.*, 1998; Jankowski *et al.*, 2009; McCain, 2009; Price *et al.*, 2014; Peters *et al.*, 2016; Quintero & Jetz, 2018), accounting for species richness in montane systems remains difficult. In particular, although climate and available energy have an imprint globally on the distribution of life (Francis & Currie, 2003; Hawkins *et al.*, 2003), they cannot predict the agglomeration of range-restricted species in tropical mountains (Rahbek & Graves, 2001; Jetz & Rahbek, 2002; Rahbek *et al.*, 2007; but see Ruggiero & Hawkins, 2008). Because the high richness and uniqueness of mountains in the tropics may instead reflect high speciation rates or low extinction rates (Jetz *et al.*, 2004; Badgley *et al.*, 2017), considering evolutionary processes is crucial to a better understanding of montane diversity (Graham *et al.*, 2014; Laiolo *et al.*, 2018; Quintero & Jetz, 2018).

Dozens of studies have used phylogenetic and population genetic perspectives to probe into evolutionary processes underlying patterns of avian diversity in the Neotropical mountains. Birds have diversified rapidly in the Andes, with pivotal roles of features of the landscape (e.g., low-lying valleys, high-elevation passes) and of climatic changes as drivers of divergence (Pérez-Emán, 2005; Weir, 2006; Cadena *et al.*, 2007; Ribas *et al.*, 2007; Sedano & Burns, 2010; Chaves *et al.*, 2011; Gutiérrez-Pinto *et al.*, 2012; Valderrama *et al.*, 2014; Benham *et al.*, 2015; Sánchez-González *et al.*, 2015; Winger & Bates, 2015; Prieto-Torres *et al.*, 2018). Allopatric differentiation of lineages separated by barriers to dispersal is predominant (Hazzi *et al.*, 2018), whereas evidence for speciation in parapatry along mountain slopes remains elusive (Patton & Smith, 1992; García-Moreno & Fjeldså, 2000; Cadena *et al.*, 2012; Caro *et al.*, 2013). Thus, the replacement of closely related species along elevational gradients, a salient geographic pattern in tropical avifaunas (Terborgh, 1971, 1977), appears to result largely from populations coming into secondary contact after allopatric divergence (Diamond, 1973; Cadena, 2007; Freeman, 2015). However, with hundreds of bird species living in the Neotropical mountains, much remains to be learned about the histories of individual clades and about how such histories collectively resulted in the patterns of diversity we observe today.

Before conducting analyses seeking to characterize and account for patterns of diversity one must have proper knowledge of what species exist and where they occur (Fine, 2015). Traditionally, the species-level taxonomy of birds was considered well-known (Scheffers *et al.*, 2012), with suggestions that the inventory of species was essentially complete by the mid 20th century (Mayr, 1946). This, however, proved incorrect: multiple avian species have been discovered and described over recent decades, and analyses of novel data sets (notably, of vocal and genetic variation) have revealed that species-level diversity was seriously underestimated (Fjeldså, 2013). The extent to which avian taxonomy will require revision depends on how one delimits species (Tobias *et al.*, 2010; Gill, 2014; Toews, 2015; Barrowclough *et al.*, 2016; Remsen, 2016), but clearly there are more species of birds than traditionally thought, particularly in the tropics. Although problems with species delimitation are unlikely to affect assessments of patterns in local (alpha) diversity of birds, inadequate knowledge of species limits may seriously influence perceptions of patterns in species turnover in space and hence regional and global patterns of diversity (beta and gamma diversity). Alternative approaches for species delimitation may also influence inferences about biogeographic history (Smith *et al.*, 2018).

Birds in which species diversity is likely greater than traditionally thought are those in which plumages vary subtly (in which case one would expect species recognition to be based more on vocal cues), and in which ecologically relevant traits (body size, habitat, dispersal ability) may be conducive to population isolation (Burney & Brumfield, 2009; Salisbury *et al.*, 2012; Smith *et al.*, 2014; Harvey *et al.*, 2017a). Here, we analyze the phylogeography of the Grey-breasted Wood-wren complex (*Henicorhina leucophrys,* Troglodytidae), a group of small, drably colored and highly vocal songbirds of forest interior, with poor dispersal abilities. Because the complex is broadly distributed from

Mexico to Bolivia and restricted to montane forest habitats, it is an appropriate system in which to ask questions
relevant to understanding the role of evolutionary processes in establishing patterns of diversity in Neotropical
mountains. We used extensive geographic sampling to reconstruct the phylogenetic relationships of populations in the
complex and to characterize patterns of genetic variation with the goals of (1) gaining insight about the tempo and
mode of evolutionary differentiation and on the role of colonization of new regions in diversification, (2) understanding
the role of geographic isolation in the differentiation of lineages, and (3) exploring the origin of elevational
replacements leading to changes in species assemblages with elevation. We also examined the extent to which current
taxonomy adequately reflects true diversity and reflected on the influence of cryptic differentiation for inferences about
diversification processes and patterns of diversity in the tropics.

Methods

Study system

Henicorhina wrens (Troglodytidae) range widely in the Neotropical region. Traditionally, taxonomists recognized two
widespread species, the White-breasted Wood-Wren (*H. leucosticta*) and the Grey-breasted Wood-Wren (*H.*
leucophrys). Two species with restricted ranges, Bar-winged Wood-Wren (*H. leucoptera*) from southern Ecuador and
northern Peru (Fitzpatrick *et al.*, 1977), and Munchique Wood-Wren (*H. negreti*) from western Colombia (Salaman *et*
al., 2003), were later described. More recently, another narrow endemic formerly considered a subspecies of *H.*
leucophrys, Hermit Wood-Wren (*H. anachoreta*) from northern Colombia, was elevated to species status (Cadena *et*
al., 2016). Preliminary data on phylogenetics and population structure of wood-wrens based on mitochondrial DNA
sequences suggest that both *H. leucosticta* and *H. leucophrys* are paraphyletic (*H. leucoptera* is nested within *H.*
leucosticta and *H. anachoreta* is nested within *H. leucophrys*), and both comprise multiple distinct lineages (Dingle *et*
al., 2006; Becker *et al.*, 2007; Caro *et al.*, 2013; Aguilar *et al.*, 2014; Smith *et al.*, 2014). However, no comprehensive
analysis of genetic variation across the range of either widespread species has been conducted.

Wood-wrens segregate ecologically by elevation. Overall, *H. leucosticta* is a lowland species replaced in montane areas
by *H. leucophrys;* their replacement is sharp and likely mediated by interspecific competition (Jankowski *et al.*, 2010).
In the isolated Cordillera del Cóndor of southern Ecuador and northern Peru, *H. leucophrys* also replaces *H.*
leucosticta in montane areas but is absent from higher elevations where *H. leucoptera* occurs (i.e. the three species
turn over along the elevation gradient; Fitzpatrick *et al.*, 1977; Dingle *et al.*, 2006). Likewise, in part of the western
slope of the Colombian Andes, *H. negreti* replaces *H. leucophrys* (subspecies *brunneiceps*) at higher elevations, and is
in turn replaced by nominate *H. leucophrys* east of the ridgeline on the eastern slope of the cordillera (Salaman *et al.*,
2003). In addition, two populations of *H. leucophrys* differing in mtDNA sequences, morphology, and songs are
parapatrically distributed along an elevational gradient in Ecuador (Dingle *et al.*, 2008; Dingle *et al.*, 2010), but
nuclear gene flow indicates they are conspecific (Halfwerk *et al.*, 2016). A similar scenario with populations differing
genetically, morphologically and vocally, and turning over along an elevational gradient exists in the Sierra Nevada de
Santa Marta, northern Colombia (Caro *et al.*, 2013; Burbidge *et al.*, 2015); because there is little to no hybridization,
these populations are now treated as separate species, with *H. anachoreta* sharply replacing *H. leucophrys* at higher
elevations (Cadena *et al.*, 2016).

Sampling

We focused on the *H. leucophrys* complex, i.e. the clade defined by the most recent common ancestor of populations
referable to *H. leucophrys* in current taxonomy and *H. anachoreta* (Dingle *et al.*, 2006; Caro *et al.*, 2013). Although *H.*
negreti was not sampled in previous molecular analyses, we consider it part of the complex based on phenotypic traits
(Salaman *et al.*, 2003) and our data (see below). The *H. leucophrys* complex is widespread in Neotropical mountains,
ranging from Mexico to Bolivia (Figure 1); as currently circumscribed, it consists of 19 taxa including *H. anachoreta*,
H. negreti, and 17 subspecies of *H. leucophrys* (Kroodsma & Brewer, 2005). For phylogeographic analyses, we sought
to sample as thoroughly as possible across geography and taxonomy. Combining sequences generated for this study
and published sequences available in GenBank (total 288 individuals), we managed to cover nearly all of the
distribution range of the complex and all named taxa, with multiple individuals and localities per taxon whenever

possible (Figure 1; Supplementary Table 1). Sampling in Middle America covered all the major montane areas where
members of the complex occur; within South America sampling was especially thorough in Venezuela, Colombia and
Ecuador, and less so in the southern part of the range (i.e. Peru and Bolivia). As outgroups for phylogenetic analyses,
we used specimens of *H. leucosticta* and *H. leucoptera,* and of species of *Microcerculus, Campylorhynchus,*
Cistothorus, Troglodytes, Cantorchilus and *Cyphorhinus* (see Barker, 2017 for an overview of relationships among
wren genera)*,* for a grand total of 300 individuals considered in analyses.

DNA extraction and sequencing

We extracted DNA using DNeasy tissue extraction kits (Qiagen, Valencia, CA) following the manufacturer's protocol.
We then amplified an 842 base-pair region of the mtDNA gene ATPase 6/8 using primers described by Joseph et al.
(2004). We chose to sequence this region because it has been employed in earlier studies of *Henicorhina* (Dingle *et al.*,
2006; Dingle *et al.*, 2008; Caro *et al.*, 2013), which enabled us to include published sequences in analyses. Fragments
were amplified via polymerase chain reaction (PCR) in 12.5 μl reactions with denaturation at 94 °C for 10 min, 40
cycles of 94 °C for 30 s, 54 °C for 45 s, and 72 °C for 2 min, followed by 10 min elongation at 72 °C and 4 °C soak.
PCR products were sequenced at the Barrick Museum of Natural History (University of Nevada, Las Vegas),
Universidad de los Andes (Bogotá, Colombia), or the High-Throughput Genomics Unit at the University of
Washington. Chromatograms were aligned using Sequencher v4.9 (GeneCodes Corporation, Ann Arbor, MI).

Analyses

Gene trees

Before phylogenetic analyses, we determined the best-fit model of evolution to be GTR + G with jModeltest v2. 1. 7
(Posada, 2008). We then used Bayesian (BEAST v1.8.4; Drummond *et al.*, 2012) and maximum-likelihood (RAxML v
8.2.4; Stamatakis, 2006) methods to estimate phylogenetic trees. For the Bayesian analyses we ran 50 million
generations, sampling trees and parameters every 5000 generations. A relaxed uncorrelated lognormal clock with a rate
of 2% and a birth-death speciation tree prior (Ritchie *et al.*, 2017) were applied. We confirmed likelihood stationarity
and adequate effective sample sizes above 200 for all estimated parameters using Tracer v1.6.0
(http://tree.bio.ed.ac.uk/software/tracer). The parameter values of the samples from the posterior distribution on the
maximum clade credibility tree were summarized after discarding the first 5 million generations (10%) as burn-in
using TreeAnnotator v1.8.4 (Drummond *et al.*, 2012). Maximum-likelihood analyses were conducted using a
GTRGAMMA model and run for 1000 nonparametric rapid bootstrap replicates to provide an assessment of nodal
support.

Both Bayesian and maximum-likelihood analyses were first done with a dataset containing all ATPase sequences (n =
300), then repeated with a reduced data set containing only individuals having unique haplotypes (n = 184), and then
reduced further by removing all non-*Henicorhina* taxa (i.e. outgroups). Trees constructed using the latter data set (n
= 178) were set aside for use in species delimitation analyses described below. To visualize and annotate trees and to
produce figures, we employed R packages ggtree (Guangchuang *et al.*, 2017) and phytools (Revell, 2012), and QGIS v
2.18.20 with the Qgis2threejs plugin (http://qgis.osgeo.org).

Species delimitation

Given uncertainty about species diversity in the *H. leucophrys* complex, we employed two coalescent approaches using
mtDNA data to identify distinct lineages which may be considered presumptive species worthy of additional study
with other sources of information (Carstens *et al.*, 2013; Sukumaran & Knowles, 2017): (1) multi-rate Poisson Tree
Processes (mPTP v0.2.0; Kapli *et al.*, 2017) and (2) a Bayesian implementation of the General Mixed Yule Coalescent
Model (bGMYC v1.0.2; Reid & Carstens, 2012). The trees generated by BEAST using the unique haplotype dataset
were used as input for these analyses.

For the mPTP analysis we ran five replicate mcmc chains of 10,000,000 generations, sampling every 10,000 with a
burn-in of 1,000,000 (10%) using the maximum clade credibility tree obtained as described above. Minimum branch
length was calculated using the minbr_auto function prior to the analysis. We ran bGMYC with 100 of the 10,000
gene trees estimated in the BEAST analysis, after removing 10% as burn-in; this approach accounts for error in gene-
tree estimation by integrating over uncertainty in tree topology and branch lengths. For each tree we ran an MCMC
chain of 50,000 steps with 40,000 steps of burn-in and a thinning interval of 100 steps. We focused our results and
discussion on lineages (i.e. presumptive species) defined using a threshold of 0.5 on probability of membership of
individuals (Gehara *et al.*, 2017). However, we also considered more conservative approaches where presumptive
species were identified as clusters in the gene tree with posterior probabilities of belonging to the same species ≥ 0.90
or ≥ 0.95, which allowed us to compare our results to similar work on other birds (Smith *et al.*, 2014; Harvey *et al.*,
2017b; Smith *et al.*, 2017).

Diversification through time

To describe patterns of lineage accumulation over time, we constructed lineage-through-time (LTT) plots and
estimated the gamma statistic (Pybus *et al.*, 2000). We accounted for phylogenetic uncertainty by performing these
analyses on a sample of credible trees in the posterior distribution obtained from the BEAST analysis. We employed
the 100 trees constructed using only unique haplotypes as input for the bGMYC analyses and trimmed them to
include 39 tips corresponding to the presumptive species recognized under the 0.5 threshold. Then we used functions
implemented in R packages ape (Paradis *et al.*, 2004) and paleotree (Bapst, 2012) to build an LTT plot with a 95%
confidence interval and to calculate the gamma statistic for each tree. We also examined diversification dynamics
employing Bayesian Analysis of Macroevolutionary Mixtures (program BAMM v 2.5.0; Rabosky *et al.*, 2013). Because
results were qualitatively similar between methods, we report only those obtained using the simpler approach
implemented in ape.

Results

We found substantial genetic differentiation among populations in the *H. leucophrys* complex. In total, we recovered
172 haplotypes among the 288 individuals analyzed. Several haplotypes were highly divergent from each other, with
uncorrected genetic distances between them reaching >12 % (i.e. between individuals from Sierra Madre del Sur of
Mexico and from the east slope of the Cordillera Occidental of Colombia). Genetic variation was highly structured
spatially, but was not readily accounted for by geographic distance among populations. We did not conduct formal
analyses of isolation by distance, but genetic distances among isolated populations from adjacent mountains were often
much greater than genetic distances observed over larger distances in more continuous ranges. For instance, mean
genetic distances among the five montane areas of Venezuela that we sampled was 7.3% (range 6.1%-8.3%), whereas
genetic distances within montane regions extending over comparable distances were much lower, e.g. reaching only
3.5% in the Sierra Madre Oriental of Mexico or 1.7% in the Talamanca-Chiriquí mountains of Costa Rica and
Panama.

Maximum-likelihood and Bayesian phylogenetic analyses recovered similar overall patterns (Figure 2., Supplementary
Figures 1-2). The deepest split in gene trees separates clades corresponding to Mexican populations from the Sierra
Madre del Sur and the western reaches of the Trans-Mexican Volcanic Belt from a large and strongly supported clade
including the remainder of populations in the complex. Within the latter clade, the earliest diverging group occurs in
eastern Mexico and consists of three subclades, each corresponding to a unique region within the Sierra Madre
Oriental. Sister to this group is a clade divided in two main groups (albeit without strong support, i.e. 0.88 posterior
probability, 68% ML bootstrap in analyses using only unique haplotypes): one includes samples from lower Central
America (Costa Rica and Panama, subspecies *collina*), whereas the other includes all South American populations of
H. leucophrys, the Colombian endemic species *H. anachoreta* and *H. negreti*, and a clade formed by samples of *H.*
leucophrys from Chiapas (Mexico), Guatemala, and El Salvador (subspecies *castanea* and *composita*). The latter clade,
of somewhat uncertain affinities within an otherwise South American group (it was recovered as sister to *H. negreti*
with 0.95 posterior probability and 56% ML bootstrap in analyses using only unique haplotypes), is the only exception
to the pattern in which Mexican and Central American populations are the earliest diverging lineages in the complex.

Basal relationships among major South American lineages were unresolved or poorly supported, yet some patterns in
the region are noteworthy. Whereas populations from some isolated montane systems (e.g. the Serranía de Perijá, or
the Venezuelan Cordillera de la Costa and Sierra de San Luis) formed distinct clades, this was not the case for the
main cordilleras of the northern Andes, resulting in complicated patterns of area relationships. For example, several
populations from the western slope of the Andes from Colombia through Ecuador and into northwestern Peru formed
a large clade (clade i. in Figure 2), but this clade was not exclusive because it also included some -but not all-
populations from the northern sector of the Cordillera Central of Colombia and did not cluster all populations from
the western Andes: *H. negreti* and lineages of *H. leucophrys* from the northern and southern sectors of the Cordillera
Occidental of Colombia and from western Ecuador occupied different positions in the tree. Likewise, birds from the
Cordillera Oriental of the Colombian Andes formed multiple distinct clades seemingly distantly related to each other,
and populations from the Central and Southern Andes (i.e. from Bolivia and Peru south of the Marañón Valley or
North Peru Low) formed at least two highly divergent clades with differing affinities. Given weak support for deep
branches in South America we do not elaborate further on relationships among major biogeographic areas, but do
emphasize the complexity of phylogeographic pattern and the strong genetic structure existing over relatively fine
spatial scales throughout the continent.

Part of the complexity in phylogeographic pattern related to occurrence of phylogenetically distant groups in the same
regions can be understood by examining elevational distributions: lineages known to replace each other along
elevational gradients in northern South America are not sister to each other (Figure 3). This was true of taxa
occurring in the Ecuadorean Andes (*H. l. hilaris* and nominate *H. l. leucophrys*), the Santa Marta mountains (*H. l.*
bangsi and *H. anachoreta*), and in the western slope of the Colombian Andes (*H. l. brunneiceps* and *H. negreti*). Our
data further revealed a previously unknown case of cryptic replacement of mtDNA lineages along an elevational
gradient in the Venezuelan Andes. The lineage occurring in the Tamá massif near the Colombia-Venezuela border
crosses the Táchira depression to the northeast into the Cordillera de Mérida where we found it from c. 1520 m to c.
1920 m. Only a small distance upslope in this range, a different lineage occupied elevations from c. 2100 m to c. 2750
m. Divergence in mtDNA sequences between lineages replacing each other with elevation was substantial, in all cases
exceeding 5% uncorrected *p* distances (Figure 3).

In addition to examples of elevational parapatry, our analyses revealed cases where lineages may meet in contact zones
along a latitudinal axis. For instance, two lineages differing in c. 7% uncorrected p-distance replace each other along
the northern sector of the Cordillera Central of the Colombian Andes. One of these lineages occurs in the northern tip
of the cordillera in Antioquia, whereas the other is also found in Antioquia, where we suspect it might range north to
the southern extreme of the Aburrá Valley in the outskirts of the city of Medellín. Individuals occupying the northern
extreme of the Cordillera Central are most closely allied to geographically distant populations from the western slope
of the Andes (i.e. subspecies *hilaris* and *brunneiceps*), which tend to occur at lower elevations and are replaced upslope
by nominate *H. leucophrys* in Ecuador or *H. negreti* in southwest Colombia. Several other examples of distinct
lineages occurring at different latitudes within mountain systems exist in the Cordillera Oriental of Colombia and
along the Andes of Ecuador (Figure 2).

Coalescent approaches to delimit species produced consistent results: both mPTP and bGMYC (the latter with a 0.50
probability threshold to define group membership) recovered *H. anachoreta* and *H. negreti* as distinct species, and
both methods identified 37 additional lineages in the *H. leucophrys* complex which may prove to be distinct species
(Figure 2, Supplementary Figure 3). Although methods did not exactly agree in how they assigned individuals to
presumptive species, congruence was remarkable. The only differences were that in the Cordillera de Mérida,
Venezuela, mPTP recognized three presumptive species while bGMYC recognized two, and that in a clade from
northern Peru and southeast Ecuador mPTP recognized a single presumptive species and bGMYC recognized two
(Figure 2). Applying more stringent probability thresholds to delimit species in bGMYC analyses resulted in the
inference of slightly lower numbers of presumptive species: 36 and 35 with 0.90 and 0.95 thresholds, respectively. In
general, presumptive species appear to have restricted ranges (Figure 4, Supplementary Figure 4); in some cases,
particular mountain systems harbor a single presumptive species (e.g. Sierra Madre del Sur and Trans-Mexican
Volcanic Belt in Mexico, Cordillera de la Costa and Sierra de San Luis in Venezuela), but more than one presumptive
species may also occur within a region (e.g. Sierra Madre Oriental of Mexico, Sierra Nevada de Santa Marta in

Colombia) and a few of them have ranges encompassing various montane areas (e.g. across cordilleras of Costa Rica
and Panama). The diversity of presumptive species is especially remarkable in northern South America, with 7-8
identified in Venezuela, 15 in Colombia, and 7-8 in Ecuador. Because our sampling was sparser in Peru and Bolivia,
our figures for these countries are likely underestimates of presumptive species richness.

Our estimates of divergence times obtained from the BEAST analysis of unique haplotypes indicate that the *H.*
leucophrys complex diverged from its sister group (i.e. the clade formed by *H. leucosticta* and *H. leucoptera*)
approximately 16.8 million years before present (13.0-20.5 95% highest posterior density, HPD), with the crown age of
extant populations dating to 10.6 m.a. (8.4-13.1 95% HPD). Over this period, the complex has diversified into
multiple lineages; we found it minimally consists of 10 lineages of at least 5 million years of age and of 26 lineages of
at least 2 million years of age (Figure 2, Supplementary Figure 3). The estimated age of the node including all South
American populations as well populations from Chiapas, Guatemala, and El Salvador is 6.2 m.a. (5.2-7.3 95% HPD),
whereas that of the node including all South American populations excluding *H. negreti* is 5.8 m.a. (4.9-6.7 95%
HPD).

Analyses of lineage accumulation over time based on presumptive species identified by bGMYC suggested that rates of
diversification in the *H. leucophrys* complex may have declined over time, with a significantly negative gamma
statistic (Figure 5). However, through much of the history of the complex, diversification appears to have been nearly
constant and exponential, with an apparent downturn in the last million years most likely reflecting that our species
delimitation analyses recognized no species younger than this age.

Discussion

Phylogeography: bridges, barriers and the distribution of genetic variation.

As evidenced by the branching pattern in the gene tree with deep splits involving populations from the north, the *H.*
leucophrys complex likely originated in the Mexican highlands ca. 8 to 13 m.a., from where it expanded south through
Central America, then colonizing South America. Other birds ranging broadly in montane forests also originated in
the northern Neotropics, including single species as well as clades which diversified in Central and South America
(Pérez-Emán, 2005; Cadena *et al.*, 2007; Weir *et al.*, 2008; Sánchez-González *et al.*, 2015). Molecular-based estimates
of when did birds colonize South America from the north vary (Bacon *et al.*, 2015; Barker *et al.*, 2015); our results
indicate colonization by the *H. leucophrys* complex occurred slightly earlier (ca. 7 to 5 m.a.) than a pulse of avian
interchange via the Isthmus of Panama 4 to 2 m.a. (Smith & Klicka, 2010). Our estimates of the age of the *H.*
leucophrys complex and of the timing of events like its colonization of South America are old relative to what one
would expect given published estimates of divergence times among wren genera (Barker, 2017). However, such
estimates were derived assuming that *Certhia* and *Troglodytes* diverged ca. 16 m.a. (Moyle *et al.*, 2016), while
analyses integrating more extensive fossil evidence suggest such divergence occurred much earlier, ca. 27 m.a.
(Claramunt & Cracraft, 2015). The time frame for wren diversification implied by the latter analysis is more
congruent with our estimated ages for nodes in *Henicorhina* and with previous work in other wren genera based on
mtDNA data (Barker, 2007).

Poorly supported relationships among South American clades associated with short internodes subtending long
branches are common to *H. leucophrys* and other birds colonizing South America from the north (e.g., Pérez-Emán,
2005; Cadena *et al.*, 2007). This indicates range expansions and ensuing rapid diversification of lineages in geographic
isolation, a pattern also documented in montane clades with South American (Chaves *et al.*, 2011) or uncertain
geographic origins (Gutiérrez-Pinto *et al.*, 2012). Rapid range expansions occurring in concert across birds may
indicate that geological changes like closure of the Isthmus of Panama and uplift of mountains increased connectivity
among formerly isolated regions, enabling subsequent diversification of various taxa over the vast South American
landscape; climatic changes driving population isolation likely facilitated such diversification (Barrantes, 2009;
Ramírez-Barahona & Eguiarte, 2013).

Genetic divergence associated with landscape features isolating montane habitats is another pattern shared by the *H.*
leucophrys complex and co-distributed clades (Weir, 2009). Such features include lowland areas in Central America
(Cadena *et al.*, 2007; Barber & Klicka, 2010), inter-Andean valleys like the Magdalena and Marañón (Gutiérrez-Pinto
et al., 2012; Benham *et al.*, 2015), and alpine areas separating slopes of cordilleras (Parra *et al.*, 2009; Valderrama *et*
al., 2014). For many Neotropical montane birds that have been studied, genetic structure across geographic barriers
coincides with plumage differences (Cadena *et al.*, 2011; Winger & Bates, 2015; Winger, 2017). Phenotypic differences
among distinct lineages of *H. leucophrys*, however, are either subtle or appear to be nonexistent (Kroodsma & Brewer,
2005). Reduced gene flow across barriers may have influenced vocal differentiation of wood-wrens more strongly, but
given their complex songs, confirming it awaits studies documenting repertoires of individuals as well as variation
within and among populations. Such data are relevant given uncertainty about species limits in the complex (see
below) because vocalizations likely play a critical role in species recognition (Caro *et al.*, 2013; but see Halfwerk *et al.*,
2016).

In sum, our study and other phylogeographic analyses point to geological and climatic dynamics of the montane
Neotropics as drivers of avian speciation both by (1) promoting dispersal across formerly isolated areas and (2)
spurring diversification linked to the origin of new habitats resulting from uplift processes and vicariance.
Furthermore, because wood-wrens live in rugged landscapes and disperse little, their populations may become isolated
and diverge even without marked geological or climatic changes (Smith *et al.*, 2014). Beyond patterns common to the
H. leucophrys complex and other tropical montane birds, two aspects appear unique to our study system. First, the
degree of genetic structure within a single recognized species we uncovered far exceeds that observed in other montane
birds. Second, our finding that distinct mtDNA lineages which likely diverged in allopatry have come into contact in
various regions and some coexist segregated by elevation is novel. Because the extreme genetic structure we uncovered
may imply that *H. leucophrys* comprises more species than traditionally thought and because secondary sympatry of
divergent populations is crucial to the buildup of species richness, these results have implications for understanding
tropical diversity and the historical and evolutionary processes generating and sustaining it.

Extreme population structure, cryptic divergence, and patterns in tropical diversity

We uncovered genetic structure in the *H. leucophrys* complex across well-known geographic barriers (Hazzi *et al.*,
2018), but also over fine scales in ways not associated with divergence in other tropical montane birds. For example,
in the Cordillera Occidental and Cordillera Central of Colombia, where other birds show little to no population
structure (Cadena *et al.*, 2007; Gutiérrez-Pinto *et al.*, 2012; Isler *et al.*, 2012; Valderrama *et al.*, 2014), we found six
mtDNA lineages of at least 1 million years of age. These lineages and others have restricted ranges, and some of their
boundaries reflect topographic or climatic breaks (Graham *et al.*, 2010; Supplementary Figure 4). Traits affecting
dispersal abilities and dependence on closed understory habitats mediate divergence across putative barriers and thus
diversification in topographically complex landscapes (Burney & Brumfield, 2009; Smith *et al.*, 2014). Because wood-
wrens are small-bodied, have small and rounded wings and live in dark forest understory, they likely disperse little
(Moore *et al.*, 2008), and this may account in part for their exceptionally strong population structure (Claramunt *et*
al., 2012; Salisbury *et al.*, 2012; but see Smith *et al.*, 2017). Deep phylogeographic structure also exists in other small-
bodied wrens (i.e. other *Henicorhina, Cistothorus, Troglodytes*; Dingle *et al.*, 2006; Campagna *et al.*, 2012; Galen &
Witt, 2014; Robbins & Nyári, 2014), suggesting that their biology predisposes populations to become isolated and
diverge.

Regardless of the ultimate causes of population structure, we discovered heretofore underappreciated diversity within a
taxon traditionally treated as a single species. Although our study employed only one molecular marker, some of the
lineages we recovered coexist as distinct phenotypic entities exhibiting behavioral barriers to hybridization (Salaman
et al., 2003; Caro *et al.*, 2013; Burbidge *et al.*, 2015), implying that several species are involved. Genetic distances
(i.e., divergence times) are not appropriate surrogates for reproductive isolation (Roux *et al.*, 2016), but we note that
wood-wren lineages demonstrating barriers to gene flow in sympatry (i.e. *anachoreta* and *l. bangsi*; *negreti* and *l.*
brunneiceps; *negreti* and nominate *leucophrys*) last shared ancestors more recently than many other lineages in the
complex. It is also remarkable that the phylogeography of *H. leucophrys* resembles that of *Atlapetes* brushfinches
(Emberizidae), which also have a northern origin and montane distribution through the Neotropics, and which have

diverged into numerous lineages upon colonizing South America (Sánchez-González *et al.*, 2015; J. L. Pérez-Emán, unpubl. data). In contrast to *H. leucophrys*, *Atlapetes* diversified extensively in plumage and this has arguably influenced taxonomy, with researchers recognizing 28 species in the group (Remsen *et al.*, 2018). Just as lineages of the *H. leucophrys* complex replace each other in space, species of *Atlapetes* are for the most part allopatric or replace each other sharply along elevational or latitudinal axes in the Andes (Remsen & Graves, 1995), with their ranges often matching those of lineages of *H. leucophrys* uncovered by our study. This comparison serves to illustrate what might be a more general situation in which clades with roughly similar ages and genetic structure (South American *Atlapetes* are actually younger than South American *H. leucophrys*) may be split to different degrees by taxonomists because of differences among clades in the traits birds employ for signaling and in the lability of such traits. In other words, birds like wood-wrens may be under split owing to their conserved plumage and because the role of vocalizations in species recognition remains understudied (see also D'Horta *et al.*, 2013).

Our coalescent analyses indeed suggest that taxonomy underestimates species diversity in the *H. leucophrys* complex: we consistently identified 39 presumptive species across methods. Even the 35 presumptive species identified by bGMYC using a more conservative probability threshold of 0.95 represents a quite remarkable figure relative to similar studies conducted in the Neotropics. In 27 clades of lowland birds ranging from Central America across the Andes through much of Amazonia and even into the Atlantic Forest (Smith *et al.*, 2014), the mean number of species identified using bGMYC also with a 0.95 threshold was 5.3 (range 1-18); figures were slightly higher in understory birds (11 clades; mean = 6.6, range 3-11 presumptive species) but still much lower than our estimates for the *H. leucophrys* complex. Moreover, in 173 taxonomic species of birds from the New World subject to phylogeographic analyses employing mtDNA data, the largest number of presumptive species identified by bGMYC with a 0.90 threshold was 23 (Harvey *et al.*, 2017b; Smith *et al.*, 2017), highlighting the *H. leucophrys* complex as a distinct outlier (Figure 6). This is despite our sparse sampling in the Peruvian and Bolivian Andes, where one would expect more lineages exist. We do not argue that all lineages we uncovered are species given existing evidence, but several are candidates for studies examining other molecular markers, morphology, voices, and behavior (Caro *et al.*, 2013; Burbidge *et al.*, 2015; Halfwerk *et al.*, 2016). The allopatric ranges of most wood-wren lineages preclude tests of intrinsic barriers to gene flow, but given postzygotic isolation in other phenotypically cryptic, old lineages of Neotropical birds (Pulido-Santacruz *et al.*, 2018), some of them may well be reproductively isolated.

Even if hypothetical species our analyses flagged represent distinct lineages not yet reaching the status of "biological" species (Carstens *et al.*, 2013; Sukumaran & Knowles, 2017), our work allows for conclusions about cryptic diversity which should be robust to analyses involving other data. First, regardless of the species concept one follows and of the criteria one uses to recognize species, there are more wood-wren species than traditionally thought. Second, the *H. leucophrys* complex comprises multiple independently evolving populations which have diverged to different degrees along the speciation continuum; if one embraces the generalized lineage concept, which views species as segments of metapopulation lineages and considers all other species "concepts" as contingent –albeit not necessary– properties of species one may use as criteria to recognize them (de Queiroz, 1998, 2007), then the complex is arguably a collection of several kinds of species, all of which represent fundamental evolutionary units. Finally, even if one takes a conservative standpoint and treats most of the lineages we uncovered as distinct populations of a single or a few species (Sukumaran & Knowles, 2017), the *H. leucophrys* complex has clearly differentiated into numerous lineages (especially following its colonization of South America) and such lineages have persisted over long periods.

Because most wood-wren lineages are not sympatric (see below), our finding that the *H. leucophrys* complex probably comprises multiple species has no influence on local estimates of diversity. However, species richness and endemism at regional scales might need to be revised. For example, current taxonomy recognizes only one species (and only five subspecies) in the complex in Venezuela (Kroodsma & Brewer, 2005), yet we uncovered 7-8 presumptive species –each endemic to a particular montane system– in the country. If lineages identified as presumptive species are elevated to species status and similar patterns exist in other taxa not yet studied, then geographic variation in population structure may alter knowledge of spatial patterns of diversity (e.g., differences in species richness among cordilleras and slopes of cordilleras of Colombia; Kattan *et al.*, 2004; Figure 4, Supplementary Figure 4). Alternative classifications also affect inferences about beta diversity; under current taxonomy, there is no species turnover along thousands of kilometers and across multiple geographic barriers from Mexico to Bolivia except for the local

replacements involving *H. anachoreta* and *H. negreti*. At the other extreme, if distinct lineages of *H. leucophrys* are
species, then spatial turnover would be substantial even over relatively short distances (e.g., in Colombia), likely
exacerbating differences among regions in beta diversity (Gaston *et al.*, 2007; Fjeldså *et al.*, 2012) and with potential
conservation implications (Socolar *et al.*, 2016). Recognizing distinct lineages as species would also alter the perceived
role of features of montane landscapes setting range limits and thus explaining spatial turnover in assemblages
(Graham *et al.*, 2010).

In addition to affecting perceptions of patterns of diversity, our results have implications for thinking about historical
processes underlying such patterns. The accumulation of biological diversity via diversification within a region like the
montane Neotropics requires that (1) populations become isolated to initiate divergence, (2) budding population
isolates persist in time, (3) populations expand their ranges and come into secondary sympatry, and (4) newly
sympatric populations are differentiated enough that they may coexist without coalescing owing to hybridization or
without excluding each other via competition (Mayr, 1942; Ricklefs & Bermingham, 2007). In the following we discuss
our results in the context of these steps.

Lineage splitting, persistence, and the origin of diversity

A leading explanation for high tropical diversity involves latitudinal differences in net diversification rates (Fischer,
1960; Schluter & Pennell, 2017). In particular, rapid diversification may explain the high species richness and
concentration of narrow-ranged species of birds in tropical mountains, which cannot be accounted for by area or
contemporary climate (Jetz *et al.*, 2004; Fjeldså *et al.*, 2012). Although evidence that diversification rates vary with
latitude remains mixed in birds (Ricklefs, 2006; Martin & Tewksbury, 2008; Jetz *et al.*, 2012; Belmaker & Jetz, 2015)
and other taxa (e.g. Pyron & Wiens, 2013; Pyron, 2014; Rolland *et al.*, 2014; Schluter, 2016; Rabosky *et al.*, 2018),
tropical mountains are indeed hotbeds of rapid diversification (Madriñán *et al.*, 2013). Furthermore, differences in the
rate at which species originate may not be as important as the rate at which they go extinct in establishing broad-
scale patterns in avian diversity (Hawkins *et al.*, 2006; Weir & Schluter, 2007; Pulido-Santacruz & Weir, 2016).
Accordingly, the high diversity and endemism of tropical montane areas may reflect low extinction rates of species
(Fjeldså *et al.*, 2012). In turn, higher diversification rates at higher elevations in montane areas (Quintero & Jetz,
2018) may reflect both high speciation and low extinction (Fjeldså & Irestedt, 2009). A complementary historical
explanation for diversity in tropical mountains which is less commonly addressed in the literature is high persistence of
budding populations, an important control of rates of speciation (Mayr, 1963; Dynesius & Jansson, 2014; Rabosky,
2016).

We found that the *H. leucophrys* complex radiated rapidly into multiple lineages, several of which have persisted for
periods exceeding millions of years. Also, LTT plots suggest nearly constant rates of exponential accumulation of
lineages over nearly 10 million years, with an apparent slowdown in diversification in the last million years. Although
LTT plots with such a shape and their associated negative gamma statistic are often considered evidence of ecological
limits to diversification (Rabosky & Hurlbert, 2015), we interpret the pattern more as an artifact of our methods
resulting from (1) using species based on a model specifying a divergence threshold separating population-level
processes (gene coalescence) from diversification dynamics (speciation and extinction) as units for analysis (Reid &
Carstens, 2012), and (2) potential limitations in geographic sampling leading to failure to identify additional
independent lineages of young age. Because the bGMYC analysis we employed to delimit presumptive species
established an age cutoff of ca. 1 m.a. defining the units included in the LTT analysis, we simply conclude that
diversification was nearly constant through much of the history of the *H. leucophrys* complex. To the extent that
similar diversification dynamics may characterize evolutionary history of other Neotropical montane birds, high rates
of lineage splitting (Harvey *et al.*, 2017b) and high persistence of such lineages over time (Smith *et al.*, 2017) have
likely contributed to diversification and probably account for the high diversity of tropical montane systems and, more
broadly, to large-scale biodiversity patterns such as latitudinal gradients in species richness.

Range dynamics, secondary sympatry and the regional buildup of diversity

Our data revealed that mtDNA lineages in the *H. leucophrys,* which likely diverged in geographic isolation, have come
into secondary sympatry. This is most evident where divergent mtDNA lineages not sister to each other segregate
with elevation. In addition to previously documented cases of elevational replacements of lineages involving distinct
taxa (i.e. different species or subspecies in the Sierra Nevada de Santa Marta, in western Colombia, and in western
Ecuador), we discovered a novel elevational replacement of distinct lineages in the Mérida Cordillera of Venezuela
where no phenotypic differences had been noted. Likewise, previous work in other wood-wrens revealed that although
H. leucoptera is nested within *H. leucosticta*, the lineage of *H. leucosticta* replaced by *H. leucoptera* at higher
elevations in the Cordillera del Cóndor east of the Andes is distantly related to it, whereas its closest relative
seemingly occurs in the Chocó region west of the Andes (Dingle *et al.*, 2006). The consistent pattern of elevational
replacements involving fairly distant relatives as opposed to sister lineages fits the hypothesis that evolutionary
divergence in tropical montane birds occurs largely in allopatry and not in parapatry along mountain slopes (Patton &
Smith, 1992; García-Moreno & Fjeldså, 2000; Caro *et al.*, 2013).

In addition to documenting elevational replacements, we found evidence of regional co-occurrence of lineages replacing
each other with latitude (e.g. along the cordilleras of Colombia and Ecuador). More fine-scaled sampling is required to
determine whether geographic gaps separate the ranges of such lineages or if they come into close contact. Part of the
observed genetic differentiation along the latitudinal axis may reflect the propensity of the linear distributions of
tropical montane birds to become fragmented (Graves, 1988). However, some lineages replacing each other with
latitude in a region are not sisters and may even be distantly related, which suggests range expansions and secondary
contact rather than primary divergence along cordilleras.

Other intriguing phylogeographic patterns aside from secondary contact of lineages in elevational or latitudinal
parapatry speak to the dynamism of geographic ranges over broad scales. For example, we found that populations of
H. leucophrys from southern Mexico (Chiapas), Guatemala and El Salvador are not closely related to other Middle
American populations; within a large, otherwise South American clade, these specimens appeared closest to *H. negreti,*
a species endemic to western Colombia whose northernmost records are ca. 1700 km south of montane El Salvador.
Likewise, the only sequence analyzed from Bolivia is a long branch more closely allied to lineages from northern South
America (Colombia and Venezuela) than to geographically much closer lineages from Peru. Because closest relatives
may occur in distant areas, spatial patterns of genetic variation are not easily accounted for by geography (e.g. by
isolation-by-distance; Seeholzer & Brumfield, 2018). Given that such patterns are unlikely evidence of long-distance
dispersal and are not unique to wood-wrens in the region (Cadena *et al.*, 2007), considering dynamics of expansion and
contraction of geographic ranges involving localized extinctions is crucial to understand biogeographic and
demographic processes underlying the distribution of genetic and species diversity in Neotropical birds.

Shifting climatic conditions affecting habitat connectivity drive changes in species ranges, thereby influencing
phylogeographic patterns and the buildup of montane diversity (Ramírez-Barahona & Eguiarte, 2013; Flantua &
Hooghiemstra, 2018). Species ranges may also experience phases of expansion and contraction linked to shifts in
ecological specialization and interactions with natural enemies (i.e. the taxon cycle; Wilson, 1959; Ricklefs &
Bermingham, 2002). Although taxon cycles are more evident in insular settings with discrete populations and areas
(e.g., Ricklefs & Bermingham, 1999; Jønsson *et al.*, 2014), they may also take place in continents (Graves, 1982). In
fact, lineages experiencing the taxon cycle may account for what one might call continental great speciators like
Henicorhina wood-wrens, which occur widely in space -revealing an ability to expand their ranges- yet split into
isolated populations at a fast rate due to cessation of gene flow (cf. Diamond *et al.*, 1976). Wood-wrens disperse little
at present because of their morphology and ecology, which arguably explain their remarkable patterns of genetic
structure reflecting long-term population isolation. However, our findings that wood-wrens dispersed throughout much
of the montane Neotropics from a northern area of origin and that several lineages achieved secondary sympatry
indicate that episodes of range expansion interspersed with periods of divergence occurred at various moments,
possibly in sync with morphological or behavioral changes influencing their abilities to disperse (Pigot & Tobias, 2015;
Hosner *et al.*, 2017). Furthermore, gaps separating the ranges of closely related lineages of wood-wrens arguably reflect
extinctions of intervening populations of formerly widespread lineages, which left vacant spaces that could, in turn,
become occupied by other expanding lineages.

Phylogeographers will often not detect range dynamics embodied in the taxon cycle because incomplete reproductive
isolation between young lineages can result in homogenization of gene pools upon secondary contact (Kearns *et al.*,
2018). Furthermore, niche similarities between incipient species achieving contact may preclude long-term sympatry
owing to interspecific competition (Pigot & Tobias, 2013). Irrespective of whether the patterns we observed resulted
from the taxon cycle, we identified aspects making the *H. leucophrys* complex well suited for further work on the
origins of tropical diversity and its accumulation over time and space. Our results and other work on the complex
reveal that the completion of reproductive isolation between lineages meeting in secondary sympatry seemingly
exhibits a continuum ranging from neutral divergence with no obvious phenotypic differences (forms in montane
Venezuela), to phenotypic and behavioral divergence with persistent interbreeding (*hilaris* and nominate *leucophrys* in
Ecuador), to completed speciation with little to no hybridization (anachoreta and negreti vs. various forms of
leucophrys in Colombia; Salaman *et al.*, 2003; Dingle *et al.*, 2008; Dingle *et al.*, 2010; Caro *et al.*, 2013; Burbidge *et*
al., 2015; Halfwerk *et al.*, 2016). Furthermore, divergence in elevational ranges occurring during periods of isolation
(Cadena, 2007; Tobias *et al.*, 2014) or arising in secondary sympatry (Diamond, 1973; Freeman, 2015) has enabled
coexistence of lineages at the landscape scale in various regions. Given that range boundaries may be maintained –and
possibly reinforced– evolutionarily by phenotypic and behavioral barriers to interbreeding and ecologically by
competition (Jankowski *et al.*, 2010), our study has uniquely captured wood-wren populations in the act of building up
diversity via divergence and persistence in allopatry, achievement of secondary sympatry, and coexistence mediated by
ecological and evolutionary divergence. Comparative work on the structure and dynamics of contact zones between
lineages should provide rich insights into the origin and maintenance of high diversity in tropical mountains.

Acknowledgements

We are grateful to the institutions allowing access to tissue samples for our study: Academy of Natural Sciences of
Drexel University, Burke Museum of Natural History and Culture at University of Washington, Colección
Ornitológica Phelps, The Field Museum, Instituto Alexander von Humboldt, Instituto de Ciencias Naturales at
Universidad Nacional de Colombia, Louisiana State University Museum of Natural Science, Museum of Vertebrate
Zoology at UC Berkeley, Peabody Museum of Natural History at Yale University, and the Zoological Museum at the
University of Copenhagen. We thank field collectors and field assistants for their crucial role in obtaining samples, as
well as researchers who conducted groundbreaking work on our study system, made their sequences available and
assisted us in various ways: Caroline Dingle, Lina Caro, and colleagues. Our work received financial support from the
Facultad de Ciencias at Universidad de los Andes, and the Centro de Desarrollo Científico y Humanístico de la
Universidad Central de Venezuela. Permits were granted by the Oficina Nacional de Diversidad Biológica, Ministerio
del Poder Popular para el Ambiente, Venezuela.

Literature Cited

Aguilar, C., De León, L.F., Loaiza, J.R., McMillan, W.O. & Miller, M.J. (2014) Extreme sequence divergence between
mitochondrial genomes of two subspecies of White-breasted Wood-wren (*Henicorhina leucosticta*, Cabanis,
1847) from western and central Panamá. *Mitochondrial DNA*, **27**, 956-957.
Bacon, C.D., Silvestro, D., Jaramillo, C., Smith, B.T., Chakrabarty, P. & Antonelli, A. (2015) Biological evidence
supports an early and complex emergence of the Isthmus of Panama. *Proceedings of the National Academy*
of Sciences of the USA, **112**, 6110-6115.
Badgley, C., Smiley, T.M., Terry, R., Davis, E.B., DeSantis, L.R.G., Fox, D.L., Hopkins, S.S.B., Jezkova, T., Matocq,
M.D., Matzke, N., McGuire, J.L., Mulch, A., Riddle, B.R., Roth, V.L., Samuels, J.X., Strömberg, C.A.E. &
Yanites, B.J. (2017) Biodiversity and topographic complexity: Modern and geohistorical perspectives. *Trends*
in Ecology and Evolution, **32**, 211-226.
Bapst, D.W. (2012) paleotree: an R package for paleontological and phylogenetic analyses of evolution. *Methods in*
Ecology and Evolution, **3**, 803-807.
Barber, B.R. & Klicka, J. (2010) Two pulses of diversification across the Isthmus of Tehuantepec in a montane
Mexican bird fauna. *Proceedings of the Royal Society of London B*, **277**, 2675-2681.

Barker, F.K. (2007) Avifaunal interchange across the Panamanian Isthmus: insights from *Campylorhynchus* wrens.
Biological Journal of the Linnean Society, **90**, 687-702.
Barker, F.K. (2017) Molecular phylogenetics of the wrens and allies (Passeriformes: Certhioidea), with comments on
the relationships of *Ferminia*. *American Museum Novitates*, **3887**, 1-27.
Barker, F.K., Burns, K.J., Klicka, J., Lanyon, S.M. & Lovette, I.J. (2015) New insights into New World biogeography:
An integrated view from the phylogeny of blackbirds, cardinals, sparrows, tanagers, warblers, and allies. *Auk*,
132, 333-348.
Barrantes, G. (2009) The role of historical and local factors in determining species composition of the highland
avifauna of Costa Rica and Western Panama. *Revista de Biología Tropical*, **57 (Suppl. 1)**, 333-349.
Barrowclough, G.F., Cracraft, J., Klicka, J. & Zink, R.M. (2016) How many kinds of birds are there and why does it
matter? *PLoS One*, **11**, e0166307.
Becker, C.D., Richardson, E.A. & Brown, S.J. (2007) MtDNA haplotypes, sequence divergence, and morphological
variation in Gray-breasted Wood Wrens (*Henicorhina leucophrys*) and their conservation implications.
Journal of Field Ornithology, **78**, 117-123.
Belmaker, J. & Jetz, W. (2015) Relative roles of ecological and energetic constraints, diversification rates and region
history on global species richness gradients. *Ecology Letters*, **18**, 563-571.
Benham, P.M., Cuervo, A.M., McGuire, J.A. & Witt, C.C. (2015) Biogeography of the Andean metaltail
hummingbirds: contrasting evolutionary histories of tree line and habitat-generalist clades. *Journal of*
Biogeography, **42**, 763–777.
Bertuzzo, E., Carrara, F., Mari, L., Altermatt, F., Rodriguez-Iturbe, I. & Rinaldo, A. (2016) Geomorphic controls on
elevational gradients of species richness. *Proceedings of the National Academy of Sciences of the United*
States of America, **113**, 1737-1742.
Burbidge, T., Parson, T., Caycedo-Rosales, P.C., Cadena, C.D. & Slabbekoorn, H. (2015) Playbacks revisited:
asymmetry in behavioural response across an acoustic boundary between two parapatric bird species.
Behaviour, **152**, 1933-1951.
Burney, C.W. & Brumfield, R.T. (2009) Ecology predicts levels of genetic differentiation in Neotropical birds.
American Naturalist, **174**, 358-368.
Cadena, C.D. (2007) Testing the role of interspecific competition in the evolutionary origin of elevational zonation: an
example with *Buarremon* brush-finches (Aves, Emberizidae) in the neotropical mountains. *Evolution*, **61**,
1120-1136.
Cadena, C.D., Klicka, J. & Ricklefs, R.E. (2007) Evolutionary differentiation in the Neotropical montane region:
molecular phylogenetics and phylogeography of *Buarremon* brush-finches (Aves, Emberizidae). *Molecular*
Phylogenetics and Evolution, **44**, 993-1016.
Cadena, C.D., Cheviron, Z.A. & Funk, W.C. (2011) Testing the molecular and evolutionary causes of a 'leapfrog'
pattern of geographical variation in coloration. *Journal of Evolutionary Biology*, **24**, 402-414.
Cadena, C.D., Caro, L.M., Caycedo, P.C., Cuervo, A.M., Bowie, R.C.K. & Slabbekoorn, H. (2016) *Henicorhina*
anachoreta (Troglodytidae), another endemic bird species for the Sierra Nevada de Santa Marta, Colombia
Ornitología Colombiana, **15**, 82-89.
Cadena, C.D., Kozak, K.H., Gómez, J.P., Parra, J.L., McCain, C.M., Bowie, R.C.K., Carnaval, A.C., Moritz, C.,
Rahbek, C., Roberts, T.E., Sanders, N.J., Schneider, C.J., VanDerWal, J., Zamudio, K.R. & Graham, C.H.
(2012) Latitude, elevational climatic zonation, and speciation in New World vertebrates. *Proceedings of the*
Royal Society of London B, **279**, 194-201.
Campagna, L., St. Clair, J.H., Lougheed, S.C., Woods, R.W., Imberti, S. & Tubaro, P.L. (2012) Divergence between
passerine populations from the Malvinas – Falkland Islands and their continental counterparts: a
comparative phylogeographical study. *Biological Journal of the Linnean Society*, **106**, 865-879.
Caro, L.M., Caycedo-Rosales, P.C., Bowie, R.C.K., Slabbekoorn, H. & Cadena, C.D. (2013) Ecological speciation
along an elevational gradient in a tropical passerine bird? *Journal of Evolutionary Biology*, **26**, 357-374.
Carstens, B., Pelletier, T.A., Reid, N.M. & Satler, J.D. (2013) How to fail at species delimitation. *Molecular Ecology*,
22, 4369-4383.
Chaves, J.A., Weir, J.T. & Smith, T.B. (2011) Diversification in *Adelomyia* hummingbirds follows Andean uplift.
Molecular Ecology, **20**, 4564-4576.

Claramunt, S. & Cracraft, J. (2015) A new time tree reveals Earth history's imprint on the evolution of modern birds.
Science Advances, **1**, e1501005.
Claramunt, S., Derryberry, E.P., Remsen, J.V., Jr. & Brumfield, R.T. (2012) High dispersal ability inhibits speciation
in a continental radiation of passerine birds. *Proceedings of the Royal Society of London B*, **279**, 1567-1574.
D'Horta, F.M., Cuervo, A.M., Ribas, C.C., Brumfield, R.T. & Miyaki, C.Y. (2013) Phylogeny and comparative
phylogeography of *Sclerurus* (Aves: Furnariidae) reveal constant and cryptic diversification in an old
radiation of rain forest understorey specialists. *Journal of Biogeography*, **40**, 37-49.
de Queiroz, K. (1998) The general lineage concept of species, species criteria, and the process of speciation. *Endless*
forms: species and speciation (ed. by D.J. Howard and S.H. Berlocher), pp. 57-75. Oxford University Press,
New York.
de Queiroz, K. (2007) Species concepts and species delimitation. *Systematic Biology*, **56**, 879-886.
Diamond, J., Gilpin, M.E. & Mayr, E. (1976) Species-distance relation for birds of the Solomon Archipelago, and the
paradox of the great speciators. *Proceedings of the National Academy of Sciences of the United States of*
America, **73**, 2160-2164.
Diamond, J.M. (1973) Distributional ecology of New Guinea birds. *Science*, **179**, 759-769.
Dingle, C., Halfwer, W. & Slabbekoorn, H. (2008) Habitat-dependent song divergence at subspecies level in the grey-
breasted wood-wren. *Journal of Evolutionary Biology*, **21**, 1079-1089.
Dingle, C., Lovette, I.J., Canaday, C. & Smith, T.B. (2006) Elevational zonation and the phylogenetic relationships of
the *Henicorhina* wood-wrens. *Auk*, **123**, 119-134.
Dingle, C., Poelstra, J.W., Halfwerk, W., Brinkhuizen, D.M. & Slabbekoorn, H. (2010) Asymmetric response patterns
to subspecies-specific song differences in allopatry and parapatry in the gray-breasted wood-wren. *Evolution*,
64, 3537-3548.
Drummond, A.J., Suchard, M.A., Xie, D. & Rambaut, A. (2012) Bayesian phylogenetics with BEAUti and the
BEAST 1.7. *Molecular Biology and Evolution*, **29**, 1969-1973.
Dynesius, M. & Jansson, R. (2014) Persistence of within-species lineages: a neglected control of speciation rates.
Evolution, **68**, 923-934.
Fine, P.V.A. (2015) Ecological and evolutionary drivers of geographic variation in species diversity. *Annual Review of*
Ecology, Evolution and Systematics, **46**, 369–392.
Fischer, A.G. (1960) Latitudinal variations in organic diversity. *Evolution*, **14**, 64-81.
Fitzpatrick, J.W., Terborgh, J.W. & Willard, D.E. (1977) A new species of wood-wren from Peru. *Auk*, **94**, 195-201.
Fjeldså, J. (2013) The discovery of new bird species. *Handbook of the Birds of the World, Special Volume: New*
Species and Global Index (ed. by J. Del Hoyo, A. Elliot, J. Sargatal and D.A. Christie), pp. 147-185. Lynx
Edicions, Barcelona.
Fjeldså, J. & Irestedt, M. (2009) Diversification of the South American avifauna: patterns and implications for
conservation in the Andes. *Annals of the Missouri Botanical Garden*, **96**, 398-409.
Fjeldså, J., Bowie, R.C.K. & Rahbek, C. (2012) The role of mountain ranges in the diversification of birds. *Annual*
Review of Ecology, Evolution and Systematics, **43**, 249-265.
Flantua, S.G.A. & Hooghiemstra, H. (2018) Historical connectivity and mountain biodiversity. *Mountains, Climate*
and Biodiversity (ed. by C. Hoorn, A. Perrigo and A. Antonelli), pp. 171-185. John Wiley & Sons.
Francis, A.P. & Currie, D.J. (2003) A globally consistent richness-climate relationship for angiosperms. *American*
Naturalist, **161**, 523-536.
Freeman, B. (2015) Competitive interactions upon secondary contact drive elevational divergence in tropical birds.
American Naturalist, **186**, 470-479.
Galen, S.C. & Witt, C.C. (2014) Diverse avian malaria and other haemosporidian parasites in Andean house wrens:
evidence for regional co-diversification by host-switching. *Journal of Avian Biology*, **45**, 374-386.
García-Moreno, J. & Fjeldså, J. (2000) Chronology and mode of speciation in the Andean avifauna. *Isolated*
Vertebrate Communities in the Tropics (ed. by G. Rheinwald), pp. 25-46. Proc. 4th Int. Symp, Bonn. Zool.
Monogr. 46, Bonn.
Gaston, K.J., Davies, R.G., Orme, C.D.L., Olson, V.A., Thomas, G.H., Ding, T.-S., Rasmussen, P.C., Lennon, J.J.,
Bennett, P.M., Owens, I.P.F. & Blackburn, T.M. (2007) Spatial turnover in the global avifauna. *Proceedings*
of the Royal Society of London B, **274**, 1567-1574.

Gehara, M., Barth, A., Oliveira, E.F., Costa, M.A., Haddad, C.F.B. & Vences, M. (2017) Model-based analyses reveal
insular population diversification and cryptic frog species in the *Ischnocnema parva* complex in the Atlantic
forest of Brazil. *Molecular Phylogenetics and Evolution*, **112**, 68-78.
Gill, F.B. (2014) Species taxonomy of birds: Which null hypothesis? *Auk*, **131**, 150-161.
Graham, C.H., Silva, N. & Velásquez-Tibatá, J. (2010) Evaluating the potential causes of range limits of birds of the
Colombian Andes. *Journal of Biogeography*, **37**, 1863-1875.
Graham, C.H., Carnaval, A.C., Cadena, C.D., Zamudio, K.R., Roberts, T.E., Parra, J.L., McCain, C.M., Bowie,
R.C.K., Moritz, C., Baines, S.B., Schneider, C.J., VanDerWal, J., Rahbek, C., Kozak, K.H. & Sanders, N.J.
(2014) The origin and maintenance of montane diversity: integrating evolutionary and ecological processes.
Ecography, **37**, 711-719.
Graves, G.R. (1982) Speciation in the carbonated flower-piercer (*Diglossa carbonaria*) complex in the Andes. *Condor*,
84, 1-14.
Graves, G.R. (1988) Linearity of geographic range and its possible effect on the population structure of Andean birds.
Auk, **105**, 47-52.
Guangchuang, Y., K., S.D., Huachen, Z., Yi, G. & Tsan-Yuk, L.T. (2017) ggtree: an r package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. *Methods in Ecology and
Evolution*, **8**, 28-36.
Gutiérrez-Pinto, N., Cuervo, A.M., Miranda, J., Pérez-Emán, J.L., Brumfield, R.T. & Cadena, C.D. (2012) Non-
monophyly and deep genetic differentiation across low-elevation barriers in a Neotropical montane bird
(*Basileuterus tristriatus*; Aves: Parulidae). *Molecular Phylogenetics and Evolution*, **64**, 156-165.
Halfwerk, W., Dingle, C., Brinkhuizen, D.M., Poelstra, J.W., Komdeur, J. & Slabbekoorn, H. (2016) Sharp acoustic
boundaries across an altitudinal avian hybrid zone despite asymmetric introgression. *Journal of Evolutionary
Biology*, **29**, 1356-1367.
Harvey, M.G., Aleixo, A., Ribas, C.C. & Brumfield, R.T. (2017a) Habitat association predicts genetic diversity and
population divergence in Amazonian birds. *American Naturalist*, **190**, 631-648.
Harvey, M.G., Seeholzer, G.F., Smith, B.T., Rabosky, D.L., Cuervo, A.M. & Brumfield, R.T. (2017b) Positive
association between population genetic differentiation and speciation rates in New World birds. *Proceedings
of the National Academy of Sciences of the United States of America*, **114**, 6328-6333.
Hawkins, B.A., Diniz-Filho, J.A.F., Jaramillo, C.A. & Soeller, S.A. (2006) Post-Eocene climate change, niche
conservatism, and the latitudinal diversity gradient of New World birds. *Journal of Biogeography*, **33**, 770-
780.
Hawkins, B.A., Field, R., Cornell, H.V., Currie, D.J., Guégan, J.-F., Kaufman, D.M., Kerr, J.T., Mittlebach, G.G.,
Oberdorff, T., O'Brien, E.M., Porter, E.E. & Turner, J.R.G. (2003) Energy, water, and broad-scale
geographic patterns of species richness. *Ecology*, **84**, 3105-3117.
Hazzi, N.A., Moreno, J.S., Ortiz-Movliav, C. & Palacio, R.D. (2018) Biogeographic regions and events of isolation and
diversification of the endemic biota of the tropical Andes. *Proceedings of the National Academy of Sciences*,
Hosner, P.A., Tobias, J.A., Braun, E.L. & Kimball, R.T. (2017) How do seemingly non-vagile clades accomplish trans-
marine dispersal? Trait and dispersal evolution in the landfowl (Aves: Galliformes). *Proceedings of the Royal
Society of London B*, **284**, 20170210.
Isler, M.L., Cuervo, A.M., Bravo, G.A. & Brumfield, R.T. (2012) An integrative approach to species-level systematics
reveals the depth of diversification in an Andean thamnophilid, the Long-tailed Antbird. *Condor*, **114**, 571-
583.
Jankowski, J.E., Robinson, S.K. & Levey, D.J. (2010) Squeezed at the top: Interspecific aggression may constrain
elevational ranges in tropical birds. *Ecology*, **91**, 1877-1884.
Jankowski, J.E., Ciecka, A.L., Meyer, N.Y. & Rabenold, K.N. (2009) Beta diversity along environmental gradients:
implications of habitat specialization in tropical montane landscapes. *Journal of Animal Ecology*, **78**, 315-327.
Jetz, W. & Rahbek, C. (2002) Geographic range size and determinants of avian species richness. *Science*, **297**, 1548-
1551.
Jetz, W., Rahbek, C. & Colwell, R.K. (2004) The coincidence of rarity and richness and the potential signature of
history in centres of endemism. *Ecology Letters*, **7**, 1180-1191.
Jetz, W., Thomas, G.H., Joy, J.B., Hartmann, K. & Mooers, A.O. (2012) The global diversity of birds in space and
time. *Nature*, **491**, 444-448.

Jønsson, K.A., Irestedt, M., Christidis, L., Clegg, S.M., Holt, B.G. & Fjeldså, J. (2014) Evidence of taxon cycles in an
Indo-Pacific passerine bird radiation (Aves: *Pachycephala*). *Proceedings of the Royal Society B: Biological*
Sciences, **281**, 20131727.
Joseph, L., Wilke, T., Bermingham, E., Alpers, D. & Ricklefs, R. (2004) Towards a phylogenetic framework for the
evolution of shakes, rattles, and rolls in *Myiarchus* tyrant-flycatchers (Aves: Passeriformes: Tyrannidae).
Molecular Phylogenetics and Evolution, **31**, 139-152.
Kapli, P., Lutteropp, S., Zhang, J., Kobert, K., Pavlidis, P., Stamatakis, A. & Flouri, T. (2017) Multi-rate Poisson
tree processes for single-locus species delimitation under maximum likelihood and Markov chain Monte Carlo.
Bioinformatics, **33**, 1630-1638.
Kattan, G.H., Franco, P., Rojas, V. & Morales, G. (2004) Biological diversification in a complex region: a spatial
analysis of faunistic diversity and biogeography of the Andes of Colombia. *Journal of Biogeography*, **31**, 1829-
1839.
Kearns, A.M., Restani, M., Szabo, I., Schrøder-Nielsen, A., Kim, J.A., Richardson, H.M., Marzluff, J.M., Fleischer,
R.C., Johnsen, A. & Omland, K.E. (2018) Genomic evidence of speciation reversal in ravens. *Nature*
Communications, **9**, 906.
Kroodsma, D.E. & Brewer, D. (2005) Family Troglodytidae (wrens). *Handbook of birds of the world: Volume 10,*
Cuckoo-shrikes to thrushes (ed. by J. Del Hoyo, A. Elliot and D. Christie), pp. 356-447. Lynx Edicions,
Barcelona.
Laiolo, P., Pato, J. & Obeso, J.R. (2018) Ecological and evolutionary drivers of the elevational gradient of diversity.
Ecology Letters, **21**, 1022-1032.
Madriñán, S., Cortés, A.J. & Richardson, J.E. (2013) Páramo is the world's fastest evolving and coolest biodiversity
hotspot. *Frontiers in Genetics*, **4**, 192.
Martin, P.R. & Tewksbury, J.J. (2008) Latitudinal variation in subspecific diversification of birds. *Evolution*, **62**, 2775-
2788.
Mayr, E. (1942) *Systematics and the origin of species.* Columbia University Press, New York.
Mayr, E. (1946) The number of species of birds. *Auk*, **63**, 64-69.
Mayr, E. (1963) *Animal species and evolution.* Harvard University Press, Cambridge, Massachusetts.
McCain, C.M. (2009) Global analysis of bird elevational diversity. *Global Ecology and Biogeography*, **18**, 346-360.
Moore, R.P., Robinson, W.D., Lovette, I.J. & Robinson, T.R. (2008) Experimental evidence for extreme dispersal
limitation in tropical forest birds. *Ecology Letters*, **11**, 960-968.
Moyle, R.G., Oliveros, C.H., Andersen, M.J., Hosner, P.A., Benz, B.W., Manthey, J.D., Travers, S.L., Brown, R.M. &
Faircloth, B.C. (2016) Tectonic collision and uplift of Wallacea triggered the global songbird radiation.
Nature Communications, **7**, 12709.
Orme, C.D.L., Davies, R.G., Burgess, M., Eigenbrod, F., Pickup, N., Olson, V.A., Webster, A.J., Ding, T.-S.,
Rasmussen, P.C., Ridgely, R.S., Stattersfield, A.J., Bennett, P.M., Blackburn, T.M., Gaston, K.J. & Owens,
I.P.F. (2005) Global hotspots of species richness are not congruent with endemism or threat. *Nature*, **436**,
1016-1019.
Paradis, E., Claude, J. & Strimmer, K. (2004) APE: Analyses of Phylogenetics and Evolution in R language.
Bioinformatics, **20**, 289-290.
Parra, J.L., Remsen Jr., J.V., Alvarez-Rebolledo, M. & McGuire, J.A. (2009) Molecular phylogenetics of the
hummingbird genus *Coeligena*. *Molecular Phylogenetics and Evolution*, **53**, 425-434.
Patterson, B.D., Stotz, D.F., Solari, S., Fitzpatrick, J.W. & Pacheco, V. (1998) Contrasting patterns of elevational
zonation for birds and mammals in the Andes of southeastern Peru. *Journal of Biogeography*, **25**, 593-607.
Patton, J.L. & Smith, M.F. (1992) mtDNA phylogeny of Andean mice: a test of diversification across ecological
gradients. *Evolution*, **46**, 174-183.
Pérez-Emán, J.L. (2005) Molecular phylogenetics and biogeography of the Neotropical redstarts (*Myioborus*; Aves,
Parulinae). *Molecular Phylogenetics and Evolution*, **37**, 511-528.
Peters, M.K., Hemp, A., Appelhans, T., Behler, C., Classen, A., Detsch, F., Ensslin, A., Ferger, S.W., Frederiksen,
S.B., Gebert, F., Haas, M., Helbig-Bonitz, M., Hemp, C., Kindeketa, W.J., Mwangomo, E., Ngereza, C.,
Otte, I., Röder, J., Rutten, G., Costa, D.S., Tardanico, J., Zancolli, G., Deckert, J., Eardley, C.D., Peters,
R.S., Rödel, M.-O., Schleuning, M., Ssymank, A., Kakengi, V., Zhang, J., Böhning-Gaese, K., Brandl, R.,
Kalko, E.K.V., Kleyer, M., Nauss, T., Tschapka, M., Fischer, M. & Steffan-Dewenter, I. (2016) Predictors of

elevational biodiversity gradients change from single taxa to the multi-taxa community level. *Nature*
Communications, **7**, 13736.
Pigot, A.L. & Tobias, J. (2013) Species interactions constrain geographic range expansion over evolutionary time.
Ecology Letters, **16**, 330-338.
Pigot, A.L. & Tobias, J.A. (2015) Dispersal and the transition to sympatry in vertebrates. *Proceedings of the Royal*
Society of London B, **282**, 20141929.
Posada, D. (2008) jModelTest: phylogenetic model averaging. *Molecular Biology and Evolution*, **25**, 1253-1256.
Price, T.D., Hooper, D.M., Buchanan, C.D., Johansson, U.S., Tietze, D.T., Alström, P., Olsson, U., Ghosh-Harihar,
M., Ishtiaq, F., Gupta, S.K., Martens, J., Harr, B., Singh, P. & Mohan, D. (2014) Niche filling slows the
diversification of Himalayan songbirds. *Nature*, **509**, 222-225.
Prieto-Torres, D.A., Cuervo, A.M. & Bonaccorso, E. (2018) On geographic barriers and Pleistocene glaciations:
Tracing the diversification of the Russet-crowned Warbler (*Myiothlypis coronata*) along the Andes. *PLoS*
One, **13**, e0191598.
Pulido-Santacruz, P. & Weir, J.T. (2016) Extinction as a driver of avian latitudinal diversity gradients. *Evolution*, **70**,
860-872.
Pulido-Santacruz, P., Aleixo, A. & Weir, J.T. (2018) Morphologically cryptic Amazonian bird species pairs exhibit
strong postzygotic reproductive isolation. *Proceedings of the Royal Society of London B*, **285**, 20172081.
Pybus, O.G., Rambaut, A. & Harvey, P.H. (2000) An integrated framework for the inference of viral population
history from reconstructed genealogies. *Genetics*, **155**, 1429-1437.
Pyron, R.A. (2014) Temperate extinction in squamate reptiles and the roots of latitudinal diversity gradients. *Global*
Ecology and Biogeography, **23**, 1126-1134.
Pyron, R.A. & Wiens, J.J. (2013) Large-scale phylogenetic analyses reveal the causes of high tropical amphibian
diversity. *Proceedings of the Royal Society of London B*, **280**, 20131622.
Quintero, I. & Jetz, W. (2018) Global elevational diversity and diversification of birds. *Nature*, **555**, 246-250.
Rabosky, D.L. (2016) Reproductive isolation and the causes of speciation rate variation in nature. *Biological Journal*
of the Linnean Society, **118**, 13-25.
Rabosky, D.L. & Hurlbert, A.H. (2015) Species richness at continental scales Is dominated by ecological limits. *The*
American Naturalist, **185**, 572-583.
Rabosky, D.L., Santini, F., Eastman, J., Smith, S.A., Sidlauskas, B., Chang, J. & Alfaro, M.E. (2013) Rates of
speciation and morphological evolution are correlated across the largest vertebrate radiation. *Nature*
Communications, **4**, 1958.
Rabosky, D.L., Chang, J., Title, P.O., Cowman, P.F., Sallan, L., Friedman, M., Kaschner, K., Garilao, C., Near, T.J.,
Coll, M. & Alfaro, M.E. (2018) An inverse latitudinal gradient in speciation rate for marine fishes. *Nature*,
559, 392-395.
Rahbek, C. & Graves, G.R. (2001) Multiscale assessment of patterns of avian species richness. *Proceedings of the*
National Academy of Sciences of the USA, **98**, 4534-4539.
Rahbek, C., Gotelli, N.J., Colwell, R.K., Entsminger, G.L., Rangel, T.F.L.V.B. & Graves, G.R. (2007) Predicting
continental-scale patterns of bird species richness with spatially explicit models. *Proceedings of the Royal*
Society of London B, **274**, 165-174.
Ramírez-Barahona, S. & Eguiarte, L.E. (2013) The role of glacial cycles in promoting genetic diversity in the
Neotropics: the case of cloud forests during the Last Glacial Maximum. *Ecology and Evolution*, **3**, 725-738.
Reid, N.M. & Carstens, B.C. (2012) Phylogenetic estimation error can decrease the accuracy of species delimitation: a
Bayesian implementation of the general mixed Yule-coalescent model. *BMC Evolutionary Biology*, **12**, 196.
Remsen, J.V., Jr. (2016) A "rapid assessment program" for assigning species rank? *Journal of Field Ornithology*, **87**,
110-115.
Remsen, J.V., Jr. & Graves, W.S. (1995) Distribution patterns and zoogeography of *Atlapetes* brush-finches
(Emberizinae) of the Andes. *Auk*, **112**, 210-224.
Remsen, J.V., Jr., Areta, J.I., Cadena, C.D., Claramunt, S., Jaramillo, A., Pacheco, J.F., Robbins, M.B., Stiles, F.G.,
Stotz, D.F. & Zimmer, K.J. (2018) *A classification of the bird species of South America, version July 7 2018*.
American Ornithologists' Union. http://www.museum.lsu.edu/~Remsen/SACCBaseline.htm.
Revell, L.J. (2012) phytools: an R package for phylogenetic comparative biology (and other things). *Methods in*
Ecology and Evolution, **3**, 217-223.

Ribas, C.C., Moyle, R.G., Miyaki, C.Y. & Cracraft, J. (2007) The assembly of montane biotas: linking Andean
tectonics and climatic oscillations to independent regimes of diversification in *Pionus* parrots. *Proceedings of*
the Royal Society of London B, **274**, 2399-2408.

Ricklefs, R.E. (2006) Global variation in the diversification rate of passerine birds. *Ecology*, **87**, 2468-2478.

Ricklefs, R.E. & Bermingham, E. (1999) Taxon cycles in the Lesser Antillean avifauna. *Ostrich*, **70**, 49-59.

Ricklefs, R.E. & Bermingham, E. (2002) The concept of the taxon cycle in biogeography. *Global Ecology and*
Biogeography, **11**, 353-361.

Ricklefs, R.E. & Bermingham, E. (2007) The causes of evolutionary radiations in archipelagoes: passerine birds in the
Lesser Antilles. *American Naturalist*, **169**, 285-297.

Ritchie, A.M., Lo, N. & Ho, S.Y.W. (2017) The Impact of the tree prior on molecular dating of data sets containing a
mixture of inter- and intraspecies sampling. *Systematic Biology*,

Robbins, M.B. & Nyári, Á.S. (2014) Canada to Tierra del Fuego: Species limits and historical biogeography of the
Sedge Wren (*Cistothorus platensis*). *Wilson Journal of Ornithology*, **126**, 649-662.

Rolland, J., Condamine, F.L., Jiguet, F. & Morlon, H. (2014) Faster speciation and reduced extinction in the tropics
contribute to the mammalian latitudinal diversity gradient. *PLoS Biology*, **12**, e1001775.

Roux, C., Fraïsse, C., Romiguier, J., Anciaux, Y., Galtier, N. & Bierne, N. (2016) Shedding light on the grey zone of
speciation along a continuum of genomic divergence. *PLoS Biology*, **14**, e2000234.

Ruggiero, A. & Hawkins, B.A. (2008) Why do mountains support so many species of birds? *Ecography*, **31**, 306-315.

Salaman, P., Coopmans, P., Donegan, T.M., Mulligan, M., Cortés, A., Hilty, S.L. & Ortega, L.A. (2003) A new
species of wood-wren (Troglodytidae: *Henicorhina*) from the Western Andes of Colombia. *Ornitología*
Colombiana, **1**, 4-21.

Salisbury, C.L., Seddon, N., Cooney, C.R. & Tobias, J.A. (2012) The latitudinal gradient in dispersal constraints:
ecological specialisation drives diversification in tropical birds. *Ecology Letters*, **15**, 847-855.

Sánchez-González, L.A., Navarro-Sigüenza, A.G., Krabbe, N.K., Fjeldså, J. & García-Moreno, J. (2015) Diversification
in the Andes: the *Atlapetes* brush-finches. *Zoologica Scripta*, **44**, 135-152.

Scheffers, B.R., Joppa, L.N., Pimm, S.L. & Laurance, W.F. (2012) What we know and don't know about Earth's
missing biodiversity. *Trends in Ecology and Evolution*, **27**, 501-510.

Schluter, D. (2016) Speciation, ecological opportunity, and latitude. *American Naturalist*, **187**, 1-18.

Schluter, D. & Pennell, M.W. (2017) Speciation gradients and the distribution of biodiversity. *Nature*, **546**, 48-55.

Sedano, R.E. & Burns, K.J. (2010) Are the Northern Andes a species pump for Neotropical birds? Phylogenetics and
biogeography of a clade of Neotropical tanagers (Aves: Thraupini). *Journal of Biogeography*, **37**, 325-343.

Seeholzer, G.F. & Brumfield, R.T. (2018) Isolation by distance, not incipient ecological speciation, explains genetic
differentiation in an Andean songbird (Aves: Furnariidae: *Cranioleuca antisiensis*, Line-cheeked Spinetail)
despite near threefold body size change across an environmental gradient. *Molecular Ecology*, **27**, 279-296.

Smith, B.T. & Klicka, J. (2010) The profound influence of the Late Pliocene Panamanian uplift on the exchange,
diversification, and distribution of New World birds. *Ecography*, **33**, 333-342.

Smith, B.T., Seeholzer, G.F., Harvey, M.G., Cuervo, A.M. & Brumfield, R.T. (2017) A latitudinal phylogeographic
diversity gradient in birds. *PLoS Biology*, **15**, e2001073.

Smith, B.T., Bryson, J., Robert W., Mauck, W.M., Chaves, J., Robbins, M.B., Aleixo, A. & Klicka, J. (2018) Species
delimitation and biogeography of the gnatcatchers and gnatwrens (Aves: Polioptilidae). *Molecular*
Phylogenetics and Evolution, **126**, 45-57.

Smith, B.T., McCormack, J.E., Cuervo, A.M., Hickerson, M.J., Aleixo, A., Cadena, C.D., Pérez-Emán, J., Burney,
C.W., Xie, X., Harvey, M.G., Faircloth, B.C., Glenn, T.C., Derryberry, E.P., Prejean, J., Fields, S. &
Brumfield, R.T. (2014) The drivers of tropical speciation. *Nature*, **515**, 406-409.

Socolar, J.B., Gilroy, J.J., Kunin, W.E. & Edwards, D.P. (2016) How should beta-diversity inform biodiversity
conservation? *Trends in Ecology and Evolution*, **31**, 67-80.

Stamatakis, A. (2006) RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and
mixed models. *Bioinformatics*, **22**, 2688-2690.

Stattersfield, A.J., Crosby, M.J., Long, A.J. & Wege, D.C. (1997) *Endemic bird areas of the world: priorities for*
biodiversity conservation. BirdLife International (Conservation Series No. 7), Cambridge, UK.

Sukumaran, J. & Knowles, L.L. (2017) Multispecies coalescent delimits structure, not species. *Proceedings of the*
National Academy of Sciences of the USA, **114**, 1607-1612.

Terborgh, J. (1971) Distribution on environmental gradients: theory and a preliminary interpretation of distributional
patterns in the avifauna of the Cordillera Vilcabamba, Peru. *Ecology*, **52**, 23-40.
Terborgh, J. (1977) Bird species diversity on an Andean elevational gradient. *Ecology*, **58**, 1007-1019.
Tobias, J.A., Seddon, N., Spottiswoode, C.N., Pilgrim, J.D., Fishpool, L.D.C. & Collar, N.J. (2010) Quantitative
criteria for species delimitation. *Ibis*, **152**, 724-746.
Tobias, J.A., Cornwallis, C.K., Derryberry, E.P., Claramunt, S., Brumfield, R.T. & Seddon, N. (2014) Species
coexistence and the dynamics of phenotypic evolution in adaptive radiation. *Nature*, **506**, 359-363.
Toews, D.P.L. (2015) Biological species and taxonomic species: Will a new null hypothesis help? (A comment on Gill
2014). *Auk*, **132**, 78-81.
Valderrama, E., Pérez-Emán, J.L., Brumfield, R.T., Cuervo, A.M. & Cadena, C.D. (2014) The influence of the
complex topography and dynamic history of the montane Neotropics on the evolutionary differentiation of a
cloud forest bird (*Premnoplex brunnescens*, Furnariidae). *Journal of Biogeography*, **41**, 1533-1546.
Velásquez-Tibatá, J., Salaman, P. & Graham, C.H. (2013) Effects of climate change on species distribution,
community structure, and conservation of birds in protected areas in Colombia. *Regional Environmental*
Change, **13**, 235-248.
Weir, J.T. (2006) Divergent timing and patterns of species accumulation in lowland and highland neotropical birds.
Evolution, **60**, 842-855.
Weir, J.T. (2009) Implications of genetic differentiation in Neotropical montane forest birds. *Annals of the Missouri*
Botanical Garden, **96**, 410-433.
Weir, J.T. & Schluter, D. (2007) The latitudinal gradient in recent speciation and extinction rates of birds and
mammals. *Science*, **315**, 1574-1576.
Weir, J.T., Bermingham, E., Miller, M.J., Klicka, J. & González, M.A. (2008) Phylogeography of a morphologically
diverse Neotropical montane species, the Common Bush-Tanager (*Chlorospingus ophthalmicus*). *Molecular*
Phylogenetics and Evolution, **47**, 650-664.
Wilson, E.O. (1959) Adaptive shift and dispersal in a tropical ant fauna. *Evolution*, **13**, 122-144.
Winger, B.M. (2017) Consequences of divergence and introgression for speciation in Andean cloud forest birds.
Evolution, **71**, 1815-1831.
Winger, B.M. & Bates, J.M. (2015) The tempo of trait divergence in geographic isolation: Avian speciation across the
Marañon Valley of Peru. *Evolution*, **69**, 772-787.

903
904

Figure 1. Geographic distribution of wood-wrens in the *Henicorhina leuocophrys* complex in the Neotropical montane
region and localities where specimens were sampled for our phylogeographic analyses. The complex currently
comprises three species: the widely distributed *H. leucophrys* ranging from Mexico to Bolivia and two narrow endemics
from Colombia (*H. anachoreta* in the Sierra Nevada de Santa Marta and *H. negreti* on the western slope of the
Cordillera Occidental). Distribution maps were obtained from BirdLife International (*H. leucophrys*) and Velásquez-
Tibatá et al. (2013; *H. negreti*), or generated for this study based on information on elevational range (*H. anachoreta*;
Cadena *et al.*, 2016). Illustrations by F. Ayerbe.

Figure 2. Phylogenetic relationships and divergence times among unique mtDNA haplotypes in the *H. leucophrys*
complex inferred using BEAST suggest (1) paraphyly of *H. leucophrys* with respect to *H. anachoreta* and *H. negreti,*
(2) a northern origin for the complex with subsequent colonization of South America, and (3) marked population
genetic structure partly attributable to geographic isolation mediated by physical barriers. Color shading on nodes
corresponds to posterior probabilities ≥ 0.85. Black vertical lines indicate geographic regions; note that all deep
branches correspond to clades from mountain regions in Mexico and Central America and that South American
populations are also strongly structured. Green vertical lines signal presumptive species identified using multi-rate
Poisson Tree Processes (mPTP, left) and the Bayesian General Mixed Yule Coalescent Model (bGMYC, right); results
of these analyses were almost identical, with only minor discrepancies in two clades marked with asterisks. Haplotypes
are numbered at the tips of the tree; information on specimens having each haplotype is provided in Supplementary
Table 1.

Figure 3. Elevational replacements involving distinct lineages of the *H. leucophrys* complex in montane South America. Lineages replacing each other with elevation in a region share colors in the map, the phylogeny (modified from Supplementary Figure 1, nodes indicated with grey dots have ≥0.85 posterior probability support), and the close-up view of mountain slopes, where different shades are used for each lineage (arrows on the map show the direction from which mountains are seen in panes A-D). In at least three regions (A-C), elevational replacements do not involve sister taxa with the only possible exception being the novel case of cryptic replacement of lineages in the Cordillera de Mérida, Venezuela (D), where affinities of lineages to each other and to those from other regions are not strongly supported. Lineages involved in elevational replacements are deeply divergent from each other (panes show mean uncorrected *p* distances in ATPase genes), occur in close proximity, and correspond to different presumptive species identified by coalescent analyses (see text).

Figure 4. Geographic locations where we sampled 39 presumptive species in the *H. leucophrys* complex identified by coalescent analyses of mtDNA sequences. Dots and numbers on the tree (modified from Figure 2) correspond to species statistically inferred by the Bayesian General Mixed Yule Coalescent Model (bGMYC) with the threshold probability used to define group membership set at 0.50. Colors correspond to species epithets as per the current three-species taxonomy. Encircled numbers on the map indicate the number of presumptive species occurring in each country. Almost identical patterns were observed using the multi-rate Poisson Tree Processes (mPTP) method. For close-up views of geographic locations where each presumptive species was sampled, see Supplementary Figure 4.

Figure 5. Lineage-through time plot showing accumulation of lineages over time in the *H. leucophrys* complex. Black
lines are estimates based on the maximum clade credibility tree and grey indicates the 95% credibility interval across
100 trees for the plot and for estimates of the gamma statistic. The shape of the curve and the associated gamma
statistic suggests that rates of lineage accumulation have declined over time, but note that because this analyses used
results of bGMYC as input no presumptive species younger than 1 million years were considered. The pattern may
also reflect incomplete sampling of young lineages particularly within South America.

Figure 6. Relative to the frequency distribution of the number of presumptive species identified by coalescent analyses of mtDNA sequence data within 173 taxonomic species of New World birds (data from Harvey et al. 2017, Smith et al. 2017), our result for the *H. leucophrys* complex is a distant outlier. Even if many lineages cannot be shown to be reproductively isolated from others, the data reveal a remarkable and previously undocumented degree of population genetic structuring. Illustration by F. Ayerbe.

Supplementary Figure 1. Phylogenetic relationships among individuals in the *H. leucophrys* complex inferred using
Bayesian analysis of sequences of the ATPase 6&8 mitochondrial genes. The phylogeny is the maximum clade
credibility tree obtained in BEAST. Nodal support (i.e, posterior probabilities ≥ 0.85) is shown using a grey scale.
Nodes with a colored outline (green = *H. leucophrys*, blue = *H. negreti*, orange = *H. anachoreta*) were also recovered
with strong support (≥ 80% boostrap) in maximum-likelihood analysis (Supplementary Figure 2).

978   **Supplementary Figure 2.** Phylogenetic relationships among individuals in the *H. leucophrys* complex inferred using
979   maximum-likelihood analysis of sequences of the ATPase 6&8 mitochondrial genes. The phylogeny is the maximum-
980   likelihood tree obtained in RAxML. Nodal support (i.e, boostrap values ≥ 80%) is shown using a grey scale. Nodes
981   with a colored outline (green = *H. leucophrys*, blue = *H. negreti*, orange = *H. anachoreta*) were also recovered with
982   strong support (≥ 0.90 posterior probability) in Bayesian analysis (Supplementary Figure 1).

Supplementary Figure 3. Results of species delimitation analysis in the *H. leucophrys* complex employing the Bayesian General Mixed Yule Coalescent Model (bGMYC). The phylogeny showing relationships among haplotypes is the maximum clade credibility obtained using BEAST and the table to the right is a sequence-by-sequence matrix in which cells are color-coded to indicate the posterior probability that each pair of sequences is conspecific.

Supplementary Figure 4. Close-up views of geographic locations where presumptive species in the *H. leucophrys* complex identified by coalescent analyses of mtDNA sequences were sampled. Maps showing known locations of each presumptive species are ordered roughly from North to South, and are numbered according to numbers on nodes in the tree in Figure 4; points on maps are colored based on the current three-species taxonomy recognizing *H. leucophrys*, *H. negreti*, and *H. anachoreta.* Some presumptive species have relatively large ranges (e.g. no. 6 across Costa Rica and Panama) whereas others appear to be much more restricted, in some cases found at single localities so far (e.g. no. 14 and 15 in the northern extreme of the Cordillera Central in Antioquia, Colombia).

Supplementary Figure 5. Number of mtDNA lineages in the *H. leucophrys* complex of various ages (from 5 to 1 million years ago [m.a.]). Vertical black lines correspond to the median number of lineages dating to at least each of the four ages (i.e. splitting from their common ancestor with other lineages before each age) observed in a sample of 100 trees in the posterior distribution obtained using BEAST; gray bars are the frequency distributions of number of lineages per age across all trees.

1012   **Supplementary Table 1.** (Provided as a separate .xlsx file). Information on specimens considered in phylogeographic

1013   analyses including museum catalogue numbers, locality data, and GenBank accession numbers when available (those

1014   for sequences generated for this study are pending). For each specimen, we also indicate the name used to refer to it in

1015   Supplementary Figures 1 and 2, the ATPase 6/8 haplotype as shown in Figure 2, and the ID of the presumptive

1016   species identified using bGMYC to which it belongs (Figure 3, Supplementary Figure 4).