Divergent receptor proteins confer responses to different karrikins in two ephemeral weeds

Yueming Kelly Sun[1,2], Jiaren Yao[1], Adrian Scaffidi[1], Kim T. Melville[1], Sabrina F Davies[1], Charles S Bond[1], Steven M Smith[3,4], Gavin R Flematti[1] and Mark T Waters[1,2,5]

1. School of Molecular Sciences, The University of Western Australia, 35 Stirling Hwy, Perth, WA 6009, Australia
2. Australian Research Council Centre of Excellence in Plant Energy Biology, The University of Western Australia, 35 Stirling Hwy, Perth, WA 6009, Australia
3. School of Natural Sciences, The University of Tasmania, Hobart, TAS 7000, Australia
4. Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, 1 West Beichen Road, Beijing 100101, PR China.
5. Author for correspondence: mark.waters@uwa.edu.au

Word count (Abstract, main text and methods): 7505 words

ABSTRACT

Wildfires can encourage the establishment of invasive plants by releasing potent germination stimulants, such as karrikins. Seed germination of *Brassica tournefortii*, a noxious weed of Mediterranean climates, is strongly stimulated by KAR_1, which is the archetypal karrikin produced from burning vegetation. In contrast, the closely-related yet non-fire-associated ephemeral *Arabidopsis thaliana* is unusual because it responds preferentially to KAR_2. The α/β-hydrolase KARRIKIN INSENSITIVE2 (KAI2) is the putative karrikin receptor identified in *Arabidopsis*. Here we show that *B. tournefortii* differentially expresses three *KAI2* homologues, and the most highly-expressed homologue is sufficient to confer enhanced responses to KAR_1 relative to KAR_2 when expressed in *Arabidopsis*. We further identify two variant amino acid residues near the KAI2 active site that explain the ligand selectivity, and show that this combination has arisen independently multiple times within dicots. Our results suggest that duplication and diversification of KAI2 proteins could confer upon weedy ephemerals and potentially other angiosperms differential responses to chemical cues produced by environmental disturbance, including fire. (162 words)

INTRODUCTION

Environmental disturbance promotes the establishment of invasive species, posing a potent threat to global biodiversity. Changing wildfire regimes, such as increasing frequency of fires, is one of the most relevant disturbance factors contributing to elevated invasion threat[1]. Wildfires create germination opportunities in part by releasing seed germination stimulants, such as karrikins, from burning vegetation[2, 3]. Karrikins comprise a family of butenolides with six known members. In samples of smoke water generated by burning grass straw, KAR_1 is the most abundant karrikin, while KAR_2 is six times less abundant[4], although these proportions may differ depending on source material, preparation method, age and storage conditions[5]. These two analogues differ only by the presence of a methyl group on the butenolide ring in KAR_1, which is absent in KAR_2 (Supplementary Fig. 1a). Invasive plant species that are responsive to karrikins could utilise natural and human-induced fires to facilitate their establishment[6, 7].

Karrikins can overcome seed dormancy and promote seed germination in a number of smoke-responsive species, as well as others that are not associated with fire regimes[8-10].

Furthermore, karrikins also influence light-dependent seedling development. Karrikins

enhance the sensitivity of *Arabidopsis thaliana* seedlings to light by inhibiting hypocotyl

elongation, stimulating cotyledon expansion, and promoting chlorophyll accumulation in a

dose-dependent manner[11]. Karrikins have been reported to enhance seedling survival and

biomass in species such as tomato, rice and maize[12, 13]. Such growth-promoting effects of

karrikins, especially at critical early stages of the life cycle, have the potential to further

encourage the establishment of invasive species after fire events.

Brassica tournefortii (Brassicaceae; Sahara mustard) is native to northern Africa and the

Middle East, but is an invasive weed that blights many ecosystems with a Mediterranean

climate and chaparral-type vegetation that are prone to wildfires in North America, Australia

and South Africa. *B. tournefortii* seeds can persist in the soil for many seasons, undergoing

wet-dry cycling that can influence dormancy and contribute to boom-bust cycles that

outcompete native species[9, 14]. *B. tournefortii* plants may radically alter fire frequency and

intensity by influencing fuel profiles[15, 16], further exacerbating the impact of fire on

susceptible native ecosystems. In addition, seeds of *B. tournefortii* are particularly responsive

to smoke-derived karrikins, and show a positive germination response to KAR_1 in the

nanomolar range[10]. Accordingly, *B. tournefortii* is particularly well positioned to invade areas

disturbed by fire events[17, 18].

The putative karrikin receptor KARRIKIN INSENSITIVE 2 (KAI2) was identified in

Arabidopsis, a weedy ephemeral that originated in Eurasia but is now widely distributed

throughout the northern hemisphere[19-21]. Arabidopsis is not known to colonise fire-prone

habitats, but nevertheless seeds germinate in response to karrikins in the micromolar range[22].

Unlike most smoke-responsive species that respond more readily to KAR_1[8, 23], Arabidopsis

responds preferentially to KAR_2[22]. KAI2 is an evolutionarily ancient α/β-hydrolase and a

paralogue of DWARF14 (D14), the receptor for strigolactones[24, 25]. Karrikins and

strigolactones are chemically similar by virtue of a butenolide moiety that is necessary for

bioactivity[26, 27]. KAI2 and D14 have dual functions as both enzyme and receptor, but the

functional significance of the enzymatic activity remains contested[28-32]. Furthermore, the

basis for ligand specificity by these two highly congruent proteins remains essentially

unknown.

Orthologues of *KAI2* are ubiquitous in land plants, and are normally present as a single gene

copy within an ancient and highly conserved "eu-KAI2" clade[33]. There is growing evidence

that, beyond its ability to mediate karrikin responses, KAI2 has a core ancestral role in

perceiving an endogenous KAI2 ligand ("KL") that regulates seed germination, seedling

development, leaf shape and cuticle development[34-36]. Since its divergence from the

Arabidopsis lineage, the tribe Brassiceae, which includes the genus *Brassica*, underwent a

whole genome triplication event 24–29 million years ago[37-39]. This process might have

allowed additional KAI2 copies to gain altered ligand specificity, potentially enhancing

perception of environmental signals such as karrikins from smoke. Here, we report that two

out of three KAI2 homologues expressed in *B. tournefortii* show distinct preferences for

different karrikins. We take advantage of the relatively recent genome triplication event in the

Brassiceae to identify two amino acids that are sufficient to explain these karrikin preferences

and confirm this by mutagenesis. Beyond demonstrating the potential ecological significance

of diversity among KAI2 homologues, our findings also reveal active site regions critical for

ligand selectivity among KAI2 receptor-enzymes that are found in all land plants.

RESULTS

B. tournefortii is most sensitive to KAR_1

To characterise the karrikin response of *B. tournefortii*, we performed multiple physiological

and molecular assays comparing KAR_1 activity with that of KAR_2. First, germination of *B.*

tournefortii seeds was consistently more responsive to KAR_1 than KAR_2 at 10 nM, 100 nM and

1 μM (Fig. 1a and Supplementary Fig. 1b-d). Second, homologues of two karrikin-responsive

transcripts, *DWARF14-LIKE2* (*BtDLK2*) and *SALT TOLERANCE HOMOLOG7* (*BtSTH7*), identified

on the basis of close sequence homology to their Arabidopsis counterparts[11, 20], were

significantly more highly expressed when treated with 1 μM KAR_1 than with 1 μM KAR_2 (Fig.

1b). These observed differences in seed response are not due to differential karrikin uptake,

since both KAR_1 and KAR_2 were taken up from solution at similar rates by *B. tournefortii* seeds

during imbibition, as was also true for Arabidopsis seeds (Supplementary Fig. 1e-f). Besides

promoting germination of primary dormant seeds, karrikins also inhibited hypocotyl

elongation in *B. tournefortii* seedlings, as is the case in Arabidopsis; again, KAR_1 showed a

stronger effect than KAR_2 (Fig. 1c-d). Levels of *BtDLK2* transcripts in seedlings were also more

responsive to KAR_1 than KAR_2 at a given concentration (Fig. 1e). Therefore, we conclude that

B. tournefortii is more sensitive to KAR_1 than to KAR_2, a ligand preference that is a feature of

seed germination in many karrikin-responsive species from ecosystems prone to fires[8, 23].

Two out of three *B. tournefortii* KAI2 homologues are functional in Arabidopsis

To establish whether there are multiple *KAI2* homologues present in *B. tournefortii*, we examined transcriptomes from seeds and seedlings. Three putative *KAI2* homologues were identified (*BtKAI2a*, *BtKAI2b*, and *BtKAI2c*; Fig. 2a; Supplementary Fig. 2 and 3). *BtKAI2a* grouped with *AtKAI2* and those of other Brassicaceae within a single clade, whereas *BtKAI2b* and *BtKAI2c* grouped within a clade unique to *Brassica*. This taxonomic distribution implies that *BtKAI2a* is most similar to *AtKAI2*, and that *BtKAI2b* and *BtKAI2c* are paralogues that arose via genome triplication during the evolution of *Brassica*. All three *BtKAI2* transcripts were expressed in *B. tournefortii* seeds, but only *BtKAI2a* and *BtKAI2b* could be detected in seedlings (Fig. 2b). In seeds and seedlings, *BtKAI2b* transcripts were the most abundant of the three, but there were no consistent effects on any of the transcripts upon treatment with 1 µM KAR$_1$. We also identified two *BtD14* homologues, at least one of which is functionally orthologous to *AtD14* (Supplementary Figs. 2 and 4).

We performed transgenic complementation of the Arabidopsis *kai2-2* null mutant by expressing each of the three isoforms as a GFP fusion protein driven by the Arabidopsis *KAI2* promoter and 5′UTR (*KAI2pro*:*GFP-BtKAI2*) to accurately reflect the native expression profile of KAI2 in Arabidopsis. Such fusions of GFP with KAI2 proteins have been used previously to analyse KAI2 activity[19, 40, 41]. Both *BtKAI2a* and *BtKAI2b* complemented the seedling and leaf phenotypes of *kai2-2*, whereas *BtKAI2c* did not (Fig. 2c-d). GFP-BtKAI2a and GFP-BtKAI2b accumulated at consistent levels when detected using both an anti-KAI2 antibody and an anti-GFP antibody (which negated the effects of sequence differences in the anti-KAI2 epitope region; Fig. 2e and Supplementary Fig. 3). However, we could not detect GFP-BtKAI2c protein in three independent transgenic lines with either antibody. We verified the *GFP-BtKAI2c* transcript sequences expressed in the transgenic plant material by RT-PCR and Sanger sequencing (Supplementary Fig. 5). Since the *GFP-BtKAI2c* mRNA is processed faithfully in Arabidopsis, and the apparent absence of protein is most likely a result of posttranslational events. We then tested whether BtKAI2c was generally poorly expressed in plant cells by transient overexpression in tobacco leaves using *Agrobacterium*-mediated infiltration and tobacco mosaic virus-based plant expression vectors[42]. Even with the extremely high levels of expression supported by this system, levels of myc-tagged BtKAI2c protein were substantially lower than equivalently-tagged BtKAI2a and BtKAI2b (Supplementary Fig. 5). This result implies that BtKAI2c protein may be inherently unstable and/or prone to degradation in plant cells.

 BtKAI2c carries a destabilising mutation at position 98

Numerous mutations identified by forward genetics destabilise AtKAI2 and render the

protein undetectable in plant extracts[43]. Ten residues in the BtKAI2c sequence are unique to

BtKAI2c within the genus *Brassica* (Supplementary Figure 3). We predicted the likely effect of

each of these residues upon protein function, relative to the *Brassica* consensus residues at

the same positions, using the PROVEAN tool[44]. Among the ten sites, R98 and E129 exceeded

the cutoff value of -2.5 and were thus predicted to be highly deleterious to protein function,

while D137 narrowly missed this threshold (Supplementary Table 2). Therefore based on this

predictive analysis, the native BtKAI2c protein carries at least two and possibly three

substitutions that have the potential to render the protein non-functional.

R98 is potentially highly deleterious because it is situated in the active site adjacent to the

catalytic serine, whereas E129 and D137 are located on the flexible hinge that links the lid and

core domains (Supplementary Fig. 6a). In AtKAI2, the G133E mutation is highly

destabilising[20, 41], suggesting that non-conserved substitutions in the hinge region have the

potential to be deleterious as well. To test the effect of these residues on BtKAI2c stability, we

constructed three variants that reverted the native BtKAI2c sequence to consensus amino

acids within *Brassica* at position 98 and/or in the hinge region: BtKAI2c^{R98V}, a quadruple

mutant BtKAI2c$^{E129D;D130V;E133Q;D137E}$ (BtKAI2c^{QUAD}), and a quintuple mutant that combined all

five substitutions (BtKAI2c^{QUINT}). Should these amino acids be crucial to protein stability, we

expected to restore BtKAI2c expression.

First, we tried expressing the native and mutated BtKAI2c variants in tobacco leaves using

Agrobacterium-mediated infiltration. The vectors were based on those used to generate the

GFP-BtKAI2c Arabidopsis transgenics, allowing us to compare expression of the same GFP

fusion proteins. As was the case for stably-transformed Arabidopsis, we found that native

GFP-BtKAI2c protein was undetectable, as was the BtKAI2c^{QUAD} variant (Supplementary Fig.

6b). However, BtKAI2c^{R98V} and BtKAI2c^{QUINT} were clearly detected. We also examined the

effect of the equivalent V96R mutation on GFP-AtKAI2, and found that this mutation

completely abolished expression compared to the native GFP-AtKAI2 control (Supplemental

Fig. 6b). These results suggest that the R98 residue causes the loss of protein expression in

BtKAI2c, and that arginine at this position is not tolerated by KAI2 proteins in general.

Early in our investigation, we had tried without success to express and purify BtKAI2c from *E.*

coli for biochemical characterisation. Having restored BtKAI2c expression in plants with the

R98V mutation, we next tried to express these proteins in bacteria. We found that SUMO-
BtKAI2c R98V was expressed at ~10-fold higher levels than native SUMO-BtKAI2c in crude
lysates, suggesting that the R98V mutation enhanced protein folding and/or stability in
bacterial cells. Although it was detectable in crude lysates, SUMO-BtKAI2c was consistently
intransigent to recovery following affinity chromatography; in contrast, the R98V and
quintuple mutant versions were stably recovered at high levels and purity (Supplemental
Figure 6c). From these results, we conclude that native BtKAI2c is non-functional due to
protein instability induced by a highly non-conservative substitution at the active site.

BtKAI2a and BtKAI2b show differential ligand specificity

We further characterised the ligand specificity of *BtKAI2* homologues by performing
physiological and molecular assays with the stable transgenic Arabidopsis lines. Primary-
dormant Arabidopsis seeds homozygous for the transgenes were tested for germination
response (Fig. 3a and Supplementary Fig. 7). None of the transgenic lines completely restored
germination levels to those of wild type, perhaps reflecting the need for additional, seed-
specific regulatory elements not included in our transgenes, or the importance of genomic
context for proper *KAI2* expression in seeds. Nevertheless, germination responses to karrikins
were evident in the transgenic seeds: germination of *BtKAI2b* seeds was more responsive to
KAR_1 than KAR_2 at 1 μM, whereas no significant difference was observed for *BtKAI2a* seeds. As
expected, *BtKAI2c* seeds were indistinguishable from the *kai2* control and insensitive to
karrikins. We also found that levels of *KUF1* transcripts, which are responsive to karrikins in
Arabidopsis seeds (Nelson 2010), were only slightly induced by karrikins in *BtKAI2a*
transgenic seeds, but were strongly enhanced by KAR_1 in *BtKAI2b* seeds (Fig. 3b).

In seedlings, both *BtKAI2a* and *BtKAI2b*, but not *BtKAI2c*, could complement the *kai2*
hypocotyl elongation phenotype (Fig 3c). Responses to karrikins in terms of hypocotyl
elongation (Fig. 3c) and levels of *DLK2* transcripts (Fig. 3d) broadly agreed with the
germination response with respect to karrikin preferences, with *BtKAI2b* transgenics showing
a clear preference for KAR_1 over KAR_2. However, *BtKAI2a* seedlings showed a preference for
KAR_2 that was not evident in seeds. From these experiments, we conclude that BtKAI2a is
similar to AtKAI2 in terms of ligand specificity, whereas BtKAI2b has a clear preference for
KAR_1 over KAR_2. As *BtKAI2b* is more highly expressed than *BtKAI2a* in *B. tournefortii* seeds
and seedlings (Fig. 2b), we conclude that the ligand specificity of BtKAI2b substantially

contributes to the enhanced KAR$_1$-responsiveness of this species at these stages of the life
cycle.

Positions 98 and 191 account for ligand specificity between BtKAI2a and BtKAI2b

To investigate interactions between BtKAI2 homologues and ligands, we performed
differential scanning fluorimetry (DSF) assays on purified recombinant proteins
(Supplementary Fig. 8). DSF has been used extensively for inferring the interaction of
strigolactone-like compounds with D14- and KAI2-related proteins[29, 41, 45-48]. Racemic GR24 is
a widely-used synthetic strigolactone analogue that consists of two enantiomers
(Supplementary Fig. 1a), of which GR24*ent-5DS* is bioactive via AtKAI2[40, 49]. Catalytically inactive
D14 and KAI2 variants do not respond to GR24 in DSF, suggesting that the shift in thermal
response results from ligand hydrolysis and a corresponding conformational change in the
receptor[29, 41, 46]. In DSF assays, AtKAI2 shows a specific response to >100 μM GR24*ent-5DS* but,
for unclear reasons, no response to karrikins[41]. Likewise, we found that both BtKAI2a and
BtKAI2b were also unresponsive to karrikins in DSF (Supplementary Fig. 9a). Therefore, we
used DSF assays with GR24*ent-5DS* as a surrogate substrate as a means to analyse differences in
ligand interactions between BtKAI2a and BtKAI2b. We used N-terminal 6xHIS-SUMO as a
solubility tag to aid expression in *E. coli* and enhance stability. We found that the presence of
the tag did not compromise the thermal destabilisation response of BtKAI2b to GR24*ent-5DS*
(Supplementary Fig. 9b), and therefore we used intact SUMO fusion proteins for all other
experiments.

We found that AtKAI2 and BtKAI2a showed similar responses to >100 μM GR24*ent-5DS*, but the
response of BtKAI2b was clear at >25 μM (Supplementary Fig. 9c). We then used a lower
range of GR24*ent-5DS* concentrations (0–50 μM) to determine the threshold for response, which
we defined as a statistically significant reduction in the maximal rate of change in
fluorescence at the melting temperature of the protein (T$_m$). Although BtKAI2a showed only a
weak and non-significant response at 40 and 50 μM GR24*ent-5DS*, BtKAI2b responded
significantly at 10 μM and above (Fig. 4a). These results suggest that BtKAI2b is more
sensitive than BtKAI2a to GR24*ent-5DS*, and that BtKAI2a is most like AtKAI2 in this respect.

BtKAI2a and BtKAI2b differ in primary amino acid sequence at just 14 positions
(Supplementary Fig. 3). We postulated that differences in ligand specificity might be
determined by amino acids in the vicinity of the ligand binding pocket. Protein structural
homology models revealed only two residues that differ in this region: V98 and V191 in

BtKAI2a, and L98 and L191 in BtKAI2b; the corresponding residues in AtKAI2 and AtD14 are valines (Supplementary Fig. 8). Residue 98 is immediately adjacent to the catalytic serine at the base of the pocket. Residue 191 is located internally on αT4 of the lid domain which, in AtD14, is associated with a major rearrangement of protein structure upon ligand binding that reduces the size of the pocket[31]. Homology modelling suggests only subtle differences in the size and shape of the primary ligand binding pocket of BtKAI2a and BtKAI2b (Supplementary Fig. 8). To determine if these residues are pertinent to ligand specificity, we replaced the two valine residues of BtKAI2a with leucine residues, generating the variant BtKAI2a$^{L98;L191}$, and *vice-versa* for BtKAI2b, generating the variant BtKAI2b$^{V98;V191}$ (Supplementary Fig. 8). In DSF assays, we found that exchanging the two residues was sufficient to switch the original responses, such that BtKAI2a$^{L98;L191}$ responded sensitively to GR24$^{ent\text{-}5DS}$, but BtKAI2b$^{V98;V191}$ did not (Fig. 4b). We also assessed the function of the stable BtKAI2c^{R98V} and BtKAI2c^{QUINT} variants by DSF against GR24$^{ent\text{-}5DS}$ ligand, and found that they were only weakly destabilised by this ligand, when compared to the highly sensitive BtKAI2b (Supplemental Fig. 6e). Overall, these results indicate that BtKAI2a and BtKAI2b have distinct response profiles, albeit to a synthetic ligand, and that residues 98 and 191 contribute to this difference.

We reasoned that the most robust and informative means to assess the effect of residues 98 and 191 upon the response to karrikins was to test their function directly *in planta*. We expressed BtKAI2a$^{L98;L191}$ and BtKAI2b$^{V98;V191}$ as GFP fusion proteins in the Arabidopsis *kai2-2* null mutant background driven by the Arabidopsis *KAI2* promoter (*KAI2pro:GFP-BtKAI2a$^{L98;L191}$* and *KAI2pro:GFP-BtKAI2b$^{V98;V191}$*), and selected two independent homozygous transgenic lines for each, on the basis of protein expression level and segregation ratio (Supplementary Fig. 10). Using two different assays, we found that substitutions between BtKAI2a and BtKAI2b at positions 98 and 191 also reversed karrikin preference in Arabidopsis seedlings both in terms of hypocotyl elongation and *DLK2* transcript levels (Fig. 4c, d; Supplementary Fig. 11). Most prominently, the clear preference of BtKAI2b for KAR$_1$ was unambiguously reversed to a preference for KAR$_2$ in *BtKAI2b$^{V98;V191}$* transgenics, effectively recapitulating the response of the native BtKAI2a protein. We also examined the responses of seeds in these transgenic lines, and found that *BtKAI2a$^{L98;L191}$* seeds germinated with a clear preference for KAR$_1$, whereas *BtKAI2b$^{V98;V191}$* seeds showed no clear preference for either karrikin (Supplementary Fig. 12). Notably, this germination response pattern is the opposite to that observed for seeds expressing the native BtKAI2 proteins, with BtKAI2a

exhibiting no preference for either karrikin and BtKAI2b a preference for KAR$_1$ (Fig. 3a,

Supplementary Fig. 7). Taken together, these results demonstrate that the residues at

positions 98 and 191 largely determine differences in karrikin specificity between BtKAI2a

and BtKAI2b, at least in the context of Arabidopsis seedlings.

Taxonomic variability in KAI2 sequences suggests functional co-dependency between residues

96 and 189

We wished to gain further insight into the functional importance of diversity at positions 96

and 189 of KAI2 proteins (based on AtKAI2 notation, equivalent to positions 98 and 191 in

BtKAI2a and BtKAI2b). We analysed 476 distinct KAI2 homologues from 441 angiosperm

species (spanning eudicots, monocots and magnoliids, but excluding *B. tournefortii*) using

data collected from the 1000 Plant Transcriptomes project[50]. We found that at positions 96

and 189, valine was by far the most common amino acid (93.1% and 93.5% of sequences,

respectively; Figure 5a and Supplemental Table 3). L96 was observed in 6.1% of sequences,

while L189 was even rarer (2.1%). The L96;V189 combination was observed in 14 sequences

(2.9%), whereas V96;L189 was observed in only one homologue from *Ternstroemia*

gymanthera (Pentaphylacaceae; 0.2%). Notably, this species also expressed a second KAI2

homologue with the canonical V96;V189 combination. If the frequencies of valine and leucine

at positions 96 and 189 were independent, the L96;L189 combination would be expected to

occur less than once among 476 sequences (0.13%). However, the coincidence of L96 and

L189 was observed in nine sequences (1.9%), representing a significant 15-fold enrichment

for the L96;L189 pair over that expected by chance alone (χ^2 = 115.6; P<0.0001).

Among the nine newly-identified KAI2$^{L96;L189}$ sequences, eight were restricted to eudicots,

distributed evenly between both the superrosids (Brassicales, Fabales and Saxifragales; four

species) and superasterids (Caryophyllales; four species). The ninth sequence, co-expressed

with another transcript encoding KAI2$^{V96;V189}$, was identified from the basal dicot *Hakea*

drupacea (syn. *Hakea suaveolens*; Proteaceae). Strikingly, this species is native to

southwestern Australia, but is recognised as an invasive weed in South Africa that relies upon

fire to liberate seeds from the canopy[51]. The L96;L189 combination therefore seems to have

arisen independently on multiple occasions. Furthermore, within the order Saxifragales,

Daphniphyllum macropodum (KAI2$^{L96;L189}$) belongs to a clade of species that all express

KAI2$^{L96;V189}$ homologues (Figure 5b). Although this clade is only represented by five

sequences and four species, this distribution pattern suggests that a V96L mutation occurred

first in the common ancestor of the Hamamelidaceae, Cercidiphyllaceae and Daphyniphyllaceae, followed by V189L in the lineage leading to *D. macropodum*. Overall, this analysis suggests that positions 96 and 189, while not in direct contact with each other in the protein tertiary structure (Supplemental Fig. 8), likely exhibit functional interdependence. Valine is most probably the ancestral state at both positions, but leucine at position 96 is also tolerated. While rare, leucine at position 189 is generally found in combination with leucine at position 96, and tentatively, a V96L mutation might potentiate a subsequent V189L mutation.

The above analysis suggested that a V96L substitution might be sufficient to confer altered substrate specificity upon KAI2 proteins. To test this hypothesis, we generated transgenic Arabidopsis seedlings expressing GFP-BtKAI2b proteins with just a single amino acid substitution from leucine to valine at either position 98 or 191 (i.e. *KAI2pro:GFP-BtKAI2b^{V98}* or *KAI2pro:GFP-BtKAI2b^{V191}*), and examined their response to KAR_1 and KAR_2 in hypocotyl elongation assays. We found that the L98V substitution was sufficient to invert the karrikin preference of GFP-BtKAI2b in three independent homozygous transgenic lines (Figure 5c; Supplemental Figure 13). Although this result does not rule out an additional contribution towards ligand specificity from L191, it is consistent with the higher frequency of L96 compared to L189 observed among the angiosperms, and suggests that substitution at position 96 may have been selected relatively frequently in the evolution of KAI2 function.

DISCUSSION

The plant α/β-hydrolases KAI2 and D14 are characterised by responsiveness towards butenolide compounds, including endogenous strigolactones and strigolactone-related compounds, abiotic karrikins derived from burnt vegetation, and synthetic strigolactone analogues with a wide array of functional groups. Direct evidence for a receptor-ligand relationship between karrikins and KAI2 homologues stems from crystallography and *in vitro* binding assays. Two crystal structures of KAR-responsive KAI2 proteins from Arabidopsis and the parasitic plant *Striga hermonthica* reveal largely similar overall protein structure, but surprisingly are non-congruent with respect to KAR_1 binding position and orientation[52, 53]. The affinity of AtKAI2 for KAR_1 is imprecisely defined, with estimates of dissociation coefficients (Kd) ranging from 4.6 μM[54], to 26 μM[55], to 148 μM[56] using isothermal calorimetry, and 9 μM using fluorescence microdialysis[52]. While variability in affinity estimates can be explained in part by different experimental conditions and techniques, it should also be

considered that, depending on the homologue under examination, KAR_1 may not be the
optimal ligand. Furthermore, binding of a hydrophobic compound to a hydrophobic pocket
does not necessarily equate to ligand recognition or receptor activation. Our data, which are
derived from clear and distinct biological responses, provide strong evidence that KAI2 is
sufficient to determine ligand specificity, which in turn strengthens the case that KAI2 is the
receptor by which karrikins are perceived.

The predominance of valine and leucine at position 96 (98 in BtKAI2 proteins), observed in
99.2% of our sample of 476 KAI2 sequences, suggests that this site is critical to protein
function, which makes sense given its immediate proximity to the catalytic serine. This
position not only contributes to ligand preference, but also to protein stability, given the
negative effects of R98 in BtKAI2c. Compared with valine and leucine, arginine is relatively
bulky with a charged side chain; indeed, in the few KAI2 sequences in which position 96 is
neither leucine nor valine, the residue is nevertheless small and hydrophobic (either
methionine or alanine; Supplementary Table 3). Interestingly, substitutions in AtKAI2 at or
near the catalytic site, such as S95F, S119F and G245D, are not tolerated and render the
protein unstable in plants[43]. The likely signalling mechanism for KAI2 – by analogy of that for
D14 – involves ligand binding and/or hydrolysis, which triggers conformational changes in
the lid domain and formation of a protein complex with MAX2 and SMAX1/SMXL2 protein
partners[30, 31, 57]. As a result of signalling, KAI2 itself is degraded[40, 43]. It is possible that KAI2 is
an inherently metastable protein that is prone to conformational change and subsequent
degradation by necessity; perhaps this explains why it is so susceptible to instability through
mutation.

Ligand specificity is a key contributor to the functional distinction between karrikin and
strigolactone receptors, and elucidating the molecular mechanisms behind ligand specificity is
a significant research challenge. In certain root-parasitic weeds in the Orobanchaceae,
substitutions of bulky residues found in AtKAI2 with smaller hydrophobic amino acids that
increase the ligand-binding pocket size have likely improved the affinity for host-derived
strigolactone ligands, as opposed to the smaller-sized karrikin-type ligands[58-60]. Moreover, lid-
loop residues that affect the rigidity and size of the ligand entry tunnel determine the ligand
selectivity between KAR_1 and *ent*-5-deoxystrigol (a strigolactone analogue with non-natural
stereochemistry) among eleven KAI2 homologues in *Physcomitrella patens*[61]. Karrikin and
strigolactone compounds also show chemical diversity within themselves, yet ligand
discrimination among KAI2 and D14 receptors is not well characterised. Our data

demonstrate that, in the case of BtKAI2 proteins, the two KAI2 homologues respond
differently to subtly different KAR analogues, and that subtle changes in pocket residues likely
account for preferences between these ligands. Position 96 appears to play a primary role in
determining ligand specificity, while position 189 may improve ligand sensitivity, protein
stability or protein conformational dynamics following a prior mutation at position 96.
Although the identified residues L98 and L191 are not predicted to change dramatically the
pocket size of BtKAI2b in comparison to BtKAI2a, these residues would make the BtKAI2b
pocket more hydrophobic, which is consistent with KAR_1 being more hydrophobic than
KAR_2[4]. Therefore, ligand specificity between highly similar chemical analogues can be
achieved through fine-tuning of pocket hydrophobicity. Similarly subtle changes have been
reported for the rice gibberellin receptor GID1: changing Ile133 to Leu or Val increases the
affinity for GA_{34} relative to the less polar GA_4, which lacks just one hydroxyl group[62].

It is possible that what we learn about ligand specificity in KAI2 proteins may also be
informative about strigolactone perception by D14. Different D14 proteins may show ligand
specificity towards diverse natural strigolactones, resulting in part from multiplicity of
biosynthetic enzymes, such as the cytochrome P450 enzymes in the MAX1 family[45, 63, 64] and
supplementary enzymes such as LATERAL BRANCHING OXIDOREDUCTASE[65]. Although the
functional reasons underlying such strigolactone diversity are still unclear, it is possible that
variation among D14 homologues yields varying affinities for different strigolactone
analogues[66]. As an increasingly refined picture emerges of the features that determine ligand
specificity for KAI2 and D14 receptors, we envision the rational design of synthetic receptor
proteins with desirable ligand specificity in the future.

Gene duplication is a common feature in plant evolutionary histories as an initial step towards
neofunctionalisation[67]. In obligate parasitic weeds, duplication and diversification of *KAI2*
genes (also referred to as *HTL* genes) have shifted ligand specificity towards strigolactones[58,
59]. Karrikins, as abiotic molecules with limited natural occurrence, are unlikely to be the
natural ligand for most KAI2 proteins, which are found throughout land plants. Instead, the
evolutionary maintenance of a highly conserved receptor probably reflects the core KAI2
function of perceiving an endogenous ligand ("KL") that regulates plant development[34-36]. This
core function is presumably retained after gene duplication events through original, non-
divergent copies that are under purifying selection[58, 59]. KAI2 diversity may also reflect
differing affinity for KL variants in different species and at different life stages. Our results are
consistent with a scenario in which both BtKAI2a and BtKAI2b have retained the core KAI2

function of perceiving KL to support plant development, because both copies complement the Arabidopsis *kai2* phenotype, especially at the seedling and later stages. However, BtKAI2b has also acquired mutations that alter its ligand specificity, which in turn enhance sensitivity to the archetypal karrikin released first discovered in smoke. The *BtKAI2b* gene is also more highly expressed than *BtKAI2a* in seeds and seedlings, consistent with an adaptation to post-fire seedling establishment. This diversity among KAI2 proteins may provide *B. tournefortii* with a selective advantage in fire-prone environments, contributing to its invasive nature.

We have shown evidence for the independent evolution of valine-to-leucine substitutions in KAI2 homologues throughout the dicots, albeit at low frequency, suggesting that these changes may have adaptive significance in various ecological contexts. It will be of interest to assess the function of such homologues in other species, to validate and extend the specific conclusions we have made here. Recent findings indicate that KAI2 is an integrator that also modulates germination in response to other abiotic environmental signals, including temperature and salinity[68]. As germination is a critical life stage for seed plants, strategies for environmental conservation, restoration and weed control will benefit from specific knowledge of KAI2 sequence diversity and expression profiles.

METHODS

Chemical synthesis

Karrikins (KAR$_1$ and KAR$_2$), ^{13}[C]$_5$-labelled karrikins and GR24 enantiomers (GR24^{5DS} and GR24$^{ent-5DS}$) were prepared as previously described[69, 70].

Plant material

Arabidopsis *kai2-2* (L*er*) and *Atd14-1* (Col-0) mutants were previously described[20]. Arabidopsis plants were grown on a 6:1:1 mixture of peat-based compost (Seedling Substrate Plus; Bord Na Mona, Newbridge, Ireland), vermiculite and perlite, respectively. Light was provided by wide-spectrum white LED panels emitting 120-150 µmol photons.m^{-2}.s^{-1} with a 16 h light/8 h dark photoperiod, a 22 °C light/16 °C dark temperature cycle, and constant 60% relative humidity. The seeds of *Brassica tournefortii* used in this work were collected in November and December 2009 from two sites in Western Australia (Kings Park in Perth, and Merridin)[71]. *Brassica tournefortii* seeds were dried to 15% relative humidity for one month prior to storage in air-tight bags at –20 °C.

Seed germination assays

Seed germination assays using Arabidopsis were performed on Phytagel as described previously[49]. *Brassica tournefortii* seeds were sowed in triplicates (35-70 seeds each) on glass microfibre filter paper (Grade 393; Filtech, NSW Australia) held in 9-cm petri dishes and supplemented with mock or karrikin treatments (3 mL of aqueous treatment solution per petri dish). Treatments were prepared by diluting acetone (mock) or karrikin stocks (dissolved in acetone) 1:1000 with ultrapure water. Because germination of *B. tournefortii* is inhibited by light[9], the seeds were imbibed in the dark at 22 °C. Numbers of germinated seeds were counted each day until the germination percentages remained unchanged.

Hypocotyl elongation assays

Arabidopsis hypocotyl elongation assays were performed under red light as described previously[20]. For *B. tournefortii*, assays were performed with the following modifications to the Arabidopsis protocol: *B. tournefortii* seeds were sowed in triplicate on glass microfibre filter paper (Filtech) held in petri-dishes supplemented with mock or karrikin treatments. The seeds were imbibed in the dark for 22 h at 24 °C before exposing to continuous red light (5 μmol photons m^{-2} s^{-1}) for 4 days.

Transcript analysis

Brassica tournefortii seeds were imbibed on glass fibre filters in the dark at 22 °C and treated with karrikins using the same procedure as described under seed germination assays, above. For transcript quantification in *B. tournefortii* seeds, samples were taken at the indicated time points, briefly blot-dried on paper towel, and frozen in liquid nitrogen. For *B. tournefortii* seedlings, two methods were used. For data shown in Fig. 1e, seeds were sowed and seedlings grown in triplicate under identical conditions to those described for hypocotyl elongation assays. For data shown in Fig. 2b, seeds were first imbibed for 24 h in the dark, and then transferred to the growth room ($\sim$120-150 μmol photons m^{-2} s^{-2} white light, 16 h light/8 h dark, 22 °C light/16 °C dark) for three days. A sample of approximately 20 seedlings was then transferred to a 250 mL conical flask containing 50 mL sterile ultrapure water supplemented with 1 μM KAR$_1$, or an equivalent volume of acetone (0.1% v/v), with three replicate samples per treatment. The flasks were then shaken for 24 h before seedlings were harvested.

Arabidopsis seedlings were grown on 0.5× MS agar under growth room conditions (wide-

spectrum white LED panels emitting 120-150 μmol photons.m^{-2}.s^{-1} with a 16 h light/8 h dark

photoperiod and a 22 °C light/16 °C dark temperature cycle) for seven days. On the seventh

day, seedlings were transferred to 3 mL liquid 0.5× MS medium in 12-well culture plates

(CORNING Costar 3513) and shaken at 70 rpm for a further 22 h under the same growth room

conditions. The medium was then removed by pipette and replaced with fresh medium

containing relevant compounds or an equivalent volume (0.1% v/v) of acetone. After a

further incubation period with shaking (8 h), the seedlings were harvested, blotted dry, and

frozen in liquid nitrogen.

RNA extraction, DNase treatment, cDNA synthesis and quantitative PCR were conducted as

previously described[72]. All oligonucleotides are listed in the Supplementary Table 4.

Cloning and mutagenesis of *Brassica tournefortii KAI2* and *D14* homologues

Full-length *BtKAI2* coding sequences (and unique 3′UTR sequences) were amplified from *B.*

tournefortii cDNA with Gateway-compatible *attB* sites using the universal forward primer

BtKAI2_universal_F and homologue-specific reverse primers RACE_R (BtKAI2a), Contig1_R

(BtKAI2b) and Contig5_R (BtKAI2c), before cloning into pDONR207 (Life Technologies). The

pDONR207 clones were confirmed by Sanger sequencing and recombined with pKAI2pro-

GFP-GW[41] to generate the binary plant expression plasmids pKAI2pro-GFP-BtKAI2a,

pKAI2pro-GFP-BtKAI2b and pKAI2pro-GFP-BtKAI2c. For transient overexpression in tobacco

(Supplementary Figure 5d), BtKAI2 coding sequences were transferred via Gateway-mediated

recombination into pSKI106, which drives very high expression of coding sequences placed

immediately downstream of a full-length cDNA clone of the tobacco mosaic virus U1 strain, all

under control of the CaMV *35S* promoter in a standard T-DNA binary vector[73]. pSKI106 also

encodes an N-terminal 3× *c*-myc tag. For transient expression in tobacco using standard

binary vectors (Supplementary Figure 6b), the *BtKAI2* coding sequences were transferred into

pMDC43, which is identical to pKAI2pro-GFP-GW except that the *AtKAI2* promoter & 5′UTR

sequences are replaced with a double CaMV *35S* promoter[74]. The *BtD14a* coding sequence

was amplified from cDNA using oligonucleotides BtD14_F and BtD14_R, cloned into

pDONR207 as above, and transferred into pD14pro-GW[41].

The full-length *BtKAI2* coding sequences (excluding 3′UTRs) were amplified from the

pDONR207 clones and reconstituted with the pE-SUMO vector by Gibson Assembly to

generate the heterologous expression plasmids pE-SUMO-BtKAI2a, -BtKAI2b and -BtKAI2c.

Site-directed mutagenesis generated pE-SUMO-BtKAI2a$^{L98;L191}$, pE-SUMO-BtKAI2b$^{V98;V191}$, and

the BtKAI2c variants. For expression of mutated versions of BtKAI2a, BtKAI2b, BtKAI2c and

AtKAI2 in tobacco and Arabidopsis, site-directed mutagenesis was performed on pDONR207

clones prior to recombination with pKAI2pro-GFP-GW or pMDC43. In the case of the double

substitutions in BtKAI2a and BtKAI2b, both targeted residues were mutated simultaneously

in one PCR product, while the remainder of the plasmid was amplified in a second PCR

product. These double mutated plasmids were reconstituted by Gibson assembly. For the

single mutants, the entire plasmid was amplified using a single pair of mutagenesis primers,

and the PCR product was self-ligated. The BtKAI2c^{QUINT} variant was generated by two

successive rounds of mutagenesis. Coding regions were confirmed by Sanger sequencing.

Plant transformation

Homozygous *kai2-2* plants were transformed by floral dip. Primary transgenic seedlings were

selected on sterile 0.5× MS medium supplemented with 20 μg/mL hygromycin B. T_2 lines

exhibiting a 3:1 ratio of hygromycin resistant-to-sensitive seedlings were propagated further

to identify homozygous lines in the T_3 generation. Experiments were performed from the T_3

generation onwards.

For transient expression in tobacco, *Agrobacterium* (GV3101) carrying pSKI106 or pMDC43

variants was grown in LB medium (25 mL) supplemented with antibiotics and 20 μM

acetosyringone until OD$_{600}$ reached 1.0. The bacteria were then harvested by centrifugation

(15 min, 5000 × *g*) and resuspended in 10 mM MgCl$_2$, 10 mM MES (pH 5.6) and 100 μM

acetosyringone. The optical density was adjusted to 0.4, and the suspension was left standing

at 22 °C overnight (approximately 14 h). Leaves of three-week-old *Nicotiana benthamiana*

were then infiltrated with a 5-mL syringe, through the abaxial leaf surface. After four days, the

leaves were collected and frozen in liquid nitrogen.

Karrikin uptake measurements

Fifteen samples of seeds were sowed for each karrikin treatment (five time points, each in

triplicate). In each sample, approximately 40 mg of *Brassica tournefortii* seeds were imbibed

in 3 mL ultrapure water for 24 h in 5-mL tubes. After centrifugation (2 min at 3220 × *g*),

excess water was removed by pipette and the volume of residual water (mostly absorbed into

the seeds) was calculated by weighing the seeds before and after imbibition. Fresh ultrapure

water was added to the fully-imbibed seeds to reach a total volume of 980 μL. Then 20 μL of

100 µM KAR$_1$ or KAR$_2$ was added to a final concentration of 2 µM. The seeds were imbibed at

22 °C in darkness. At the indicated time point (0, 2, 4, 8 or 24 h post-treatment), 500 µL of the

imbibition solution was removed and combined with 100 ng of either ^{13}[C]$_5$-KAR$_1$ or ^{13}[C]$_5$-

KAR$_2$ (100 µL at 1 µg/mL) as an internal standard for quantification purposes. The sample

was then extracted once with ethyl acetate (500 µL), and 1 µL of this organic layer was

analysed using GC-MS in selective ion monitoring (SIM) mode as previously described[75].

The amount of KAR$_1$ in each sample was calculated by the formula $\frac{A(Ion150)}{A(Ion155)} \times 100$ ng and

converted to moles, where *A(Ion150)* indicates the peak area of the ion 150 (KAR$_1$ to be

measured), *A(Ion151)* indicates the peak area of the ion 151 (^{13}[C]$_5$-KAR$_1$), and 100 ng is the

amount of ^{13}[C]$_5$-KAR$_1$ spiked in before the ethyl acetate extraction. Similarly, the amount of

KAR$_2$ in each sample was calculated by the formula $\frac{A(Ion136)}{A(Ion141)} \times 100$ ng and converted to moles.

The uptake percentage adjusted to 40 mg of *B. tournefortii* seeds was calculated by the

formula: $(1 - \frac{N(x)}{N(0)}) \times \frac{40\ mg}{m\ (seed)}$, where N(0) indicates moles of karrikins at time 0, *N(x)* indicates

moles of karrikins at time point *x*, and *m*(*seed*) indicates the dry weight (mg) of seeds tested in

each replicate. For Arabidopsis seeds, the procedure was scaled down for a smaller mass of

seeds (20 mg).

Transcriptome assembly and analysis

Twenty milligrams of dry *B. tournefortii* seeds were imbibed in water for 24 h and incubated

at 22 °C in the dark. The seeds were collected by centrifugation, blotted dry, and frozen in

liquid nitrogen. A separate sample of seeds was sown on glass filter paper, imbibed for 24 h as

above, and then incubated for 96 h under continuous red light (20 µmol m^{-2} s^{-1}) with a 22 °C

(16 h)/16 °C (8h) temperature cycle. A single sample of seedlings (50 mg fresh weight) was

harvested and frozen in liquid nitrogen. Total RNA was extracted from both seed and seedling

samples using the Spectrum Plant RNA kit (Sigma-Aldrich), including an on-column DNase

step. PolyA$^+$ mRNA was purified using oligo(dT) magnetic beads (Illumina), and cDNA

libraries for sequencing were generated as described[76]. Sequencing was performed on the

Illumina HiSeq 2000 platform at the Beijing Genomic Institute, Shenzhen, China. Raw reads

were filtered to remove adapters and low-quality reads (those with >5% unknown

nucleotides, or those in which greater than 20% of base calls had quality scores ≤10). After

filtering, both libraries generated reads with >99.9% of nucleotides attaining Q20 quality

score. Transcriptome *de novo* assembly was performed with Trinity[77]. For each library,

contigs were assembled into Unigenes; Unigenes from both libraries were then combined,

yielding a total of 45,553 predicted coding region sequences with a mean length of 1011 nt.

The combined Unigenes were then interrogated for homology to *AtKAI2*, *AtD14*, *AtDLK2*,

AtSTH7 and *AtCACS* using BLASTn searches.

Phylogenetic analysis and protein sequence analysis

KAI2 and *D14* homologues in *Brassica* species were identified from BLAST searches using

Arabidopsis coding sequences as a query. Additional sequences were sampled from an

existing phylogenetic analysis[33]. Multiple sequence alignments were performed using MAFFT

plugin implemented in Geneious R10 (Biomatters). The alignment was trimmed slightly at the

5′ end to remove non-aligned regions of monocot D14 sequences. Maximum likelihood

phylogenies were generated using PHYML (GTR +G +I substitution model, combination of NNI

and SPR search, and 100 bootstraps). The choice of substitution model was guided by Smart

Model Selection in PhyML[78] (http://www.atgc-montpellier.fr/sms). A list of all sequences, and

their sources, is provided in Supplementary Table 1.

For analysis of BtKAI2c using PROVEAN, all ten positions that were unique to BtKAI2c when

compared to AtKAI2 and other *Brassica* sequences (Supplementary Fig. 3) were manually

edited to the consensus identity at each position. This was the "reverted" sequence against

which each of the native BtKAI2c amino acid substitutions were tested using the PROVEAN

server (http://provean.jcvi.org) using the default cut-off score of -2.5. The PROVEAN output is

presented in Supplementary Table 2.

KAI2 homologues from The 1000 Plant Transcriptomes Project[50] were identified by TBLASTN

searches using *Arabidopsis thaliana* KAI2 protein sequence as a query against "onekp

database v5", which comprises 1328 individual samples (https://db.cngbdb.org). The search

was configured to return a maximum of 500 target sequences with Expect values <0.01, using

a word size of 6 and the BLOSUM62 scoring matrix. Nucleotide sequences were collated, and

open reading frames (ORFs) >700 nucleotides were predicted in Geneious software. A single

ORF was selected for each sequence; those with multiple ORFs were screened manually for

the one with homology to AtKAI2. The ORFs were translated and aligned using the MAFFT

multiple alignment tool. A few misaligned or mis-translated sequences were removed from

the alignment manually, resulting in a dataset of 476 unique sequences. Extracted parts of this

alignment centred on positions 96 and 189 were used to generate WebLogos

(http://weblogo.threeplusone.com[79]). A list of identified KAI2 homologues and their

positional analysis are presented in Supplementary Table 3.

Protein homology modelling

KAI2 structures were modelled using the SWISS-MODEL server

(https://swissmodel.expasy.org) using the alignment mode[80] and the *Arabidopsis thaliana*

KAI2 structure 3w06 as a template[54]. Figures of protein structure and homology models were

generated using PyMOL v1.3 (Schrödinger LLC). Cavity surfaces were visualised using the

"cavities & pockets (culled)" setting in PyMOL and a cavity detection cut-off value of four

solvent radii. Cavity volumes were calculated using the CASTp server v3.0[81]

(http://sts.bioe.uic.edu/castp) with a probe radius of 1.4Å. Values indicate Connolly's solvent-

excluded volumes. Cavities were inspected visually using Chimera v1.12

(https://www.cgl.ucsf.edu/chimera/). For both BtKAI2a and BtKAI2b models, CASTp

erroneously included surface residues near the primary pocket entrance in the calculation of

the pocket volumes. This issue was resolved by the artificial placement of a free alanine

residue adjacent to the cavity entrance, as described previously[82].

Protein expression and purification

599 BtKAI2 proteins were generated as N-terminal 6×HIS-SUMO fusion proteins. All proteins were

600 expressed in BL21 Rosetta DE3 pLysS cells (Novagen) and purified using IMAC as described in

601 detail previously[41].

Differential scanning fluorimetry

603 DSF was performed in 384-well format and thermal shifts were quantified as described

604 previously[41]. Reactions (10 μL) contained 20 μM protein, 20 mM HEPES pH 7.5, 150 mM NaCl,

605 1.25% (v/v) glycerol, 5× SYPRO Tangerine dye (Molecular Probes) and varying

606 concentrations of ligand that resulted in a final concentration of 5% (v/v) acetone.

Statistical analysis

Data were analysed using one- or two-way ANOVA ($\alpha = 0.05$, with Tukey's multiple

comparisons test). For Figure 5a, in which data from three experimental replicates were

combined, data were analysed using a mixed effects model with experimental replicate as a

random effect, and genotype and treatment as fixed effects. Prior to ANOVA, germination data

were arcsine-transformed, and gene expression data were log-transformed. Tests were

implemented in GraphPad Prism version 7.0 or 8.0 (GraphPad Software, graphpad.com).

DATA AVAILABILITY

All relevant data and materials are available from the authors upon reasonable request.

Sequence data are available at NCBI Genbank under the following accessions: *BtKAI2a*,

MG783328; *BtKAI2b*, MG783329; *BtKAI2c*, MG783330; *BtD14a*, MG783331; *BtD14b*,

MG783332; *BtDLK2*, MG783333; *BtSTH7*, MK756121; *BtCACS*, MK756122. Raw RNA sequence

data from *Brassica tournefortii* seed and seedlings are available in the NCBI SRA database

under accession SRP128835. The original data underlying the following figures are provided

as a Source Data file: Figs 1a,b,d,e; 2b,e; 3a–d; 4a–d; 5a–c; Supplementary Figs 1b–f; 2; 3; 4c;

5b–d; 6b,c,e; 7; 8a–e; 9a–c; 10a; 11; 12; 13. A reporting summary for this Article is available as

a Supplementary Information File.

REFERENCES

1.   Early R, *et al.* Global threats from invasive alien species in the twenty-first century and
626   national response capacities. *Nat Commun* **7**, 12485-12485 (2016).

627
628   2.   Flematti GR, *et al.* Karrikin and cyanohydrin smoke signals provide clues to new
629   endogenous plant signaling compounds. *Molecular Plant* **6**, 29-37 (2013).

630
631   3.   Keeley JE, Pausas JG. Evolution of 'smoke' induced seed germination in pyroendemic
632   plants. *South African Journal of Botany* **115**, 251-255 (2018).

633
634   4.   Flematti GR, Ghisalberti EL, Dixon KW, Trengove RD. Identification of Alkyl Substituted
635   2 H-Furo[2,3- c]pyran-2-ones as Germination Stimulants Present in Smoke. *Journal of*
636   *Agricultural and Food Chemistry* **57**, 9475-9480 (2009).

637
638   5.   Hrdlička J, *et al.* Quantification of karrikins in smoke water using ultra-high
639   performance liquid chromatography-tandem mass spectrometry. *Plant Methods* **15**, 81
640   (2019).

641
642   6.   Long RL, Stevens JC, Griffiths EM, Adamek M, Powles SB, Merritt DJ. Detecting
643   karrikinolide responses in seeds of the Poaceae. *Australian Journal of Botany* **59**, 610-
644   620 (2011).

7.   Milberg P, Lamont BB. Fire enhances weed invasion of roadside vegetation in southwestern Australia. *Biological Conservation* **73**, 45-49 (1995).

8.   Flematti GR, Goddard-Borger ED, Merritt DJ, Ghisalberti EL, Dixon KW, Trengove RD. Preparation of 2H-furo[2,3-c]pyran-2-one derivatives and evaluation of their germination-promoting activity. *Journal of Agricultural and Food Chemistry* **55**, 2189-2194 (2007).

9.   Long RL, *et al.* Seeds of Brassicaceae weeds have an inherent or inducible response to the germination stimulant karrikinolide. *Annals of Botany* **108**, 933-944 (2011).

10. Stevens J, Merritt D, Flematti G, Ghisalberti E, Dixon K. Seed germination of agricultural weeds is promoted by the butenolide 3-methyl-2H-furo[2,3-c]pyran-2-one under laboratory and field conditions. *Plant Soil* **298**, 113-124 (2007).

11. Nelson DC, Flematti GR, Riseborough J-A, Ghisalberti EL, Dixon KW, Smith SM. Karrikins enhance light responses during germination and seedling development in Arabidopsis thaliana. *Proceedings of the National Academy of Sciences* **107**, 7095-7100 (2010).

12. Kulkarni MG, Sparg SG, Light ME, Staden JV. Stimulation of Rice (Oryza sativa L.) Seedling Vigour by Smoke‐water and Butenolide. *J Agron Crop Sci* **192**, 395-398 (2006).

13. Staden Jv, Sparg SG, Kulkarni MG, Light ME. Post-germination effects of the smoke-derived compound 3-methyl-2H-furo[2,3-c]pyran-2-one, and its potential as a preconditioning agent. *Field Crop Res* **98**, 98-105 (2006).

14. Bangle DN, Walker LR, Powell EA. Seed germination of the invasive plant Brassica tournefortii (Sahara mustard) in the Mojave Desert. *Western North American Naturalist* **68**, 334-342 (2008).

15. Brooks ML, *et al.* Effects of Invasive Alien Plants on Fire Regimes. *BioScience* **54**, 677 (2004).

16. Sanders R, Minnich R. Brassica tournefortii. In: *Invasive Plants of California's Wildlands* (eds Bossard CC, Randall JM, Hoshovsky MC). University of California Press (2000).

17. Schiermeier Q. Pall hangs over desert's future as alien weeds fuel wildfires. *Nature* **435**, 724 (2005).

686
687   18.   Steers RJ. Invasive Plants, Fire Succession, and Restoration of Creosote Bush Scrub in
688   Southern California.). UNIVERSITY OF CALIFORNIA RIVERSIDE (2008).

689
690   19.   Sun X-D, Ni M. HYPOSENSITIVE TO LIGHT, an Alpha/Beta Fold Protein, Acts
691   Downstream of ELONGATED HYPOCOTYL 5 to Regulate Seedling De-Etiolation.
692   *Molecular Plant* **4**, 116-126 (2011).

693
694   20.   Waters MT*, et al.* Specialisation within the DWARF14 protein family confers distinct
695   responses to karrikins and strigolactones in Arabidopsis. *Development* **139**, 1285-1295
696   (2012).

697
698   21.   Durvasula A*, et al.* African genomes illuminate the early history and transition to
699   selfing in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences* **114**,
700   5213-5218 (2017).

701
702   22.   Nelson DC*, et al.* Karrikins discovered in smoke trigger Arabidopsis seed germination
703   by a mechanism requiring gibberellic acid synthesis and light. *Plant Physiology* **149**,
704   863-873 (2009).

705
706   23.   Flematti GR, Ghisalberti EL, Dixon KW, Trengove RD. A compound from smoke that
707   promotes seed germination. *Science* **305**, 977 (2004).

708
709   24.   Waters MT, Gutjahr C, Bennett T, Nelson DC. Strigolactone Signaling and Evolution.
710   *Annual Review of Plant Biology* **68**, 291-322 (2017).

711
712   25.   Yao R, Chen L, Xie D. Irreversible strigolactone recognition: a non-canonical
713   mechanism for hormone perception. *Current Opinion in Plant Biology* **45**, 155-161
714   (2018).

715
716   26.   Scaffidi A*, et al.* Exploring the molecular mechanism of karrikins and strigolactones.
717   *Bioorganic & Medicinal Chemistry* **22**, 3743-3746 (2012).

718
719   27.   Zwanenburg B, Pospíšil T. Structure and Activity of Strigolactones: New Plant
720   Hormones with a Rich Future. *Molecular Plant* **6**, 38-62 (2013).

721
722   28.   de Saint Germain A*, et al.* An histidine covalent receptor and butenolide complex
723   mediates strigolactone perception. *Nature Chemical Biology* **12**, 787-794 (2016).

724
725   29.   Seto Y*, et al.* Strigolactone perception and deactivation by a hydrolase receptor
726   DWARF14. *Nat Commun* **10**, 191 (2019).

30. Shabek N, Ticchiarelli F, Mao H, Hinds TR, Leyser O, Zheng N. Structural plasticity of D3–D14 ubiquitin ligase in strigolactone signalling. *Nature* **563**, 652-656 (2018).

31. Yao R*, et al.* DWARF14 is a non-canonical hormone receptor for strigolactone. *Nature* **536**, 469-473 (2016).

32. Marzec M, Brewer P. Binding or Hydrolysis? How Does the Strigolactone Receptor Work? *Trends in Plant Science* **24**, 571-574 (2019).

33. Bythell-Douglas R*, et al.* Evolution of strigolactone receptors by gradual neo-functionalization of KAI2 paralogues. *BMC Biology* **15**, 52 (2017).

34. Conn CE, Nelson DC. Evidence that KARRIKIN-INSENSITIVE2 (KAI2) Receptors may Perceive an Unknown Signal that is not Karrikin or Strigolactone. *Frontiers in Plant Science* **6**, 1219 (2015).

35. Li W*, et al.* The karrikin receptor KAI2 promotes drought resistance in Arabidopsis thaliana. *PLoS Genetics* **13**, e1007076 (2017).

36. Sun YK, Flematti GR, Smith SM, Waters MT. Reporter Gene-Facilitated Detection of Compounds in Arabidopsis Leaf Extracts that Activate the Karrikin Signaling Pathway. *Frontiers in Plant Science* **7**, 1799 (2016).

37. Liu S*, et al.* The Brassica oleracea genome reveals the asymmetrical evolution of polyploid genomes. *Nat Commun* **5**, 3930 (2014).

38. Lysak MA, Koch MA, Pecinka A, Schubert I. Chromosome triplication found across the tribe Brassiceae. *Genome Res* **15**, 516-525 (2005).

39. Wang X*, et al.* The genome of the mesopolyploid crop species Brassica rapa. *Nature Genetics* **43**, 1035-1039 (2011).

40. Waters MT, Scaffidi A, Flematti G, Smith SM. Substrate-Induced Degradation of the α/β-Fold Hydrolase KARRIKIN INSENSITIVE2 Requires a Functional Catalytic Triad but Is Independent of MAX2. *Molecular Plant* **8**, 814-817 (2015).

41. Waters MT, Scaffidi A, Moulin SLY, Sun YK, Flematti GR, Smith SM. A Selaginella moellendorffii Ortholog of KARRIKIN INSENSITIVE2 Functions in Arabidopsis Development but Cannot Mediate Responses to Karrikins or Strigolactones. *Plant Cell* **27**, 1925-1944 (2015).

768
769 42. Kagale S*, et al.* TMV-Gate vectors: Gateway compatible tobacco mosaic virus based
770   expression vectors for functional analysis of proteins. *Scientific Reports* **2**, 874 (2012).

771
772 43. Yao J*, et al.* An allelic series at the KARRIKIN INSENSITIVE 2 locus of Arabidopsis
773   thaliana decouples ligand hydrolysis and receptor degradation from downstream
774   signalling. *Plant J* **96**, 75-89 (2018).

775
776 44. Choi Y, Chan AP. PROVEAN web server: a tool to predict the functional effect of amino
777   acid substitutions and indels. *Bioinformatics* **31**, 2745-2747 (2015).

778
779 45. Abe S*, et al.* Carlactone is converted to carlactonoic acid by MAX1 in Arabidopsis and its
780   methyl ester can directly interact with AtD14 in vitro. *Proceedings of the National
781   Academy of Sciences of the United States of America* **111**, 18084-18089 (2014).

782
783 46. Hamiaux C*, et al.* DAD2 Is an α/β Hydrolase Likely to Be Involved in the Perception of
784   the Plant Branching Hormone, Strigolactone. *Current Biology* **22**, 2032-2036 (2012).

785
786 47. Hamiaux C*, et al.* Inhibition of strigolactone receptors by N-phenylanthranilic acid
787   derivatives: structural and functional insights. *Journal of Biological Chemistry* **293**,
788   6530-6543 (2018).

789
790 48. Végh A*, et al.* Comprehensive Analysis of DWARF14-LIKE2 (DLK2) Reveals Its
791   Functional Divergence from Strigolactone-Related Paralogs. *Frontiers in Plant Science*
792   **8**, 1641 (2017).

793
794 49. Scaffidi A*, et al.* Strigolactone hormones and their stereoisomers signal through two
795   related receptor proteins to induce different physiological responses in Arabidopsis.
796   *Plant Physiology* **165**, 1221-1232 (2014).

797
798 50. Leebens-Mack JH*, et al.* One thousand plant transcriptomes and the phylogenomics of
799   green plants. *Nature* **574**, 679-685 (2019).

800
801 51. Richardson DM, Wilgen BWV, Mitchell DT. Aspects of the reproductive ecology of four
802   australian Hakea species (Proteaceae) in South Africa. *Oecologia* **71**, 345-354 (1987).

803
804 52. Guo Y, Zheng Z, La Clair JJ, Chory J, Noel JP. Smoke-derived karrikin perception by the
805   α/β-hydrolase KAI2 from Arabidopsis. *Proceedings of the National Academy of Sciences
806   of the United States of America* **110**, 8284-8289 (2013).

808 53. Xu Y*, et al.* Structural basis of unique ligand specificity of KAI2-like protein from
809 parasitic weed Striga hermonthica. *Scientific Reports* **6**, 31386 (2016).

810
811 54. Kagiyama M*, et al.* Structures of D14 and D14L in the strigolactone and karrikin
812 signaling pathways. *Genes to Cells* **18**, 147-160 (2013).

813
814 55. Toh S, Holbrook-Smith D, Stokes ME, Tsuchiya Y, McCourt P. Detection of parasitic
815 plant suicide germination compounds using a high-throughput Arabidopsis HTL/KAI2
816 strigolactone perception system. *Chem Commun* **21**, 988-998 (2014).

817
818 56. Lee I*, et al.* A missense allele of KARRIKIN-INSENSITIVE2 impairs ligand-binding and
819 downstream signaling in Arabidopsis thaliana. *Journal of Experimental Botany* **69**,
820 3609-3623 (2018).

821
822 57. Yao J, Waters MT. Perception of karrikins by plants: a continuing enigma. *Journal of
823 Experimental Botany*, 10.1093/jxb/erz1548 (2019).

824
825 58. Conn CE*, et al.* Convergent evolution of strigolactone perception enabled host detection
826 in parasitic plants. *Science* **349**, 540-543 (2015).

827
828 59. Toh S*, et al.* Structure-function analysis identifies highly sensitive strigolactone
829 receptors in Striga. *Science* **350**, 203-207 (2015).

830
831 60. Xu Y*, et al.* Structural analysis of HTL and D14 proteins reveals the basis for ligand
832 selectivity in Striga. *Nat Commun* **9**, 3947 (2018).

833
834 61. Bürger M*, et al.* Structural Basis of Karrikin and Non-natural Strigolactone Perception
835 in *Physcomitrella patens*. *Cell Reports* **26**, 855-865 (2019).

836
837 62. Shimada A*, et al.* Structural basis for gibberellin recognition by its receptor GID1.
838 *Nature* **456**, 520-523 (2008).

839
840 63. Zhang Y*, et al.* Rice cytochrome P450 MAX1 homologs catalyze distinct steps in
841 strigolactone biosynthesis. *Nature Chemical Biology* **10**, 1028-1033 (2014).

842
64. Challis RJ, Hepworth J, Mouchel C, Waites R, Leyser O. A Role for MORE AXILLARY
GROWTH1 (MAX1) in Evolutionary Diversity in Strigolactone Signaling Upstream of
MAX2. *Plant Physiology* **161**, 1885-1902 (2013).

65. Brewer PB, *et al. LATERAL BRANCHING OXIDOREDUCTASE* acts in the final stages of
strigolactone biosynthesis in *Arabidopsis*. *Proceedings of the National Academy of*
Sciences **113**, 6301-6306 (2016).

850
851 66. Yoneyama K, *et al.* Which are the major players, canonical or non-canonical
852   strigolactones? *Journal of Experimental Botany* **69**, 2231-2239 (2018).

853
854 67. Panchy N, Lehti-Shiu MD, Shiu S-H. Evolution of gene duplication in plants. *Plant*
855   *Physiology* **171**, 2294-2316 (2016).

856
857 68. Wang L, Waters MT, Smith SM. Karrikin-KAI2 signalling provides Arabidopsis seeds
858   with tolerance to abiotic stress and inhibits germination under conditions
859   unfavourable for seedling establishment. *New Phytologist* **219**, 605-618 (2018).

860
861 69. Goddard-Borger ED, Ghisalberti EL, Stick RV. Synthesis of the germination stimulant 3-
862   methyl-2H-furo[2,3-c]pyran-2-one and analogous compounds from carbohydrates.
863   *European Journal of Organic Chemistry*, 3925-3934 (2007).

864
865 70. Mangnus E, Vanvliet LA, Vadenput D, Zwanenburg B. Structural modifications of strigol
866   analogs - influence of the B and C rings on the bioactivity of the germination stimulant
867   GR24. *Journal of Agricultural and Food Chemistry* **40**, 1222-1229 (1992).

868
869 71. Gorecki MJ, Long RL, Flematti GR, Stevens JC. Parental environment changes the
870   dormancy state and karrikinolide response of Brassica tournefortii seeds. *Annals of*
871   *Botany* **109**, 1369-1378 (2012).

872
873 72. Waters MT, Smith SM. KAI2- and MAX2-mediated responses to karrikins and
874   strigolactones are largely independent of HY5 in Arabidopsis seedlings. *Molecular*
875   *Plant* **6**, 63-75 (2013).

876
877 73. Kagale S, *et al.* TMV-Gate vectors: gateway compatible tobacco mosaic virus based
878   expression vectors for functional analysis of proteins. *Scientific Reports* **2**, 874 (2012).

879
880 74. Curtis MD, Grossniklaus U. A Gateway Cloning Vector Set for High-Throughput
881   Functional Analysis of Genes in Planta. *Plant Physiology* **133**, 462-469 (2003).

882
883 75. Flematti GR, Scaffidi A, Dixon KW, Smith SM, Ghisalberti EL. Production of the seed
884   germination stimulant karrikinolide from combustion of simple carbohydrates. *Journal*
885   *of Agricultural and Food Chemistry* **59**, 1195-1198 (2011).

887   76.   Liang C, Liu X, Yiu S-M, Lim BL. De novo assembly and characterization of Camelina
888   sativa transcriptome by paired-end sequencing. *BMC Genomics* **14**, 146 (2013).

889
890   77.   Grabherr MG*, et al.* Full-length transcriptome assembly from RNA-Seq data without a
891   reference genome. *Nature Biotechnology* **29**, 644-652 (2011).

892
893   78.   Lefort V, Longueville J-E, Gascuel O. SMS: Smart Model Selection in PhyML. *Molecular
894   Biology and Evolution* **34**, 2422-2424 (2017).

895
896   79.   Crooks GE, Hon G, Chandonia J-M, Brenner SE. WebLogo: A Sequence Logo Generator.
897   *Genome Res* **14**, 1188-1190 (2004).

898
899   80.   Bordoli L, Kiefer F, Arnold K, Benkert P, Battey J, Schwede T. Protein structure
900   homology modeling using SWISS-MODEL workspace. *Nature Protocols* **4**, 1-13 (2009).

901
902   81.   Dundas J, Ouyang Z, Tseng J, Binkowski A, Turpaz Y, Liang J. CASTp: computed atlas of
903   surface topography of proteins with structural and topographical mapping of
904   functionally annotated residues. *Nucleic Acids Research* **34**, W116-W118 (2006).

905
906   82.   Lopez-Obando M*, et al.* Structural modelling and transcriptional responses highlight a
907   clade of PpKAI2-LIKE genes as candidate receptors for strigolactones in Physcomitrella
908   patens. *Planta* **243**, 1441-1453 (2016).

ACKNOWLEDGEMENTS

This work was supported by funding from the Australian Research Council (DP130103646
to SMS and GRF; DP140104567 to SMS; FT150100162 to MTW, DP160102888 to GRF and
MTW). Y.K.S was recipient of a Research Training Program Scholarship from the Australian
Government. We thank Dr Rowena Long for the original provision of *B. tournefortii* seeds, Dr
Rohan Bythell-Douglas for advice on homology modelling, and Mr Yongjie Meng with
assistance with RNA extractions. We are grateful to Dr Kevin Rozwadowski (Agriculture and
Agri-Food Canada) for the provision of pSKI106.

CONTRIBUTIONS

Y.K.S., S.M.S., C.S.B., G.R.F. and M.T.W. conceived and designed the research. Y.K.S. and M.T.W. performed the majority of experiments with assistance from J.Y. (hypocotyl elongation assays and screening of transgenic lines), A.S. (chemical synthesis and seed germination assays), K.T.M. and S.F.D. (protein expression and purification). Y.K.S. and M.T.W. analysed the data. Y.K.S., S.M.S and M.T.W. wrote the manuscript.

ADDITIONAL INFORMATION

Supplementary Figure 1 Germination of *Brassica tournefortii* seed treated with karrikins

Supplementary Figure 2 Extended phylogeny of eu-KAI2 and D14 proteins in angiosperms

Supplementary Figure 3 Alignment of Brassica KAI2 sequences

Supplementary Figure 4 BtD14a is functionally homologous to AtD14

Supplementary Figure 5 The *GFP-BtKAI2c* transgene is faithfully transcribed in Arabidopsis

Supplementary Figure 6 The R98V mutation restores stability of BtKAI2c

Supplementary Figure 7 Germination profiles of transgenic Arabidopsis seeds expressing native BtKAI2 homologues

Supplementary Figure 8 BtKAI2 homology models and SDS-PAGE of SUMO-BtKAI2 fusion proteins used for DSF

Supplementary Figure 9 BtKAI2a and BtKAI2b do not respond to karrikins in DSF assays

Supplementary Figure 10 Stable transgenic expression of BtKAI2 valine–leucine double exchange proteins in Arabidopsis

Supplementary Figure 11 Three experimental replicates of hypocotyl elongation assays with BtKAI2 Arabidopsis transgenics (double exchange of residues 98 and 191)

Supplementary Figure 12 Germination profiles of transgenic Arabidopsis seed expressing BtKAI2 homologues

FIGURE LEGENDS

Figure 1. *Brassica tournefortii* is highly sensitive to KAR$_1$, the major karrikin analogue isolated from plant-derived smoke

a, Germination responses of *B. tournefortii* seed to KAR$_1$ and KAR$_2$. Data are cumulative germination after 11 days (mean ± SE; n = 3 biological replicates per treatment, ≥35 seeds per replicate). **b**, Levels of karrikin-responsive transcripts *BtDLK2* and *BtSTH7* in *B. tournefortii* seed. Seed were imbibed for 24 hours in the dark supplemented with KAR$_1$ and KAR$_2$. Transcripts were normalised to *BtCACS* reference transcripts. Data are means ± SE, n = 3 biological replicates, ≥50 seeds per replicate. **c**, **d**, Hypocotyl elongation responses of *B. tournefortii* seedlings grown for four days under continuous red light on water-saturated glass filter paper containing 0.1% acetone (mock), KAR$_1$ or KAR$_2$. Data are means ± 95% CI of n = 18 to 24 seedlings. Scale bar: 10 mm. **e**, Levels of *BtDLK2* transcripts in *B. tournefortii* seedlings grown under the same conditions as for **d**. Data are means ± SE, n = 3 biological replicates, ≥20 seedlings per replicate. In all panels, asterisks denote significance levels (ANOVA) between indicated conditions: * $P<0.05$; ** $P<0.01$; *** $P<0.001$; **** $P<0.0001$. Source data are provided as a Source Data file.

Figure 2. Two differentially expressed *B. tournefortii KAI2* homologues are functional in *Arabidopsis*

a, Maximum likelihood phylogeny of KAI2 homologues in the Brassicaceae, based on nucleotide data. Node values represent bootstrap support from 100 replicates. A eu-KAI2 sequence from *Gossypium raimondii* (Malvaceae) serves as an outgroup. Tree shown is a subset of a larger phylogeny in Supplementary Fig. 2. **b**, Transcript levels of the three *BtKAI2* homologues in *B. tournefortii* seeds imbibed for 24 h (left) and four-day-old seedlings (right) treated with 0.1% acetone or with 1 μM KAR_1 for 24 h. **c, d**, Seedling and rosette phenotypes of two independent transgenic lines of Arabidopsis homozygous for *KAI2pro:GFP-BtKAI2* transgenes. Scale bars: 5 mm (**c**); 50 mm (**d**). **e**, Immunoblots of soluble proteins challenged with antibodies against KAI2 (upper panel), GFP (middle panel) or actin as a loading control (lower panel). Anti-KAI2 detects both the native AtKAI2 protein (30 kDa) and the GFP-BtKAI2 fusion proteins (58 kDa) but detects BtKAI2b relatively poorly. Non-specific bands are marked with asterisks. Protein was isolated from pools of approximately fifty 7-day-old seedlings. Source data are provided as a Source Data file.

Figure 3. Functional divergence between BtKAI2 homologues

a, Germination responses of primary dormant *Arabidopsis* seed homozygous for *KAI2pro:GFP-BtKAI2* transgenes in the *kai2-2* background. Germination values were determined 120 h after sowing. Extended germination curves are shown in Supplementary Fig. 7. Data are means ± SE, *n* = 3 independent seed batches, 75 seed per batch. **b**, Levels of *KUF1* transcripts in *KAI2pro:GFP-BtKAI2* seeds treated with 1 μM KAR_1 or KAR_2. Expression was normalised to *CACS* reference transcripts and scaled to the value for mock-treated seed within each genotype. Data are means ± SE of n = 3 biological replicates. **c**, Hypocotyl elongation responses of *KAI2pro:GFP-BtKAI2* seedlings treated with KAR_1 or KAR_2. Data are means ± SE of *n* = 3 biological replicates, 12–18 seedlings per replicate. **d**, Levels of *DLK2* transcripts in 8-day-old *KAI2pro:GFP-BtKAI2* seedlings treated with KAR_1 or KAR_2 for eight hours. Expression was normalised to *CACS* reference transcripts and scaled to the value for mock-treated seedlings within each genotype. Data are means ± SE of *n* = 3 biological replicates. Pairwise significant differences: * $P < 0.05$ ** $P < 0.01$ *** $P < 0.001$; ns, $P > 0.05$ (ANOVA). Source data are provided as a Source Data file.

Figure 4. Two residues account for ligand specificity between BtKAI2a and BtKAI2b

a, DSF curves of SUMO-BtKAI2a and SUMO-BtKAI2b fusion proteins treated with 0-50 μM GR24$^{ent-5DS}$, a KAI2-bioactive ligand. Each curve is the average of three sets of reactions, each

comprising four technical replicates. Insets plot the minimum value of –(dF/dT) at the melting

point of the protein as determined in the absence of ligand (means ± SE, n = 3). Significant

differences from untreated control: * $P < 0.05$ ** $P < 0.01$ *** $P < 0.001$ (ANOVA). **b**, DSF curves

of SUMO-BtKAI2a$^{L98;L191}$ and SUMO-BtKAI2b$^{V98;V191}$ fusion proteins treated with 0-50 μM

GR24$^{ent\text{-}5DS}$. **c**, Hypocotyl elongation responses of Arabidopsis expressing GFP-BtKAI2a$^{L98;L191}$

and GFP-BtKAI2b$^{V98;V191}$ fusion proteins and treated with KAR$_1$ or KAR$_2$. Data are a summary

of three experimental replicates performed on separate occasions, each comprising 25-40

seedlings per genotype/treatment combination. Data for each replicate are shown in

Supplementary Figure 11. Error bars are SE, n = 3 experimental replicates; each dot

corresponds to the mean value derived from each replicate. Asterisks denote significant

differences: * P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001 (linear mixed model with

experimental replicate as a random effect; specific pairwise comparisons using Tukey's HSD

correction). **d**, Levels of *DLK2* transcripts in the same transgenic lines as above treated with 1

μM KAR$_1$, 1 μM KAR$_2$, or 0.1% acetone and harvested 8 hours later. Expression was

normalised to *CACS* reference transcripts. Data are means ± SE of n = 3 pools of ~50 seedlings

treated in parallel. Asterisks denote significant differences as above (two-way ANOVA;

specific pairwise comparisons using Tukey's HSD correction). Source data are provided as a

Source Data file.

Figure 5. Diversity at positions 96 and 189 in KAI2 homologues across angiosperms

a, Sequence logos generated from 476 aligned KAI2 homologues sampled from eudicots,

monocots and magnoliids and centred on position 96 (left) and 189 (right). b, Species tree of

the Saxifragales based upon 410 single-copy nuclear genes estimated using the ASTRAL gene

tree method[50]. Branch support values are shown only where less than 1. The four-letter taxon

prefix corresponds to the 1000 Plant Transcriptomes project Sample ID. Taxa are highlighted

by colour according to identity of amino acids of KAI2 orthologues at positions 96 and 189

respectively: black, VV; blue, LV; red, LL; magenta, other. Postulated ancestral identities at

specific nodes are indicated with filled circles. c, Hypocotyl elongation responses to karrikins

in three independent transgenic lines homozygous for *GFP-BtKAI2a$^{L98;V191}$* and three lines for

GFP-BtKAI2b$^{V98;L191}$. Data shown are a summary of three experimental replicates performed

on separate occasions, each comprising ~20 seedlings per genotype/treatment combination.

Data from each replicate are presented in Supplementary Figure 13. Error bars are SE, n = 3

experimental replicates; each dot corresponds to the mean value derived from each replicate.

Asterisks denote significant differences: * P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001

(linear mixed model with experimental replicate as a random effect; specific pairwise

comparisons using Tukey's HSD correction). Source data are provided as a Source Data file.

SUPPLEMENTARY FIGURE LEGENDS

Supplementary Figure 1. Germination of *Brassica tournefortii* seed treated with karrikins

a, Structures of KAR_1, KAR_2 and two enantiomers of GR24. **b**, Germination response of the "Merridin" batch of *B. tournefortii* seed treated with KAR_1 and KAR_2 after three days. **c-d**, Germination response of the "Perth" batch to KAR_1 (**c**) and KAR_2 (**d**). Data in Figure 1a are derived from the data shown in **c**. **e-f**, Uptake of KAR_1 and KAR_2 by imbibed seed, as determined by GC-MS. All error bars are mean ± SE of n = 3 batches of 75 seed (**b-d**) or 3 samples of 40 mg (**e**) or 20 mg (**f**) seed as described in Methods. Source data are provided as a Source Data file.

Supplementary Figure 2. Extended phylogeny of eu-KAI2 and D14 proteins in angiosperms

Maximum likelihood phylogeny of KAI2 and D14 homologues in the Brassicaceae and monocots, based on nucleotide data. Node values represent bootstrap support from 100 replicates. A KAI2 sequence from *Selaginella moellendorffii* (SmKAI2a) serves as an outgroup for the eu-KAI2 clade. Source data are provided as a Source Data file.

Supplementary Figure 3. Alignment of Brassica KAI2 sequences

Full length protein coding regions of *KAI2* homologues from four *Brassica* species (*Brassica tournefortii*, *B. rapa*, *B. nigra* and *B. oleracea*) were translated from database nucleic acid sequences and aligned to *Arabidopsis thaliana* KAI2 using MAFFT[1] implemented in Geneious R10 software (Biomatters Ltd). Amino acid residues are coloured according to polarity: yellow, non-polar (G, A, V, L, I, F, W, M, P); green, polar & uncharged (S, T, C, Y, N, Q); red, polar & acidic (D, E); blue, polar & basic (K, R, H). Residues that are unique to BtKAI2c but otherwise invariant are highlighted with asterisks (*); residues 98 and 191 are highlighted with red boxes. Source data are provided as a Source Data file.

Supplementary Figure 4. BtD14a is functionally homologous to AtD14

Three independent transgenic lines of Arabidopsis *Atd14-1* were analysed for functional complementation of the mutant phenotype by an *AtD14pro:BtD14a* transgene. **a**, Rosette and

leaf morphology at 31 days post-germination. **b**, Plant height and number of primary rosette branches at 45 days post-germination. **c**, Quantification of height and branching parameters, *n* = 10 plants per genotype. Scale bars: 50 mm. Source data are provided as a Source Data file.

Supplementary Figure 5. The GFP-BtKAI2c transgene is faithfully transcribed in Arabidopsis

a, Structure of the *AtKAI2pro:mGFP6-BtKAI2c* transgene. Primers used for RT-PCR are shown with arrows. The promoter and 5′UTR are derived from *At4g37470* (*AtKAI2*). attB1, Gateway recombination site that links *mGFP6* and *BtKAI2c* regions; *nos ter*, nopaline synthase terminator. Not drawn to scale.

b, RT-PCR analysis of *GFP-BtKAI2c* transcripts after 35 cycles of amplification. Primers MW406 + BK010 target the transgene, while a second primer pair (MW275 + MW278) serves as a control and spans five introns of *At1g03055*. *kai2-2* serves as a non-transgenic control genotype. Templates: R, total RNA; D, genomic DNA; W, water only. RT, reverse transcriptase (Supercript III). DNA size standards (in base pairs) are indicated on the left, with anticipated PCR product sizes shown on the right and defined in the table.

c, The RT-PCR product generated with proof-reading polymerase (Q5, New England Biolabs) and primers MW406 and BK010 was cloned into pCR4-TOPO (Life Technologies). Five dideoxy sequence traces were aligned against the *GFP-BtKAI2c* transgene reference. No disagreements with the reference sequence were observed. Red bars indicate trimmed regions of sequence traces to remove low quality data.

d, Transient expression of BtKAI2a, BtKAI2b and BtKAI2c proteins in tobacco. Plasmids encoding N-terminal, c-myc-tagged proteins were transferred to Agrobacterium, and the resulting strains used to infiltrate tobacco leaves. After 96 h, samples were harvested in triplicate (two to three leaves per sample). Mock-treated leaves were transformed with a plasmid encoding a non-tagged protein. Sixty micrograms of total protein were separated by SDS-PAGE, blotted and challenged with anti-c-myc antibody (Genscript A00704). Band intensity was measured using ImageJ, and expression was normalised to intensity of the Rubisco large subunit (RbcL) band on the "stain-free" gel imaged under UV light. Error bars indicate SE, n = 3 replicates. Source data are provided as a Source Data file.

Supplementary Figure 6. The R98V mutation restores stability of BtKAI2c

a, Homology model of the native BtKAI2c protein. Highlighted in yellow are five residues and their corresponding mutations that were assessed for effects on BtKAI2c stability. The combination of E129D, D130V, E133Q and D173E in the hinge region between the lid domain (orange) and core domain (green) is the quadruple ("quad") mutant; combining this with R98V yields the quintuple ("quint") mutant. Red residues are the Ser-His-Asp catalytic triad. **b**, Transient expression of AtKAI2 and BtKAI2c variants in tobacco leaves following infiltration by *Agrobacterium* strains encoding GFP-KAI2 variants driven by the CaMV *35S* promoter. Three leaves were infiltrated on each of two different plants. After five days, 20 μg soluble protein was electrophoresed, blotted and challenged with antibodies against either anti-GFP (top) or anti-actin (bottom). **c**, Immunoblot of crude bacterial lysates expressing SUMO fusion proteins and challenged with anti-SUMO antibody. Negative values above each lane indicate $\log_{10}$ dilution factors of lysate; U, lysate from uninduced bacterial culture expressing SUMO-AtKAI2. **d**, SDS-PAGE analysis of protein purification runs of SUMO-BtKAI2c, -BtKAI2c^{R98V} and -BtKAI2c^{QUINT} harvested from 900 mL bacterial culture. Lys, clarified lysate; FT, column flow through after immobilisation on Co^{2+} affinity column; W, wash with 10 mM imidazole. Proteins were eluted with 200 mM imidazole in eight successive fractions. **e**, DSF curves of SUMO fusion proteins treated with 0-200 μM GR24$^{ent\text{-}5DS}$. Insets plot the minimum value of − (dF/dT) at the melting point of the protein as determined in the absence of ligand (means ± SE, n = 3). Significant differences from untreated control: * P < 0.05 ** P < 0.01 *** P < 0.001 (ANOVA). Source data are provided as a Source Data file.

Supplementary Figure 7. Germination profiles of transgenic Arabidopsis seeds expressing BtKAI2 homologues

Freshly harvested seed (three batches per genotype, each batch harvested from four plants) were removed from freezer storage, surface-sterilised and sown on 1% Phytagel supplemented with 0.1% acetone (mock), 1 μM KAR$_1$ or 1 μM KAR$_2$. Seed were incubated under constant light at 25 °C. Seed were examined for germination (radicle protrusion) 72 h after sowing and every 24 h thereafter. Data are means ± SE of three independent seed batches and 75 seed per batch. For each transgene, two independent, homozygous transgenic lines were analysed. Data presented in Figure 3 of the main manuscript are derived from these data. Source data are provided as a Source Data file.

Supplementary Figure 8. BtKAI2 homology models and SDS-PAGE of SUMO-BtKAI2 fusion proteins used for DSF

a-d, Solved structure of AtKAI2 (PDB: 3w06, ref. 2), AtD14 (PDB: 4IH4, ref. 3) and predicted homology models of BtKAI2a and BtKAI2b. Coloured surfaces depict internal cavities; values indicate the volumes of the primary ligand-binding cavities, adjacent to the catalytic Ser-His-Asp residues. Also shown is a variable secondary pocket, to the left of the primary pocket in these images.

e, To assess purity after affinity chromatography, five micrograms of each purified protein was electrophoresed on a 12% acrylamide gel containing 2,2,2-trichloroethanol and visualised under UV light. Protein size standards at 75 and 25 kDa (Bio-Rad Precision Plus Dual Colour) fluoresce strongly under UV light. Source data are provided as a Source Data file.

Supplementary Figure 9. BtKAI2a and BtKAI2b do not respond to karrikins in DSF assays

a, Differential scanning fluorimetry curves of SUMO-BtKAI2a and SUMO-BtKAI2b in presence of 0–200 μM KAR_1 or KAR_2. **b**, DSF responses of SUMO-tagged BtKAI2b (left) or untagged BtKAI2b (right) to 0–200 μM GR24*ent-5DS*. **c**, DSF responses to 0–200 μM of the two enantiomers of GR24. Data are means of eight (**a** and **c**) or four (**b**) technical replicates at each concentration of ligand. Source data are provided as a Source Data file.

Supplementary Figure 10. Stable transgenic expression of BtKAI2 valine–leucine exchange proteins in Arabidopsis

a, Immunoblots of total soluble protein extracted from 7-day-old seedlings of independent transgenic lines segregating in a 3:1 ratio for hygromycin resistance. Transgene expression in six lines expressing *GFP-BtKAI2a^{L98;L191}* (upper panels) and five expressing *GFP-BtKAI2b^{V98;V191}* (lower panels) were compared to a representative unmodified control (*GFP-BtKAI2a* 1F and *GFP-BtKAI2b* 9A respectively). Based on expression level two lines of each construct (3-5 and 3-14; 6-4 and 6-13) were selected and brought to homozygosity for further experiments. Protein blots were challenged with anti-GFP antibody. Equal gel loading was assessed by imaging total protein prior to blotting; the RbcL band is shown. **b**, Rosette phenotypes of homozygous individuals expressing native and modified *GFP-BtKAI2* transgenes. Plants were 25 days old and grown under long day conditions as described in Methods. Scale bar: 50 mm. Source data are provided as a Source Data file.

Supplementary Figure 11. Three experimental replicates of hypocotyl elongation assays with BtKAI2a and BtKAI2b transgenics (double exchange of residues 98 & 191)

Each panel depicts data from an independent experiment performed on a separate date,

which are shown in summarised format in Figure 4. Data are means ± SE, n=24 to 40

seedlings. Each dot corresponds to an individual seedling. Source data are provided as a

Source Data file.

Supplementary Figure 12. Germination profiles of transgenic Arabidopsis seeds

expressing BtKAI2 homologues (double exchange of residues 98 and 191)

Freshly harvested seeds (three batches per genotype, each batch harvested from three plants)

were removed from freezer storage, surface-sterilised and sown on 0.7% agar supplemented

with 0.02% acetone (mock), 1 µM KAR_1 or 1 µM KAR_2. Seeds were incubated under constant

light at 25 °C. Seeds were examined for germination (radicle protrusion) 72 h after sowing

and every 24 h thereafter. Data are means ± SE of three independent seed batches and 100

seed per batch. For the $BtKAI2a^{L98;L191}$ and $BtKAI2b^{V98;V191}$ transgenes, two independent,

homozygous transgenic lines were analysed. Source data are provided as a Source Data file.

Supplementary Figure 13. Three experimental replicates of hypocotyl elongation

assays with BtKAI2b transgenics (individual exchange of residues 98 and 191)

Each panel depicts data from a separate experiment performed on the indicated date, which

are shown in summarised format in Figure 5. Data are means ± SE, n = 19-21 seedlings. Each

dot corresponds to an individual seedling. Source data are provided as a Source Data file.

Figure 1. *Brassica tournefortii* is most sensitive to KAR$_1$, the major karrikin analogue isolated from plant-derived smoke
a, Germination responses of *B. tournefortii* seed to KAR$_1$ and KAR$_2$. Data are cumulative germination after 11 days (means ± SE; *n* = 3 biological replicates per treatment, ≥35 seeds per replicate). **b**, Levels of karrikin-responsive transcripts *BtDLK2* and *BtSTH7* in *B. tournefortii* seed. Seed were imbibed for 24 hours in the dark supplemented with KAR$_1$ and KAR$_2$. Transcripts were normalised to *BtCACS* reference transcripts. Data are means ± SE, *n* = 3 biological replicates, ≥50 seeds per replicate. **c, d**, Hypocotyl elongation responses of *B. tournefortii* seedlings grown for four days under continuous red light on water-saturated glass filter paper containing 0.1% acetone (mock), KAR$_1$ or KAR$_2$. Data are means ± 95% CI of *n* = 18 to 24 seedlings. Scale bar: 10 mm. **e**, Levels of *BtDLK2* transcripts in *B. tournefortii* seedlings grown under the same conditions as for d. Data are means ± SE, *n* = 3 biological replicates, ≥20 seedlings per replicate. In all panels, asterisks denote significance levels (ANOVA) between indicated conditions: * *P*<0.05; ** *P*<0.01; *** *P*<0.001; **** *P*<0.0001. Source data are provided as a Source Data file.

Figure 2. Two differentially expressed *B. tournefortii* KAI2 homologues are functional in Arabidopsis

a, Maximum likelihood phylogeny of KAI2 homologues in the Brassicaceae, based on nucleotide data. Node values represent bootstrap support from 100 replicates. A *KAI2* sequence from *Gossypium raimondii* (Malvaceae) serves as an outgroup. Tree shown is a subset of a larger phylogeny in Supplementary Figure 2. **b**, Transcript levels of the three *BtKAI2* homologues in *B. tournefortii* seed imbibed for 24 h (left) and four-day-old *B. tournefortii* seedlings (right) treated with 0.1% acetone or 1 μM KAR$_1$ for 24 h. Data are means ± SE, n = 3 biological replicates. **c, d,** Seedling and rosette phenotypes of two independent transgenic lines of Arabidopsis homozygous for *KAI2pro:GFP-BtKAI2* transgenes. Scale bars: 5 mm (**c**); 50 mm (**d**).**e**, Immunoblots of soluble proteins challenged with antibodies against KAI2 (upper panel), GFP (middle panel) or actin as a loading control (lower panel). Anti-KAI2 detects both the native AtKAI2 protein (30 kDa) and the GFP-BtKAI2 fusion proteins (58 kDa) but detects BtKAI2b relatively poorly. Non-specific bands are marked with asterisks. Protein was isolated from pools of approximately fifty 7-day-old seedlings. Source data are provided as a Source Data file.

Figure 3. Functional divergence between BtKAI2 homologues in seeds and seedlings of transgenic Arabidopsis
a, Germination responses of primary dormant Arabidopsis seed homozygous for *KAI2pro:GFP-BtKAI2* transgenes in the *kai2-2* background. Germination values were determined 120 h after sowing. Extended germination curves are shown in Supplementary Fig. 7. Data are means ± SE of n = 3 independent seed batches, 75 seed per batch. **b**, Levels of *KUF1* transcripts in *KAI2pro:GFP-BtKAI2* seeds treated with 1 μM KAR$_1$ or KAR$_2$. Expression was normalised to *CACS* reference transcripts and scaled to the value for mock-treated seed within each genotype. Data are means ± SE of n = 3 biological replicates. **c**, Hypocotyl elongation responses of *KAI2pro:GFP-BtKAI2* seedlings treated with KAR$_1$ or KAR$_2$. Data are means ± SE of n = 3 biological replicates, 12–18 seedlings per replicate. **d**, Levels of *DLK2* transcripts in 8-day-old *KAI2pro:GFP-BtKAI2* seedlings treated with KAR$_1$ or KAR$_2$ for eight hours. Expression was normalised to *CACS* reference transcripts and scaled to the value for mock-treated seedlings within each genotype. Data are means ± SE of n = 3 biological replicates. Pairwise significant differences: * P < 0.05 ** P < 0.01 *** P < 0.001; ns, P > 0.05 (two-way ANOVA). Source data are provided as a Source Data file.

Figure 4. Two residues contribute to ligand specificity between BtKAI2a and BtKAI2b

a, DSF curves of SUMO-BtKAI2a and SUMO-BtKAI2b fusion proteins treated with 0-50 μM GR24$^{ent\text{-}5DS}$, a KAI2-bioactive ligand. Each curve is the average of three sets of reactions, each comprising four technical replicates. Insets plot the minimum value of –(dF/dT) at the melting point of the protein as determined in the absence of ligand (means ± SE, n = 3). Significant differences from untreated control: * $P < 0.05$ ** $P < 0.01$ *** $P < 0.001$ (ANOVA). **b**, DSF curves of SUMO-BtKAI2a$^{L98;L191}$ and SUMO-BtKAI2b$^{V98;V191}$ fusion proteins treated with 0-50 μM GR24$^{ent\text{-}5DS}$. **c**, Hypocotyl elongation responses of Arabidopsis expressing GFP-BtKAI2a$^{L98;L191}$ and GFP-BtKAI2b$^{V98;V191}$ fusion proteins and treated with KAR$_1$ or KAR$_2$. Data are a summary of three experimental replicates performed on separate occasions, each comprising 25-40 seedlings per genotype/treatment combination. Data for each replicate are shown in Supplementary Figure 11. Error bars are SE, n = 3 experimental replicates; each dot corresponds to the mean value derived from each replicate. Asterisks denote significant differences: * P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001 (linear mixed model with experimental replicate as a random effect; specific pairwise comparisons using Tukey's HSD correction). **d**, Levels of *DLK2* transcripts in the same transgenic lines as above treated with 1 μM KAR$_1$, 1 μM KAR$_2$, or 0.1% acetone and harvested 8 hours later. Expression was normalised to *CACS* reference transcripts. Data are means ± SE of n = 3 pools of ~50 seedlings treated in parallel. Asterisks denote significant differences as above (two-way ANOVA; specific pairwise comparisons using Tukey's HSD correction). Source data are provided as a Source Data file.

Figure 5. Diversity at positions 96 and 189 in KAI2 homologues across angiosperms

a, Sequence logos generated from 476 aligned KAI2 homologues sampled from eudicots, monocots and magnoliids and centred on position 96 (left) and 189 (right). **b**, Species tree of the Saxifragales based upon 410 single-copy nuclear genes estimated using the ASTRAL gene tree method (Leebens-Mack et al., 2019). Branch support values are shown only where less than 1. The four-letter taxon prefix corresponds to the 1000 Plant Transcriptomes project Sample ID. Taxa are highlighted by colour according to identity of amino acids of KAI2 orthologues at positions 96 and 189 respectively: black, VV; blue, LV; red, LL; magenta, other. Postulated ancestral identities at specific nodes are indicated with filled circles. **c**, Hypocotyl elongation responses to karrikins in three independent transgenic lines homozygous for *GFP-BtKAI2a*[L98;V191] and three lines for *GFP-BtKAI2b*[V98;L191]. Data shown are a summary of three experimental replicates performed on separate occasions, each comprising ~20 seedlings per genotype/treatment combination. Data from each replicate are presented in Supplementary Figure 13. Error bars are SE, n = 3 experimental replicates; each dot corresponds to the mean value derived from each replicate. Asterisks denote significant differences: * P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001 (linear mixed model with experimental replicate as a random effect; specific pairwise comparisons using Tukey's HSD correction). Source data are provided as a Source Data file.

Supplementary Figure 1. Germination of *Brassica tournefortii* seeds treated with karrikins
a, Structures of KAR$_1$, KAR$_2$ and two enantiomers of GR24. **b,** Germination response of the "Merridin" batch of *B. tournefortii* seed treated with KAR$_1$ and KAR$_2$ after three days. **c-d,** Germination response of the "Perth" batch to KAR$_1$ (**c**) and KAR$_2$ (**d**). Data in Figure 1a are derived from the data shown in **c** and **d. e-f**, Rates of KAR$_1$ and KAR$_2$ uptake by imbibed seed, as determined by GC-MS. All error bars are mean ± SE of n = 3 batches of 75 seed (b-d) or 3 samples of 40 mg (e) or 20 mg (f) seed as described in Methods. Source data are provided as a Source Data file.

Supplementary Figure 2. Extended phylogeny of eu-KAI2 and D14 proteins in angiosperms
Maximum likelihood phylogeny of KAI2 and D14 homologues in the Brassicaceae and monocots, based on nucleotide data. Node values represent bootstrap support from 100 replicates. A KAI2 sequence from *Selaginella moellendorffii* (SmKAI2a) serves as an outgroup for the eu-KAI2 clade. Source data are provided as a Source Data file.

Supplementary Figure 3. Alignment of *Brassica* KAI2 sequences

Full length protein coding regions of KAI2 homologues from four *Brassica* species (*Brassica tournefortii*, *B. rapa*, *B. nigra* and *B. oleracea*) were translated from database nucleic acid sequences and aligned to Arabidopsis KAI2 using MAFFT (ref. 1) implemented in Geneious R10 software (Biomatters Ltd). Amino acid residues are coloured according to polarity: yellow, non-polar (G, A, V, L, I, F, W, M, P); green, polar & uncharged (S, T, C, Y, N, Q); red, polar & acidic (D, E); blue, polar & basic (K, R, H). Residues that are unique to BtKAI2c but otherwise invariant are highlighted with asterisks (*); residues 98 and 191 are highlighted with red boxes. The underlined region indicates the epitope used to raise the antibody against AtKAI2. Source data are provided as a Source Data file.

Supplementary Figure 4. BtD14a is functionally homologous to AtD14

Three independent transgenic lines of Arabidopsis *Atd14-1* were analysed for functional complementation of the mutant phenotype by an *AtD14pro:BtD14a* transgene. **a**, Rosette and leaf morphology at 31 days post-germination. **b**, Plant height and number of primary rosette branches at 45 days post-germination. **c**, Quantification of height and branching parameters, means ± SE, *n* = 10 plants per genotype. Scale bars: 50 mm. Source data are provided as a Source Data file.

a

b

Primers	cDNA (R)	gDNA (D)
MW406 + BK010	1721	1721
MW275 + MW278	425	1191

c

d

Supplementary Figure 5. The *GFP-BtKAI2c* transgene is faithfully transcribed in Arabidopsis

a, Structure of the *AtKAI2pro:mGFP6-BtKAI2c* transgene. Primers used for RT-PCR are shown with arrows. The promoter and 5′UTR are derived from *At4g37470* (*AtKAI2*). attB1, Gateway recombination site that links mGFP6 and BtKAI2c regions; *nos* ter, nopaline synthase terminator. Not drawn to scale.

b, RT-PCR analysis of *GFP-BtKAI2c* transcripts after 35 cycles of amplification. Primers MW406 + BK010 target the transgene, while a second primer pair (MW275 + MW278) serves as a control and spans five introns of *At1g03055*. *kai2-2* serves as a non-transgenic control genotype. Templates: R, total RNA; D, genomic DNA; W, water only. RT, reverse transcriptase (Supercript III). DNA size standards (in base pairs) are indicated on the left, with anticipated PCR product sizes shown on the right and defined in the table.

c, The RT-PCR product generated with proof-reading polymerase (Q5, New England Biolabs) and primers MW406 and BK010 was cloned into pCR4-TOPO (Life Technologies). Five dideoxy sequence traces were aligned against the *GFP-BtKAI2c* transgene reference. No disagreements with the reference sequence were observed. Red bars indicate trimmed regions of sequence traces to remove low quality data.

d, Transient expression of BtKAI2a, BtKAI2b and BtKAI2c proteins in tobacco. Plasmids encoding N-terminal, c-myc-tagged proteins were transferred to Agrobacterium, and the resulting strains used to infiltrate tobacco leaves. After 96 h, samples were harvested in triplicate (two to three leaves per sample). Mock-treated leaves were transformed with a plasmid encoding a non-tagged protein. Sixty micrograms of total protein were separated by SDS-PAGE, blotted and challenged with anti-c-myc antibody (Genscript A00704). Band intensity was measured using ImageJ, and expression was normalised to intensity of the Rubisco large subunit (RbcL) band on the "stain-free" gel imaged under UV light. Error bars indicate SE, *n* = 3 replicates. Source data are provided as a Source Data file.

Supplementary Figure 6. The R98V mutation restores stability of BtKAI2c

a, Homology model of the native BtKAI2c protein. Highlighted in yellow are five residues and their corresponding mutations that were assessed for effects on BtKAI2c stability. The combination of E129D, D130V, E133Q and D173E in the hinge region between the lid domain (orange) and core domain (green) is the quadruple ("quad") mutant; combining this with R98V yields the quintuple ("quint") mutant. Red residues are the Ser-His-Asp catalytic triad. **b**, Transient expression of AtKAI2 and BtKAI2c variants in tobacco leaves following infiltration by Agrobacterium strains encoding GFP-KAI2 variants driven by the CaMV *35S* promoter. Three leaves were infiltrated on each of two different plants. After five days, 20 μg soluble protein was electrophoresed, blotted and challenged with antibodies against either anti-GFP (top) or anti-actin (bottom). **c**, Immunoblot of crude bacterial lysates expressing SUMO fusion proteins and challenged with anti-SUMO antibody. Negative values above each lane indicate $\log_{10}$ dilution factors of lysate; U, lysate from uninduced bacterial culture expressing SUMO-AtKAI2.**d**, SDS-PAGE analysis of protein purification runs of SUMO-BtKAI2c, -BtKAI2c^{R98V} and -BtKAI2c^{QUINT} harvested from 900 mL bacterial culture. Lys, clarified lysate; FT, column flow through after immobilisation on Co^{2+} affinity column; W, wash with 10 mM imidazole. Proteins were eluted with 200 mM imidazole in eight successive fractions. **e**, DSF curves of SUMO fusion proteins treated with 0-200 μM GR24$^{ent-5DS}$. Insets plot the minimum value of –(dF/dT) at the melting point of the protein as determined in the absence of ligand (means ± SE, n = 3). Significant differences from untreated control: * P < 0.05 ** P < 0.01 *** P < 0.001 (ANOVA). Source data are provided as a Source Data file.

Supplementary Figure 7. Germination profiles of transgenic Arabidopsis seeds expressing native BtKAI2 homologues

Freshly harvested seed (three batches per genotype, each batch harvested from four plants) were removed from freezer storage, surface-sterilised and sown on 1% Phytagel supplemented with 0.1% acetone (mock), 1 μM KAR$_1$ or 1 μM KAR$_2$. Seed were incubated under constant light at 25 °C. Seed were examined for germination (radicle protrusion) 72 h after sowing and every 24 h thereafter. Data are means ± SE of three independent seed batches and 75 seed per batch. For each transgene, two independent, homozygous transgenic lines were analysed. Data presented in Figure 3 of the main manuscript are derived from these data. Source data are provided as a Source Data file.

Supplementary Figure 8. BtKAI2 homology models and SDS-PAGE of SUMO-BtKAI2 fusion proteins used for DSF.
a-d, Solved structure of AtKAI2 (PDB: 3w06, ref. 2), AtD14 (PDB: 4IH4, ref. 3) and predicted homology models of BtKAI2a and BtKAI2b. Coloured surfaces depict internal cavities; values indicate the volumes of the primary ligand-binding cavities, adjacent to the catalytic Ser-His-Asp residues. Also shown is a variable secondary pocket, to the left of the primary pocket in these images.
e, To assess purity after affinity chromatography, five micrograms of each purified protein was electrophoresed on a 12% acrylamide gel containing 2,2,2-trichloroethanol and visualised under UV light. Protein size standards at 75 and 25 kDa (Bio-Rad Precision Plus Dual Colour) fluoresce strongly under UV light. Source data are provided as a Source Data file.

Supplementary Figure 9. BtKAI2a and BtKAI2b do not respond to karrikins in DSF assays

a, Differential scanning fluorimetry curves of SUMO-BtKAI2a and SUMO-BtKAI2b in presence of 0–200 µM KAR_1 or KAR_2. **b**, DSF responses of SUMO-tagged BtKAI2b (left) or untagged BtKAI2b (right) to 0–200 µM GR24$^{ent-5DS}$. c, DSF responses to 0–200 µM of the two enantiomers of GR24. Data are means of eight (a and c) or four (b) technical replicates at each concentration of ligand. Source data are provided as a Source Data file.

Supplementary Figure 10. Stable transgenic expression of BtKAI2 valine–leucine double exchange proteins in Arabidopsis

a, Immunoblots of total soluble protein extracted from 7-day-old seedlings of independent transgenic lines segregating in a 3:1 ratio for hygromycin resistance. Transgene expression in six lines expressing GFP-BtKAI2a$^{L98;L191}$ (upper panels) and five expressing GFP-BtKAI2b$^{V98;V191}$ (lower panels) were compared to a representative unmodified control (GFP-Bt-KAI2a 1F and GFP-BtKAI2b 9A respectively). Based on expression level two lines of each construct (3-5 and 3-14; 6-4 and 6-13) were selected and brought to homozygosity for further experiments. Protein blots were challenged with anti-GFP antibody. Equal gel loading was assessed by imaging total protein prior to blotting; the RbcL band is shown. **b**, Rosette phenotypes of homozygous individuals expressing native and modified *GFP-BtKAI2* transgenes. Plants were 25 days old and grown under long day conditions as described in Methods. Scale bar: 50 mm. Source data are provided as a Source Data file.

Supplemental Figure 11. Three experimental replicates of hypocotyl elongation assays with BtKAI2a and BtKAI2b transgenics (double exchange of residues 98 & 191)
Each panel depicts data from a separate experiment performed on the indicated date, which are shown in summarised format in Figure 4. Data are means ± SE, n=24 to 40 seedlings. Each dot corresponds to an individual seedling. Source data are provided as a Source Data file.

Supplementary Figure 12. Germination profiles of transgenic Arabidopsis seeds expressing BtKAI2 homologues (double exchange of residues 98 and 191)

Freshly harvested seeds (three batches per genotype, each batch harvested from three plants) were removed from freezer storage, surface-sterilised and sown on 0.7% agar supplemented with 0.02% acetone (mock), 1 µM KAR_1 or 1 µM KAR_2. Seeds were incubated under constant light at 25 °C. Seeds were examined for germination (radicle protrusion) 72 h after sowing and every 24 h thereafter. Data are means ± SE of three independent seed batches and 100 seed per batch. For the $BtKAI2a^{L98;L191}$ and $BtKAI2b^{V98;V191}$ transgenes, two independent, homozygous transgenic lines were analysed. Source data are provided as a Source Data file.

Supplemental Figure 13. Three experimental replicates of hypocotyl elongation assays with BtKAI2b transgenics (individual exchange of residues 98 & 191)

Each panel depicts data from a separate experiment performed on the indicated date, which are shown in summarised format in Figure 5. Data are means ± SE, n = 19-21 seedlings. Each dot corresponds to an individual seedling. Source data are provided as a Source Data file.