Divergent receptor proteins confer responses to different karrikins in two ephemeral weeds

Yueming Kelly Sun[1,2], Adrian Scaffidi[1], Jiaren Yao[1], Kim Melville[1], Steven M Smith[3,4], Gavin R Flematti[1] and Mark T Waters[1,2,5]

1. School of Molecular Sciences, The University of Western Australia, 35 Stirling Hwy, Perth, WA 6009, Australia
2. Australian Research Council Centre of Excellence in Plant Energy Biology, The University of Western Australia, 35 Stirling Hwy, Perth, WA 6009, Australia
3. School of Natural Sciences, The University of Tasmania, Hobart, TAS 7000, Australia
4. Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, 1 West Beichen Road, Beijing 100101, PR China.
5. Author for correspondence: mark.waters@uwa.edu.au

Word count (Abstract and main text): 2396 words

Abstract

Seed germination is responsive to environmental chemical cues, such as karrikins, which are germination stimulants produced from wildfires[1,2]. Seeds of *Brassica tournefortii* – a widespread problematic weed associated with post-fire and disturbance sites – are highly sensitive to KAR_1, the most abundant karrikin, whereas *Arabidopsis thaliana* responds more sensitively to KAR_2, a less abundant karrikin[3,4]. The mechanism for these differential responses is unclear. KARRIKIN INSENSITIVE2 (KAI2), a paralogue of the strigolactone receptor DWARF14 (D14), is the putative karrikin receptor identified from *Arabidopsis*[4,5]. Here we show that *B. tournefortii* exhibits greater sensitivity for KAR_1 relative to KAR_2 in multiple biological assays. We demonstrate that *B. tournefortii* differentially expresses three *KAI2* homologues, two of which rescue the *Arabidopsis kai2-2* null mutant. Notably, the most abundantly expressed homologue in *B. tournefortii* is sufficient to confer enhanced responses to KAR_1 relative to KAR_2 when expressed in *Arabidopsis*. We further identify two variant amino acid residues that support differential ligand preferences between the two *B. tournefortii* homologues. Our findings provide clear evidence that KAI2 is the receptor for karrikins, since specific amino acid substitutions near the active site change ligand specificity. Our results also suggest that duplication and diversification of KAI2 proteins could provide weedy ephemerals with the ability to respond sensitively to different karrikins or related compounds produced by environmental disturbance.

Main

Recovery of vegetation after wildfires results in part from the germination of dormant seed in the soil. Plant-derived smoke contains organic and inorganic compounds that cue the seed of many species to germinate under post-fire conditions[6,7]. Among the organic compounds are karrikins, a family of butenolides that comprises six known members. In samples of smoke water generated by burning straw, KAR_1 is the most abundant karrikin, followed by KAR_2[8]. These two analogues differ by the presence or absence, respectively, of a methyl group on the butenolide ring (Supplementary Fig. 1a).

Karrikin mode of action has been elucidated by studies using *Arabidopsis thaliana*, a weedy ephemeral widely distributed in temperate climates throughout the northern hemisphere. Although seed germination in *A. thaliana* is stimulated by karrikins[3], this species is not

specifically associated with fire-prone habitats. *A. thaliana* is unusual in that it responds preferentially to KAR_2, whereas most karrikin-responsive species, including fire-followers, respond preferentially to KAR_1[1,9]. To understand the molecular basis for differential sensitivity to distinct karrikins, we chose to investigate a different member of the Brassicaceae that responds preferentially to KAR_1.

Brassica tournefortii (Brassicaceae; sahara mustard) is an invasive weed native to the Mediterranean and Middle East that blights many semi-arid areas in North America, Australia and South Africa. Infestations with *B. tournefortii* lead to dramatic reductions in plant cover of native annuals, which suffer reduced stem and flower production, especially after relatively wet winters[10]. *B. tournefortii* seed can persist in the soil for many seasons, undergoing wet-dry cycling that can influence dormancy and contribute to boom-bust cycles that outcompete native species[11-13]. Semi-arid Mediterranean-type ecosystems are also prone to wildfires. As an invasive species, *B. tournefortii* may radically alter fire frequency and intensity by influencing fuel profiles[14,15], further exacerbating the impact of fire on susceptible native ecosystems. In addition, *B. tournefortii* is particularly responsive to KAR_1 and shows a positive germination response in the nanomolar range[16]. Accordingly, although *B. tournefortii* is not dependent on fire to spread, it is particularly well positioned to invade areas disturbed by fire events[17,18].

To characterise the karrikin response of *B. tournefortii*, we performed multiple physiological and molecular assays comparing KAR_1 with KAR_2. First, germination of *B. tournefortii* seeds was consistently more sensitive to KAR_1 than KAR_2 at 10 nM, 100 nM and 1 µM (Fig. 1a and Supplementary Fig. 1b-d). Second, expression of two karrikin-responsive genes, *BtDLK2* and *BtSTH7*, required ten-fold higher concentrations of KAR_2 (1 µM) compared with KAR_1 (100 nM) for equivalent responses (Fig. 1b). Third, we found that KAI2 protein in *B. tournefortii* (BtKAI2) degraded upon karrikin treatment, as is the case for *Arabidopsis* KAI2 (AtKAI2)[19]. Degradation of BtKAI2 in *B. tournefortii* seed was approximately 10-fold more sensitive to KAR_1 than to KAR_2, such that after 48 h and 72 h treatment, 100 nM KAR_1 and 1 µM KAR_2 induced similar levels of KAI2 degradation (Fig. 1c). To rule out differential karrikin uptake to account for these differences, we tested the absorption of KAR_1 and KAR_2 and found that both were taken up at similar rates from solution by seeds during imbibition (Supplementary Fig. 1e-f). Besides promoting germination of primary dormant seeds, karrikins also inhibited hypocotyl elongation in *B. tournefortii* seedlings, as is the case in *Arabidopsis*; again, KAR_1 showed a stronger effect than KAR_2, most clearly at 1 µM (Fig. 1d). Therefore, we conclude

Figure 1: B. *tournefortii* is highly sensitive to KAR$_1$, the major karrikin analogue isolated from smoke

a, Germination responses of *B. tournefortii* seed to KAR$_1$ and KAR$_2$. Data are means ± SE; *n* = 3 biological replicates per treatment, ≥35 seeds per replicate. **b**, Levels of karrikin-responsive transcripts *BtDLK2* and *BtSTH7* in *B. tournefortii* seed. Seed were imbibed for 24 hours in the dark supplemented with KAR$_1$ and KAR$_2$. Transcripts were normalised to *BtCACS* reference transcripts. Data are means ± SE, *n* = 3 biological replicates, ≥50 seeds per replicate. **c**, Degradation of BtKAI2 protein levels in *B. tournefortii* seeds treated with KAR$_1$ and KAR$_2$ for 24, 48, and 72 hours. For each sample, 30 µg total protein isolated from imbibed seeds was separated by SDS-PAGE, blotted, and challenged with anti-KAI2 antibody. The reference panel shows the blot stained by amido black. Samples from *A. thaliana* ecotype L*er* and *kai2-2* seedlings serve as controls for antibody specificity. **d**, Hypocotyl elongation responses of *B. tournefortii* seedlings treated with KAR$_1$ and KAR$_2$. Data are means ± SE of *n* = 3 biological replicates, ≥12 seedlings per replicate. Scale bar: 10 mm.

that *B. tournefortii* is more sensitive to KAR$_1$ than to KAR$_2$. This pattern of ligand preference is

true of many karrikin-responsive species from ecosystems prone to wildfires[1,9], but notably is

opposite to that of *Arabidopsis*[3,4].

Orthologues of *KARRIKIN INSENSITIVE2/HYPOSENSITIVE TO LIGHT* (*KAI2/HTL*) are

ubiquitous in land plants, and are normally present as a single gene copy within an ancient

and highly conserved "eu-KAI2" clade[20]. There is growing evidence that, beyond its ability to

mediate karrikin responses, KAI2 has a core ancestral role in perceiving an endogenous

karrikin-like ligand ("KL") that regulates seed germination, seedling development, leaf shape

and cuticle development[21-23]. The genus *Brassica* has undergone a whole genome triplication

event since its divergence from the Arabidopsis lineage[24-26]. This process might have allowed

additional KAI2 copies to gain altered substrate specificity, potentially enhancing perception

of environmental signals such as karrikins from smoke.

To establish whether there are multiple *KAI2* homologues present in *B. tournefortii*, we

examined transcriptomes from seeds and seedlings. Three putative *KAI2* homologues were

identified (*BtKAI2a*, *BtKAI2b*, and *BtKAI2c*; Fig. 2a; Supplementary Fig. 2 and 3). *BtKAI2a*

grouped with *AtKAI2* and those of other Brassicaceae within a single clade, whereas *BtKAI2b*

and *BtKAI2c* grouped within a clade unique to *Brassica*. This phylogenetic pattern implies that

BtKAI2a is the ancestral copy of *KAI2*, whereas *BtKAI2b* and *BtKAI2c* arose more recently via

genome triplication within *Brassica*. All three *BtKAI2* transcripts were expressed in *B.*

tournefortii seeds, but only *BtKAI2a* and *BtKAI2b* could be detected in seedlings (Fig. 2b). In

both tissues, *BtKAI2b* transcripts were the most abundantly expressed of the three. We also

identified two *BtD14* homologues, at least one of which is functionally orthologous to *AtD14*

(Supplementary Figs. 2 and 4).

We performed transgenic complementation of the *Arabidopsis kai2-2* null mutant by

expressing each of the three isoforms as a GFP fusion protein driven by the *Arabidopsis KAI2*

promoter (*KAI2pro*:*GFP-BtKAI2*). Such fusions of GFP with KAI2 proteins have been used

previously with to analyse KAI2 activity[19,27,28]. Both *BtKAI2a* and *BtKAI2b* complemented the

seedling and leaf phenotypes of *kai2-2*, whereas *BtKAI2c* did not, despite being expressed at

the transcriptional level (Fig. 2c,d and Supplementary Fig. 5c). We could not detect GFP-

BtKAI2c protein in three independent transgenic lines, while GFP-BtKAI2a and GFP-BtKAI2b

accumulated at consistent levels (Fig. 2e). Following transient expression in tobacco leaves,

levels of BtKAI2c protein were considerably lower than for BtKAI2a and BtKAI2b, suggesting

Figure 2 Two differentially expressed *B. tournefortii* KAI2 homologues are functional in Arabidopsis
a, Maximum likelihood phylogeny of KAI2 homologues in the Brassicaceae, based on nucleotide data. Node values represent bootstrap support from 100 replicates. A *KAI2* sequence from *Gossypium raimondii* (Malvaceae) serves as an outgroup. Tree shown is a subset of a larger phylogeny in Supplementary Figure 2. **b**, Transcript levels of the three *BtKAI2* homologues in seed imbibed for 24 h (left) and four-day-old seedlings treated with or without 1 µM KAR_1 for 24 h. **c, d,** Seedling and rosette phenotypes of two independent transgenic lines of Arabidopsis homozygous for *KAI2pro:GFP-BtKAI2x* transgenes. **e**, Immunoblots of soluble proteins challenged with antibodies against KAI2 (upper panel), GFP (middle panel) or actin (lower panel). Non-specific bands are marked with asterisks. Protein was isolated from pools of approximately fifty 7-day-old seedlings.

that BtKAI2c may be poorly expressed in plant cells (Supplementary Fig. 5b). BtKAI2c carries
several unique amino acid substitutions, which together may compromise protein folding and
stability (Supplementary Fig. 3). Therefore, we conclude that BtKAI2a and BtKAI2b are
functionally orthologous to AtKAI2 in regulating plant development, whereas BtKAI2c is non-
functional and may constitute a pseudogene.

We further characterised the ligand preference of *BtKAI2* homologues by performing
physiological and molecular assays with the transgenic lines. Primary-dormant Arabidopsis
seeds homozygous for the transgenes were tested for germination response. Germination of
BtKAI2b transgenic seeds was more responsive to KAR_1 than KAR_2 at 1 μM, whereas no
significant difference was observed for *BtKAI2a* transgenic seeds (Fig. 3a and Supplementary
Fig. 6). *BtKAI2c* transgenic seeds were insensitive to karrikins. Hypocotyl elongation
responses (Fig. 3b) and accumulation patterns of *DLK2* transcripts (Fig. 3c) agreed with the
germination response with respect to karrikin preferences. Therefore, and consistent with
their relative phylogenetic positions (Fig. 2a), we conclude that *BtKAI2a* has similarity to
AtKAI2 in terms of ligand preference, whereas *BtKAI2b* has a preference for KAR_1 over KAR_2.
As *BtKAI2b* is more highly expressed than *BtKAI2a* in *B. tournefortii* seed and seedlings, we
suggest that the ligand preference of BtKAI2b accounts for the enhanced KAR_1-
responsiveness of this species.

To investigate the ligand preferences of BtKAI2a and BtKAI2b in more detail, we performed
differential scanning fluorimetry (DSF) assays on purified recombinant proteins. DSF has
been used extensively for inferring the interaction of strigolactone-like compounds with D14-
and KAI2-related proteins[28-32]. Racemic GR24 is a widely-used synthetic strigolactone
analogue that consists of two enantiomers, of which GR24[ent-5DS] is bioactive via AtKAI2[33].
Catalytically inactive D14 and KAI2 variants do not respond to GR24 in DSF, suggesting that
the shift in thermal response results from ligand hydrolysis and a corresponding
conformational change in the receptor[28-30]. In DSF assays, AtKAI2 shows a specific response to
>100 μM GR24[ent-5DS] but, for unclear reasons, no response at all to karrikins[28]. Likewise, we
found that both BtKAI2a and BtKAI2b were also unresponsive to karrikins in DSF
(Supplementary Fig. 7a). Therefore, we used GR24[ent-5DS] as a surrogate substrate in DSF
assays to infer BtKAI2–ligand interactions. We found that AtKAI2 and BtKAI2a showed similar
responses to >100 μM GR24[ent-5DS], but the BtKAI2b response was clear at >25 μM
(Supplementary Fig. 7b). We then used a lower range of GR24[ent-5DS] concentrations (0–50 μM)
to determine the threshold for response, which we defined as a statistically significant

Figure 3. Functional divergence between BtKAI2 homologues

a, Germination responses of primary dormant Arabidopsis seed homozygous for *KAI2pro:GFP-BtKAI2* transgenes in the *kai2-2* background. Germination values were determined 120 h after sowing. Extended germination curves are shown in Supplementary Fig. 6. Data are means ± SE of *n* = 3 independent seed batches, 75 seed per batch. **b**, Hypocotyl elongation responses of *KAI2pro:GFP-BtKAI2x* seedlings treated with KAR$_1$ or KAR$_2$. Data are means ± SE of *n* = 3 biological replicates, 12–18 seedlings per replicate. **c**, Levels of *DLK2* transcripts in 8-day-old *KAI2pro:GFP-BtKAI2x* seedlings treated with KAR$_1$ or KAR$_2$ for eight hours. Expression was normalised to *CACS* reference transcripts and scaled to the value for mock-treated seedlings within each genotype. Data are means ± SE of *n* = 3 biological replicates. Pairwise significant differences: * $P < 0.05$ ** $P < 0.01$ *** $P < 0.001$; ns, $P > 0.05$ (ANOVA).

reduction in the maximal rate of change in fluorescence at the melting temperature of the
protein (T_m). Although BtKAI2a showed only a weak and non-significant response at 40 and
50 µM GR24$^{ent\text{-}5DS}$, BtKAI2b responded significantly at 10 µM and above (Fig. 4a). These
results suggest that BtKAI2b is more sensitive than BtKAI2a to GR24$^{ent\text{-}5DS}$, and that BtKAI2a is
most like AtKAI2 in this respect.

BtKAI2a and BtKAI2b differ in primary amino acid sequence at just 14 positions
(Supplementary Fig. 4). We postulated that differences in ligand preference might be
determined by amino acids in the vicinity of the ligand binding pocket. Protein structural
homology models revealed only two residues that differ in this region: V98 and V191 in
BtKAI2a, and L98 and L191 in BtKAI2b; the corresponding residues in AtKAI2 are V96 and
V189 (Fig. 4b-d). Residue 98 is immediately adjacent to the catalytic serine at the base of the
pocket. Residue 191 is located internally on αT4 of the lid domain which, in AtD14, is
associated with a major rearrangement of protein structure upon ligand binding that reduces
the size of the pocket[34]. Homology modelling suggests only subtle differences in the size and
shape of the primary ligand binding pocket of BtKAI2a and BtKAI2b (Fig. 4b-d), but L98
would make the BtKAI2b pocket more hydrophobic, which is consistent with KAR$_1$ being
more hydrophobic than KAR$_2$[8]. To determine if these residues are pertinent to ligand
response, we replaced the two valine residues of BtKAI2a with leucine residues, generating
the variant BtKAI2a(b), and vice-versa for BtKAI2b, generating the variant BtKAI2b(a). In DSF
assays, we found that exchanging the two residues was sufficient to switch the original
responses, such that BtKAI2a(b) responded sensitively to GR24$^{ent\text{-}5DS}$, but BtKAI2b(a) did not
(Fig. 4e). Finally, to validate the different effects observed with GR24$^{ent\text{-}5DS}$ upon BtKAI2a and
BtKAI2b *in vitro*, we examined the response of these proteins to GR24$^{ent\text{-}5DS}$ *in planta*. We
found that, in terms of induction of *DLK2* transcripts, *GFP-BtKAI2b* transgenic seedlings
responded more strongly than *GFP-BtKAI2a* and wild type L*er* seedlings to 0.1 µM GR24$^{ent\text{-}5DS}$
(Fig. 4f). Overall, these results suggest that two variant residues can account for differences in
substrate preference between BtKAI2a and BtKAI2b. Given the importance of these residues,
it is notable that the non-functional BtKAI2c carries a highly non-conservative Leu-to-Arg
substitution at position 98 (Supplementary Fig. 3).

Two crystal structures of KAI2 proteins from *Arabidopsis thaliana* and *Striga hermonthica* are
non-congruent with respect to KAR$_1$ binding position and orientation[5,35]. In addition, the
affinity of AtKAI2 for KAR$_1$ is poorly defined, with estimates of dissociation coefficients (Kd)
ranging from 4.6 µM[36] to 148 µM[37] using isothermal calorimetry; for comparison, the Kd value

Figure 4. Two residues account for ligand specificity between BtKAI2a and BtKAI2b

a, DSF curves of SUMO-BtKAI2a and SUMO-BtKAI2b fusion proteins treated with 0-50 μM GR24$^{ent\text{-}5DS}$, a KAI2-bioactive ligand. Each curve is the average of three sets of reactions, each comprising four technical replicates. Insets plot the minimum value of –(dF/dT) at the melting point of the protein as determined in the absence of ligand (means ± SE, *n* = 3). Significant differences from untreated control: * *P* < 0.05 ** *P* < 0.01 *** *P* < 0.001 (ANOVA). **b–d**, Solved structure of AtKAI2 (PDB: 3w06, Kagiyama et al 2013) (b) and derived homology models (c–d) of BtKAI2a and BtKAI2b. Coloured surfaces depict internal cavities; values indicate the volumes of the primary ligand-binding cavities, adjacent to the catalytic Ser-His-Asp residues. Also shown is a variable secondary pocket, to the left of the primary pocket in these images. **e**, DSF curves of SUMO-BtKAI2a(b) and SUMO-BtKAI2b(a) fusion proteins treated with 0-50 μM GR24$^{ent\text{-}5DS}$. **f**, Levels of *DLK2* transcripts (normalized to *CACS* reference transcripts) in two independent lines of *KAI2pro:GFP-BtKAI2a* and *-BtKAI2b* seedlings treated for two hours with 0.1 μM GR24$^{ent\text{-}5DS}$. Data are means ± SE; *n* = 3 biological replicates. Letters indicate significantly different groupings (ANOVA; P < 0.05).

of AtD14 for GR24 is 0.3 μM using the same technique[36]. As such, it has been difficult to
establish a definitive receptor-ligand relationship for KAI2 and karrikins. Our results provide
strong evidence that KAI2 has the capacity to confer karrikin specificity, which in turn
demonstrates that KAI2 is the receptor for karrikins. However, we cannot rule out the
possibility that karrikins interact with the receptor differently from GR24*ent-5DS*, which may
account for the inactivity of karrikins in DSF assays.

The ligand binding pocket of KAI2 is generally hydrophobic, but the key residues that confer
ligand specificity are not clear. In certain root-parasitic weeds in the Orobanchaceae,
substitutions of bulky residues found in AtKAI2 with smaller hydrophobic amino acids have
potentially improved the affinity for host-derived strigolactone ligands with different
structural configurations[38,39]. In the case of BtKAI2 proteins, the more subtle changes in
pocket residues likely account for fine-tuning between very similar ligands. Similarly subtle
changes have been reported for the rice gibberellin receptor GID1: changing Ile133 to Leu or
Val increases the affinity for GA_{34} relative to GA_4, which differ by just one hydroxyl group[40].

Gene duplication is a common feature in plant evolutionary histories that can allow genes to
acquire new functions[41]. In obligate parasitic weeds, duplication and diversification of
KAI2/HTL genes have shifted ligand specificity towards strigolactones. Crucially, the core
KAI2 function of perceiving an endogenous ligand ("KL") is presumably retained through
ancestral, non-divergent copies that are under purifying selection[38,39]. KAI2 diversity may also
reflect differing affinity for KL in different species and at different life stages. Our results are
consistent with a scenario in which both BtKAI2a and BtKAI2b have retained the core KAI2
function of perceiving KL to support plant development. However, BtKAI2b has also acquired
mutations that alter its substrate specificity, which in turn enhance sensitivity to the more
abundant karrikin released after fire. This diversity among KAI2 proteins potentially provides
B. tournefortii with a selective advantage in fire-prone environments, assisting its invasive
nature.

Recent findings indicate that KAI2 is an integrator that also modulates germination in
response to abiotic environmental signals, including temperature and salinity[42]. As seed
germination is a critical life stage that contributes to the invasiveness of a species, strategies
for weed control will benefit from specific knowledge of KAI2 sequence diversity and
expression profiles.

METHODS

Chemical synthesis

Karrikins (KAR$_1$ and KAR$_2$), ^{13}[C]$_5$-labelled karrikins and GR24 enantiomers (GR24^{5DS} and GR24$^{ent\text{-}5DS}$) were prepared as previously described[1,9,43-45].

Plant material

Arabidopsis *kai2-2* (L*er*) and *Atd14-1* (Col-0) mutants were previously described[4]. The seeds of *Brassica tournefortii* used in this work were collected in November and December 2009 from two sites in Western Australia (Kings Park in Perth, and Merridin) as described previously[46]. Seeds were dried to 15% relative humidity for one month prior to storage in air-tight bags at –20 °C.

Seed germination assays

Seed germination assays using Arabidopsis were performed on Phytagel as described previously[33]. *Brassica tournefortii* seeds were sowed in triplicates (35-70 seeds each) on glass microfibre filter paper (Grade 393; Filtech, NSW Australia) held in petri dishes and supplemented with mock or karrikin treatments. Treatments were prepared by diluting acetone (mock) or karrikin stocks (dissolved in acetone) 1:1000 with ultrapure water. The seeds were imbibed in the dark at 22 °C. Numbers of germinated seeds were counted each day until the germination percentages remained unchanged.

Hypocotyl elongation assays

Arabidopsis hypocotyl elongation assays were performed under red light as described previously[4]. For *B. tournefortii*, assays were performed with the following modifications to the Arabidopsis protocol: *B. tournefortii* seeds were sowed in triplicate on glass microfibre filter paper (Filtech) held in petri-dishes supplemented with mock or karrikin treatments. The seeds were imbibed in the dark for 22 h at 22 °C before exposing to continuous red light (20 μmol photons m^{-2} s^{-1}) for 3 days.

KAI2 degradation assays

B. tournefortii seeds were imbibed on glass fibre filters as described above and incubated in the dark at 22°C. Imbibition solutions contained the indicated concentration of karrikins.

Seeds were harvested after 24, 48 and 72 h and frozen in liquid nitrogen. Protein extraction
and immunoblotting were conducted as previously described[28]. The reference panels show a
representative region of each blot stained with amido black solution (0.03% w/v in 1:10
acetic acid:methanol).

Transcript analysis

Brassica tournefortii seeds were imbibed on glass fibre filters in the dark at 22 °C as described
above For transcript quantification in *B. tournefortii* seeds, samples were taken at the
indicated time points, briefly blot-dried on paper towel, and frozen in liquid nitrogen. For *B.*
tournefortii seedlings, seeds imbibed for 24 h were transferred to the growth room (~120-
150 μmol photons m^{-2} s^{-2}, 16 h light/8 h dark, 22 °C light/16 °C dark) for three days. A sample
of approximately 20 seedlings was then transferred to a 250 mL conical flask containing 50
mL ultrapure water supplemented with 1 μM KAR_1, or an equivalent volume of acetone (0.1%
v/v), with three replicate samples per treatment. The flasks were then shaken for 24 h before
seedlings were harvested.

Arabidopsis seedlings were grown on 0.5× MS agar under the same growth room conditions
for seven days. On the seventh day, seedlings were transferred to 3 mL liquid 0.5× MS
medium in 12-well culture plates (CORNING Costar 3513) and shaken at 70 rpm for a further
22 h. The medium was then removed by pipette and replaced with fresh medium containing
relevant compounds or an equivalent volume (0.1% v/v) acetone. After a further incubation
period with shaking (between 2 and 8 h), the seedlings were harvested, blotted dry, and
frozen in liquid nitrogen.

RNA extraction, DNase treatment, cDNA synthesis and quantitative PCR were conducted as
previously described [47]. Oligonucleotides are listed in the Supplementary Table 2.

Cloning and mutagenesis of *Brassica tournefortii KAI2* and *D14* homologues

The Gateway-compatible destination vector pMDC43[48] was digested with *Hind*III and *Kpn*I to
release the 35S promoter, which was replaced with the 776-bp *KAI2* promoter and 5′UTR as
defined previously [28], yielding the binary destination vector pKAI2pro-GFP-GW. Due to high
sequence similarity among BtKAI2 and BtD14 paralogues in the coding region, unique
transcripts were identified on the basis of distinct 3′UTRs. Full-length *BtKAI2* coding
sequences (and unique 3′UTR sequences) were amplified from *B. tournefortii* cDNA with the
attB sites using the universal forward primer BtKAI2_universal_F and homologue-specific

reverse primers RACE_R (BtKAI2a), Contig1_R (BtKAI2b) and Contig5_R (BtKAI2c), before cloning into pDONR207 (Life Technologies). The pDONR207 clones were confirmed by Sanger sequencing and recombined with pKAI2pro-GFP-GW to generate the binary plant expression plasmids pKAI2pro-GFP-BtKAI2a, pKAI2pro-GFP-BtKAI2b and pKAI2pro-GFP-BtKAI2c. For transient expression in tobacco, BtKAI2 coding sequences were transferred via Gateway-mediated recombination into pSKI106, which drives very high expression and encodes an N-terminal 3× *c*-myc tag[49]. The *BtD14a* coding sequence was amplified from cDNA using oligonucleotides BtD14_F and BtD14_R, cloned into pDONR207 as above, and transferred into pD14pro-GW [28].

The full-length *BtKAI2* coding sequences (excluding 3′UTR) were amplified from the pDONR207 clones and reconstituted with the pE-SUMO vector by Gibson Assembly to generate the heterologous expression plasmids pE-SUMO-BtKAI2a, pE-SUMO-BtKAI2b and pE-SUMO-BtKAI2c. Site-directed mutagenesis was performed on pE-SUMO-BtKAI2a and pE-SUMO-BtKAI2b clones to generate the heterologous expression plasmids pE-SUMO-BtKAI2a(b) and pE-SUMO-BtKAI2b(a). Both targeted residues were mutated simultaneously in one PCR product, while the remainder of the plasmid was amplified in a second PCR product. The mutated plasmids were reconstituted by Gibson assembly, and coding regions confirmed by Sanger sequencing.

Plant transformation

Homozygous *kai2-2* plants were transformed by floral dip. Primary transgenic seedlings were selected on sterile 0.5× MS medium supplemented with 20 μg/mL hygromycin B. T_2 lines exhibiting a 3:1 ratio of hygromycin resistant-to-sensitive seedlings were propagated further to identify homozygous lines in the T_3 generation. Experiments were performed from the T_3 generation onwards.

For transient expression in tobacco, *Agrobacterium* (GV3101) carrying pSKI106 variants was grown in LB medium (25 mL) supplemented with antibiotics and 20 μM acetosyringone until OD_{600} reached 1.0. The bacteria were then harvested by centrifugation (15 min, $5000 \times g$) and resuspended in 10 mM $MgCl_2$, 10 mM MES (pH 5.6) and 100 μM acetosyringone. The optical density was adjusted to 0.4, and the suspension was left standing at 22 °C overnight (approximately 14 h). Leaves of three-week-old *Nicotiana bethamiana* were then infiltrated with a 5-mL syringe, through the abaxial leaf surface. After four days, the leaves were collected and frozen in liquid nitrogen.

 Karrikin uptake measurements

Fifteen samples of seed were sowed for each karrikin treatment (five time points, each in
triplicate). In each sample, approximately 40 mg of *Brassica tournefortii* seeds were imbibed
in 3 mL ultrapure water for 24 h in 5-mL tubes. After centrifugation (2 min at 3220 × *g*),
excess water was removed by pipette and the volume of residual water (mostly absorbed into
the seeds) was calculated by weighing the seeds before and after imbibition. Fresh ultrapure
water was added to the fully-imbibed seeds to reach a total volume of 980 μL. Then 20 μL of
100 μM KAR_1 or KAR_2 was added to a final concentration of 2 μM. The seeds were imbibed at
22 °C in darkness. At the indicated time point (0, 2, 4, 8 or 24 h post-treatment), 500 μL of the
imbibition solution was removed and combined with 100 ng of either $^{13}[C]_5$-KAR_1 or $^{13}[C]_5$-
KAR_2 (100 μL at 1 μg/mL) as an internal standard for quantification purposes. The sample
was then extracted once with ethyl acetate (500 μL), and 1 μL of this organic layer was
analysed using GC-MS in selective ion monitoring (SIM) mode as previously described[50].

The amount of KAR_1 in each sample was calculated by the formula $\frac{A(Ion150)}{A(Ion155)}\times100$ ng and
converted to moles, where *A(Ion150)* indicates the peak area of the ion 150 (KAR_1 to be
measured), *A(Ion151)* indicates the peak area of the ion 151 ($^{13}[C]_5$-KAR_1), and 100 ng is the
amount of $^{13}[C]_5$-KAR_1 spiked in before the ethyl acetate extraction. Similarly, the amount of
KAR_2 in each sample was calculated by the formula $\frac{A(Ion136)}{A(Ion141)}\times100$ ng and converted to moles.

The uptake percentage adjusted to 40 mg of *B. tournefortii* seeds was calculated by the
formula: $(1 - \frac{N(x)}{N(0)}) \times \frac{40\ mg}{m\ (seed)}$, where N(0) indicates moles of karrikins at time 0, *N*(*x*) indicates
moles of karrikins at time point *x*, and *m*(*seed*) indicates the dry weight (mg) of seeds tested in
each replicate.

For Arabidopsis seeds, the procedure was scaled down for a smaller mass of seeds (20 mg).

Transcriptome assembly and analysis

Twenty milligrams of dry *B. tournefortii* seed was imbibed in water for 24 h and incubated at
22 °C in the dark. The seeds were collected by centrifugation, blotted dry, and frozen in liquid
nitrogen. A separate sample of seed was sown on glass filter paper, imbibed for 24 h as above,
and then incubated for 96 h under continuous red light (20 μmol m^{-2} s^{-1}) with a 22 °C (16
h)/16 °C (8h) temperature cycle. A single sample of seedlings (50 mg fresh weight) was

harvested and frozen in liquid nitrogen. Total RNA was extracted from both seed and seedling samples using the Spectrum Plant RNA kit (Sigma-Aldrich), including an on-column DNase step. PolyA$^+$ mRNA was purified using oligo(dT) magnetic beads (Illumina), and cDNA libraries for sequencing were generated as described[51]. Sequencing was performed on the Illumina HiSeq 2000 platform at the Beijing Genomic Institute, Shenzhen, China. Raw reads were filtered to remove adapters and low quality reads (those with >5% unknown nucleotides, or those in which greater than 20% of base calls had quality scores ≤10). After filtering, both libraries generated reads with >99.9% of nucleotides attaining Q20 quality score. Transcriptome *de novo* assembly was performed with Trinity[52]. For each library, contigs were assembled into Unigenes; Unigenes from both libraries were then combined, yielding a total of 45,553 predicted coding region sequences with a mean length of 1011 nt. The combined Unigenes were then interrogated for homology to AtKAI2 and AtD14 using BLASTn searches.

Phylogenetic analysis

KAI2 and *D14* homologues in *Brassica* species were identified from BLAST searches using Arabidopsis coding sequences as a query. Additional sequences were sampled from an extended phylogenetic analysis[20]. Multiple sequence alignments were performed using MAFFT plugin implemented in Geneious R10 (Biomatters). The alignment was trimmed slightly at the 5′ end to remove non-aligned regions of monocot D14 sequences. Maximum likelihood phylogenies were generated using PHYML (GTR +G +I substitution model, combination of NNI and SPR search, and 100 bootstraps). The choice of substitution model was guided by Smart Model Selection in PhyML (http://www.atgc-montpellier.fr/sms; [53]). A list of all sequences, and their sources, is provided in Supplementary Table 1.

Protein homology modelling

KAI2 structures were modelled using the SWISS-MODEL server (https://swissmodel.expasy.org) using the alignment mode[54] and the *Arabidopsis thaliana* KAI2 structure 3w06 as a template[36]. Figures of protein structure and homology models were generated using PyMOL v1.3 (Schrödinger LLC). Cavity surfaces were visualised using the "cavities & pockets (culled)" setting in PyMOL and a cavity detection cut-off value of four solvent radii. Cavity volumes were calculated using the CASTp server v3.0[55] (http://sts.bioe.uic.edu/castp) with a probe radius of 1.4Å. Values indicate Connolly's solvent-excluded volumes. Cavities were inspected visually using Chimera v1.12

360 (https://www.cgl.ucsf.edu/chimera/). For both BtKAI2a and BtKAI2b models, CASTp

361 erroneously included surface residues near the primary pocket entrance in the calculation of

362 the pocket volumes. This issue was resolved by the artificial placement of a free alanine

363 residue adjacent to the cavity entrance, as described previously[56].

Protein expression and purification

365 BtKAI2 proteins were generated as N-terminal 6×HIS-SUMO fusion proteins. All proteins were

366 expressed in BL21 Rosetta DE3 pLysS cells (Novagen) and purified using IMAC as described

367 previously[28].

Differential scanning fluorimetry

369 DSF was performed and thermal shifts were quantified as described previously[28]. Reactions

370 (10 μL) contained 20 μM protein, 20 mM HEPES pH 7.5, 150 mM NaCl, 1.25% (v/v) glycerol,

371 5× SYPRO Tangerine dye (Molecular Probes) and varying concentrations of ligand that

372 resulted in a final concentration of 5% (v/v) acetone.

Statistical analysis

374 Data were analysed using one- or two-way ANOVA ($\alpha = 0.05$, with Holm-Sidak's multiple

375 comparisons test). Germination data were arcsin transformed prior to ANOVA. Tests were

376 implemented in GraphPad Prism version 7.0 (GraphPad Software, graphpad.com).

Sequence data

378 Sequence data are available at NCBI Genbank under the following accessions: *BtKAI2a*,

379 MG783328; *BtKAI2b*, MG783329; *BtKAI2c*, MG783330; *BtD14a*, MG783331; *BtD14b*,

380 MG783332; *BtDLK2*, MG783333. Raw RNA sequence data from *Brassica tournefortii* seed and

381 seedlings are available in the NCBI SRA database under accession SRP128835.

REFERENCES

1. Flematti, G. R., Ghisalberti, E. L., Dixon, K. W. & Trengove, R. D. A compound from smoke
that promotes seed germination. *Science* **305,** 977 (2004).
2. Nelson, D. C., Flematti, G. R., Ghisalberti, E. L., Dixon, K. W. & Smith, S. M. Regulation of
seed germination and seedling growth by chemical signals from burning vegetation.
Ann Rev Plant Biol **63,** 107–130 (2012).
3. Nelson, D. C. *et al.* Karrikins discovered in smoke trigger Arabidopsis seed germination

by a mechanism requiring gibberellic acid synthesis and light. *Plant Physiol* **149,** 863–
873 (2009).

4. Waters, M. T. *et al.* Specialisation within the DWARF14 protein family confers distinct
responses to karrikins and strigolactones in Arabidopsis. *Development* **139,** 1285–1295
(2012).

5. Guo, Y., Zheng, Z., La Clair, J. J., Chory, J. & Noel, J. P. Smoke-derived karrikin perception
by the α/β-hydrolase KAI2 from Arabidopsis. *Proc Natl Acad Sci USA* **110,** 8284–8289
(2013).

6. Flematti, G. R. *et al.* Karrikin and Cyanohydrin Smoke Signals Provide Clues to New
Endogenous Plant Signaling Compounds. *Mol Plant* **6,** 29–37 (2013).

7. Keeley, J. E. & Pausas, J. G. Evolution of 'smoke' induced seed germination in
pyroendemic plants. *South African Journal of Botany* **115,** 251–255 (2018).

8. Flematti, G. R., Ghisalberti, E. L., Dixon, K. W. & Trengove, R. D. Identification of Alkyl
Substituted 2 H-Furo[2,3- c]pyran-2-ones as Germination Stimulants Present in Smoke.
J Agric Food Chem **57,** 9475–9480 (2009).

9. Flematti, G. R. *et al.* Preparation of 2H-furo[2,3-c]pyran-2-one derivatives and
evaluation of their germination-promoting activity. *J Agric Food Chem* **55,** 2189–2194
(2007).

10. Barrows, C. W., Allen, E. B., Brooks, M. L. & Allen, M. F. Effects of an invasive plant on a
desert sand dune landscape. *Biol Invasions* **11,** 673–686 (2009).

11. Bangle, D. N., Walker, L. R. & Powell, E. A. Seed germination of the invasive plant
Brassica tournefortii (Sahara mustard) in the Mojave Desert. *Western North American*
Naturalist **68,** 334–342 (2008).

12. Long, R. L. *et al.* Prior hydration of Brassica tournefortii seeds reduces the stimulatory
effect of karrikinolide on germination and increases seed sensitivity to abscisic acid.
Ann Bot **105,** 1063–1070 (2010).

13. Long, R. L. *et al.* Seeds of Brassicaceae weeds have an inherent or inducible response to
the germination stimulant karrikinolide. *Ann Bot* **108,** 933–944 (2011).

14. Brooks, M. L. *et al.* Effects of Invasive Alien Plants on Fire Regimes. *BioScience* **54,** 677
(2004).

15. Sanders, R. & Minnich, R. in *Invasive Plants of California's Wildlands* (eds. Bossard, C. C.,
Randall, J. M. & Hoshovsky, M. C.) 360 (University of California Press, 2000). at
<http://www.cal-ipc.org/resources/library/booksandcds/ipcw/online/>

16. Stevens, J., Merritt, D., Flematti, G., Ghisalberti, E. & Dixon, K. Seed germination of
agricultural weeds is promoted by the butenolide 3-methyl-2H-furo[2,3-c]pyran-2-one
under laboratory and field conditions. *Plant Soil* **298,** 113–124 (2007).

17. Schiermeier, Q. Pall hangs over desert's future as alien weeds fuel wildfires. *Nature*
435, 724 (2005).

18. Steers, R. J. Invasive Plants, Fire Succession, and Restoration of Creosote Bush Scrub in
Southern California. 1–210 (UNIVERSITY OF CALIFORNIA RIVERSIDE, 2008).

19. Waters, M. T., Scaffidi, A., Flematti, G. & Smith, S. M. Substrate-Induced Degradation of
the α/β-Fold Hydrolase KARRIKIN INSENSITIVE2 Requires a Functional Catalytic Triad
but Is Independent of MAX2. *Mol Plant* **8,** 814–817 (2015).

20. Bythell-Douglas, R. *et al.* Evolution of strigolactone receptors by gradual neo-
functionalization of KAI2 paralogues. *BMC Biol* **15,** 52 (2017).

21. Conn, C. E. & Nelson, D. C. Evidence that KARRIKIN-INSENSITIVE2 (KAI2) Receptors
may Perceive an Unknown Signal that is not Karrikin or Strigolactone. *Frontiers in Plant*
Science **6,** 1219 (2015).

22. Sun, Y. K., Flematti, G. R., Smith, S. M. & Waters, M. T. Reporter Gene-Facilitated
Detection of Compounds in Arabidopsis Leaf Extracts that Activate the Karrikin
Signaling Pathway. *Frontiers in Plant Science* **7,** 1799 (2016).

23. Li, W. *et al.* The karrikin receptor KAI2 promotes drought resistance in Arabidopsis
thaliana. *PLoS Genet* **13,** e1007076 (2017).
24. Martin A Lysak, M. A. K. A. P. I. S. Chromosome triplication found across the tribe
Brassiceae. *Genome Res.* **15,** 516–525 (2005).
25. Wang, X. *et al.* The genome of the mesopolyploid crop species Brassica rapa. *Nat Genet*
43, 1035–1039 (2011).
26. Liu, S. *et al.* The Brassica oleracea genome reveals the asymmetrical evolution of
polyploid genomes. *Nat Comms* **5,** 3930 (2014).
27. Sun, X.-D. & Ni, M. HYPOSENSITIVE TO LIGHT, an Alpha/Beta Fold Protein, Acts
Downstream of ELONGATED HYPOCOTYL 5 to Regulate Seedling De-Etiolation. *Mol*
Plant **4,** 116–126 (2011).
28. Waters, M. T. *et al.* A Selaginella moellendorffii Ortholog of KARRIKIN INSENSITIVE2
Functions in Arabidopsis Development but Cannot Mediate Responses to Karrikins or
Strigolactones. *Plant Cell* **27,** 1925–1944 (2015).
29. Hamiaux, C. *et al.* DAD2 Is an α/β Hydrolase Likely to Be Involved in the Perception of
the Plant Branching Hormone, Strigolactone. *Curr Biol* **22,** 2032–2036 (2012).
30. Abe, S. *et al.* Carlactone is converted to carlactonoic acid by MAX1 in Arabidopsis and its
methyl ester can directly interact with AtD14 in vitro. *Proc Natl Acad Sci USA* **111,**
18084–18089 (2014).
31. Végh, A. *et al.* Comprehensive Analysis of DWARF14-LIKE2 (DLK2) Reveals Its
Functional Divergence from Strigolactone-Related Paralogs. *Frontiers in Plant Science* **8,**
(2017).
32. Hamiaux, C. *et al.* Inhibition of strigolactone receptors by N-phenylanthranilic acid
derivatives: structural and functional insights. *Journal of Biological Chemistry* (2018).
doi:10.1074/jbc.RA117.001154
33. Scaffidi, A. *et al.* Strigolactone hormones and their stereoisomers signal through two
related receptor proteins to induce different physiological responses in Arabidopsis.
Plant Physiol **165,** 1221–1232 (2014).
34. Yao, R. *et al.* DWARF14 is a non-canonical hormone receptor for strigolactone. *Nature*
536, 469–473 (2016).
35. Xu, Y. *et al.* Structural basis of unique ligand specificity of KAI2-like protein from
parasitic weed Striga hermonthica. *Sci. Rep.* **6,** 31386 (2016).
36. Kagiyama, M. *et al.* Structures of D14 and D14L in the strigolactone and karrikin
signaling pathways. *Genes Cells* **18,** 147–160 (2013).
37. Lee, I. *et al.* Functional Analysis of a Missense Allele of KARRIKIN-INSENSITIVE2 that
Impairs Ligand-Binding and Downstream Signaling in Arabidopsis thaliana. *Journal of*
Experimental Botany (2018). doi:10.1093/jxb/ery164
38. Conn, C. E. *et al.* Convergent evolution of strigolactone perception enabled host
detection in parasitic plants. *Science* **349,** 540–543 (2015).
39. Toh, S. *et al.* Structure-function analysis identifies highly sensitive strigolactone
receptors in Striga. *Science* **350,** 203–207 (2015).
40. Shimada, A. *et al.* Structural basis for gibberellin recognition by its receptor GID1.
Nature **456,** 520 (2008).
41. Panchy, N., Lehti-Shiu, M. D. & Shiu, S.-H. Evolution of gene duplication in plants. *Plant*
Physiol pp.00523.2016 (2016). doi:10.1104/pp.16.00523
42. Wang, L., Waters, M. T. & Smith, S. M. Karrikin-KAI2 signalling provides Arabidopsis
seeds with tolerance to abiotic stress and inhibits germination under conditions
unfavourable for seedling establishment. *New Phytol* (2018). doi:10.1111/nph.15192
43. Goddard-Borger, E. D., Ghisalberti, E. L. & Stick, R. V. Synthesis of the germination
stimulant 3-methyl-2H-furo[2,3-c]pyran-2-one and analogous compounds from
carbohydrates. *Eur J Org Chem* 3925–3934 (2007). doi:10.1002/ejoc.200700334

44.   Mangnus, E., Dommerholt, F., Jong, R. de & Zwanenburg, B. Improved synthesis of strigol
492   analog GR24 and evaluation of the biological-activity of its diastereomers. *J Agric Food*
493   *Chem* **40,** 1230–1235 (1992).
494   45.   Scaffidi, A. *et al.* The synthesis and biological evaluation of labelled karrikinolides for
495   the elucidation of the mode of action of the seed germination stimulant. *Tetrahedron*
496   **67,** 152–157 (2011).
497   46.   Gorecki, M. J., Long, R. L., Flematti, G. R. & Stevens, J. C. Parental environment changes
498   the dormancy state and karrikinolide response of Brassica tournefortii seeds. *Ann Bot*
499   (2012). doi:10.1093/aob/mcs067
500   47.   Waters, M. T. & Smith, S. M. KAI2- and MAX2-Mediated Responses to Karrikins and
501   Strigolactones Are Largely Independent of HY5 in Arabidopsis Seedlings. *Mol Plant* **6,**
502   63–75 (2013).
503   48.   Curtis, M. D. & Grossniklaus, U. A gateway cloning vector set for high-throughput
504   functional analysis of genes in planta. *Plant Physiol* **133,** 462–469 (2003).
505   49.   Kagale, S. *et al.* TMV-Gate vectors: gateway compatible tobacco mosaic virus based
506   expression vectors for functional analysis of proteins. *Sci. Rep.* **2,** 874 (2012).
507   50.   Flematti, G. R., Scaffidi, A., Dixon, K. W., Smith, S. M. & Ghisalberti, E. L. Production of the
508   seed germination stimulant karrikinolide from combustion of simple carbohydrates. *J*
509   *Agric Food Chem* **59,** 1195–1198 (2011).
510   51.   Liang, C., Liu, X., Yiu, S.-M. & Lim, B. L. De novo assembly and characterization of
511   Camelina sativa transcriptome by paired-end sequencing. *BMC Genomics 2013 14:1* **14,**
512   146 (2013).
513   52.   Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data without a
514   reference genome. *Nat Biotechnol* **29,** 644–652 (2011).
515   53.   Lefort, V., Longueville, J.-E. & Gascuel, O. SMS: Smart Model Selection in PhyML. *Mol Biol*
516   *Evol* **34,** 2422–2424 (2017).
517   54.   Bordoli, L. *et al.* Protein structure homology modeling using SWISS-MODEL workspace.
518   *Nat Protoc* **4,** 1–13 (2009).
519   55.   Dundas, J. *et al.* CASTp: computed atlas of surface topography of proteins with
520   structural and topographical mapping of functionally annotated residues. *Nucleic Acids*
521   *Res* **34,** W116–W118 (2006).
522   56.   Lopez-Obando, M. *et al.* Structural modelling and transcriptional responses highlight a
523   clade of PpKAI2-LIKE genes as candidate receptors for strigolactones in Physcomitrella
524   patens. *Planta* **243,** 1441–1453 (2016).
525

ACKNOWLEDGEMENTS

This work was supported by funding from the Australian Research Council (DP130103646 to SMS and GRF; DP140104567 to SMS; FT150100162 to MTW, DP160102888 to GRF and MTW). Y.K.S was recipient of a Research Training Program Scholarship from the Australian Government. We thank Dr Rowena Long for the original provision of *B. tournefortii* seeds, Dr Rohan Bythell-Douglas for advice on homology modelling, and Mr Yongjie Meng with assistance with RNA extractions. We are grateful to Dr Kevin Rozwadowski (Agriculture and Agri-Food Canada) for the provision of pSKI106.

CONTRIBUTIONS

Y.K.S., S.M.S., G.R.F. and M.T.W. conceived and designed the research; Y.K.S., A.S., J.Y., K.M. and M.T.W. performed the experiments; Y.K.S. and M.T.W. analysed the data; Y.K.S., S.M.S. and M.T.W. wrote the manuscript.

SUPPLEMENTARY INFORMATION

Supplementary Figure 1 Germination of *Brassica tournefortii* seed treated with karrikins

Supplementary Figure 2 Extended phylogeny of eu-KAI2 and D14 proteins in angiosperms

Supplementary Figure 3 Alignment of Brassica KAI2 sequences

Supplementary Figure 4 BtD14a is functionally homologous to AtD14

Supplementary Figure 5 Expression of BtKAI2 homologues in Arabidopsis and tobacco

Supplementary Figure 6 Germination profiles of transgenic Arabidopsis seeds expressing BtKAI2 homologues

Supplementary Figure 7 BtKAI2a and BtKAI2b do not respond to karrikins in DSF assays

Supplementary Table 1 List of sequences identified in this study

Supplementary Table 2 List of oligonucleotides

FIGURE LEGENDS

Figure 1. *Brassica tournefortii* is highly sensitive to KAR$_1$, the major karrikin analogue isolated from smoke

a, Germination responses of *B. tournefortii* seed to KAR$_1$ and KAR$_2$. Data are means ± SE; n = 3 biological replicates per treatment, ≥35 seeds per replicate. **b**, Levels of karrikin-responsive transcripts *BtDLK2* and *BtSTH7* in *B. tournefortii* seed. Seed were imbibed for 24 hours in the dark supplemented with KAR$_1$ or KAR$_2$. Transcripts were normalised to *BtCACS* reference transcripts. Data are means ± SE, n = 3 biological replicates, ≥50 seeds per replicate. **c**, Degradation of BtKAI2 protein levels in *B. tournefortii* seeds treated with KAR$_1$ or KAR$_2$ for 24,

48, and 72 hours. For each sample, 30 μg total protein isolated from imbibed seeds was

separated by SDS-PAGE, blotted, and challenged with anti-KAI2 antibody. The reference panel

shows the blot stained by amido black. Samples from *A. thaliana* ecotype L*er* and *kai2-2*

seedlings serve as controls for antibody specificity. **d**, Hypocotyl elongation responses of *B.*

tournefortii seedlings treated with KAR_1 or KAR_2. Data are means ± SE of *n* = 3 biological

replicates, ≥12 seedlings per replicate. Scale bar: 10 mm.

Figure 2. Two differentially expressed *B. tournefortii KAI2* homologues are functional in

Arabidopsis

a, Maximum likelihood phylogeny of KAI2 homologues in the Brassicaceae, based on

nucleotide data. Node values represent bootstrap support from 100 replicates. A eu-KAI2

sequence from *Gossypium raimondii* (Malvaceae) serves as an outgroup. Tree shown is a

subset of a larger phylogeny in Supplementary Fig. 2. **b**, Transcript levels of the three *BtKAI2*

homologues in seed imbibed for 24 h (left) and four-day-old seedlings (right) treated with or

without 1 μM KAR_1 for 24 h. **c**, Seedling and rosette phenotypes of two independent

transgenic lines of Arabidopsis homozygous for *KAI2pro:GFP-BtKAI2x* transgenes. Scale bars:

5 mm (upper); 50 mm (lower). **d**, Immunoblots of soluble proteins challenged with antibodies

against KAI2 (upper panel), GFP (middle panel) or actin (lower panel). Non-specific bands are

marked with asterisks. Protein was isolated from pools of approximately fifty 7-day-old

seedlings.

Figure 3. Functional divergence between BtKAI2 homologues

a, Germination responses of primary dormant *Arabidopsis* seed homozygous for *KAI2pro:GFP-*

BtKAI2x transgenes in the *kai2-2* background. Germination values were determined 120 h

after sowing. Extended germination curves are shown in Supplementary Fig. 6. Data are

means ± SE, *n* = 3 independent seed batches, 75 seed per batch. **b**, Hypocotyl elongation

responses of *KAI2pro:GFP-BtKAI2x* seedlings treated with KAR_1 or KAR_2. Data are means ± SE

of *n* = 3 biological replicates, 12–18 seedlings per replicate. **c**, Levels of *DLK2* transcripts in 8-

day-old *KAI2pro:GFP-BtKAI2x* seedlings treated with KAR_1 or KAR_2 for eight hours. Expression

was normalised to *CACS* reference transcripts and scaled to the value for mock-treated

seedlings within each genotype. Data are means ± SE of *n* = 3 biological replicates. Pairwise

significant differences: * $P < 0.05$ ** $P < 0.01$ *** $P < 0.001$; ns, $P > 0.05$ (ANOVA).

Figure 4. Two residues account for ligand specificity between BtKAI2a and BtKAI2b

a, DSF curves of SUMO-BtKAI2a and SUMO-BtKAI2b fusion proteins treated with 0-50 μM GR24$^{ent-5DS}$, a KAI2-bioactive ligand. Each curve is the average of three sets of reactions, each comprising four technical replicates. Insets plot the minimum value of –(dF/dT) at the melting point of the protein as determined in the absence of ligand (means ± SE, n = 3). Significant differences from untreated control: * $P < 0.05$ ** $P < 0.01$ *** $P < 0.001$ (ANOVA). **b–d**, Solved structure of AtKAI2 (PDB: 3w06 [36]) (**b**) and derived homology models (**c–d**) of BtKAI2a and BtKAI2b. Coloured surfaces depict internal cavities; values indicate the volumes of the primary ligand-binding cavities, adjacent to the catalytic Ser-His-Asp residues. Also shown is a variable secondary pocket, to the left of the primary pocket in these images. **e**, DSF curves of SUMO-BtKAI2a(b) and SUMO-BtKAI2b(a) fusion proteins treated with 0-50 μM GR24$^{ent-5DS}$. **f**, Levels of *DLK2* transcripts (normalized to *CACS* reference transcripts) in two independent lines of *KAI2pro:GFP-BtKAI2a* and *-BtKAI2b* seedlings treated for two hours with 0.1 μM GR24$^{ent-5DS}$. Data are means ± SE; n = 3 biological replicates. Letters indicate significantly different groupings (ANOVA; $P < 0.05$).

SUPPLEMENTARY FIGURE LEGENDS

Supplementary Figure 1. Germination of *Brassica tournefortii* seed treated with karrikins

a, Structures of KAR$_1$, KAR$_2$ and two enantiomers of GR24. **b**, Germination response of the "Merridin" batch of *B. tournefortii* seed treated with KAR$_1$ and KAR$_2$ after three days. **c-d**, Germination response of the "Perth" batch to KAR$_1$ (**c**) and KAR$_2$ (**d**). Data in Figure 1a are derived from the data shown in **c**. **e-f**, Rates of KAR$_1$ and KAR$_2$ uptake by imbibed seed, as determined by GC-MS.

Supplementary Figure 2. Extended phylogeny of eu-KAI2 and D14 proteins in angiosperms

Maximum likelihood phylogeny of KAI2 and D14 homologues in the Brassicaceae and monocots, based on nucleotide data. Node values represent bootstrap support from 100 replicates. A KAI2 sequence from *Selaginella moellendorffii* (SmKAI2a; Waters et al. 2015) serves as an outgroup for the eu-KAI2 clade.

Supplementary Figure 3. Alignment of Brassica KAI2 sequences

Full length protein coding regions of *KAI2* homologues from four *Brassica* species (*Brassica tournefortii*, *B. rapa*, *B. nigra* and *B. oleracea*) were translated from database nucleic acid

620 sequences and aligned to Arabidopsis *KAI2* using MAFFT (Katoh et al, 2002) implemented in

621 Geneious R10 software (Biomatters Ltd). Residues that are unique to BtKAI2c but otherwise

622 invariant are highlighted with asterisks (*); residues 98 and 191 are highlighted with red

623 boxes.

Supplementary Figure 4. BtD14a is functionally homologous to AtD14

625 Three independent transgenic lines of Arabidopsis *Atd14-1* were analysed for functional

626 complementation of the mutant phenotype by an *AtD14pro:BtD14a* transgene. **a**, Rosette and

627 leaf morphology at 31 days post-germination. **b**, Plant height and number of primary rosette

628 branches at 45 days post-germination. **c**, Quantification of height and branching parameters, *n*

629 = 10 plants per genotype. Scale bars: 50 mm.

Supplementary Figure 5. Expression of BtKAI2 homologues in Arabidopsis and tobacco

a, Alignment of the protein regions corresponding to the peptide epitope TTNPDYFDFDRYSN

of AtKAI2, against which the anti-KAI2 peptide was raised. BtKAI2b, with a more divergent

epitope, is relatively poorly detected by the anti-KAI2 antibody (see Figure 2d). **b**, Transient

expression of BtKAI2a, BtKAI2b and BtKAI2c proteins in tobacco. Plasmids encoding N-

terminal, *c*-myc-tagged proteins were transferred to *Agrobacterium*, and the resulting strains

used to infiltrate tobacco leaves. After 96 h, samples were harvested in triplicate (two to three

leaves per sample). Mock-treated leaves were transformed with a plasmid encoding a non-

tagged protein. Sixty micrograms of total protein were separated by SDS-PAGE, blotted and

challenged with anti-*c*-myc antibody (Genscript A00704). Band intensity was measured using

ImageJ, and expression was normalised to intensity of the Rubisco large subunit (RbcL) band

on the "stain-free" gel imaged under UV light. Error bars indicate SE, *n* = 3 replicates. **c**,

Quantitative RT-PCR analysis of *GFP* and *DLK2* transcripts, normalised to *CACS* reference

transcripts. Data are derived from a single biological replicate with four technical PCR

replicates. RNA was isolated from approximately 50 seven-day-old seedlings per genotype,

and are the same samples as those shown in the immunoblots in Figure 2d.

Supplementary Figure 6. Germination profiles of transgenic Arabidopsis seeds
expressing BtKAI2 homologues

Freshly harvested seed (three batches per genotype, each batch harvested from four plants)

were removed from freezer storage, surface-sterilised and sown on 1% Phytagel

supplemented with 0.1% acetone (mock), 1 μM KAR_1 or 1 μM KAR_2. Seed were incubated

under constant light at 25 °C. Seed were examined for germination (radicle protrusion) 72 h

after sowing and every 24 h thereafter. Data are means ± SE of three independent seed

batches and 75 seed per batch. For each transgene, two independent, homozygous transgenic

lines were analysed. Data presented in Figure 3 of the main manuscript are derived from

these data.

Supplementary Figure 7. BtKAI2a and BtKAI2b do not respond to karrikins in DSF

assays

Differential scanning fluorimetry curves of BtKAI2a and BtKAI2b in presence of 0–200 μM

KAR_1 or KAR_2 (**a**) or the two enantiomers of GR24 (**b**). Data are means of eight technical

replicates at each concentration of ligand.

Supplementary Figure 1. Germination of *Brassica tournefortii* seed treated with karrikins
a, Structures of KAR$_1$, KAR$_2$ and two enantiomers of GR24. **b,** Germination response of the "Merridin" batch of *B. tournefortii* seed treated with KAR$_1$ and KAR$_2$ after three days. **c-d,** Germination response of the "Perth" batch to KAR$_1$ (**c**) and KAR$_2$ (**d**). Data in Figure 1a are derived from the data shown in **c. e-f,** Rates of KAR$_1$ and KAR$_2$ uptake by imbibed seed, as determined by GC-MS.

Supplementary Figure 2. Extended phylogeny of eu-KAI2 and D14 proteins in angiosperms
Maximum likelihood phylogeny of KAI2 and D14 homologues in the Brassicaceae and monocots, based on nucleotide data. Node values represent bootstrap support from 100 replicates. A KAI2 sequence from *Selaginella moellendorffii* (SmKAI2a; Waters et al. 2015) serves as an outgroup for the eu-KAI2 clade.

Supplementary Figure 3. Alignment of *Brassica* KAI2 sequences

Full length protein coding regions of KAI2 homologues from four *Brassica* species (*Brassica tournefortii*, *B. rapa*, *B. nigra* and *B. oleracea*) were translated from database nucleic acid sequences and aligned to Arabidopsis KAI2 using MAFFT (Katoh et al, 2002) implemented in Geneious R10 software (Biomatters Ltd). Residues that are unique to BtKAI2c but otherwise invariant are highlighted with asterisks (*); residues 98 and 191 are highlighted with red boxes.

Supplementary Figure 4. BtD14a is functionally homologous to AtD14

Three independent transgenic lines of Arabidopsis *Atd14-1* were analysed for functional complementation of the mutant phenotype by an *AtD14pro:BtD14a* transgene. **a**, Rosette and leaf morphology at 31 days post-germination. **b**, Plant height and number of primary rosette branches at 45 days post-germination. **c**, Quantification of height and branching parameters, *n* = 10 plants per genotype. Scale bars: 50 mm.

a

b

c

Supplementary Figure 5. Expression of BtKAI2 homologues in Arabidopsis and tobacco

a, Alignment of the protein regions corresponding to the peptide epitope TTNPDYFDFDRYSN of AtKAI2, against which the anti-KAI2 peptide was raised. BtKAI2b, with a more divergent epitope, is relatively poorly detected by the anti-KAI2 antibody (see Figure 2). **b**, Transient expression of BtKAI2a, BtKAI2b and BtKAI2c proteins in tobacco. Plasmids encoding N-terminal, c-myc-tagged proteins were transferred to Agrobacterium, and the resulting strains used to infiltrate tobacco leaves. After 96 h, samples were harvested in triplicate (two to three leaves per sample). Mock-treated leaves were transformed with a plasmid encoding a non-tagged protein. Sixty micrograms of total protein were separated by SDS-PAGE, blotted and challenged with anti-c-myc antibody (Genscript A00704). Band intensity was measured using ImageJ, and expression was normalised to intensity of the Rubisco large subunit (RbcL) band on the "stain-free" gel imaged under UV light. Error bars indicate SE, *n* = 3 replicates. **c**, Quantitative RT-PCR analysis of *GFP* and *DLK2* transcripts, normalised to *CACS* reference transcripts. Data are derived from a single biological replicate with four technical PCR replicates. RNA was isolated from approximately 50 seven-day-old seedlings per genotype, and are the same samples as those shown in the immunoblots in Figure 2.

Supplementary Figure 6. Germination profiles of transgenic Arabidopsis seeds expressing BtKAI2 homologues
Freshly harvested seed (three batches per genotype, each batch harvested from four plants) were removed from freezer storage, surface-sterilised and sown on 1% Phytagel supplemented with 0.1% acetone (mock), 1 µM KAR_1 or 1 µM KAR_2. Seed were incubated under constant light at 25 °C. Seed were examined for germination (radicle protrusion) 72 h after sowing and every 24 h thereafter. Data are means ± SE of three independent seed batches and 75 seed per batch. For each transgene, two independent, homozygous transgenic lines were analysed. Data presented in Figure 3 of the main manuscript are derived from these data.

Supplementary Figure 7. BtKAI2a and BtKAI2b do not respond to karrikins in DSF assays
Differential scanning fluorimetry curves of BtKAI2a and BtKAI2b in presence of 0–200 μM KAR_1 or KAR_2 (**a**) or the two enantiomers of GR24 (**b**). Data are means of eight technical replicates at each concentration of ligand.

Supplementary Table 1. List of sequences identified in this study

KAI2

Species	Gene name	Sequence ID	Source	Reference
Arabidopsis lyrata	*AlKAI2*	AL7G13320.t1	Phytozome	This work
Arabidopsis thaliana	*AtKAI2*	At3g37470	TAIR	Waters et al (2012)
Brassica tournefortii (saharan mustard)	*BtKAI2a*	MG783328	GenBank/NCBI	This work
Brassica tournefortii (saharan mustard)	*BtKAI2b*	MG783329	GenBank/NCBI	This work
Brassica tournefortii (saharan mustard)	*BtKAI2c*	MG783330	GenBank/NCBI	This work
Brassica rapa (field mustard, AA genome)	*BrKAI2a*	XM_009111126.2	GenBank/NCBI	This work
Brassica rapa (field mustard, AA genome)	*BrKAI2b*	XM_009140247	GenBank/NCBI	This work
Brassica rapa (field mustard, AA genome)	*BrKAI2c*	XM_009144693	GenBank/NCBI	This work
Brassica nigra (black mustard, BB genome)	*BnKAI2a*	LFLV01000699.1	GenBank/NCBI	This work
Brassica nigra (black mustard, BB genome)	*BnKAI2b*	LFLV01001772.1	GenBank/NCBI	This work
Brassica oleracea (cabbage, CC genome)	*BoKAI2a*	XM_013766720.1	Phytozome	This work
Brassica oleracea (cabbage, CC genome)	*BoKAI2b*	XM_013738288.1	Phytozome	This work
Brassica oleracea (cabbage, CC genome)	*BoKAI2c*	XM_013778816.1	Phytozome	This work
Boechera stricta	*BsKAI2*	Bostr.30440s0001.1	Phytozome	This work
Capsella rubella	*CrKAI2*	Carubv10005485m	Phytozome	This work
Capsella grandiflora	*CgKAI2*	Cagra.1232s0005.1	Phytozome	This work
Eutrema salsugineum	*EsKAI2*	Thalv10025969m	Phytozome	Bythell-Douglas et al (2017)
Gossypium raymondii	*GrKAI2*	Gorai.003G028500.1	Phytozome	Bythell-Douglas et al (2017)
Prunus persica	*PpKAI2*	Ppa009957m	Phytozome	Bythell-Douglas et al (2017)
Brachypodium distachyon	*BdKAI2*	Bradi1g15880.1	Phytozome	Bythell-Douglas et al (2017)
Oryza sativa	*OsKAI2*	Os03g32270.1	Phytozome	Bythell-Douglas et al (2017)
Sorghum bicolor	*SbKAI2*	Sorbic.001G330000.1	Phytozome	Bythell-Douglas et al (2017)
Selaginella moellendorffii	*SmKAI2a*	Selmo_441991	Phytozome	Waters et al (2015)

D14

Species	Gene name	Sequence ID	Source	Reference
Arabidopsis lyrata	*AlD14*	AL3G13900.t1	Phytozome	This work
Arabidopsis thaliana	*AtD14*	AT3G03990.1	TAIR	Waters et al (2012)
Brassica oleracea (cabbage, CC genome)	*BoD14a*	LOC106343216	Phytozome	This work
Brassica oleracea (cabbage, CC genome)	*BoD14b*	LOC106343689	Phytozome	This work
Brassica nigra (black mustard, BB genome)	*BnD14a*	LFLV01002090	GenBank/NCBI	This work
Brassica nigra (black mustard, BB genome)	*BnD14b*	LFLV01000699	GenBank/NCBI	This work
Brassica rapa (field mustard, AA genome)	*BrD14a*	Brara.A03790	Phytozome	This work
Brassica rapa (field mustard, AA genome)	*BrD14b*	Brara.E03476	Phytozome	This work
B. tournefortii (saharan mustard)	*BtD14a*	MG783331	Phytozome	This work
B. tournefortii (saharan mustard)	*BtD14b*	MG783332	Phytozome	This work
Boechera stricta	*BsD14*	Bostr.2570s0310	Phytozome	This work
Capsella grandiflora	*CgD14*	Cagra.15970s0001	Phytozome	This work
Capsella rubella	*CrD14*	Carubv10014401m	Phytozome	This work
Eutrema salsugineum	*EsD14*	Thhalv10021292m	Phytozome	Bythell-Douglas et al (2017)
Gossypium raymondii	*GrD14*	Gorai.010G025600.1	Phytozome	Bythell-Douglas et al (2017)
Prunus persica	*PpD14*	ppa010005m	Phytozome	Bythell-Douglas et al (2017)
Brachypodium distachyon	*BdD14*	Bradi1g70930.3	Phytozome	Bythell-Douglas et al (2017)
Oryza sativa	*OsD14*	Os03g10620.1	Phytozome	Bythell-Douglas et al (2017)
Sorghum bicolor	*SbD14*	Sobic.001G465100.1	Phytozome	Bythell-Douglas et al (2017)

DLK2

Species	Gene name	Sequence ID	Source	Reference
B. tournefortii (saharan mustard)	*BtDLK2*	MG783333	GenBank/NCBI	This work

Supplementary Table 2. Oligonucleotides used in this study

Oligonucleotide	Sequence	Notes
Cloning		
BtD14a_F	AAAAAAGCAGGCTTCCATTATGAGTCAACACAACAT	BtD14a
BtD14a_R	CAAGAAAGCTGGGTTTTAAAGTCACCGAGGAAG	BtD14a
BtKAI2_universal_F	GGGGACAAGTTTGTACAAAAAAGCAGGCTTCATGGGAGGTGTGGTAGAGGA	All BtKAI2
RACE_R	GGGGACCACTTTGTACAAGAAAGCTGGGTCATTCTCAATAGACAATAGAC	BtKAI2a
Contig1_R	GGGGACCACTTTGTACAAGAAAGCTGGGTCCGAAACGACCCACTCACTTACC	BtKAI2b
Contig5_R	GGGGACCACTTTGTACAAGAAAGCTGGGTCCGAAGCAGAACCAAGCTAA	BtKAI2c
BtKAI2-SUMO-Gib_F	TGAGGCTCACCGCGAACAGATTGGAGGTATGGGAGGTGTGGTAGAGGAAG	pSUMO-BtKAI2a / BtKAI2b
BtKAI2a-SUMO-Gib_R	CGGATCTCAGTGGTGGTGGTGGTGGTGCTCAGACAGCGATGTCATTACGA	pSUMO-BtKAI2a
BtKAI2b-SUMO-Gib_R	CGGATCTCAGTGGTGGTGGTGGTGGTGCTCATACAGCAATGTCGTTGCGG	pSUMO-BtKAI2b
BtKAI2b_L98V_F1	ATCTTTGTTGGTCACTCTGTCTCCGCCATG	For BtKAI2b(a) mutagenesis
BtKAI2b_L191V_R1	TTTGGAAAATGGTCTGAGCCACGGAGAGTG	For BtKAI2b(a) mutagenesis
BtKAI2b_L191V_F2	TCTCCGTGGCTCAGACCATTTTCCAAAGC	For BtKAI2b(a) mutagenesis
BtKAI2b_L98V_R2	CATGGCGGAGACAGAGTGACCAACAAAG	For BtKAI2b(a) mutagenesis
BtKAI2a_V98L_F1	ATCTTTGTTGGTCACTCTCTCTCCGCCATG	For BtKAI2a(b) mutagenesis
BtKAI2a_V191L_R1	TTTGGAAAATGGTCTGAGCCAGGGAGAGAG	For BtKAI2a(b) mutagenesis
BtKAI2a_V191L_F2	TCTCCCTGGCTCAGACCATTTTCCAAAG	For BtKAI2a(b) mutagenesis
BtKAI2a_V98L_R2	CATGGCGGAGAGAGAGTGACCAACAAAG	For BtKAI2a(b) mutagenesis
Quantitative PCR		
STH7_qF	CATCTCCGGTTCTCTCTCACTTCT	Designed against At4g39070
STH7_qR	CATTCTCTGCATAGTATTGCTCTGTC	Designed against At4g39070
BtDLK2_qF	GCTGCTTCTCCCAGGTATATAA	BtDLK2
BtDLK2_qR	GAAAGCAACCGCCCAAGCC	BtDLK2
AtDLK2_qF	GCTGCTTCTCCAAGGTATATAA	Designed against At3g24420
AtDLK2_qR	GAAATCAACCGCCCAAGCT	Designed against At3g24420
KUF1_qF	AACCCGTCAGTCCGGTTTATGTG	Designed against At1g31350
KUF1_qR	AACGACGGATGACGGTAAAGAATCC	Designed against At1g31350
IAA6_qF	TTCGGCTGTCTTGGCATAGGAG	Designed against At1g52830
IAA6_qR	TTCACGATCCTCAGCCTCTTGC	Designed against At1g52830
BtKAI2_universal_qF	GGTCATCTTCCTCAGCTTAG	All BtKAI2
BtKAI2_a_qR1	CCATTAATTTAATTTAAAATCACTTCCC	BtKAI2a
BtKAI2_b_qR1	GCATTTAATTTCGCTAAAACTCTTG	BtKAI2b
BtKAI2_c_qR1	GAAACGATATACTCAGTACTTACCAC	BtKAI2c
CACS_qF	ACTCAGGAAGGTGTACGGTCA	Designed against At5g46630
CACS_qR	TGCATTTGGAACAGGTTTGT	Designed against At5g46630