Transdiagnostic characterization of neuropsychiatric disorders by hyperexcitation-induced

immaturity (100 characters)

Tomoyuki Murano[1,2,3], Hideo Hagihara[1], Katsunori Tajinda[4], Mitsuyuki Matsumoto[5],

Tsuyoshi Miyakawa[1]*

[1]Division of Systems Medical Science, Institute for Comprehensive Medical Science,

Fujita Health University, Japan

[2]Department of Physiological Science, School of Life Science, The Graduate

University for Advanced Studies, Japan [SOKENDAI]

[3]Division of Cell Signaling, National Institute for Physiological Sciences, Japan

[4] Neuroscience, La Jolla Laboratory, Astellas Research Institute of America LLC,

USA

[5] Drug Discovery Research, Astellas Pharma Inc., Japan

*Correspondence: miyakawa@fujita-hu.ac.jp

**Abstract**

Biomarkers are needed to improve the diagnosis of neuropsychiatric disorders.

Promising candidates are imbalance of excitation and inhibition in the brain, and maturation

abnormalities. Here, we characterized different disease conditions by mapping changes in the

expression patterns of maturation-related genes whose expression was altered by experimental

neural hyperexcitation in published studies. This revealed two gene expression patterns:

decreases in maturity markers and increases in immaturity markers. These two groups of genes

were characterized by the overrepresentation of genes related to synaptic function and

chromosomal modification, respectively. We used these two groups in a transdiagnostic

analysis of 80 disease datasets for eight neuropsychiatric disorders and 12 datasets from

corresponding animal models, and found that transcriptomic pseudoimmaturity inducible by

neural hyperexcitation is shared by multiple neuropsychiatric disorders, such as schizophrenia,

Alzheimer disorders, and ALS. Our results suggest that this endophenotype may serve as a

basis for transdiagnostic characterization of these disorders.

(149/150 words)

## Introduction

Neuropsychiatric disorders—such as schizophrenia, bipolar disorder, major depression disorders, and autism spectrum disorder—are common, with over a third of the population in most countries being diagnosed with at least one such disorder at some point in their life[1]. Almost all neuropsychiatric disorders are currently classified mainly on the basis of clinical signs and symptoms. However, there is evidence that patients with different clinical diagnoses share similar biological features, such as genetic mutations, molecular expression, or brain activity[2–6]. Recently, psychiatry has undergone a tectonic shift to incorporate the concepts of modern biology. There have been recent attempts to reclassify psychiatric disorders according to biological domains (e.g. genes, neural circuits, behavior), such as through the Research Domain Criteria (RDoC) initiative[7]. Therefore, identifying appropriate biomarkers which can be used for transdiagnostic assessment of neuropsychiatric disorders is essential for better classification of these diseases and understanding of their biological basis.

Using coexpression network analysis, a recent study revealed that cross-disorder gene expression overlaps could be used to characterize five major neuropsychiatric disorders[8]. Some of these overlapping gene groups were biologically well-characterized by Gene Ontology enrichment or cell-type specificity, but the biological properties of other gene groups were rather unclear. Thus, non-biased coexpression network analyses do not necessarily detect modules that extract the biological features of neuropsychiatric disorders. Then, to achieve better characterization of neuropsychiatric disorders, it might be helpful to detect modules of coexpressed genes and conduct gene expression analysis based on the findings derived from researches on animal models of neuropsychiatric disorders.

To date, we have screened more than 180 strains of genetically engineered mice using a large-scale and comprehensive battery of behavioral tests, and identified several strains with abnormal behaviors related to neuropsychiatric disorders such as schizophrenia, bipolar disorder, and intellectual disability[9]. We discovered common endophenotypes in the brains of multiple strains of these genetically engineered mice with behavioral abnormalities. We termed one such endophenotype in the hippocampus of adult mice the "immature dentate gyrus (iDG)" phenotype[10–13]. In this phenotype, the molecular and electrophysiological properties of adult DG neurons in the genetically engineered mice were similar to those of immature DG neurons in typically developing infants. For example, the expression of calbindin, a marker of maturity in DG neurons, was decreased and the expression of calretinin, a marker of immaturity, was increased[10–15]. Similar molecular changes to some of those found in mice with iDG have been observed in the postmortem brains of patients with schizophrenia[16], bipolar disorder[16], and epilepsy[17–19]. Furthermore, there is growing evidence that changes in molecular markers of

pseudoimmaturity are also observed in other brain areas of patients with schizophrenia[20-28],

bipolar disorder[26], autism[26], and alcoholism[29]. Therefore, we proposed that pseudoimmaturity

of the brain could potentially be a useful transdiagnostic biomarker[9].

Pseudoimmaturity of the brain can be induced in adulthood. Previously, we found

that chronic fluoxetine treatment reverses the maturation status of DG neurons in adult

wild-type mice, a phenomenon that we termed "dematuration"[30,31]. Likewise, recent studies

suggest that several maturation-related genes and electrophysiological properties in the DG of

wild-type adult mice assume an immature-like status after treatment with pilocarpine or

electroconvulsive stimulation[16,32]. As mentioned above, an iDG-like phenotype has been found

in patients with epilepsy[17–19]. Therefore, we hypothesized that the hyperexcitation of neurons

may be a cause of pseudoimmaturity of the brain in adulthood.

Some studies suggest that hyperexcitation of neurons may underlie abnormalities

related to certain types of neuropsychiatric disorders. Individuals with epilepsy are at increased

risk of developing schizophrenia, and vice versa[33,34] , and patients with epilepsy can also

display psychotic symptoms that resemble those found in patients with schizophrenia[35].

Imbalances in excitatory and inhibitory brain circuitry have been proposed to be involved in

the pathogenesis and pathophysiology of schizophrenia[36–39]. Hyperactive action-potential

firing has also been observed in hippocampal granule-cell-like neurons derived from induced

pluripotent stem cells (iPSCs) of patients with bipolar disorder[40]. Recent studies suggested that

human patients with Alzheimer's disease and temporal lobe epilepsy may harbor common

underlying mechanisms[17,41–43]. Considering these findings, we hypothesized that the

immature-like gene expression patterns induced by neural hyperexcitation may overlap with

the abnormal gene expression patterns in the brains of patients with neuropsychiatric disorders

and the related animal models. If this is the case, we hypothesized that this overlap can be used

to perform transdiagnostic characterization of neuropsychiatric disorders.

To test this hypothesis, we first performed a meta-analysis of microarray datasets,

comparing the changes in gene expression in rat DG after seizure induction with the

differences in gene expression in infant mice versus adult mice. To assess consistency across

species, we also conducted a similar comparison with human fetal hippocampus. The overlap

between gene sets was estimated using the Running Fisher test[44], which is a nonparametric

rank-based statistical method developed by BSCE. This method enables us to statistically

assess the pairwise correlations between any two datasets, including datasets from different

species and organs[28,29,45,46]. The gene expression patterns in rat DG after seizure induction

significantly overlapped with those specific to immature mouse DG and also with those

specific to early-stage human fetal hippocampus. From the set of overlapping genes, we

defined two groups; maturity-marker genes and immaturity-marker genes that are inducible by

neural hyperexcitation. We assessed the expression patterns of these two groups of

maturation-related genes in 80 public gene-expression datasets derived from the postmortem

brains of patients with various neuropsychiatric disorders and from neural cells derived from

patient iPSCs, and in a further 12 datasets from the brains of related animal models. Through

this analysis, we characterized the expression patterns of maturation-related genes that are

inducible by neural hyperexcitation across different disease conditions.

(932 words)

**Results**

*Neural hyperexcitation induces immature-like gene expression patterns in DG*

117           To examine the developmental changes in gene expression patterns in the rodent DG,

we created a microarray dataset from postnatal day 8, 11, 14, 17, 21, 25, and 29 infant mice

(GSE113727) and compared it with a dataset from 33-week-old adult mice (GSE42778)[12].

Within the entire mice DG dataset, the largest overlap for changes in gene expression after

pilocarpine injection was for the comparison between day 8 infant and 33-week-old adult mice

(Figure S1a). We included the dataset from postnatal day 8 infant mice for subsequent analysis.

The expression levels of 6552 genes were increased in the DG of infant mice compared with

adult mice, whereas the expression levels of 8637 genes were decreased (absolute fold change

$> 1.2$ and t-test $P < 0.05$). Next, we assessed the changes in gene expression induced by neural

hyperexcitation in a rodent model. We obtained publicly available microarray datasets from the

DG of adult rats after seizures induced by injection of pilocarpine (GSE47752)[47]. The

expression levels of 7073 genes were significantly changed in the DG of epileptic-seizure rats

1 day after pilocarpine injection compared with rats treated with saline (absolute fold change $>$

1.2, $P < 0.05$).

131           To investigate whether the neuronal hyperexcitation datasets contain immature-like

gene expression patterns, we assessed the overlap between the set of genes with altered

expression in immature mice and the set of genes with altered expression in adult

seizure-model rats, using the Running Fisher algorithm on the BaseSpace platform to

determine the significance of the overlap (see Supplementary Methods for Details). We found a

striking degree of similarity: 2807 genes showed changes in expression in both datasets

(overlap $P = 3.8 \times 10^{-11}$) (Figure 1a). Among these 2807 genes, we named the 726 genes whose

expression levels decreased in both datasets "hyperexcitation-induced maturity-related genes"

(hiM genes (mouse): green bar in Figure 1a) and the 938 genes whose expression levels

increased in both datasets "hyperexcitation-induced immaturity-related genes" (hiI genes

(mouse): red bar in Figure 1a). The comprehensive gene lists of hiM and hiI genes are in Table

S1 (Table S1: hiM/hiI_GeneList). The overlap for genes with positively correlated expression

(red and green bars) was larger than the overlap for genes with negatively correlated expression

(light and dark yellow bars), indicating that the direction of expressional changes in the two

datasets are more alike than they are different. These results suggest that neuronal

hyperexcitation induces a pattern of immature-like gene expression in the adult DG.

147           Next, we compared the changes in expression during development in the human fetal

hippocampus with those in rats after seizure induction to assess consistency across species. We

obtained publicly available microarray datasets for the human fetal hippocampus during

development (GSE25219)[48]. Within the entire fetal hippocampal dataset, the largest overlap for

changes in gene expression after pilocarpine injection was for the comparison between

8–9-week fetuses and 19–23-week fetuses (Figure S1b). We again found a striking degree of

similarity: 2043 genes showed changes in expression in both datasets (overlap $P = 1.8 \times 10^{-12}$)

(Figure 1b). Among these 2043 genes, we termed the 579 genes whose expression decreased in

both datasets "hiM genes (human)" (green bar in Figure 1b) and the 716 genes whose

expression increased in both datasets "hiI genes (human)" (red bar in Figure 1b). The overlap

for genes with positively correlated expression (red and green bars) were larger than those with

negatively correlated expression (light and dark yellow bars), suggesting that, similar to the

results in mice, the gene expression changes in rat DG after seizure induction are comparable to

the reverse of the changes that occur as the human hippocampus develops.

***Hyperexcitation-induced maturity- and immaturity-related genes exhibit different biological***

***properties***

To characterize the biological features associated with the hiM and hiI gene groups in

mouse and human, we conducted pathway enrichment analyses in BaseSpace. The 20

biogroups with the most significant overlap with hiM and hiI genes are listed in Table 1a and

1b. Among mouse hiM genes, 4 out of the top 20 biogroups are associated with synapse and

channel activity (e.g., transmission of nerve impulse, synapse, and synaptic transmission)

(Table 1a), whereas among human hiM genes, 6 out of the top 20 biogroups are also associated

with synapse and channel activity (e.g., transmission of nerve impulse, synaptic transmission,

axon, and synapse) (Table 1b).

Among the mouse hiI genes, 4 out of the top 20 biogroups were associated with the

nucleus (e.g., genes involved in the cell cycle and genes involved in DNA replication) (Table

1a). Among the human hiI genes, 15 out of the top 20 biogroups were associated with the

nucleus (e.g., genes involved in the cell cycle, chromosomes, and response to DNA damage

stimulus) (Table 1b). It is noteworthy that there is little overlap in the top 20 biogroups for the

hiM and hiI genes (Table 1a, 1b). Thus, the biogroups related to the hiM and hiI genes are

likely to be functionally different.

We also compared datasets from the DG of typically developing infants with datasets

from rat DG at three different timepoints after seizure induction by injection of pilocarpine or

kainite (day 1, day 3, and day 10) ,and performed principal component analysis on the changes

in mouse hiM/hiI genes at different timepoints (Figure S2a, S2b; Supplementary Results). The

time-course of changes in the mouse hiM genes after seizure induction was different from the

time-course of changes in the mouse hiI genes. In addition, we conducted a spatial pattern
analysis of the mouse hiM/hiI genes, which indicated that their protein products have slightly
different patterns of subcellular localization (Figure S2c; Supplementary Results). The mouse
hiM genes tend to be strongly expressed at the plasma membrane, with expression changes
stabilizing by the third day after seizure induction. By contrast, the hiI genes tend to be
expressed in the nucleus and changes in expression after seizure induction are slower to
stabilize. Together, these results indicate that the hiM/hiI genes have different spatiotemporal
patterns of changes in expression.
192
***Gene expression patterns in patients can be characterized in terms of hiM/hiI genes***
Next, we investigated whether and to what extent the expression changes in
maturation-related genes induced by hyperexcitation overlap with gene expression patterns in
various neuropsychiatric disorders. As above, we evaluated similarities between the changes in
gene expression patterns in different groups using overlap *P*-values calculated by Running
Fisher algorithm (Figure 2a). Similarity indexes for each comparison were defined as the −log
of the overlap *P*-values with hiM or hiI genes, denoted by hiM-index or hiI-index, respectively.
High values in hiM-/hiI-index represent that there is large overlap between the dataset analyzed
and hiM/hiI genes. We obtained the hiM-/hiI-indexes for the datasets from human patients and
plotted them in two-dimensional (2-D) space to show the extent of overlap between datasets
and hiM/hiI genes (Figure 2a).
We initially performed this 2-D analysis on a dataset containing the expression
profile of the prefrontal cortex in the postmortem brains of patients with schizophrenia
(schizophrenia dataset #1: details in Table S2) (Figure 2b). The expression of 1744 genes
differed between patients and healthy controls (significance level of 0.05). The numbers of hiM
and hiI genes with altered expression in schizophrenia dataset #1 were 87 and 76, respectively,
and the overlap *P*-values were $2.4 \times 10^{-10}$ and $6.9 \times 10^{-7}$. The hiM-index and hiI-index for this
dataset were 9.62 (= −log($2.4\times10^{-10}$)) and 6.16 (= −log($6.9\times10^{-7}$)). This result corresponds to a
point in 2-D space (Figure 2b). Points that fall below the unity line (dashed line) indicate
datasets in which changes in hiM genes are dominant, whereas points above the unity line
indicate datasets with dominant changes in hiI genes. The angle from unity line indicates the
degree of the hiM-/hiI- dominance. The same analysis was performed for other schizophrenia
datasets (schizophrenia datasets #2–#16), including ones obtained from different areas of the
postmortem brain and from cultured neurons derived from the iPSCs of patients. Scatter plots
of the results from the schizophrenia datasets are shown in Figure 3a. Thirteen out of sixteen
points were below the unity line. Most of the schizophrenia datasets exhibited hiM-index

dominant patterns, showing high hiM-index values and low hiI-index values (Figure 3a).

We extended the same analysis to 80 disease datasets from seven other

neuropsychiatric diseases (amyotrophic lateral sclerosis (ALS), Alzheimer's disease (AD),

autism spectrum disorder (ASD), Parkinson's disease (PD), bipolar disorder (BPD),

Huntington's disease (HD), and major depressive disorder (MDD); Table S2). Results of each

dataset are shown in Figure 3b-3h. Overall distribution patterns of each disease are shown in

Figure 3i. The ALS datasets tended to show higher hiI-index values than hiM-index values,

indicating hiI-index-dominant pattern (Figure 3b). The AD datasets showed different patterns

in the hiM-/hiI-index depending on the type of sample; datasets from the postmortem brains of

patients with AD tended to show high values only in the hiM-index, and datasets from patient

iPSCs tended to show high values only in the hiI-index (Figure 3c). Datasets from ASD did not

show any dominant patterns either in the hiM-index or hiI-index (Figure 3d). Most datasets

from patients with PD, BPD, HD, and MDD did not show pronounced values in the hiM-index

or the hiI-index (Figure 3e, 3f, 3g, 3h).

Thus, the 2-D analysis revealed that some neuropsychiatric diseases have

characteristic patterns in the hiM-/hiI-indexes: for example, most datasets from patients with

schizophrenia exhibited a higher hiM-index than hiI-index, whereas ALS datasets showed

hiI-index-dominant pattern (Figure 3i). Meanwhile, different diseases sometimes show similar

changes in the hiM- or hiI-index; for example, some of the schizophrenia, ASD, and AD

datasets shared a high hiM-index, and some of the ALS, and AD datasets shared a high

hiI-index. The other four diseases—PD, BPD, HD, and MDD—did not show pronounced

changes in hiM-/hiI-indexes, suggesting that these diseases may not share endophenotype as

pseudoimmaturity inducible by neural hyperexcitation. These results raise the possibility that

there are patterns of gene-expression perturbations that are shared and distinct across these

neuropsychiatric disorders.

***Genetic and environmental risk factors induce changes in the pattern of expression of***

***hiM/hiI genes***

Previous studies suggest that many genetic risk and environmental factors, such as

seizure, hypoxia, and infection, contribute to the development of neuropsychiatric

disorders[2,49,50]. We next applied the 2-D analysis technique to datasets from genetic animal

models of disorders and from animals that experienced risk events.

First, we obtained publicly available datasets for mice that had experienced putative

risk events for schizophrenia, bipolar disorders, and Alzheimer's disease, including seizure

(#1: GSE49030, #2: GSE4236)[51,52], ischemia (#1: GSE32529, #2: GSE35338)[1–3], and infection

(mimicked by CpG; GSE32529)[53,54]. All the studies used here included datasets for different
time points after the risk event; hence, we were able to examine the time-course of changes in
the hiM- and hiI-indexes to reveal the short- and long-term effect of risk event on expression
patterns of hiM/hiI genes. The results showed that datasets from mouse hippocampus treated
with kainite, seizure-inducing drug, exhibited time-course changes in the hiM- and hiI-indexes:
the hiM-index tended to be dominant in the early stage after seizure induction, and the
hiI-index became more dominant in the late stages (Figure 4a: seizure #1). The results from
other datasets on seizure, ischemia, and infection showed roughly similar time-course pattern
changes in the hiM- and hiI-indexes as those observed in seizure dataset #1, being relatively
hiM-index-dominant in the early stage, and then relatively hiI-index-dominant in the later
stage (Figure 4a: seizure, ischemia, and CpG infection). These results indicate that different
types of putative risk events for neuropsychiatric disorders induce roughly similar time-course
changes in the expression of maturation-related genes induced by neural hyperexcitation.

Next, we obtained datasets from animal models with a genetic risk of a
neurodegenerative disease: mice with transgenic expression of a G93A mutant form of human
SOD1, as a model of ALS (#1: GSE46298, #2: GSE18597)[55,56]; transgenic mice with mutant
human amyloid precursor protein (APP) and presenilin1 (PSEN1) genes, which cause familial
Alzheimer's disease (#1: GSE64398, #2: GSE64398)[57]; and Df16(A) heterozygous mice
carrying a chromosome 16 deletion syntenic to human 22q11.2 microdeletions, as a model of
schizophrenia (GSE29767)[58]. We also obtained datasets from Schnurri-2 (Shn-2) knockout
mice as a model of schizophrenia[12] and intellectual disability[14,59,60] and from mice with
heterozygous knockout of the alpha-isoform of calcium/calmodulin-dependent protein kinase
II (alpha-CaMKII+/−)[10,13] as a model of bipolar disorder. We performed 2-D analysis on these
datasets and evaluated the changes in the hiM-/hiI-indexes of these model mice. Both datasets
from transgenic mice with a SOD1(G93A) mutation exhibited a higher hiI-index than
hiM-index in the later stages of disease progression (Figure 4b). These hiI-index-dominant
patterns were also observed in the results derived from human patients with ALS (Figure 3b).
In the mice with mutant human APP and PSEN1, both the hiM- and hiI-indexes increased in
dataset from hippocampus and only hiI-index increased in dataset from cortex during the
course of disease progression (Figure 4c). These patterns are neutral or hiI-index-dominant.
These patterns partially mimic the results from human patients with Alzheimer's disease
(Figure 3c). The Df16(A) heterozygous mice and alpha-CaMKII+/− mice showed
hiM-index-dominant patterns, which are similar to results from human patients with
schizophrenia (Figure 4d). Shn-2 KO mice showed high values for both the hiM- and
hiI-indexes (Figure 4d). Thus, the results from the 2-D analysis of animal models are to some

extent consistent with the results from human patients, indicating that these model mice share

similar patterns of pseudoimmaturity induced by neural hyperexcitation with those of human

patients.

292              (2236 words)

**Discussion**

In this study, we demonstrated that neural hyperexcitation induces changes in the

pattern of gene expression in the DG that are significantly similar to the immature

hippocampus of typically developing human fetuses. From the pool of genes, we identified two

groups of genes, and found that these are shared by multiple neuropsychiatric disorders, such

as schizophrenia, Alzheimer disorders, and ALS.

Many of the datasets from patients with schizophrenia and from the postmortem

brains of patients with Alzheimer's disease exhibited hiM-index-dominant pattern changes.

The hiM genes include a GABA receptor, voltage-dependent calcium channel, glutamate

receptor, and voltage-dependent sodium channel (Table S1). These genes have been reported to

be implicated in the pathological changes in the brains of patients with schizophrenia and

Alzheimer's disease[61–64]. Thus, many of the synaptic genes that changed in the brains of

patients with schizophrenia or Alzheimer's disease could be genes whose expression increases

during maturation and decreases with neural hyperexcitation. Although reductions in the

expression of some synaptic genes in these disorders are well documented, our results are the

first to raise the possibility that neuronal hyperexcitation may also induce reductions in such

synaptic molecules.

Most of the datasets from patients with ALS and Alzheimer's disease exhibited

hiI-index-dominant patterns. The hiI genes include DNA methyltransferase, cyclin,

cyclin-dependent kinase, integrin beta 3 binding protein, and tumor protein p53 (Table S1).

These genes are known to be important in chromosomal modification and DNA repair, and

abnormal functions of these systems have been observed in patients with ALS and Alzheimer's

disease[65–69]. Thus, some of the genes that are considered to be important in the development of

these disorders are immaturity-related genes, whose expressions decrease during maturation

and can be increased by neural hyperexcitation.

As for the datasets from patients with PD, BPD, HD, and MDD, most of them did not

show significant overlap either hiM or hiI genes, indicating that there might not be pathological

changes of pseudoimmaturity inducible by neural hyperexcitation in the datasets of these four

diseases. Thus, we suggest that gene expression analysis based on the findings derived from

shared endophenotypes are helpful to conduct transdiagnostic characterization of

neuropsychiatric disorders.

Our study has some limitations. First, the number of available datasets was limited.

All the datasets except the one for mouse development were obtained from the BaseSpace

Correlation Engine. On this platform, vast numbers (over 21,000) of complex biological and

clinical datasets are available. Even though we used all the gene expression dataset hits from

our keyword query to avoid sampling bias, the number of datasets was still small, from 8

datasets for ALS to 16 datasets for schizophrenia. Further accumulation of the studies will

improve the reliability of our results. Another limitation is that the datasets used in this study

are from different types of sample, including various central nervous system areas, such as the

hippocampus, prefrontal cortex, striatum, and the spinal cord. The gene expression

abnormalities in patients could differ depending on the brain area[48]. We also used datasets from

cultured neurons differentiated from the iPSCs of human subjects, and it is controversial

whether the pattern of gene expression in these neurons is comparable with that of neurons in

the patients' brains[70,71]. It is also possible that the altered gene expression in the postmortem

brains is due to the effects of medication rather than pathological changes from the disease

itself[72]. Other conditions that were not controlled in this study include the age at death, storage

conditions of the samples, genetic background of animals, and animal housing conditions. For

these reasons, we need to be careful in interpreting the results of the analyses. It is noteworthy,

however, that despite the variety of sample types used, we were able to identify some shared

and distinct patterns of gene expression.

Recent attempts such as RDoC initiative have tried to reclassify psychiatric disorders

according to biological domains (e.g. genes, neural circuits, behavior)[7]. While Gandal *et al*

conducted non-biased coexpression analyses[8], in this study, we utilized gene groups which

were derived from the findings based on the studies of animal models of neuropsychiatric

disorders. Characterization by these gene groups enabled us to extract novel biological features

of some neuropsychiatric disorders which are related to pseudoimmaturity inducible by neural

hyperexcitation. Detecting such domains that extract the biological features of each

neuropsychiatric disorder will move this diagnostic framework forward, from criteria based on

signs and symptoms to those including biological dimensions.

In conclusion, biological domain, which is pseudoimmaturity inducible by neural

hyperexcitation, are common endophenotype among several neuropsychiatric disorders.

Future studies are needed to find translational indices that correspond to these features and can

be applicable to human patients for better diagnosis of these neuropsychiatric disorders. Our

findings here may promote the development of novel biomarkers, leading to better diagnosis of

neuropsychiatric disorders.

(793 words)

(Main Text Total: 3967 words)

**Methods**

*Microarray experiments to examine mouse DG development*

Wild-type mouse DGs were sampled at postnatal day 8, 11, 14, 17, 21, 25, 29

(C57BL/6J × BALB/cA background; n = 5)[73], and microarray experiments were performed as

previously described[10]. The microarray data, including those used in this study, were deposited

in the GEO database under accession number GSE113727. We also obtained a dataset for

33-week-old wild-type mice, which we previously reported (C57BL/6J × BALB/cA

background) (GSE42778)[12]. We integrated these two datasets into one to construct the dataset

for the development of wild-type mouse DG used in this study (P8 versus adult, fold change >

1.2, $P < 0.05$).

*Data collection and processing*

Except for the mouse DG developmental dataset mentioned above, the 91 gene

expression datasets used in this study were obtained from publicly available databases (listed

in Table S2). All gene expression datasets were analyzed with the BaseSpace Correlation

Engine          (BSCE;          formally          known          as          NextBio)

(https://japan.ussc.informatics.illumina.com/c/nextbio.nb; Illumina, Cupertino, CA), a

database of biomedical experiments. BaseSpace is a repository of analyzed gene expression

datasets that allows researchers to search expression profiles and other results[44]. The datasets

registered in BaseSpace undergo several preprocessing, quality control, and organization

stages. Quality control ensures the integrity of the samples and datasets and includes

evaluations of pre- and postnormalization boxplots, missing value counts, and *P*-value

histograms (after statistical testing) with false-discovery rate analysis to establish whether the

number of significantly altered genes is larger than that expected by chance. Other microarray

data processing was performed in MAS5 (Affymetrix, Santa Clara, CA, USA)[44].

Genes with a *P*-value < 0.05 (without correction for multiple testing) and an absolute

fold change > 1.2 were included in the differentially expressed gene datasets. This sensitivity

threshold is typically the lowest used with commercial microarray platforms and the default

criterion in BaseSpace analyses[44]. All data from the Affymetrix GeneChip series were

downloaded from the NCBI GEO database. Affymetrix Expression Console software

(specifically the robust multiarray average algorithm) was used to preprocess the data.

We used the expression values (on a log base-2 scale) to calculate the fold changes

and *P*-values between two conditions (infants–adults and patients–healthy controls). To

determine the fold changes, the expression values of the probes/genes in the test data sets were

divided by those of the control data sets. If the fold change was < 1.0, these values were

converted into the negative reciprocal or $-1/$(fold change). Genes with an absolute fold change

> 1.2 and a $t$-test $P$-value < 0.05 were imported into BSCE according to the instructions

provided by the manufacturer. The rank order of these genes was determined by their absolute

fold change. We compared the signatures in two given gene sets using BSCE. All statistical

analyses were performed in BaseSpace and similarities between any two datasets were

evaluated as overlap $P$-values using the Running Fisher algorithm[44].

(472words)

**Funding and Disclosures**

Competing financial interests: K.T. and M.M. are employees of Astellas Pharma Inc. Other authors report no biomedical financial interests.

**Acknowledgments**

We thank Wakako Hasegawa, Yumiko Mobayashi, Misako Murai, Tamaki Murakami, Miwa Takeuchi, Yoko Kagami, Harumi Mitsuya, Aki Miyakawa and other members of Miyakawa lab for their support. This work was supported by JSPS Grant-in-Aid for Scientific Research on Innovative Areas Grant Number 25116526, 15H01297 ("Microendophenotype"), 16H06462 ("Scrap & Build"), JSPS KAKENHI Grant Number 25242078, and grant from Astellas Pharma Inc.

**Author Contribution Statement**

T.M. generated the data, performed the analysis, prepared all figures and wrote the manuscript. All authors reviewed the manuscript.

    **Figure legends**

    *Figure 1.*

        The patterns of changes in gene expression in rat DG 1 day after pilocarpine

    treatment compared with developmental changes in mouse DG and human hippocampus. Venn

    diagrams illustrating the overlap in genome-wide gene-expression changes between rat DG

    after seizure induction (GSE47752) and the DG of typically developing mouse infants

    (GSE113727; P8 infants compared with 33-week adults) (a) or the hippocampus of typically

    developing human fetuses (GSE25219: 19–23 week fetuses compared with 8–9 week fetuses)

    (b). Bar graphs illustrate the –log of the overlap *P*-values for genes upregulated (red arrows) or

    downregulated (green arrows) by each condition. The Bonferroni correction was used to adjust

    the significance level according to the number of dataset pairs (see the Methods section and

    Supplementary Method). Genes that were downregulated in both conditions were defined as

    mouse hiM genes (green bar in (a)), and genes that were upregulated in both conditions are

    defined as mouse hiI genes (red bar in (a)). Similarly, human hiM genes and human hiI genes

    were defined as the groups of genes with positive correlation between the two conditions,

    development and seizure (b).

    *Figure 2.*

        Overview of the two-dimensional analysis (2-D analysis) for disease datasets. (a)

    Genes with expression changes in the disease datasets are compared with the hiM and hiI gene

    groups. The hiM- and hiI-indexes were defined as the –log of the overlap *P*-values with the

    hiM and hiI genes. The gene expression patterns of the disease datasets are plotted in

    two-dimensional coordinates, in which the x-/y-axes are defined by the hiM-/hiI-indexes. Each

    dataset is characterized as hiM- or hiI- dominant by the ratio of hiM-/hiI-indexes, and the

    degree of the hiM-/hiI-dominance are evaluated by deviation from the unity line. Distance of

    each dataset from the origin show the degree of overlap with hiM-/hiI-genes. (b) 2-D analysis

    applied to a dataset of postmortem brains (prefrontal cortex) from patients with schizophrenia

    (schizophrenia dataset #1). The expression levels of 1744 genes were significantly changed in

    this disease dataset. Of these, 87 and 76 genes overlap with the hiM/hiI genes. The overlap

    *P*-values between the disease dataset and the hiM/hiI genes were $2.4 \times 10^{-10}$ and $6.9 \times 10^{-7}$.

    The hiM- and hiI-indexes for the disease dataset were therefore 9.62 and 6.16, indicating that

    this dataset is hiM-dominant. The results of the 2-D analysis for this dataset are plotted in the

    two-dimensional coordinates defined by the hiM- and hiI-indexes.

*Figure 3.*

Two-dimensional analysis (2-D analysis) for disease datasets from various

neuropsychiatric disorders. Each point corresponds to the result from one independent study.

(a-h) Results of the 2-D analysis of datasets for schizophrenia (a), ALS (b), Alzheimer's

disease (c), autism (d), Parkinson's disease (e), bipolar disorder (f), Huntington's disease (g),

and major depression (h). Filled points indicate datasets from the postmortem brain or spinal

cord (ALS) of patients, and open points indicate those from cultured neural cells from patient

iPSCs. (i) The distribution patterns of hiM- and hiI-index for all diseases analyzed. The extent

of the changes in hiM-/hiI-indexes is assessed by the average distance of all datasets in each

disease from origins. Four diseases whose average distance from origin are over 5.0 are shown

as circular sectors, and others are shown as points. The radii of circular sector indicate the

average distance of all datasets in each disease from origins, and the central angles of circular

sector are average deviation $\pm$ sem from unity line. Each point indicates their average

distance from origin and average deviation from unity line.

*Figure 4.*

Time-dependent changes in the hiM-/hiI-indexes in animals subjected to various

putative risk events for neuropsychiatric disorders and in genetic mouse models of

schizophrenia, bipolar disorder, ALS, and Alzheimer's disease. (a) Pattern of changes in the

hiM- and hiI-indexes in mouse and rat hippocampus after treatment with kainite (seizure #1

(GSE1831) and seizure #2 (GSE4236)), in mouse cortex and astrocytes after middle cerebral

artery occlusion (MCAO; ischemia #1 (GSE32529), ischemia #2 (GSE35338)), and in mouse

cortex after CpG infection (GSE32529). (b) Pattern of changes in the hiM- and hiI-indexes of

the spinal cord of an ALS mouse model with the SOD1(G93A) mutation. (c) Pattern of changes

in the hiM- and hiI-indexes of the hippocampus and cortex of an Alzheimer's disease mouse

model with mutations in APP and PSEN1. (d) hiM- and hiI-indexes in mouse models of

schizophrenia and bipolar disorder.

*Table 1.*

Summary of results from the pathway analyses of hiM/hiI genes. (a) The 20

biogroups with the most significant similarities to mouse hiM genes and mouse hiI genes.

Green columns indicate biogroups that are related to the plasma membrane. Red columns

indicate biogroups that are related to reactions in the nucleus. (b) The 20 biogroups with the

most significant similarities to human hiM genes and human hiI genes.

(790 words, 4 figures, 1 table)

**References**

1.      Cross-national comparisons of the prevalences and correlates of mental

disorders. WHO International Consortium in Psychiatric Epidemiology. *Bull World Health*

*Organ* **78,** 413–426 (2000).

2.      Cardno, A. G. & Owen, M. J. Genetic Relationships Between

Schizophrenia, Bipolar Disorder, and Schizoaffective Disorder. *Schizophr Bull* **40,** 504–515

(2014).

3.      Hall, J., Trent, S., Thomas, K. L., O'Donovan, M. C. & Owen, M. J.

Genetic Risk for Schizophrenia: Convergence on Synaptic Pathways Involved in Plasticity.

*Biological Psychiatry* **77,** 52–58 (2015).

4.      Forero, D. A. *et al.* A network of synaptic genes associated with

schizophrenia and bipolar disorder. *Schizophrenia Research* **172,** 68–74 (2016).

5.      Douaud, G. *et al.* A common brain network links development, aging, and

vulnerability to disease. *PNAS* **111,** 17648–17653 (2014).

6.      Argyelan, M. *et al.* Resting-State fMRI Connectivity Impairment in

Schizophrenia and Bipolar Disorder. *Schizophr Bull* **40,** 100–110 (2014).

7.      Insel, T. R. & Cuthbert, B. N. Brain disorders? Precisely. *Science* **348,**

499–500 (2015).

8.      Gandal, M. J. *et al.* Shared molecular neuropathology across major

psychiatric disorders parallels polygenic overlap. *Science* **359,** 693–697 (2018).

9.      Hagihara, H., Takao, K., Walton, N. M., Matsumoto, M. & Miyakawa, T.

Immature Dentate Gyrus: An Endophenotype of Neuropsychiatric Disorders. *Neural Plasticity*

**2013,** (2013).

10.      Yamasaki, N. *et al.* Alpha-CaMKII deficiency causes immature dentate

gyrus, a novel candidate endophenotype of psychiatric disorders. *Molecular Brain* **1,** 6 (2008).

11.      Ohira, K. *et al.* Synaptosomal-associated protein 25 mutation induces

immaturity of the dentate granule cells of adult mice. *Molecular Brain* **6,** 12 (2013).

12.      Takao, K. *et al.* Deficiency of Schnurri-2, an MHC Enhancer Binding

Protein, Induces Mild Chronic Inflammation in the Brain and Confers Molecular, Neuronal,

and Behavioral Phenotypes Related to Schizophrenia. *Neuropsychopharmacology* **38,**

1409–1425 (2013).

13.      Hagihara, H. *et al.* Circadian Gene Circuitry Predicts Hyperactive

Behavior in a Mood Disorder Mouse Model. *Cell Reports* **14,** 2784–2796 (2016).

14.      Nakao, A. *et al.* Immature morphological properties in subcellular-scale

structures in the dentate gyrus of Schnurri-2 knockout mice: a model for schizophrenia and

intellectual disability. *Molecular Brain* **10,** 60 (2017).

15.      Hagihara, H., Fujita, M., Umemori, J., Hashimoto, M. & Miyakawa, T.

Immature-like molecular expression patterns in the hippocampus of a mouse model of

dementia with Lewy body-linked mutant β-synuclein. (2018).

16.      Shin, R. *et al.* The immature dentate gyrus represents a shared phenotype

of mouse models of epilepsy and psychiatric disease. *Bipolar Disord* **15,** 405–421 (2013).

17.      You, J. C. *et al.* Epigenetic suppression of hippocampal calbindin-D28k by

ΔFosB drives seizure-related cognitive deficits. *Nature Medicine* **23,** 1377 (2017).

18.      Maglóczky, Z., Halász, P., Vajda, J., Czirják, S. & Freund, T. F. Loss of

Calbindin-D28K immunoreactivity from dentate granule cells in human temporal lobe

epilepsy. *Neuroscience* **76,** 377–385 (1997).

19.      Karádi, K. *et al.* Correlation between calbindin expression in granule cells

of the resected hippocampal dentate gyrus and verbal memory in temporal lobe epilepsy.

*Epilepsy & Behavior* **25,** 110–119 (2012).

20.      Zhang, Z. J. & Reynolds, G. P. A selective decrease in the relative density

of parvalbumin-immunoreactive neurons in the hippocampus in schizophrenia. *Schizophrenia*

*Research* **55,** 1–10 (2002).

21.      Hashimoto, T. *et al.* Gene Expression Deficits in a Subclass of GABA

Neurons in the Prefrontal Cortex of Subjects with Schizophrenia. *J. Neurosci.* **23,** 6315–6326

(2003).

22.      Fung, S. J. *et al.* Expression of Interneuron Markers in the Dorsolateral

Prefrontal Cortex of the Developing Human and in Schizophrenia. *AJP* **167,** 1479–1488

(2010).

23.      Pantazopoulos, H., Woo, T.-U. W., Lim, M. P., Lange, N. & Berretta, S.

Extracellular Matrix-Glial Abnormalities in the Amygdala and Entorhinal Cortex of Subjects

Diagnosed With Schizophrenia. *Arch Gen Psychiatry* **67,** 155–166 (2010).

24.      Berretta, S. Extracellular matrix abnormalities in schizophrenia.

*Neuropharmacology* **62,** 1584–1597 (2012).

25.      Hagihara, H., Ohira, K., Takao, K. & Miyakawa, T. Transcriptomic

evidence for immaturity of the prefrontal cortex in patients with schizophrenia. *Molecular*

*Brain* **7,** 41 (2014).

26.      Gandal, M. J., Nesbitt, A. M., McCurdy, R. M. & Alter, M. D. Measuring

the Maturity of the Fast-Spiking Interneuron Transcriptional Program in Autism,

Schizophrenia, and Bipolar Disorder. *PLOS ONE* **7,** e41215 (2012).

27.      Mellios, N. *et al.* Molecular Determinants of Dysregulated GABAergic

Gene Expression in the Prefrontal Cortex of Subjects with Schizophrenia. *Biological*

*Psychiatry* **65,** 1006–1014 (2009).

28.     Hashimoto, T. *et al.* Alterations in GABA-related transcriptome in the

dorsolateral prefrontal cortex of subjects with schizophrenia. *Mol. Psychiatry* **13,** 147–161

(2008).

29.     Murano, T., Koshimizu, H., Hagihara, H. & Miyakawa, T. Transcriptomic

immaturity of the hippocampus and prefrontal cortex in patients with alcoholism. *Scientific*

*Reports* **7,** srep44531 (2017).

30.     Kobayashi, K. *et al.* Reversal of hippocampal neuronal maturation by

serotonergic antidepressants. *Proceedings of the National Academy of Sciences* **107,**

8434–8439 (2010).

31.     Ohira, K., Takeuchi, R., Iwanaga, T. & Miyakawa, T. Chronic fluoxetine

treatment reduces parvalbumin expression and perineuronal netsin gamma-aminobutyric

acidergic interneurons of the frontal cortex in adultmice. *Molecular Brain* **6,** 43 (2013).

32.     Imoto, Y., Segi-Nishida, E., Suzuki, H. & Kobayashi, K. Rapid and stable

changes in maturation-related phenotypes of the adult hippocampal neurons by

electroconvulsive treatment. *Molecular Brain* **10,** 8 (2017).

33.     Chang, Y.-T. *et al.* Bidirectional relation between schizophrenia and

epilepsy: A population-based retrospective cohort study. *Epilepsia* **52,** 2036–2042 (2011).

34.     Hyde, T. M. & Weinberger, D. R. Seizures and schizophrenia. *Schizophr*

*Bull* **23,** 611–622 (1997).

35.     Cascella, N. G., Schretlen, D. J. & Sawa, A. SCHIZOPHRENIA AND

EPILEPSY: IS THERE A SHARED SUSCEPTIBILITY? *Neuroscience research* **63,** 227

(2009).

36.     Coyle, J. T. Glutamate and Schizophrenia: Beyond the Dopamine

Hypothesis. *Cell Mol Neurobiol* **26,** 363–382 (2006).

37.     Mouri, A., Noda, Y., Enomoto, T. & Nabeshima, T. Phencyclidine animal

models of schizophrenia: Approaches from abnormality of glutamatergic neurotransmission

and neurodevelopment. *Neurochemistry International* **51,** 173–184 (2007).

38.     Kehrer, C., Maziashvili, N., Dugladze, T. & Gloveli, T. Altered

Excitatory-Inhibitory Balance in the NMDA-Hypofunction Model of Schizophrenia. *Front*

*Mol Neurosci* **1,** (2008).

39.     O'Donnell, C., Gonçalves, J. T., Portera-Cailliau, C. & Sejnowski, T. J.

Beyond excitation/inhibition imbalance in multidimensional models of neural circuit changes

in brain disorders. *eLife* **6,**

597  40.  Mertens, J. *et al.* Differential responses to lithium in hyperexcitable

598 neurons from patients with bipolar disorder. *Nature* **527,** 95–99 (2015).

599  41.  Vossel, K. A. *et al.* Incidence and impact of subclinical epileptiform

600 activity in Alzheimer's disease. *Annals of Neurology* **80,** 858–870

601  42.  Lam, A. D. *et al.* Silent Hippocampal Seizures and Spikes Identified by

602 Foramen Ovale Electrodes in Alzheimer's Disease. *Nat Med* **23,** 678–680 (2017).

603  43.  Vossel, K. A., Tartaglia, M. C., Nygaard, H. B., Zeman, A. Z. & Miller, B.

604 L. Epileptic activity in Alzheimer's disease: causes and clinical relevance. *The Lancet*

605 *Neurology* **16,** 311–322 (2017).

606  44.  Kupershmidt, I. *et al.* Ontology-Based Meta-Analysis of Global

607 Collections of High-Throughput Public Data. *PLOS ONE* **5,** e13066 (2010).

608  45.  Takao, K. & Miyakawa, T. Genomic responses in mouse models greatly

609 mimic human inflammatory diseases. *PNAS* **112,** 1167–1172 (2015).

610  46.  Ryan, S. D. *et al.* Isogenic Human iPSC Parkinson's Model Shows

611 Nitrosative Stress-Induced Dysfunction in MEF2-PGC1α Transcription. *Cell* **155,** 1351–1364

612 (2013).

613  47.  Dingledine, R. *et al.* Data Descriptor: Transcriptional profile of

614 hippocampal dentate granule cells in four rat epilepsy models. *Scientific Data* **4,** (2017).

615  48.  Kang, H. J. *et al.* Spatio-temporal transcriptome of the human brain.

616 *Nature* **478,** 483–489 (2011).

617  49.  van Os, J., Kenis, G. & Rutten, B. P. F. The environment and

618 schizophrenia. *Nature* **468,** 203–212 (2010).

619  50.  DeMichele-Sweet, M. a. A. *et al.* Genetic risk for schizophrenia and

620 psychosis in Alzheimer disease. *Molecular Psychiatry* (2017). doi:10.1038/mp.2017.81

621  51.  Wilson, D. N. *et al.* Microarray analysis of postictal transcriptional

622 regulation of neuropeptides. *J Mol Neurosci* **25,** 285 (2005).

623  52.  Hermey, G. *et al.* Genome-Wide Profiling of the Activity-Dependent

624 Hippocampal Transcriptome. *PLOS ONE* **8,** e76903 (2013).

625  53.  Vartanian, K. B. *et al.* LPS preconditioning redirects TLR signaling

626 following stroke: TRIF-IRF3 plays a seminal role in mediating tolerance to ischemic injury.

627 *Journal of Neuroinflammation* **8,** 140 (2011).

628  54.  Stevens, S. L. *et al.* Multiple Preconditioning Paradigms Converge on

629 Interferon Regulatory Factor-Dependent Signaling to Promote Tolerance to Ischemic Brain

630 Injury. *J. Neurosci.* **31,** 8456–8463 (2011).

631  55.  Nardo, G. *et al.* Transcriptomic indices of fast and slow disease

progression in two mouse models of amyotrophic lateral sclerosis. *Brain* **136,** 3305–3332

(2013).

56.  Lerman, B. J. *et al.* Deletion of galectin-3 exacerbates microglial

activation and accelerates disease progression and demise in a SOD1 G93A mouse model of

amyotrophic lateral sclerosis. *Brain and Behavior* **2,** 563–575 (2012).

57.  Matarin, M. *et al.* A Genome-wide Gene-Expression Analysis and

Database in Transgenic Mice during Development of Amyloid or Tau Pathology. *Cell Reports*

**10,** 633–644 (2015).

58.  Xu, B., Hsu, P.-K., Stark, K. L., Karayiorgou, M. & Gogos, J. A.

Derepression of a Neuronal Inhibitor due to miRNA Dysregulation in a Schizophrenia-Related

Microdeletion. *Cell* **152,** 262–275 (2013).

59.  Srivastava, S. *et al.* Loss-of-function variants in *HIVEP2* are a cause of

intellectual disability. *European Journal of Human Genetics* **24,** 556–561 (2016).

60.  Steinfeld, H. *et al.* Mutations in HIVEP2 are associated with

developmental delay, intellectual disability, and dysmorphic features. *Neurogenetics* **17,**

159–164 (2016).

61.  Harrison, P. J. The hippocampus in schizophrenia: a review of the

neuropathological evidence and its pathophysiological implications. *Psychopharmacology* **174,**

151–162 (2004).

62.  Glausier, J. R. & Lewis, D. A. Dendritic spine pathology in schizophrenia.

*Neuroscience* **251,** 90–107 (2013).

63.  Fatemi, S. H., Folsom, T. D. & Thuras, P. D. GABAA and GABAB

receptor dysregulation in superior frontal cortex of subjects with schizophrenia and bipolar

disorder. *Synapse* **71,** (2017).

64.  Lisman, J. E. *et al.* Circuit-based framework for understanding

neurotransmitter and risk gene interactions in schizophrenia. *Trends Neurosci.* **31,** 234–242

(2008).

65.  Chestnut, B. A. *et al.* Epigenetic Regulation of Motor Neuron Cell Death

through DNA Methylation. *J. Neurosci.* **31,** 16619–16636 (2011).

66.  Martin, L. J. & Wong, M. Aberrant Regulation of DNA Methylation in

Amyotrophic Lateral Sclerosis: A New Target of Disease Mechanisms. *Neurotherapeutics* **10,**

722–733 (2013).

67.  Nguyen, M. D. *et al.* Cell Cycle Regulators in the Neuronal Death Pathway

of Amyotrophic Lateral Sclerosis Caused by Mutant Superoxide Dismutase 1. *J. Neurosci.* **23,**

2131–2140 (2003).

68.      Dorszewska, J. *et al.* Mutations of TP53 Gene and Oxidative Stress in
Alzheimer's Disease Patients. *Advances in Alzheimer's Disease* **03,** 24 (2014).

69.      Moskalev, A. A. *et al.* Gadd45 proteins: Relevance to aging, longevity and
age-related pathologies. *Ageing Research Reviews* **11,** 51–66 (2012).

70.      Brennand, K. J. *et al.* Modelling schizophrenia using human induced
pluripotent stem cells. *Nature* **473,** 221–225 (2011).

71.      Wen, Z. *et al.* Synaptic dysregulation in a human iPS cell model of mental
disorders. *Nature* **515,** 414–418 (2014).

72.      Chan, M. K. *et al.* Evidence for disease and antipsychotic medication
effects in post-mortem brain from schizophrenia patients. *Mol Psychiatry* **16,** 1189–1202
(2011).

73.      Hagihara, H., Toyama, K., Yamasaki, N. & Miyakawa, T. Dissection of
hippocampal dentate gyrus from adult mouse. *J Vis Exp* (2009). doi:10.3791/1543

**a**

hiM genes (human)

Disease dataset

hiI genes (human)

Definition of hiM–/hiI– index for Disease dataset

hiM-index = –log(overlap *P*-value with[hiM genes])
hiI-index = –log(overlap *P*-value with[hiI genes])

Unity Line

hiI-index dominant

hiM-index dominant

Disease dataset
(hiM-index, hiI-index)

hiI-index

hiM-index

**b**

Schizophrenia data set #1
(GSE21138)

hiM genes          hiI genes

579          1744          716

87          76

Overlap *P*-value
= 2.4x10$^{-10}$

Overlap *P*-value
= 6.9x10$^{-7}$

Schizophrenia data set #1

(hiM-index, hiI-index)
= (9.62, 6.16)

hiI-index

hiM-index

**a** Mouse DG development vs Rat DG Seizure (Pilocarpine day1)

**b** Human Hippocampus development vs Rat DG Seizure (Pilocarpine day1)

## a

| hiM genes (mouse) | | | |
|---|---|---|---|
| Biogroup name | direction | Common genes | p–value |
| multicellular organismal signaling | down | 56 | 4.00E–23 |
| transmission of nerve impulse | down | 51 | 3.10E–19 |
| axon | down | 45 | 7.30E–19 |
| neuron differentiation | down | 58 | 4.70E–17 |
| synapse | down | 57 | 5.70E–17 |
| neuron development | down | 50 | 1.50E–16 |
| neuronal cell body | down | 45 | 5.20E–16 |
| cell junction | down | 56 | 3.00E–15 |
| cell part morphogenesis | down | 43 | 4.70E–15 |
| neuron projection development | down | 42 | 6.20E–15 |
| cell projection part | down | 56 | 6.50E–15 |
| cell morphogenesis involved in neuron differentiation | down | 35 | 2.90E–14 |
| cell morphogenesis involved in differentiation | down | 40 | 6.70E–14 |
| cellular chemical homeostasis | down | 53 | 2.50E–13 |
| cell–cell signaling | down | 44 | 2.60E–13 |
| synaptic transmission | down | 37 | 3.60E–13 |
| regulation of neurological system process | down | 31 | 1.10E–12 |
| regulation of transmission of nerve impulse | down | 30 | 1.40E–12 |
| Genes involved in Neuronal System | down | 31 | 1.70E–12 |
| cellular ion homeostasis | down | 49 | 1.80E–12 |

| hiI genes (mouse) | | | |
|---|---|---|---|
| Biogroup name | direction | Common genes | p–value |
| response to wounding | up | 60 | 1.10E–16 |
| positive regulation of developmental process | up | 66 | 9.30E–16 |
| cardiovascular system development | up | 61 | 7.70E–15 |
| circulatory system development | up | 61 | 7.70E–15 |
| proteinaceous extracellular matrix | up | 32 | 7.80E–15 |
| Genes involved in Cell Cycle | up | 47 | 1.00E–14 |
| extracellular matrix | up | 37 | 1.30E–14 |
| Genes involved in Cell Cycle, Mitotic | up | 42 | 6.30E–14 |
| protein domain specific binding | up | 60 | 7.00E–14 |
| positive regulation of signal transduction | up | 58 | 8.70E–14 |
| Genes involved in DNA Replication | up | 32 | 9.20E–14 |
| cell cycle | up | 67 | 3.00E–13 |
| Genes involved in Adaptive Immune System | up | 56 | 3.20E–13 |
| Focal adhesion | up | 31 | 5.60E–13 |
| kinase binding | up | 44 | 5.50E–12 |
| cytoskeleton organization | up | 54 | 5.60E–12 |
| neuron differentiation | up | 52 | 5.80E–12 |
| basement membrane | up | 19 | 7.50E–12 |
| regulation of cell migration | up | 36 | 1.10E–11 |
| MAPKinase Signaling Pathway | up | 20 | 1.40E–11 |

## b

| hiM genes (human) | | | |
|---|---|---|---|
| Biogroup name | direction | Common genes | p–value |
| multicellular organismal signaling | up | 87 | 4.40E–58 |
| transmission of nerve impulse | up | 83 | 1.40E–54 |
| synaptic transmission | up | 75 | 6.70E–50 |
| neuron projection | up | 70 | 2.60E–44 |
| axon | up | 42 | 2.90E–33 |
| synapse | up | 51 | 2.60E–32 |
| neuron development | up | 58 | 1.60E–30 |
| Genes involved in Neuronal System | up | 41 | 3.40E–30 |
| Genes involved in Transmission across Chemical Synapses | up | 32 | 1.50E–29 |
| regulation of neurological system process | up | 35 | 6.50E–29 |
| regulation of transmission of nerve impulse | up | 34 | 9.20E–29 |
| cell projection part | up | 53 | 1.80E–27 |
| passive transmembrane transporter activity | up | 44 | 1.80E–26 |
| ion channel activity | up | 43 | 2.20E–26 |
| cell part morphogenesis | up | 49 | 1.80E–24 |
| behavior | up | 38 | 2.90E–24 |
| single–organism behavior | up | 35 | 3.70E–24 |
| cell morphogenesis involved in neuron differentiation | up | 44 | 4.30E–24 |
| neuron projection development | up | 47 | 4.40E–24 |
| dendrite | up | 37 | 2.00E–23 |

| hiI genes (human) | | | |
|---|---|---|---|
| Biogroup name | direction | Common genes | p–value |
| chromosome | down | 65 | 1.30E–32 |
| response to DNA damage stimulus | down | 60 | 4.80E–32 |
| Genes involved in Cell Cycle | down | 50 | 3.60E–31 |
| Genes involved in Cell Cycle, Mitotic | down | 43 | 1.20E–30 |
| Genes involved in DNA Replication | down | 34 | 2.00E–30 |
| interphase | down | 47 | 8.20E–30 |
| interphase of mitotic cell cycle | down | 46 | 5.10E–29 |
| Genes involved in Mitotic M–M/G1 phases | down | 29 | 2.60E–26 |
| Cell cycle | down | 25 | 7.60E–23 |
| DNA repair | down | 40 | 2.30E–22 |
| wound healing | down | 50 | 7.40E–22 |
| response to ionizing radiation | down | 21 | 1.20E–20 |
| nuclear division | down | 35 | 8.00E–20 |
| mitosis | down | 35 | 8.00E–20 |
| cell division | down | 38 | 2.20E–19 |
| G1/S transition of mitotic cell cycle | down | 26 | 1.70E–18 |
| cardiovascular system development | down | 46 | 3.20E–18 |
| circulatory system development | down | 46 | 3.20E–18 |
| S phase | down | 22 | 3.80E–18 |
| blood coagulation | down | 41 | 6.10E–18 |