Exchange catalysis during anaerobic methanotrophy revealed by ^{12}CH$_2$D$_2$ &

^{13}CH$_3$D in methane

Jeanine L. Ash[1*], Matthias Egger[2,#], Tina Treude[3,4], Issaku Kohl[3], Barry Cragg[4], R. John

Parkes[4], Caroline P. Slomp[5], Barbara Sherwood Lollar[6], Edward D. Young[3]*

[1]Department of Earth, Environmental and Planetary Sciences, Rice University, Houston

Texas, USA

[2]Center for Geomicrobiology, Aarhus University, Aarhus, Denmark

[3]Department of Earth, Planetary, and Space Sciences, UCLA, Los Angeles CA, USA

[4]School of Earth and Ocean Sciences, Cardiff University, Cardiff, UK.

[5]Department of Earth Sciences-Geochemistry, Utrecht University, Utrecht, The

Netherlands

[6]Department of Earth Sciences, University of Toronto, Toronto Ontario, Canada

[#]now at The Ocean Cleanup Foundation, Delft, The Netherlands

*correspondence to jeanine.ash@rice.edu, eyoung@epss.ucla.edu

The anaerobic oxidation of methane (AOM) is a crucial component of the methane cycle,

but its enzymatic versatility under environmental conditions remains poorly understood.

We use sediment samples collected during IODP Expedition 347 to the Baltic Sea to

show that relative abundances of ^{12}CH$_2$D$_2$ and ^{13}CH$_3$D molecules in methane gas trace the

reversibility of methyl-coenzyme M reductase during AOM by driving methane towards

internal, thermodynamic isotopic equilibrium. These data suggest that $^{12}CH_2D_2$ and $^{13}CH_3D$ together can identify the influence of methanotrophy in environments where conventional bulk isotope ratios are ambiguous, and these findings may lead to new insights regarding the global significance of enzymatic back-flux in the methane cycle.

Quantifying the subsurface carbon cycle by linking biogeochemistry to deep biosphere metabolisms remains a long-standing challenge compounded by slow rates of reaction (WEBSTER et al., 2015; WAITE et al., 2017). Even in well-characterized environments, determining how methanogens and anaerobic methanotrophs influence organic matter degradation can be challenging (REGNIER et al., 2011; BEULIG et al., 2018a). These challenges are due in part to how methane produced by methanogens inherits characteristics from its substrates (both a range of organic and inorganic carbon) (CONRAD, 2005), and they are further complicated when methane is itself the substrate during anaerobic methane oxidation (AOM) (WHITICAR, 1999). Our understanding of how anaerobic methanotrophs respond to varying environmental conditions (HOLLER et al., 2009; HOLLER et al., 2011; YOSHINAGA et al., 2014) and how they might contribute to "cryptic cycles" (the rapid production and consumption of reactive intermediates) in the marine subsurface is constantly evolving (EGGER et al., 2018).

AOM is thought to be carried out by a consortium of methane-oxidizing archaea and sulphate-reducing bacteria performing the net reaction $CH_4 + SO_4^{2-} \rightarrow HCO_3^- + HS^- + H_2O$ (BOETIUS et al., 2000). Evidence suggests that these processes can be decoupled (MILUCKA et al., 2012; SCHELLER et al., 2016), and metal oxides may also act as terminal

electron acceptors (BEAL *et al.*, 2009; CAI *et al.*, 2018). In all cases, the first step of AOM

involves C-H bond activation by a modified methyl-coenzyme M reductase (Mcr), the

terminal enzyme in methanogenesis (MILUCKA *et al.*, 2012; SCHELLER *et al.*, 2016).

Anaerobic methanotrophs use enzymes homologous to those used by methanogens,

prompting questions regarding the directionality of these metabolisms (LLOYD *et al.*,

2011; BEULIG *et al.*, 2018b).

Coupling exploration of methane enzymatic chemistry with isotope geochemistry can

provide powerful insights. Here, we interrogate subsurface methane metabolisms by

precisely determining concentrations of molecules containing two heavy isotopes,

referred to as "clumped" isotopologues (EILER, 2007). The relative proportions of

$^{13}CH_3D$ and $^{12}CH_2D_2$ (reported as $\Delta^{13}CH_3D$ and $\Delta^{12}CH_2D_2$ values relative to a random

distribution of isotopes among all CH_4 isotopologues; see Supplementary Information)

have known temperature-sensitive equilibrium concentrations and thus are sensitive

indicators of reversibility in reactions (PIASECKI *et al.*, 2016). For instance, axenic

cultures of methanogens produce methane with depletions in clumped isotopes relative to

the random distribution that reflect kinetic processes (STOLPER *et al.*, 2015; WANG *et al.*,

2015; DOUGLAS *et al.*, 2016; YOUNG *et al.*, 2017; GRUEN *et al.*, 2018). In some cases, the

clumped isotope composition of environmental microbial methane is consistent with the

axenic cultures, but in others it displays a clumped isotope composition reflective of

equilibration within the environment of formation (WANG *et al.*, 2015; YOUNG *et al.*,

2017). This span in clumped-isotope distributions suggests that enzymatic reactions

associated with methanogenesis and methanotrophy are capable of ranging from

reversible to kinetic. Previous work using one rare isotopologue ($\Delta^{13}CH_3D$) suggests that

that rate of methanogenesis controls the degree of molecular isotopic equilibrium

(STOLPER et al., 2015; WANG et al., 2015). However, the inability to produce equilibrated

methane from axenic cultures has led to speculation that AOM may be responsible for

equilibrated environmental methane (STOLPER et al., 2015; WANG et al., 2016; YOUNG et

al., 2017; GRUEN et al., 2018). The use of a second rare isotopologue, $^{12}CH_2D_2$, aids in

interpreting the causes of intra-methane disequilibrium and suggests that microbial

methanogenesis does not lead to isotopologue equilibrium among methane molecules.

Here, we examine microbial methane from the Bornholm Basin (Baltic Sea; see

Supplementary Information) to test the hypothesis that enzymatic-back reaction during

AOM drives methane to thermodynamic equilibrium. We compare methane clumped

isotope composition to geochemical profiles (ANDRÉN et al., 2015; EGGER et al., 2017)

and present-day rates of methanogenesis and methanotrophy calculated with a

multicomponent diagenetic model (DIJKSTRA et al., 2018) to determine parameters

controlling the isotopic composition of methane (Fig 1A-D). Methane $\delta^{13}C$ and δD,

$\delta^{13}C_{DIC,}$ $\delta^{13}C_{TOC}$ and δD_{H2O} vary non-monotonically over 30 meters below seafloor (Fig.

1H-G, Fig. S1) while $\Delta^{13}CH_3D$ and $\Delta^{12}CH_2D_2$ values increase with depth (Fig. 1E-F).

Comparisons of the bulk isotopic composition of methane to carbon and hydrogen

reservoirs at this location are complicated by the change from limnic to marine deposition

(EGGER et al., 2017). For instance, the hydrogen isotopes in methane are seemingly not

equilibrated with porewater that changes by >100‰ in this interval (Fig. S1), obscuring

the history of this reservoir. Instead, we will focus on the intra-molecular isotopic

equilibrium of the methane itself using clumped isotopes, thus sidestepping the

complications in bulk isotope ratios. We introduce $\xi_{^{12}CH_2D_2}$, a parameter that quantifies

the degree of thermodynamic disequilibrium for a methane system. $\xi_{^{12}CH_2D_2}$ is analogous

to the symbol for the reaction progress variable ($\xi_{^{12}CH_2D_2} = \Delta^{12}CH_2D_{2,\text{ measured}} - \Delta^{12}CH_2D_{2,}$

$_\text{equilibrium,}$ see Supplementary Information). Values for $\xi_{^{12}CH_2D_2}$ in upper sediments as low

as -7.5‰ are significantly different (>6σ) from 0‰ (*i.e.* thermodynamic equilibrium)

which is approached with depth (Fig. 2). In order to determine how methanogenesis and

methanotrophy influence this transition, we consider evidence for and against

reversibility in each metabolism.

Methanogenesis in intra-molecular isotopic thermodynamic equilibrium requires

reversibility at the final hydrogen-addition step. Trace methane oxidation (TMO) does

occur during methanogenesis, but altering the concentration of H_2 or terminal electron

acceptors necessary for oxidizing methane does not increase TMO and has never been

shown to consume greater than ~3% of the CH_4 produced (MORAN *et al.*, 2005; MORAN

et al., 2007). Although some of the first steps of methanogenesis may be reversible

(VALENTINE *et al.*, 2004), the final transfer of the methyl group to coenzyme M and the

subsequent final hydrogen addition are believed to be irreversible (GÄRTNER *et al.*, 1994;

THAUER, 2011). Work on determining the biochemical pathways of Fe (III) reducing

ANME has so far required genetically modifying a methanogen, *M. acetivorans* with the

Mcr from an ANME-2 group organism because unmodified *M. acetivorans* are unable to

be cultured on methane without the modified Mcr (Soo *et al.*, 2016; Yan *et al.*, 2018).

In contrast, AOM is known to have an enzymatic back reaction that suggests the first step

of anaerobic methanotrophy is partially reversible, providing a potential mechanism for

equilibrating methane isotopologues with environmental temperatures.   This back

reaction is sensitive to concentrations of the terminal electron acceptor SO_4^{2-} (Holler *et*

al., 2011; Yoshinaga *et al.*, 2014).   At high concentrations of SO_4^{2-}, the back reaction

produces 3-7% of the CH_4 consumed by AOM, and at low concentrations of SO_4^{2-} (below

0.5 mM), back reaction can produce as much as 78% of the CH_4 consumed by AOM

(Yoshinaga *et al.*, 2014; Timmers *et al.*, 2017). Intracellular exposure of methane to

Mcr can lead to isotope exchange catalysis during bond rupture and reformation

(Marlow *et al.*, 2017), and methane that diffuses back out of the cell could move an

environmental reservoir of methane towards thermodynamic equilibrium. Using

estimates of AOM rates from Dijkstra et al., (2018), we calculate the timescale of this

equilibration (see Supplemental Information) to be on the order of 10^3-10^4 years,

consistent with the estimated age of methane at 20 MCD of <~8000 years [Fig. S1-S2].

Such timescales would vary in environments with differing rates of AOM.

Our findings suggest that when terminal electron acceptors are limiting, equilibrium bond

ordering may be used to identify AOM activity even when the bulk isotopic composition

of reservoir material is changing and ambiguous. Methane from Kidd Creek Mine

(Ontario, Canada) also shows methane isotopologues transitioning with time from non-

equilibrated to equilibrated (Fig. 3), suggesting that as energy-rich fractures open, AOM

consumes the abiotically produced methane and drives it towards thermodynamic

equilibrium (YOUNG *et al.*, 2017).

Methane driven to equilibrium by AOM has a distinct $\Delta^{13}CH_3D$ and $\Delta^{12}CH_2D_2$

composition when compared to methane produced through other known mechanisms

(thermogenesis, microbial methanogenesis and abiotic methanogenesis) (Fig. 3). AOM

metabolisms have been shown to be active from -1°C to 70°C (NIEMANN *et al.*, 2006;

HOLLER *et al.*, 2011), a range of temperatures that is unique and distinguishable from

equilibrated thermogenic methane forming in the 100°C to 250°C gas window. Abiotic

methane may form also form at low temperatures, yet even when it is apparently

equilibrated in $\Delta^{13}CH_3D$, it may exhibit large depletions in $\Delta^{12}CH_2D_2$ that distinguish it

from methane that has undergone exchange catalysis during AOM. We suggest that

when combined (as in the $\xi_{^{12}CH_2D_2}$ parameter), $\Delta^{12}CH_2D_2$ and $\Delta^{13}CH_3D$ values are

sensitive indicators of the degree of thermodynamic equilibrium that may be useful in

determining the role of enzymatic back-flux during AOM in the global methane cycle.

Figure 1: Geochemical profiles (±2σ) from Bornholm Basin from 3-35 MCD.

Horizontal grey bar denotes sediments deposited during lacustrine conditions overlain

with sediments deposited during brackish marine conditions. Samples were not recovered

above the shallow SMTZ. Pink dashed lines denote a shallow (3.3 MCD) and deep (19.8

MCD) SMTZ; only the shallow SMTZ is shown in the bottom panel for clarity. Dark

grey vertical bars in **E** and **F** denote equilibrium isotopologue compositions for the

subsurface temperature of 7.8 ± 0.6°C. Modelled CH_4 production, Fe-AOM and SO_4-

AOM rates are shown in panel **D** (DIJKSTRA *et al.*, 2018). CH_4 production (purple) and

SO_4-AOM (gold) rates correspond with the upper x-axis and Fe-AOM rates (red)

correspond with the lower x-axis. SO_4-AOM rates above the upper SMTZ are calculated

using bottom-water SO_4 concentrations of 15.0 mmol. Additional porewater data shown

in Supplemental Information.

169
170 **Figure 2:** $\xi_{^{12}CH_2D_2}$ values ($\pm 2\sigma$) are the most negative at shallow MCD, indicating the

171 greatest departure from equilibrium and approach zero (violet dashed line) with

172 increasing MCD, indicating an approach towards intra-species thermodynamic

173 equilibrium. ΔG values suggest either metabolism is capable of reversibility (See

174 Supplemental Information).

Figure 3: $\Delta^{12}CH_2D_2$ is plotted versus $\Delta^{13}CH_3D$. The solid black line represents theoretical thermodynamic equilibrium abundances (with dots representing 100°C increments from 0-1000°C). Methane produced by thermogenesis, high-temperature abiotic reactions, microbial methanogenesis and low-temperature abiotic reactions inhabit unique zones in double-isotopologue space. AOM, equilibrating through exchange catalysis during enzymatic back reaction, also inhibits a unique zone: low-temperature intra-species thermodynamic equilibrium. Bornholm Basin data (magenta diamonds) span between the methanogenesis and AOM zones, while data from Kidd Creek Mine (black to white symbols)(YOUNG *et al.*, 2017) span from the abiotic to AOM zone. See Supplemental Information for additional sample description.

References

Andrén, T., Jorgensen, B., Cotterill, C., Green, S., Andrén, E., Ash, J., Bauersachs, T., Cragg, B., Fanget, A., Fehr, A. (2015) Expedition 347 Baltic Sea Paleoenvironment. *Proceedings of the Integrated Ocean Drilling Program* 347, 1-66.

Beal, E.J., House, C.H., Orphan, V.J. (2009) Manganese-and iron-dependent marine methane oxidation. *Science* 325, 184-187.

Beulig, F., Røy, H., Glombitza, C., Jørgensen, B. (2018a) Control on rate and pathway of anaerobic organic carbon degradation in the seabed. *Proceedings of the National Academy of Sciences* 115, 367-372.

Beulig, F., Røy, H., McGlynn, S., Jørgensen, B. (2018b) Cryptic CH4 cycling in the sulfate-methane transition of marine sediments apparently mediated by ANME-1 archaea. *The ISME journal*.

Boetius, A., Ravenschlag, K., Schubert, C.J., Rickert, D., Widdel, F., Gieseke, A., Amann, R., Jørgensen, B.B., Witte, U., Pfannkuche, O. (2000) A marine microbial consortium apparently mediating anaerobic oxidation of methane. *Nature* 407, 623-626.

Cai, C., Leu, A.O., Xie, G.-J., Guo, J., Feng, Y., Zhao, J.-X., Tyson, G.W., Yuan, Z., Hu, S. (2018) A methanotrophic archaeon couples anaerobic oxidation of methane to Fe (III) reduction. *The ISME journal*, 1.

Conrad, R. (2005) Quantification of methanogenic pathways using stable carbon isotopic signatures: a review and a proposal. *Organic Geochemistry* 36, 739-752.

Dijkstra, N., Hagens, M., Egger, M., Slomp, C.P. (2018) Post-depositional vivianite formation alters sediment phosphorus records. *Biogeosciences* 15, 861-883.

Douglas, P., Stolper, D., Smith, D., Anthony, K.W., Paull, C., Dallimore, S., Wik, M., Crill, P.M., Winterdahl, M., Eiler, J. (2016) Diverse origins of Arctic and Subarctic methane point source emissions identified with multiply-substituted isotopologues. *Geochimica Et Cosmochimica Acta* 188, 163-188.

Egger, M., Hagens, M., Sapart, C.J., Dijkstra, N., van Helmond, N.A., Mogollón, J.M., Risgaard-Petersen, N., van der Veen, C., Kasten, S., Riedinger, N. (2017) Iron oxide reduction in methane-rich deep Baltic Sea sediments. *Geochimica Et Cosmochimica Acta* 207, 256-276.

Egger, M., Riedinger, N., Mogollón, J.M., Jørgensen, B.B. (2018) Global diffusive fluxes of methane in marine sediments. *Nature Geoscience* 11, 421.

Eiler, J.M. (2007) "Clumped-isotope" geochemistry -- The study of naturally-occurring,
multiply-substituted isotopologues. *Earth and Planetary Science Letters* 262, 309-
327.

Gärtner, P., Weiss, D.S., Harms, U., Thauer, R.K. (1994) N5﹘
Methyltetrahydromethanopterin: Coenzyme M Methyltransferase from
Methanobacterium thermoautotrophicum. *The FEBS Journal* 226, 465-472.

Gruen, D.S., Wang, D.T., Könneke, M., Topçuoğlu, B.D., Stewart, L.C., Goldhammer,
T., Holden, J.F., Hinrichs, K.-U., Ono, S. (2018) Experimental investigation on
the controls of clumped isotopologue and hydrogen isotope ratios in microbial
methane. *Geochimica Et Cosmochimica Acta* 237, 339-356.

Holler, T., Wegener, G., Knittel, K., Boetius, A., Brunner, B., Kuypers, M.M., Widdel, F.
(2009) Substantial 13C/12C and D/H fractionation during anaerobic oxidation of
methane by marine consortia enriched in vitro. *Environmental microbiology*
reports 1, 370-376.

Holler, T., Wegener, G., Niemann, H., Deusner, C., Ferdelman, T.G., Boetius, A.,
Brunner, B., Widdel, F. (2011) Carbon and sulfur back flux during anaerobic
microbial oxidation of methane and coupled sulfate reduction. *Proceedings of the*
National Academy of Sciences 108, E1484-E1490.

Lloyd, K.G., Alperin, M.J., Teske, A. (2011) Environmental evidence for net methane
production and oxidation in putative ANaerobic MEthanotrophic (ANME)
archaea. *Environmental Microbiology* 13, 2548-2564.

Marlow, J.J., Steele, J.A., Ziebis, W., Sheller, S., Case, D., Reynard, L.M., Orphan, V.J.
(2017) Monodeuterated Methane, an Isotopic Tool To Assess Biological
Metabolism Rates. *mSphere* 2, e00309-17.

Milucka, J., Ferdelman, T.G., Polerecky, L., Franzke, D., Wegener, G., Schmid, M.,
Lieberwirth, I., Wagner, M., Widdel, F., Kuypers, M.M. (2012) Zero-valent
sulphur is a key intermediate in marine methane oxidation. *Nature* 491, 541-546.

Moran, J.J., House, C.H., Freeman, K.H., Ferry, J.G. (2005) Trace methane oxidation
studied in several Euryarchaeota under diverse conditions. *Archaea* 1, 303-309.

Moran, J.J., House, C.H., Thomas, B., Freeman, K.H. (2007) Products of trace methane
oxidation during nonmethyltrophic growth by Methanosarcina. *Journal of*
Geophysical Research: Biogeosciences 112.

Niemann, H., Lösekann, T., De Beer, D., Elvert, M., Nadalig, T., Knittel, K., Amann, R.,
Sauter, E.J., Schlüter, M., Klages, M. (2006) Novel microbial communities of the
Haakon Mosby mud volcano and their role as a methane sink. *Nature* 443, 854-
858.

Piasecki, A., Sessions, A., Peterson, B., Eiler, J. (2016) Prediction of equilibrium
distributions of isotopologues for methane, ethane and propane using density
functional theory. *Geochimica Et Cosmochimica Acta* 190, 1-12.

Regnier, P., Dale, A.W., Arndt, S., LaRowe, D., Mogollón, J., Van Cappellen, P. (2011)
Quantitative analysis of anaerobic oxidation of methane (AOM) in marine
sediments: a modeling perspective. *Earth-Science Reviews* 106, 105-130.

Scheller, S., Yu, H., Chadwick, G.L., McGlynn, S.E., Orphan, V.J. (2016) Artificial
electron acceptors decouple archaeal methane oxidation from sulfate reduction.
Science 351, 703-707.

Soo, V.W., McAnulty, M.J., Tripathi, A., Zhu, F., Zhang, L., Hatzakis, E., Smith, P.B.,
Agrawal, S., Nazem-Bokaee, H., Gopalakrishnan, S. (2016) Reversing
methanogenesis to capture methane for liquid biofuel precursors. *Microbial cell*
factories 15, 11.

Stolper, D.A., Martini, A.M., Clog, M., Douglas, P.M., Shusta, S.S., Valentine, D.L.,
Sessions, A.L., Eiler, J.M. (2015) Distinguishing and understanding thermogenic
and biogenic sources of methane using multiply substituted isotopologues.
Geochimica Et Cosmochimica Acta 161, 219-247.

Thauer, R.K. (2011) Anaerobic oxidation of methane with sulfate: on the reversibility of
the reactions that are catalyzed by enzymes also involved in methanogenesis from
CO_2. *Current Opinion in Microbiology* 14, 292-299.

Timmers, P.H., Welte, C.U., Koehorst, J.J., Plugge, C.M., Jetten, M.S., Stams, A.J.
(2017) Reverse Methanogenesis and respiration in methanotrophic archaea.
Archaea 2017.

Valentine, D.L., Chidthaisong, A., Rice, A., Reeburgh, W.S., Tyler, S.C. (2004) Carbon
and hydrogen isotope fractionation by moderately thermophilic methanogens.
Geochimica Et Cosmochimica Acta 68, 1571-1590.

Waite, J.H., Glein, C.R., Perryman, R.S., Teolis, B.D., Magee, B.A., Miller, G., Grimes,
J., Perry, M.E., Miller, K.E., Bouquet, A. (2017) Cassini finds molecular
hydrogen in the Enceladus plume: Evidence for hydrothermal processes. *Science*
356, 155-159.

Wang, D.T., Gruen, D.S., Lollar, B.S., Hinrichs, K.-U., Stewart, L.C., Holden, J.F.,
Hristov, A.N., Pohlman, J.W., Morrill, P.L., Könneke, M. (2015) Nonequilibrium
clumped isotope signals in microbial methane. *Science* 348, 428-431.

Wang, D.T., Welander, P.V., Ono, S. (2016) Fractionation of the methane isotopologues
13 CH 4, 12 CH 3 D, and 13 CH 3 D during aerobic oxidation of methane by
Methylococcus capsulatus (Bath). *Geochimica Et Cosmochimica Acta* 192, 186-
202.

Webster, C.R., Mahaffy, P.R., Atreya, S.K., Flesch, G.J., Mischna, M.A., Meslin, P.-Y.,
Farley, K.A., Conrad, P.G., Christensen, L.E., Pavlov, A.A. (2015) Mars methane
detection and variability at Gale crater. *Science* 347, 415-417.

Whiticar, M.J. (1999) Carbon and hydrogen isotope systematics of bacterial formation
and oxidation of methane. *Chemical Geology* 161, 291-314.

Yan, Z., Joshi, P., Gorski, C.A., Ferry, J.G. (2018) A biochemical framework for
anaerobic oxidation of methane driven by Fe (III)-dependent respiration. *Nature
Communications* 9.

Yoshinaga, M.Y., Holler, T., Goldhammer, T., Wegener, G., Pohlman, J.W., Brunner, B.,
Kuypers, M.M.M., Hinrichs, K.U., Elvert, M. (2014) Carbon isotope equilibration
during sulphate-limited anaerobic oxidation of methane. *Nature Geoscience* 7,
190-194.

Young, E., Kohl, I., Lollar, B.S., Etiope, G., Rumble, D., Li, S., Haghnegahdar, M.,
Schauble, E., McCain, K., Foustoukos, D. (2017) The relative abundances of
resolved $^{12}CH_2D_2$ and $^{13}CH_3D$ and mechanisms controlling isotopic bond ordering
in abiotic and biotic methane gases. *Geochimica Et Cosmochimica Acta* 203, 235-
264.
314