Exchange catalysis during anaerobic methanotrophy revealed by ^{12}CH$_2$D$_2$ &

^{13}CH$_3$D in methane

Jeanine L. Ash[1*], Matthias Egger[2,#], Tina Treude[3,4], Issaku Kohl[3], Barry Cragg[4], R. John

Parkes[4], Caroline P. Slomp[5], Barbara Sherwood Lollar[6], Edward D. Young[3*]

[1]Department of Earth, Environmental and Planetary Sciences, Rice University, Houston

Texas, USA

[2]Center for Geomicrobiology, Aarhus University, Aarhus, Denmark

[3]Department of Earth, Planetary, and Space Sciences, UCLA, Los Angeles CA, USA

[4]School of Earth and Ocean Sciences, Cardiff University, Cardiff, UK.

[5]Department of Earth Sciences-Geochemistry, Utrecht University, Utrecht, The

Netherlands

[6]Department of Earth Sciences, University of Toronto, Toronto Ontario, Canada

[#]now at The Ocean Cleanup Foundation, Delft, The Netherlands

*correspondence to jeanine.ash@rice.edu, eyoung@epss.ucla.edu

The anaerobic oxidation of methane (AOM) is a crucial component of the methane cycle,

but its enzymatic versatility under environmental conditions remains poorly understood.

We use sediment samples collected during IODP Expedition 347 to the Baltic Sea to

show that relative abundances of ^{12}CH$_2$D$_2$ and ^{13}CH$_3$D molecules in methane gas trace the

reversibility of methyl-coenzyme M reductase during AOM by driving methane towards

internal, thermodynamic isotopic equilibrium. These data suggest that $^{12}CH_2D_2$ and

$^{13}CH_3D$ together can identify the influence of methanotrophy in environments where

conventional bulk isotope ratios are ambiguous, and these findings may lead to new

insights regarding the global significance of enzymatic back-flux in the methane cycle.

____

Recent discoveries of dynamically varying methane in the Martian atmosphere and

molecular hydrogen in the plumes of Enceladus have renewed excitement about the role

of reduced gases of astrobiological potential and the possibility that habitability may not

be unique to Earth (WEBSTER *et al.*, 2015; WAITE *et al.*, 2017).   Methane can be

produced in a number of biological and abiotic reactions, precluding our ability to

determine unequivocally its sources on extraterrestrial bodies (MCCOLLOM AND

SEEWALD, 2007; ETIOPE AND SHERWOOD LOLLAR, 2013). Even on Earth, where direct

measurements of methane and substrate characteristics are possible, distinguishing

microbial from other methanogenesis pathways can be problematic (WHITICAR, 1990;

MARTINI *et al.*, 1996; MCCOLLOM AND SEEWALD, 2006; POHLMAN *et al.*, 2009).

Additional complexity in sourcing methane is introduced by methanotrophy, the aerobic

or anaerobic consumption of methane by microbes. This process can alter parameters

that are traditionally useful in sourcing methane such as bulk isotope ratios (WHITICAR,

1999). Perhaps nowhere is this complexity greater than in the deep biosphere, where

linking biogeochemistry to metabolisms remains a long-standing challenge compounded

by slow rates of reaction (EDWARDS *et al.*, 2012; LAU *et al.*, 2016; MARLOW *et al.*, 2017).

Coupling exploration of enzymatic chemistry that drives these cycles with isotope

geochemistry can provide powerful insights. Here, we focus on AOM, estimated to

consume ~90% of methane produced in marine sediments (KNITTEL AND BOETIUS, 2009).

AOM is traditionally thought to be carried out by a consortium of methane-oxidizing

archaea   and   sulphate-reducing   bacteria   performing   the   net   reaction

$CH_4 + SO_4^{2-} \rightarrow HCO_3^- + HS^- + H_2O$ (BOETIUS *et al.*, 2000). Evidence suggests that these

processes can be decoupled (MILUCKA *et al.*, 2012; SCHELLER *et al.*, 2016), and metal

oxides may also act as terminal electron acceptors (BEAL *et al.*, 2009). In all cases, the

first step of AOM involves interaction between methane and modified methyl-coenzyme

M reductase (*mcr*), the terminal enzyme in methanogenesis (KRÜGER *et al.*, 2003;

SCHELLER *et al.*, 2010; TIMMERS *et al.*, 2017). Methanogenesis and AOM are linked via

this homologous enzyme, prompting questions regarding the reversibility of these

metabolisms (LLOYD *et al.*, 2011).

Developments in isotope-ratio mass spectrometry permit the interrogation of these

metabolisms by precisely determining concentrations of molecules containing two heavy

isotopes, referred to as "clumped" isotopologues. The relative proportions of $^{13}CH_3D$ and

$^{12}CH_2D_2$ (reported as $\Delta^{13}CH_3D$ and $\Delta^{12}CH_2D_2$ values relative to a random distribution of

isotopes among all CH_4 isotopologues; see Supplementary Information) have known

temperature-sensitive equilibrium concentrations and thus are sensitive indicators of

reversibility   in   reactions.   Microbial   methane   displays   clumped   isotopologue

concentrations that range from those consistent with low-temperature equilibration within

the environment of formation (INAGAKI *et al.*, 2015) to those with depletions in $\Delta^{13}CH_3D$

and $\Delta^{12}CH_2D_2$ reflecting kinetic processes (WANG *et al.*, 2015; YOUNG *et al.*, 2017). This

span in clumped-isotope distributions suggests that enzymatic reactions associated with

methanogenesis and methanotrophy are capable of ranging from reversible to kinetic.

Previous work using one rare isotopologue ($\Delta^{13}CH_3D$) suggests that that rate of

methanogenesis controls the degree of molecular isotopic equilibrium (STOLPER *et al.*,

2015; WANG *et al.*, 2015).

Here, we test the hypothesis that enzymatic-back reaction during AOM drives methane to

thermodynamic equilibrium by measuring the isotopic composition of sedimentary

methane from Bornholm Basin (Baltic Sea), where methane produced in sediments

deposited under brackish-marine conditions diffuses down into lacustrine sediments (see

Supplementary Information). Metagenomic analyses reveal that anaerobic methanotrophs

(ANME-1) and relatives of the methanogen *Methanohalobium evestigatum* are present,

and methyl coenzyme reductase subunit A (*mcr*A) exists throughout the interval

(MARSHALL *et al.*, 2017). We compare these data to geochemical profiles (ANDRÉN *et al.*,

2015; EGGER *et al.*, 2017) and present-day rates of methanogenesis and methanotrophy

calculated with a multicomponent diagenetic model (DIJKSTRA *et al.*, 2018) to determine

parameters controlling the isotopic composition of methane.

Methane concentrations decrease from 10.1 to 1.4 mM over 3.5 to 30.0 meters composite

depth [MCD, akin to meters below seafloor (ANDRÉN *et al.*, 2015)] and sulphate

concentrations are <0.5 mM throughout the profile. Two sulphate methane transition

zones (SMTZ) are noted at 3.3 MCD and 19 MCD (Fig. 1A-B). In sediments that are

sulphate poor but rich in iron oxides (*e.g.* 10-20 MCD [Fig 1C]), metal oxides may play a

role in methane consumption (SIVAN *et al.*, 2011). At 3.3 MCD, methane $\delta^{13}C$ is -

76.16‰, then increases to a maximum of -71.81‰ by 8.1 MCD, followed by a decrease

downcore to -75.59‰ (Fig. 1G). Methane δD at 3.3 MCD is the most negative hydrogen

isotope value throughout the core at -244.0‰ and downcore, δD in methane increases to -

238.2‰ with a small reversal present between 10-12 MCD before the trend of increasing

δD resumes (Fig 1H). In contrast, $\Delta^{13}CH_3D$ and $\Delta^{12}CH_2D_2$ values show trends of

consistent increase downcore (Fig. 1E-F). At the shallowest measured depth of 3.5 MCD,

values of 4.7‰ and 7.9‰ for $\Delta^{13}CH_3D$ and $\Delta^{12}CH_2D_2$, respectively, are lower than

predicted equilibrium values of 6.3‰ and 22.3‰, respectively, for the average

subsurface temperature over this interval of 7.8±0.6°C. Activity measurements imply that

methanogenesis rates decrease with depth (Table S1). Modelling based on porewater and

solid phase profiles shows a methane production peak at 2.7 MCD that decreases

downcore and peak Fe-AOM consumption at 12 MCD [Fig. 1D]. The SMTZ at 19 MCD

is an additional methane sink not considered in the model by Dijkstra et al., (2018).

Therefore, we consider the present-day Fe-AOM rates derived from modelling as a most

conservative possible estimate for total AOM activity at these depths.

Comparing the measured $\Delta^{12}CH_2D_2$ values to the equilibrium $\Delta^{12}CH_2D_2$ values associated

with measured $\Delta^{13}CH_3D$ values provides a novel parameter capable of demonstrating the

degree of thermodynamic disequilibrium for a methane system. $\xi_{^{12}CH_2D_2}$ is analogous to

the symbol for the reaction progress variable ($\xi_{^{12}CH_2D_2} = \Delta^{12}CH_2D_{2,\ measured} - \Delta^{12}CH_2D_{2,}$

equilibrium, see Supplementary Information). Values for $\xi_{^{12}CH_2D_2}$ in upper sediments as low

as -7.5‰ are clear evidence for disequilibrium, while 0‰ (*i.e.* thermodynamic

equilibrium) is approached with increasing depth (Fig. 2). In order to determine how

methanogenesis and methanotrophy influence this transition, we discuss evidence for

reversibility in each metabolism.

Methanogenesis in intra-species, thermodynamic equilibrium requires reversibility at the

final hydrogen-addition step. If methanogenesis is reversible at this terminal step, then its

directionality should be influenced by concentrations of products and reactants. For

example, in the net reaction $CO_2+4H_2 \rightarrow CH_4+2H_2O$, H_2 concentrations would control

the rate of CH_4 oxidation via the reverse reaction, trace methane oxidation (TMO). TMO

does occur during methanogenesis, but altering the concentration of H_2 or terminal

electron acceptors necessary for oxidizing methane does not increase TMO and has never

been shown to consume greater than 0.5% of the CH_4 produced (MORAN *et al.*, 2005).

Although some of the first steps of methanogenesis may be reversible (VALENTINE *et al.*,

2004), the final transfer of the methyl group to coenzyme M and the subsequent final

hydrogen addition are believed to be irreversible (GÄRTNER *et al.*, 1994; THAUER, 2011).

Moreover, with one exception, all clumped-isotope measurements of methane produced

by axenic lab cultures have shown methane in intra-species isotopic disequilibrium

(STOLPER *et al.*, 2015; WANG *et al.*, 2015; YOUNG *et al.*, 2017), indicating terminal

irreversibility in this metabolism. A thermophilic methanogen grown on an H_2/CO_2

substrate produced methane with a clumped isotope temperature consistent with its

growth conditions (STOLPER *et al.*, 2014), but this result has yet to be reproduced.

On the other hand, AOM has been shown to have an enzymatic back reaction that

suggests the first step of anaerobic methanotrophy is partially reversible, providing a

potential mechanism for equilibrating methane isotopologues with environmental

temperatures. This back reaction is sensitive to concentrations of the terminal electron

acceptor SO_4^{2-} (HOLLER *et al.*, 2011; YOSHINAGA *et al.*, 2014). At high concentrations of

SO_4^{2-}, the back reaction produces 3-7% of the CH_4 consumed by AOM, and at low

concentrations of SO_4^{2-} (below 0.5 mM), back reaction can produce as much as 78% of

the CH_4 consumed by AOM (TIMMERS *et al.*, 2017). Intracellular exposure of methane to

mcr can lead to isotope exchange catalysis during bond rupture and reformation, and

methane that diffuses back out of the cell could move a reservoir towards thermodynamic

equilibrium. Using estimates of AOM rates from Dijkstra et al., (2018), we calculate the

timescale of this equilibration (see Supplemental Information) to be on the order of 10^3-

10^4 years, consistent with the estimated age of methane at 20 MCD of <~8000 years [Fig.

S1-S2]. Such timescales would vary in environments with differing rates of AOM.

Conventional isotopic experiments suggest that AOM back-flux is capable of reversing

the canonical stable isotope trend expected during methanotrophy (YOSHINAGA *et al.*,

2014); during lab incubations where sulphate is limiting, the methane residue is

progressively depleted in heavy isotopes of carbon rather than enriched. Such a reversal

is present in the $\delta^{13}C_{CH_4}$ profile of the downward diffusing methane of Bornholm Basin,

but interpreting such signals can be difficult where methane produced in modern-late

Holocene brackish-marine sediments is transported into glacial-lacustrine sediments with

each sedimentary package having unique carbon and hydrogen reservoirs (EGGER *et al.*,

2017). Our findings suggest that when terminal electron acceptors are limiting,

equilibrium bond ordering may be used to identify AOM activity even when the bulk

isotopic composition of reservoir material is changing and ambiguous. Methane from

Kidd Creek Mine (Ontario, Canada) also shows methane isotopologues transitioning with

time from non-equilibrated to equilibrated (Fig. 3), suggesting that as energy-rich

fractures open, AOM consumes the abiotically produced methane and drives it towards

thermodynamic equilibrium (YOUNG *et al.*, 2017).

Methane driven to equilibrium by AOM has a distinct $\Delta^{13}CH_3D$ and $\Delta^{12}CH_2D_2$

composition when compared to methane produced through other known mechanisms

(thermogenesis, microbial methanogenesis and abiotic methanogenesis) (Fig. 3). AOM

metabolisms have been shown to be active from -1°C to 70°C (NIEMANN *et al.*, 2006;

HOLLER *et al.*, 2011), a range of temperatures that is unique and distinguishable from

equilibrated thermogenic methane forming in the 100°C to 250°C gas window. Abiotic

methane may form also form at low temperatures, yet even when it is apparently

equilibrated in $\Delta^{13}CH_3D$, it may exhibit large depletions in $\Delta^{12}CH_2D_2$ that distinguish it

from methane that has undergone exchange catalysis during AOM. We suggest that

when combined, $\Delta^{12}CH_2D_2$ and $\Delta^{13}CH_3D$ values are sensitive indicators of the degree of

thermodynamic equilibrium that may be useful in determining the role of enzymatic

back-flux during AOM in the global methane cycle.

Figure 1: Geochemical profiles (±1σ) from Bornholm Basin from 3-35 MCD. Horizontal grey bar denotes sediments deposited during lacustrine conditions overlain with sediments deposited during brackish marine conditions. Samples were not recovered above the shallow SMTZ. Pink dashed lines denote the shallow (3.3 MCD) and deep (19.8 MCD) SMTZ; only the shallow SMTZ is shown in the bottom panel for clarity. Dark grey vertical bars in **E** and **F** denote equilibrium isotopologue compositions for the

190   subsurface temperature of $7.8 \pm 0.6°C$. Modelled CH_4 production, Fe-AOM and SO_4-

191   AOM rates are shown in panel **D** (DIJKSTRA *et al.*, 2018). CH_4 production (purple) and

192   SO_4-AOM (gold) rates correspond with the upper x-axis and Fe-AOM rates (red)

193   correspond with the lower x-axis. SO_4-AOM rates above the upper SMTZ are calculated

194   using bottom-water SO_4 concentrations of 15.0 mmol.

Figure 2: $\xi_{^{12}CH_2D_2}$ values ($\pm 1\sigma$) are the most negative at shallow MCD, indicating the greatest departure from equilibrium and approach zero (violet dashed line) with increasing MCD, indicating an approach towards intra-species thermodynamic equilibrium. ΔG values suggest either metabolism is capable of reversibility (See Supplemental Information).

Figure 3: $\Delta^{12}CH_2D_2$ is plotted versus $\Delta^{13}CH_3D$. The solid black line represents

theoretical thermodynamic equilibrium abundances (with dots representing 100°C

increments from 0-1000°C). Methane produced by thermogenesis, high-temperature

abiotic reactions, microbial methanogenesis and low-temperature abiotic reactions inhabit

unique zones in double-isotopologue space. AOM, equilibrating through exchange

catalysis during enzymatic back reaction, also inhibits a unique zone: low-temperature

intra-species thermodynamic equilibrium. Bornholm Basin data (magenta diamonds)

span between the methanogenesis and AOM zones, while data (YOUNG *et al.*, 2017) from

Kidd Creek Mine (black to white symbols) span from the abiotic to AOM zone.

References

Andrén, T., Jorgensen, B., Cotterill, C., Green, S., Andrén, E., Ash, J., Bauersachs, T.,
Cragg, B., Fanget, A., Fehr, A. (2015) Expedition 347 Baltic Sea
Paleoenvironment. *Proceedings of the Integrated Ocean Drilling Program* 347, 1-
66.

Beal, E.J., House, C.H., Orphan, V.J. (2009) Manganese-and iron-dependent marine
methane oxidation. *Science* 325, 184-187.

Boetius, A., Ravenschlag, K., Schubert, C.J., Rickert, D., Widdel, F., Gieseke, A.,
Amann, R., Jørgensen, B.B., Witte, U., Pfannkuche, O. (2000) A marine
microbial consortium apparently mediating anaerobic oxidation of methane.
Nature 407, 623-626.

Dijkstra, N., Hagens, M., Egger, M., Slomp, C.P. (2018) Post-depositional vivianite
formation alters sediment phosphorus records. *Biogeosciences* 15, 861-883.

Edwards, K.J., Becker, K., Colwell, F. (2012) The deep, dark energy biosphere:
intraterrestrial life on earth. *Annual Review of Earth and Planetary Sciences* 40,
551-568.

Egger, M., Hagens, M., Sapart, C.J., Dijkstra, N., van Helmond, N.A., Mogollón, J.M.,
Risgaard-Petersen, N., van der Veen, C., Kasten, S., Riedinger, N. (2017) Iron
oxide reduction in methane-rich deep Baltic Sea sediments. *Geochimica Et*
Cosmochimica Acta 207, 256-276.

Etiope, G., Sherwood Lollar, B. (2013) Abiotic Methane on Earth. *Reviews of Geophysics*
51, 276-299.

Gärtner, P., Weiss, D.S., Harms, U., Thauer, R.K. (1994) N5‐
Methyltetrahydromethanopterin: Coenzyme M Methyltransferase from
Methanobacterium thermoautotrophicum. *The FEBS Journal* 226, 465-472.

Holler, T., Wegener, G., Niemann, H., Deusner, C., Ferdelman, T.G., Boetius, A.,
Brunner, B., Widdel, F. (2011) Carbon and sulfur back flux during anaerobic
microbial oxidation of methane and coupled sulfate reduction. *Proceedings of the*
National Academy of Sciences 108, E1484-E1490.

Inagaki, F., Hinrichs, K.-U., Kubo, Y., Bowles, M.W., Heuer, V.B., Hong, W.-L.,
Hoshino, T., Ijiri, A., Imachi, H., Ito, M. (2015) Exploring deep microbial life in
coal-bearing sediment down to~ 2.5 km below the ocean floor. *Science* 349, 420-
424.

Knittel, K., Boetius, A. (2009) Anaerobic oxidation of methane: progress with an
unknown process. *Annual review of microbiology* 63, 311-334.

Krüger, M., Meyerdierks, A., Glöckner, F.O., Amann, R., Widdel, F., Kube, M.,
Reinhardt, R., Kahnt, J., Böcher, R., Thauer, R.K. (2003) A conspicuous nickel
protein in microbial mats that oxidize methane anaerobically. *Nature* 426, 878-
881.

Lau, M.C.Y., Kieft, T.L., Kulouo, O., Linage-Alvarez, B., van Heerden, E., Lindsay,
M.R., Magnabosco, C., Wang, W., Wiggins, J.B., Guo, L., Perlman, D.H., Kyin,
S., Wei, S., Li, L., Sherwood Lollar, B., Onstott, T. (2016) An oligotrophic deep-
subsurface community dependent on syntrophy is dominated by sulfur-driven
autotrophic denitrifiers. *Proceedings of the National Academy of Sciences* 113,
E7927-E7936.

Lloyd, K.G., Alperin, M.J., Teske, A. (2011) Environmental evidence for net methane
production and oxidation in putative ANaerobic MEthanotrophic (ANME)
archaea. *Environmental Microbiology* 13, 2548-2564.

Marlow, J.J., Steele, J.A., Ziebis, W., Sheller, S., Case, D., Reynard, L.M., Orphan, V.J.
(2017) Monodeuterated Methane, an Isotopic Tool To Assess Biological
Metabolism Rates. *mSphere* 2, e00309-17.

Marshall, I.P., Karst, S.M., Nielsen, P.H., Jørgensen, B.B. (2017) Metagenomes from
deep Baltic Sea sediments reveal how past and present environmental conditions
determine microbial community composition. *Marine genomics*.

Martini, A.M., Budai, J.M., Walter, L.M., Schoell, M. (1996) Microbial generation of
economic accumulations of methane within a shallow organic-rich shale. *Nature*
383, 155-158.

McCollom, T.M., Seewald, J.S. (2006) Carbon isotope composition of organic
compounds produced by abiotic synthesis under hydrothermal conditions. *Earth*
and Planetary Science Letters 243, 74-84.

McCollom, T.M., Seewald, J.S. (2007) Abiotic synthesis of organic compounds in deep-
sea hydrothermal environments. *Chemical Reviews* 107, 382-401.

Milucka, J., Ferdelman, T.G., Polerecky, L., Franzke, D., Wegener, G., Schmid, M.,
Lieberwirth, I., Wagner, M., Widdel, F., Kuypers, M.M. (2012) Zero-valent
sulphur is a key intermediate in marine methane oxidation. *Nature* 491, 541-546.

Moran, J.J., House, C.H., Freeman, K.H., Ferry, J.G. (2005) Trace methane oxidation
studied in several Euryarchaeota under diverse conditions. *Archaea* 1, 303-309.

Niemann, H., Lösekann, T., De Beer, D., Elvert, M., Nadalig, T., Knittel, K., Amann, R.,
Sauter, E.J., Schlüter, M., Klages, M. (2006) Novel microbial communities of the
Haakon Mosby mud volcano and their role as a methane sink. *Nature* 443, 854-
858.

Pohlman, J., Kaneko, M., Heuer, V., Coffin, R., Whiticar, M. (2009) Methane sources
and production in the northern Cascadia margin gas hydrate system. *Earth and*
Planetary Science Letters 287, 504-512.

Scheller, S., Goenrich, M., Boecher, R., Thauer, R.K., Jaun, B. (2010) The key nickel
enzyme of methanogenesis catalyses the anaerobic oxidation of methane. *Nature*
465, 606.

Scheller, S., Yu, H., Chadwick, G.L., McGlynn, S.E., Orphan, V.J. (2016) Artificial
electron acceptors decouple archaeal methane oxidation from sulfate reduction.
Science 351, 703-707.

Sivan, O., Adler, M., Pearson, A., Gelman, F., Bar-Or, I., John, S.G., Eckert, W. (2011)
Geochemical evidence for iron‐mediated anaerobic oxidation of methane.
Limnology and Oceanography 56, 1536-1544.

Stolper, D.A., Martini, A.M., Clog, M., Douglas, P.M., Shusta, S.S., Valentine, D.L.,
Sessions, A.L., Eiler, J.M. (2015) Distinguishing and understanding thermogenic
and biogenic sources of methane using multiply substituted isotopologues.
Geochimica Et Cosmochimica Acta 161, 219-247.

Stolper, D.A., Sessions, A.L., Ferreira, A.A., Neto, E.V.S., Schimmelmann, A., Shusta,
S.S., Valentine, D.L., Eiler, J.M. (2014) Combined ^{13}C-D and D-D clumping in
methane: Methods and preliminary results. *Geochimica Et Cosmochimica Acta*
126, 169-191.

Thauer, R.K. (2011) Anaerobic oxidation of methane with sulfate: on the reversibility of
the reactions that are catalyzed by enzymes also involved in methanogenesis from
CO_2. *Current Opinion in Microbiology* 14, 292-299.

Timmers, P.H., Welte, C.U., Koehorst, J.J., Plugge, C.M., Jetten, M.S., Stams, A.J.
(2017) Reverse Methanogenesis and respiration in methanotrophic archaea.
Archaea 2017.

Valentine, D.L., Chidthaisong, A., Rice, A., Reeburgh, W.S., Tyler, S.C. (2004) Carbon
and hydrogen isotope fractionation by moderately thermophilic methanogens.
Geochimica Et Cosmochimica Acta 68, 1571-1590.

Waite, J.H., Glein, C.R., Perryman, R.S., Teolis, B.D., Magee, B.A., Miller, G., Grimes,
J., Perry, M.E., Miller, K.E., Bouquet, A. (2017) Cassini finds molecular
hydrogen in the Enceladus plume: Evidence for hydrothermal processes. *Science*
356, 155-159.

Wang, D.T., Gruen, D.S., Lollar, B.S., Hinrichs, K.-U., Stewart, L.C., Holden, J.F.,
Hristov, A.N., Pohlman, J.W., Morrill, P.L., Könneke, M. (2015) Nonequilibrium
clumped isotope signals in microbial methane. *Science* 348, 428-431.

Webster, C.R., Mahaffy, P.R., Atreya, S.K., Flesch, G.J., Mischna, M.A., Meslin, P.-Y.,
Farley, K.A., Conrad, P.G., Christensen, L.E., Pavlov, A.A. (2015) Mars methane
detection and variability at Gale crater. *Science* 347, 415-417.

Whiticar, M.J. (1990) A geochemial perspective of natural gas and atmospheric methane.
Organic Geochemistry 16, 531-547.

Whiticar, M.J. (1999) Carbon and hydrogen isotope systematics of bacterial formation
and oxidation of methane. *Chemical Geology* 161, 291-314.

Yoshinaga, M.Y., Holler, T., Goldhammer, T., Wegener, G., Pohlman, J.W., Brunner, B.,
Kuypers, M.M.M., Hinrichs, K.U., Elvert, M. (2014) Carbon isotope equilibration
during sulphate-limited anaerobic oxidation of methane. *Nature Geoscience* 7,
190-194.

Young, E., Kohl, I., Lollar, B.S., Etiope, G., Rumble, D., Li, S., Haghnegahdar, M.,
Schauble, E., McCain, K., Foustoukos, D. (2017) The relative abundances of
resolved $^{12}CH_2D_2$ and $^{13}CH_3D$ and mechanisms controlling isotopic bond ordering
in abiotic and biotic methane gases. *Geochimica Et Cosmochimica Acta* 203, 235-
264.
339