Title

Modelling cancer immunomodulation using epithelial organoid cultures

Authors

Yotam E. Bar-Ephraim[1,2,8], Kai Kretzschmar[1,2,8], Priyanca Asra[1,2], Evelien de Jongh[1,2], Kim E.

Boonekamp[1,2], Jarno Drost[3], Joost van Gorp[4], Apollo Pronk[5], Niels Smakman[5], Inez J. Gan[6], Zsolt

Sebestyén[6], Jürgen Kuball[6], Robert G.J. Vries[7], Hans Clevers[1,2,3*]

[1]Oncode Institute, Hubrecht Institute, Royal Netherlands Academy for Arts and Sciences (KNAW)

and University Medical Centre (UMC) Utrecht, the Netherlands

[2]UMC Utrecht, Cancer Genomics Netherlands, Utrecht, the Netherlands

[3]Princess Máxima Centre for Paediatric Oncology, Utrecht, the Netherlands

[4]Department of Pathology, Diakonessenhuis hospital, Utrecht, the Netherlands

[5]Department of Surgery, Diakonessenhuis hospital, Utrecht, the Netherlands

[6]Laboratory of Translational Immunology, Department of Immunology, UMC, Utrecht, the

Netherlands

[7]Foundation Hubrecht Organoid Technology (HUB), Utrecht, the Netherlands

[8]These authors contributed equally: Yotam E. Bar-Ephraim, Kai Kretzschmar

*Corresponding author, e-mail: h.clevers@hubrecht.eu

Introduction

Colorectal cancer (CRC) is among the most common cancers worldwide[1]. While early CRC stages

are highly treatable by surgical removal, later stages are usually incurable[2]. CRC arises through a

multi-step process from small lesions of the epithelium of the large intestine. These lesions grow

into adenomas with low grade dysplasia that progress into high grade dysplasia, eventually giving

rise to infiltrating carcinomas[3]. Genetic mutations in signalling pathways such as the canonical

Wnt signalling are the molecular basis of CRC[4]. However, the interaction of the tumour with its

microenvironment is another critical hallmark[5]. Cancer cells remodel their microenvironment (e.g.

fibroblasts, the vasculature and immune cells) to support tumour growth[6]. Infiltrating immune cells

(ICs) such as cytotoxic T cells (CTLs) or macrophages play a crucial role by generating different

immune responses such as anti-tumour cytotoxicity (the former) or tumour-promoting chronic

inflammation (the latter)[7]. As such, escape from the surveilling immune system has been

recognised as one of the hallmarks of cancer[5]. Cancer cells undergo a process called

immunoediting and silence anti-tumour responses, for example, by preventing T-cell activation

through stimulation of inhibitory cell surface receptors such as CTL-associated antigen (CTLA)-

4 or programmed death (PD)1[6,8]. Overcoming this active immunomodulation by tumour cells has

become a major therapeutic target[9]. However, tumour heterogeneity, such as differential CTL

infiltration, could influence therapeutic efficiency of anti-tumour drugs by mediating drug

resistance[6]. Developing *ex vivo* model systems to characterise the communication of the tumour

with its environment is therefore of great importance. Organoid cultures grown from different

epithelial tissues serve as an excellent tool to study tissue homeostasis and disease[10]. Furthermore,

organoid biobanks of multiple epithelial organ systems have been established and tumor-derived

organoids have successfully been used as platforms for screenings of different drugs to predict

patient response[11]. Here we describe the establishment of a method to model antigen-specific

epithelial-cell killing and cancer immunomodulation and *in vitro* using CRC organoids co-cultured

with CTLs.

Results

CRC organoids express immunomodulatory molecules

We first assessed whether CRC organoids expressed immunomodulatory molecules in established

long-term expanded cultures. Therefore, we compared gene expression of T-cell-specific

immunomodulators in CRC organoids to the expression levels found in normal colon organoids

using a transcriptome dataset generated using our 'living organoid biobank' of CRC patients[12]. On

average, transcription of genes associated with T-cell stimulation such as *TNFSF4* or *TNFSF9* was

not altered in CRC organoids compared to normal colon organoids (Fig. 1a). However, expression

of human leukocyte antigen (HLA) genes *HLA-A* and *HLA-C*, encoding major histocompatibility

complex class (MHC)-I molecules that present antigens to T cells, were significantly

downregulated in CRC organoids (Fig. 1a), a well-described phenomenon found in cancers[13].

Expression of genes associated with inhibition of T-cell function was either significantly

upregulated such as *BTLA*, significantly downregulated such as *CD80, CD86* or LGALS9 or not

altered at all such as *CD274* (encoding PD-L1), *PDCD1LG2* (encoding PD-L2) (Fig. 1a). When

assessing expression levels of immunomodulatory molecules on individual organoids, CRC

organoids largely clustered together showing heterogeneous down regulation of *HLA-A*, *HLA-C*

and *LGALS9* compared to healthy colon organoids (Fig. 1b). However, expression of

immunoinhibitory genes *CD274* and *PDCD1LG2*, for instance, was highly upregulated in some

CRC organoids in comparison to the matched normal colon organoid cultures, reflecting

previously reported preservation of tumour heterogeneity in organoids[12] (Fig. 1b). These molecular

signatures provide a basis for further investigation of tumour immunogenicity and its association

with other characteristics of the tumour.

Four of the most commonly mutated genes in CRC are *APC*, *P53*, *KRAS* and *SMAD4*,

reflecting the stepwise progression of the normal intestinal epithelium into a metastatic

carcinoma[14]. Introduction of these cancer mutations into human intestinal organoid cultures using

clustered regularly interspaced short palindromic repeats (CRISPR)/Cas9 demonstrated that this

process can be mimicked *in vitro* and upon xenotransplantation into mice[15,16]. Using colon

organoids carrying one or more of these cancer mutations, we investigated whether up-regulation

of PD-L1 was associated with a certain mutational status. Additionally, we exposed mutant

organoids and their wild-type control organoid line to Interferon (IFN)-γ, which is secreted by T

cells and can trigger increased expression of immunomodulatory molecules such as PD-L1[17].

Subsequently, we assessed PD-L1 expression by quantitative polymerase chain reaction (qPCR)

and flow cytometry (Fig. 1 c,d). In the absence of IFN-γ, organoids carrying triple ($APC^{KO/KO}$,

$P53^{KO/KO}$, $KRAS^{G12D/+}$) and quadruple mutations ($APC^{KO/KO}$, $P53^{KO/KO}$, $KRAS^{G12D/+}$ and

$SMAD4^{KO/KO}$) showed lower *CD274* gene expression in comparison to control wild-type organoids

(Fig. 1 c). Overall, PD-L1 expression was low in untreated organoid lines (Fig. 1 c,d). However,

PD-L1 expression was dramatically upregulated in IFN-γ-treated organoids both on transcript and

protein level (Fig. 1 c,d). These data demonstrate that CRC organoids express immunomodulators

and that this expression is regulated in a similar way as previously shown for tissue *in vivo*.

CRC organoids as tools for assessment of antigen specific killing by CD8+ T cells

We next aimed at establishing a co-culture system for CRC organoids and CTLs to model antigen-

specific killing of tumour cells *in vitro*. For this, we used αβ T cells carrying a transgenic T-cell

receptor (TCR) recognizing an HLA-A2-restricted Wilms tumour (WT)1-derived peptide[18,19].

Therefore, we first screened CRC organoids from the 'living biobank'[12] as well as newly generated

CRC organoids for HLA-A2 expression using flow cytometry. Interestingly, we found three CRC

organoid lines that showed partial downregulation of HLA-A2 (Supplementary Fig. 1a). We were

able to purify HLA-A2$^+$ and HLA-A2$^-$ CRC organoids and successfully established cultures from

both populations (Fig. 2b). We confirmed stable MHC-I downregulation in HLA-A2$^-$ CRC

organoids, as IFN-γ stimulation did not trigger HLA-A2 re-expression (Supplementary Fig. 1b).

Next, we pulsed these CRC organoid lines with WT1 peptide and, subsequently, co-cultured them

for 48 hours with peptide-specific T cells. Following co-culture, we found that HLA-A2$^-$ CRC

organoids did survive irrespective of whether pulsed with the peptide or not (Fig. 2c). However,

only the HLA-A2$^+$ CRC organoids without prior peptide incubation survived co-culture (Fig. 2c).

Peptide-pulsed HLA-A2$^+$ CRC organoids were effectively killed by the peptide-specific T cells

providing a proof-of-principle that organoids can be utilised to study anti-tumour response by

cytotoxic T cells *in vitro*.

To further confirm antigen-specificity in our 'killing' assay system, we improved our co-

culture method by transfecting HLA-A2$^+$ CRC organoids with a construct expressing

mNeonGreen-tagged histone H2B and staining T cells with CellTracker violet to allow for long-

term tracking of both cell types (Methods). We then pulsed HLA-A2$^+$ CRC organoids with either

the WT1 peptide or with an EBV-derived peptide (Methods) and co-cultured the organoids with T

cells carrying either a WT1- or an EBV-specific TCR. Here, only organoids pulsed with the

cognate peptide were efficiently killed by the T cells (Fig. 2d, Supplementary Movies 1 and 2).

Testing for IFN-γ production by the T cells in the co-culture using enzyme-linked immunosorbent

assay (ELISA) confirmed antigen-specific organoid killing by the T cells (Fig. 2e). In order to

better follow the kinetics of the organoid killing, we applied to a fluorescent dye (NucRed Dead

647, Methods), which specifically stains apoptotic cells, and performed live confocal imaging on

the co-culture (Fig. 2f). We then quantified organoid killing by assessing co-localization of

NucRed Dead dye with H2B-mNeonGreen (Methods). Significant co-localisation of both labels

and, hence, organoid killing, was only observed when peptide-pulsed HLA-A2$^+$ CRC organoids

were co-culture with the respective peptide-specific T cells (Fig. 2g). Furthermore, T cells

infiltrating into the epithelium of the organoids could be readily detected in this co-culture

condition (Fig. 2h). In conclusion, T cells efficiently killed co-cultured CRC organoids in an

antigen-specific manner.

125 Discussion

Here we demonstrate that epithelial organoids can be used faithfully to recapitulate the interaction

between tumour tissue and the immune system. Also, using our co-culture assay, we set a first step

in rebuilding the tumour microenvironment *in vitro*. Further addition of other components of this

microenvironment (such as fibroblasts, natural killer cells, myeloid-derived suppressor cells, B

cells) may shed light on the complex interactions between the different cell types leading to

immune evasion of the tumour. Lastly, this co-culture system can be used as a scaffold for T-cell

expansion for cancer immunotherapy or for drug-screens testing applicability of certain

immunotherapies, for instance, chimeric antigen receptor (CAR)- or TCR transgenic T cells,

antibody-dependent cell-mediated cytotoxicity (ADCC) or antibody-dependent cellular

phagocytosis (ADCP) inducing antibodies directed at the tumour, to different tumours and

different patients.

138 Methods

Procurement of human material and informed consent

Colonic tissues (both normal colon and tumour tissue) were obtained from the Departments of

Surgery and Pathology of the Diakonessenhuis hospital, Utrecht, the Netherlands. All patients

included in this study were diagnosed with CRC. Informed consent was signed by all included

patients. Collection of tissue was approved by the medical ethical committee (METC) of the

Diakonessenhuis hospital, in agreement with the declaration of Helsinki and according to Dutch

and European Union legislation.

Organoid generation and cultures

Epithelial organoid lines were derived from healthy colon or tumor tissue as previously

described[12,20]. In brief, healthy colonic crypts were isolated by digestion of the colonic mucosa in

chelation solution (5.6 mM Na_2HPO_4, 8.0 mM KH_2PO_4, 96.2 mM NaCl, 1.6 mM KCl, 43.4 mM

Sucrose, and 54.9 mM *D*-Sorbitol, Sigma) supplemented with dithiotreitol (0.5 mM, Sigma) and

EDTA (2 mM, in-house), for 30 minutes at 4°C. Colon crypts were subsequently plated in

basement membrane extract (BME; Cultrex PC BME RGF type 2, Amsbio) and organoids were

grown in human intestinal stem cell medium (HISC), which is composed of Advanced Dulbecco's

modified Eagle medium/F12 supplemented with penicillin/streptomycin, 10 mM HEPES and

Glutamax (all Gibco, Thermo Fisher Scientific) with 50% Wnt3a conditioned medium (in-house),

20% R-Spondin1 conditioned medium (in-house), 10% Noggin conditioned medium (in-house), 1

x B27, 1,25 mM n-acetyl cysteine, 10 mM nicotinamide, 50 ng/mL human EGF, 10 nM Gastrin,

500 nM A83-01, 3 μM SB202190, 10 nM prostaglandine E2 and 100 μg/mL Primocin (Invivogen).

Tumor specimens were digested to single cells in collagenase II (1 mg/mL, Gibco, Thermo

Scientific), supplemented with hyaluronidase (10 μg/mL) and LY27632 (10 μM) for 30 minutes

at 37°C while shaking. Single tumor cells were plated in BME and organoids were cultured in

HICS minus Wnt conditioned medium and supplemented with 10 μM LY27632 at 37°C.

Organoid transfection

CRC organoids were dissociated into small clumps using TrypLE and then transduced with H2B-

mNeonGreen (pLV-H2B-mNeonGreen-ires-Puro), as previously described[21].

T cells

Generation of αβ T cells carrying a transgenic TCR recognizing an HLA-A2-restricted WT1-

derived peptide were described elsewhere[18]. Briefly, TCRα and β chains were cloned from raised

tetramer positive T cell clones. Subsequently, CD8$^+$ αβ TCR T cells were transduced using

retroviral supernatant from Phoenix-Ampho packaging cells that were transfected with gag-pol,

env, and pBullet retroviral constructs containing the cloned TCR genes.

Organoid-T cell co-culture and live cell imaging

Organoids stably transfected with H2B-mNeonGreen were split and digested a 5 to 7 days prior to

co-culture and seeded at a density of 5000 cells per 10 μL of BME (25,000 cells per well in a 12-

well cell culture plate). Two days prior to co-culture, T cells were starved from IL-2. One day prior

to co-culture, organoids were stimulated with IFN-γ at indicated concentrations.

Organoids were pulsed with TCR-specific peptide (ProImmune) for 2 hours at 37°C prior to co-

culture. Organoids and T cells were harvested and taken up in T cell medium, supplemented with

10% BME, 100 IU/mL IL-2 and NucRed Dead 647 (Thermo Fischer). Where indicated, anti-PD1

blocking antibodies (2 μg/mL) were added to the co-culture. Cells were plated in glass-bottom 96-

well plates and co-cultures were imaged using an SP8X confocal microscope (Leica).

Flow cytometry

APC-labelled pentamers to the EBV-derived, HLA-2:02 restricted peptide FLYALALLL

(ProImmune) where used to sort pentamer[+] CD8[+] CD3[+] T cells from PBMCs isolated from

buffycoats from healthy individuals. Cells were sorted as single cells into 96-well plates using a

BD FACS Aria (BD Biosciences) cytometer. For flow cytometry, the following antibodies were

used (all anti-human): CD8–PE (clone RPA-T8), CD45–PerCP-Cy5.5 (2D1), CD274 (PD-L1)–

APC (MIH1) (all BD Biosciences), CD279 (PD-1)–PE (EH12.2H7, Biolegend), HLA-A2–PE

(BB7.2, Santa Cruz).

Quantitative polymerase chain reaction (qPCR)

For qPCR analysis, RNA was isolated from organoids using the RNAeasy kit (QIAGEN)

according to the manufacturer's protocol. PCR analysis was performed using the SYBR Green

Reagent (Biorad). PCR reactions were performed in duplicate with a standard curve for every

primer. Primers were designed using the NCBI primer design tool. Primers used in this study:

GAPDH forward (GTC GGA GTC AAC GGA TT), *GAPDH* reverse (AAG CTT CCC GTT CTC

AG), *HPRT* forward (GGC GTC GTG ATT AGT GAT), *HPRT* reverse (AGG GCT ACA ATG

TGA TGG), *CD274* forward (TGC AGG GCA TTC CAG AAA GAT), *CD274* reverse (CCG

TGA CAG TAA ATG CGT TCAG).

Transcriptional profiling

Microarray analysis of biobank organoids was performed as described elsewhere[12].

Enzyme linked immunosorbent assays (ELISA)

Culture supernatants were kept at −20°C and ELISA was performed for indicated cytokines using

ELISA MA Standard (Biolegend) according to manufacturer's protocol.

Image analysis

Image analysis was done using Imaris software package (Bitplane). In brief, threshold for positive

staining was set on negative controls. A co-localization channel was made for H2B-neon and

NucRed Dead 647 signals. Cell death was quantified as percentage of H2B-mNeonGreen[+] voxels

co-localising with NucRed Dead signal.

Bioinformatics analysis

220 Bioinformatics analysis of gene-expression data from microarray experiments[12] was performed

221 using standard packages (*i.e.* gplots) in R version 3.4.0 (R Foundation, https://www.r-project.org)

222 and RStudio version 1.0.143 (https://www.rstudio.com).

Statistical analysis

225 All experiments were repeated at least three times unless otherwise indicated. All data were shown

226 as mean ± SEM. Statistical significance was analyzed by either ANOVA or two-tailed Student's

227 *t*-test using either Graphpad Prism 6 or Microsoft Excel 2010.

References

1. Ferlay, J. *et al.* GLOBOCAN 2012 v1.0, Cancer Incidence and Mortality Worldwide:
IARC CancerBase No. 11. *International Agency for Research on Cancer* (2013).

2. Markowitz, S. D., Dawson, D. M., Willis, J. & Willson, J. K. Focus on colon cancer.
Cancer cell **1**, 233-236 (2002).

3. Vries, R. G., Huch, M. & Clevers, H. Stem cells and cancer of the stomach and intestine.
Molecular oncology **4**, 373-384, doi:10.1016/j.molonc.2010.05.001 (2010).

4. Markowitz, S. D. & Bertagnolli, M. M. Molecular origins of cancer: Molecular basis of
colorectal cancer. *The New England journal of medicine* **361**, 2449-2460,
doi:10.1056/NEJMra0804588 (2009).

5. Hanahan, D. & Weinberg, R. A. Hallmarks of cancer: the next generation. *Cell* **144**, 646-
674, doi:10.1016/j.cell.2011.02.013 (2011).

6. Junttila, M. R. & de Sauvage, F. J. Influence of tumour micro-environment heterogeneity
on therapeutic response. *Nature* **501**, 346-354, doi:10.1038/nature12626 (2013).

7. Gajewski, T. F., Schreiber, H. & Fu, Y. X. Innate and adaptive immune cells in the tumor
microenvironment. *Nature immunology* **14**, 1014-1022, doi:10.1038/ni.2703 (2013).

8. Dunn, G. P., Koebel, C. M. & Schreiber, R. D. Interferons, immunity and cancer
immunoediting. *Nature reviews. Immunology* **6**, 836-848, doi:10.1038/nri1961 (2006).

9.   Zitvogel, L., Galluzzi, L., Smyth, M. J. & Kroemer, G. Mechanism of action of
248   conventional and targeted anticancer therapies: reinstating immunosurveillance. *Immunity*
249   **39**, 74-88, doi:10.1016/j.immuni.2013.06.014 (2013).

250   10.   Kretzschmar, K. & Clevers, H. Organoids: Modeling Development and the Stem Cell
251   Niche in a Dish. *Developmental cell* **38**, 590-600, doi:10.1016/j.devcel.2016.08.014
252   (2016).

253   11.   Drost, J. & Clevers, H. Organoids in cancer research. *Nature reviews. Cancer* **18**, 407-418,
254   doi:10.1038/s41568-018-0007-6 (2018).

255   12.   van de Wetering, M. *et al.* Prospective derivation of a living organoid biobank of colorectal
256   cancer patients. *Cell* **161**, 933-945, doi:10.1016/j.cell.2015.03.053 (2015).

257   13.   Leone, P. *et al.* MHC class I antigen processing and presenting machinery: organization,
258   function, and defects in tumor cells. *Journal of the National Cancer Institute* **105**, 1172-
259   1187, doi:10.1093/jnci/djt184 (2013).

260   14.   Fearon, E. R. & Vogelstein, B. A genetic model for colorectal tumorigenesis. *Cell* **61**, 759-
261   767 (1990).

262   15.   Drost, J. *et al.* Sequential cancer mutations in cultured human intestinal stem cells. *Nature*
263   **521**, 43-47, doi:10.1038/nature14415 (2015).

264   16.   Fumagalli, A. *et al.* Genetic dissection of colorectal cancer progression by orthotopic
265   transplantation of engineered cancer organoids. *Proceedings of the National Academy of*
266   *Sciences of the United States of America* **114**, E2357-E2364,
267   doi:10.1073/pnas.1701219114 (2017).

268   17.   Kryczek, I. *et al.* Cutting edge: IFN-gamma enables APC to promote memory Th17 and
269   abate Th1 cell development. *Journal of immunology* **181**, 5842-5846 (2008).

270   18.   Kuball, J. *et al.* Facilitating matched pairing and expression of TCR chains introduced into
271   human T cells. *Blood* **109**, 2331-2338, doi:10.1182/blood-2006-05-023069 (2007).

272   19.   Sebestyen, Z. *et al.* RhoB Mediates Phosphoantigen Recognition by Vgamma9Vdelta2 T
273   Cell Receptor. *Cell reports* **15**, 1973-1985, doi:10.1016/j.celrep.2016.04.081 (2016).

274   20.   Sato, T. *et al.* Long-term expansion of epithelial organoids from human colon, adenoma,
275   adenocarcinoma, and Barrett's epithelium. *Gastroenterology* **141**, 1762-1772,
276   doi:10.1053/j.gastro.2011.07.050 (2011).

277   21.   Bolhaqueiro, A. C. F. *et al.* Live imaging of cell division in 3D stem-cell organoid cultures.
278   *Methods in cell biology* **145**, 91-106, doi:10.1016/bs.mcb.2018.03.016 (2018).
279   22.   Sachs, N. *et al.* A Living Biobank of Breast Cancer Organoids Captures Disease
280   Heterogeneity. *Cell* **172**, 373-386 e310, doi:10.1016/j.cell.2017.11.010 (2018).

Acknowledgements

We thank T. Aarts, S. van den Brink, A. Cleven, T. Mizutani, M. Schiffler, M. van de Wetering
for technical assistance and experimental advice, A. de Graaff and the Hubrecht Imaging Facility
for help with microscopy, S. van der Elst, R. van den Linden and T. Poplonski for help with flow
cytometry. This work was supported by the European Research Council (Advanced Grant ERC-
AdG 67013-Organoid to H.C.) and a VENI grant from the Netherlands Organisation for Scientific
Research (NWO-ZonMW, 016.166.140 to K.K.). This work is part of the Oncode Institute, which
is partly funded by the Dutch Cancer Society. K.K. is a long-term fellow of the Human Frontier
Science Program Organization (HFSPO, LT771/2015) and was a long-term fellow of the European
Molecular Biology Organisation (EMBO, ALTF 839-2014).

Author contributions

Y.B.E. and K.K. designed, performed and analysed the experiments and wrote the manuscript.
Y.B.E. performed image analysis. K.K. performed bioinformatics analysis. P.A., E.d.J. and K.E.B.
assisted with experiments. J.D. generated cancer gene-mutant organoid lines. J.v.G. isolated
tumour and normal tissue from resected material. A.P. and N.S. performed surgery. I.J.G., Z.S.
and J.K. provided WT1 peptide and WT1 peptide-specific transgenic TCR $\alpha\beta$ T cells. R.G.J.V.
organised tissue collection. K.K. and H.C. acquired funding. H.C. supervised the project and wrote
the manuscript. All of the authors commented on the manuscript.

Fig. 1 | CRC organoids express immunomodulatory molecules. a,**b**, Normal colon and CRC organoid lines were generated in a patient-specific manner and RNA was extracted and analysed using Affymetrix single transcript microarrays. Average gene expression of different immunomodulators in normal colon and CRC organoid lines; n.s., non-significant; *, $p < 0.05$ (**a**). Hierarchical clustering of the individual normal colon and CRC organoid lines in the 'living biobank' displaying gene expression of selected immunomodulators. Color gradients represent z valued of each row (gene transcripts). **c-d**, Human colon organoid lines genetically engineered to carry one or more mutations found in CRCs. Expression levels of CD274 (PD-L1) in organoid lines (n = 2) at steady state (Ctrl) and upon stimulation with 20 ng/mL recombinant human IFN-γ assessed by quantitative PCR (**c**) and flow cytometry (**d**). A, $APC^{KO/KO}$; N.D., not detected; K, $KRAS^{G12D/+}$: P, $P53^{KO/KO}$; S, $SMAD4^{KO/KO}$, WT, wild-type.

316
317

Fig. 2 | CRC organoids as tools for assessment of antigen specific killing by CD8$^+$ T cells. a,
Experimental scheme. **b**, Flow cytometry analysis of HLA-A2 expression in cloned HLA-A2$^+$ and
HLA-A2$^-$ lines. **c**, Brightfield images of HLA-A2$^+$ and HLA-A2$^-$ CRC organoids co-cultured
with transgenic T cells expressing the WT1 peptide-specific T-cell receptor for 48 hours; scale
bars: 1 mm. **d,** Images showing HLA-A2$^+$ CRC organoids transfected with H2B-neon and pulsed
with indicated peptides at the beginning of co-culture (left panels) and after 18-hour co-culture
with indicated peptide-specific T cells (right panels); scale bars: 70 µm. **e**, IFN-γ production by
WT1 (top) and EBV (bottom) peptide-specific T cells as measured by ELISA of supernatants
collected after 18-hour co-culture with HLA-A2$^+$ CRC organoids pulsed with indicated peptides.
f, Live-cell imaging stills of a co-culture experiment with H2B-mNeonGreen-transfected HLA-
A2$^+$ CRC organoids pulsed with EBV peptide and co-cultured for 14 hours with an EBV-specific
T-cell clone. Images were taken at the indicated time points. **g**, Quantification of CRC organoid
killing by specific T cells as assessed by co-localization of H2B-mNeonGreen and NucRed Dead
dye. Graphs are representative of multiple repeated experiments with either EBV peptide and EBV
T-cell- or WT1 peptide and WT1 T-cell co-cultures. **h**, Representative projection image of T cells

333   (blue) infiltrating a peptide-pulsed CRC organoid as recorded during the live-cell imaging

334   experiments.