1

Sympatric and allopatric evolutionary contexts shape differential immune response in

***Biomphalaria / Schistosoma* interaction.**

Anaïs Portet[1], Silvain Pinaud[1,#], Cristian Chaparro[1], Richard Galinier[1], Guillaume M.

Charriere[2], Jean-François Allienne[1], David Duval[1], Benjamin Gourbal[1§]

[1] Univ. Perpignan Via Domitia, Interactions Hôtes Pathogènes Environnements UMR 5244,

CNRS, IFREMER, Univ. Montpellier, F-66860 Perpignan, France

[2] Interactions Hôtes-Pathogènes-Environnements (IHPE), UMR 5244, CNRS, Ifremer,

Université de Perpignan Via Domitia, Université de Montpellier, Montpellier, 34095, France.

present address : Wellcome Trust Sanger Institute, Hinxton, CB10 1SA, UK
17

19 [§] corresponding author:

20 Benjamin Gourbal

21 UMR5244 IHPE, Université de Perpignan

22 58 avenue Paul Alduy, 66860 Perpignan

23 benjamin.gourbal@univ-perp.fr

Abstract

Selective pressures between hosts and their parasites can result in reciprocal evolution or adaptation of specific life history traits. Local adaptation of resident hosts and parasites should lead to host-parasite systems performing better in sympatry when compared to allopatry. Between-population variations in parasite infectivity/virulence and host defence/resistance, referred to as compatibility phenotype, were often the proxy used to analyse sympatric or allopatric adaptation. Nevertheless, some reported cases exist where allopatric host-parasite systems demonstrate compatibility phenotypes similar or greater than the one observed in sympatry. In these cases, the role of local adaptation is worth considering. Here, we study the interaction between *Schistosoma* and its vector snail *Biomphalaria* in which such a discrepancy in local versus foreign compatibility phenotype has been observed. Herein, we developed an integrative approach to investigate sympatric and allopatric interaction processes and link the underlying molecular mechanisms to the resulting phenotypes. Using comparative "omics" approaches joined to analysis of life history traits (immune cellular response, mortality, prevalence and compatibility) we tried to bridge the gap of knowledge that exists for connecting local adaptation observations to molecular phenotypes in *Schistosoma/Biomphalaria* interactions.

We found that despite displaying similar prevalence phenotypes, parasite infection triggered an immune suppression in snails living in sympatry, while it activated an immune response for those living in allopatry. Dual-comparative molecular analyses revealed that parasite infection causes immune suppression in sympatry. miRNAs were used to hijack the host's immune response, allowing sympatric parasites to initiate their developmental program earlier and more efficiently.

We show that despite having similar prevalence phenotypes, sympatric and allopatric snail-*Schistosoma* interactions displayed a strongly different immunobiological molecular dialogue. The ability of allopatric pathogens to adapt rapidly and efficiently to new hosts could have critical consequences on disease emergence and risk of schistosomiasis outbreaks. These observations would have important consequences in term of schistosomiasis disease control.

Keywords: sympatry, allopatry, host-parasite compatibility, RNA-seq, immunity

Introduction

Schistosomiasis is the second most widespread human parasitic disease after malaria and affects more than 200 million people worldwide (1). *Schistosoma mansoni* (Platyhelminthes, Lophotrochozoa) causes intestinal schistosomiasis. *Schistosoma* needs a mollusc (belonging to *Biomphalaria, Bulinus* or *Planorbarius* genus) to act as its first intermediate host, where it undergoes part of its life cycle before infecting humans. Patently infected snails support the continuous production of thousands of cercariae, the free-living human-infective stage of the parasite, until dying. As such, vector snails appear to be a central actor in the parasite transmission and thus have deserved greater attention as a potential target for schistosomiasis control. To achieve this objective, it is necessary to understand snail-parasite immunobiological interactions and characterize the molecular mechanisms by which snails and *Schistosoma* interact.

Biomphalaria/Schistosoma compatibility is based on a complex phenotype-to-phenotype or matching-phenotype model (2-7). The success or failure of the infection of *B. glabrata* by *S. mansoni* reflects a complex interplay between the host's defence mechanisms and the parasite's infective strategies. In the past 15 years, the molecular basis of this compatibility polymorphism has been investigated at the genomic level (8-10), the transcriptomic level (6, 11-15), the proteomic level (16-23) and the epigenomic level (24-29). These studies have revealed that various molecules and pathways involved in immune recognition (snail immune receptors versus parasite antigens), immune effector/anti-effector systems and immune regulation/activation participate in a complex interplay that governs the match or mismatch of host and parasite phenotypes (30). This complex phenotype-by-phenotype interaction (compatibility polymorphism) varies between populations and individuals resulting in "multi-parasite susceptibility" or "multi-host infectivity" phenotypes (5) that reflect between-population variations in parasite infectivity/virulence and host defence/resistance (31, 32).

In *B. glabrata/Schistosoma sp.* models, numerous sympatric and allopatric combinations have been tested: (i) different *B. glabrata* laboratory strains (or isolates) show various degrees of susceptibility to *S. mansoni* infection, and (ii) different strains of *S. mansoni* display different levels of infectivity toward a particular strain of snail host (2, 4, 5, 33, 34). In most cases, interaction in *B.*

glabrata/Schistosoma models has been investigated by comparing, (i) sympatric/compatible and (ii) allopatric/incompatible host-parasite associations. Indeed, the general assumption is that the parasites, which have a shorter generation time, larger population sizes and higher reproductive outputs, are ahead in the co-evolutionary conflict against their host and are therefore more likely to locally adapt to them (35, 36). This is exactly what is stated by the "host-parasite local adaptation hypothesis", in which the sympatric parasite is supposed to perform better than the allopatric one (36-39). However, it has been observed that compatibility phenotypes between *Biomphalaria* and *Schistosoma* were the same comparing sympatric and allopatric combinations or even greater for allopatric parasites, and thus did not fulfil the "local versus foreign" criterion of the "host-parasite local adaptation hypothesis" (33, 40, 41). Such compatibility phenotypes have been mostly investigated in terms of parasite infectivity or pathogenicity toward snail hosts, but to our knowledge very few studies have investigated the molecular basis of allopatric compatible interactions from the perspective of both sides of the interaction, the host and parasite (42, 43).

Hence, in order to bridge this gap, we herein study sympatric/allopatric interactions displaying similar compatibilities using an integrative approach that links the underlying molecular mechanisms to the resulting phenotypes, based on comparative molecular approaches on both host snails and *Schistosoma* parasites. We set out to analyse the compatibility between South American snail strains (from Recife Brazil and Guacara Venezuela) in combination with three different parasite isolates that displayed nearly the same compatibility phenotype: (i) *S. mansoni* Recife Brazil, (ii) *S. mansoni* Guacara Venezuela (both sympatric and allopatric, narrow geographic scale) and (iii) *S. rodhaini* from Burundi Africa (allopatric, large geographic and phylogenetic scales). Thus, even if compatibility phenotypes (prevalence/intensity) are similar, the characterization of the underlying molecular mechanisms demonstrates a very different immunobiological dialogue between *B. glabrata* vector snails and their sympatric or allopatric *Schistosoma* parasites at the cellular and molecular scales. To the best of our knowledge very few studies have investigated the molecular basis of local adaptation in host/parasite interactions (44, 45). Here we used a powerful global approach to identify the differentially regulated transcripts in snails of a single *B. glabrata* strain under infection by three sympatric or allopatric compatible strains or species of the parasite, *Schistosoma sp.*

Moreover, our results may be of critical importance in terms of schistosomiasis control. Understanding

the mechanisms whereby allopatric parasites efficiently infect host snails, would be essential to

develop new tools to predict or quantify risks of schistosomiasis disease outbreaks (46).

Materials and Methods

Ethical statements

Our laboratory holds permit # A66040 for experiments on animals, which was obtained from the

French Ministry of Agriculture and Fisheries and the French Ministry of National Education,

Research, and Technology. The housing, breeding and care of the utilized animals followed the ethical

requirements of our country. The experimenter possesses an official certificate for animal

experimentation from both of the above-listed French ministries (Decree # 87–848, October 19, 1987).

The various protocols used in this study have been approved by the French veterinary agency of the

DRAAF Languedoc-Roussillon (Direction Régionale de l'Alimentation, de l'Agriculture et de la

Forêt), Montpellier, France (authorization # 007083).

Biological materials

Two snail strains of *Biomphalaria glabrata* were used in this study: the albino Brazilian strain,

(BgBRE) and the Venezuelan strain, (BgVEN). They were exposed to infection by a Brazilian

(SmBRE) or Venezuelan (SmVEN) strain of *Schistosoma mansoni*, or by another *Schistosoma*

species, *S. rodhaini* (Srod). All host and parasite strains of each different geographical origin were

recovered in their native locality and maintained in the laboratory always on their sympatric snail

hosts to maintain reciprocal selective pressures and reciprocal sympatric strain adaptation. We housed

snails in tanks filled with pond water at 25°C with a 12:12 hour light:dark cycle and supplied *ad*

libitum with fresh lettuce. The Brazilian strain originates from the locality of Recife (east Brazil), the

Venezuelan strains of snail and parasite were recovered from the locality of Guacara (north

Venezuela) and the African species *Schistosoma rodhaini* originates from Burundi and was obtained

from the London British Museum. These *Schistosoma* isolates/species have been selected because they

exhibited similar infectivity toward BgBRE or BgVEN strains (see prevalence and intensity in

supplementary table S1). Prevalence of SmBRE and SmVEN for the African vector snail

Biomphalaria pfeifferi from Senegal (BpSEN), and prevalence of the corresponding parasite SmSEN

on South American snails were also tested (supplementary table S1).

RNAseq experimental protocol

In order to investigate the molecular response of snails against sympatric and allopatric parasites, a

global comparative transcriptomic approach was conducted. One hundred and twenty BgBRE snails

were infected with SmBRE, SmVEN or Srod. Each snail was individually exposed for 12 h to 10

miracidia in 5mL of pond water. For each experimental infection, 20 snails were recovered at 24 and

96 h after infection. The samples were named as follows: BB24, BB96 for BgBRE infected with

SmBRE; BV24, BV96 for BgBRE infected with SmVEN; and BR24, BR96 for BgBRE infected with

Srod. Two pools of 30 uninfected BgBRE snails (Bre1 and Bre2) were used as controls.

148   - <u>Whole-snail RNA extraction and sequencing</u>

149   Total RNA was extracted using TRIZOL$^{®}$ (Sigma Life Science, USA) according to the

150   manufacturer's instructions. Sequencing was performed using paired-end 72-bp read lengths on

151   Illumina Genome Analyzer II (MGX-Montpellier GenomiX, Montpellier, France).

152   - <u>De novo transcriptome assembly</u>

153   *De novo* transcriptome assembly, using all time points, was performed using an in-house pipeline

154   created with the Velvet (1.2.02), Oases (v0.2.04) and CD-HIT-EST (v4.5.4) programs. The assembly

155   of the consensus reference transcriptome was optimized using various parameters, including k-mer

156   length, insert length and expected coverage, as previously described (47, 48). A de novo transcriptome

157   was created that contained 159,711 transcripts.

158   - <u>Differential expression analysis</u>

High-quality reads (Phred score >29) were aligned to the de novo transcriptome using Bowtie2

(v2.0.2), which was run locally on a Galaxy server. To compare the host responses during the

sympatric or allopatric interactions, we used the DESeq2 (v2.12) software to analyse the differential

transcript representation between *BgBRE* control strains (uninfected *Bg*BRE1 and *Bg*BRE2) to the

sympatric and allopatric conditions (p-value < 0.1) (48). A Venn diagram was generated using the

Venny 2.1 software to highlight which differentially expressed transcripts were specific or common to

the different interactions. A heatmap was then created to analyse transcript expression patterns, using

log2 Fold Change with Hierarchical Ascending Clustering (HAC) and Pearson correlation

(uncentered) as applied by the Cluster (v3.0) and Java TreeView (v1.1.6r4) software packages. The

differentially represented transcripts were functionally classified using a BlastX analysis with the cut-

off set to e-value < $1e^{-3}$ (NCBI dataset; thanks to the Roscoff Data center Cluster, UPMC) and gene

ontology was assigned using an automatic annotation, Blast2GO (v3.0.8) (supplementary table S2).

Based on automatic annotation and functional domains and according to the literature we identified the

potential immune transcripts involved in snail immunity.

Schistosoma intra-molluscal stage transcriptome analysis: Dual RNA-seq

Host response against pathogens constitutes solely one side of the molecular dialogue occurring in

host/parasite interactions. To analyse all the complexity of such dialogue, the parasite response into

host tissue also needs to be investigated. Herein a dual RNA-seq approach was thus conducted to gain

a broader understanding of sympatric and allopatric host/parasite interactions.

- Schistosome read selection

The *Biomphalaria* (v1) and *Schistosoma* (v5.2) genomes have been concatenated

(https://www.vectorbase.org/organisms/biomphalaria-glabrata;

http://www.sanger.ac.uk/resources/downloads/helminths/schistosoma-mansoni.html).   High-quality

reads (Phred score > 29) were mapped against these concatenated genomes using Bowtie2 (v2.0.2),

which was run locally on the Galaxy project server. The reads that mapped only once and exclusively

to the *Schistosoma* genome were collected as corresponding to *Schistosoma* reads; reads that mapped

to the *Biomphalaria* genome or more than once to either genomes were removed from the analysis.

- Gene analysis

The above-selected *Schistosoma* reads were mapped against the concatenate genome to identify intra-

molluscal stage-specific *Schistosoma* genes. In order to select the relevant genes, the reads mapped in

all experimental conditions were summed. Only genes with a minimal sum of 10 reads were kept for

the analysis. A heatmap was generated to analyse *Schistosoma* gene expression patterns using

Hierarchical Ascending Clustering (HAC) with Pearson correlation (uncentered) as applied by the

Cluster (v3.0) and Java TreeView (v1.1.6r4) software packages. Functional annotation of the genes

was assigned using BlastX with the cut-off set to e-value $< 1e^{-3}$ (NCBI dataset, local cluster) and gene

ontology was performed using Blast2GO (v4.0.7) (supplementary table S3).

Innate immune cellular response analysis: microscopy and flow cytometry

Hemocytes appeared as the main cells supporting the *Biomphalaria* snail immune response. Thus, to

go further in the description of snail response against parasites, quantitative and qualitative changes in

hemocyte populations were investigated. For this purpose, BgBRE and BgVEN snails were used.

Snails were infected as described above, using either SmBRE or SmVEN parasites. For each

experimental infection, snails were recovered at 24 and 96 h after infection and designated as follows:

BB24 and BB96 for BgBRE infected with SmBRE; BV24 and BV96 for BgBRE infected with

SmVEN; VV24 and VV96 for BgVEN infected with SmVEN; and VB24 and VB96 for BgVEN

infected with SmBRE. Snails of each strain, BgBRE and BgVEN, were recovered and used as

controls.

- Hemocyte proliferation analysis: microscopy

Microscopic inspection of hemocyte proliferation was conducted using 12 infected BgBRE (6

BgBRExSmBRE and 6 BgBRExSmVEN) and 3 uninfected BgBRE snails. The hemocytes of 3 snails

(biological replicates) were counted for each condition at 24 h and 96 h after infection. The

proliferation of circulating hemocytes was studied by using a Click-iT EdU Alexa Fluor 488 Flow

Imaging Kit (Molecular Probes). At each time point, circulating hemocytes were recovered by direct

puncture after foot retraction and 1mM of EdU solution was added to the hemolymph. Three hours

later, the amount of EdU incorporated by the circulating hemocytes was visualized in vitro after

fixation of the cells and performing a covalent coupling of Alexa Fluor 488 to the EdU residues trough

a click chemistry reaction flowing manufacturer indications. Next, nuclei of hemocytes were

counterstained with DAPI (Biotum) staining, and the sample was analysed on a confocal microscope

using a Zeiss LSM 700, with 4 lasers (405, 488, 555 and 633 nm). Positive cells were counted and

between-sample differences in the percentage of proliferation were tested using a Fisher exact test,

with significance accepted at p-value < 0.05.

- Hemocyte proliferation and population profiles analysis: flow cytometry

Qualitative changes in hemocyte populations following infection by sympatric or allopatric parasites

were studied using a flow cytometry approach. For this, 72 infected BgBRE or BgVEN (36 infected

by SmBRE and 36 infected by SmVEN) and 18 uninfected BgBRE or BgVEN snails were used. Six

biological replicates (pools of three snails per replicate) were performed for each condition. Flow

cytometry was used to profile and assess the proliferation of circulating hemocytes using Click-

iTEdUAlexa Fluor 647 labelling (Molecular Probes). At each time point, 1mM of EdU solution was

injected into pericardial cavity of each snail. Three hours later six replicates of three snails were

collected, and the hemolymph was extracted from the head-foot according to standard procedures (49).

The hemolymph was pooled from the three snails, and 100 µl were subjected to analysis with the

above listed kit, according to the manufacturer's instructions. The percentage of proliferative cells was

calculated by flow cytometry.

The hemocytes were profiled along the course of infection using Side Scatter Chanel (SSC) to

estimate cell granularity and Forward Scatter Chanel (FSC) to estimate cell size. The cell repartition

along these two parameters enables the identification of cell subpopulations. The flow cytometry was

performed using a FACS Canto from BD Biosciences (RIO Imaging Platform, Montpellier, France).

For each sample, 10,000 events were counted. The results were analysed with the FlowJo V 10.0.8

software. Between-group differences in the percent of proliferation were tested using the Mann-

Whitney U-test, with significance accepted at p-value<0.05.

Histological procedures

A histological approach was conducted in order to investigate differences in miracidia to sporocyst

development, while comparing sympatric and allopatric parasite growth, development and maturation

into snail tissues. BgBRE snails (n=8 per group) were infected as described above with either SmBRE

(sympatric) or SmVEN (allopatric) parasites. At 24 and 96 h after infection, two snails were fixed in

Halmi's fixative (4.5% mercuric chloride, 0.5% sodium chloride, 2% trichloroacetic acid, 20% formol,

4% acetic acid and 10% picric acid-saturated aqueous solution). Both, embedding in paraffin and

transverse histological sections (3-μm) were performed using the RHEM platform (Montpellier,

France) facilities. The slides were stained using Heidenhain's azan trichromatic staining solution as

follows: (i) serial re-hydration was performed in toluene followed by 95%, 70% and 30% ethanol and

then distilled water; (ii) coloration was performed using azocarmine G (70% ethanol, 1% aniline, 1%

acetic alcohol, distilled water, 5% phosphotungstic acid, distilled water, Heidenhain's azan) and (iii)

serial dehydration was performed using 95% ethanol, absolute ethanol, and toluene. The preparations

were then mounted with Entellan (Sigma Life Science, St. Louis Missouri, USA) and subjected to

microscopic examination. The parasite sizes were measured with the ImageJ software (v2.0.0), and

between-group size differences were tested using the Mann-Whitney U-test with statistical

significance accepted at a p-value<0.05 based on the mean of the three largest serial cuts of each

sporocyst in sympatric or allopatric conditions.

In-silico *characterization of Schistosoma miRNAs*

Parasites may communicate or interfere with their host using different strategies based mainly on

excreted/secreted products released into hemolymph. In this context, miRNAs appeared as the most

relevant means of communication that can be used by parasites. To test for such a hypothesis *S.*

mansoni miRNAs were analysed *in-silico* by comparing the relevant miRNA database (miRBase) to

our RNAseq libraries generated at the 24 h following sympatric or allopatric infections. S. *mansoni*

precursor sequences were downloaded from miRBase (http://www.mirbase.org, 03/09/2017), and

high-quality reads from naive (BgBRE) and 24 h post-infection samples (BB24, BV24, BR24) were

aligned against a *S. mansoni* sub-database of miRBase, as previously described (50). The identified

precursors were confirmed by alignment of high-scoring reads onto precursor and mature miRNAs

from miRBase. Only reads with 100% identity were retained for analysis. The localization of each

read against miRNA sequence allowed us to identify either the precursor or just the mature miRNA.

Precursors found under both naive and infected conditions were excluded to retain exclusively the

miRNAs present in samples from infected snails and avoid cross-species contamination because of the

potential conserved features of miRNAs from *B. glabrata* and *S. mansoni*.

Putative miRNA targets were predicted from among the differentially represented immune-

related transcripts (Fig. 1) using Miranda tools (using parameters: Miranda

input_miRinput_Transcriptome -out results.txt -quiet -sc 140 -en -15) (51). Because mature miRNAs

may exist in two forms depending on which strand (5'-3') of the precursor stem-loop is maturated the

predicted interactions could involve the 5' and/or 3' forms, as noted. The results were extracted using

the awk tool, listed in supplementary files (Supplementary Table 4), and used to generate a Venn

diagram. To confirm the ability of a selected pre-miRNA to produce the stem-loop necessary to

produce the mature form, the secondary structures of precursor were predicted using a RNA structure

Web tool (http://rna.urmc.rochester.edu/RNAstructureWeb, 03/09/2017) using default parameters.

Snail survival analysis

Allopatric or sympatric parasites could have different levels of virulence or impacts on their host that

could impair snail survival. To test for such discrepancy we investigated the mortality rates of infected

snails over the course of sympatric or allopatric infections. One hundred and sixty BgBRE snails were

infected as described above with SmBRE or SmVEN strains (n = 50), and 60 non-infected BgBRE

snails were retained as controls. The numbers of dead snails were compiled weekly for 14 weeks. A

Kaplan-Meier estimator was used to estimate the survival function from lifetime data. Survival curves

were generated using the xlstats Mac software and the log-rank test was applied with significance

accepted at $p < 0.05$.

Results

A whole-snail transcriptomic approach for investigating the molecular basis of the innate immune

response in the sympatric and allopatric contexts

The global transcript representation was analysed using the previously described RNAseq pipeline

developed in our laboratory (6, 47, 48). The *Bg*BRE snail host had the same prevalence and intensity

phenotypes when exposed to *Sm*BRE, *Sm*VEN or *S*rod parasites (see Supplementary Table 1). Of the

159,711 transcripts of the *Bg*BRE transcriptome, 3,865 (2.4%) were differentially represented in all

sympatric and allopatric conditions compared to naive snails (Table 1, Supplementary Fig. 1). We

performed automatic Blast2GO annotation, discarded the non-annotated transcripts, and retained 1,017

annotated transcripts (26.3% of the differentially represented transcripts). Finally, as we were

interested in the innate immune response of the snail to the sympatric and allopatric parasites, we

focused on transcripts known to have immune-related functions. Of the annotated differentially

expressed transcripts, 336 were identified as being immune-related; this subset corresponded to 8.7%

of all differentially represented transcripts. The heatmap generated for all differentially represented

transcripts is presented in Supplementary Fig. 1.

Venn diagrams were used to highlight the relationships of the transcripts that were

differentially represented in the *B. glabrata* and *Schistosoma sp.* interactions. Overall, 189, 180 and

164 transcripts were differentially represented in the BB (BgBRE/SmBRE, sympatric), BV

(BgBRE/SmVEN, allopatric) and BR (BgBRE/Srod, allopatric) interactions, respectively (Fig. 1A).

Of the selected transcripts, 40 were shared by the BB, BV and BR interactions (Fig. 1A), and

displayed globally the same differentially expressed pattern of expression (Fig. 1B, cluster 1). These

transcripts may correspond to the snail's basal immune response against *Schistosoma* parasites.

Of the 140 transcripts exclusive to the sympatric (BB) interaction, which represented 74.1% of

the immune transcripts found to be differentially represented in this interaction (Fig. 1A), 100% were

underrepresented at 24 h post-infection, and 74.6% of these transcripts were observed exclusively at

this time point (Fig. 1B, cluster 5).

Very similar transcript expression patterns were observed for the two allopatric interactions

(BV and BR). Clusters 6 and 7, comprising only 28 and 11 transcripts, were exclusive to BV and BR,

respectively. In BV interaction (Fig. 1B, cluster 6), at 96 h after infection 96.5% of the transcripts

were expressed: 22% over-represented. In BR interaction (Fig. 1B, cluster 7), 100% of the transcripts

were expressed at 96 h after infection: 82% overrepresented. Of the 108 transcripts shared by BV and

BR (cluster 3, representing 60% and 65.9 % of the transcripts, respectively) (Fig. 1A; Fig. 1B, cluster

3), 98.1% were expressed at 96 h post-infection: 28.2% over-represented (Fig. 1B, cluster 3).

Altogether, these results show that the host immune-related transcriptomic response differed

widely between the sympatric and allopatric interactions. All of the differentially represented

transcripts were under expressed at 24 h after infection in sympatry, while the differentially

represented transcripts in allopatry were mostly over expressed at 96 h after infection. Moreover, the

sympatric interaction was associated with a strong immunosuppression (Fig. 1B, cluster 5), whereas

allopatry yielded an activation of the immune response (Fig. 1B, cluster 3).

We decided to explore the function of differentially regulated transcripts between the

sympatric and allopatric interactions in order to investigate if these differentially regulated transcripts

belong to different functional categories. For that, we performed an automatic annotation with

Blast2GO (Supplementary Table 2). Based on automatic annotation and according to the literature we

distributed the relevant differentially expressed immune transcripts into three groups: (i) immune

recognition molecules, (ii) immune effectors and (iii) immune signalling molecules (Fig. 1C). Then

these three groups were subdivided into functional categories (Fig. 2). In the BB sympatric interaction,

immune recognition molecules, immune effectors and immune signalling pathways correspond to

13.1%, 18.3% and 4.2%, respectively (sums of all relevant transcripts at all infection time points).

Among the transcripts distributed into these groups, 89.0% were underrepresented. In the BV

allopatric interaction, the three groups corresponded respectively to 10.8%, 12.9% and 8.6% and

among them 65.9% were underrepresented. Finally, in the BR allopatric interaction, they

corresponded to 12%, 12.5% and 7.8% and 62.5% of them were underrepresented. When we

compared the percentage of each immunological group in the sympatric and allopatric interactions, no

specific functional subset was particularly repressed in the BB sympatric interaction (Fig. 1C; Fig. 2).

In fact, most of the time, the same immune functions were affected in sympatric and allopatric

infections but different immune transcripts (grey and black diamond in Fig. 2) showed differential

regulation following infections (Fig. 2).

The differentially regulated transcripts belonging to the three immunological groups (Fig. 2)

were largely involved in immune cellular responses, cell adhesion, extracellular matrix component,

cell migration, cell differentiation and cell proliferation. All of these functions were under-represented

in sympatric interaction (76%), whereas many were overrepresented in allopatric ones (39%) at 24 h

after infections (Fig. 2).

Immune cellular responses in the sympatric and allopatric contexts

As the transcriptomic approach indicated that multiple immunobiological functions seem to be altered in the host response during the interaction with the parasites, we next investigated whether differences in immune responses in sympatric versus allopatric infections could be seen at the cellular level. Hemocytes, the snail immune cells, participate directly in the immune response against the parasites, and immune cell activation under an immunological challenge can translate into cell proliferation and/or cell morphology modifications.

Thus, to investigate such immune cell activation, cell proliferation was quantified using EdU nuclear labelling (Fig. 3). EdU is a nucleoside analogue of thymine incorporated into DNA during DNA synthesis. This incorporation reflects the mitotic activity of hemocytes. As the two tested allopatric interactions appeared extremely similar at the transcriptomic level, we compared only the BV (allopatric) interaction and BB (sympatric) interaction for further analyses. Circulating hemocytes recovered in snail hemolymph were EdU-labelled in vitro to identify mitotic circulating hemocytes (Fig. 3A). EdU-positive hemocytes were quantified using confocal microscopy and showed that hemocyte proliferation was induced threefold following *Schistosoma* infection in BV compared to BB (Fig. 3B). Twenty-four hours after infection, proliferative circulating cells increase significantly in BV, with 5.2% of proliferative cells compared to BB with 2.6% (Fisher exact test two-tailed p = 7.6 e10^{-6}) (Fig. 3B). Moreover, this result demonstrates for the first time that "circulating" hemocytes are able to proliferate following *Schistosoma* infections.

Hemocyte proliferation was assessed using flow cytometry after in vivo EdU-labelling (Fig. 4A). At 24 h after infection, the rate of proliferating cells was increased significantly (Mann Whitney U test: U=36; z = -2.8; p = 0.0022) by more than twofold between the allopatric and sympatric interactions, 4.2% compared to 1.8% respectively. At 96 h after infection, there were fewer proliferating cells: the percentage of proliferating hemocytes was similar in sympatric BB interaction compared to the control (1%, Mann Whitney U test: U = 17; z = -0.27; p = 0.3936), while remaining somewhat higher in the allopatric BV interaction (2.3%, Mann Whitney U test: U = 0; z = 2.65; p = 0.009).

To confirm that our observations were related to the type of interaction rather than to a strain effect, we performed the same experiments using another *Biomphalaria glabrata* strain, BgVEN as the

host and SmVEN and SmBRE as the sympatric and the allopatric parasite, respectively (Fig. 4B). The rate of proliferating hemocytes increased significantly (Mann Whitney U test: U = 36; z = -2.8; p = 0.0022) by more than threefold between the allopatric (VB: BgVEN/SmBRE) and the sympatric (VV: BgVEN/SmVEN) interactions at 24 h after infection (6.8% compared to 2.0% respectively). At 96 h after infection, both rates were decreased, albeit more for VV (0.1%, Mann Whitney U test: U = 2; z = 2.48; p = 0.013) than for VB (2.7%, Mann Whitney U test: U = 36; z = 2.8; p = 0.0022). These results similar to those observed for BgBRE interactions, confirmed that the effects observed at the cellular level were related to the type of interactions (sympatric versus allopatric) rather than to the host or parasite strains used.

In addition to the mitotic activity, hemocyte morphology was estimated using flow cytometry by measuring size and granularity. We analysed the circulating hemocyte populations in noninfected versus infected snails in two *B. glabrata* snail strains (BgBRE and BgVEN) in sympatric and allopatric interactions with parasites (Fig. 4C and 4D). Morphology and heterogeneity of circulating hemocytes varied in a similar manner between BgBRE and BgVEN snails subjected to infection (Fig. 4C and 4D) and we observed infection-related differences in cell morphology. In noninfected snails, the content of circulating hemocytes was very heterogeneous but appeared to represent a single population that exhibited a continuous gradient of size and granularity (Fig. 4C and 4D), which is typical of *B. glabrata* hemocytes (52). However, hemocyte population heterogeneity changed quickly after infection. In allopatric interactions, 24 h after infection (Fig. 4C, BV24, and 4D, VB24) two populations could be distinguished: a population P1 (corresponding to that seen in noninfected snails) and a population P2 (a new population). P2 cells exhibited increased granularity, retained a high degree of size variability, and showed a mitotic activity, as indicated by EdU labelling (Fig. 4C and 4D, red dots). This profile was transitory, with the P2 population disappearing by 96 h after infection (Fig. 4C, BV96, and 4D, VB96). Altogether, these results show that, upon infection, the snail circulating immune cells exhibit a particular population dynamics with transient increase of the mitotic activity associated with morphology modifications that are likely to be involved in the host defence response against the parasite during the interaction. Moreover, this response of the immune cells appears to be exacerbated in the allopatric interaction when compared to the sympatric interaction.

Schistosoma growth and development in Biomphalaria tissues

<u>Parasite development</u>

To investigate the development of *S. mansoni* in *B. glabrata* tissues, we examined the fate of sporocyst in sympatric and allopatric compatible interactions using a histological approach. For both interactions, miracidia were able to penetrate, transform into primary sporocysts (SpI) and develop. At 24 h after infection, we observed a significant difference (Mann Whitney U test: U = 40; z = 4.33; p = 1.42 e10^{-6}) in the size of sporocyst from sympatric parasites (11,838 μm^2 average size) versus allopatric parasites (7,402 μm^2 average size) (Fig. 5). This difference was no longer seen 96 h after infection (41,413 μm^2 and 36,920 μm^2, respectively, Mann Whitney U test: U = 280; z = -1.31; p = 0.1917) (Fig. 5). These results show that during the early events following infection, the parasites develop more slowly in the allopatric compared to the sympatric interaction; thereafter, however, the allopatric parasites catch up quickly, such that no difference is seen at 96 h post-infection (Fig. 5).

<u>Parasite transcript expression analysis</u>

In order to investigate the biological responses of the parasites within their host, dual RNAseq analyses allowed us to identify regulated transcripts from *Schistosoma* parasites during their intra-molluscal development. The parasite RNAseq data at 24 h after infection revealed five clusters of differentially expressed genes from the sympatric (SmBRE) and the allopatric (SmVEN, Srod) parasite responses (Fig. 6). Cluster 1 corresponds to genes highly expressed and cluster 5 weakly expressed for all parasite strains. Cluster 2 represents genes overexpressed in SmBRE versus SmVEN and Srod. Cluster 3 contained genes overexpressed in SmBRE and SmVEN versus Srod and cluster 4 SmBRE and Srod versus SmVEN. In all clusters, the gene expression levels in SmBRE sympatric parasite are always greater than in the other allopatric parasites. Blast2GO annotation helps in characterizing 70% of the 351 genes shown in the five clusters (Supplementary Table 3). According to the global Gene Ontology (GO): 70% of the annotated genes were involved in general metabolism and growth, translation processes, regulation of cellular processes and RNA biosynthesis; 25% were involved in molecular transport or cell organization; and 5% were involved in organism defence or

response to stimuli. In all these clusters, we identified six parasite gene products that had been involved in parasite modulation or suppression of snail immunity. These molecules correspond to heat shock proteins (Fig. 6, Clusters 1 and 2) (27); glutathione-S-transferase, NADH dehydrogenase subunit, and calreticulin (Fig. 6, Cluster 2) (20, 53, 54); Alpha-2-macroglobulin (Fig. 6, Cluster 4) (55); von Willebrand factor type EGF with pentraxin domain (Fig. 6, Cluster 5) (56) (see Supplementary Table 3). These particular genes could be involved in the strategy used by the parasite to ensure immunosuppression and favour its development and growth within host tissues.

In allopatric interactions, an activation of the immune response is observed. Interestingly, allopatric parasites did not overexpress any transcripts that could have immunosuppressive function or impair the activation of the immune response (Fig. 6 and Supplementary Table 3). Furthermore, a variant of a glycerol-3-phosphate acyl-transferase (Schisto_mansoni.Chr_3.5623) is highly overexpressed in SmVEN and Srod compared to SmBRE (Cluster 1 Supplementary Table 3). This molecule is known to participate in the biosynthesis of phosphatidic acid, itself involved in macrophage activation and regulation of inflammatory signalling (57, 58). This would thus indirectly enhance the cellular immune response of *Biomphalaria glabarata* against those allopatric parasites.

Parasite microRNAs analysis

The microRNAs (miRNAs) are known as noncoding small RNA (<24nt) able to regulate gene expressions. As we identified deep modifications of transcriptional or posttranscriptional activities between sympatric and allopatric interactions, we investigated, in silico, the potential presence of *Schistosoma mansoni* miRNAs in our parasite RNAseq data. At 24 h post infections, we identified 54 miRNA precursors from miRBase with high-quality alignment scores against the different RNAseq read libraries (naïve BgBRE, BB24, BV24, BR24). To avoid cross-species misidentifications, we selected precursors that were exclusively identified in infected and never identified in uninfected snails (naive BgBRE). Eleven miRNA precursors corresponding to *Schistosoma mansoni* were identified (Fig. 7A). Eight of the parasite miRNA precursors were specific to the Brazil-infected libraries (BB24); two were specific of the Venezuela-infected libraries (BV24); and one was shared across the three infected conditions (BB24, BV24 and BR24). Although we identified 49 miRNA

precursor sequences specific to *S. mansoni* (Fig. 7B), we decided to select only miRNAs covered by

100% nucleotide similarity that allowed us to predict 11 miRNAs in mature (e.g., sma-mir-2d-3p,

sma-mir-190-3p) or precursor (sma-mir-8431) forms. Then, in order to identify candidate sequences

that could represent putative miRNA targets, we used the Miranda tool (Supplementary Table 4). Only

RNA-RNA interactions that showed good scores for pairing (> 140) and enthalpy (< 15 Kcal) were

considered. The number of targets pertaining to the differentially expressed immune-related transcripts

identified in Fig. 1 that were found for the identified miRNAs ranged from two targets for sma-mir-

8456,to 50 targets for sma-mir-2d of the differentially expressed immune-related transcripts identified

in Fig. 1.

The miRNAs identified under the sympatric condition (SmBRE) were predicted to target 43.5%

of the differentially represented immune-related transcripts identified in the RNAseq data (Fig. 1B,

Fig. 7) whereas 6.8% and 5.1% were targeted in allopatric conditions, SmVEN and Srod, respectively

(Fig. 1B, Fig. 7).To explain the similar prevalence between sympatric and allopatric infections we

focus our attention on miRNAs that were shared between sympatric and allopatric interactions and we

identified one miRNA:sma-miR-190-3p (Fig. 7 and Supplementary Table 4). This miRNA was

predicted to bind 17 different targets among which we identified different variants of the fibrinogen-

related protein (FREP) family and a cytotoxic/cytolytic humoral factor the biomphalysin. To go

further, we look at the expression of those candidates following infection. If FREP transcripts were

downregulated in sympatric interactions, it is not always the case in allopatry. However, interestingly

all biomphalysin transcripts were underrepresented in sympatric and allopatric interactions.

Altogether, these data suggest that the parasites might hijack the host immune response using

dedicated miRNAs.

Survival of snail following infection

To examine the potential impact of allopatric or sympatric parasites on snail survival, we investigated

the mortality rates of infected snails over 4 months. The survival rate was significantly higher for

noninfected snails compared to infected snails (sympatric interaction Kaplan-Meier Log Rank test p =

$1.39 \ e10^{-5}$ and allopatric interaction p = 0.0005). However, there was no significant difference in the

mortality rates of snails subjected to sympatric versus allopatric interactions: at the end of the

experiment, the survival rates were 72% and 65% for the allopatric and sympatric interactions,

respectively (Kaplan-Meier Log Rank test p = 0.243) (Supplementary Fig. 2).

Discussion

In the natural environment, it is assumed that the parasitic genes responsible for infectivity will evolve

alongside the host defence genes, resulting in adaptation of the interactions between local host and

parasite populations (59, 60). In this context, local/sympatric parasites were expected to display

greater infectiveness, reproductive success and virulence in host populations compared to

foreign/allopatric parasites (38, 39, 61, 62). However, in some cases this rule may be contradicted, as

certain allopatric parasite-host interactions have been reported to be significantly more compatible

than the corresponding sympatric combinations (40, 63). It appears that certain

Biomphalaria/Schistosoma interactions (64, 65) do not fulfil the "host-parasite local adaptation

hypothesis" (38, 39, 61, 62), in which the sympatric parasite is expected to perform better than the

allopatric one.

Using field data, Morand et al. (1996) and Prugnolle et al. (2006) showed that although sympatric

parasite-host combinations of schistosomes and snails do tend to be more compatible, exceptions exist

wherein particular allopatric combinations are equally or significantly more compatible. Similar

results were obtained when comparing the interactions of Brazilian and Guadeloupean snails versus

Schistosoma infections (42). The authors found that allopatric Guadeloupean parasites were not able to

infect Brazilian snails; but Brazilian parasites were able to infect the allopatric Guadeloupean snails.

Furthermore, this work demonstrated the presence of local adaptation between reactive oxygen species

(ROS) and ROS scavengers in this system (42). Based on these observations, we propose that it would

be important to develop integrative analysis to depict and understand the precise molecular crosstalk

(immunobiological interactions) occurring in such highly compatible sympatric and allopatric systems.

Thus, dual-comparative approaches were used herein to simultaneously analyse the responses of

Biomphalaria snails and *Schistosoma* parasites into sympatric or allopatric interactions displaying

similar compatibilities.

The present RNAseq analysis demonstrated that in sympatric interaction (BB) a huge

immunosuppression occurs. Twenty-four hours after the infection, the three immunological processes

were downregulated: (i) immune recognition, (ii) effector and (iii) signalling pathways (Fig. 1 and 2).

Conversely, in allopatric interactions (BV and BR), host immune response was activated (Fig. 1 and

2). Differentially regulated transcripts mostly belong to immune cellular activation, migration,

proliferation or differentiation (Fig. 2). An EdU-labelling was used to detect proliferation and to

confirm that more hemocyte proliferation was observed in allopatry compared to sympatry (Fig. 3, 4A

and 4B). Using flow cytometry, we demonstrated that a new hemocyte subpopulation (named P2) was

observed exclusively following allopatric infection at 24 h (Fig. 4C and 4D). P2 was EdU-positive and

characterized by an increase in granularity, indicating that the new P2 cell subtype could proliferate.

In Biomphalaria snails, we know three main hemocyte morphotypes: the Blast-like cells, the

type I hyalinocytes and the granulocytes (66). However, in absence of specific hemocyte markers, it is

impossible to analyse which hemocyte morphotypes are proliferating in our flow cytometry results

(Figs. 4C, 4D). The P2 subpopulation would thus originate from either a morphological change in an

existing subset (correlating potentially with a decline in the P1 population), or represent cells that are

migrating from tissues or a hematopoietic organ to reach the hemolymph. Indeed the P2 population

reflects newly proliferated cells that present higher EdU positive cells than the P1 population (Fig. 4C,

4D). Further investigations will be necessary to determine the origin of the P2 population. However,

hemocyte proliferation decreased more rapidly in sympatric versus allopatric interactions (Fig. 3 and

4). By comparing different sympatric and allopatric interactions, we demonstrated that the cellular or

molecular phenotype observed may refers to reciprocal evolution or adaptation rather to a simple host

or parasite strain effect.

The strong immunosuppression observed in sympatric interaction compared to the immune

cell activation in allopatry certainly results in differences in the capabilities of sympatric and allopatric

parasites to infect and grow in such hosts. This hypothesis was confirmed by our histological analysis

of sporocyst development in snail tissues. We observed a significant difference in sporocyst size 24 h

after infection (Fig. 5), when sympatric sporocysts were one-third bigger than allopatric sporocysts. At

96 h after infection, in contrast, there was no size difference between sympatric and allopatric

parasites (Fig. 5). This indicates that the allopatric parasites seem to overcome this early delay in

development. However, the consequences of this delay for duration of cercariae development, number

of cercariae produced, cercariae infectivity and pathogenicity in vertebrate host, will deserves further

investigation to determine the potential fitness cost between sympatric and allopatric parasites.

To go further and understand how parasites immunosuppress the host or circumvent the host

immune system, we used a dual-RNAseq approach to investigate transcripts expression of the

sympatric and allopatric parasite intra-molluscal stages (Fig. 6). As the histological differences were

solely observed at 24 h after infection, we used dual-RNAseq to investigate the parasite expression

patterns at the same time point of infection. Most of the parasite transcripts belonged to the processes

of nucleotide metabolism, transcription, translation and cell differentiation, development and growth.

We also identified some transcripts with GO terms or functions related to immunomodulation or

immunosuppression (Fig. 6 and Supplementary Table 4). Nearly all of the identified transcripts were

overrepresented in the sympatric interaction compared to the allopatric interactions. Our results

therefore suggest that the installation, development and growth of the parasite occurred much more

rapidly in the sympatric combination, as sympatric parasites seemed to interfere more efficiently with

the host immune system. However, RNAseq data did not give any clear information about how

allopatric parasites succeed in circumventing the host immune system. We thus next examined the

generated dual-RNAseq libraries in an effort to identify whether sympatric and/or allopatric

schistosomes could hijack the host immune system using microRNAs. To achieve this goal, we

confronted the dual-RNAseq data to the *Schistosoma mansoni* subset of miRBase to identify the

presence of microRNAs (miRNAs) in our datasets. miRNAs are known to regulate numerous

biological processes, including key immune response genes (67, 68). Recent work has demonstrated

that circulating small noncoding RNAs from parasites have hijack roles against host metabolism,

especially in the interaction of schistosomes with their vertebrate hosts (69-71). Such noncoding

RNAs could act as exogenous miRNAs to interfere with or circumvent the host immune system. At 24

h after infection, several differentially expressed parasite miRNAs were identified. We investigated

the potential role played by these miRNAs on host immune response. We identified some predicted

targets of such schistosome miRNAs in the *Biomphalaria* immune reference transcriptome. We found

that schistosome miRNAs identified in the snails 24 h after sympatric infection may target 43.5% of

the differentially regulated immune transcripts (Fig. 7). In contrast, far fewer correspondences were

identified for the allopatric interactions (Fig. 7). The higher proportion of targeted genes in the

sympatric interaction may result in the observed efficient immunosuppression, demonstrating a

specific co-evolution or adaptation in the transcriptional regulation between sympatric host and

parasite. However, even if more host immune genes appeared to be targeted in the sympatric

combination compared to the allopatric combination (Fig. 7), both sympatric and allopatric

interactions displayed the same phenotype of compatibility. In the allopatric condition, despite the

activation of the snail immune response, the parasite is able to develop and succeed to infect the host.

This similarity in compatibility phenotype between sympatric and allopatric parasites could potentially

results from their ability to target host immune weapons or host genes that regulate the snail innate

cellular response. In this context, we were expecting to find miRNAs shared between sympatric and

allopatric parasites, that is, miRNAs that tend to weaken the immune system. A unique miRNA was

found to be shared by allopatric and sympatric parasites, sma-miR-190-3p, which has been identified

and predicted to bind different targets among FREPs and biomphalysin. The FREP family members

are known as pathogen recognition receptors (64, 65)). Moreover, FREP knockdown was associated

with an increase of snail compatibility toward *Schistosoma* infections (72, 73). Biomphalysin is a key

humoral factor involved in cytotoxic/cytolytic activities against *Schistosoma* parasites with the ability

to bind miracidia and sporocyst surfaces (17, 74). Moreover, the associated snail transcripts to these

molecules are mostly down represented in sympatric and allopatric interactions. The miRNA sma-

miR-190-3p is perhaps essential for parasite compatibility. Parasites expressing such miRNA would

thus be considered as highly virulent parasites with strong infecting capabilities. By producing

dedicated miRNAs, the parasites were potentially able to regulate transcriptional, post-transcriptional,

translational and protein stability processes that might help them to subvert the snail's immune

defences.

Compatibility reflects the outcome of complex immunobiological interactions and depends on:

(i) the ability of the snail immune system to recognize and kill the parasite; and (ii) the ability of the

parasite to circumvent or evade the host immune response (20, 53, 75). Based on the present

observations, we propose that sympatric and allopatric interactions trigger totally different responses. In the sympatric interaction, the parasite is able to induce a host immunosuppression within the first day of infection, enabling it to infect quickly the host and readily begins its development. In the allopatric interaction, the parasite is not able to neutralize quickly the host immune system, and therefore the parasite is recognized by the host defence system, which mounts a potent immune response. The need for allopatric parasites to resist the immune system seems to disrupt the activation of their developmental program during the first day of infection. However, they seemed to be rapidly able to protect themselves against the host immune response and develop normally in snail tissues 96 h post-infection. Thereafter, in the medium- or long-term, there are no observable differences in the prevalence, intensity or snail survival comparing sympatric and allopatric interactions (Supplementary Table 1, Supplementary Fig. 2).

Thus, we show that despite having similar prevalence phenotypes, sympatric and allopatric snail-*Schistosoma* interactions displayed a very different immunobiological dialogue at the molecular level. Intriguingly, these different immunobiological interactions seem to have no repercussions upon parasite growth at longer term or to host survival. As differences at the molecular level do not correspond apparently to any ecologically meaningful changes in term of fitness, it is not straightforward to demonstrate local adaptation in such systems. However, we do not know if fitness costs could affect other biological traits in sympatric and allopatric interactions, for example, sporocyst II or cercariae production, or cercariae infectivity and pathogenicity towards the vertebrate host. Demonstrating local adaptation would thus appear extremely complex and would indeed deserve further investigation. It is hard to draw the line as to when local adaptation is or is not present. However, our results argue that the differences found at the molecular level may ultimately contribute to the evolution of local adaptation at an ecological level. Nevertheless, the ability for allopatric pathogens to adapt rapidly and efficiently to new hosts could have critical consequences on disease emergence and risk of schistosomiasis outbreaks.

Past events of allopatric parasites reaching new areas of transmission, even in a large geographic scale dispersion, have been largely documented, the most famous example being the schistosomiasis colonization of South America since the slave trade of the 16th through 19th centuries

(76, 77). *Schistosoma* originated in Asia, reached Africa 12 to 19 million years ago (MYA), and gave rise to all *Schistosoma* species known in Africa (77). *S. mansoni* diverged from *S. rodhaini* around 2.8 MYA (76, 78), and thereafter, 400 to 500 years ago, colonized South America (76, 77). This colonization of South America by *S. mansoni* from Africa was rendered possible by the presence of the snail host *Biomphalaria glabrata*. All African species of *Biomphalaria* are monophyletic and seem to have originated from paraphyletic South American clades (79-81). The ancestor of *B. glabrata* appears to have colonized Africa 1 to 5 MYA, giving rise to all 12 species of *Biomphalaria* known today in Africa (82). In South America and the Caribbean islands, *S. mansoni* infects *B. glabrata*; in Africa, it infects mostly *B. pfeifferi* and *B. alexandrina*. We found that South American *S. mansoni* parasites are highly compatible with their sympatric South American snail hosts, whereas African *S. mansoni* parasites display low compatibility phenotype with South American snail hosts (Supplementary Table 1). Interestingly, the South American parasites did not lose their compatibility for African snail hosts; that is, prevalence is similar to African parasites when confronted with African snails (Supplementary Table 1). The recent African origin of South American *Schistosoma* parasites (introduction in South America 400 to 500 years ago) may explain that they have not diverged sufficiently in South America to lose their compatibility for African snail hosts. In this case, the transfer of allopatric parasites from Africa to South American snail hosts have been successful and resulted in the emergence of schistosomiasis in South America.

More recently another case of compatible allopatric parasite emergence have been observed when schistososmiasis reached Europe (83, 84). Here, humans infected in Senegal have imported a hybrid between *Schistosoma haematobium* and *Schistosoma bovis* into Corsica. In this case, urogenital schistosomiasis could be introduced and easily and rapidly spread into this novel area of south Corsica because *Bulinus truncatus* the vector snail of *S. haematobium* was endemic in the Corsica Cavu River (83, 84). However, this allopatric African hybrid parasite was able to adapt efficiently to the new Corsican *B. truncatus* host. If parasite hybridization can potentially have a putative role in increasing the colonization potential of such *S. haematobium*, it would be particularly interesting to analyse and depict the molecular support of such allopatric interactions to predict the potential risk of schistosomiasis outbreaks in other European areas, or other potential transmission foci.

Conclusions

669   If we hope to draw conclusions regarding the existence of emerging or outbreak risks, we need to develop integrative approaches to explore fine-scale patterns of host-parasite interactions. We must consider the spatial scale at which comparisons are conducted, the patterns of disease occurrence, the population genetics, and the involvement of physiological, immunological and molecular processes. Studying the relevant factors at the relevant timing would be of critical importance in terms of schistosomiasis control. Understanding further, how these allopatric parasites efficiently infect host snails would be mandatory to identify markers and develop new tools to predict or to quantify risks of schistosomiasis outbreaks.

Acknowledgements

We thank Cécile Saint-Béat, Nathalie Arancibia and Anne Rognon for their work and diligence in helping generate some of the data described herein.

Declarations

Ethics approval and consent to participate

Our laboratory holds permit # A66040 for experiments on animals, which was obtained from the French Ministry of Agriculture and Fisheries and the French Ministry of National Education, Research, and Technology. The housing, breeding, and care of the utilized animals followed the ethical requirements of our country. The experimenter possesses an official certificate for animal experimentation from both of the above-listed French ministries (Decree # 87–848, October 19, 1987). The various protocols used in this study have been approved by the French veterinary agency of the DRAAF Languedoc-Roussillon (Direction Régionale de l'Alimentation, de l'Agriculture et de la Forêt), Montpellier, France (authorization # 007083).

Availability of data and material

The datasets generated and/or analysed during the current study are not publicly available due to ongoing analysis of transcriptome but are available from the corresponding author on reasonable request.

Raw read fastq files were submitted to the Sequence Read Archive at NCBI (http://trace.ncbi.nlm.nih.gov/Traces/sra/) under the reference PRJNA213050. The reference transcriptome is publicly available from the transcriptomic database of *Biomphalaria glabrata* at the IHPE website (http://ihpe.univ-perp.fr/acces-aux-donnees/)

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the French National Agency for Research (ANR) through the Invimory project grant [ANR-13-JSV7-0009] to BG. The funding agency had no role in study design, collection, analysis or interpretation of data and in writing of the manuscript.

Authors' contributions

Conceived and designed the experiments: AP, SP, RG, DD, BG. Performed the experiments: AP, SP, CC, GMC, JFA. Analysed the data: AP, SP, CC, RG, GMC, DD, BG. AP, RG, BG led the manuscript writing. All authors participated in manuscript writing, editing and critical reviewing. All authors read and approved the final manuscript.

Authors' information

SP current position Welcome Trust Sanger Institute, CB10 1SA Hinxton, United Kingdom

References:

1. WHO. 2010. Schistosomiasis *Fact Sheet No. 115. World Health Organisation.*

2. Basch, P. F. 1976. Intermediate host specificity in Schistosoma mansoni. *Experimental parasitology* 39:150-169.

3. Mitta, G., C. M. Adema, B. Gourbal, E. S. Loker, and A. Theron. 2012. Compatibility polymorphism in snail/schistosome interactions: From field to theory to molecular mechanisms. *Developmental and comparative immunology* 37:1-8.

4. Theron, A., J. R. Pages, and A. Rognon. 1997. Schistosoma mansoni: distribution patterns of miracidia among Biomphalaria glabrata snail as related to host susceptibility and sporocyst regulatory processes. *Experimental parasitology* 85:1-9.

5. Theron, A., A. Rognon, B. Gourbal, and G. Mitta. 2014. Multi-parasite host susceptibility and multi-host parasite infectivity: a new approach of the Biomphalaria glabrata/Schistosoma mansoni compatibility polymorphism. *Infection Genetics and Evolution* 26:80-88.

6. Galinier, R., E. Roger, Y. Mone, D. Duval, A. Portet, S. Pinaud, C. Chaparro, C. Grunau, C. Genthon, E. Dubois, A. Rognon, N. Arancibia, B. Dejean, A. Theron, B. Gourbal, and G. Mitta. 2017. A multistrain approach to studying the mechanisms underlying compatibility in the interaction between Biomphalaria glabrata and Schistosoma mansoni. *PLoS neglected tropical diseases* 11:e0005398.

7. Mitta, G., B. Gourbal, C. Grunau, M. Knight, J. M. Bridger, and A. Theron. 2017. The Compatibility Between Biomphalaria glabrata Snails and Schistosoma mansoni: An Increasingly Complex Puzzle. *Adv Parasitol* 97:111-145.

8. Bonner, K. M., C. J. Bayne, M. K. Larson, and M. S. Blouin. 2012. Effects of Cu/Zn superoxide dismutase (sod1) genotype and genetic background on growth, reproduction and defense in Biomphalaria glabrata. *PLoS neglected tropical diseases* 6:e1701.

9. Goodall, C. P., R. C. Bender, J. K. Brooks, and C. J. Bayne. 2006. Biomphalaria glabrata cytosolic copper/zinc superoxide dismutase (SOD1) gene: association of SOD1 alleles with resistance/susceptibility to Schistosoma mansoni. *Molecular and biochemical parasitology* 147:207-210.

10. Tennessen, J. A., A. Theron, M. Marine, J. Y. Yeh, A. Rognon, and M. S. Blouin. 2015. Hyperdiverse gene cluster in snail host conveys resistance to human schistosome parasites. *PLoS genetics* 11:e1005067.

11. Ittiprasert, W., A. Miller, J. Myers, V. Nene, N. M. El-Sayed, and M. Knight. 2010. Identification of immediate response genes dominantly expressed in juvenile resistant and susceptible Biomphalaria glabrata snails upon exposure to Schistosoma mansoni. *Molecular and biochemical parasitology* 169:27-39.

12. Lockyer, A. E., A. M. Emery, R. A. Kane, A. J. Walker, C. D. Mayer, G. Mitta, C. Coustau, C. M. Adema, B. Hanelt, D. Rollinson, L. R. Noble, and C. S. Jones. 2012. Early differential gene expression in haemocytes from resistant and susceptible Biomphalaria glabrata strains in response to Schistosoma mansoni. *PLoS One* 7:e51102.

13. Lockyer, A. E., L. R. Noble, D. Rollinson, and C. S. Jones. 2004. Schistosoma mansoni: resistant specific infection-induced gene expression in Biomphalaria glabrata identified by fluorescent-based differential display. *Experimental parasitology* 107:97-104.

769   14.   Lockyer, A. E., J. N. Spinks, A. J. Walker, R. A. Kane, L. R. Noble, D. Rollinson, E. Dias-
770   Neto, and C. S. Jones. 2007. Biomphalaria glabrata transcriptome: identification of
771   cell-signalling, transcriptional control and immune-related genes from open reading
772   frame expressed sequence tags (ORESTES). *Developmental and comparative*
773   *immunology* 31:763-782.
774   15.   Roger, E., C. Grunau, R. J. Pierce, H. Hirai, B. Gourbal, R. Galinier, R. Emans, I. M.
775   Cesari, C. Cosseau, and G. Mitta. 2008. Controlled Chaos of Polymorphic Mucins in a
776   Metazoan Parasite (Schistosoma mansoni) Interacting with Its Invertebrate Host
777   (Biomphalaria glabrata). *PLoS neglected tropical diseases* 2:e330.
778   16.   Bender, R. C., C. P. Goodall, M. S. Blouin, and C. J. Bayne. 2007. Variation in
779   expression of Biomphalaria glabrata SOD1: a potential controlling factor in
780   susceptibility/resistance to Schistosoma mansoni. *Developmental and comparative*
781   *immunology* 31:874-878.
782   17.   Galinier, R., J. Portela, Y. Mone, J. F. Allienne, H. Henri, S. Delbecq, G. Mitta, B.
783   Gourbal, and D. Duval. 2013. Biomphalysin, a new beta pore-forming toxin involved
784   in Biomphalaria glabrata immune defense against Schistosoma mansoni. *PLoS*
785   *pathogens* 9:e1003216.
786   18.   Hahn, U. K., R. C. Bender, and C. J. Bayne. 2001. Involvement of nitric oxide in killing
787   of Schistosoma mansoni sporocysts by hemocytes from resistant Biomphalaria
788   glabrata. *Journal of Parasitology* 87:778-785.
789   19.   Larson, M. K., R. C. Bender, and C. J. Bayne. 2014. Resistance of Biomphalaria
790   glabrata 13-16-R1 snails to Schistosoma mansoni PR1 is a function of haemocyte
791   abundance and constitutive levels of specific transcripts in haemocytes. *International*
792   *journal for parasitology* 44:343-353.
793   20.   Mone, Y., B. Gourbal, D. Duval, L. Du Pasquier, S. Kieffer-Jaquinod, and G. Mitta.
794   2010. A large repertoire of parasite epitopes matched by a large repertoire of host
795   immune receptors in an invertebrate host/parasite model. *PLoS neglected tropical*
796   *diseases* 4.
797   21.   Myers, J., W. Ittiprasert, N. Raghavan, A. Miller, and M. Knight. 2008. Differences in
798   cysteine protease activity in Schistosoma mansoni-resistant and -susceptible
799   Biomphalaria glabrata and characterization of the hepatopancreas cathepsin B Full-
800   length cDNA. *Journal of Parasitology* 94:659-668.
801   22.   Pila, E. A., M. A. Gordy, V. K. Phillips, A. L. Kabore, S. P. Rudko, and P. C. Hanington.
802   2016. Endogenous growth factor stimulation of hemocyte proliferation induces
803   resistance to Schistosoma mansoni challenge in the snail host. *Proceedings of the*
804   *National Academy of Sciences of the United States of America* 113:5305-5310.
805   23.   Roger, E., G. Mitta, Y. Mone, A. Bouchut, A. Rognon, C. Grunau, J. Boissier, A. Theron,
806   and B. E. Gourbal. 2008. Molecular determinants of compatibility polymorphism in
807   the Biomphalaria glabrata/Schistosoma mansoni model: New candidates identified
808   by a global comparative proteomics approach. *Molecular and biochemical*
809   *parasitology* 157:205-216.
810   24.   Bridger, J. M., H. D. Arican-Gotkas, H. A. Foster, L. S. Godwin, A. Harvey, I. R. Kill, M.
811   Knight, I. S. Mehta, and M. H. Ahmed. 2014. The non-random repositioning of whole
812   chromosomes and individual gene loci in interphase nuclei and its relevance in
813   disease, infection, aging, and cancer. *Advances in experimental medicine and biology*
814   773:263-279.

25. Ittiprasert, W., and M. Knight. 2012. Reversing the resistance phenotype of the
Biomphalaria glabrata snail host Schistosoma mansoni infection by temperature
modulation. *PLoS pathogens* 8:e1002677.
26. Ittiprasert, W., R. Nene, A. Miller, N. Raghavan, F. Lewis, J. Hodgson, and M. Knight.
2009. Schistosoma mansoni infection of juvenile Biomphalaria glabrata induces a
differential stress response between resistant and susceptible snails. *Experimental*
parasitology 123:203-211.
27. Knight, M., O. Elhelu, M. Smith, B. Haugen, A. Miller, N. Raghavan, C. Wellman, C.
Cousin, F. Dixon, V. Mann, G. Rinaldi, W. Ittiprasert, and P. J. Brindley. 2015.
Susceptibility of Snails to Infection with Schistosomes is influenced by Temperature
and Expression of Heat Shock Proteins. *Epidemiology* 5:pii: 189. Epub.
28. Nelson, M. K., B. C. Cruz, K. L. Buena, H. Nguyen, and J. T. Sullivan. 2016. Effects of
abnormal temperature and starvation on the internal defense system of the
schistosome-transmitting snail Biomphalaria glabrata. *Journal of invertebrate*
pathology.
29. Perrin, C., J. M. Lepesant, E. Roger, D. Duval, S. Fneich, V. Thuillier, J. F. Alliene, G.
Mitta, C. Grunau, and C. Cosseau. 2012. Schistosoma mansoni mucin gene
(SmPoMuc) expression: epigenetic control to shape adaptation to a new host. *PLoS*
pathogens 9:e1003571.
30. Portet, A., S. Pinaud, G. Tetreau, R. Galinier, C. Cosseau, D. Duval, C. Grunau, G.
Mitta, and B. Gourbal. 2017. Integrated multi-omic analyses in Biomphalaria-
Schistosoma dialogue reveal the immunobiological significance of FREP-SmPoMuc
interaction. *Developmental and comparative immunology* doi:
10.1016/j.dci.2017.02.025.
31. Osnas, E. E., P. J. Hurtado, and A. P. Dobson. 2015. Evolution of pathogen virulence
across space during an epidemic. *The American naturalist* 185:332-342.
32. Tack, A. J., P. H. Thrall, L. G. Barrett, J. J. Burdon, and A. L. Laine. 2012. Variation in
infectivity and aggressiveness in space and time in wild host-pathogen systems:
causes and consequences. *Journal of evolutionary biology* 25:1918-1936.
33. Prugnolle, F., T. de Meeus, J. P. Pointier, P. Durand, A. Rognon, and A. Theron. 2006.
Geographical variations in infectivity and susceptibility in the host-parasite system
Schistosoma mansoni/Biomphalaria glabrata: no evidence for local adaptation.
Parasitology 133:313-319.
34. Richards, C. S. 1975. Genetic factors in susceptibility of Biomphalaria glabrata for
different strains of Schistosoma mansoni. *Parasitology* 70:231-241.
35. Frank, S. A. 1996. Models of parasite virulence. *The Quarterly review of biology*
71:37-78.
36. Kawecki, T., and D. Ebert. 2004. Conceptual issues in local adaptation. *Ecology letters*
7:1225-1241.
37. Kalbe, M., C. Eizaguirre, J. P. Scharsack, and P. J. Jakobsen. 2016. Reciprocal cross
infection of sticklebacks with the diphyllobothriidean cestode Schistocephalus solidus
reveals consistent population differences in parasite growth and host resistance.
Parasites & vectors 9:130.
38. Ebert, D. 1994. Virulence and local adaptation of horizontally transmitted parasite.
Science 265:1084-1086.
39. Lively, C. M. 1989. Adaptation by a Parasitic Trematode to Local Populations of Its
Snail Host. *Evolution; international journal of organic evolution* 43(8).

40. Morand, S., S. D. Manning, and M. E. Woolhouse. 1996. Parasite-host coevolution
and geographic patterns of parasite infectivity and host susceptibility. *Proceedings of*
the Royal Society B 263:119-128.
41. Osnas, E. E., and C. M. Lively. 2004. Parasite dose, prevalence of infection and local
adaptation in a host-parasite system. *Parasitology* 128:223-228.
42. Mone, Y., A. C. Ribou, C. Cosseau, D. Duval, A. Theron, G. Mitta, and B. Gourbal. 2011.
An example of molecular co-evolution: reactive oxygen species (ROS) and ROS
scavenger levels in Schistosoma mansoni/Biomphalaria glabrata interactions.
International journal for parasitology 41:721-730.
43. Penczykowski, R. M., A. L. Laine, and B. Koskella. 2015. Understanding the ecology
and evolution of host-parasite interactions across scales. *Evolutionary applications*
9:37-52.
44. Robertson, S., J. E. Bradley, and A. D. MacColl. 2017. No evidence of local adaptation
of immune responses to Gyrodactylus in three-spined stickleback (Gasterosteus
aculeatus). *Fish and Shellfish Immunology* 60:275-281.
45. Bankers, L., P. Fields, K. E. McElroy, J. L. Boore, J. M. Logsdon, Jr., and M. Neiman.
2018. Genomic evidence for population-specific responses to co-evolving parasites in
a New Zealand freshwater snail. *Mol Ecol* 26:3663-3675.
46. Kincaid-Smith, J., O. Rey, E. Toulza, A. Berry, and J. Boissier. 2017. Emerging
Schistosomiasis in Europe: A Need to Quantify the Risks. *Trends in Parasitology*
33:600-609.
47. Dheilly, N. M., D. Duval, G. Mouahid, R. Emans, J. F. Allienne, R. Galinier, C. Genthon,
E. Dubois, L. Du Pasquier, C. M. Adema, C. Grunau, G. Mitta, and B. Gourbal. 2015. A
family of variable immunoglobulin and lectin domain containing molecules in the
snail Biomphalaria glabrata. *Developmental and comparative immunology* 48:234-
243.
48. Pinaud, S., J. Portela, D. Duval, F. C. Nowacki, M. A. Olive, J. F. Allienne, R. Galinier, N.
M. Dheilly, S. Kieffer-Jaquinod, G. Mitta, A. Theron, and B. Gourbal. 2016. A Shift
from Cellular to Humoral Responses Contributes to Innate Immune Memory in the
Vector Snail Biomphalaria glabrata. *PLoS pathogens* 12:e1005361.
49. Sminia, T., and L. Barendsen. 1980. A comparative morphological and enzyme
histochemical study on blood cells of the freshwater snails, *Lymnaea stagnalis*,
Biomphalaria glabrata, and *Bulinus truncatus*. *Journal of Morphology* 165:31-39.
50. Zhao, X., H. Yu, L. Kong, S. Liu, and Q. Li. 2016. High throughput sequencing of small
RNAs transcriptomes in two Crassostrea oysters identifies microRNAs involved in
osmotic stress response. *Scientific reports* 6:22687.
51. Enright, A. J., B. John, U. Gaul, T. Tuschl, C. Sander, and D. S. Marks. 2003. MicroRNA
targets in Drosophila. *Genome Biology* 5:R1.
52. Baeza Garcia, A., R. J. Pierce, B. Gourbal, E. Werkmeister, D. Colinet, J. M. Reichhart,
C. Dissous, and C. Coustau. 2010. Involvement of the cytokine MIF in the snail host
immune response to the parasite Schistosoma mansoni. *PLoS pathogens* 6:e1001115.
53. Guillou, F., E. Roger, Y. Mone, A. Rognon, C. Grunau, A. Theron, G. Mitta, C. Coustau,
and B. E. Gourbal. 2007. Excretory-secretory proteome of larval Schistosoma
mansoni and Echinostoma caproni, two parasites of Biomphalaria glabrata. *Molecular*
and biochemical parasitology.
54. Raghavan, M., S. J. Wijeyesakere, L. R. Peters, and N. Del Cid. 2013. Calreticulin in the
immune system: ins and outs. *Trends in Immunology* 34:13-21.

55. Ryu, S. J., and S. C. Park. 2009. Targeting major vault protein in senescence-
associated apoptosis resistance. *Expert Opinion on Therapeutic Targets* 13:479-484.
56. Gilges, D., M. A. Vinit, I. Callebaut, L. Coulombel, V. Cacheux, P. H. Romeo, and I.
Vigon. 2000. Polydom: a secreted protein with pentraxin, complement control
protein, epidermal growth factor and von Willebrand factor A domains. *Biochemical*
Journal 352 Pt 1:49-59.
57. Bohdanowicz, M., D. Schlam, M. Hermansson, D. Rizzuti, G. D. Fairn, T. Ueyama, P.
Somerharju, G. Du, and S. Grinstein. 2013. Phosphatidic acid is required for the
constitutive ruffling and macropinocytosis of phagocytes. *Molecular Biology of the*
Cell 24:1700-1712, S1701-1707.
58. Grkovich, A., and E. A. Dennis. 2009. Phosphatidic acid phosphohydrolase in the
regulation of inflammatory signaling. *Advances in Enzyme Regulation* 49:114-120.
59. Dybdahl, M. F., and A. Storfer. 2003. Parasite local adaptation: Red Queen versus
suicide king. *TRENDs in Ecology and Evolution* 18:523-530.
60. Osnas, E. E., and C. M. Lively. 2005. Immune response to sympatric and allopatric
parasites in a snail-trematode interaction. *Frontiers in Zoology* 2:8.
61. Ebert, D. 1998. Experimental evolution of parasites. *Science* 282:1432-1435.
62. Minchella, D. J. 1985. Host life-history variation in response to parasitism.
Parasitology 90:205-216.
63. Greischar, M. A., and B. Koskella. 2007. A synthesis of experimental work on parasite
local adaptation. *Ecology letters* 10:418-434.
64. Adema, C. M., L. A. Hertel, R. D. Miller, and E. S. Loker. 1997. A family of fibrinogen-
related proteins that precipitates parasite-derived molecules is produced by an
invertebrate after infection. *Proceedings of the National Academy of Sciences of the*
United States of America 94:8691-8696.
65. Zhang, H., L. Song, C. Li, J. Zhao, H. Wang, Q. Gao, and W. Xu. 2007. Molecular cloning
and characterization of a thioester-containing protein from Zhikong scallop Chlamys
farreri. *Molecular Immunology* 44:3492-3500.
66. Cavalcanti, M. G., F. C. Filho, A. M. Mendonca, G. R. Duarte, C. C. Barbosa, C. M. De
Castro, L. C. Alves, and F. A. Brayner. 2012. Morphological characterization of
hemocytes from Biomphalaria glabrata and Biomphalaria straminea. *Micron* 43:285-
291.
67. Li, S., L. Shen, L. Sun, J. Xu, P. Jin, L. Chen, and F. Ma. 2017. Small RNA-Seq analysis
reveals microRNA-regulation of the Imd pathway during Escherichia coli infection in
Drosophila. *Developmental and comparative immunology* 70:80-87.
68. Zawada, A. M., L. Zhang, I. E. Emrich, K. S. Rogacev, N. Krezdorn, B. Rotter, D. Fliser, Y.
Devaux, L. Ziegler-Heitbrock, and G. H. Heine. 2017. MicroRNA profiling of human
intermediate monocytes. *Immunobiology* 222:587-596.
69. Meningher, T., G. Lerman, N. Regev-Rudzki, D. Gold, I. Z. Ben-Dov, Y. Sidi, D. Avni, and
E. Schwartz. 2017. Schistosomal MicroRNAs Isolated From Extracellular Vesicles in
Sera of Infected Patients: A New Tool for Diagnosis and Follow-up of Human
Schistosomiasis. *Journal of Infectious Diseases* 215:378-386.
70. Nowacki, F. C., M. T. Swain, O. I. Klychnikov, U. Niazi, A. Ivens, J. F. Quintana, P. J.
Hensbergen, C. H. Hokke, A. H. Buck, and K. F. Hoffmann. 2015. Protein and small
non-coding RNA-enriched extracellular vesicles are released by the pathogenic blood
fluke Schistosoma mansoni. *Journal of Extracellular Vesicles* 4:28665.

71. Walzer, K. A., and J. T. Chi. 2017. Trans-kingdom small RNA transfer during host-
pathogen interactions: The case of P. falciparum and erythrocytes. *RNA Biology*:1-8.

72. Hanington, P. C., M. A. Forys, and E. S. Loker. 2012. A somatically diversified defense
factor, FREP3, is a determinant of snail resistance to schistosome infection. *PLoS*
neglected tropical diseases 6:e1591.

73. Jiang, Y., E. S. Loker, and S. M. Zhang. 2006. In vivo and in vitro knockdown of FREP2
gene expression in the snail Biomphalaria glabrata using RNA interference.
Developmental and comparative immunology 30:855-866.

74. Tetreau, G., S. Pinaud, A. Portet, R. Galinier, B. Gourbal, and D. Duval. 2017. Specific
Pathogen Recognition by Multiple Innate Immune Sensors in an Invertebrate. *Front*
Immunol 8:1249.

75. Van Der Knaap, W. P. W., and E. S. Loker. 1990. Immune mechanisms in trematod-
snail interactions. *Parasitology Today* 6:175-182.

76. Crellen, T., F. Allan, S. David, C. Durrant, T. Huckvale, N. Holroyd, A. M. Emery, D.
Rollinson, D. M. Aanensen, M. Berriman, J. P. Webster, and J. A. Cotton. 2016. Whole
genome resequencing of the human parasite Schistosoma mansoni reveals
population history and effects of selection. *Scientific reports* 6:20954.

77. Lockyer, A. E., P. D. Olson, P. Ostergaard, D. Rollinson, D. A. Johnston, S. W. Attwood,
V. R. Southgate, P. Horak, S. D. Snyder, T. H. Le, T. Agatsuma, D. P. McManus, A. C.
Carmichael, S. Naem, and D. T. Littlewood. 2003. The phylogeny of the
Schistosomatidae based on three genes with emphasis on the interrelationships of
Schistosoma Weinland, 1858. *Parasitology* 126:203-224.

78. Morgan, J. A., R. J. Dejong, G. O. Adeoye, E. D. Ansa, C. S. Barbosa, P. Bremond, I. M.
Cesari, N. Charbonnel, L. R. Correa, G. Coulibaly, P. S. D'Andrea, C. P. De Souza, M. J.
Doenhoff, S. File, M. A. Idris, R. N. Incani, P. Jarne, D. M. Karanja, F. Kazibwe, J. Kpikpi,
N. J. Lwambo, A. Mabaye, L. A. Magalhaes, A. Makundi, H. Mone, G. Mouahid, G. M.
Muchemi, B. N. Mungai, M. Sene, V. Southgate, L. A. Tchuente, A. Theron, F. Yousif,
E. M. Zanotti-Magalhaes, G. M. Mkoji, and E. S. Loker. 2005. Origin and diversification
of the human parasite Schistosoma mansoni. *Molecular ecology* 14:3889-3902.

79. Campbell, G., C. S. Jones, A. E. Lockyer, S. Hughes, D. Brown, L. R. Noble, and D.
Rollinson. 2000. Molecular evidence supports an african affinity of the neotropical
freshwater gastropod, Biomphalaria glabrata, say 1818, an intermediate host for
Schistosoma mansoni. *Proceedings of the Royal Society B* 267:2351-2358.

80. DeJong, R. J., J. A. Morgan, W. L. Paraense, J. P. Pointier, M. Amarista, P. F. Ayeh-
Kumi, A. Babiker, C. S. Barbosa, P. Bremond, A. Pedro Canese, C. P. de Souza, C.
Dominguez, S. File, A. Gutierrez, R. N. Incani, T. Kawano, F. Kazibwe, J. Kpikpi, N. J.
Lwambo, R. Mimpfoundi, F. Njiokou, J. Noel Poda, M. Sene, L. E. Velasquez, M. Yong,
C. M. Adema, B. V. Hofkin, G. M. Mkoji, and E. S. Loker. 2001. Evolutionary
relationships and biogeography of Biomphalaria (Gastropoda: Planorbidae) with
implications regarding its role as host of the human bloodfluke, Schistosoma
mansoni. *Molecular biology and evolution* 18:2225-2239.

81. Theron, A. 2005. Les schistosomes et leurs hôtes : apport des marqueurs
moléculaires à la connaissance de leur phylogéographie, épidémiologie et écologie.
Bulletin de la Société Zoologique de France 130 205-218.

82. Morgan, J. A., R. J. Dejong, S. D. Snyder, G. M. Mkoji, and E. S. Loker. 2001.
Schistosoma mansoni and Biomphalaria: past history and future trends. *Parasitology*
123 Suppl:S211-228.

83.   Boissier, J., S. Grech-Angelini, B. L. Webster, J. F. Allienne, T. Huyse, S. Mas-Coma, E.
1003   Toulza, H. Barre-Cardi, D. Rollinson, J. Kincaid-Smith, A. Oleaga, R. Galinier, J. Foata,
1004   A. Rognon, A. Berry, G. Mouahid, R. Henneron, H. Mone, H. Noel, and G. Mitta. 2016.
1005   Outbreak of urogenital schistosomiasis in Corsica (France): an epidemiological case
1006   study. *The Lancet Infectious Diseases* 16:971-979.
1007   84.   Boissier, J., H. Mone, G. Mitta, M. D. Bargues, D. Molyneux, and S. Mas-Coma. 2015.
1008   Schistosomiasis reaches Europe. *The Lancet Infectious Diseases* 15:757-758.
1009
1010
1011

 Figure Legends

1013
FIGURE 1. Dual-RNAseq of Biomphalaria immune-related transcripts. Among the differentially represented

transcripts, Blast2GO functional annotation allowed us to identify 336 transcripts that appeared to be related to

the *Biomphalaria* immune response. Abbreviations and colours: blue BB, sympatric interaction between BgBRE

and SmBRE; green BV, allopatric interaction between BgBRE and SmVEN; and red BR, allopatric interaction

between BgBRE and Srod. For each interaction 40 whole-snails are used, 20 pooled at 24 h and 20 at 96 h post-

infection. A) Venn diagram showing the relationships among the immune transcripts found to be differentially

expressed in the sympatric and allopatric interactions. B) Clustering of differentially represented immune

transcripts. Heatmap representing the profiles of the 336 differentially represented immune-related transcripts in

the BB, BV or BR interactions along the kinetic of infection (at 24 and 96 h). Each transcript is represented

once, and each line represents one transcript. Colours: yellow, overrepresented transcripts; purple,

underrepresented transcripts; and black, unchanged relative to levels in control naïve snails. C) Pie chart

showing the distribution of the selected immune-related transcripts across three immunological processes:

immune recognition (pink), immune effectors (brown) and immune signalling (blue). For each category and

interaction, the respective proportion of transcripts and the direction of the effect (over- or under-expression) are

indicated.

1029
FIGURE 2. Differentially represented immune-related transcripts in sympatric and allopatric interactions.

Cumulative expression [Log2FC (fold change) from DESeq2 analysis] of the immune-related transcripts

identified as being differentially represented following sympatric or allopatric infection. Transcripts were

grouped into the three immunological groups described in Fig. 1, and from there into functional categories. The

yellow histograms correspond to cumulatively overrepresented transcripts, while the purple histograms show

underrepresented transcripts. The black (overrepresented) and gray (underrepresented) diamonds correspond to

the number of transcripts analysed in each functional category. Abbreviations: BB, BgBRE/SmBRE interaction;

BV, BgBRE/SmVEN interaction; and BR, BgBRE/Srod interaction. A. Immune transcript expression at 24 h

post-infection. B. Immune transcript expression at 96 h post-infection.

1039
FIGURE 3. Microscopic analysis of snail hemocyte proliferations. In vitro EdU-labelling of hemocytes was

conducted for sympatric and allopatric interactions. A) Hemocytes were collected at 24 h post-infection for in

vitro analysis. Confocal microscopy of EdU-labeled hemocytes from snails subjected to the allopatric BV

interaction (BgBRE/SmVEN). Colours: blue, DAPI; and green, EdU. B) Microscopic counting of EdU-labelled

hemocytes from naïve control snails (BgBRE) (n = 1,811) and those subjected to the sympatric interaction (BB:

BgBRE/SmBRE) (n=2,064) or an allopatric interaction (BV: BgBRE/SmVEN) (n = 1,366) recovered from three

individual snails by condition. Colours: green, EdU-positive cells; and blue, EdU-negative cells. Between-group

differences in the percentage of proliferation were tested using a Fisher exact test, with statistical significance

accepted at p < 0.05. The "a" indicates a significant difference between the naïve and infective conditions, while

"b" indicates a significant difference between the infective conditions.

1050
1051 **FIGURE 4.** Flow cytometry analysis of the hemocyte response in sympatric and allopatric interactions. A) Flow

1052 cytometry was used to count in vivo EdU-labelled hemocytes at 24 h and 96 h after infection in sympatric and

1053 allopatric interactions. A total number of hemocytes of n = 10,000 were recovered for six biological replicates of

three snails. Control naïve snails (BgBRE, yellow) were compared to those subjected to the sympatric interaction (BB, BgBRE/SmBRE, blue) or an allopatric interaction (BV, BgBRE/SmVEN, green). B) The experiment described in A was repeated using the BgVEN snail strain. Control naïve snails (BgVEN, yellow) were compared to those subjected to the sympatric interaction (VV, BgVEN/SmVEN, green) or an allopatric interaction (VB, BgVEN/SmBRE, blue). C) FSC (forward-scattered light, representing cell size) and SSC (side-scattered light, representing cell granularity) circulating hemocyte patterns in BgBRE snails under the naïve condition (yellow) or 24 h and 96 h after infections in sympatry (BB24/96, BgBRE/SmBRE, blue) or allopatry (BV24/96, BgBRE/SmVEN, green). D) FSC and SSC circulating hemocyte patterns in BgVEN snails under the naïve condition (yellow) or 24 h and 96 h after infections in sympatry (VV24/96, BgVEN/SmVEN, blue) or allopatry (VB24/96, BgVEN/SmBRE, green). The red dots correspond to EdU-positive hemocytes. Between-group differences in the percentage of proliferation were tested using the Mann-Whitney U-test, with statistical significance accepted at $p < 0.05$. The "a" indicates a significant difference between the naïve and infective condition, "b" indicates a significant difference between the infective conditions at 24h, and "c" indicates a significant difference between the infective conditions at 96h.

FIGURE 5. Development of parasites into snail tissues. A histological approach was used to monitor parasite size along the course of snail infection. The sympatric interaction (BgBRE x SmBRE) is shown in blue, and the allopatric interaction (BgBRE x SmVEN) is shown in green. For each experimental interaction, the parasite sizes were quantified at 24 and 96 h after infection. Morpho-anatomical aspects of the parasite are depicted to highlight a potential difference in parasite survival. N = 7 to 10 sporocystes were used. Between-group parasite size differences were assessed using the Mann-Whitney U-test, with significance accepted at $p < 0.05$ (indicated by "a" on the histograms).

FIGURE 6. Clustering of intra-molluscalSchistosoma expression patterns. RNAseq library mapping enabled us to identify 351 genes expressed by *Schistosoma* parasites in *Biomphalaria* snail tissues. Colours: blue, *S. mansoni* Brazil (SmBRE); green, *S. mansoni* Venezuela (SmVEN); and red, *S. rodhaini* (Srod). The heatmap represents the profiles of the 351 genes expressed by the different parasites at 24 h after infection. Each transcript is represented once, and each line represents one transcript. The expression level is highlighted by the different shades of blue.

FIGURE 7. In silico identification of parasite miRNAs. miRNAs were assessed using libraries obtained from naïve snails and snails infected for 24 h under the various interaction conditions (BB24, BgBRE x SmBRE; BV24, BgBRE x SmVEN; BR24, BgBRE x Srod). A) Table highlighting the precursor miRNAs that may have targets among the immune-related snail transcripts selected in the present work. They include eight precursors specifically recovered in BB24, two in BV24, and one shared across the three infected conditions. The total numbers of potential targets in each condition are indicated. B) Venn diagram showing the potential targets according to the sympatric or allopatric interactions. Shown is an example miRNA stem-loop precursor that presents the highest number of potential host targets.

SUPPLEMENTARY FIGURE 1. Clustering of all differential represented transcripts. Clustering of differentially represented transcripts. Heatmap representing the profiles of the 1,895 differentially represented immune-related transcripts in the BB, BV or BR interactions along the time course of infection (at 24 and 96 h).

1096 Each transcript is represented once, and each line represents one transcript. Colours: yellow, overrepresented

1097 transcripts; purple, underrepresented transcripts; and black, unchanged relative to levels in control naïve snails.

1098

SUPPLEMENTARY FIGURE 2. Mortality of *B. glabrata* snails after infections. The survival rates of *B.*

glabrata subjected to infection by different *S. mansoni* strains were observed over 14 weeks. Kaplan Meier

graphs were generated using xlstat, and the log-rank test ($p < 0.05$) was used to test for significant between-

group differences. Coloured curves indicate the mortality rates of naïve snails (yellow) (n = 60), snails infected

by the sympatric parasite (BB, BgBRE/SmBRE, blue) (n = 50), and snails infected by the allopatric parasite

(BV, BgBRE/SmVEN, green) (n=50). The difference in mortality between naïve and infected snails was

significant ($p < 0.05$), whereas that between the two infected conditions was not ($p = 0.243$).

1106

1107
1108

1109
1110
1111
1112 **TABLE 1.** Number of transcripts in each step of transcriptomic analysis

1113

Analysis of transcriptomic data		Analyses of Blast2GO	Analyses of annotations	
Full transcriptome	Differentially expressed	Informative annotation	Immune transcripts	Non-immune transcripts
159,711	3,865	1,017	336	681

1114
1115

1116

1117

1118

1119

SUPPLEMENTARY TABLE 1. *Biomphalaria* and *Schistosoma* compatibility between African and
South American strains

			Biomphalaria		
			glabrata		*pfeifferi*
			BRE	VEN	SEN
Schistosoma mansoni	BRE	P %	**100**	70	92
		I	**3.6**	3.5	2.9
	VEN	P %	100	**95**	100
		I	3.2	**3.8**	4
	SEN	P %	27	13	**100**
		I	1.3	1	**2.5**
Schistosoma rodhaini		P %	80	*NA*	*NA*
		I	2	*NA*	*NA*

The prevalence (P %: percentage of snail infected) and intensity (I: number of parasites per
infected host) of infection are presented for each experimental infection. The indicated values
correspond to 20 miracidia. Each pairwise combination of *Biomphalaria* (BgBRE, BgVEN,
BpSEN) and *Schistosomamansoni* (SmBRE, SmVEN, SmSEN) or *SchistosomaRodhaini*
(Srod) were tested for compatibility. *NA*: non-available data.

    Figure 1

A

B

1150 Figure 3
1151

1158   Figure 4
1159

1160
1161
1162
1163

1164   Figure 5
1165

1173   Figure 6
1174

1180 Figure 7
1181

A

Precursor	BgBRE duplicates		BB24	BV24	BR24	Targets
sma-mir-8460	0	0	1	0	0	7
sma-mir-8463	0	0	1	0	0	9
sma-mir-2a	0	0	1	0	0	20
sma-mir-2d	0	0	1	0	0	50
sma-mir-3492	0	0	1	0	0	10
sma-mir-8404	0	0	1	0	0	4
sma-mir-8414	0	0	1	0	0	14
sma-mir-8446	0	0	1	0	0	23
sma-mir-8456	0	0	0	1	0	2
sma-mir-8431	0	0	0	1	0	4
sma-mir-190	0	0	1	1	1	17
Sum of potential Bg targets	**0**	**0**	**154**	**23**	**17**	

SmBRE specific SmVEN specific Shared

B

5'-UAAUUUGAUGUUAUUAAGG-3'

5'-GUUCUUAAUGUUGUAUUAGUU-3'

sma-mir-8431
E=-29.9
4 putative targets

Lowest free energy structure
Probability >= 99%
99% > Probability >= 95%
95% > Probability >= 90%
90% > Probability >= 80%
80% > Probability >= 70%
70% > Probability >= 60%
60% > Probability >= 50%
50% > Probability

Prediction from ~/RNA structure web

SmVEN specific

Srod specific

SmBRE specific

0
0
6 17 0
0
123

Shared

sma-mir-2d-3p
E=-29.6
50 putative targets

sma-mir-190-3p
E=-43.8
17 putative targets

5'-UAUCACAGUCCUGCUUAGGUGA-3'

5'-CAGUGACCAGACAUAUCCCU-3'

1188   Supplementary Figure 1
1189

1190
1191
1192
1193

1194   Supplementary Figure 2
1195

1196
1197