ARTICLE

Sympatric and allopatric evolutionary contexts shape differential immune response in

***Biomphalaria / Schistosoma* interaction.**

Anaïs PORTET[1], Silvain PINAUD[1], Cristian CHAPARRO[1], Richard GALINIER[1],

Guillaume M. CHARRIERE[2], Jean-François ALLIENNE[1], David DUVAL[1], Benjamin

GOURBAL[1§]

[1] Univ. Perpignan Via Domitia, Interactions Hôtes Pathogènes Environnements UMR 5244,

CNRS, IFREMER, Univ. Montpellier, F-66860 Perpignan, France

[2] Interactions Hôtes-Pathogènes-Environnements (IHPE), UMR 5244, CNRS, Ifremer,

Université de Perpignan Via Domitia, Université de Montpellier, Montpellier, 34095, France.

19 [§] corresponding author:

20 Benjamin Gourbal

21 UMR5244 IHPE, Université de Perpignan

22 58 avenue Paul Alduy, 66860 Perpignan

23 benjamin.gourbal@univ-perp.fr

Abstract

Background:

Selective pressures between hosts and their parasites can results in reciprocal evolution or adaptation of specific life history traits. Local adaptation of resident hosts and parasites should lead to host-parasite systems performing better in sympatry when compared to allopatry. Between-population variations in parasite infectivity/virulence and host defense/resistance, referring as compatibility phenotype, were often the proxy used to analyse sympatric or allopatric adaptation. Nevertheless, some reported cases exist where allopatric host-parasite systems demonstrate compatibility phenotypes similar or greater than the one observed in sympatry. In these cases, the role of local adaptation is questioning. Here, we study the interaction between *Schistosoma* and its vector snail *Biomphalaria* in which such discrepancy in local versus foreign compatibility phenotype has been observed. Herein, we developed an integrative approach to investigate sympatric and allopatric interaction processes and link the underlying molecular mechanisms to the resulting phenotypes. Using comparative "omics" approaches joined to analysis of life history traits (immune cellular response, mortality, prevalence and compatibility) we try to bridge the gap of knowledge that exists for connecting local adaptation observations to molecular phenotypes in *Schistosoma/Biomphalaria* interactions.

Results:

We found that despite displaying similar prevalence phenotypes, parasite infection triggered an immune suppression in snails living in sympatry, while it activated an immune response for those living in allopatry. Dual-comparative molecular analyses revealed that parasite infection causes immune suppression in sympatry. miRNAs were used to hijack host's immune response, allowing sympatric parasites to initiate its developmental program earlier and more efficiently.

Conclusions:

We show that despite having similar prevalence phenotypes, sympatric and allopatric snail-*Schistosoma* interactions displayed a strongly different immunobiological molecular dialogue. The ability for allopatric pathogens to adapt rapidly and efficiently to new hosts could have critical consequences on disease emergence and risk of Schistosomiasis outbreaks. These observations would have important consequences in term of Schistosomiasis disease control.

Keywords: sympatry, allopatry, host-parasite compatibility, RNA-seq, immunity

Introduction

Schistosomiasis is the second most widespread human parasitic disease after malaria and affects over 200 million people worldwide (WHO, 2010). *Schistosoma mansoni* (Platyhelminthes, Lophotrochozoa) causes intestinal schistosomiasis. *Schistosoma* needs a mollusc (belonging to *Biomphalaria, Bulinus,* or *Planorbarius* genus) to act as its first intermediate host, where it undergoes part of its life cycle before infecting humans. Patently infected snails support the continuous production of thousands of cercariae, the free-living human-infective stage of the parasite, until dying. Vector snails appeared thus as a central actor of the parasite transmission and thus have deserved greater attention as potential target for Schistosomiasis control. To deserve this objective it is necessary to understand snail-parasite immunobiological interactions and to characterize the molecular mechanisms by which snails and *Schistosoma* interact.

Biomphalaria / Schistosoma compatibility is based on a complex phenotype-to-phenotype or matching-phenotype model (Basch, 1976; Galinier *et al.*, 2017; Mitta *et al.*, 2012; Mitta *et al.*, 2017; Theron *et al.*, 1997; Theron *et al.*, 2014). The success or failure of the infection of *B. glabrata* by *S. mansoni* reflects a complex interplay between the host's defense mechanisms and the parasite's infective strategies. In the past 15 years, the molecular basis of this compatibility polymorphism has been investigated at the genomic (Bonner *et al.*, 2012; Goodall *et al.*, 2006; Tennessen *et al.*, 2015), transcriptomic (Galinier *et al.*, 2017; Ittiprasert *et al.*, 2010; Lockyer *et al.*, 2012; Lockyer *et al.*, 2004; Lockyer *et al.*, 2007; Roger *et al.*, 2008a), proteomic (Bender *et al.*, 2007; Galinier *et al.*, 2013; Hahn *et al.*, 2001; Larson *et al.*, 2014; Mone *et al.*, 2010; Myers *et al.*, 2008; Pila *et al.*, 2016; Roger *et al.*, 2008b) and epigenomic levels (Bridger *et al.*, 2014; Ittiprasert, Knight, 2012; Ittiprasert *et al.*, 2009; Knight *et al.*, 2015; Nelson *et al.*, 2016; Perrin *et al.*, 2012). These studies have revealed that various molecules and pathways involved in immune recognition (snail immune receptors versus parasite antigens), immune effector/anti-effector systems, and immune regulation/activation participate in a complex interplay that governs the match or mismatch of host and parasite phenotypes (Portet *et al.*, 2017). This complex phenotype-by-phenotype interaction (compatibility polymorphism) varies between populations and individuals resulting in a "multi-parasite susceptibility" or "multi-host

infectivity" phenotypes (Theron *et al.*, 2014) that reflect between-population variations in parasite

infectivity/virulence and host defense/resistance (Osnas *et al.*, 2015; Tack *et al.*, 2012).

In *B. glabrata/Schistosoma sp.* models, numerous sympatric and allopatric combinations have been

tested; (i) different *B. glabrata* laboratory strains (or isolates) show various degrees of susceptibility to

S. mansoni infection and (ii) different strains of *S. mansoni* display different levels of infectivity

towards a particular strain of snail host (Basch, 1976; Prugnolle *et al.*, 2006; Richards, 1975; Theron

et al., 1997; Theron *et al.*, 2014). In most cases, interaction in *B.glabrata/Schistosoma* models has

been investigated by comparing, (i) sympatric/compatible and (ii) allopatric/incompatible host-parasite

associations. Indeed, the general assumption is that the parasites, which have a shorter generation

time, larger population sizes and higher reproductive outputs, are ahead in the co-evolutionary conflict

against their host and are therefore more likely to locally adapt to them (Frank, 1996; Kawecki, Ebert,

2004). This is exactly what is stated by the "host-parasite local adaptation hypothesis", in which the

sympatric parasite is supposed to perform better than the allopatric one (Ebert, 1994; Kalbe *et al.*,

2016; Kawecki, Ebert, 2004; Lively, 1989). However, it has been observed that compatibility

phenotypes between *Biomphalaria* and *Schistosoma* were the same comparing sympatric and

allopatric combinations or even greater for allopatric parasites, and thus did not fulfil the "local versus

foreign" criterion of the "host-parasite local adaptation hypothesis" (Morand *et al.*, 1996; Osnas,

Lively, 2004; Prugnolle *et al.*, 2006). Such compatibility phenotypes have been mostly investigated in

terms of parasite infectivity or pathogenicity toward snail hosts, but to our knowledge very few studies

have investigated the molecular basis of allopatric compatible interactions from the perspective of

both side of the interaction, the host and parasite (Mone *et al.*, 2011; Penczykowski *et al.*, 2015).

Hence, in order to bridge this gap, we herein study sympatric/allopatric interactions displaying similar

compatibilities using an integrative approach that links the underlying molecular mechanisms to the

resulting phenotypes, based on comparative molecular approaches on both host snails and *Schistosoma*

parasites. We set out to analyze the compatibility between South American snail strains (from Recife

Brazil and Guacara Venezuela) in combination with three different parasite isolates that displayed

nearly the same compatibility phenotype: (i) *S. mansoni* Recife Brazil, (ii) *S. mansoni* Guacara

Venezuela (both sympatric and allopatric, narrow geographic scale), and (iii) *S. rodhaini* from Burundi

Africa (allopatric, large geographic and phylogenetic scales). Thus even if compatibility phenotypes

(prevalence/intensity) are similar, the characterization of the underlying molecular mechanisms

demonstrates a very different immunobiological dialogue between *B. glabrata* vector snails and their

sympatric or allopatric *Schistosoma* parasites at the cellular and molecular scales.

To the best of our knowledge very few studies have investigated the molecular basis of local

adaptation in host/parasite interactions (Bankers *et al.*, 2018; Robertson *et al.*, 2017). Here we used a

powerful global approach to identify the differentially regulated transcripts in snails of a single *B.*

glabrata strain under infection by three sympatric or allopatric compatible strains or species of the

parasite, *Schistosoma sp.*

Moreover, our results may be of critical importance in terms of Schistosomiasis control.

Understanding the mechanisms whereby allopatric parasites efficiently infect host snails, would be

mandatory to develop new tools to predict or quantify risks of Schistosomiasis disease outbreaks

(Kincaid-Smith *et al.*, 2017).

Materials and Methods

Ethical statements

Our laboratory holds permit # A66040 for experiments on animals, which was obtained from the

French Ministry of Agriculture and Fisheries and the French Ministry of National Education,

Research, and Technology. The housing, breeding, and care of the utilized animals followed the

ethical requirements of our country. The experimenter possesses an official certificate for animal

experimentation from both of the above-listed French ministries (Decree # 87–848, October 19, 1987).

The various protocols used in this study have been approved by the French veterinary agency of the

DRAAF Languedoc-Roussillon (Direction Régionale de l'Alimentation, de l'Agriculture et de la

Forêt), Montpellier, France (authorization # 007083).

Biological materials

Two snail strains of *Biomphalaria glabrata* were used in this study: the albino Brazilian strain,

(BgBRE) and the Venezuelan strain, (BgVEN). They were exposed to infection by a Brazilian

(SmBRE) or Venezuelan (SmVEN) strains of *Schistosoma mansoni*, or by another *Schistosoma*

species, *S. rodhaini* (Srod). All host and parasite strains of each different geographical origin were

recovered in their native locality and maintain in the laboratory always on their sympatric snail hosts

to maintain reciprocal selective pressures and reciprocal sympatric strain adaptation. We housed snails

in tanks filled with pond water at 25°C with a 12:12 hour light:dark cycle and supplied ad libitum with

fresh lettuce. The Brazilian strain originates from the locality of Recife (east Brazil), the Venezuelan

strains of snail and parasite were recovered from the locality of Guacara (north Venezuela) and the

African species *Schistosoma rodhaini* originates from Burundi and was obtained from the London

British Museum. These *Schistosoma* isolates/species have been selected because they exhibited similar

infectivity toward BgBRE or BgVEN strains (see prevalence and intensity in supplementary table S1).

Prevalence of SmBRE and SmVEN for the African vector snail *Biomphalaria pfeifferi* from Senegal

(BpSEN), and prevalence of the corresponding parasite SmSEN on South American snails were also

tested (supplementary table S1).

RNAseq experimental protocol

In order to investigate the molecular response of snails against sympatric and allopatric parasites, a

global comparative transcriptomic approach was conducted. One hundred and twenty BgBRE snails

were infected with SmBRE, SmVEN or Srod. Each snail was individually exposed for 12 h to 10

miracidia in 5mL of pond water. For each experimental infection, 20 snails were recovered at 24 and

96 h after infection. The samples were named as follows: BB24, BB96 for BgBRE infected with

SmBRE; BV24, BV96 for BgBRE infected with SmVEN; and BR24, BR96 for BgBRE infected with

Srod. Two pools of 30 uninfected BgBRE snails (Bre1 and Bre2) were used as controls.

- Whole-snail RNA extraction and sequencing

Total RNA was extracted using TRIZOL® (Sigma Life Science, USA) according to the

manufacturer's instructions. Sequencing was performed using in paired-end 72-bp read lengths on

Illumina Genome Analyzer II (MGX-Montpellier GenomiX, Montpellier, France).

- De novo transcriptome assembly

De novo transcriptome assembly, using all time points, was performed using an in-house pipeline

created with the Velvet (1.2.02), Oases (v0.2.04), and CD-HIT-EST (v4.5.4) programs. The assembly

of the consensus reference transcriptome was optimized using various parameters, including k-mer

length, insert length and expected coverage, as previously described (Dheilly *et al.*, 2015; Pinaud *et*

al., 2016). A *de novo* transcriptome was created and contained 159,711 transcripts.

- Differential expression analysis

High-quality reads (Phred score >29) were aligned to the *de novo* transcriptome using Bowtie2

(v2.0.2), which was run locally on a Galaxy server. To compare the host responses during the

sympatric or allopatric interactions, we used the DESeq2 (v2.12) software to analyse the differential

transcript representation between *BgBRE* control strains (uninfected *Bg*BRE1 and *Bg*BRE2) to the

sympatric and allopatric conditions (p-value < 0.1) (Pinaud *et al.*, 2016). A Venn diagram was

generated using the Venny 2.1 software to highlight which differentially expressed transcripts were

specific or common to the different interactions. A heatmap was then created to analyze transcript

expression patterns, using log2 Fold Change with Hierarchical Ascending Clustering (HAC) and

Pearson correlation (uncentered) as applied by the Cluster (v3.0) and Java TreeView (v1.1.6r4)

softwares packages. The differentially represented transcripts were functionally classified using a

BlastX analysis with the cut-off set to e-value < $1e^{-3}$ (NCBI dataset; thanks to the Roscoff Data center

Cluster, UPMC) and gene ontology was assigned using an automatic annotation, Blast2GO (v3.0.8)

(supplementary table S2). Based on automatic annotation and functional domains and according to the

literature we identified the potential immune transcripts involved in snail immunity.

Schistosoma intra-molluscal stage transcriptome analysis: Dual RNA-seq

Host response against pathogens constitutes solely one side of the molecular dialogue occurring in

host/parasite interactions. To analyse all the complexity of such dialogue, the parasite response into

host tissue needs also to be investigated. Herein a dual RNA-seq approach was thus conducted to gain

in a broader understanding of sympatric and allopatric host/parasite interactions.

- Schistosome read selection

The *Biomphalaria* (v1) and *Schistosoma* (v5.2) genomes have been concatenated

(https://www.vectorbase.org/organisms/biomphalaria-glabrata;

http://www.sanger.ac.uk/resources/downloads/helminths/schistosoma-mansoni.html). High quality

reads (Phred score >29) were mapped against these concatenated genomes using Bowtie2 (v2.0.2),

which was run locally on the Galaxy project server. The reads that mapped only once and exclusively

to the *Schistosoma* genome were collected as corresponding to *Schistosoma* reads; reads that mapped

to the *Biomphalaria* genome or more than once to either genomes were removed from the analysis.

- *Gene analysis*

The above-selected *Schistosoma* reads were mapped against the concatenate genome to identify intra-

molluscal stage-specific *Schistosoma* genes. In order to select the relevant genes, the reads mapped in

all experimental conditions were summed. Solely genes with a minimal sum of 10 reads were kept for

the analysis. A heatmap was generated to analyse *Schistosoma* gene expression patterns using

Hierarchical Ascending Clustering (HAC) with Pearson correlation (uncentered) as applied by the

Cluster (v3.0) and Java TreeView (v1.1.6r4) software packages. Functional annotation of the genes

was assigned using BlastX with the cut-off set to e-value $< 1e^{-3}$ (NCBI dataset, local cluster) and gene

ontology was performed using Blast2GO (v4.0.7) (supplementary table S3).

Innate immune cellular response analysis: microscopy and flow cytometry

Hemocytes appeared as the main cells supporting *Biomphalaria* snail immune response. Thus, to go

further in the description of snail response against parasites, quantitative and qualitative changes in

hemocyte populations were investigated. For this purpose, BgBRE and BgVEN snails were used.

Snails were infected as described above, using either SmBRE or SmVEN parasites. For each

experimental infection, snails were recovered at 24 and 96 h after infection, and designated as follows:

BB24 and BB96 for BgBRE infected with SmBRE; BV24 and BV96 for BgBRE infected with

SmVEN; VV24 and VV96 for BgVEN infected with SmVEN; and VB24 and VB96 for BgVEN

infected with SmBRE. Snails of each strain, BgBRE and BgVEN, were recovered and used as

controls.

- Hemocyte proliferation analysis: microscopy

Microscopic inspection of hemocyte proliferation was conducted using 12 infected BgBRE (6

BgBRExSmBRE and 6 BgBRExSmVEN) and 3 uninfected BgBRE snails. The hemocytes of 3 snails

(biological replicates) were counted for each condition at 24h and 96h after infection. The

proliferation of circulating hemocytes was studied by using a Click-iT EdU Alexa Fluor 488 Flow

Imaging Kit (Molecular Probes). At each time point, circulating hemocytes were recovered by direct

puncture after foot retraction and 1mM of EdU solution was added to the hemolymph. Three hours

later, the amount of EdU incorporated by the circulating hemocytes was visualized *in-vitro* after

fixation of the cells and performing a covalent coupling of Alexa Fluor 488 to the EdU residues trough

a click chemistry reaction flowing manufacturer indications, then nuclei of hemocytes were

counterstained with DAPI (Biotum) staining, and the sample was analysed on a confocal microscope

using a Zeiss LSM 700, with 4 lasers (405, 488, 555 and 633 nm). Positive cells were counted and

between-sample differences in the percentage of proliferation were tested using a Fisher exact test,

with significance accepted at p-value<0.05.

- Hemocyte proliferation and population profiles analysis: flow cytometry

Qualitative changes in hemocyte populations following infection by sympatric or allopatric parasites

were studied using a flow cytometry approach. For this 72 infected BgBRE or BgVEN (36 infected by

SmBRE and 36 infected by SmVEN) and 18 uninfected BgBRE or BgVEN snails were used. Six

biological replicates (pools of 3 snails per replicate) were performed for each condition. Flow

cytometry was used to profile and assess the proliferation of circulating hemocytes using Click-

iTEdUAlexa Fluor 647 labelling (Molecular Probes). At each time point, 1mM of EdU solution was

injected into pericardial cavity of each snail. Three hours later six replicates of 3 snails were collected,

and the hemolymph was extracted from the head-foot according to standard procedures (Sminia,

Barendsen, 1980). The hemolymph was pooled from the three snails, and 100 µl were subjected to

analysis with the above-listed kit, according to the manufacturer's instructions. The percentage of

proliferative cells was calculated by flow cytometry.

The hemocytes were profiled along the course of infection using Side Scatter Chanel (SSC) to

estimate cell granularity and Forward Scatter Chanel (FSC) to estimate cell size. The cell repartition

along these two parameters enables to identify cell sub-populations. The flow cytometry was

performed using a FACS Canto from BD Biosciences (RIO Imaging Platform, Montpellier, France).

For each sample, 10,000 events were counted. The results were analyzed with the FlowJo V 10.0.8

software. Between-group differences in the percent of proliferation were tested using the Mann-

Whitney U-test, with significance accepted at p-value<0.05.

Histological procedures

A histological approach was conducted in order to investigate differences in miracidia to sporocyst

development, while comparing sympatric and allopatric parasite growth, development and maturation

into snail tissues. BgBRE snails (n=8 per group) were infected as described above with either SmBRE

(sympatric) or SmVEN (allopatric) parasites. At 24 and 96 h after infection, two snails were fixed in

Halmi's fixative (4.5% mercuric chloride, 0.5% sodium chloride, 2% trichloroacetic acid, 20% formol,

4% acetic acid and 10% picric acid-saturated aqueous solution). Embedding in paraffin and transverse

histological sections (3-μm) were performed using the RHEM platform (Montpellier, France)

facilities. The slides were stained using Heidenhain's azan trichromatic staining solution as follows:

(i) serial re-hydration was performed in toluene followed by 95%, 70%, and 30% ethanol and then

distilled water; (ii) coloration was performed using azocarmine G (70% ethanol, 1% aniline, 1% acetic

alcohol, distilled water, 5% phosphotungstic acid, distilled water, Heidenhain's azan) and (iii) serial

dehydration was performed using 95% ethanol, absolute ethanol, and toluene. The preparations were

then mounted with Entellan (Sigma Life Science, St. Louis Missouri, USA) and subjected to

microscopic examination. The parasite sizes were measured with the ImageJ software (v2.0.0), and

between-group size differences were tested using the Mann-Whitney U-test with statistical

significance accepted at a p-value<0.05 based on the mean of the three largest serial cuts of each

sporocyst in sympatric or allopatric conditions.

In-silico *characterization of* Schistosoma *miRNAs*

Parasites may communicate or interfere with their host using different strategies based mainly on

excreted/secreted products released into hemolymph. In this context, miRNAs appeared as the most

relevant mean of communication that can be used by parasites. To test for such hypothesis *S. mansoni*

miRNAs were analyzed *in-silico* by comparing the relevant miRNA database (miRBase) to our

RNAseq libraries generated at the 24h following sympatric or allopatric infections. S. *mansoni*

precursor sequences were downloaded from miRBase (http://www.mirbase.org, 03/09/2017), and

high-quality reads from naive (BgBRE) and 24 h post-infection samples (BB24, BV24, BR24) were

aligned against a *S. mansoni* sub-database of miRBase, as previously described (Zhao *et al.*, 2016).

The identified precursors were confirmed by alignment of high-scoring reads onto precursor and

mature miRNAs from miRBase. Solely reads with 100% identity were retained for analysis. The

localization of each read against miRNA sequence allowed us to identify either the precursor or just

the mature miRNA. Precursors found under both naive and infected conditions were excluded to retain

exclusively the miRNAs present in samples from infected snails and avoid cross-species

contamination because of the potential conserved features of miRNAs from *B. glabrata* and *S.*

mansoni.

Putative miRNA targets were predicted from among the differentially represented immune-related

transcripts (figure 1) using Miranda tools (using parameters: Miranda input_miRinput_Transcriptome

-out results.txt -quiet -sc 140 -en -15) (Enright *et al.*, 2003). Because mature miRNAs may exist in two

forms depending on which strand (5'-3') of the precursor stem-loop is maturated the predicted

interactions could involve the 5' and/or 3' forms, as noted. The results were extracted using the awk

tool, listed in supplementary files (supplementary table S4), and used to generate a Venn diagram. To

confirm the ability of a selected pre-miRNA to produce the stem-loop necessary to produce the mature

form, the secondary structures of precursor were predicted using RNA structure Web tool

(http://rna.urmc.rochester.edu/RNAstructureWeb, 03/09/2017) using default parameters.

Snail survival analysis

Allopatric or sympatric parasites could have different levels of virulence or impacts on their host that

could impair snail survival. To test for such discrepancy we investigated the mortality rates of infected

snails over the course of sympatric or allopatric infections. One hundred and sixty BgBRE snails were

infected as described above with SmBRE or SmVEN strains (n=50), and 60 non-infected BgBRE

snails were retained as controls. The numbers of dead snails were compiled weekly for 14 weeks. A

Kaplan-Meier estimator was used to estimate the survival function from lifetime data. Survival curves

were generated using the xlstats Mac software and the log-rank test was applied with significance

accepted at p<0.05.

Results

A whole-snail transcriptomic approach for investigating the molecular basis of the innate immune

response in the sympatric and allopatric contexts

The global transcript representation was analyzed using the previously described RNAseq pipeline

developed in our laboratory (Dheilly *et al.*, 2015; Galinier *et al.*, 2017; Pinaud *et al.*, 2016). The

*Bg*BRE snail host had the same prevalence and intensity phenotypes when exposed to *Sm*BRE,

*Sm*VEN or *S*rod parasites (see Table S1). Of the 159,711 transcripts of the *Bg*BRE transcriptome,

3,865 (2.4%) were differentially represented in all sympatric and allopatric conditions compared to

naive snails (table 1, supplementary fig. S1). We performed automatic Blast2GO annotation, discarded

the non-annotated transcripts, and retained 1,017 annotated transcripts (26.3% of the differentially

represented transcripts). Finally, as we were interested in the innate immune response of the snail to

the sympatric and allopatric parasites, we focused on transcripts known to have immune-related

functions. Of the annotated differentially expressed transcripts, 336 were identified as being immune-

related; this subset corresponded to 8.7% of all differentially represented transcripts. The heatmap

generated for all differentially represented transcripts is presented in supplementary fig. S1.

Venn diagrams were used to highlight the relationships of the transcripts that were differentially

represented in the *B. glabrata* and *Schistosoma sp.* interactions. Overall, 189, 180, and 164 transcripts

were differentially represented in the BB (BgBRE/SmBRE, sympatric), BV (BgBRE/SmVEN,

allopatric), and BR (BgBRE/Srod, allopatric) interactions, respectively (fig. 1A). Of the selected

transcripts, 40 were shared by the BB, BV and BR interactions (fig. 1A), and displayed globally the

same differentially expressed pattern of expression (fig. 1B, cluster 1). These transcripts may

correspond to the snail's basal immune response against *Schistosoma* parasites.

Of the 140 transcripts exclusive to the sympatric (BB) interaction, which represented 74.1% of the

immune transcripts found to be differentially represented in this interaction (fig. 1A), 100% were

under-represented at 24 h post-infection, and 74.6% of these transcripts were observed exclusively at this time point (fig. 1B, cluster 5).

Very similar transcript expression patterns were observed for the two allopatric interactions (BV and BR). Clusters 6 and 7, comprising only 28 and 11 transcripts, were exclusive to BV and BR, respectively. In BV interaction (fig. 1B, cluster 6), at 96 h after infection 96.5% of the transcripts were expressed: 22% over-represented. In BR interaction (fig. 1B, cluster 7), 100% of the transcripts were expressed at 96 h after infection: 82% over-represented. Of the 108 transcripts shared by BV and BR (cluster 3, representing 60% and 65.9 % of the transcripts, respectively) (fig. 1A; fig. 1B, cluster 3), 98.1% were expressed at 96 h post-infection: 28.2% over-represented (fig. 1B, cluster 3).

Altogether, these results show that the host immune-related transcriptomic response differed widely between the sympatric and allopatric interactions. All of the differentially represented transcripts were under-expressed at 24 h after infection in sympatry, while the differentially represented transcripts in alloparty were mostly over-expressed at 96 h after infection. Moreover, the sympatric interaction was associated with a strong immunosuppression (fig. 1B, cluster 5), whereas allopatry yielded an activation of the immune response (fig. 1B, cluster 3).

We decided to explore the function of differentially regulated transcripts between the sympatric and allopatric interactions in order to investigate if these differentially regulated transcripts belong to different functional categories. For that, we performedan automatic annotation with Blast2GO (supplementary table S2). Based on automatic annotation and according to the literature we distributed the relevant differentially expressed immune transcripts into three groups: (i) immune recognition molecules, (ii) immune effectors, and (iii) immune signalling molecules (fig. 1C). Then these three groups were subdivided into functional categories (fig. 2). In the BB sympatric interaction, immune recognition molecules, immune effectors and immune signalling pathways correspond to 13.1%, 18.3% and 4.2%, respectively (sums of all relevant transcripts at all infection time points). Among the transcripts distributed into these groups, 89.0% were under-represented. In the BV allopatric interaction, the three groups corresponded respectively to 10.8%, 12.9% and 8.6% and among them 65.9% were under-represented. Finally, in the BR allopatric interaction, they correspond to 12%, 12.5% and 7.8% and 62.5% of them were under-represented. When we compared the percentage of

each immunological group in the sympatric and allopatric interactions, no specific functional subset

was particularly repressed in the BB sympatric interaction (fig. 1C; fig. 2). In fact, most of the time the

same immune functions were affected in sympatric and allopatric infections but different immune

transcripts (grey and black diamond in fig. 2) showed differential regulation following infections (fig.

2).

The differentially regulated transcripts belonging to the three immunological groups (fig. 2) were

largely involved in immune cellular responses, cell adhesion, extra cellular matrix component, cell

migration, cell differentiation and cell proliferation. All of these functions were under-represented in

sympatric interaction (76%), whereas many were over-represented in allopatric ones (39%) at 24h

after infections (fig. 2).

Immune cellular responses in the sympatric and allopatric contexts

As the transcriptomic approach indicated that multiple immunobiological functions seem to be altered

in the host response during the interaction with the parasites, we next investigated whether differences

in immune responses in sympatric versus allopatric infections could be seen at the cellular level.

Hemocytes, the snail immune cells, participate directly in the immune response against the parasites,

and immune cell activation under an immunological challenge can translate into cell proliferation

and/or cell morphology modifications.

Thus to investigate such immune cell activation, cell proliferation was quantified using EdU nuclear

labelling (fig. 3). EdU is a nucleoside analogue of thymine incorporated into DNA during DNA

synthesis. This incorporation reflects the mitotic activity of hemocytes. As the two tested allopatric

interactions appeared extremely similar at the transcriptomic level, we compared only the BV

(allopatric) interaction and BB (sympatric) interaction for further analyses. Circulating hemocytes

recovered in snail hemolymph were EdU-labelled *in-vitro* to identify mitotic circulating hemocytes

(fig. 3A). Edu-positive hemocytes were quantified using confocal microscopy and showed that

hemocyte proliferation was induced by 3 fold following *Schistosoma* infection in BV compared to BB

(fig. 3B). Twenty-four hours after infection, proliferative circulating cells increase significantly in BV,

with 5.2% of proliferative cells compared to BB with 2.6% (Fisher exact test two-tailed p = 7.6 e10^{-6})

(fig. 3B). Moreover, this result demonstrates for the first time that "circulating" hemocytes are able to proliferate following *Schistosoma* infections.

Hemocyte proliferation was assessed using flow cytometry after *in-vivo* EdU-labelling (fig. 4A). At 24h after infection, the rate of proliferating cells was increased significantly (Mann Whitney U test: U=36; z = -2.8; p = 0.0022) by more than two fold between the allopatric and sympatric interactions, 4.2% compared to 1.8% respectively. At 96 h after infection, there were fewer proliferating cells: the percentage of proliferating hemocytes was similar in sympatric BB interaction compared to the control (1%, Mann Whitney U test: U=17; z=-0.27; p = 0.3936), while remaining somewhat higher in allopatric BV interaction (2.3%, Mann Whitney U test: U=0; z=2.65; p = 0.009).

To confirm that our observations were related to the type of interaction rather than to a strain effect, we performed the same experiments using another *Biomphalaria glabrata* strain, BgVEN as the host and SmVEN and SmBRE as the sympatric and the allopatric parasite, respectively (fig. 4B). The rate of proliferating hemocytes increased significantly (Mann Whitney U test: U=36; z=-2.8; p = 0.0022) by more than 3 fold between the allopatric (VB: BgVEN/SmBRE) and the sympatric (VV: BgVEN/SmVEN) interactions at 24 h after infection (6.8% compared to 2.0% respectively). At 96 h after infection, both rates were decreased, albeit more for VV (0.1%, Mann Whitney U test: U=2; z=2.48; p = 0.013) than for VB (2.7%, Mann Whitney U test: U=36; z=2.8; p = 0.0022). These results similar to the one observed for BgBRE interactions confirmed that the effects observed at the cellular level were related to the type of interactions (sympatric versus allopatric) rather than to the host or parasite strains used.

In addition to the mitotic activity, hemocyte morphology was estimated using flow cytometry by measuring their size and granularity. We analyzed the circulating hemocyte populations in non-infected versus infected snails in two *B. glabrata* snail strains (BgBRE and BgVEN) in sympatric and allopatric interactions with parasites (fig. 4C and 4D). Morphology and heterogeneity of circulating hemocytes varied in a similar manner between BgBRE and BgVEN snails subjected to infection (fig. 4C and 4D) and we observed infection-related differences in cell morphology. In non-infected snails,

the content of circulating hemocytes was very heterogeneous, but appeared to represent a single

population that exhibited a continuous gradient of size and granularity (fig. 4C and 4D), which is

typical of *B. glabrata* hemocytes (Baeza Garcia *et al.*, 2010). However, hemocyte population

heterogeneity changed quickly after infection. In allopatric interactions, 24 h after infection (fig. 4C,

BV24, and 4D, VB24) two populations could be distinguished: a population P1 (corresponding to that

seen in non-infected snails) and a population P2 (a new population). P2 cells exhibited increased

granularity, retained a high degree of size variability, and showed a mitotic activity, as indicated by

EdU labeling (fig. 4C and 4D, red dots). This profile was transitory, with the P2 population

disappearing by 96 h after infection (fig. 4C, BV96, and 4D, VB96). Altogether, these results show

that, upon infection, the snail circulating immune cells exhibit a particular population dynamics with

transient increase of the mitotic activity associated with morphology modifications that are likely to be

involved in the host defence response against the parasite during the interaction. Moreover, this

response of the immune cells appears to be exacerbated in the allopatric interaction when compared to

the sympatric interaction.

Schistosoma growth and development in Biomphalaria tissues

<u>Parasite development</u>

To investigate the development of *S. mansoni* in *B. glabrata* tissues, we examined the fate of

sporocyst in sympatric and allopatric compatible interactions using a histological approach. For both

interactions, miracidia were able to penetrate, transform into primary sporocysts (SpI), and develop.

At 24 h after infection, we observed a significant difference (Mann Whitney U test: U=40; z=4.33; p =

1.42 e10^{-6}) in the size of sporocyst from sympatric parasites (11,838 μm^2 average size) versus

allopatric parasites (7,402 μm^2 average size) (fig. 5). This difference was no longer seen 96 h after

infection (41,413 μm^2 and 36,920 μm^2, respectively, Mann Whitney U test: U=280; z=-1.31; p =

0.1917) (fig. 5). These results show that during the early events following infection, the parasites

develop more slowly in the allopatric compared to the sympatric interaction; thereafter, however, the

allopatric parasites catch up quickly, such that no difference is seen at 96 h post-infection (fig. 5).

<u>Parasite transcript expression analysis</u>

In order to investigate the biological responses of the parasites within their host, dual RNAseq

analyses allowed us to identify regulated transcripts from *Schistosoma* parasites during their intra-

molluscal development. The parasite RNAseq data at 24 h after infection, revealed five clusters of

differentially expressed genes from the sympatric (SmBRE) and the allopatric (SmVEN, Srod)

parasite responses (fig. 6). Cluster 1 corresponds to genes highly expressed and cluster 5 weakly

expressed for all parasite strains. Cluster 2 represents genes over-expressed in SmBRE versus SmVEN

and Srod. Cluster 3 contained genes over-expressed in SmBRE and SmVEN versus Srod and cluster 4

SmBRE and Srod versus SmVEN. In all clusters, the gene expression levels in SmBRE sympatric

parasite are always greater than in the other allopatric parasites. Blast2GO annotation helps in

characterizing 70% of the 351 genes shown in the five clusters (supplementary table S3). According to

the global Gene Ontology (GO): 70% of the annotated genes were involved in general metabolism and

growth, translation processes, regulation of cellular processes and RNA biosynthesis; 25% were

involved in molecular transport or cell organization; and 5% were involved in organism defence or

response to stimuli. In all these clusters, we identified 6 parasite gene products been involved in

parasite modulation or suppression of snail immunity. These molecules correspond to heat shock

proteins (fig. 6, clusters 1 and 2) (Knight *et al.*, 2015); glutathione-S-transferase, NADH

dehydrogenase subunit, and calreticulin (fig. 6, cluster 2) (Guillou *et al.*, 2007; Mone *et al.*, 2010;

Raghavan *et al.*, 2013); Alpha-2-macroglobulin (fig. 6, cluster 4) (Ryu, Park, 2009); von willebrand

factor type EGF with pentraxin domain (fig. 6, cluster 5) (Gilges *et al.*, 2000) (see supplementary table

S3). These particular genes could be involved in the strategy used by the parasite to ensure

immunosuppression and favour its development and growth within host tissues.

In allopatric interactions, an activation of the immune response is observed. Interestingly, allopatric

parasites did not over express any transcripts that could have immunosuppression function or impair

the activation of the immune response (fig. 6 and supplementary table S3). Furthermore, a variant of a

glycerol-3-phosphate acyl-transferase (Schisto_mansoni.Chr_3.5623) is highly over expressed in

SmVEN and Srod compared to SmBRE (cluster 1 supplementary table S3). This molecule is known to

participate in the biosynthesis of phosphatidic acid, itself involved in macrophage activation and

regulation of inflammatory signalling (Bohdanowicz *et al.*, 2013; Grkovich, Dennis, 2009). This

would thus indirectly enhance the cellular immune response of *Biomphalaria glabarata* against those

allopatric parasites.

<u>Parasite microRNAs analysis</u>

The microRNAs (miRNAs) are known as non-coding small RNA (<24nt) able to regulate gene

expressions. As we identified deep modifications of transcriptional or post-transcriptional activities

between sympatric and allopatric interactions, we investigated, *in-silico,* the potential presence of

Schistosoma mansoni miRNAs in our parasite RNAseq data. At 24 h post-infections, we identified 54

miRNA precursors from miRBase with high quality alignment scores against the different RNAseq

read libraries (naïve BgBRE, BB24, BV24, BR24). To avoid cross-species misidentifications, we

selected precursors that were exclusively identified in infected and never identified in uninfected

snails (naive BgBRE). Eleven miRNA precursors corresponding to *Schistosoma mansoni* were

identified (fig. 7A). Eight of the parasite miRNA precursors were specific to the Brazil-infected

libraries (BB24); two were specific of the Venezuela-infected libraries (BV24); and one was shared

across the three infected conditions (BB24, BV24 and BR24). Although we identify 49 miRNA

precursor sequences specific to *S. mansoni* (fig. 7B), we decided to select only miRNAs covered by

100% nucleotide similarity that allowed to predict 11 miRNAs in mature (eg. sma-mir-2d-3p, sma-

mir-190-3p) or precursor (sma-mir-8431)forms. Then, in order to identify candidate sequences that

could represent putative miRNA targets, we used the Miranda tool (supplementary table S4). Only

RNA-RNA interactions that showed good scores for pairing (>140) and enthalpy (<15 Kcal) were

considered. The number of targets pertaining to the differentially expressed immune-related transcripts

identified in Fig. 1 that were found for the identified miRNAs ranged from 2 targets for sma-mir-

8456,to 50 targets for sma-mir-2d of the differentially expressed immune-related transcripts identified

in Fig. 1.

The miRNAs identified under the sympatric condition (SmBRE) were predicted to target 43.5%of the

differentially represented immune-related transcripts identified in the RNAseq data (fig. 1B, fig. 7)

whereas 6.8% and 5.1% were targeted in allopatric conditions, SmVEN and Srod, respectively (figure 1B, fig. 7).To explain the similar prevalence between sympatric and allopatric infections we focus our attention on miRNAs that were shared between sympatric and allopatric interactions and we identified one miRNA:sma-miR-190-3p (fig. 7 and supp. Table 4). This miRNA was predicted to bind 17 different targets among which, we identified different variants of the Fibrinogen Related Protein (FREP) family and a cytotoxic/cytolytic humoral factor the biomphalysin. To go further, we look at the expression of those candidates following infection. If FREP transcripts were down regulated in sympatric interaction, it is not always the case in allopatry. However, interestingly all biomphalysin transcripts were under-represented in sympatric and allopatric interactions. Altogether, these data suggest that the parasites might hijack the host immune response using dedicated miRNAs.

Survival of snail following infection

To examine the potential impact of allopatric or sympatric parasites on snail survival, we investigated the mortality rates of infected snails over 4 months. The survival rate was significantly higher for non-infected snails compared to infected snails (sympatric interaction Kaplan-Meier Log Rank test p = 1.39 e10^{-5} and allopatric interaction p = 0.0005). However, there was no significant difference in the mortality rates of snails subjected to sympatric versus allopatric interactions: at the end of the experiment, the survival rates were 72% and 65% for the allopatric and sympatric interactions, respectively (Kaplan-Meier Log Rank test p = 0.243) (supplementary fig. S2).

Discussion

In the natural environment, it is assumed that the parasitic genes responsible for infectivity will evolve alongside the host defence genes, resulting in adaptation of the interactions between local host and parasite populations (Dybdahl, Storfer, 2003; Osnas, Lively, 2005). In this context, local/sympatric parasites were expected to display greater infectiveness, reproductive success, and virulence in host populations compared to foreign/allopatric parasites (Ebert, 1994; Ebert, 1998; Lively, 1989; Minchella, 1985). However, in some cases this rule may be contradicted, as certain allopatric parasite-host interactions have been reported to be significantly more compatibles than the corresponding

sympatric combinations (Greischar, Koskella, 2007; Morand *et al.*, 1996), it appears that certain

Biomphalaria/Schistosoma interactions *(Adema et al., 1997; Zhang et al., 2007)* do not fulfil the

"host-parasite local adaptation hypothesis"(Ebert, 1994; Ebert, 1998; Lively, 1989; Minchella, 1985),

in which the sympatric parasite is expected to perform better than the allopatric one.

Using field data, Morand et al. (1996) and Prugnolle et al. (2006) showed that although sympatric

parasite-host combinations of schistosomes and snails do tend to be more compatible, exceptions exist

wherein particular allopatric combinations are equally or significantly more compatibles. Similar

results were obtained when comparing the interactions of Brazilian and Guadeloupean snails versus

Schistosoma infections(Mone *et al.*, 2011). The authors found that allopatric Guadeloupean parasites

were not able to infect Brazilian snails; but Brazilian parasites were able to infect the allopatric

Guadeloupean snails. Furthermore, this work demonstrated the presence of local adaptation between

reactive oxygen species (ROS) and ROS scavengers in this system (Mone *et al.*, 2011). Based on these

observations, we propose that it would be important to develop integrative analysis to depict and

understand the precise molecular crosstalk (immuno-biological interactions) occurring in such highly

compatible sympatric and allopatric systems. Thus, dual-comparative approaches were used herein to

simultaneously analyze the responses of *Biomphalaria* snails and *Schistosoma* parasites into sympatric

or allopatric interactions displaying similar compatibilities.

The present RNAseq analysis demonstrated that in sympatric interaction (BB) a huge

immunosuppression occurs. Twenty-four hours after the infection, the three immunological processes:

(i) immune recognition, (ii) effector and (iii) signaling pathways (fig. 1 and 2) were down regulated.

Conversely, in allopatric interactions (BV and BR), host immune response was activated (fig. 1 and 2).

Differentially regulated transcripts mostly belong to immune cellular activation, migration,

proliferation, or differentiation (fig. 2). An EdU labelling was used to detect proliferation and to

confirm that more hemocyte proliferation was observed in allopatry compared to sympatry (fig. 3, 4A

and 4B). Using flow cytometry we demonstrated that a new hemocyte subpopulation (named P2) was

observed exclusively following allopatric infection at 24h (fig. 4C and 4D). P2 was EdU-positive and

characterized by an increased in granularity, indicating that the new P2 cell subtype could proliferate.

In Biomphalaria snails, we know 3 main hemocyte morphotypes, the Blast-like cells, the type I

hyalinocytes and the granulocytes (Cavalcanti *et al.*, 2012). However, in absence of specific hemocyte

markers, it is impossible to analyze which hemocyte morphotype are proliferating in our flow

cytometry results (fig. 4C, D). The P2 subpopulation would thus originates from either a

morphological change in an existing subset (correlating potentially with a decline in the P1

population), or represents cells that are migrating from tissues or hematopoietic organ to reach the

hemolymph. Indeed P2 population reflects newly proliferated cells that present higher EdU positive

cells than the P1 population (Fig. 4C, D). Further investigations will be necessary to conclude on the

origin of P2 population. However, hemocyte proliferation decreased more rapidly in sympatric rather

than in allopatric interactions (fig. 3 and 4). By comparing different sympatric and allopatric

interactions, we demonstrate that the cellular or molecular phenotype observed refers to reciprocal

evolution or adaptation rather to a simple host or parasite strain effect.

The strong immunosuppression observed in sympatric interaction compared to the immune cell

activation in allopatry would certainly results in differences in the capabilities of sympatric and

allopatric parasites to infect and grow in such hosts. This hypothesis was confirmed by our

histological analysis of sporocyst development in snail tissues. We observed a significant difference in

sporocyst size 24h after infection (fig. 5), when sympatric sporocysts were one-third bigger than

allopatric sporocysts. At 96h after infection, in contrast, there was no size difference between

sympatric and allopatric parasites (fig. 5). This indicates that the allopatric parasites seem to overcome

this early delay in development. However, the consequences of this delay for duration of cercariae

development, number of cercariae produced, cercariae infectivity and pathogenicity in vertebrate host,

will deserve further investigation to conclude about a potential fitness cost between sympatric and

allopatric parasites.

To go further and understand how parasites immunosuppress the host or circumvent the host immune

system, we used a dual-RNAseq approach to investigate transcripts expression of the sympatric and

allopatric parasite intra-molluscal stages (fig. 6). As the histological differences were solely observed

at 24h after infection, we used dual-RNAseq to investigate the parasite expression patterns at the same

time point of infection. Most of the parasite transcripts belonged to the processes of nucleotide

metabolism, transcription, translation and cell differentiation, development, and growth. We also identified some transcripts with GO terms or functions related to immuno-modulation or immuno-suppression (fig. 6 and supplementary table S4). Nearly all of the identified transcripts were over-represented in the sympatric interaction compared to the allopatric ones. Our results therefore suggest that the installation, development and growth of the parasite occurred much more rapidly in the sympatric combination, as sympatric parasites seemed to interfere more efficiently with the host immune system. However, RNAseq data did not give any clear information about how allopatric parasites succeed in circumventing the host immune system. We thus next examined the generated dual-RNAseq libraries in an effort to identify whether sympatric and/or allopatric schistosomes could hijack the host immune system using microRNAs. To achieve this goal, we confronted the dual-RNAseq data to the *Schistosoma mansoni* subset of miRBase to identify the presence of microRNAs (miRNAs) in our datasets. miRNAs are known to regulate numerous biological processes, including key immune response genes (Li *et al.*, 2017; Zawada *et al.*, 2017). Recent work has demonstrated that circulating small non-coding RNAs from parasites have hijack roles against host metabolism, specially in the interaction of schistosomes with their vertebrate hosts (Meningher *et al.*, 2017; Nowacki *et al.*, 2015; Walzer, Chi, 2017). Such non-coding RNAs could act as exogenous miRNAs to interfere with or circumvent the host immune system. At 24h after infection, several differentially expressed parasite miRNAs were identified. We investigated the potential role played by these miRNAs on host immune response. We identified some predicted targets of such schistosome miRNAs in the *Biomphalaria* immune reference transcriptome. We found that schistosome miRNAs identified in the snails 24h after sympatric infection may target 43.5% of the differentially regulated immune transcripts (fig. 7). In contrast, far fewer correspondences were identified for the allopatric interactions (fig. 7). The higher proportion of targeted genes in the sympatric interaction may result in the observed efficient immunosuppression, demonstrating a specific co-evolution or adaptation in the transcriptional regulation between sympatric host and parasite. However, even if more host immune genes appeared to be targeted in the sympatric combination compared to the allopatric one's (fig. 7), both sympatric and allopatric interactions displayed the same phenotype of compatibility. In allopatric condition, despite the activation of the snail immune response, the parasite is able to develop and succeed to

infect the host. This similarity in compatibility phenotype between sympatric and allopatric parasites

could potentially results from their ability to target host immune weapons or host genes that regulate

innate cellular response. In this context, we were expecting to find miRNAs shared between sympatric

and allopatric parasites, miRNAs that do tend to weaken the immune system. A unique miRNA was

found to be shared by allopatric and sympatric parasites, sma-miR-190-3p, which has been identified

and predicted to bind different targets among which the Fibrinogen Related Protein (FREP) and

biomphalysin. The FREP family members are known as pathogen recognition receptors (Adema *et al.*,

1997; Zhang *et al.*, 2007))**.** Moreover, FREP knockdown was associated with an increase of snail

compatibility toward *Schistosoma* infections (Hanington *et al.*, 2012; Jiang *et al.*, 2006). Biomphalysin

is a key humoral factor involved in cytotoxic/cytolytic activities against *Schistosoma* parasites with the

ability to bind miracidia and sporocyst surfaces (Galinier *et al.*, 2013; Tetreau *et al.*, 2017). Moreover,

the associate snail-transcripts to these molecules are mostly down represented in sympatric and

allopatric interactions. The miRNA sma-miR-190-3p is perhaps essential for parasite compatibility.

Parasites expressing such miRNA would thus be considered as highly virulent parasites with strong

infecting capabilities. By producing dedicated miRNAs, the parasites were potentially able to regulate

transcriptional, post-transcriptional, translational and protein stability processes that might help them

to subvert the snail's immune defenses.

Compatibility reflects the outcome of complex immunobiological interactions and depends on: (i) the

ability of the snail immune system to recognize and kill the parasite; and (ii) the ability of the parasite

to circumvent or evade the host immune response (Guillou *et al.*, 2007; Mone *et al.*, 2010; Van Der

Knaap, Loker, 1990). Based on the present observations, we propose that sympatric and allopatric

interactions trigger totally different responses. In the sympatric interaction, the parasite is able to

induce a host immunosuppression within the first day of infection enabling it to quickly infect the host

and readily begins its development. In the allopatric interaction, the parasite is not able to quickly

neutralize the host immune system, and as a consequence the parasite is recognized by host defense

system that mounts a potent immune response. The need for allopatric parasites to resist the immune

system seems to disrupt the activation of their developmental program during the first day of infection.

However, they seemed to be rapidly able to protect themselves against the host immune response and

develop normally in snail tissues 96h post-infection. Thereafter, in the medium- or long-term, there are

no observable differences in the prevalence, intensity, or snail survival comparing sympatric and

allopatric interactions (supplementary table S1, supplementary fig. S2).

Thus, we show that despite having similar prevalence phenotypes, sympatric and allopatric snail-

Schistosoma interactions displayed a very different immunobiological dialogue at the molecular level.

Intriguingly, these different immunobiological interactions seem to have no repercussions upon

parasite growth at longer term or to host survival. As differences at the molecular level do not

correspond apparently to any ecologically meaningful changes in term of fitness, it is not

straightforward to demonstrate local adaptation in such systems. However, we do not know if fitness

costs could affect other biological traits in sympatric and allopatric interactions, as for example

sporocyst II or cercariae production, or cercariae infectivity and pathogenicity towards the vertebrate

host. Demonstrating local adaptation would thus appear extremely complex and would indeed deserve

further investigation. It is hard to draw the line as to when local adaptation is or is not present.

However, our results argue that the differences find at the molecular level may ultimately contribute to

the evolution of local adaptation at an ecological level.

Nevertheless, the ability for allopatric pathogens to adapt rapidly and efficiently to new hosts could

have critical consequences on disease emergence and risk of Schistosomiasis outbreaks.

Past events of allopatric parasites reaching new areas of transmission, even in large-geographic scale

dispersion, have been largely documented. The most famous example being the Schistosomiasis

colonization of South America since the slave trade of the 16th-19th Centuries (Crellen *et al.*, 2016;

Lockyer *et al.*, 2003). *Schistosoma* originated in Asia, reached Africa 12 to 19 million years ago

(MYA), and gave rise to all *Schistosoma* species known in Africa (Lockyer *et al.*, 2003). *S. mansoni*

diverged from *S. rodhaini* around 2.8MYA (Crellen *et al.*, 2016; Morgan *et al.*, 2005), and

thereafter,400 to 500 years ago, colonized South America (Crellen *et al.*, 2016; Lockyer *et al.*, 2003).

This colonization of South America by *S. mansoni* from Africa was rendered possible by the presence

of the snail host: *Biomphalaria glabrata*. All African species of *Biomphalaria* are monophyletic and

seem to have originated from paraphyletic South American clades (Campbell *et al.*, 2000; DeJong *et*

al., 2001; Theron, 2005). The ancestor of *B. glabrata* appears to have colonized Africa 1 to 5 MYA,

giving rise to all 12 species of *Biomphalaria* known today in Africa (Morgan *et al.*, 2001). In South

America and Caribbean Island, *S. mansoni* infects *B. glabrata*; in Africa, it infects mostly *B. pfeifferi*

and *B. alexandrina*. We found that South American *S. mansoni* parasites are highly compatible with

their sympatric South American snail hosts, whereas African *S. mansoni* parasites display low

compatibility phenotype with South American snail hosts (supplementary table S1). Interestingly, the

South American parasites did not lose their compatibility for African snail hosts; i.e., prevalences are

similar to African parasites when confronted to African snails (supplementary table S1). The recent

African origin of South American *Schistosoma* parasites (introduction in South America 400 to 500

years ago) may explain that they have not diverged sufficiently in South America to lose their

compatibility for African snail hosts. In this case, the transfer of allopatric parasites from Africa to

South American snail hosts have be successful and result in the emergence of Schsistosomiaisi in

South America.

More recently another case of compatible allopatric parasite emergence have been observed when

Schistososmiasis have reach Europe (Boissier *et al.*, 2016; Boissier *et al.*, 2015). Here, humans

infected in Senegal have imported a hybrid between *Schistosoma haematobium* and *Schistosoma bovis*

into Corsica. In this case urogenital schistosomiasis could be introduced and easily and rapidly spread

into this novel area of south Corsica because *Bulinus truncatus* the vector snail of *S. haematobium* was

endemic in the Corsica Cavu River (Boissier *et al.*, 2016; Boissier *et al.*, 2015). However, this

allopatric African hybrid parasite was able to adapt efficiently to the Corsican new *B. truncatus* host.

If parasite hybridization can potentially have a putative role in increasing the colonization potential of

such *S. haematobium*, it would be particularly interesting to analyze and depict the molecular support

of such allopatric interactions to predict the potential risk of schistosomiasis outbreaks in other

European areas, or other potential transmission foci.

Conclusions

If we hope to draw conclusions regarding the existence of emerging or outbreak risks, we need

to develop integrative approaches to explore fine-scale patterns of host-parasite interactions. We must

consider the spatial scale at which comparisons are conducted, the patterns of disease occurrence, the

population genetics, and the involvement of physiological, immunological, and molecular processes.

Studying the relevant factors at the relevant timing would be of critical importance in terms of

Schistosomiasis control. Understanding further, how these allopatric parasites efficiently infect host

snails would be mandatory to identify markers and develop new tools to predict or to quantify risks of

Schistosomiasis outbreaks.

Acknowledgements

We thank Ms. Cécile Saint-Béat, Ms. Nathalie Arancibia, and Ms. Anne Rognon for their

work and diligence in helping generate some of the data described herein.

707
708
709

Declarations

Ethics approval and consent to participate

Our laboratory holds permit # A66040 for experiments on animals, which was obtained from the French Ministry of Agriculture and Fisheries and the French Ministry of National Education, Research, and Technology. The housing, breeding, and care of the utilized animals followed the ethical requirements of our country. The experimenter possesses an official certificate for animal experimentation from both of the above-listed French ministries (Decree # 87–848, October 19, 1987). The various protocols used in this study have been approved by the French veterinary agency of the DRAAF Languedoc-Roussillon (Direction Régionale de l'Alimentation, de l'Agriculture et de la Forêt), Montpellier, France (authorization # 007083).

Availability of data and material

The datasets generated and/or analysed during the current study are not publicly available due to ongoing analysis on transcriptome but are available from the corresponding author on reasonable request.

Raw read fastq files were submitted to the Sequence Read Archive at NCBI (http://trace.ncbi.nlm.nih.gov/Traces/sra/) under the reference PRJNA213050. The reference transcriptome is publicly available from the transcriptomic database of *Biomphalaria glabrata* at the 2ei website (http://2ei.univ-perp.fr/?page_id=89 or http://ihpe.univ-perp.fr/acces-aux-donnees/)

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the French National Agency for Research (ANR) through the Invimory project grant [ANR-13-JSV7-0009] to BG.

Funding agency has no role in study design, collection, analysis, and interpretation of data and in writing of the manuscript.

Authors' contributions

Conceived and designed the experiments: AP, SP, RG, DD, BG. Performed the experiments: AP, SP, CC, GMC, JFA. Analyzed the data: AP, SP, CC, RG, GMC, DD, BG. AP, RG, BG led the manuscript writing. All authors participated to manuscript writing, editing, and critical reviewing. All authors read and approved the final manuscript.

Authors' information

SP current position Welcome Trust Sanger Institute, CB10 1SA Hinxton, United Kingdom

References:

Adema CM, Hertel LA, Miller RD, Loker ES (1997) A family of fibrinogen-related proteins that precipitates parasite-derived molecules is produced by an invertebrate after infection. *Proc Natl Acad Sci U S A* **94**, 8691-8696.

Baeza Garcia A, Pierce RJ, Gourbal B*, et al.* (2010) Involvement of the cytokine MIF in the snail host immune response to the parasite Schistosoma mansoni. *PLoS Pathogens* **6**, e1001115.

Bankers L, Fields P, McElroy KE*, et al.* (2018) Genomic evidence for population-specific responses to co-evolving parasites in a New Zealand freshwater snail. *Mol Ecol* **26**, 3663-3675.

Basch PF (1976) Intermediate host specificity in Schistosoma mansoni. *Experimental Parasitology* **39**, 150-169.

Bender RC, Goodall CP, Blouin MS, Bayne CJ (2007) Variation in expression of Biomphalaria glabrata SOD1: a potential controlling factor in susceptibility/resistance to Schistosoma mansoni. *Dev Comp Immunol* **31**, 874-878.

Bohdanowicz M, Schlam D, Hermansson M*, et al.* (2013) Phosphatidic acid is required for the constitutive ruffling and macropinocytosis of phagocytes. *Molecular Biology of the Cell* **24**, 1700-1712, S1701-1707.

Boissier J, Grech-Angelini S, Webster BL*, et al.* (2016) Outbreak of urogenital schistosomiasis in Corsica (France): an epidemiological case study. *The Lancet Infectious Diseases* **16**, 971-979.

Boissier J, Mone H, Mitta G*, et al.* (2015) Schistosomiasis reaches Europe. *The Lancet Infectious Diseases* **15**, 757-758.

Bonner KM, Bayne CJ, Larson MK, Blouin MS (2012) Effects of Cu/Zn superoxide dismutase (sod1) genotype and genetic background on growth, reproduction and defense in Biomphalaria glabrata. *PLoS Negl Trop Dis* **6**, e1701.

Bridger JM, Arican-Gotkas HD, Foster HA*, et al.* (2014) The non-random repositioning of whole chromosomes and individual gene loci in interphase nuclei and its relevance in disease, infection, aging, and cancer. *Adv Exp Med Biol* **773**, 263-279.

Campbell G, Jones CS, Lockyer AE*, et al.* (2000) Molecular evidence supports an african affinity of the neotropical freshwater gastropod, Biomphalaria glabrata, say 1818, an intermediate host for Schistosoma mansoni. *Proceedings of the Royal Society B* **267**, 2351-2358.

Cavalcanti MG, Filho FC, Mendonca AM*, et al.* (2012) Morphological characterization of hemocytes from Biomphalaria glabrata and Biomphalaria straminea. *Micron* **43**, 285-291.

Crellen T, Allan F, David S*, et al.* (2016) Whole genome resequencing of the human parasite Schistosoma mansoni reveals population history and effects of selection. *Sci Rep* **6**, 20954.

DeJong RJ, Morgan JA, Paraense WL*, et al.* (2001) Evolutionary relationships and biogeography of Biomphalaria (Gastropoda: Planorbidae) with implications regarding its role as host of the human bloodfluke, Schistosoma mansoni. *Mol Biol Evol* **18**, 2225-2239.

Dheilly NM, Duval D, Mouahid G*, et al.* (2015) A family of variable immunoglobulin and lectin domain containing molecules in the snail Biomphalaria glabrata. *Dev Comp Immunol* **48**, 234-243.

Dybdahl MF, Storfer A (2003) Parasite local adaptation: Red Queen versus suicide king.
TRENDs in Ecology and Evolution **18**, 523-530.
Ebert D (1994) Virulence and local adaptation of horizontally transmitted parasite. *Science*
265, 1084-1086.
Ebert D (1998) Experimental evolution of parasites. *Science* **282**, 1432-1435.
Enright AJ, John B, Gaul U*, et al.* (2003) MicroRNA targets in Drosophila. *Genome Biology* **5**,
R1.
Frank SA (1996) Models of parasite virulence. *Q Rev Biol* **71**, 37-78.
Galinier R, Portela J, Mone Y*, et al.* (2013) Biomphalysin, a new beta pore-forming toxin
involved in Biomphalaria glabrata immune defense against Schistosoma mansoni.
PLoS Pathogens **9**, e1003216.
Galinier R, Roger E, Mone Y*, et al.* (2017) A multistrain approach to studying the mechanisms
underlying compatibility in the interaction between Biomphalaria glabrata and
Schistosoma mansoni. *PLoS Negl Trop Dis* **11**, e0005398.
Gilges D, Vinit MA, Callebaut I*, et al.* (2000) Polydom: a secreted protein with pentraxin,
complement control protein, epidermal growth factor and von Willebrand factor A
domains. *Biochemical Journal* **352 Pt 1**, 49-59.
Goodall CP, Bender RC, Brooks JK, Bayne CJ (2006) Biomphalaria glabrata cytosolic
copper/zinc superoxide dismutase (SOD1) gene: association of SOD1 alleles with
resistance/susceptibility to Schistosoma mansoni. *Mol Biochem Parasitol* **147**, 207-
210.
Greischar MA, Koskella B (2007) A synthesis of experimental work on parasite local
adaptation. *Ecol Lett* **10**, 418-434.
Grkovich A, Dennis EA (2009) Phosphatidic acid phosphohydrolase in the regulation of
inflammatory signaling. *Advances in Enzyme Regulation* **49**, 114-120.
Guillou F, Roger E, Mone Y*, et al.* (2007) Excretory-secretory proteome of larval Schistosoma
mansoni and Echinostoma caproni, two parasites of Biomphalaria glabrata. *Mol*
Biochem Parasitol.
Hahn UK, Bender RC, Bayne CJ (2001) Involvement of nitric oxide in killing of Schistosoma
mansoni sporocysts by hemocytes from resistant Biomphalaria glabrata. *Journal of*
Parasitology **87**, 778-785.
Hanington PC, Forys MA, Loker ES (2012) A somatically diversified defense factor, FREP3, is a
determinant of snail resistance to schistosome infection. *PLoS Negl Trop Dis* **6**, e1591.
Ittiprasert W, Knight M (2012) Reversing the resistance phenotype of the Biomphalaria
glabrata snail host Schistosoma mansoni infection by temperature modulation. *PLoS*
Pathogens **8**, e1002677.
Ittiprasert W, Miller A, Myers J*, et al.* (2010) Identification of immediate response genes
dominantly expressed in juvenile resistant and susceptible Biomphalaria glabrata
snails upon exposure to Schistosoma mansoni. *Molecular and Biochemical*
Parasitology **169**, 27-39.
Ittiprasert W, Nene R, Miller A*, et al.* (2009) Schistosoma mansoni infection of juvenile
Biomphalaria glabrata induces a differential stress response between resistant and
susceptible snails. *Experimental Parasitology* **123**, 203-211.
Jiang Y, Loker ES, Zhang SM (2006) In vivo and in vitro knockdown of FREP2 gene expression
in the snail Biomphalaria glabrata using RNA interference. *Dev Comp Immunol* **30**,
855-866.

Kalbe M, Eizaguirre C, Scharsack JP, Jakobsen PJ (2016) Reciprocal cross infection of
sticklebacks with the diphyllobothriidean cestode Schistocephalus solidus reveals
consistent population differences in parasite growth and host resistance. *Parasit*
Vectors **9**, 130.

Kawecki T, Ebert D (2004) Conceptual issues in local adaptation. *Ecology letters* **7**, 1225-
1241.

Kincaid-Smith J, Rey O, Toulza E, Berry A, Boissier J (2017) Emerging Schistosomiasis in
Europe: A Need to Quantify the Risks. *Trends in Parasitology* **33**, 600-609.

Knight M, Elhelu O, Smith M*, et al.* (2015) Susceptibility of Snails to Infection with
Schistosomes is influenced by Temperature and Expression of Heat Shock Proteins.
Epidemiology **5**, pii: 189. Epub.

Larson MK, Bender RC, Bayne CJ (2014) Resistance of Biomphalaria glabrata 13-16-R1 snails
to Schistosoma mansoni PR1 is a function of haemocyte abundance and constitutive
levels of specific transcripts in haemocytes. *International Journal for Parasitology* **44**,
343-353.

Li S, Shen L, Sun L*, et al.* (2017) Small RNA-Seq analysis reveals microRNA-regulation of the
Imd pathway during Escherichia coli infection in Drosophila. *Dev Comp Immunol* **70**,
80-87.

Lively CM (1989) Adaptation by a Parasitic Trematode to Local Populations of Its Snail Host.
Evolution **43(8)**.

Lockyer AE, Emery AM, Kane RA*, et al.* (2012) Early differential gene expression in
haemocytes from resistant and susceptible Biomphalaria glabrata strains in response
to Schistosoma mansoni. *PLoS One* **7**, e51102.

Lockyer AE, Noble LR, Rollinson D, Jones CS (2004) Schistosoma mansoni: resistant specific
infection-induced gene expression in Biomphalaria glabrata identified by fluorescent-
based differential display. *Experimental Parasitology* **107**, 97-104.

Lockyer AE, Olson PD, Ostergaard P*, et al.* (2003) The phylogeny of the Schistosomatidae
based on three genes with emphasis on the interrelationships of Schistosoma
Weinland, 1858. *Parasitology* **126**, 203-224.

Lockyer AE, Spinks JN, Walker AJ*, et al.* (2007) Biomphalaria glabrata transcriptome:
identification of cell-signalling, transcriptional control and immune-related genes
from open reading frame expressed sequence tags (ORESTES). *Developmental and*
Comparative Immunology **31**, 763-782.

Meningher T, Lerman G, Regev-Rudzki N*, et al.* (2017) Schistosomal MicroRNAs Isolated
From Extracellular Vesicles in Sera of Infected Patients: A New Tool for Diagnosis and
Follow-up of Human Schistosomiasis. *Journal of Infectious Diseases* **215**, 378-386.

Minchella DJ (1985) Host life-history variation in response to parasitism. *Parasitology* **90**,
205-216.

Mitta G, Adema CM, Gourbal B, Loker ES, Theron A (2012) Compatibility polymorphism in
snail/schistosome interactions: From field to theory to molecular mechanisms. *Dev*
Comp Immunol **37**, 1-8.

Mitta G, Gourbal B, Grunau C*, et al.* (2017) The Compatibility Between Biomphalaria glabrata
Snails and Schistosoma mansoni: An Increasingly Complex Puzzle. *Adv Parasitol* **97**,
111-145.

Mone Y, Gourbal B, Duval D*, et al.* (2010) A large repertoire of parasite epitopes matched by
a large repertoire of host immune receptors in an invertebrate host/parasite model.
PLoS Negl Trop Dis **4**.

Mone Y, Ribou AC, Cosseau C*, et al.* (2011) An example of molecular co-evolution: reactive
oxygen species (ROS) and ROS scavenger levels in Schistosoma
mansoni/Biomphalaria glabrata interactions. *International Journal for Parasitology*
41, 721-730.

Morand S, Manning SD, Woolhouse ME (1996) Parasite-host coevolution and geographic
patterns of parasite infectivity and host susceptibility. *Proceedings of the Royal*
Society B **263**, 119-128.

Morgan JA, Dejong RJ, Adeoye GO*, et al.* (2005) Origin and diversification of the human
parasite Schistosoma mansoni. *Mol Ecol* **14**, 3889-3902.

Morgan JA, Dejong RJ, Snyder SD, Mkoji GM, Loker ES (2001) Schistosoma mansoni and
Biomphalaria: past history and future trends. *Parasitology* **123 Suppl**, S211-228.

Myers J, Ittiprasert W, Raghavan N, Miller A, Knight M (2008) Differences in cysteine
protease activity in Schistosoma mansoni-resistant and -susceptible Biomphalaria
glabrata and characterization of the hepatopancreas cathepsin B Full-length cDNA.
Journal of Parasitology **94**, 659-668.

Nelson MK, Cruz BC, Buena KL, Nguyen H, Sullivan JT (2016) Effects of abnormal temperature
and starvation on the internal defense system of the schistosome-transmitting snail
Biomphalaria glabrata. *J Invertebr Pathol*.

Nowacki FC, Swain MT, Klychnikov OI*, et al.* (2015) Protein and small non-coding RNA-
enriched extracellular vesicles are released by the pathogenic blood fluke
Schistosoma mansoni. *Journal of Extracellular Vesicles* **4**, 28665.

Osnas EE, Hurtado PJ, Dobson AP (2015) Evolution of pathogen virulence across space during
an epidemic. *Am Nat* **185**, 332-342.

Osnas EE, Lively CM (2004) Parasite dose, prevalence of infection and local adaptation in a
host-parasite system. *Parasitology* **128**, 223-228.

Osnas EE, Lively CM (2005) Immune response to sympatric and allopatric parasites in a snail-
trematode interaction. *Frontiers in Zoology* **2**, 8.

Penczykowski RM, Laine AL, Koskella B (2015) Understanding the ecology and evolution of
host-parasite interactions across scales. *Evol Appl* **9**, 37-52.

Perrin C, Lepesant JM, Roger E*, et al.* (2012) Schistosoma mansoni mucin gene (SmPoMuc)
expression: epigenetic control to shape adaptation to a new host. *PLoS Pathogens* **9**,
e1003571.

Pila EA, Gordy MA, Phillips VK*, et al.* (2016) Endogenous growth factor stimulation of
hemocyte proliferation induces resistance to Schistosoma mansoni challenge in the
snail host. *Proc Natl Acad Sci U S A* **113**, 5305-5310.

Pinaud S, Portela J, Duval D*, et al.* (2016) A Shift from Cellular to Humoral Responses
Contributes to Innate Immune Memory in the Vector Snail Biomphalaria glabrata.
PLoS Pathogens **12**, e1005361.

Portet A, Pinaud S, Tetreau G*, et al.* (2017) Integrated multi-omic analyses in Biomphalaria-
Schistosoma dialogue reveal the immunobiological significance of FREP-SmPoMuc
interaction. *Developmental and Comparative Immunology* **doi:**
10.1016/j.dci.2017.02.025.

Prugnolle F, de Meeus T, Pointier JP*, et al.* (2006) Geographical variations in infectivity and
susceptibility in the host-parasite system Schistosoma mansoni/Biomphalaria
glabrata: no evidence for local adaptation. *Parasitology* **133**, 313-319.

Raghavan M, Wijeyesakere SJ, Peters LR, Del Cid N (2013) Calreticulin in the immune system:
ins and outs. *Trends in Immunology* **34**, 13-21.

Richards CS (1975) Genetic factors in susceptibility of Biomphalaria glabrata for different
strains of Schistosoma mansoni. *Parasitology* **70**, 231-241.
Robertson S, Bradley JE, MacColl AD (2017) No evidence of local adaptation of immune
responses to Gyrodactylus in three-spined stickleback (Gasterosteus aculeatus). *Fish*
and Shellfish Immunology **60**, 275-281.
Roger E, Grunau C, Pierce RJ*, et al.* (2008a) Controlled Chaos of Polymorphic Mucins in a
Metazoan Parasite (Schistosoma mansoni) Interacting with Its Invertebrate Host
(Biomphalaria glabrata). *PLoS Negl Trop Dis* **2**, e330.
Roger E, Mitta G, Mone Y*, et al.* (2008b) Molecular determinants of compatibility
polymorphism in the Biomphalaria glabrata/Schistosoma mansoni model: New
candidates identified by a global comparative proteomics approach. *Mol Biochem*
Parasitol **157**, 205-216.
Ryu SJ, Park SC (2009) Targeting major vault protein in senescence-associated apoptosis
resistance. *Expert Opinion on Therapeutic Targets* **13**, 479-484.
Sminia T, Barendsen L (1980) A comparative morphological and enzyme histochemical study
on blood cells of the freshwater snails, *Lymnaea stagnalis*, *Biomphalaria glabrata*,
and *Bulinus truncatus*. *Journal of Morphology* **165**, 31-39.
Tack AJ, Thrall PH, Barrett LG, Burdon JJ, Laine AL (2012) Variation in infectivity and
aggressiveness in space and time in wild host-pathogen systems: causes and
consequences. *J Evol Biol* **25**, 1918-1936.
Tennessen JA, Theron A, Marine M*, et al.* (2015) Hyperdiverse gene cluster in snail host
conveys resistance to human schistosome parasites. *PLoS Genet* **11**, e1005067.
Tetreau G, Pinaud S, Portet A*, et al.* (2017) Specific Pathogen Recognition by Multiple Innate
Immune Sensors in an Invertebrate. *Front Immunol* **8**, 1249.
Theron A (2005) Les schistosomes et leurs hôtes : apport des marqueurs moléculaires à la
connaissance de leur phylogéographie, épidémiologie et écologie. *Bulletin de la*
Société Zoologique de France **130** 205-218.
Theron A, Pages JR, Rognon A (1997) Schistosoma mansoni: distribution patterns of
miracidia among Biomphalaria glabrata snail as related to host susceptibility and
sporocyst regulatory processes. *Experimental Parasitology* **85**, 1-9.
Theron A, Rognon A, Gourbal B, Mitta G (2014) Multi-parasite host susceptibility and multi-
host parasite infectivity: a new approach of the Biomphalaria glabrata/Schistosoma
mansoni compatibility polymorphism. *Infection Genetics and Evolution* **26**, 80-88.
Van Der Knaap WPW, Loker ES (1990) Immune mechanisms in trematod-snail interactions.
Parasitology Today **6**, 175-182.
Walzer KA, Chi JT (2017) Trans-kingdom small RNA transfer during host-pathogen
interactions: The case of P. falciparum and erythrocytes. *RNA Biology*, 1-8.
WHO (2010) Schistosomiasis *Fact Sheet No. 115. World Health Organisation.*
Zawada AM, Zhang L, Emrich IE*, et al.* (2017) MicroRNA profiling of human intermediate
monocytes. *Immunobiology* **222**, 587-596.
Zhang H, Song L, Li C*, et al.* (2007) Molecular cloning and characterization of a thioester-
containing protein from Zhikong scallop Chlamys farreri. *Molecular Immunology* **44**,
3492-3500.
Zhao X, Yu H, Kong L, Liu S, Li Q (2016) High throughput sequencing of small RNAs
transcriptomes in two Crassostrea oysters identifies microRNAs involved in osmotic
stress response. *Sci Rep* **6**, 22687.
983

984
985

Figure Legends

Fig. 1: Dual-RNAseq of Biomphalaria immune-related transcripts

Among the differentially represented transcripts, Blast2GO functional annotation allowed us to identify 336 transcripts that appeared to be related to the *Biomphalaria* immune response. Abbreviations and colors: blue BB, sympatric interaction between BgBRE and SmBRE; green BV, allopatric interaction between BgBRE and SmVEN; and red BR, allopatric interaction between BgBRE and Srod. For each interaction 40 whole-snails are used, 20 pooled at 24h and 20 at 96h post-infection. A) Venn diagram showing the relationships among the immune transcripts found to be differentially expressed in the sympatric and allopatric interactions. B) Clustering of differentially represented immune transcripts. Heatmap representing the profiles of the 336 differentially represented immune-related transcripts in the BB, BV, or BR interactions along the kinetic of infection (at 24 and 96 h). Each transcript is represented once and each line represents one transcript. Colors: yellow, over-represented transcripts; purple, under-represented transcripts; and black, unchanged relative to levels in control naïve snails. C) Pie chart showing the distribution of the selected immune-related transcripts across three immunological processes: immune recognition (pink), immune effectors (brown), and immune signaling (blue). For each category and interaction, the respective proportion of transcripts and the direction of the effect (over- or underexpression) are indicated.

Fig. 2: Differentially represented immune-related transcripts in sympatric and allopatric interactions.

Cumulative expression [Log2FC (fold change) from DESeq2 analysis] of the immune-related transcripts identified as being differentially represented following sympatric or allopatric infection. Transcripts were grouped into the three immunological groups described in Fig. 1, and from there into functional categories. The yellow histograms correspond to cumulatively over-represented transcripts, while the purple histograms show under-represented transcripts. The black (over-represented) and gray (under-represented) diamonds correspond to the number of transcripts analyzed in each functional category. Abbreviations: BB, BgBRE/SmBRE interaction; BV, BgBRE/SmVEN interaction; and BR, BgBRE/Srod interaction. A. Immune transcript expression at 24 h post-infection. B. Immune transcript expression at 96 h post-infection.

Fig. 3: Microscopic analyse of snail hemocyte proliferations

In vitro EdU labeling of hemocytes was conducted for sympatric and allopatric interactions A) Hemocytes were collected at 24 h post-infection for *in vitro* analysis. Confocal microscopy of EdU-labeled hemocytes from snails subjected to the allopatric BV interaction (BgBRE/SmVEN). Colors: blue, DAPI; and green, EdU. B) Microscopic counting of EdU-labeled hemocytes from naïve control snails (BgBRE) (n=1,811) and those subjected to the sympatric interaction (BB: BgBRE/SmBRE) (n=2,064) or an allopatric interaction (BV: BgBRE/SmVEN) (n=1,366) recovered from 3 individual snails by condition. Colors: green, EdU-positive cells; and blue, EdU-negative cells. Between-group differences in the percentage of proliferation were tested using a Fisher exact test, with statistical significance accepted at p<0.05. The "a" indicates a significant difference between the naïve and infective conditions, while "b" indicates a significant difference between the infective conditions.

Fig. 4: Flow cytometry analyse of the hemocyte response in sympatric and allopatric interactions

A) Flow cytometry was used to count *in vivo* EdU-labeled hemocytes at 24 h and 96 h after infection in sympatric and allopatric interactions. A total number of hemocytes of n=10,000were recoveredfor6 biological replicates of 3 snails. Control naïve snails (BgBRE, yellow) were compared to those subjected to the sympatric interaction (BB, BgBRE/SmBRE, blue) or an allopatric interaction (BV, BgBRE/SmVEN, green).

B) The experiment described in A was repeated using the BgVEN snail strain. Control naïve snails (BgVEN, yellow) were compared to those subjected to the sympatric interaction (VV, BgVEN/SmVEN, green) or an allopatric interaction (VB, BgVEN/SmBRE, blue).

C) FSC (forward-scattered light, representing cell size) and SSC (side-scattered light, representing cell granularity) circulating hemocyte patterns in BgBRE snails under the naïve condition (yellow) or 24 h and 96 h after infections in sympatry (BB24/96, BgBRE/SmBRE, blue) or allopatry (BV24/96, BgBRE/SmVEN, green).

D) FSC and SSC circulating hemocyte patterns in BgVEN snails under the naïve condition (yellow) or 24 h and 96 h after infections in sympatry (VV24/96, BgVEN/SmVEN, blue) or allopatry (VB24/96, BgVEN/SmBRE, green). The red dots correspond to EdU-positive hemocytes.

Between-group differences in the percentage of proliferation were tested using the Mann-Whitney U-test, with statistical significance accepted at p<0.05. The "a" indicates a significant difference between the naïve and infective condition, "b" indicates a significant difference between the infective conditions at 24h, and "c" indicates a significant difference between the infective conditions at 96h.

Fig. 5: Development of parasites into snail tissues

A histological approach was used to monitor parasite size along the course of snail infection. The sympatric interaction (BgBRE x SmBRE) is shown in blue, and the allopatric interaction (BgBRE x SmVEN) is shown in green. For each experimental interaction, the parasite sizes were quantified at 24 and 96 h after infection. Morpho-anatomical aspects of the parasite are depicted to highlight a potential difference in parasite survival. N=7 to 10sporocystes were used as indicated in the figure.Between-group parasite size differences were assessed using the Mann-Whitney U-test, with significance accepted at p<0.05 (indicated by "a" on the histograms).

Fig. 6: Clustering of intra-molluscalSchistosoma expression patterns

RNAseq library mapping enabled us to identify 351 genes expressed by *Schistosoma* parasites in *Biomphalaria* snail tissues. Colors: blue, *S. mansoni* Brazil (SmBRE); green, *S. mansoni* Venezuela (SmVEN); and red, *S. rodhaini* (Srod). The heatmap represents the profiles of the 351 genes expressed by the different parasites at 24 h after infection. Each transcript is represented once and each line represents one transcript. The expression level is highlighted by the different shades of blue.

Fig. 7: In-silico identification of parasite miRNAs

miRNAs were assessed using libraries obtained from naïve snails and snails infected for 24 h under the various interaction conditions (BB24, BgBRE x SmBRE; BV24, BgBRE x SmVEN; BR24, BgBRE x Srod). A) Table highlighting the precursor miRNAs that may have targets among the immune-related snail transcripts selected in the present work. They include eight precursors specifically recovered in BB24, two in BV24, and one shared across the three infected conditions. The total numbers of potential targets in each condition are indicated. B) Venn diagram showing the potential targets according to the sympatric or allopatric interactions. Shown is an example miRNA stem-loop precursor that presents the highest number of potential host targets.

Supplementary fig. S1: Clustering of all differential represented transcripts

Clustering of differentially represented transcripts. Heatmap representing the profiles of the 1,895 differentially represented immune-related transcripts in the BB, BV, or BR interactions along the time course of infection (at 24 and 96 h). Each transcript is represented once and each line represents one transcript. Colors: yellow, over-represented transcripts; purple, under-represented transcripts; and black, unchanged relative to levels in control naïve snails.

1

Supplementary fig. S2: Mortality of B. glabrata snails after infections

The survival rates of *B. glabrata* subjected to infection by different *S. mansoni* strains were observed over 14 weeks. Kaplan Meier graphs were generated using xlstat, and the log-rank test (p < 0.05) was used to test for significant between-group differences. Colored curves indicate the mortality rates of naïve snails (yellow) (n=60), snails infected by the sympatric parasite (BB, BgBRE/SmBRE, blue) (n=50), and snails infected by the allopatric parasite (BV, BgBRE/SmVEN, green) (n=50). The difference in mortality between naïve and infected snails was significant (p<0.05), whereas that between the two infected conditions was not (p = 0.243).

 Table 1: Number of transcripts in each step of transcriptomic analysis

Analysis of transcriptomic data		Analyses of Blast2GO	Analyses of annotations	
Full Transcriptome	Differentially expressed	Informative annotation	Immune Transcripts	Non Immune Transcripts
159,711	3,865	1,017	336	681

1095

Supplementary table S1: *Biomphalaria* and *Schistosoma* compatibility between African and South-

American strains

			Biomphalaria		
			glabrata		*pfeifferi*
			BRE	VEN	SEN
Schistosoma mansoni	BRE	P %	**100**	70	92
		I	**3.6**	3.5	2.9
	VEN	P %	100	**95**	100
		I	3.2	**3.8**	4
	SEN	P %	27	13	**100**
		I	1.3	1	**2.5**
Schistosoma rodhaini		P %	80	*NA*	*NA*
		I	2	*NA*	*NA*

1098
1099
The prevalence (P %: percentage of snail infected) and intensity (I: number of parasites per
infected host) of infection are presented for each experimental infection. The indicated values
correspond to 20 miracidia. Each pairwise combination of *Biomphalaria* (BgBRE, BgVEN,
BpSEN) and *Schistosomamansoni* (SmBRE, SmVEN, SmSEN) or *SchistosomaRodhaini*
(Srod) were tested for compatibility. *NA*: non-available data.

1105
1106
1107

    Figure 1

   Figure 2

1126   Figure 3
1127

1134   Figure 4

1135

1136
1137
1138
1139

1140   Figure 5
1141

1149   Figure 6
1150

A

Precursor	BgBRE duplicates		BB24	BV24	BR24	Targets
sma-mir-8460	0	0	1	0	0	7
sma-mir-8463	0	0	1	0	0	9
sma-mir-2a	0	0	1	0	0	20
sma-mir-2d	0	0	1	0	0	50
sma-mir-3492	0	0	1	0	0	10
sma-mir-8404	0	0	1	0	0	4
sma-mir-8414	0	0	1	0	0	14
sma-mir-8446	0	0	1	0	0	23
sma-mir-8456	0	0	0	1	0	2
sma-mir-8431	0	0	0	1	0	4
sma-mir-190	0	0	1	1	1	17
Sum of potential Bg targets	**0**	**0**	**154**	**23**	**17**	

SmBRE specific | SmVEN specific | Shared

B

sma-mir-8431
E=-29.9
4 putative targets

SmVEN specific

Srod specific

SmBRE specific

Shared

sma-mir-2d-3p
E=-29.6
50 putative targets

sma-mir-190-3p
E=-43.8
17 putative targets

Lowest free energy structure
Probability >= 99%
99% > Probability >= 95%
95% > Probability >= 90%
90% > Probability >= 80%
80% > Probability >= 70%
70% > Probability >= 60%
60% > Probability >= 50%
50% > Probability

Prediction from ~/RNA structure web

5'-UAAUUUGAUGUUAUUAAGG-3'
5'-GUUCUUAAUGUUGUAUUAGUU-3'
5'-UAUCACAGUCCUGCUUAGGUGA-3'
5'-CAGUGACCAGACAUAUCCCU-3'

0   0   0
6   17   0
123 0

1164   Figure S1
1165

1166
1167
1168
1169

1170   Figure S2
1171

Naïve   Sympatric interaction   Allopatric interaction

1172
1173