Short-term insurance versus long-term bet-hedging strategies as adaptations to variable environments

Thomas Ray Haaland[1], Jonathan Wright[1], Jarle Tufto[2] and Irja Ida Ratikainen[1]

1: Department of Biology, Centre for Biodiversity Dynamics, Norwegian University of Science and Technology

2: Department of Mathematical Sciences, Centre for Biodiversity Dynamics, Norwegian University of Science and Technology

Corresponding Author: Thomas Ray Haaland, Department of Biology, NTNU, Høgskoleringen 5, Trondheim, N-7491 Norway

Abstract

Understanding how organisms adapt to environmental variation is a key challenge of biology. Central to this are bet-hedging strategies that maximize geometric mean fitness across generations, either by being conservative or diversifying phenotypes. Theoretical models of bet-hedging and the multiplicative fitness effects of environmental variation across generations have traditionally assumed that environmental conditions are constant within lifetimes. However, behavioral ecology has revealed adaptive responses to additive fitness effects of environmental variation within lifetimes, either through insurance or risk-sensitive strategies. Here we explore whether the effects of adaptive insurance interact with the evolution of bet-hedging by varying the position and skew of fitness functions within and between lifetimes. When insurance causes the optimal phenotype to shift from the peak to down the less steeply decreasing side of the fitness function, then conservative bet-hedging does not generally evolve on top of this, even if diversifying bet-hedging can. Canalization to reduce phenotypic variation within a lifetime is almost always favored, except when the tails of the fitness function are steeply convex and produce a novel risk-sensitive increase in phenotypic variance akin to diversifying bet-hedging. Importantly, using skewed fitness functions, we provide the first example of how conservative and diversifying bet-hedging strategies might coexist.

Keywords

Fluctuating selection; environmental stochasticity; variance-sensitivity; geometric mean fitness; cliff-edge effect; phenotypic canalization.

Introduction

How organisms adapt to unpredictable fluctuations in the environment has been an intriguing and important problem for many years in evolutionary biology, and especially recently when predicting adaptive responses to environmental change. Conditions may vary over different time scales, selecting for adaptations that maximize fitness in the face of environmental stochasticity in everything from labile behavioral traits within a lifetime (e.g. variance-sensitive foraging, Stephens 1981) to cross-generational effects of life-history traits (e.g. bet-hedging; Simons 2011; Starrfelt and Kokko 2012). Thus, the phenotypes we observe in organisms today have likely been shaped by environmental variation experienced across longer timescales during their evolutionary history, and trait values may not necessarily appear optimal when considering just short-term current environmental conditions (Nadeau et al. 2017). Environmental variation itself is expected to be a strong selective agent, since genotypic rather than individual fitness determines optimal strategies that are produced over evolutionary time in stochastic environments (Lewontin and Cohen 1969; McNamara 1998).

Asymmetric fitness functions pose an additional challenge to evolutionary biologists seeking to understand genotypic adaptations in variable environments (Yoshimura and Shields 1987; Urban et al. 2013). Skew in the function relating a single, continuous phenotypic trait to fitness is commonly seen in nature, occurring whenever costs and benefits differ in how they relate to increasing versus decreasing values of the phenotype, or when the strength of selection acting on the two sides of the phenotypic distribution differs. Common examples are thermal performance curves (Angilletta 2009), optimal clutch or litter sizes (Mountford 1968; Boyce and Perrins 1987), and reproductive benefits versus viability costs of sexually selected ornaments (Andersson and Iwasa 1996). In these types of scenarios, uncertainty across instances in any component determining individual fitness will cause the optimal trait value to differ from the trait value at the peak of the fitness function (Yoshimura and Shields 1987; Parker and Smith 1990). Such uncertainty in fitness pay-offs across instances is also almost ubiquitous in biological systems. Across lifetimes, phenotypic differences among individuals (as instances) of the same genotype may arise due to developmental instability creating random (uncanalized) variation in phenotypes and thus also in their fitness, and avoiding such variation may be costly (Zhang and Hill 2005). Within lifetimes, uncertainty may occur in individual energetic state on short (e.g. behavioral) timescales, due to stochastic probabilities of resource acquisition such as prey captures. In addition, the fitness effects of the phenotype itself (i.e. the shape or position of the fitness function) may be uncertain, for example

due to micro-environmental variability, or variation occurring over short time scales, such as in social
environments.

With a skewed fitness function, any stochastic environmentally induced variation in fitness pay-offs will
select for apparently suboptimal phenotypes with trait values away from the peak of the deterministic
fitness function when selection maximizes arithmetic mean fitness (Fig. 1). Finding the (arithmetic)
mean fitness in such cases involves multiplying the phenotype-specific fitnesses with the frequencies of
the different phenotypes (Mountford 1968). This phenomenon is sometimes described as the cliff-edge
effect (Vercken et al. 2012; Mitteroecker et al. 2016), and is commonly encountered as 'insurance'
strategies in fields such as behavioral ecology (Dall 2010). A well-known example is the small bird in
winter (Brodin 2007). Facing a starvation-predation trade-off, the small passerine bird wants to be as
light as possible to nimbly avoid predators during the day, but needs to store fat before nightfall, which
it metabolizes to stay warm during the night. The small bird in winter will therefore adaptively store
more fat when temperatures are more variable (Bednekoff et al. 1994), and/or when food supply is
more uncertain (Krams et al. 2010; Ratikainen and Wright 2013)

Across generations, however, the fitness of a lineage is determined by its geometric mean fitness rather
than the arithmetic mean, due to reproduction being an inherently multiplicative process (Lewontin and
Cohen 1969; Simons 2011). When environmental conditions are constant between generations (i.e. the
function relating the trait in question to fitness is exactly the same each generation), arithmetic and
geometric mean fitness are equal. However, once some aspect of the fitness function differs between
generations, creating variation in realized fitness between individuals with the same genotype, then the
geometric mean will be lower than arithmetic mean. Crucially, a change in strategy that lowers variance
in realized fitness at the genotype level may increase geometric mean fitness, and thus be selectively
favored. If such a strategy that increases geometric mean fitness at the genotype level also involves a
simultaneous decrease in arithmetic mean fitness, it is defined as bet-hedging (Philippi and Seger 1989).
Two main types of bet-hedging are usually considered, diversifying bet-hedging (DBH) and conservative
bet-hedging (CBH) (Philippi and Seger 1989; Simons 2011; Starrfelt and Kokko 2012). DBH increases
phenotypic variance and thus reduces the correlations in fitness between individuals of the same
genotype, such that not all individuals are affected by the environment in the same correlated way. DBH
strategies can include both producing two discrete 'types' of individuals, such as dry and wet-year
specialists, or dormant and active life stages in response to good and bad years (Lewontin and Cohen
1969; Venable 2007; Graham et al. 2014), and continuous variation in a trait among individuals, such as

size of offspring, or timing of reproduction and entering or exiting dormancy (Simons 2009; Devaux and

Lande 2010; Lof et al. 2012). In contrast, CBH reduces variance in realised fitness at the individual level,

such that each individual within a genotype will perform moderately well across a range of

environments. However, there is nothing stopping a strategy from combining the two types, and

Starrfelt and Kokko (2012) argue that DBH, reducing only among-individual fitness variance, and CBH,

reducing only each individual's fitness variance, are actually two ends of a continuum with strategic

combination of DBH and CBH possible in between. Despite this, neither Starrfelt and Kokko (2012) nor

later authors exploring similar models (e.g. Crowley et al. 2016) explain how such a combination of DBH

and CBH would work. These papers examine models with two discrete environments, and a suggested

conservative bet-hedger (acting as a generalist coping moderately well with both environments) is never

able to outperform a diversified bet-hedger (producing specialists to each environment in the

proportions that they occur).

We present a different interpretation of CBH, which potentially allows for both CBH and DBH to coexist

within the same model. Considering a continuous trait with an asymmetric fitness function that

fluctuates between generations, CBH can be envisioned as having a cliff-edge effect in the same way as

insurance (see above). Organisms would thus be 'playing it safe' by shifting the mean trait value away

from the fitness function maximum, towards the less steeply decreasing side. In such a scenario, we

expect insurance to maximize arithmetic mean fitness within a generation. An additional shift in the

optimal trait value even further away from the cliff edge might then be selected for if it lowers fitness

variance between generations (despite lowering arithmetic mean fitness in a single generation). Such an

effect would essentially constitute a CBH strategy. Phenological features such as breeding date,

migration date or egg laying date for temperate birds are examples of traits with such asymmetric

fitness functions. The strength of selection may differ for the underlying selection pressures, for

example if being too late leads to lower offspring competitive ability, but being too early leads to a much

more severe mismatch with the food peak resulting in complete reproductive failure (Gienapp 2012).

Whether breeding after the peak in the fitness function represents insurance or CBH depends upon

whether the mismatch between the mean trait value and the peak of the asymmetric fitness function is

the result of individuals maximizing arithmetic mean fitness within their lifetime versus lineages being

favored that maximize geometric mean fitness over long time periods (see Lof et al. 2012). Despite

having much in common and some confusion between the terms in the literature, CBH has rarely been

placed in the same theoretical framework as insurance, and insurance has been all but absent as part of

the bet-hedging literature.

Here we investigate the relative importance of insurance and CBH in coping with stochastically

fluctuating environments within and between generations when the fitness function is skewed. We use

a single, continuous trait and calculate the means and variances in phenotype that maximize arithmetic

or geometric mean fitness under different magnitudes of fluctuations in the optimal trait value.

Mechanisms regulating the phenotypic variance expressed within a genotype, such as DBH increasing

such variance or phenotypic canalization decreasing it, are expected to interact with insurance and/or

CBH. Previous theoretical work has shown that DBH will adaptively increase variance in trait values once

the variance in the phenotypic optimum exceeds the squared width of the fitness function, whereas

smaller environmental variance favors the opposite mechanism, canalization of the trait towards the

value that maximizes fitness in the mean environment (Slatkin and Lande 1976; Bull 1987). Intuitively,

greater stochastic variation in trait values should require there to be more insurance or CBH modifying

the mean trait value, but these different components have not previously been placed in a common

framework. By using a skewed fitness function to illustrate the effects of insurance versus CBH, we are

able to examine these interactions, whilst modeling DBH alongside CBH in such a way allows us to

formally explore Starrfelt and Kokko's (2012) suggestion regarding an adaptive continuum between

these two potentially coexisting forms of bet-hedging.

Model description

The skew normal fitness function

A wide variety of fitness functions have been used to characterize asymmetrical relationships between phenotype and fitness (Martin and Huey 2008; Vasseur et al. 2014), but the results we demonstrate here can arise from any function with nonzero skew. We base our skewed fitness function on the density function of the skew normal distribution (O'Hagan and Leonard 1976), omitting the normalizing constant such that individual fitness takes a value of one for $z = \theta$. Fitness as a function of the phenotype z is then given by

$$w(z; \theta, \omega, \alpha) = 2e^{-\frac{(z-\theta)^2}{2\omega^2}} \Phi\left(\alpha \frac{z-\theta}{\omega}\right), \tag{1}$$

which is a Gaussian function multiplied by a term involving the cumulative distribution function Φ of the standard normal distribution. The parameter θ specifies the location, ω the width, and α the skew of the fitness function. $\alpha > 0$ gives a right-skewed function (positive skewness), $\alpha < 0$ a left-skewed function, and $\alpha = 0$ a symmetric Gaussian fitness function. Importantly, α also changes the position of the maximum value of w (despite location parameter θ being kept constant), so we will write θ_0 as the value of θ that provides maximum fitness for a trait value of zero. To examine the effects of skewed fitness functions on trait values we will use $\alpha > 0$ and $\theta = \theta_0$, so that adaptations in terms of phenotypic values shifted away from the fitness function maximum (due to insurance or CBH) become positive and easily interpretable relative to zero (i.e. the value of z simply becomes the distance from the peak, or the 'amount' of insurance or CBH).

Genotypic fitness when phenotypes vary within genotype

There may be variation in the phenotype z produced by a certain genotype k, due either to some (adaptive or non-adaptive) instability in individual development, or due to different individuals experiencing different microclimates. Following Bull (1987), we assume this variation to follow a normal distribution $f_k(z)$, with a mean μ_k and variance σ^2_k. We are interested in the joint evolution of the two underlying genotypic values μ_k and σ_k, and assume no genetic linkage or pleiotropic effects between them.

The mean fitness of all individuals with the genotype k (with genotypic values μ_k and σ^2_k) in any given environment θ (or a constant environment over time) then becomes

$$\overline{w}(\mu_k, \sigma_k | \theta, \omega, \alpha) = \int_{z=-\infty}^{\infty} w(z; \theta, \omega, \alpha) f_k(z) dz = \frac{2\omega}{\sqrt{\omega^2 + \sigma^2_k}} e^{\left(-\frac{(\mu_k - \theta)^2}{2(\omega^2 + \sigma^2_k)}\right)} \Phi\left(\frac{\omega\alpha}{\sqrt{\omega^2 + \sigma^2_k(1+\alpha^2)}} \frac{\mu_k - \theta}{\sqrt{\omega^2 + \sigma^2_k}}\right),$$

$\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad$ (2)

which is akin to eq. (3) in Bull (1987). The resulting function $\overline{w}$ has the same form as (1) but with a larger

width parameter and a smaller skew parameter. When $\sigma^2_k = 0$, the functions are identical. Choosing a

constant phenotypic variance $\sigma^2_k > 0$, we can use numerical optimization over μ_k and compare the

difference in maxima of w and $\overline{w}$, to find the amount of insurance needed to maximize arithmetic mean

fitness across all individuals of the genotype (i.e. the optimal shift in mean phenotype away from the

fitness function maximum). Fig. 1 shows w (black) together with $\overline{w}(k; \theta_0, 1, 5)|\sigma_k$ (reds) for increasing

values of σ_k. In this case, since the maximum of the fitness function is at zero, the optimal amount of

insurance is simply the value of the phenotype that gives the highest fitness, $\text{argmax}(\overline{w})$. The larger the

Figure 1: Asymmetric fitness functions and phenotypic variation produce insurance. Black line: the skew

normal fitness function with location $\theta = \theta_0$, width $\omega = 1$ and skew $\alpha = 5$. Red lines: arithmetic mean fitness

$\overline{w}(k; \theta_0, 1, 5)$ across individuals for genotypes with a fixed σ_k, from darkest to lightest, $\sigma_k = \{0.5, 1, 1.5, 2\}$.

Maxima of these functions are indicated by the dotted lines, and approximate to 0.002, 0.044, 0.153, 0.297

and 0.458, respectively.

phenotypic variance $\sigma^2{}_k$, the more insurance is needed to maximize genotype fitness. We also note that

genotype fitness strongly declines with increasing phenotypic variance – except when the mean

phenotype is far away from the fitness function peak.

Long-term fitness in a fluctuating environment

In a fluctuating environment it is not just individual fitness but the fitness of a genotype that will differ in

different environments, and long-term fitness in such cases is determined by geometric mean rather

than arithmetic mean fitness (Lewontin and Cohen 1969; Simons 2011). In the case of no fluctuations,

the geometric mean is simply equal to the arithmetic mean, and equation (2) is valid. If we let the

optimum position Θ follow a normal distribution f with a mean of Θ_0 and a variance σ_Θ across

generations, long-term arithmetic mean fitness can be found by taking the expectation of equation (2)

across the environmental fluctuations,

$$W_{arit} = \int_{\theta=-\infty}^{\infty} \bar{w}(\mu_k, \sigma_k | \theta, \omega, \alpha) f(\theta | \theta_0, \sigma_\theta)\, d\theta, \qquad (3a)$$

and for geometric mean fitness we can take the exponential of log fitnesses integrated across all

different environmental conditions,

$$W_{geom} = \exp\{\int_{\theta=-\infty}^{\infty} \ln[\bar{w}(\mu_k, \sigma_k | \theta, \omega, \alpha)]\, f(\theta | \theta_0, \sigma_\theta)\, d\theta\} \qquad (3b)$$

Maximizing both eq. 3a and 3b over μ_k and σ_k gives the strategy that provides the highest long-term

arithmetic or geometric mean fitness, respectively, for genotype k. Comparing them allows us to tease

apart bet-hedging effects (those maximizing geometric mean fitness at the expense of arithmetic mean

fitness) from non-bet-hedging effects (those maximizing only arithmetic mean fitness). Maximizing 3a

for a fixed σ_k gives the optimal amount of insurance (since the peak of the individual fitness function is

at zero), equal to argmax$_\mu$ $W_{arit}|\sigma_k$. Maximizing 3b for the same σ_k then reveals whether any additional

shift in mean phenotype can be attributed to bet-hedging rather than just insurance, with the optimal

amount of CBH then being equal to argmax$_\mu$[$W_{arit}|\sigma_k$] - argmax$_\mu$ [$W_{arit}|\sigma_k$].

Numerical integration was carried out in R version 3.3.1 (R Core Team 2016) and the code is available

upon request.

Results

Figures 2 and 3 show the long-term fitness of genotype k (consisting of the gene for mean phenotype, μ_k, and for variance in the phenotype, σ^2_k) measured in either arithmetic or geometric mean fitness in an environment with increasing fluctuations in the position parameter Θ of the individual fitness function (increasing σ_Θ). In Fig. 2 the individual fitness function is symmetrical and in Fig. 3 it is skewed. With a stable environment across generations (σ_Θ=0) the arithmetic and geometric mean fitness are equal (top and bottom panels the same). The fitness surface in these cases peaks at μ_k=0 and σ^2_k=0, that is, the

Figure 2: Fitness surfaces for a symmetrical fitness function for genotypes with different values of the mean phenotype μ_k and variation in phenotype σ_k. Contour lines show long-term arithmetic mean fitness (top row) and geometric mean fitness (bottom row). The position Θ of the fitness function fluctuates stochastically between generations, Θ~N(0, σ_Θ), the magnitude σ_Θ of environmental fluctuations increases successively (from 0 to 3) in the different columns from left to right. Irrespective of the scale of these fluctuations and the phenotypic variation (σ_k), fitness is always maximized by a peak in the contours in the middle of the x-axis, corresponding to a mean phenotypic value (μ_k) of zero, because the individual fitness function is a symmetrical normal distribution with a mean of 0 and width of ω=1.

optimal genotype is a trait value phenotypically canalized (i.e. with as little variation as possible) at the

peak of the individual fitness function. As environmental fluctuations increase (σ_θ>0), the differences

between arithmetic and geometric mean fitness increase. Notably, in Fig. 2 the peak geometric mean

fitness contours move upwards in the bottom panels as the environmental fluctuations increase. This

adaptive increase in phenotypic variation (σ_k) within the genotype represents DBH, and since arithmetic

mean fitness strictly declines with increasing σ_k (top panels), the necessary requirement that bet-

hedging involves a lowering of arithmetic mean fitness is fulfilled. In accordance with Bull's (1987) result,

this selection for increased phenotypic variation only appears once the environmental variance σ^2_θ is

Figure 3: Fitness surfaces for an asymmetrical fitness function for genotypes with different values of the mean phenotype μ_k and phenotypic variation σ_k. Contour lines show long-term arithmetic mean fitness (top row) and geometric mean fitness (bottom row). The position θ of the fitness function fluctuates stochastically between generations, θ~N(0, σ_θ), the magnitude σ_θ of the fluctuations increasing from left to right. In contrast to Fig. 2, fitness is always maximized here by a peak in the contours to the right of the middle of the x-axis (where μ_k=0), because the individual fitness function is a skew normal distribution with a mode of 0, width of ω=1 and skew of α=5.

larger than the squared width of the fitness function, and the optimal σ^2_k is then equal to $\sigma^2_\theta - \omega^2$. For

the symmetric Gaussian distribution (Fig. 2) this threshold is simply $\omega^2 = 1$, and DBH appears when $\sigma^2_\theta >$

1. In Fig. 3 asymmetry is introduced into the fitness function (results are shown for $\alpha = 5$, which matches

the fitness function in Fig. 1), but all other parameters remain as in Fig. 2. The variance of the fitness

function decreases as the skew increases, so this scenario also produces DBH (fitness surface peak with

$\sigma^2_k > 0$) under lower levels of environmental variance. The skew also leads to the fitness surface peaks

shifting to positive values of μ_k, away from the steeply decreasing side of the individual fitness function.

This shift is seen both in the top and bottom rows of Fig. 3, and in each case the shift occurs to the same

extent for the maximization of arithmetic and geometric mean fitness. Therefore, this represents only

insurance and there is no extra shift that can be attributed to a CBH effect if μ_k and σ_k are jointly

maximized.

However, geometric mean fitness is maximized for higher mean trait values μ_k than arithmetic mean

fitness when phenotypic variation σ_k is constrained at low values - see Figs 4 & 5a. Fig. 4 shows this

difference in mean phenotype (argmax$_\mu$ W_{geom} - argmax$_\mu$ W_{arit}), which is attributable to CBH (y-axis), for

different amounts of phenotypic variation σ_k (x-axis) and environmental variation σ_θ (line color). Note

that this CBH effect is only detectable when phenotypic variation σ_k is small, and environmental

fluctuations σ_θ are large enough that DBH would provide a much greater fitness gain (the steepest

incline on the fitness surface comes by moving upwards along the σ_k axis). The selection gradient will

thus adaptively increase the phenotypic variance, and not the mean phenotype *per se*.

For phenotypically canalized traits (i.e. traits that have experienced selection in σ_k towards zero), we can

infer that the fitness functions of these traits have fluctuated less across generations (smaller σ_θ), as we

otherwise would not see have seen this canalization. For these values of σ_θ, geometric and arithmetic

mean fitness peaks at more similar μ_k values when $\sigma_k = 0$ (fig. 4, bluer lines). As σ_k increases, mean

fitness $\overline{w}(k)$ becomes less skewed (see eq. 2, skew parameter α decreases with increasing σ_k) and

therefore CBH will not shift μ_k on top of any insurance already occurring. We note that if σ_k is

constrained to only exhibit a limited amount of DBH, then CBH and DBH will co-occur (e.g. the

dependence of optimal μ on σ in the lower right panel of fig. 3; redder lines showing CBH > 0 at $\sigma_k = 1$ in

fig. 4), but there are no cases in Figs. 3 and 4 where CBH co-occurs with optimal amounts of DBH. Once

environmental fluctuations have caused DBH ($\sigma_\theta > \omega$) and phenotypic variation σ_k increases to its new

optimum, geometric and arithmetic mean are maximized at the same mean phenotype, and the shift in

mean phenotype relative to the fitness function peak is then attributed to insurance rather than CBH.

Figure 4: Conservative bet-hedging (CBH), defined as the difference in the mean phenotype (μ_k) maximizing long-term arithmetic versus geometric mean genotype fitness, for given phenotypic standard deviation (σ_k). Lines represent results for different values environmental variation σ_Θ, as in Fig. 3. The individual fitness function w has a width of ω=1 and skew of α=5, also as in Fig. 3.

Figure 5b illustrates a similar case of an apparent bet-hedging effect being instead attributable to simply

maximizing arithmetic mean fitness, and thus not necessarily representing bet-hedging at all. In this

case, if the mean phenotype is constrained at a suboptimal value, such as may be the case in a climate

change scenario shifting the position of the fitness function, a positive amount of phenotypic variance is

adaptive (i.e. a risk-prone strategy due to risk sensitivity (Caraco et al. 1980; Stephens 1981), later

termed variance sensitivity - see Discussion). This result can be understood, like DBH, as the different

individuals of the genotype being phenotypically different (so that at least some are well adapted to the

current conditions) rather than everyone being possibly maladapted. However, there is no need here to

invoke a geometric mean (bet-hedging) argument because this diversification of phenotypes simply

maximizes arithmetic mean fitness across the individuals of the genotype. This type of adaptive

phenotypic variance is often attributed to bet-hedging without considering whether the trait specifically

increases geometric mean fitness at the cost of a decrease in arithmetic mean fitness (Mountford 1968).

Figure 5: Cross-sections of fitness surfaces in the third column of Fig. 3 (σ_Θ=1). The red line is the fitness surface for arithmetic mean fitness, the blue line is the fitness surface for geometric mean fitness. Dotted vertical lines show the maxima for the respective functions. A: a horizontal cross-section taken in the trait (μ_k) dimension at σ_k = 0, hence the individual fitness function (black line) peaks at zero, but fluctuates over time (black arrows), which causes mean fitness to be maximized at positive values of mean phenotype for both arithmetic and geometric mean. B: a vertical cross-section taken in the phenotypic variance (σ_k) dimension at μ_k = -2, and so diversifying bet-hedging is favored by geometric mean fitness, but arithmetic mean fitness also increases with phenotypic variance, because of Jensen's inequality (the individual fitness function is strongly convex at μ_k = -2) and hence what is known as a variance-prone strategy — see text for details.

Discussion

Among the various types of adaptive strategies to cope with environmental stochasticity, many have typically been considered from a within-individual perspective in the tradition of behavioral ecology. These use optimality theory to maximize some fitness proxy using the arithmetic mean across instances within a single generation, such as energy intake per time (Davies et al. 2012). An example is optimal foraging, a large body of the behavioral ecology literature that deals with such within-individual traits, including risk-sensitivity, state-dependent energy budgets, adaptive levels of energy reserves, and information sampling of foraging options (Stephens et al. 2007). Some rather different strategies have been considered to operate among individuals and over many generations. This long-term perspective has been in the tradition of evolutionary theory, which has identified key concepts such as phenotypic canalization and bet-hedging (Slatkin and Lande 1976; Bull 1987; Philippi and Seger 1989; Frank and Slatkin 1990). In an attempt to reconcile these contrasting views, we have calculated both long-term arithmetic and geometric mean fitnesses for combinations of trait means and variances under different levels of stochastic environmental fluctuations. By comparing the results obtained when maximizing arithmetic versus geometric mean fitness, we have illustrated some possible similarities between the two approaches, and the discrepancies that arise when considering the effects of either of these two measures of fitness in isolation.

Crucially, we use skewed fitness functions to demonstrate that shifting the mean phenotype away from the steeply decreasing side of the fitness function may provide a more useful and realistic case of conservative bet-hedging (CBH). This fulfills the definition of bet-hedging, in that it increases geometric mean fitness at a cost of lower arithmetic mean fitness (i.e. it provides lower fitness in the average environment, but also a lower variance in fitness across environments). Such a type of CBH has not been formally modelled previously, rather theoretical treatments of CBH have been limited to models with two discrete environments, where CBH has been envisioned as a canalized phenotype providing a compromise between the peaks of the fitness functions for the two environments (Starrfelt and Kokko 2012; Crowley et al. 2016). Such a CBH strategy is unsatisfactory since it always loses to a DBH strategy (producing specialists for the two environments) and is therefore not compatible with the concept of a 'continuum' between CBH and DBH and thus some sort of coexistence of the two strategies.

Various empirical studies of traits based on skewed fitness functions have invoked CBH arguments of the type we model here (e.g. Boyce and Perrins 1987; Simons and Johnston 2003). However, this shift is often also adaptive from an arithmetic mean fitness point of view in terms of an 'insurance' strategy

(Dall 2010). With a skewed fitness function, the cliff-edge effect entails that if individuals with the same

genotype differ stochastically in their phenotypes (or the fitness value of their phenotypes, e.g. due to

inhabiting different microenvironments), their average fitness is maximized if the mean phenotype is

shifted away from the peak of the fitness function, towards the less steeply decreasing side (Mountford

1968; Vercken et al. 2012; Mitteroecker et al. 2016). The same is also true for a single individual

experiencing uncertainty about its current state (i.e. regarding its phenotype or position on the x-axis on

the fitness function) or uncertainty about its current microenvironment (i.e. the position of the fitness

function on the x-axis relative to its phenotype). Its average fitness is therefore also maximized by

'playing it safe' and moving its phenotype away from the peak down the shallow slope of the skewed

fitness function. This shift (insurance) in the mean phenotype thus increases both arithmetic and

geometric mean fitness and is not simply a bet-hedging strategy. We initially hypothesized that CBH

(lowering the variance in expected fitness for each individual) might shift mean phenotype even more

away from the peak of the fitness function than insurance alone. However, our analysis shows that

maximizing geometric mean fitness generally does not require a different (i.e. a further shifted)

phenotype as compared to the phenotype that maximizes arithmetic mean fitness, but rather there is an

exact alignment of fitness interests for short-term and long-term strategies.

An exception to this is in cases where there is suboptimally little phenotypic variance, σ_k. The effect is

shown in Fig. 5A, where geometric mean fitness (blue line) is maximized for a higher phenotypic value

(the individual fitness function peaks at zero and has its steepest decline for negative values, see Fig. 1)

than arithmetic mean fitness (red line). This difference in optimum phenotypic values for canalized traits

stems solely from a fitness variance-reducing benefit and can thus be attributed to bet-hedging, and

perhaps CBH specifically. However, in this case the diversification bet-hedging (DBH) effect of increasing

phenotypic variance instead increases fitness much more than does any possible CBH effect shifting the

canalized phenotype to more positive values further down that shallow side of the fitness function. We

would therefore expect selection to favor this DBH mechanism to reduce fitness variance (Lande and

Arnold 1983), and again no additional CBH is expected to evolve beyond that already captured by the

effect of adaptive insurance.

We also conclude that, given the opportunity for insurance, there is very little scope for a single trait to

exhibit both DBH and CBH as an additional adaptation on top of any adaptive insurance already being

selected for. We do see DBH alongside insurance, but not CBH. An obvious reason for this is that in our

model we used the same pattern of environmental stochasticity at both the individual (short-term) and

genotypic (long-term) levels. Hence, any adaptive solution at the individual level that maximizes

arithmetic mean fitness also accounts for the same regime of environmental stochasticity experienced

at the genotype level, leaving nothing left for any geometric mean fitness effects to cope with. This

'alignment of fitness interests' maximizing both arithmetic and geometric mean fitnesses in the face of

the same regime of environmental fluctuations and skewed fitness functions at the two levels is

intriguing, but it does not promise to make empirical evidence for conservative bet-hedging any less

'elusive' (Childs et al. 2010; Simons 2011). Whilst the structure of environmental variation in our model

might reflect the general pattern expected of environmental stochasticity in nature, regimes of

environmental stochasticity may sometimes differ between the within lifetime variation at the individual

level versus the across lifetime variation at the genotype level (e.g. seasonal variation versus El Niño

events, as experienced by annual organisms). Therefore, understanding adaptations to environmental

stochasticity at different levels, such as insurance versus CBH, requires that we appreciate how patterns

of the stochasticity in question align and differ at the different levels of organismal experience.

Our result in this regard brings into focus the ecological relevance of previous work. For example, Lof et

al. (2012) used a stochastic dynamic model of timing of reproduction in great tits when the timing of the

food peak fluctuates between years. They assumed an asymmetric fitness function of laying date

relative to the food peak and showed that maximizing expected (arithmetic mean) fitness does indeed

produce an adaptive mismatch with the food peak, in the direction away from the steeply decreasing

side of the fitness function (an 'insurance' result). They acknowledge that "there might be additional

benefits of adaptive mismatch in terms of reductions in fitness variance" - i.e. maximizing geometric

rather than arithmetic mean fitness might yield a different result if fitness variance is lowest for a

different laying date than the observed outcome. However, their forward simulations (using the optimal

decision matrix from the dynamic model) "suggest that the variation in fitness often exhibited a

minimum close to the observed optimal laying dates" (Lof et al. 2012). Hence, there is little scope for

CBH to shift the optimal laying date any further away from the cliff edge. However, geometric mean

fitness benefits resulting from a reduction in fitness variance across generations can provide an added

selection pressure towards the same optimum phenotypic values as insurance. This result appears in our

model in that the geometric mean fitness surfaces are more peaked around the maxima than the

arithmetic mean fitness surfaces (Figs 2 & 3) - i.e. the selection pressure toward the same insurance

optimum becomes stronger due to also producing lowest fitness variance (i.e. the CBH effect) across

generations at this same optimum. All of which may provide reason for optimism with regards to species

survival in a period when human-induced environmental change may produce sudden increases in

environmental stochasticity that are too rapid for effective evolutionary responses (Barrett and Hendry

2012; Nadeau et al. 2017). This is because in the case of asymmetric fitness functions then any currently

adaptive insurance strategy will already have selected for the appropriate phenotype, and no extra

evolutionary conservative bet-hedging (CBH) response will be needed in terms of additional changes to

the mean phenotype.

The importance of environmental stochasticity at the individual versus genotypic level is highlighted in a

large body of previous work on bet-hedging (Levins 1962; Cohen 1966; Gillespie 1974). These identify

the 'grain' of the environmental variation as a strong determinant of whether bet-hedging strategies will

evolve (Starrfelt and Kokko 2012; Crowley et al. 2016). If individuals in a population experience very

different environments within a generation (i.e. the environment is 'fine-grained' or 'locally variable'),

the fitness correlations between individuals of the same genotype will be low and the scope for DBH is

reduced. Assuming a continuous distribution of environmental fluctuations, such as in our model, a very

fine-grained environment also implies that a larger proportion of the total environmental variation is

experienced by individuals of the same genotype within each generation. The between-generation

fluctuations in mean environment therefore become smaller, and a smaller proportion of the variance in

fitness is experienced at the genotype level. Only when the environmental conditions are common to a

large proportion of the population every generation (i.e. a 'coarse-grained' environment, featuring

'global variation') is there a large variance in genotype fitness between generations, which can be

ameliorated by adaptive bet-hedging strategies.

While environmental 'grain' is not explicitly modeled here, it is interesting to consider the σ_k gene as any

trait that interacts with the grain of the environment. DBH traits, such as offspring dispersal and

variation in dormancy duration (i.e. dispersal in time and space), in effect respond to and modify the

grain of the environment as it is experienced. Hence, adaptive dispersal (in time or space) leads such

genotypes to experience a more fine-grained environment (Gourbière and Menu 2009; Scheiner 2014).

We therefore see why our model suggests that the possibility of DBH makes additional CBH on top of

insurance unnecessary – the correlation in fitness between individuals of the same genotype can always

evolve to be low enough (via DBH) such that arithmetic mean fitness is a good determinant of long-term

fitness.

An early model that hinted at this point involved three distinct adaptations for 'reducing risk in variable

environments' in seed production in desert plants: seed size, dispersal and dormancy (Venable and

Brown 1988). Seed size increases survival in bad environments, but does not affect survival in good

environments. Given the trade-off between seed number and size, small seeds are optimal in good

environments, but large seeds have a lower variance in expected fitness per individual. Whether

increasing seed size represents insurance or CBH depends on the proportion of the environmental

variation that is experienced by the genotype within versus between each generation (i.e. the grain of

the environment). With no dispersal or dormancy, the environment is coarse-grained, fitness is purely a

multiplicative process and increased seed size is clearly a bet-hedging trait increasing geometric mean

fitness at the cost of a lower arithmetic mean fitness. But increased seed size may also maximize

arithmetic mean fitness across fine-grained environments, which is the appropriate fitness measure if

the genotype is sufficiently spread in space (or time, in this case) to experience the full range of

environmental variation in each generation (Levins 1962). Thus, dispersal and dormancy are not only

diversifying bet-hedging (DBH) traits, but also determine the grain of the environment and the need for

conservative bet-hedging (CBH) versus insurance. Venable and Brown (1988) show that these

adaptations to reduce risk are essentially substitutable and that a decrease in the value of either of the

traits away from the optimum leads to evolutionary compensation in an increased value of the other

traits. However, note that the DBH traits here, dispersal and dormancy, are inherently different from a

conceivable fourth risk-reducing strategy: variation in seed size itself. In discrete environments ('good'

and 'bad') the optimal DBH strategy maximizing geometric mean fitness is to produce seeds with

optimal size for each of the environments (small and large) with the probabilities of those respective

environments occurring, while in continuously varying environment seed size should vary around a

mean trait value, as in our model. Here we showed that in a continuously varying environment, this type

of DBH (which Venable and Brown (1988) do not explore) generally provides a greater benefit than CBH.

However, we do not rule out that with other types of environmental variation and environment-specific

fitness functions then changing the mean trait value may be a better strategy. We also point out again

that both of these risk-reducing strategies are only effective once the grain of the environment causes

selection to maximize geometric rather than arithmetic mean fitness (Venable and Brown 1988;

Scheiner 2014).

In our current model, the only time arithmetic mean fitness would be higher with more phenotypic

variance (i.e. for an individual, rather than a DBH genotype increasing among-individual variation) would

be if its fitness function were strongly convex around its current phenotype (Fig. 5b), such as at the tails

of a Gaussian fitness function. This is the same adaptive gambling effect that produces risk-sensitivity

(Caraco et al. 1980; Stephens 1981), aka variance sensitivity (Smallwood 1996; Stephens et al. 2007),

which is an important concept from economics used to explain foraging decisions and other behaviors

when there are more or less variable options. The fitness advantage of variance sensitivity follows

directly from Jensen's inequality: if the fitness function f of some utilized resource or trait x is convex,

then the mean fitness gained over a sequence of events with variable reward x will be larger than the

fitness gained from the mean reward x: $E[f(x)] > f(E[x])$. This is an arithmetic mean fitness maximizing

argument, and the benefit of increasing phenotypic variance can therefore be seen in our calculations of

arithmetic mean fitness as well (top rows of Fig. 2 & 3; Fig. 5b). Essentially, for a constant μ_k sufficiently

far from the fitness function peak, arithmetic mean fitness is maximized at an intermediate value $\sigma_k > 0$.

This similarity between variance-sensitivity and DBH in producing variable phenotypes but at different

adaptive timescales has not been reported before, and it is made explicit here through our comparison

of trait means and variances maximizing long-term arithmetic or geometric mean fitness.

We have demonstrated several results linking theory concerning individual-level strategies from

behavioral ecology with genotype-level adaptations from evolutionary biology in context of

environmental uncertainty. The next step is now to connect these theoretical studies to real world

examples and quantitative studies of organisms in the lab and in the wild, especially if we are to

understand how populations might respond to current human-induced rapid environmental change. In

order to apply our genotype-level view to predictive statements concerning evolutionary responses we

would require extensive data on past climate fluctuations, clear links between trait values and individual

fitness, as well as detailed knowledge of the genetic mechanisms underlying the traits. While this might

seem an insurmountable task, the empirical evidence for bet-hedging in the wild has shown that long-

term studies on natural populations can provide answers to these types of questions (Simons 2011). We

hope that our results here act as a motivation to both empirical and theoretical studies on adaptations

to stochastic environments to compare and contrast individual versus genotype perspectives and the

alternative adaptive currencies of arithmetic versus geometric mean fitness.

Acknowledgements

443   This work was supported by the Research Council of Norway through its Young Talented Researchers

444   funding scheme FRIMEDBIO, project number 240008 to I.I.R., and partly through its Centres of

445   Excellence funding scheme, project number 223257, to Centre for Biodiversity Dynamics at the

446   Norwegian University of Science and Technology.

Author contributions

IIR, JW and JT conceived of the ideas and all authors contributed to the development of the ideas. JT and TRH wrote the code and did the calculations. TRH wrote the manuscript with contributions from all authors. All authors gave final approval for publication.

Code accessibility

The code is available upon request, and will be uploaded to Dryad upon publication.

References

References

Andersson, M., and Y. Iwasa. 1996. Sexual selection. Trends Ecol. Evol. 11:53–58.

Angilletta, M. J. 2009. Thermal adaptation: A theoretical and empirical synthesis. 1st ed. Oxford
University Press, Oxford.

Barrett, R. D. H., and A. P. Hendry. 2012. Evolutionary rescue under environmental change? Pp. 216–233
in U. Candolin and B. B. M. Wong, eds. Behavioural Responses to a Changing World. Oxford
University Press, Oxford.

Bednekoff, P. A., H. Biebach, and J. Krebs. 1994. Great tit fat reserves under unpredictable
temperatures. J. Avian Biol. 25:156–160.

Boyce, M. S., and C. M. Perrins. 1987. Optimizing great tit clutch cize in a fluctuating environment.
Ecology 68:142–153.

Brodin, A. 2007. Theoretical models of adaptive energy management in small wintering birds. Philos.
Trans. R. Soc. B Biol. Sci. 362:1857–1871.

Bull, J. J. 1987. Evolution of phenotypic variance. Evolution. 41:303–315.

Caraco, T., S. Martindale, and T. S. Whittam. 1980. An empirical demonstration of risk-sensitive foraging
preferences. Anim. Behav. 28:820–830.

Childs, D. Z., C. J. E. Metcalf, and M. Rees. 2010. Evolutionary bet-hedging in the real world: empirical
evidence and challenges revealed by plants. Proc. R. Soc. B Biol. Sci. 277:3055–3064.

Cohen, D. 1966. Optimizing reproduction in a randomly varying environment. J. Theor. Biol. 12:119–129.

Crowley, P. H., S. M. Ehlman, E. Korn, and A. Sih. 2016. Dealing with stochastic environmental variation
in space and time: bet hedging by generalist, specialist, and diversified strategies. Theor. Ecol.
9:149–161.

Dall, S. R. X. 2010. Managing risk: The perils of uncertainty. Pp. 194–206 *in* D. F. Westneat and C. W. Fox,
eds. Evolutionary Behavioral Ecology. Oxford University Press, New York.

Davies, N. B., J. R. Krebs, and S. A. West. 2012. An Introduction to Behavioural Ecology. Fourth edi.
Oxford University Press, Oxford.

Devaux, C., and R. Lande. 2010. Selection on variance in flowering time within and among individuals.
Evolution. 64:1311–1320.

Frank, S. A., and M. Slatkin. 1990. Evolution in a variable environment. Am. Nat. 136:244–260.

Gienapp, P. 2012. Migration. Pp. 80–92 *in* U. Candolin and B. Wong, eds. Behavioural Responses to a
Changing World. Oxford University Press, Oxford.

Gillespie, J. H. 1974. Natural selection for within-generation variance in offspring number. Genetics
76:601–6.

Gourbière, S., and F. Menu. 2009. Adaptive dynamics of dormancy duration variability: Evolutionary
trade-off and priority effect lead to suboptimal adaptation. Evolution. 63:1879–1892.

Graham, J. K., M. L. Smith, and A. M. Simons. 2014. Experimental evolution of bet hedging under
manipulated environmental uncertainty in *Neurospora crassa*. Proc. R. Soc. B-Biological Sci. 281.

Krams, I., D. Cirule, V. Suraka, T. Krama, M. J. Rantala, and G. Ramey. 2010. Fattening strategies of
wintering great tits support the optimal body mass hypothesis under conditions of extremely low
ambient temperature. Funct. Ecol. 24:172–177.

Lande, R., and S. J. Arnold. 1983. The measurement of selection on correlated characters. Evolution.
37:1210–1226.

Levins, R. 1962. Theory of fitness in a heterogeneous environment . I . The fitness set and adaptive
function. Am. Nat. 96:361–373.

Lewontin, R. C., and D. Cohen. 1969. On population growth in a randomly varying environment. Proc.
Natl. Acad. Sci. 62:1056–1060.

Lof, M. E., T. E. Reed, J. M. McNamara, and M. E. Visser. 2012. Timing in a fluctuating environment:
environmental variability and asymmetric fitness curves can lead to adaptively mismatched avian
reproduction. Proc. R. Soc. B Biol. Sci. 279:3161–3169.

Martin, T. L., and R. B. Huey. 2008. Why "suboptimal" is optimal: Jensen's inequality and ectotherm
thermal preferences. Am. Nat. 171:E102-18.

McNamara, J. M. 1998. Phenotypic plasticity in fluctuating environments: consequences of the lack of
individual optimization. Behav. Ecol. 9:642–648.

Mitteroecker, P., S. M. Huttegger, B. Fischer, and M. Pavlicev. 2016. Cliff-edge model of obstetric
selection in humans. Proc. Natl. Acad. Sci. 113:201612410.

Mountford, M. D. 1968. The significance of litter-size. J. Anim. Ecol. 37:363–367.

Nadeau, C. P., M. C. Urban, and J. R. Bridle. 2017. Climates past, present, and yet-to-come shape climate
change vulnerabilities. Trends Ecol. Evol. 32:786–800. Elsevier Ltd.

O'Hagan, A., and T. Leonard. 1976. Bayes estimation subject to uncertainty about parameter constraints.
Biometrika 63:201–203.

Parker, G. A., and J. M. Smith. 1990. Optimality theory in evolutionary biology. Nature 348:27–33.

Philippi, T., and J. Seger. 1989. Hedging one's evolutionary bets, revisited. Trends Ecol. Evol. 4:41–44.

R Core Team. 2016. R: A language and environment for statistical computing. R Foundation for Statistical
Computing, Vienna, Austria.

Ratikainen, I. I., and J. Wright. 2013. Adaptive management of body mass by Siberian jays. Anim. Behav.
85:427–434. Elsevier Ltd.

Scheiner, S. M. 2014. Bet-hedging as a complex interaction among developmental instability,
environmental heterogeneity, dispersal, and life-history strategy. Ecol. Evol. 4:505–515.

Simons, A. M. 2009. Fluctuating natural selection accounts for the evolution of diversification bet
hedging. Proc. R. Soc. B Biol. Sci. 276:1987–1992.

Simons, A. M. 2011. Modes of response to environmental change and the elusive empirical evidence for
bet hedging. Proc. R. Soc. B Biol. Sci. 278:1601–1609.

Simons, A. M., and M. O. Johnston. 2003. Suboptimal timing of reproduction in *Lobelia inflata* may be a
conservative bet-hedging strategy. J. Evol. Biol. 16:233–243.

Slatkin, M., and R. Lande. 1976. Niche width in a fluctuating environment - density independent model.
Am. Nat. 110:31–55.

Smallwood, P. D. 1996. An introduction to risk sensitivity: The use of Jensen's inequality to clarify
evolutionary arguments of adaptation and constraint. Am. Zool. 36:392–401.

Starrfelt, J., and H. Kokko. 2012. Bet-hedging - a triple trade-off between means, variances and

correlations. Biol. Rev. 87:742–755.

Stephens, D. W. 1981. The logic of risk-sensitive foraging preferences. Anim. Behav. 29:628–629.

Stephens, D. W., J. S. Brown, and R. C. Ydenberg (eds). 2007. Foraging: Behavior and ecology. 1st ed.
University of Chicago Press.

Urban, M. C., R. Bürger, and D. I. Bolnick. 2013. Asymmetric selection and the evolution of extraordinary
defences. Nat. Commun. 4.

Vasseur, D. A., J. P. Delong, B. Gilbert, H. S. Greig, C. D. G. Harley, K. S. Mccann, V. Savage, T. D. Tunney,
M. I. O. Connor, D. A. Vasseur, J. P. Delong, B. Gilbert, H. S. Greig, C. D. G. Harley, K. S. Mccann, V.
Savage, T. D. Tunney, and M. I. O. Connor. 2014. Increased temperature variation poses a greater
risk to species than climate warming. Proc. R. Soc. B Biol. Sci. 281.

Venable, D. L. 2007. Bet Hedging in a Guild of Desert Annuals. Ecology 88:1086–1090.

Venable, D. L., and J. S. Brown. 1988. The selective interactions of dispersal, dormancy, and seed size as
adaptations for reducing risk in variable environments. Am. Nat. 131:360–384.

Vercken, E., M. Wellenreuther, E. I. Svensson, and B. Mauroy. 2012. Don't fall off the adaptation cliff:
When asymmetrical fitness selects for suboptimal traits. PLoS One 7:e34889.

Yoshimura, J., and W. M. Shields. 1987. Probabilistic optimization of phenotype distributions: a general
solution for the effects of uncertainty on natural selection? Evol. Ecol. 1:125–138.

Zhang, X. S., and W. G. Hill. 2005. Evolution of the environmental component of the phenotypic
variance: Stabilizing selection in changing environments and the cost of homogeneity. Evolution.
59:1237–1244.

Standard deviation of phenotypic optimum, σ_θ

Mean phenotype (μ_k)