Title
Local adaptation of a dominant coastal tree to freshwater availability and solar radiation
suggested by genomic and ecophysiological approaches

Authors
Mariana Vargas Cruz[1,2], Gustavo Maruyama Mori[3], Caroline Signori Müller[1], Carla Cristina
da Silva[2], Dong-Ha Oh[4], Maheshi Dassanayake[4], Maria Imaculada Zucchi[5], Rafael Silva
Oliveira[1] and Anete Pereira de Souza[1,2]*
8
[1]Department of Plant Biology, Institute of Biology, University of Campinas (Unicamp),
Campinas, SP 13083-863, Brazil.
[2]Center for Molecular Biology and Genetic Engineering, University of Campinas (Unicamp),
Campinas, SP 13083-875
[3]Institute of Biosciences, São Paulo State University (Unesp), São Vicente, SP 11330-900,
Brazil.
[4]Department of Biological Sciences, Louisiana State University (LSU), Louisiana, LA 70803,
United States.
[5]São Paulo Agency for Agribusiness Technology (APTA), Piracicaba, SP 13400-790, Brazil.
*Author for correspondence. Tel: +55 19 3521 1132 Email: anete@unicamp.br

19
20
Word count for the main text: 4,573
Abstract: 199
Introduction: 622
Materials and Methods: 1,843
Results: 1,710
Discussion: 1,702
Conclusions: 264
Acknowledgements: 190
Author Contributions: 89
Figures: 6 figures, all in colour
Tables: 1 table
Supplementary information: Supplementary Figures S1-S7 (all in colour, except for S5);
Supplementary Tables S1-S13; Supplementary Methods; Supplementary Note,
Supplementary Data File.

Abstract

Local adaptation is often a product of environmental variations in the geographical space and
has implications for biodiversity conservation. We investigated the role of latitudinal
heterogeneity in climate on the organization of genetic and phenotypic variation in the
dominant coastal tree, *Avicennia schaueriana*. In a common garden experiment, samples
from an equatorial region, marked by rainy/dry seasons, accumulated less biomass, showed
lower stomatal conductance and transpiration, narrower xylem vessels, smaller leaves and
higher reflectance of long wavelengths (red light) on the stem epidermis, than samples from a
subtropical region, marked by warm/cold seasons. Transcriptome differences identified
between trees sampled under field conditions at equatorial and subtropical sites, were
enriched in functional categories as responses to temperature, solar radiation, water deficit,
photosynthesis and cell wall biosynthesis. The diversity based on thousands of SNP loci
revealed a north-south genetic structure. Remarkably, signatures of selection were identified
in loci associated with photosynthesis, anthocyanin accumulation and the responses to
osmotic and hypoxia stresses. Our results suggest the existence of divergence in key
resource-use characteristics, likely driven by climate seasonality, based on water-deficit and
solar radiation. These findings provide a basis for conservation plans and for predictions for
coastal plant responses to climate change.

Keywords
Avicennia schaueriana (Black Mangrove); climate change; coastal ecosystems; comparative
transcriptomics; genome scans for selection; local adaptation; plant ecophysiology; resource-
use.

Introduction

Adaptation is frequently a consequence of spatial variations in selective environmental forces acting on phenotypic diversity[1,2]. As selective forces operate, they may reduce the heritable variation within a population, leading to the specialization of individuals[2]. Conversely, in highly stochastic environments, selection can increase the species potential for phenotypic plasticity[3]. Therefore, the investigation of adaptation is fundamental for understanding the ability of a species to respond to environmental changes[4,5].

Alarmingly high rates of environmental changes have been observed over the last decades, particularly affecting coastal ecosystems[6–9], for instance, bleaching of coral reefs worldwide[10], large-scale dieback of mangrove communities[11] and increased mortality of seagrass beds[12]. Further environmental changes and coastal biodiversity loss threaten critical socioeconomic and ecological services provided by coastal ecosystems, including their pivotal role as blue carbon stocks[13]. In this context, studies on the organization of adaptive variation of coastal habitat-forming species are necessary to improve predictions on the impacts of future climate conditions on marine ecosystems[14] and to support efficient conservation practices[15].

Coastal habitat-forming species are frequently water-dispersed over long distances[16,17] and often occur across wide latitudinal ranges. The connectivity among populations of these species may be influenced, not only by their dispersion capabilities but also by means of natural selection, caused by varying establishing success over the broad environmental heterogeneity across latitudes[18]. For instance, distinct coastal species in the North Atlantic show a north-south organisation of diversity with evidences of selection for different thermal regimes[19–21]. Similarly, in the Southwest Atlantic, an overlapping north-south structure of the genetic diversity, with reduced gene flow between northern and southern populations, has been observed in phylogenetically distant habitat-forming species[22–24]. In this context, one could expect northerly and southerly populations of these species to adapt differently to contrasting environments over their latitudinal ranges, especially due to their large population sizes[2,25]. However, the neutrality of the molecular markers used in previous studies has precluded inferences regarding adaptive variation. The existence of local adaptation remains virtually unknown in species that play a central role in sustaining the coastal biodiversity in the South Atlantic. This limited knowledge about the organization of non-neutral variation compromises accurate predictions and suitable conservation efforts for sustainable management.

Here, we tested the hypothesis that latitudinal heterogeneity in climate variables shape the variation of genotypes and phenotypes involved in the optimisation of resource-use in widely distributed coastal species. To reduce the potential for incorrect conclusions about selection[26], we integrated three independent, but complementary approaches, making predictions as follows: (1) using a common garden experiment, individuals from contrasting latitudes would show genetically-based phenotypic divergence in ecophysiological traits; (2) under contrasting latitudes, transcriptomic changes would be detected in genes involved in responses to environmental variation; and (3) signatures of selection along the species distribution would be detected in genes involved in responses to latitudinal variation in air temperature, solar radiation and freshwater availability determined by air water vapour pressure deficit (VPD), rainfall and tidal regime. We considered the new-world coastal tree species, *Avicennia schaueriana* Stapf & Leechman ex Moldenke, found in the Lower Lesser Antilles and widespread from Venezuela to the southernmost temperate mangroves on the Atlantic coast of South America (~28 °S)[27] as a model to test this hypothesis. The broad latitudinal range of *A. schaueriana*, spanning wide environmental variation (Fig. 1), and the previously detected north-south structure of neutral variation[22], facilitate ecological predictions and the accumulation of divergence, motivating this choice. Our results indicated

that fluctuations in VPD, rainfall and solar radiation may be associated with the observed
phenotypic and genotypic variation as well as the regulation of gene expression in this
dominant coastal tree. We discuss the implications of our results for the persistence of coastal
biodiversity in the context of climate change.

Materials and Methods
Propagule sampling
Mature propagules were collected from 15 *Avicennia schaueriana* Stapf & Leechman
ex Moldenke mother trees that were at least 100 m apart from one another, at the following
two genetically contrasting populations[22]: (1) the southernmost range limit of American
mangroves, in the subtropical region, and (2) an equatorial site in one of the world's largest
macrotidal mangrove forests[28,29], near the northernmost limit of the species range (Fig. 1).
We refer to samples collected in the former site as "subtropical" and those in the latter as
"equatorial" throughout this work. Sampling in more localities was impracticable due to the
absence of mature propagules during fieldwork. A detailed characterisation of each of these
contrasting sites can be found in Table 1, in the Supplementary Methods and in the
Supplementary information Figure S1.
123
Comparative ecophysiology using a common garden experiment
Propagules were germinated as described for *Avicennia germinans*[30]. We grew
propagules in trays with local mangrove soil for two months. After this period, 44 similar-
sized seedlings from 30 distinct mother trees, 15 from equatorial and 15 from subtropical
sites — with an average height of 18 cm, most with three leaf pairs and senesced cotyledons
— were transplanted to 6 L pots filled with topsoil and sand (1:1). Seedlings were cultivated
for seven months under homogenous conditions in a glasshouse at the University of
Campinas, São Paulo, Brazil (22°49' S 47°04' W), where automatic sensors coupled with a
data logger (Onset Computer Corp.) measured the atmospheric humidity and temperature
every 15 minutes. Seedlings were irrigated daily at 10 a.m. and 5 p.m. with a 3-minute
freshwater spray. Twice a week, nutrients were added to the soil using 600 mL of 0.4X
Hoagland solution with 15.0 g L^{-1} NaCl per pot. Pots were rotated weekly to reduce the
effects of environmental heterogeneity. The environmental conditions in the glasshouse were
different from those at each sampling site, as is shown in the Supplementary information Fig.
S2.
The light reflectance of stems was measured in ten plants from each sampling site
using a USB4000 spectrophotometer (OceanOptics, Inc.) coupled to a deuterium-halogen
light source (DH-2000; OceanOptics) using a light emission range from 200-900 nm.
Photosynthesis, stomatal conductance and transpiration rates were measured every 2.0-2.5
hours in five six-month-old individuals from each sampling site on two different days using a
Li-Cor 6400 XT (Li-Cor Corp.).
After harvest, three plants without flowers or flower buds from each sampling site
were split into leaves, stems and roots, washed with distilled water, dried for 7 days at 70 °C
and weighed. The individual leaf area, total leaf area and leaf lamina angle per plant were
measured through photographic analyses using ImageJ[31]. The specific leaf area (SLA, cm^2
leaf area kg^{-1} leaf dry biomass) was also calculated for these samples. Stems were fixed in
FAA (Formaldehyde Alcohol Acetic acid), stored in 70% alcohol for wood anatomy analysis
and cut into 30-μm thick transverse sections. Sections were stained with a mixture of 1%
Astra Blue and 50% alcohol (1:1) followed by 1% Safranin O. Micrographs were taken using
an Olympus BX51 microscope coupled to an Olympus DP71 camera (Olympus Corp.). The
following wood traits were quantified using ImageJ and R v.4.0.0: vessel lumen area (A),

vessel density in xylem (number of vessels/xylem area), vessel grouping index (mean number
of vessels per vessel grouping) and vessel lumen area in sapwood (vessel lumen
area/sapwood area). The vessel arithmetic diameter (D) was estimated according to Scholz et
al.[32].

Shapiro-Wilk's normality tests and Fisher's F-tests of equality of variances were
performed in R v.4.0.0 to determine the suitability of group comparison statistical tests. In
comparisons between equatorial and subtropical samples, we used the unpaired Student's T-
tests or, alternatively, Mann-Whitney-Wilcoxon tests. Multiple-group comparisons were
conducted for the analysis of environmental conditions in the field vs. in the glasshouse using
one-way analysis of variance (ANOVA) with Tukey's post hoc honest significant difference
(HSD) tests. A significance level of 0.05 was used as the alpha for all tests. The Bonferroni
adjustment of P-values was conducted for multiple comparisons.

Plant material for RNA extraction and RNA-sequencing
Plant material used for RNA-sequencing (RNA-Seq) was collected from the
equatorial and subtropical sites described in the "Propagule sampling" section and
immediately stored in RNAlater (Ambion, Inc.). Because precise species identification
requires the analysis of both vegetative branches and flower morphology, opportunities of
sampling were limited by the reproductive phenology at each visited site. Leaves, stems and
flowers from three adult trees at least 100 m apart were collected from July-August of 2014,
which corresponds to the end of winter at the subtropical site and the beginning of the dry
season at the equatorial site. To minimize the detection of differential transcripts expression
due to circadian changes, sampling was conducted during low tide and from 10:30 AM to
4:00 PM. A detailed description of the environmental conditions at the time of sampling is
available in the Supplementary information Table S1.
We extracted RNA according to Oliveira et al.[33] and evaluated its integrity and purity
using agarose gel electrophoresis and a NanoVue spectrophotometer (GE Healthcare Life
Sciences). lllumina TruSeq RNA Sample Preparation kits (Illumina, Inc.) were used to
construct cDNA libraries. cDNA quality was assessed using the Agilent 2100 Bioanalyzer
(Agilent Technologies) and concentrations were quantified by qPCR using the Sequencing
Library qPCR Quantification kit (Illumina, Inc.). Sequencing was performed using two 36-
cycle TruSeq SBS paired-end kits (Illumina, Inc.) and the Genome Analyzer IIx platform
(Illumina, Inc.).

Assembly and characterisation of the <u>A. schaueriana</u> transcriptome
Adapter sequences were trimmed, and 72-bp paired-end reads were filtered by quality
(Phred score ≥ 20 for at least 70% of the read length) using the NGS QC Toolkit v.2.3[34].
High-quality reads were used for transcriptome assembly in the CLC Genomics Workbench
(https://www.qiagenbioinformatics.com/). We used the default settings, except for the
distance between pairs (300-500 bp) and k-mer size (45 bp).
Reads were mapped to transcripts using bowtie1[35] in single-read mode using the
default parameters. Transcripts without read-mapping support were removed. Functional
annotation was performed using blastx v.2.2.31[36] with an e-value < 1^{-10}. NCBI RefSeq[37], The
Arabidopsis Information Resource (TAIR)[38] and NCBI non-redundant (nr) databases were
used as reference databases. We excluded transcripts that were exclusively similar to non-
plant sequences. Protein family domains were identified using HMMER3[39], which iteratively
searched transcripts against the Pfam database. To assign Gene Ontology (GO) terms to
transcripts, we used the *Arabidopsis thaliana* gene association file from the GO Consortium[40]
and retrieved the information for transcripts showing similar coding sequences in the *A.*
thaliana genome. Redundant transcripts were clustered with Cd-hit-est v.4.6.1[41] using local

alignment mode with 95% identity and 70% coverage of the shortest sequence thresholds.
Open reading frames (ORFs) were identified using Transdecoder (http://transdecoder.sf.net).
We reduced the redundancy of transcripts in the final assembly by retaining for each CD-
HIT-EST cluster either the sequence with the longest ORF or, in the absence of sequences
containing ORF, the longest sequence.
The completeness of the final transcriptome was assessed using BUSCO[42].
Additionally, a reciprocal blastn alignment using an e-value threshold of 10^{-10} and a
minimum alignment length of 100 nucleotides with at least 70% identity was used to
compare the *A. schaueriana* transcriptome with other publicly available transcriptomes of
congeneric species.
215
Comparative transcriptomics using RNA-sequencing
Tissue-specific count data were obtained from the number of reads uniquely mapped
to each transcript of the non-redundant transcriptome using bowtie1[35] and normalised using
edgeR[43]. Differentially expressed transcripts (DETs) between equatorial and subtropical tree
tissue-specific samples were detected using the exact test for negative binomial distributions
with an FDR < 0.05 (Benjamini-Hochberg). GO term enrichment analyses of the DETs were
performed using GOseq[44] with the Wallenius approximation method and P-value < 0.05.
Differential expression results were verified using reverse transcription real-time PCR (qRT-
PCR) (Supplementary Methods).
225
Detection of candidate adaptive loci
We sampled leaves from 79 adult plants at ten locations spanning most of the
distribution of *A. schaueriana* Stapf & Leechman ex Moldenke (Fig. 1, Supplementary
information Table S2). We isolated DNA using the DNeasy Plant Mini Kit (QIAGEN) and
NucleoSpin Plant II (Macherey Nagel) following the manufacturers' instructions. DNA
quality and quantity were assessed using 1% agarose electrophoresis and the QuantiFluor
dsDNA System with a Quantus fluorometer (Promega). Nextera-tagmented reductively-
amplified DNA (nextRAD) libraries[45] were prepared and sequenced by SNPsaurus
(SNPsaurus) on a HiSeq 2500 (Illumina, Inc.) with 100-bp single-end chemistry. Genomic
DNA fragmentation and short adapter ligation were performed with the Nextera reagent
(Illumina, Inc.) followed by amplification with one of the primers matching the adapter and
extending nine arbitrary nucleotides into the genomic DNA. Assembly, mapping and single
nucleotide polymorphic loci (SNP) identification were performed using custom scripts
(SNPsaurus), which created a reference catalogue of abundant reads across the combined
sample set and mapped reads to this reference, allowing two mismatches and retaining
biallelic loci present in at least 10% of the samples. We further filtered markers by allowing
no more than 65% of missing data, Phred score > 30, 8x minimum coverage, only one SNP
per locus and a minor allele frequency ≥ 0.05 using vcftools v.0.1.12b[46]. To reduce paralogy
or low-quality genotype calls, we used a maximum read coverage of 56 (the average read
depth times 1.5 standard deviation).
After excluding plants morphologically identified as *A. schaueriana* with genomic
signs of hybridisation with *A. germinans* (L.) L., we assessed the genetic structure
considering all SNPs, using the discriminant analysis of principal components (DAPC)[47] and
ADMIXTURE v.1.3.0[48]. For DAPC analyses, we considered the number of groups (K)
varying from 1-50 and the Bayesian information criteria for inferring K. Additionally, we
used the optim.a.score function to avoid overfitting during the discrimination steps. For the
ADMIXTURE analyses, we performed three separate runs for K varying from 1-15 using the
block-relaxation method for point estimation; computing was terminated when estimates
increased by < 0.0001, and the most likely K was determined by cross-validation.

We used two programs to minimise false-positive signs of natural selection:
LOSITAN[49], assuming an infinite allele model of mutation, using a confidence interval of
0.99, a false-discovery rate (FDR) of 0.1, the neutral mean FST and forcing the mean FST
options; and pcadapt 3.0.4[50], which simultaneously identifies the population structure and the
loci excessively related to this structure, using an FDR < 0.1.
Putative evidence of selection was considered only for SNP loci that were
conservatively identified by pcadapt and five independent runs of LOSITAN to avoid false-
positives[51]. As selection is presumably stronger in coding regions of the genome and there is
no reference genome for the species, we used the *de novo* assembled transcriptome,
characterized herein, as a reference to identify candidate loci within putative coding regions.
We performed a reciprocal alignment between nextRAD sequences (75 bp) and longer
expressed sequences (≈ 300-11600 bp) using blastn v.2.2.31[36], with a threshold of at least 50
aligned nucleotides, a maximum of one mismatch and no gaps.
268
269 *Data accessibility*
270   Expression data and sequences that support the findings have been deposited in
271 GenBank with the primary accession code GSE116060. A Variant Call Format file and its
272 complementary file, both required for all of the genome-wide SNP diversity analyses are
273 provided as Supplementary Files (Supplementary Datasets 1-2).
274
275

Results
Comparative physiology using a common garden experiment
Seedlings from equatorial and subtropical sites diverged in key ecophysiological traits
in the common garden experiment (Fig. 2-4). The inclination angle of the leaf lamina and the
average size of individual leaves were smaller in equatorial than in subtropical plants, but
total leaf area and specific leaf area did not differ between the groups (Fig. 2, Supplementary
information Table S3, Supplementary information Fig. S3). Additionally, equatorial plants
showed lower stomatal conductance and transpiration rates than did subtropical plants (Fig.
4). Subtropical plants accumulated more biomass in leaves and roots than did equatorial
plants. However, the stem dry mass ratio (MR) (stem dry biomass/plant dry biomass) was
greater in equatorial plants, whereas the leaf MR (leaf dry biomass/plant dry biomass) was
greater in subtropical plants (Fig. 2). The dry biomass accumulated in the stems and the root
MR (root dry biomass/plant dry biomass) were indistinguishable between the groups
(Supplementary information Table S3). Unexpectedly, 63% of the equatorial plants flowered
after six months of growth (Supplementary information Fig. S3g). Since this was not
observed in any subtropical plant, flowering plants were not used in the biomass allocation
analyses.
Mean stem vessel diameter was smaller in the equatorial seedlings than in the
subtropical seedlings, indicating enhanced hydraulic safety (Fig. 2, Supplementary
information Fig. S3h-i). However, the vessel density, vessel grouping index, vessel lumen
area in sapwood and total hydraulic conductivity of the stems were not significantly different
between the groups (P-value > 0.05) (Supplementary information Table S3). Plants from
contrasting latitudes exhibited different stem epidermis pigmentation, with equatorial
seedlings reflecting more long wavelengths red light (635-700 nm) and less medium
wavelengths green light (520-560 nm) than subtropical seedlings (Fig. 3).
301
302 *Characterisation of the A. schaueriana transcriptome*

In the absence of a reference genome, we obtained the first reference transcriptome for *A. schaueriana* from leaves, stems and flowers of adult individuals under field conditions (Supplementary information Fig. S4, Supplementary information Table S1). Over 209 million paired-end 72-bp reads showing high quality were *de novo* assembled into a reference, non-redundant transcriptome containing 49,490 sequences, of which 30,227 (61%) were putative protein-coding sequences. Over 91.9% of these reads were mapped to a single transcript, indicating minimum redundancy and a wide representation of sequenced data (Supplementary information Table S4). Moreover, 91.8% of universal plant orthologous genes were present in the reference transcriptome (Supplementary information Table S4). Sequences with complete ORFs represented approximately 42% (12,796) of all putative protein-coding transcripts (Supplementary information Table S5, Supplementary information Fig. S5). Most of these putative protein-coding sequences (94.33%) showed significant similarity (e-value < 1e-10) to proteins in the Plant RefSeq and TAIR databases (Supplementary information Fig. S5c). More than 80% of protein-coding sequences matched proteins from *Sesamum indicum* or *Erythranthe guttata*, which, as *A. schaueriana*, belong to the order Lamiales (Supplementary information Fig. S6d). We identified 27,658, 18,325 and 13,273 putative orthologs between the *A. schaueriana* reference transcriptome and transcriptomes derived from *A. marina* leaves[52] and *A. officinalis* leaves[53] and roots[54], respectively (Supplementary information Table S6).

Comparative transcriptomics between trees from contrasting latitudes

To identify the effects of environmental variation on gene expression and, thus on phenotypes, at contrasting source sites in the field, we compared expression levels in trees under field conditions as comparable as possible. Sampling was conducted during the low tide, from 10:30 AM to 4:00 PM, at the end of winter at the subtropical site and at the beginning of the dry season at the equatorial site (Supplementary information Table S1). As expected, we observed a consistent source-site pattern in overall transcript expression levels from leaves and stems (Supplementary information Fig. S6a-b). However, flowers transcript expression levels did not show a clear pattern among samples of the same origin (Supplementary information Fig. S6c), leading to the identification of only one DET. Thus, we did not include flowers in the subsequent analyses (Supplementary information Fig. S6f). Conversely, 1,837 and 904 transcripts showed significantly different (FDR < 0.05) relative abundances between equatorial and subtropical samples in leaves and stems, respectively (Supplementary information Fig. S6d-e). Among the total 2,741 DETs, 1,150 (41.91%) were putative protein-coding transcripts.

The assignment of transcripts to GO terms was possible for 25,184 (83.31%) of the 30,227 putative protein-coding sequences, allowing GO enrichment analyses. Analyses were conducted separately for leaves and stems and for each of the following two sets of DETs: one showing higher expression levels in equatorial than in subtropical samples (which we refer to as "DET-Eq") and the other showing higher abundance in subtropical than in equatorial samples (which are referred as "DET-St"). We focused on the biological processes associated with key aspects of the responses to contrasting climate conditions between the equatorial and subtropical sites (Table 1, Supplementary information Fig. S1). The enriched processes among the DET sets included photosynthesis; cell wall biosynthesis; plant responses to ultraviolet radiation (UV), temperature stimulus and water stress (Supplementary information Tables S7-S11, Supplementary information Fig. S6i-l).

Photosynthesis: Among the DET-St set, we observed various putative genes participating in the biosynthesis of the photosynthetic apparatus, chlorophyll and photoreceptors; the function of electron transporters and chloroplast movement coordination. Contrastingly, the DET-Eq set showed enrichment of transcripts similar to proteins required

for disassembling the light-harvesting complex of photosystem II in thylakoid membranes[55] (Supplementary information Table S11).

Cell wall biosynthesis: Transcripts similar to 33 distinct proteins and transcription factors that play central roles in the biosynthesis and deposition of cell wall components, such as cellulose, hemicellulose, lignin and pectin, were identified among DET-Eq (Supplementary information Table S10).

Response to UV: Both the DET-St and DET-Eq sets showed enrichment of functions related to responses to UV radiation; however, the transcript annotations differed between these sets. The DET-St set included putative UV-B protectants and genes involved in UV-B-induced antioxidant biosynthesis, such as plastid copper/zinc superoxide dismutases, photosystem II repair proteins, and L-ascorbic acid. In contrast, the DET-Eq set showed enrichment of transcripts associated with photo-oxidative damage reduction and the positive regulation of anthocyanin biosynthesis in response to UV. Antioxidants induced by UV irradiation[56], such as putative iron superoxide dismutases and pyridoxine biosynthesis genes, were among the DET-Eq set (Supplementary information Table S11).

Response to temperature: In the DET-St set, we observed putative genes presenting critical roles in chloroplast protein translation during cold acclimation and that provide tolerance to chilling temperatures[57,58]. There included transcripts similar to the *GLYCINE-RICH RNA-BINDING* (RZ1A), which has a chaperone activity during cold acclimation[59], and to the cold-inducible ATP-dependent *DNA HELICASE* ATSGS1, required for DNA damage-repair[60]. Interestingly, DET-St included a putative *AGAMOUS-LIKE 24* (AGL24) transcription factor that is involved in vernalisation-induced floral transition[61]. Contrastingly, various transcripts similar to heat shock-inducible chaperones and to *ADENINE NUCLEOTIDE ALPHA HYDROLASE-LIKE* (AT3G53990), involved in chaperone-mediated protein folding[62], were among the DET-Eq set, potentially enhancing tolerance to heat in equatorial plants. Additionally, a transcript similar to the *ETHYLENE-RESPONSIVE ELEMENT BINDING* (RAP2.3), which confers resistance to heat and hydrogen peroxide[63], was observed in this group (Supplementary information Table S11).

Response to water stress: Transcripts associated with the response and tolerance to water deficits and with cellular ion homeostasis and osmotic adjustment were enriched among DET-Eq; for instance, a transcript similar to the *ETHYLENE-RESPONSIVE TRANSCRIPTION FACTOR* (RAP2.4), which regulates the expression of several drought-responsive genes, including aquaporins[64,65]. Accordingly, a putative aquaporin *PLASMA MEMBRANE INTRINSIC (*PIP1;4)[66] was found in this set. In DET-Eq, we observed putative genes participating in the synthesis of raffinose, an osmoprotective soluble trisaccharide[67], as well as transcripts similar to osmosensitive ion channels belonging to the *EARLY-RESPONSIVE TO DEHYDRATION STRESS FAMILY*. Correspondingly, we observed an ion channel, *SLAC1 HOMOLOGUE 3* (SLAH3), required for stomatal closure induced by drought stress[68], and the putative *NINE-CIS-EPOXYCAROTENOID DIOXYGENASE 3* (NCED3), which increases plant resistance to water deficits through the accumulation of abscisic acid (ABA), leading to stomatal closure. Possibly as a consequence of decreased stomatal conductance, a transcript similar to the photorespiration gene, *ALANINE-GLYOXYLATE AMINOTRANSFERASE 2 HOMOLOG 3* (AT3G08860)[69], was also observed among the DET-Eq (Supplementary information Table S11).

We confirmed the results obtained from RNA-Seq data computational analyses by qRT-PCR using ten DETs from leaf samples (Supplementary information Fig. S7, Supplementary information Table S12, Supplementary Note).

Detection of SNPs with signs of selection

RNA-Seq enabled the assembly of a reference transcriptome and the identification of
biological processes influenced by the environmental divergence over the South American
coastline. To complement these analyses, as one cannot disentangle the effects of plasticity
and adaptive selection in the differential expression[70,71], we searched for gene sequence
variations among trees sampled along the Atlantic coast of South America (Fig. 1,
Supplementary information Table S2). After quality filtering of the sequenced data, we
selected 77 individuals without evidence of interspecific hybridisation with *A. germinans* for
downstream analyses. We identified a set of 6,170 high-quality unlinked biallelic SNP loci
with a minor allele frequency ≥ 0.05 and $\geq 8x$ coverage. The overall genetic structure of
genome-wide SNPs corroborated a previous study based on putatively neutral loci[22], dividing
the species into two populations: north and south of the northeast extremity of South America
(Fig. 5).
We observed 122 loci showing considerable departures from neutral expectations of
interpopulational divergence, as conservatively detected[72] by pcadapt and LOSITAN. Fifteen
of these loci aligned to *A. schaueriana* transcripts that were similar to gene models in *A.*
thaliana and *S. indicum* (Supplementary information Table S13), enabling screening for their
potential functional roles. However, five reference proteins did not have informative
functions described for the model plant, hindering functional inferences. Conversely, among
the remaining annotated candidates, we found five putative genes involved in processes
related to contrasting equatorial and subtropical environmental variables (Fig. 6). One
candidate locus was detected in the putative transcription factor *RAP2.4,* which is induced in
response to water and salt stress[64] and regulates the expression of aquaporins[65]. Two other
candidates showed similarity to the transcription factors *ZINC-FINGER PROTEIN 1* (ZFN1),
involved in the regulation of the expression of several water stress genes[73], and the *HYPOXIA*
RESPONSE ATTENUATOR (HRA1), which is strongly induced in response to low oxygen
levels[74]. A putative *UDP GLUCOSYL TRANSFERASE*, an enzyme that catalyses the final
step of anthocyanin biosynthesis wherein pigmentation is produced[75], also showed evidences
of positive selection. Additionally, one candidate locus was in a transcript similar to the
TETRATRICOPEPTIDE REPEAT gene (AT2G20710, TPR), which might play a role in the
biogenesis of the photosynthetic apparatus[76].

Discussion
In this study, we integrated genomic and ecophysiological approaches to investigate
the foundations of adaptive variations in a dominant tree from the Atlantic coast of South
America. We tested the hypothesis that latitudinal variations in climate regimens shape
genetic and phenotypic variation involved in resource-use strategies in widespread coastal
trees, using the mangrove *Avicennia schaueriana* as a model. Overall, our results supported
this hypothesis. Using a common garden experiment, we detected differences between plants
from contrasting latitudes in key ecophysiological traits involved in carbon and water
balances, indicating an inheritable basis of trait divergence. Accordingly, transcriptomic
changes between plants sampled in contrasting latitudes showed enrichment in processes
associated with central aspects of the environmental divergence across latitudes, such as
responses to temperature, solar radiation, water deficit, photosynthesis and cell wall
biosynthesis (Supplementary information Fig. S6i-l). The relevance of these biological
processes in the field was corroborated by the identification of SNP loci with signs of
selection present within putative coding transcripts similar to genes involved in responses to
osmotic stress, accumulation of anthocyanins and photosynthesis, unveiling a genetic basis
for environmental adaptation. Figure 6 summarises the integration of three independent

approaches that converge to suggest population adaptation to contrasting environments over
the species range.
451
Evidence of a conservative resource-use strategy in the equatorial population of <u>A.</u>
<u>schaueriana</u>
Traits exhibited by equatorial plants compared to subtropical plants during the
common garden experiment, such as a smaller leaf size and angle, higher levels of red light-
reflecting pigments, narrower vessels and lower rates of stomatal conductance, limit carbon
acquisition[77] and may have imposed constraints on carbon gain in equatorial plants.
Accordingly, these plants also accumulated less biomass than subtropical plants (Fig. 2-4,
Supplementary information Fig. S3). Despite causing limitations in growth, these
characteristics allow plants to maintain suitable leaf temperature for photosynthesis while
reducing the risk of cavitation, UV exposure and water loss through the minimisation of
evaporative cooling[77]. Given the relevance of these traits to water and carbon balances,
especially under high salinity environments, the prevalence of more conservative resource-
use traits among equatorial samples in the common garden experiment suggested a selection
favouring drought tolerance, likely being advantageous during hot-dry seasons in equatorial
wetlands (August-December). These seasons frequently present a combination of stressful
conditions, including high UV exposure; highly fluctuating soil salinity, resulting from wide
daily tidal range; and high vapour pressure deficit (VPD), resulting from high temperature (>
30 °C) and an air humidity below 60% (Figure 1, Table 1, Supplementary information Fig.
S1). Accordingly, equatorial plants also showed lower transpiration rates than subtropical
plants in the common garden (Fig. 4). Additionally, 63% of the six-month-old equatorial
plants flowered from July-August in the common garden experiment (Supplementary
information Fig. S3g), whereas subtropical plants did not flower. This period is consistent
with phenological observations reported for *A. schaueriana* in both equatorial[78] and southern
subtropical forests[79], and could indicate a genetic basis for the observed variation. Early
flowering is a phenotype with complex genetic components, rarely studied in non-model
organisms, but is renowned as an adaptive mechanism for maximising chances of
reproduction under stressful environments[80].
In their native environment, equatorial plants showed increased expression levels of
putative heat-shock proteins, drought-induced ion transporters, aquaporins and genes that
play central roles in stomatal closure and photorespiration compared to subtropical plants,
likely contributing to heat and drought tolerance. These findings provided multiple lines of
evidence of heat and water stress responses. Higher expression of aquaporins and genes
involved in the accumulation of organic solutes may contribute to lowering the cellular water
potential, while improving drought tolerance in equatorial plants compared to subtropical
plants[67,81]. Additionally, higher expression of several transcripts associated with secondary
cell wall biosynthesis and thickening, may enhance the rigidity of cells and reduce the risk of
collapse during dehydration-rehydration cycles in equatorial trees[82]. Moreover, equatorial
plants showed lower expression of several putative photosynthesis genes than subtropical
plants. In response to drought, high-light and heat stress plants minimise the photooxidative
damage by reducing the photosynthetic activity via repression of photosynthesis genes[83–86].
Remarkably, evidences of selection were detected in two putative transcription factors,
RAP2.4 and *ZFN1*, which play key roles in the regulation of osmotic stress-response[64,73].
These findings support the hypothesis that dry seasons marked by low rainfall and high VPD,
which are caused by the combination of high temperatures and low air humidity[87] in the
equatorial region, induce a pivotal selective pressure for coastal trees populations.
497

Evidence of an acquisitive resource-use strategy in the subtropical population of A. schaueriana

Subtropical plants showed higher stomatal conductance and transpiration rates, higher levels of green light-reflecting pigments, larger leaf area, wider leaf lamina angle and larger xylem vessel diameter than equatorial plants in the common garden experiment (Fig. 2-4, Supplementary information Fig. S3). These characteristics enhance light energy absorbance and carbon acquisition, at the expense of a higher cavitation risk[88,89]. Conversely, this apparent riskier strategy may compensate for seasonal declines in growth resulting from decreasing temperature, photoperiod and light quality at higher-latitudes[90] (Table 1, Supplementary information Fig. S1). Although low temperatures reduce enzymatic activity[91] and, thus, plant growth, the severity of low-temperature stress in the southernmost subtropical mangrove forests of the Americas is likely insufficient to favour the selection of freezing-tolerant adaptations, in contrast to results reported for mangroves at their northernmost edge on the Atlantic coast of the Americas[92]. At the southernmost limit of American mangrove forests, the minimum air temperature does not drop below 0 °C (Table 1, Supplementary information Fig. S1) and remains within the expected mangrove physiological thresholds[93]. Additionally, the small population size of *A. schaueriana* at this location[27] and the arrival of maladapted propagules from northerly populations likely reduce the potential strength of selection favouring cold-adaptive traits.

Corroborating the observed differences in morphophysiological traits, we found divergence at the molecular level that may also be related to the increasing amplitude of variation in photoperiod and light quality towards high latitudes. Plants optimise the use of light energy and adjust photosynthetic activity through the regulation of light-harvesting and photosystem-component genes[83]. Thus, the higher expression levels of transcripts associated with photosynthesis in subtropical than in equatorial plants may have facilitated the absorption of light energy in subtropical plants during winter. Although solar irradiance levels were indistinguishable between sampling sites at the time of sampling (Supplementary information Table S1), transcriptomic changes in putative UV-inducible antioxidant and photodamage repair genes suggest the use of distinct strategies to respond to differential seasonality in photoperiod and solar radiation levels between subtropical and equatorial latitudes. Notably, we observed signatures of selection in two transcripts, one showing similarity to a *UDP-GLUCOSYL TRANSFERASE*, a key enzyme in the anthocyanin biosynthesis pathway, and the other in a transcript similar to a TPR gene, required for chlorophyll biosynthesis and for the biogenesis of the photosynthetic apparatus[76]. These results imply that solar radiation, in addition to VPD, may act as an environmental factor driving selection in *A. schaueriana*.

Although soil oxygen availability affects plant growth in intertidal areas, we did not focus our experiments on the relevance of hypoxia in shaping adaptive divergence in coastal wetlands[94]. However, evidence of selection was detected in a transcription factor highly induced by oxygen deprivation (HRA1)[74]. Additionally, the HRA1 putative homolog also showed a 1.75-fold higher expression in subtropical leaves relative to that in equatorial leaves (Supplementary information Table S1), even though sampling was conducted during the low tide at both sites. As tidal amplitude decreases with increasing latitude along the Atlantic coast of South America[95] (Table 1, Fig. 6), trees are exposed to increasing anoxia conditions southwards. These findings suggest that subtropical populations may have better stress preparedness for hypoxia than equatorial populations.

Climate change and conservation implications

Our results provide compelling evidence that adaptations to contrasting seasonality in freshwater availability and solar radiation are involved in the organisation of diversity of the

dominant coastal tree *A. schaueriana*. The functional divergence described herein might
differentially affect the sensitivity of northerly and southerly populations to a rapidly
changing climate. It has been suggested that the southernmost mangrove forests in the
Americas could expand polewards in the near future[96]. We expect that the observed
acquisitive resource-use characteristics may indeed favour subtropical tree growth under
increased CO_2 concentrations, temperatures and rainfall, as predicted for this region by
2100[97]. However, a great landward relocation of subtropical populations will be necessary to
their persistence in the face of rising sea level[98], due to the narrowing of intertidal zones
towards mid-latitude areas on the Atlantic coast of South America (Fig. 1). Even though
subtropical populations apparently show a better preparedness for hypoxia than equatorial
populations, this scenario is aggravated by the dense coastal occupation by urban and
agricultural areas, which may preclude the landward migration of subtropical mangroves in
South America. Contrastingly, equatorial populations frequently have wider plains that are
potentially available for expansion and lower human occupation, but might be threatened by
reduced rainfall and increased VPD during more intense El Niño-Southern Oscillation
events[97]. Increased temperature will stimulate both respiration[99] and photorespiration[100] and
may eventually offset benefits in carbon acquisition caused by increased CO_2
concentrations[101]. The critical temperature threshold for photosynthesis may be overcome
more frequently, possibly reducing carbon assimilation and productivity[102]. With a
conservative resource-use strategy, further limitations in net carbon assimilation could, in
extreme events, lead to biomass loss or tree mortality triggered by carbon starvation[103]. This
scenario could be especially severe to the semiarid northeast South American coast,
particularly in the face of intense human use of adjacent plains and decreasing trends in
terrestrial freshwater availability and coastal freshwater input[104,105].
Our results corroborate previous studies of marine species from the Northern Atlantic
that associated limited north-south population connectivity with adaptation to latitudinal
environmental dissimilarity[19–21]. As widespread trees distributed along the Atlantic coast of
South America show an overlapping north-south organisation of genetic variation[22–24], we
hypothesise that they may also show divergent adaptations to heterogeneous resource
availability over their distribution ranges. Conservation efforts of coastal ecosystems should
focus on resilient, habitat-forming species that give shelter and act as climate rescuers for
several stress-sensitive species[106]. We recommend that populations of coastal trees occurring
north and south from the northeast extremity of South America should warrant attention as
distinct conservation management units[107] for the long-term persistence of coastal
biodiversity and the ecosystem services they provide.
583
Conclusions
Studies on local adaptations in tropical trees are frequently limited by difficulties in
implementing traditional approaches, such as reciprocal transplants, due to their long
generation time and the lack of basic biological information, as knowledge of their
evolutionary history, reproductive mode or phenological patterns. Despite limitations, we
provided integrated ecophysiological and genomic evidences that support the hypothesis that
seasonal environmental heterogeneity in solar radiation and freshwater availability may shape
the variation of traits associated with resource-use in *A. schaueriana*, a widespread tree along
the Atlantic coastline of South America. The non-neutral trait diversity derived from the
interdisciplinary approach used here provides a perspective into the molecular and
phenotypic scales at which environmental selection may shape functional variation of this
dominant species, on a continental scale. Integrating high-throughput sequencing and
ecophysiological approaches, has shown to be a promising strategy in the study of adaptation
in non-model, long-lived species. To provide more realistic predictions of how dominant

coastal trees may respond to current global climate changes we encourage the development of further studies accounting for phenotypically[14] and genetically[108] informed distribution modelling. Since freshwater availability has been decreasing in coastal areas worldwide[104], strongly compromising the productivity of coastal plants[105], such studies should focus on biological variation involved in the balance between carbon gain and water loss. This knowledge can improve predictions on the future of ecosystems they form and generate key information for forest conservation and management efforts[14,109]. Without the realization and dissemination of such studies, the success of conservation plans for tropical forests and their potential to mitigate climate change will likely be seriously compromised.

Acknowledgements

We thank Prof. Michel G. Vincentz and Prof. Juliana S. Mayer for advising and making their laboratory available for our training and performing of RNA extraction and slide preparation for the wood anatomy analyses. We acknowledge Ilmarina C. Menezes and Stephanie K. Bajay for assistance in the field. We thank Stephanie K. Bajay for providing assistance in the germination of the propagules used in the common garden experiment. Moreover, we thank Greg Mellers and Alessandro Alves-Pereira for critically reading the manuscript. M.V.C. received fellowships from the São Paulo Research Foundation - FAPESP (PhD 2013/26793-7) and from the Coordination for the Improvement of Higher Education Personnel - CAPES Computational Biology Program (PhD SWE 8084/2015-07, PhD 88887.177158/2018-00). G.M.M. received fellowships from FAPESP (PD 2013/08086-1, PD BEPE 2014/22821-9) and a grant from the Brazilian National Council for Scientific and Technological Development - CNPq (PD 448286/2014-9). C.C.S. received fellowships from FAPESP (PD 2015/24346-9) and CAPES Computational Biology Program (88881.185134/2018-01). R.S.O. received a CNPq productivity scholarship and a grant from Microsoft research-FAPESP (2011/52072-0). A.P.S. received a grant from CAPES - Computational Biology Program (88882.160095/2013-01), a research fellowship from CNPq (309661/2014-5) and a CNPq productivity scholarship.

Author contributions

A.P.S., R.S.O., G.M.M. and M.V.C. designed the study. M.V.C. and G.M.M. conducted fieldwork. M.V.C. and C.S.M. cultivated seedlings and performed analyses of morphophysiological data. M.V.C. prepared samples and performed RNA-sequencing. M.V.C., M.D., D.H.O. and G.M.M. analysed RNA-Seq data. C.C.S. and M.V.C. verified RNA-Seq data through qRT-PCR. G.M.M. prepared samples and performed genotyping of genome-wide SNP. G.M.M. and M.V.C. analysed nextRAD results. A.P.S., M.I.Z., G.M.M. and R.S.O. contributed material/reagents/analysis tools. M.V.C. and G.M.M. wrote the manuscript. All authors discussed the results and contributed to the manuscript.

Competing interests

The authors declare no competing interests.

References

1.   Hereford, J. A Quantitative Survey of Local Adaptation and Fitness Trade-Offs. *Am. Nat.* **173**, 579–588 (2009).

2.   Kawecki, T. J. & Ebert, D. Conceptual issues in local adaptation. *Ecol. Lett.* **7**, 1225–

1241 (2004).

3.   Albert, C. H., Grassein, F., Schurr, F. M., Vieilledent, G. & Violle, C. When and how
should intraspecific variability be considered in trait-based plant ecology? *Perspect.*
Plant Ecol. Evol. Syst. **13**, 217–225 (2011).

4.   Aitken, S. N., Yeaman, S., Holliday, J. A., Wang, T. & Curtis-McLane, S. Adaptation,
migration or extirpation: climate change outcomes for tree populations. *Evol. Appl.* **1**,
95–111 (2008).

5.   Jump, A. S. & Peñuelas, J. Running to stand still : adaptation and the response of
plants to rapid climate change. *Ecol. Lett.* **8**, 1010–1020 (2005).

6.   Hoegh-Guldberg, O. & Bruno, J. F. The Impact of Climate Change on the World's
Marine Ecosystems. *Science (80-.).* **328**, 1523–1528 (2010).

7.   Loarie, S. R. *et al.* The velocity of climate change. *Nature* **462**, 1052–1055 (2009).

8.   Duke, N. C. *et al.* A World Without Mangroves? *Science (80-.).* **317**, 41 (2007).

9.   Polidoro, B. A. *et al.* The loss of species: Mangrove extinction risk and geographic
areas of global concern. *PLoS One* **5**, e10095 (2010).

10.   Normile, D. El Niño's warmth devastating reefs worldwide. *Science (80-.).* **352**, 15–
16 (2016).

11.   Duke, N. *et al.* Large-scale dieback of mangroves in Australia. *Mar. Freshw. Res.* **68**,
1816–1829 (2017).

12.   Marbà, N. & Duarte, C. M. Mediterranean warming triggers seagrass (*Posidonia*
oceanica) shoot mortality. *Glob. Chang. Biol.* **16**, 2366–2375 (2010).

13.   Kauffman, J. B. *et al.* Carbon stocks of mangroves and salt marshes of the Amazon
region, Brazil. *Biol. Lett.* **14**, 20180208 (2018).

14.   Moran, E. V., Hartig, F. & Bell, D. M. Intraspecific trait variation across scales:
Implications for understanding global change responses. *Glob. Chang. Biol.* **22**, 137–
150 (2016).

15.   Wee, A. K. S. *et al.* The integration and application of genomic information in
mangrove conservation. *Conserv. Biol.* **33**, 206–209 (2018).

16.   Takayama, K., Tamura, M., Tateishi, Y., Webb, E. L. & Kajita, T. Strong genetic
structure over the American continents and transoceanic dispersal in the mangrove
genus *Rhizophora* (Rhizophoraceae) revealed by broad-scale nuclear and chloroplast
DNA analysis. *Am. J. Bot.* **100**, 1191–1201 (2013).

17.   Mori, G. M., Zucchi, M. I., Sampaio, I. & Souza, A. P. Species distribution and
introgressive hybridization of two *Avicennia* species from the Western Hemisphere
unveiled by phylogeographic patterns. *BMC Evol. Biol.* **15**, 1–15 (2015).

18.   Marshall, D. J., Monro, K., Bode, M., Keough, M. J. & Swearer, S. Phenotype-
environment mismatches reduce connectivity in the sea. *Ecol. Lett.* **13**, 128–140
(2010).

19.   Jeffery, N. W. *et al.* Range-wide parallel climate-associated genomic clines in Atlantic
salmon. *R. Soc. Open Sci.* **4**, 171394 (2017).

20.   Bradbury, I. R. *et al.* Parallel adaptive evolution of Atlantic cod on both sides of the
Atlantic Ocean in response to temperature. *Proc. R. Soc. B Biol. Sci.* **277**, 3725–3734
(2010).

21.   Chu, N. D., Kaluziak, S. T., Trussell, G. C. & Vollmer, S. V. Phylogenomic analyses
reveal latitudinal population structure and polymorphisms in heat stress genes in the
North Atlantic snail *Nucella lapillus*. *Mol. Ecol.* **23**, 1863–1873 (2014).

22.   Mori, G. M., Zucchi, M. I. & Souza, A. P. Multiple-geographic-scale genetic structure
of two mangrove tree species: the roles of mating system, hybridization, limited
dispersal and extrinsic factors. *PLoS One* **10**, e0118710 (2015).

23.   Francisco, P. M., Mori, G. M., Alves, F. M., Tambarussi, E. & Souza, A. P. De.

691		Population genetic structure, introgression, and hybridization in the genus *Rhizophora*
692		along the Brazilian coast. *Ecol. Evol.* **8**, 3491–3504 (2018).
693	24.	Takayama, K., Tateishi, Y., Murata, J. & Kajita, T. Gene flow and population
694		subdivision in a pantropical plant with sea-drifted seeds *Hibiscus tiliaceus* and its
695		allied species: Evidence from microsatellite analyses. *Mol. Ecol.* **17**, 2730–2742
696		(2008).
697	25.	Savolainen, O., Pyhäjärvi, T. & Knürr, T. Gene Flow and Local Adaptation in Trees.
698		*Annu. Rev. Ecol. Evol. Syst.* **38**, 595–619 (2007).
699	26.	Barrett, R. D. H. & Hoekstra, H. E. Molecular spandrels: tests of adaptation at the
700		genetic level. *Nat. Rev. Genet.* **12**, 767–780 (2011).
701	27.	Soares, M. L. G., Estrada, G. C. D., Fernandez, V. & Tognella, M. M. P. Southern
702		limit of the Western South Atlantic mangroves: Assessment of the potential effects of
703		global warming from a biogeographical perspective. *Estuar. Coast. Shelf Sci.* **101**, 44–
704		53 (2012).
705	28.	Souza-Filho, P. W. M. *et al.* Holocene coastal evolution and facies model of the
706		Bragança macrotidal flat on the Amazon Mangrove Coast, northern Brazil. *J. Coast.*
707		*Res.* **1**, 306–310 (2006).
708	29.	Kjerfve, B. *et al.* Morphodynamics of muddy environments along the Atlantic coasts
709		of North and South America. in *Muddy coasts of the World: Processes, Deposits and*
710		*Function* (eds. Healy, T., Wang, Y. & Healy, J.-A.) **4**, 479–532 (Elsevier Science
711		B.V., 2002).
712	30.	Reef, R. *et al.* The effect of atmospheric carbon dioxide concentrations on the
713		performance of the mangrove *Avicennia germinans* over a range of salinities. *Physiol.*
714		*Plant.* **154**, 358–368 (2015).
715	31.	Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of
716		image analysis. *Nat. Methods* **9**, 671–675 (2012).
717	32.	Scholz, A., Klepsch, M., Karimi, Z. & Jansen, S. How to quantify conduits in wood?
718		*Front. Plant Sci.* **4**, 1–11 (2013).
719	33.	Oliveira, R. R., Viana, A. J. C., Reátegui, A. C. E. & Vincentz, M. G. A. An efficient
720		method for simultaneous extraction of high-quality RNA and DNA from various plant
721		tissues. *Genet. Mol. Res.* **14**, 18828–18838 (2015).
722	34.	Patel, R. K. & Jain, M. NGS QC toolkit: A toolkit for quality control of next
723		generation sequencing data. *PLoS One* **7**, e30619 (2012).
724	35.	Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. Ultrafast and memory-efficient
725		alignment of short DNA sequences to the human genome. *Genome Biol.* **10**, R25
726		(2009).
727	36.	Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**,
728		421 (2009).
729	37.	O'Leary, N. A. *et al.* Reference sequence (RefSeq) database at NCBI: Current status,
730		taxonomic expansion, and functional annotation. *Nucleic Acids Res.* **44**, D733–D745
731		(2016).
732	38.	Berardini, T. Z. *et al.* The Arabidopsis Information Resource: Making and Mining the
733		"Gold Standard" Annotated Reference Plant Genome. *Genesis* **53**, 474–485 (2015).
734	39.	Finn, R. D. *et al.* Pfam: The protein families database. *Nucleic Acids Res.* **42**, 222–230
735		(2014).
736	40.	Blake, J. A. *et al.* Gene ontology consortium: Going forward. *Nucleic Acids Res.* **43**,
737		D1049–D1056 (2015).
738	41.	Li, W. & Godzik, A. Cd-hit: A fast program for clustering and comparing large sets of
739		protein or nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).
740	42.	Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M.

BUSCO: Assessing genome assembly and annotation completeness with single-copy
orthologs. *Bioinformatics* **31**, 3210–3212 (2015).

43.   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**,
139–140 (2010).

44.   Young, M. D., Wakefield, M. J., Smyth, G. K. & Oshlack, A. Gene ontology analysis
for RNA-seq: accounting for selection bias. *Genome Biol.* **11**, R14 (2010).

45.   Russello, M. A., Waterhouse, M. D., Etter, P. D. & Johnson, E. A. From promise to
practice: pairing non-invasive sampling with genomics in conservation. *PeerJ* **3**, e1106
(2015).

46.   Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–
2158 (2011).

47.   Jombart, T., Devillard, S. & Balloux, F. Discriminant analysis of principal
components: a new method for the analysis of genetically structured populations. *BMC*
Genet. **11**, 94 (2010).

48.   Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry
in unrelated individuals. *Genome Res.* **19**, 1655–1664 (2009).

49.   Antao, T., Lopes, A., Lopes, R. J., Beja-Pereira, A. & Luikart, G. LOSITAN: A
workbench to detect molecular adaptation based on a Fst -outlier method. *BMC*
Bioinformatics **9**, 1–5 (2008).

50.   Luu, K., Bazin, E. & Blum, M. G. B. pcadapt: An R package to perform genome scans
for selection based on principal component analysis. *Mol. Ecol. Resour.* **33**, 67–77
(2016).

51.   Lotterhos, K. E. & Whitlock, M. C. The relative power of genome scans to detect local
adaptation depends on sampling design and statistical method. *Mol. Ecol.* **24**, 1031–
1046 (2015).

52.   Huang, J. *et al.* Transcriptome Sequencing and Analysis of Leaf Tissue of *Avicennia*
marina Using the Illumina Platform. *PLoS One* **9**, e108785 (2014).

53.   Lyu, H., Li, X., Guo, Z., He, Z. & Shi, S. *De novo* assembly and annotation of the
Avicennia officinalis L. transcriptome. *Mar. Genomics* **38**, 17–20 (2017).

54.   Krishnamurthy, P. *et al.* Transcriptomics analysis of salt stress tolerance in the roots of
the mangrove *Avicennia officinalis*. *Nat. Sci. Reports* **7**, 1–19 (2017).

55.   Park, S.-Y. *et al.* The Senescence-Induced Staygreen Protein Regulates Chlorophyll
Degradation. *Plant Cell* **19**, 1649–1664 (2007).

56.   Myouga, F. *et al.* A Heterocomplex of Iron Superoxide Dismutases Defends
Chloroplast Nucleoids against Oxidative Stress and Is Essential for Chloroplast
Development in *Arabidopsis*. *Plant Cell Online* **20**, 3148–3162 (2008).

57.   Wang, S. *et al.* Chloroplast RNA-Binding Protein RBD1 Promotes Chilling Tolerance
through 23S rRNA Processing in *Arabidopsis*. *PLoS Genet.* **12**, 1–21 (2016).

58.   Goulas, E. *et al.* The chloroplast lumen and stromal proteomes of *Arabidopsis thaliana*
show differential sensitivity to short- and long-term exposure to low temperature.
Plant J. **47**, 720–734 (2006).

59.   Kim, Y. O., Kim, J. S. & Kang, H. Cold-inducible zinc finger-containing glycine-rich
RNA-binding protein contributes to the enhancement of freezing tolerance in
Arabidopsis thaliana. *Plant J.* **42**, 890–900 (2005).

60.   Hartung, F., Suer, S. & Puchta, H. Two closely related RecQ helicases have
antagonistic roles in homologous recombination and DNA repair in *Arabidopsis*
thaliana. *Proc. Natl. Acad. Sci.* **104**, 18836–18841 (2007).

61.   Michaels, S. D. *et al.* AGL24 acts as a promoter of flowering in *Arabidopsis* and is
positively regulated by vernalization. *Plant J.* **33**, 867–874 (2003).

62.   Jung, Y. J. *et al.* Universal Stress Protein Exhibits a Redox-Dependent Chaperone
792   Function in *Arabidopsis* and Enhances Plant Tolerance to Heat Shock and Oxidative
793   Stress. *Front. Plant Sci.* **6**, 1–11 (2015).

794   63.   Ogawa, T. *et al.* Functional Analysis of *Arabidopsis* Ethylene-Responsive Element
795   Binding Protein Conferring Resistance to Bax. *Plant Physiol.* **138**, 1436–1445 (2005).

796   64.   Lin, R. C., Park, H. J. & Wang, H. Y. Role of *Arabidopsis* RAP2.4 in regulating light-
797   and ethylene-mediated developmental processes and drought stress tolerance. *Mol.*
798   *Plant* **1**, 42–57 (2008).

799   65.   Rae, L., Lao, N. T. & Kavanagh, T. A. Regulation of multiple aquaporin genes in
800   *Arabidopsis* by a pair of recently duplicated DREB transcription factors. *Planta* **234**,
801   429–444 (2011).

802   66.   Alexandersson, E. *et al.* Whole gene family expression and drought stress regulation of
803   aquaporins. *Plant Mol. Biol.* **59**, 469–484 (2005).

804   67.   Nishizawa, A., Yabuta, Y. & Shigeoka, S. Galactinol and Raffinose Constitute a Novel
805   Function to Protect Plants from Oxidative Damage. *Plant Physiol.* **147**, 1251–1263
806   (2008).

807   68.   Zhang, A. *et al.* S-Type Anion Channels SLAC1 and SLAH3 Function as Essential
808   Negative Regulators of Inward K^+ Channels and Stomatal Opening in Arabidopsis.
809   *Plant Cell* **28**, 949–965 (2016).

810   69.   Liepman, A. H. & Olsen, L. J. Alanine aminotransferase homologs catalyze the
811   glutamate:glyoxylate aminotransferase reaction in peroxisomes of *Arabidopsis*. *Plant*
812   *Physiol.* **131**, 215–27 (2003).

813   70.   Wolf, J. B. W., Lindell, J. & Backstrom, N. Speciation genetics: current status and
814   evolving approaches. *Philos. Trans. R. Soc. B Biol. Sci.* **365**, 1717–1733 (2010).

815   71.   Gould, B. A., Chen, Y. & Lowry, D. B. Gene Regulatory Divergence Between Locally
816   Adapted Ecotypes in Their Native Habitats. *Mol. Ecol.* **27**, 4174–4188 (2018).

817   72.   Ahrens, C. W. *et al.* The search for loci under selection: trends, biases and progress.
818   *Mol. Ecol.* **27**, 1342–1356 (2018).

819   73.   Sakamoto, H., Araki, T., Meshi, T. & Iwabuchi, M. Expression of a subset of the
820   *Arabidopsis* Cys2/His2-type zinc-finger protein gene family under water stress. *Gene*
821   **248**, 23–32 (2000).

822   74.   Giuntoli, B. *et al.* A trihelix DNA binding protein counterbalances hypoxia-responsive
823   transcriptional activation in *Arabidopsis*. *PLoS Biol.* **12**, (2014).

824   75.   Tohge, T. *et al.* Functional genomics by integrated analysis of metabolome and
825   transcriptome of *Arabidopsis* plants over-expressing an MYB transcription factor.
826   *Plant J.* **42**, 218–235 (2005).

827   76.   Bohne, A. V., Schwenkert, S., Grimm, B. & Nickelsen, J. Roles of Tetratricopeptide
828   Repeat Proteins in Biogenesis of the Photosynthetic Apparatus. *Int. Rev. Cell Mol.*
829   *Biol.* **324**, 187–227 (2016).

830   77.   Reef, R. & Lovelock, C. E. Regulation of water balance in Mangroves. *Ann. Bot.* **115**,
831   385–395 (2015).

832   78.   Menezes, M. P. M. de, Berger, U. & Mehlig, U. Mangrove vegetation in Amazonia: a
833   review of studies from the coast of Pará and Maranhão States, north Brazil. *Acta Amaz.*
834   **38**, 403–420 (2008).

835   79.   De Alvarenga, A. M. S. B., Botosso, P. C. & Soffiatti, P. Stem growth and phenology
836   of three subtropical mangrove tree species. *Brazilian J. Bot.* **40**, 907–914 (2017).

837   80.   Kazan, K. & Lyons, R. The link between flowering time and stress tolerance. *J. Exp.*
838   *Bot.* **67**, 47–60 (2016).

839   81.   Lovelock, C. E. *et al.* The vulnerability of Indo-Pacific mangrove forests to sea-level
840   rise. *Nature* **526**, 559–563 (2015).

82. Gall, H. Le *et al.* Cell Wall Metabolism in Response to Abiotic Stress. *Plants* **4**, 112–
166 (2015).
83. Kimura, M. *et al.* Identification of *Arabidopsis* Genes Regulated by High Light –
Stress Using cDNA Microarray. *Photochem. Photobiol.* **77**, 226–233 (2003).
84. Wang, D. *et al.* Genome-wide temporal-spatial gene expression profiling of drought
responsiveness in rice. *BMC Genomics* **12**, 149 (2011).
85. Moumeni, A. *et al.* Comparative analysis of root transcriptome profiles of two pairs of
drought-tolerant and susceptible rice near-isogenic lines under different drought stress.
BMC Plant Biol. **11**, 174 (2011).
86. Wei, W. *et al.* Melatonin enhances plant growth and abiotic stress tolerance in soybean
plants. *J. Exp. Bot.* **66**, 695–707 (2015).
87. McRae, G. J. A Simple Procedure for Calculating Atmospheric Water Vapor
Concentration. *J. Air Pollut. Control Assoc.* **30**, 394–394 (1980).
88. Stuart, S. A., Choat, B., Martin, K. C., Holbrook, N. M. & Ball, M. C. The role of
freezing in setting the latitudinal limits of mangrove forests. *New Phytol.* **173**, 576–583
(2007).
89. Carlson, J. E., Holsinger, K. E. & Prunier, R. Plant responses to climate in the cape
floristic region of South Africa: Evidence for adaptive differentiation in the
proteaceae. *Evolution (N. Y).* **65**, 108–124 (2011).
90. Pecot, S. D., Horsley, S. B., Battaglia, M. A. & Mitchell, R. J. The influence of
canopy, sky condition, and solar angle on light quality in a longleaf pine woodland.
Can. J. For. Res. **35**, 1356–1366 (2005).
91. Arcus, V. L. *et al.* On the Temperature Dependence of Enzyme-Catalyzed Rates.
Biochemistry **55**, 1681–1688 (2016).
92. Cavanaugh, K. C. *et al.* Poleward expansion of mangroves is a threshold response to
decreased frequency of extreme cold events. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 723–7
(2014).
93. Osland, M. J. *et al.* Climatic controls on the global distribution, abundance, and
species richness of mangrove forests. *Ecol. Monogr.* **87**, 341–359 (2017).
94. Colmer, T. D. & Flowers, T. J. Flooding tolerance in halophytes. *New Phytol.* **179**,
964–974 (2008).
95. Schaeffer-Novelli, Y., Cintrón-Molero, G., Adaime, R. R. & de Camargo, T. M.
Variability of mangrove ecosystems along the Brazilian coast. *Estuaries* **13**, 204–218
(1990).
96. Godoy, M. D. P. & De Lacerda, L. D. Mangroves Response to Climate Change: A
Review of Recent Findings on Mangrove Extension and Distribution. *Ann. Brazilian*
Acad. Sci. **87**, 651–667 (2015).
97. IPCC. *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II*
and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate
Change. *Cambridge University Press* (IPCC, 2014).
98. Ellison, J. C. Vulnerability assessment of mangroves to climate change and sea-level
rise impacts. *Wetl. Ecol. Manag.* **23**, 115–137 (2015).
99. Heskel, M. A. *et al.* Convergence in the temperature response of leaf respiration across
biomes and plant functional types. *Proc. Natl. Acad. Sci.* **113**, 3832–3837 (2016).
100. Jordan, D. & Ogren, W. The CO2/O2 specificity of ribulose 1,5-bisphosphate
carboxylase/oxygenase. *Planta* **161**, 308–313 (1984).
101. Drake, B. G., Gonzàlez-Meler, M. A. & Long, S. P. More efficient plants: a
consequence of rising atmospheric CO2? *Annu. Rev. Plant Physiol. Plant Mol. Biol.*
48, 609–639 (1997).
102. Saenger, P. & Moverly, J. Vegetative phenology along the Queensland coastline. *Proc.*

Ecol. Soc. Aust. **13**, 257–265 (1985).

103. Doughty, C. E. *et al.* Drought impact on forest carbon dynamics and fluxes in
Amazonia. *Nature* **519**, 78–82 (2015).

104. Rodell, M. *et al.* Emerging trends in global freshwater availability. *Nature* **557**, 651–
659 (2018).

105. Osland, M. J. *et al.* Climate and plant control on soil organic matter in coastal
wetlands. *Glob. Chang. Biol.* (2018). doi:10.1111/gcb.14376

106. Bulleri, F. *et al.* Harnessing positive species interactions as a tool against climate-
driven loss of coastal biodiversity. *Plos Biol.* **16**, (2018).

107. Moritz, C. Defining 'Evolutionarily Significant Units' for conservation. *Trends Ecol.*
Evol. **9**, 373–375 (1994).

108. Ikeda, D. H. *et al.* Genetically informed ecological niche models improve climate
change predictions. *Glob. Chang. Biol.* **23**, 164–176 (2017).

109. Holliday, J. A. *et al.* Advances in ecological genomics in forest trees and applications
to genetic resources conservation and breeding. *Mol. Ecol.* 706–717 (2017).
doi:10.1111/mec.13963

110. Vestbo, S., Obst, M., Quevedo Fernandez, F. J., Intanai, I. & Funch, P. Present and
Potential Future Distributions of Asian Horseshoe Crabs Determine Areas for
Conservation. *Front. Mar. Sci.* **5**, (2018).

111. Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G. & Jarvis, A. Very high
resolution interpolated climate surfaces for global land areas. *Int. J. Climatol.* **25**,
1965–1978 (2005).

112. Fick, S. E. & Hijmans, R. J. WorldClim 2: new 1-km spatial resolution climate
surfaces for global land areas. *Int. J. Climatol.* **4315**, 4302–4315 (2017).

113. Sbrocco, E. J. & Barber, P. H. MARSPEC: ocean climate layers for marine spatial
ecology. *Ecology* **94**, 979–979 (2013).

114. Forsythe, W. C., Rykiel, E. J., Stahl, R. S., Wu, H. i. & Schoolfield, R. M. A model
comparison for daylength as a function of latitude and day of year. *Ecol. Modell.* **80**,
87–95 (1995).

Tables
Table 1. Characterisation of subtropical and equatorial sampling sites of propagules used in
the common garden experiment and of plant material used for RNA-sequencing. SD:
standard deviation.

	Subtropical	Equatorial
Köppen-Geiger climate characterisation	Temperate oceanic with hot summer, without a dry season (Cfa)	Tropical monsoon (Am)
Latitude (°)	28 S	0
Annual Average tidal amplitude (m)[†]	0.45 (Microtidal)	4.00 (Macrotidal)
Annual mean temperature (°C)[‡]	20.1	26.4
Minimum temperature of the coldest month (°C)[‡]	11.8	22.0
Maximum temperature of the	28.7	31.1

warmest month (°C)[‡]		
Annual precipitation (mm)[‡]	1,435	2,216
Precipitation of the driest month (mm)[‡]	88	4
Precipitation of the wettest month (mm)[‡]	162	452
Mean air vapour pressure deficit (kPa)[¶]	-0.436	-0.625
Maximum air vapour pressure deficit (kPa)[¶]	-1.459	-1.799
Mean irradiance (KJ m^{-2} day^{-1})[‡]	14,270	17,414
Maximum irradiance (KJ m^{-2} day^{-1})[‡]	20,802	21,671
Minimum irradiance (KJ m^{-2} day^{-1})[‡]	8,201	13,874
Mean sea surface salinity (g/kg)[§]	35.50	34.96
Sea surface salinity in the saltiest month (g/kg)[§]	36.24	36.87
Sea surface salinity in the freshest month (g/kg)[§]	33.73	32.54
Average day length (hours (± SD))[⧉]	12.103 (±1.251)	12.115 (±0.033)
True mangrove species in the area	*Avicennia schaueriana* *Laguncularia racemosa*	*Avicennia germinans* *Avicennia schaueriana* *Laguncularia racemosa* *Rhizophora mangle* *Rhizophora racemosa* *Rhizophora harriisoni*

[†]Source: Vestbo *et al.* (2018). [‡]Source: WorldClim[111,112]. [§]Source: MARSPEC[113]. [¶]Water vapour pressure deficit was calculated from WorldClim 'water vapour pressure' and 'average' and 'maximum temperature' raster layers, as the actual water vapour pressure minus the saturation water vapour pressure. [⧉]Source: 'daylength' function from the R package 'geosphere'[114].

Figures Legends

Figure 1. Geographical locations of *Avicennia schaueriana* sampling sites. Characterisation of *Avicennia schaueriana* sampling sites along a latitudinal gradient on the Atlantic coast of South America. The green-shaded area represents the presence of the species; coloured points represent sampling sites of propagules, used in a common garden experiment, and of plant tissues, used for both DNA- and RNA-sequencing; black points represent sampling sites of plant tissues, used for genomic DNA-sequencing only; arrows represent the flow directions of major oceanic currents. Environmental variation across sampling sites was obtained from public databases of climate ('WorldClim'[111,112]) and tide variables ('Environment Climate Data Sweden'[110]).

942
Figure 2. Pairwise comparison of morphological traits between *Avicennia schaueriana*
seedlings from equatorial and subtropical sites grown using a common garden
experiment. Violin plots represent the distribution of observations for plants from equatorial
(red) and subtropical (blue) sampling sites. Box plots represent the mean, standard error, and
maximum and minimum values. (a) Leaf inclination angle (n = 15 leaves per group, 5 plants
per group); (b) individual leaf area (n = 250 leaves per group, 3 plants per group); (c) vessel
lumen diameter (n = 180 vessels per group, 3 plants per group); (d-f) leaf, stem and root dry
weight (n = 15 plants per group); and (g-i) leaf, stem and root dry mass ratio (organ-specific
dry biomass/plant dry biomass) (n = 15 plants per group). For all variables represented in the
figure, the absence of difference between groups was rejected using the unpaired Student's T-
test or the Mann-Whitney-Wilcoxon U-tests (b and c), at a significance threshold of 0.05.
954
955 **Figure 3. Light reflectance of the stem epidermis of five-month-old** *Avicennia*
956 *schaueriana* **seedlings grown using a common garden experiment.** Grey lines represent
957 the mean reflectance, and colour-shaded areas represent the standard error for each seedling
958 source site (red: equatorial; blue: subtropical, n = 10 plants per group). The visible light
959 spectrum range is highlighted in the figure.
960
Figure 4. Daily curves of gas exchange in leaves from seven-months-old *Avicennia*
schaueriana **seedlings grown using a common garden experiment.** * Represents rejection
of the null hypothesis of the absence of a difference based on the unpaired Student's t-test, at
a significance threshold of 0.05 after Bonferroni correction (n = 10 plants per group). Error
bars represent the standard error. Red line: mean values for equatorial samples; blue line:
mean values for subtropical samples. (a) Transpiration rate; (b) stomatal conductance; (c) net
CO_2 assimilation rate.
968
Figure 5. Population structure of *A. schaueriana* **inferred from genome-wide genotyping**
of single nucleotide polymorphic (SNP) loci. (a) Density of individuals on the first retained
discriminant function, calculated using a multivariate method, the discriminant analysis of
principal components (DAPC); different colours represent inferred populations; (b) map
showing the distribution of the species (green shaded area), the geographical location of
sampling sites (points) and barplots in which each colour denotes an ancestral cluster and
each bar represents an individual as inferred by the model-based method implemented in
Admixture 1.3.0.
977
Figure 6. Graphical summary of results from ecophysiological and molecular
approaches performed in this work. Under homogeneous conditions, plants from
contrasting latitudes showed key divergences related to the use of water and to the acquisition
of carbohydrates in ecophysiological traits. Differences associated with the response to
contrasting environmental conditions were detected in transcripts expression levels of trees
sampled in their natural habitats. Additionally, a north-south genetic divergence was
observed in 77 trees sampled along the latitudinal gradient using genotypic data from over 6
thousand polymorphic loci. We also identified signatures of differential selective pressures
on specific loci associated with the accumulation of anthocyanins (UDP-
GLUCOSYLTRANSFERASE), with photosynthesis (TPR), and with the response to osmotic
stress (RAP2.4 and ZFN1) and to hypoxia (HRA1). Our results suggest that the establishing
success of propagules over the latitudinal gradient of the Atlantic coast of South America
may play a role in shaping the diversity of genotypes and phenotypes in the widespread

mangrove, *Avicennia schaueriana*. E: transpiration rate; g_S: stomatal conductance; R_L: xylem
vessel lumen resistivity; K_H: xylem vessel conductivity.
993
994