Title

Local adaptation to water-deficit and light limitation suggested from genomic and

ecophysiological approaches in a coastal habitat-forming tree

Authors

Mariana Vargas Cruz[1], Gustavo Maruyama Mori[2], Caroline Signori Müller[1], Carla Cristina

da Silva[1], Dong-Ha Oh[3], Maheshi Dassanayake[3], Maria Imaculada Zucchi[4], Rafael Silva

Oliveira[1] and Anete Pereira de Souza[1]*

[1]Department of Plant Biology, Institute of Biology, University of Campinas, Campinas, SP

13083-863, Brazil.

[2]Institute of Biosciences, São Paulo State University (Unesp), São Vicente, SP 11330-900,

Brazil.

[3]Department of Biological Sciences, Louisiana State University, Louisiana, LA 70803,

United States.

[4]São Paulo Agency for Agribusiness Technology, Piracicaba, SP 13400-790, Brazil.

*Author for correspondence. Tel: +55 19 3521 1132 Email: anete@unicamp.br

Total word count for the main body of the text: 6491

Introduction: 737

Materials and Methods: 1788

Results: 1789

Discussion: 1838

Conclusions: 193

Acknowledgements: 138

Figures: 4 figures, all coloured

Tables: 1 table

Supporting information: Figures S1-S9 (all coloured, except for S6); Tables S1-S13;

Methods S1-S2; Notes S1

Summary

• Local adaptation is often a product of environmental variation in the geographical space

and has important implications for species' response to climate change. Ongoing changes

in climate are predicted to particularly affect coastal ecosystems, which frequently span

broad latitudinal ranges and remarkable environmental variation. We investigated the

local adaptation to contrasting environments over the distribution of a coastal habitat-

forming tree, based on analyses of key resource-use characteristics.

• We integrated results from comparative ecophysiology under a common garden

experiment, with transcriptome sequencing and genome-wide scans for selection in one

of the most widely distributed trees in wetlands of the Atlantic coast of South America,

Avicennia schaueriana.

• We observed differences related to water and carbon balance between plants from

contrasting latitudes, supported by gene expression under field conditions and

morphophysiological traits under common garden. Additionally, we found signatures of

selection in putative genes associated with photosynthesis, anthocyanin accumulation and

the response to osmotic and hypoxia stresses.

• Our results strongly suggest the existence of a north-south divergence in key resource-use

characteristics, likely driven by seasonality in water-deficit and light availability. These

findings have implications for conservation and provide the basis for more realistic

predictions on coastal plants responses to a rapidly changing climate.

Keywords

Avicennia schaueriana (Black Mangrove); climate change; coastal ecosystems; comparative

transcriptomics; genome scans for selection; local adaptation; plant ecophysiology; resource-

use.

Introduction

Local adaptation is an ubiquitous consequence of spatial variation in selective environmental forces acting on phenotypic variation (Kawecki & Ebert, 2004; Hereford, 2009), which can result from plasticity or changes in allele frequencies within populations (Albert *et al.*, 2011). In this process, as selection forces reduce the heritable variation within a population, they lead to the specialisation of individuals (Kawecki & Ebert, 2004). Therefore, local adaptation represents an important component of the response of species to environmental changes (Jump & Peñuelas, 2005; Aitken *et al.*, 2008).

Alarmingly high rates of environmental changes have been observed in the last decades, particularly affecting coastal ecosystems (Duke *et al.*, 2007; Loarie *et al.*, 2009; Hoegh-Guldberg & Bruno, 2010; Polidoro *et al.*, 2010). For instance, bleaching of coral reefs worldwide (Normile, 2016), large-scale dieback of mangrove communities (Duke *et al.*, 2017) and increased mortality of seagrass beds (Marbà & Duarte, 2010) have been attributed to ongoing changes in climate. Further environmental changes and coastal biodiversity loss might substantially affect critical socio-economical and ecological services provided by coastal ecosystems, including their role as blue carbon stocks (Costanza *et al.*, 1998; Hoegh-Guldberg & Bruno, 2010; Barbier, 2011; Donato *et al.*, 2011). Thus, studies on the structure of adaptive variation of costal habitat-forming species are necessary to improve predictions on the impacts of future climate conditions on coastal ecosystems (Moran *et al.*, 2016) and to support efficient conservation practices (Wee *et al.*, 2018).

Coastal habitat-forming species are generally water-dispersed over long distances and often occur across wide latitudinal ranges, thus spanning strong environmental variation. The connectivity among populations of these species may be influenced, not only by their dispersion capabilities, but also by means of natural selection, caused by varying establishing success over the broad environmental heterogeneity across latitudes (Marshall *et al.*, 2010). For instance, distinct coastal species in the North Atlantic show a north-south organisation of diversity with evidences of selection for different thermal regimes (Bradbury *et al.*, 2010; Chu *et al.*, 2014; Jeffery *et al.*, 2017). Similarly, in the Southwest Atlantic, an overlapping north-south structure of the genetic diversity, with reduced gene flow between populations, has been observed in phylogenetically distant habitat-forming trees (Takayama *et al.*, 2008; Mori *et al.*, 2015; Francisco *et al.*, 2018). In this context, one could expect northerly and southerly populations of these species to adapt differently to contrasting environments over their latitudinal ranges, especially due to their large population sizes (Kawecki & Ebert, 2004; Savolainen *et al.*, 2007). However, the neutrality of the molecular markers used in

previous works has precluded inferences regarding adaptive variation. The existence of
locally adapted populations remains virtually unknown in coastal foundation species that
sustain biodiversity in the South Atlantic, compromising accurate predictions and suitable
conservation efforts for sustainable management.

Here, we tested the hypothesis that latitudinal heterogeneity in selective forces, such
as light and air temperature and humidity seasonal regimens shape the variation of genotypes
and phenotypes involved in the optimisation of resource-use in widely distributed coastal
habitat-forming trees. To reduce the potential for incorrect conclusions about selection
(Barrett & Hoekstra, 2011), we integrated three independent, but complementary approaches.
We predicted that: (1) Under a common garden experiment, plants from contrasting latitudes
would show phenotypic divergence in key ecophysiological traits; (2) under contrasting
latitudes, transcriptomic changes would be detected in genes involved in the response to the
variation in resources availability; (3) signatures of selection along the species distribution
would be detected in genes involved in the response to the latitudinal variation in resource
availability. We considered the new-world coastal foundation species, *Avicennia schaueriana*
Stapf & Leechman ex Moldenke, found in the Lower Lesser Antilles and widespread from
Venezuela to the southernmost temperate mangroves of the Atlantic coast of South America
(~28 °S) (Soares *et al.*, 2012) as a model to test this hypothesis. The particularly low species
richness in South American mangroves (Duke *et al.*, 1998; Ellison *et al.*, 1999), the
endemism of *A. schaueriana* to this region and its broad latitudinal range, spanning wide
environmental variation (Fig. 1, Table 1, Supporting information Fig. S1), decrease
confounding effects and facilitate ecological predictions and evolutionary inferences,
motivating this choice. We provide unprecedented insights into the effects of fluctuations in
atmospheric water-deficit and light limitation on the phenotypic and genotypic variation and
on the regulation of gene expression of a coastal habitat-forming tree. We discuss the
implications of our results for the persistence of coastal biodiversity in the context of climate
change.

Materials and Methods
Propagule sampling
Mature propagules were collected from ten *Avicennia schaueriana* Stapf & Leechman
ex Moldenke mother trees that were at least 100 m apart from each other, from each of two
natural populations described in a previous work (Mori *et al.*, 2015): (1) the southernmost

range limit of American mangroves, in the subtropical region, and (2) one equatorial site in
one of the world's largest macrotidal mangrove forests (Kjerfve *et al.*, 2002; Souza-Filho *et*
al., 2006), near the northernmost limit of the species range (Fig. 1). We refer to samples
collected in the former site as "subtropical" and those in the latter as "equatorial" throughout
this work. A detailed characterisation of each site can be found in Table 1, Supporting
information 1 and Supporting information Fig. S1.
758
Comparative ecophysiology in a common garden experiment
Propagules sampled from the equatorial and subtropical sites were germinated as
described for *Avicennia germinans* (Reef *et al.*, 2014). After two months, 44 similar-sized
seedlings from each sampling site—with an average height of 18 cm, most with three leaf
pairs and senesced cotyledons—were transplanted to 6 L pots filled with topsoil and sand
(1:1). Seedlings were cultivated for seven months under homogenous conditions in a
glasshouse at the University of Campinas, São Paulo, Brazil (22°49' S 47°04' W), where
automatic sensors coupled with a data logger (Onset Computer Corp.) measured the
atmospheric humidity and temperature every 15 minutes. Seedlings were irrigated daily at 10
a.m. and 5 p.m. with a 3-minute freshwater spray. Twice a week, nutrients were added to the
soil using 600 mL of 0.4X Hoagland solution with 15.0 g L^{-1} NaCl per pot. Pots were rotated
weekly to reduce the effects of environmental heterogeneity. Because the environmental
conditions in the glasshouse differed markedly from those at each sampling site (Supporting
information Fig. S2), none of the individuals benefitted from conditions corresponding to
those of their site of origin.
The light reflectance of stems was measured in ten plants from each sampling site
using a USB4000 spectrophotometer (OceanOptics, Inc.) coupled to a deuterium-halogen
light source (DH-2000; OceanOptics), using a light emission range from 200-900 nm.
Photosynthesis, stomatal conductance and transpiration rates were measured every 2.0-2.5
hours in five six-month-old individuals from each sampling site on two different days using a
Li-Cor 6400 XT (Li-Cor Corp.).
After harvest, three plants without flowers or flower buds from each sampling site
were split into leaves, stems and roots, washed with distilled water, dried for 7 days at 70 °C
and weighed. The individual leaf area, total leaf area and leaf lamina angle per plant were
measured through photographic analyses using ImageJ (Schneider *et al.*, 2012). The specific
leaf area (SLA, cm^2 leaf area kg^{-1} leaf dry biomass) was also calculated for these samples.
Stems were fixed in FAA (Formaldehyde Alcohol Acetic acid), stored in 70% alcohol for

wood anatomy analysis and cut into 30 µm thick transverse sections. Sections were stained
with a mixture of 1% Astra Blue and 50% alcohol (1:1) followed by 1% Safranin O.
Micrographs were taken using an Olympus BX51 microscope coupled to an Olympus DP71
camera (Olympus Corp.). The following wood traits were quantified using ImageJ and R
v.4.0.0: vessel lumen area (A), vessel density in xylem (number of vessels/xylem area),
proportion of solitary vessels (number of solitary vessels/total number of vessels), vessel
grouping index (mean number of vessels per vessel grouping), vessel lumen area ratio in
xylem (vessel lumen area/xylem area) and vessel lumen area in sapwood (vessel lumen
area/sapwood area). The vessel arithmetic diameter (D), vessel hydraulic conductivity (K_H)
and lumen resistivity (R_L) were estimated according to Scholz et al. (Scholz *et al.*, 2013).

Statistical comparisons between equatorial and subtropical samples were performed in
R 4.0.0 using the Mann-Whitney-Wilcoxon unpaired test for non-parametric distributions and
unpaired Student's t-test for parametric distributions with 5% significance level. Multiple-
group comparisons were conducted using one-way analysis of variance (ANOVA) with post
hoc Tukey honest significant difference (HSD) tests.

Plant material for RNA extraction and RNA-sequencing

Plant material used for RNA-sequencing (RNA-Seq) was collected in the sites
described in the "Propagule sampling" section and immediately stored in RNAlater (Ambion,
Inc.). Leaves, stems and flowers from three adult trees at least 100 m apart were collected
from July-August of 2014, corresponding to the end of winter at the subtropical site and the
beginning of the dry season at the equatorial site. A detailed description of environmental
conditions at the time of sampling is available in Supporting information Table S1.

We extracted RNA according to Oliveira et al. (Oliveira *et al.*, 2015) and evaluated its
integrity and purity using agarose gel electrophoresis and a NanoVue spectrophotometer (GE
Healthcare Life Sciences). lllumina TruSeq RNA Sample Preparation kits (Illumina, Inc.)
were used to construct cDNA libraries. cDNA quality was assessed using the Agilent 2100
Bioanalyzer (Agilent Technologies) and concentrations were quantified by qPCR, using the
Sequencing Library qPCR Quantification kit (Illumina, Inc.). Sequencing was performed
using two 36-cycle TruSeq SBS paired-end kits (Illumina, Inc.) and a Genome Analyzer IIx
platform (Illumina, Inc.).

Assembly and characterisation of the transcriptome of <u>A. schaueriana</u>

Adapter sequences were trimmed, and 72 bp paired-end reads were filtered by quality

(Phred score ≥ 20 for at least 70% of read length) using the NGS QC Toolkit v.2.3 (Patel &

Jain, 2012). High-quality reads were used for transcriptome assembly in the CLC Genomics

Workbench (https://www.qiagenbioinformatics.com/). We used the default settings, except

for the distance between pairs (300-500 bp) and k-mer size (45 bp).

Reads were mapped to transcripts using bowtie1 (Langmead *et al.*, 2009) in the

single-read mode using default parameters. Transcripts without read-mapping support were

removed. Functional annotation was performed using blastx v.2.2.31 (Camacho *et al.*, 2009)

with an e-value $< 1^{-10}$. NCBI RefSeq (O'Leary *et al.*, 2016), The Arabidopsis Information

Resource (TAIR) (Berardini *et al.*, 2015) and NCBI non-redundant (nr) databases were used

as reference databases. To minimise contaminants, we excluded transcripts that were

exclusively similar to non-plant sequences. Protein family domains were identified using

HMMER3 (Finn *et al.*, 2014), which iteratively searched transcripts against the Pfam

database. To assign Gene Ontology (GO) terms to transcripts, we used the *Arabidopsis*

thaliana gene association file from the Gene Ontology Consortium (Blake *et al.*, 2015) and

retrieved the information for transcripts showing similar coding sequences in the genome of

A. thaliana. Redundant transcripts were clustered using CD-HIT-EST v.4.6.1 (Li & Godzik,

2006) using the local alignment mode with 95% identity and 70% coverage of the shortest

sequence thresholds. Open reading frames (ORF) were identified using Transdecoder

(http://transdecoder.sf.net). We reduced the redundancy of transcripts in the final assembly

by retaining for each CD-HIT-EST cluster either the sequence with the longest ORF or, in the

absence of sequences containing ORF, the longest sequence.

The completeness of the final transcriptome was assessed using BUSCO (Simão *et*

al., 2015). Additionally, a reciprocal blastn alignment using an e-value threshold of 10^{-10} and

a minimum alignment length of 100 nucleotides with at least 70% identity was used to

compare the *A. schaueriana* transcriptome with other publicly available transcriptomes of

congeneric species.

Comparative transcriptomics using RNA-sequencing

Tissue-specific count data were obtained from the number of reads uniquely mapped

to each transcript of the non-redundant transcriptome using bowtie1 (Langmead *et al.*, 2009)

and normalised using edgeR (Robinson *et al.*, 2010). Differentially expressed transcripts

(DETs) between tissue-specific samples of equatorial and subtropical trees were detected

using the exact test for negative binomial distributions with an adjusted P-value < 0.05. GO

term enrichment analyses of the DETs were performed using GOseq (Young *et al.*, 2010)

with the Wallenius approximation method and P-value < 0.05. Differential expression results

were verified using reverse transcription real-time PCR (qRT-PCR) (Methods S2).

Detection of candidate adaptive loci

We sampled leaves from 79 adult plants at ten locations spanning most of the

distribution of *A. schaueriana* Stapf & Leechman ex Moldenke (Fig. 1, Supporting

information Table S2). We isolated DNA using the DNeasy Plant Mini Kit (QIAGEN) and

NucleoSpin Plant II (Macherey Nagel) following the manufacturers' instructions. DNA

quality and quantity were assessed using 1% agarose electrophoresis and the QuantiFluor

dsDNA System with the Quantus fluorometer (Promega). Nextera-tagmented reductively-

amplified DNA (nextRAD) libraries (Russello *et al.*, 2015) were prepared and sequenced by

SNPsaurus (SNPsaurus) in a HiSeq 2500 (Illumina, Inc.) with 100 bp single-end chemistry.

Genomic DNA fragmentation and short adapter ligation were performed with Nextera

reagent (Illumina, Inc.) followed by amplification with one of the primers matching the

adapter and extending nine arbitrary nucleotides into the genomic DNA. Assembly, mapping

and single nucleotide polymorphic loci (SNP) identification were performed using

proprietary custom scripts (SNPsaurus), which created a reference catalogue of abundant

reads across the combined sample set and mapped reads to this reference, allowing two

mismatches and retaining biallelic loci present in at least 10% of the samples. We further

filtered markers by allowing no more than 65% of missing data, Phred score > 30, 8x

minimum coverage, only one SNP per locus and a minor allele frequency $\geq$ 0.05 using

vcftools v.0.1.12b (Danecek *et al.*, 2011). To reduce paralogy or low-quality genotype calls,

we used a maximum read coverage of 56 (the average read depth times 1.5 standard

deviation).

After excluding plants morphologically identified as *A. schaueriana* with genomic

signs of hybridisation with *A. germinans* (L.) L. (data not published), we assessed the genetic

structure considering all SNPs, using the discriminant analysis of principal components

(DAPC) (Jombart *et al.*, 2010) and ADMIXTURE v.1.3.0 (Alexander *et al.*, 2009). For

DAPC analyses, we considered the number of groups (K) varying from 1-50 and the

Bayesian information criteria for inferring K. Additionally, we used the optim.a.score

function to avoid over-fitting during the discrimination steps. For the ADMIXTURE

analyses, we performed three separate runs for K varying from 1-15, using the block-

relaxation method for point estimation; computing was terminated when estimates increased
by < 0.0001, and the most likely K was determined by cross-validation.

We used two programs to minimise false-positive signs of natural selection:
LOSITAN (Antao *et al.*, 2008), assuming an infinite allele model of mutation, using a
confidence interval of 0.99, a false-discovery rate (FDR) of 0.1, the neutral mean FST and
forcing the mean FST options; and pcadapt 3.0.4 (Luu *et al.*, 2016), which simultaneously
identifies the population structure and the loci excessively related to this structure, using FDR
< 0.1.

Putative evidence of selection was considered only for SNP loci identified
simultaneously by pcadapt and five independent runs of LOSITAN. A reciprocal alignment
between short nextRAD sequences (75 bp) and longer expressed sequences assembled from
RNA-Seq data (≈ 300-11600 bp) was performed to identify putative evidences of selection
within expressed sequences. The alignment was conducted using blastn v.2.2.31 (Camacho *et*
al., 2009), with a threshold of at least 50 aligned nucleotides, a maximum of one mismatch
and no gaps.

Results
Comparative physiology in a common garden experiment
Seedlings from equatorial and subtropical sites (Fig. 2 and 3) expressed functional
divergence in key ecophysiological traits in the common garden experiment. Equatorial
plants showed lower stomatal conductance and transpiration rates than did subtropical plants
(Supporting information Fig. S3). Additionally, the inclination angle of the leaf lamina and
the average size of individual leaves were smaller in equatorial than in subtropical plants, but
total leaf area and specific leaf area did not differ between groups (Fig. 2, Supporting
information Table S3, Supporting information Fig. S4). Subtropical plants accumulated more
biomass in all vegetative organs (leaves, stems and roots) than did equatorial plants.
However, the stem dry mass ratio (MR) (stem dry biomass/plant dry biomass) was greater in
equatorial plants, whereas the leaf MR (leaf dry biomass/plant dry biomass) was greater in
subtropical plants. The root MR (root dry biomass/plant dry biomass) was indistinguishable
between groups (Supporting information Table S3). Unexpectedly, 63% of the equatorial
plants flowered after six months of growth (Supporting information Fig. S4g). Since this was
not observed in any subtropical plant, flowering plants were not used in biomass allocation
analyses.

Mean stem vessel diameter and the fraction of sapwood invested in vessel lumen area

were lower in equatorial seedlings compared to subtropical seedlings, indicating higher

investment in hydraulic safety (Fig. 2, Supporting information Fig. S4h-i). However, vessel

density was not different between groups (Supporting information Table S3). Vessels of

subtropical plants showed higher hydraulic specific conductivity, at a detriment of hydraulic

resistance, than did those of equatorial plants (Fig. 2). Total hydraulic conductivity of the

stems was not significantly different between groups. Plants from contrasting latitudes

exhibited different stem epidermis pigmentation, with equatorial seedlings reflecting more

red light of long wavelengths (635-700 nm) and less green light of medium wavelengths

(520-560 nm) than did subtropical seedlings (Fig. 3).

Characterisation of the first transcriptome of <u>A.</u> <u>schaueriana</u>

In the absence of a reference genome, we obtained the first transcriptome for *A.*

schaueriana from leaves, stems and flowers of adult individuals under field conditions

(Supporting information Fig. S5, Supporting information Table S1). Over 209 million paired-

end 72 bp reads showing high quality were *de novo* assembled into a reference, non-

redundant transcriptome containing 49,490 sequences, of which 30,227 (61%) were putative

protein-coding. Over 91.9% of these reads were mapped to a single transcript, indicating

minimum redundancy and wide representation of sequenced data (Supporting information

Table S4). Moreover, 91.8% of universal plant orthologous genes were present in the

reference transcriptome (Supporting information Table S4). Sequences with complete ORF

represented approximately 42% (12,796) of all putative protein-coding transcripts

(Supporting information Table S5, Supporting information Fig. S6). Most of these putative

protein-coding sequences (94.33%) showed significant similarity to proteins in the Plant

RefSeq and TAIR databases (Supporting information Fig. S6c). More than 80% of protein-

coding sequences matched proteins from *Sesamum indicum* or *Erythranthe guttata*, which, as

A. schaueriana, belong to the order Lamiales (Supporting information Fig. 6d). We identified

27,658, 18,325 and 13,273 putative orthologs between the *A. schaueriana* reference

transcriptome and transcriptomes derived from *A. marina* leaves (Huang *et al.*, 2014), *A.*

officinalis leaves (Lyu *et al.*, 2017) and roots (Krishnamurthy *et al.*, 2017), respectively

(Supporting information Table S6).

Comparative transcriptomics between trees from contrasting latitudes

To identify environmental variables that could influence transcriptomic changes and,
thus phenotypes, at contrasting source sites, we compared transcripts expression levels in
trees under field conditions during the end of winter at the subtropical site and at the
beginning of the dry season at the equatorial site (Supporting information Table S1). We
observed a consistent source-site pattern in overall transcript expression levels from leaves
and stems (Supporting information Fig. S7a-b). However, flowers transcript expression levels
did not show a clear pattern among samples from the same origin (Supporting information
Fig. S7c), leading to the identification of only one DET. Thus, we did not include flowers in
the subsequent analyses (Supporting information Fig. S7f). Conversely, 1,837 and 904
transcripts showed significantly different (FDR < 0.05) relative abundance between
equatorial and subtropical samples in leaves and stems, respectively (Supporting information
Fig. S7d-e). Among the total 2,741 DETs, 1,150 (41.91%) were putative protein-coding
transcripts.

The assignment of transcripts to GO terms was possible for 25,184 (83.31%) of
30,227 putative protein-coding sequences, allowing GO enrichment analyses. Analyses were
conducted separately for leaves and stems and for each of two sets of DETs: one showing
higher expression levels in equatorial than in subtropical samples (which we refer to as
"DET-Eq") and the other showing higher abundance in subtropical than in equatorial samples
(which are referred as "DET-St"). We focused on biological processes potentially regulating
the responses of *A. schaueriana* to contrasting climatic variables in the equatorial and
subtropical sites (Table 1, Supporting information Fig. S1). The enriched processes among
the sets of DET included photosynthesis; cell wall biosynthesis; plant responses to UV,
temperature stimulus and water stress (Supporting information Tables S7-S11, Supporting
information Fig. S7i-l).

Photosynthesis: Among the DET-St, we observed various putative genes participating
in the biosynthesis of the photosynthetic apparatus, chlorophyll and photoreceptors; the
function of electron transporters and chloroplast movement coordination. Contrastingly, the
DET-Eq set showed enrichment in transcripts similar to proteins required for disassembling
the light-harvesting complex of photosystem II in thylakoid membranes (Park *et al.*, 2007)
(Supporting information Table S11).

Cell wall biosynthesis: Transcripts similar to 33 distinct proteins and transcription
factors that play central roles in the biosynthesis and deposition of cell wall components, such
as cellulose, hemicellulose, lignin and pectin, were identified among DET-Eq (Supporting
information Table S10).

Response to UV: Both the DET-St and DET-Eq sets showed enrichment in functions
related to the response to UV-light, however, the transcript annotations differed between
these sets. The DET-St set included putative UV-B protectants and genes involved in UV-B-
induced antioxidants biosynthesis, such as plastid copper/zinc superoxide dismutases,
photosystem II repair proteins, and L-ascorbic acid. In contrast, the DET-Eq set showed
enrichment of transcripts associated with photo-oxidative damage reduction and the positive
regulation of anthocyanin biosynthesis in response to UV. Antioxidants induced by UV
irradiation (Myouga *et al.*, 2008), such as putative iron superoxide dismutases and pyridoxine
biosynthesis genes, were among the DET-Eq (Supporting information Table S11).

Response to temperature: In the DET-St set, we observed putative genes presenting
critical roles in chloroplast protein translation during cold acclimation and that provide
tolerance to chilling temperatures (Goulas *et al.*, 2006; Wang *et al.*, 2016). For instance,
transcripts similar to the *GLYCINE-RICH RNA-BINDING PROTEIN* (*RZ1A*), which has a
chaperone activity during cold acclimation (Kim *et al.*, 2005), and to the cold-inducible ATP-
dependent *DNA HELICASE* ATSGS1, required for DNA damage-repair (Hartung *et al.*,
2007). Interestingly, DET-St included a putative *AGAMOUS-LIKE 24* (*AGL24*) transcription
factor, which is involved in vernalisation-induced floral transition (Michaels *et al.*, 2003).
Contrastingly, various transcripts similar to heat shock-inducible chaperones and to
ADENINE NUCLEOTIDE ALPHA HYDROLASE-LIKE PROTEIN (*AT3G53990*), involved in
chaperone-mediated protein folding (Jung *et al.*, 2015), were among the DET-Eq set,
potentially enhancing tolerance to heat in equatorial plants. Additionally, a transcript similar
to the *ETHYLENE-RESPONSIVE ELEMENT BINDING PROTEIN* (*RAP2.3*), which confers
resistance to heat and hydrogen peroxide (Ogawa *et al.*, 2005), was observed in this group
(Supporting information Table S11).

Response to water stress: Transcripts associated with the response and tolerance to
water deficits and with cellular ion homeostasis and osmotic adjustment were enriched
among DET-Eq. For instance, a transcript similar to the *ETHYLENE-RESPONSIVE*
TRANSCRIPTION FACTOR (*RAP2.4*), which regulates the expression of several drought-
responsive genes, including aquaporins (Lin *et al.*, 2008; Rae *et al.*, 2011). Accordingly, a
putative aquaporin *PLASMA MEMBRANE INTRINSIC PROTEIN (PIP1;4)* (Alexandersson
et al., 2005) was found in this set. We observed in DET-Eq putative genes participating in the
synthesis of raffinose, an osmoprotective soluble trisaccharide (Nishizawa *et al.*, 2008), and
also transcripts similar to osmosensitive ion channels belonging to the *EARLY-RESPONSIVE*
TO DEHYDRATION STRESS FAMILY. Correspondingly, we observed an ion channel,

SLAC1 HOMOLOGUE 3 (*SLAH3*), required for stomatal closure induced by drought stress

(Zhang *et al.*, 2016), and the putative *NINE-CIS-EPOXYCAROTENOID DIOXYGENASE 3*

(*NCED3*), which increases plant resistance to water deficit through the accumulation of

abscisic acid (ABA), leading to stomatal closure. Possibly as a consequence of decreased

stomatal conductance, a transcript similar to the photorespiration gene, *ALANINE-*

GLYOXYLATE AMINOTRANSFERASE 2 HOMOLOG 3 (*AT3G08860*) (Liepman & Olsen,

2003), was also observed among the DET-Eq (Supporting information Table S11).

We confirmed the results obtained in the computational analyses of RNA-Seq data by

qRT-PCR of ten DET in leaf samples (Supporting information Fig. S8, Supporting

information Table S12, Supporting information Notes S1).

1029

Detection of SNP with signs of selection

To complement analyses of differential gene expression, which could result from

plasticity and adaptive selection (Wolf *et al.*, 2010; Gould *et al.*, 2018), we searched for gene

sequence variation among trees sampled along the Atlantic coast of South America (Fig. 1,

Supporting information Table S2). After quality filtering of sequenced data, we selected 77

individuals without evidence of interspecific hybridisation with *A. germinans* for downstream

analyses. We identified a set of 6,170 high-quality unlinked biallelic SNP loci with a minor

allele frequency ≥ 0.05 and ≥ 8x coverage. The overall genetic structure of genome-wide

SNPs corroborated a previous study based on putatively neutral loci (Mori *et al.*, 2015),

dividing the species into two populations: north and south of the northeast extremity of South

America (Supporting information Fig. S9).

We observed 122 loci showing significant departures from neutral expectations of

interpopulational divergence, as conservatively detected (Ahrens *et al.*, 2018) by pcadapt and

LOSITAN. Fifteen out of these loci aligned to *A. schaueriana* transcripts that were similar to

gene models in *A. thaliana* and *S. indicum* (Supporting information Table S13), enabling

screening for their potential functional roles. However, five reference proteins did not have

informative functions described for the model plant, hindering functional inferences.

Conversely, among the remaining annotated candidates, we found five putative genes

involved in processes related to contrasting equatorial and subtropical environmental

variables (Fig. 4). One candidate locus was detected in the putative transcription factor

RAP2.4, which is induced in response to water and salt stress (Lin *et al.*, 2008) and regulates

the expression of aquaporins (Rae *et al.*, 2011). Two other candidates showed similarity to

the transcription factors *ZINC-FINGER PROTEIN 1* (*ZFN1*), involved in the regulation of the

expression of several water stress genes (Sakamoto *et al.*, 2000)*,* and the *HYPOXIA*

RESPONSE ATTENUATOR (*HRA1*), strongly induced in response to low oxygen levels

(Giuntoli *et al.*, 2014). A putative *UDP GLUCOSYL TRANSFERASE*, an enzyme that

catalyses the final step of anthocyanin biosynthesis wherein pigmentation is produced (Tohge

et al., 2005), also showed evidences of positive selection. Additionally, one candidate locus

was in a transcript similar to a *TETRATRICOPEPTIDE REPEAT PROTEIN* (*AT2G20710*,

TPR), which might play a role in the biogenesis of the photosynthetic apparatus (Bohne *et al.*,

2016).

Discussion

Studies that integrate genomic and ecophysiological approaches to investigate local

adaptation in dominant tropical trees are virtually absent in the academic literature. We used

complementary ecophysiological and molecular approaches to test the hypothesis that

latitudinal heterogeneity in VPD and light regimens shape genetic and phenotypic variation

involved in resource-use strategies in coastal habitat-forming trees, using the mangrove

Avicennia schaueriana as a model. Overall, our results supported this hypothesis. In a

common garden experiment we detected differences between plants from contrasting

latitudes in key ecophysiological traits involved in carbon and water balances, indicating an

inheritable basis of trait divergence. Accordingly, transcriptomic changes between plants

sampled in contrasting latitudes showed enrichment in processes such as photosynthesis, cell

wall biosynthesis, plant responses to water stress, temperature and UV light (Supporting

information Fig. S7i-l). The relevance of these biological processes in the field was further

supported by the identification of SNP loci with signs of selection within transcripts

associated with response to osmotic stress, accumulation of anthocyanins and photosynthesis,

unveiling a genetic basis of local adaptation. Figure 4 summarises the integration of three

independent approaches that converges to suggest population adaptation to contrasting

environments over the species range.

1079

Evidences of a conservative resource-use strategy in the equatorial population of <u>A.</u>

<u>schaueriana</u>

Traits exhibited by equatorial plants compared to subtropical plants in the common

garden experiment, such as smaller leaf size and angle, higher levels of red light-reflecting

pigments, narrower vessels and lower rates of stomatal conductance, limit carbon acquisition

(Reef & Lovelock, 2015) and may have imposed constraints to carbon gain in equatorial
plants. Accordingly, these plants also accumulated less biomass than subtropical plants (Fig.
2-3, Supporting information Fig. S3-S4). Despite causing limitations in growth, such
characteristics allow plants to maintain suitable leaf temperature for photosynthesis while
reducing the risk of cavitation, UV exposure and water loss through the minimisation of
evaporative cooling (Reef & Lovelock, 2015). Given the relevance of these traits to water
and carbon balances under highly saline environments, the prevalence of more conservative
resource-use traits among equatorial samples suggest a strong selection pressure favouring
drought tolerance, likely being advantageous during hot-dry seasons in equatorial wetlands
(August-December). These seasons frequently present a combination of high light and high
vapour pressure deficit (VPD), resulting from high temperature (> 30 °C) and air humidity
below 60% (Table 1, Supporting information Fig. S1). Accordingly, equatorial plants also
showed lower transpiration rates than subtropical plants in the common garden (Supporting
information Fig. S3). In addition, 63% of the six-month-old equatorial plants flowered from
July-August in the common garden experiment (Supporting information Fig. S4g), whereas
subtropical plants did not flower. This period is consistent with phenological observations
reported for *A. schaueriana* in both equatorial (Menezes *et al.*, 2008) and southern
subtropical forests (De Alvarenga *et al.*, 2017), and could indicate a genetic basis for the
observed variation in this trait. Early flowering is a phenotype with complex genetic
components, rarely studied in non-model organisms, but is renowned as an adaptive
mechanism for maximising chances of reproduction under stressful environments (Kazan &
Lyons, 2016).

In their native environment, equatorial plants showed increased expression levels of
putative heat-shock proteins, drought-induced ion transporters, aquaporins and genes that
play central roles in stomatal closure and photorespiration compared to subtropical plants,
likely contributing to heat and drought tolerance. These findings provided multiple lines of
evidence of heat and water stress responses. Higher expression of aquaporins and genes
involved in the accumulation of organic solutes may contribute to lowering the cellular water
potential, while improving drought tolerance in equatorial plants compared to subtropical
plants (Nishizawa *et al.*, 2008; Lovelock *et al.*, 2015). Additionally, higher expression of
several transcripts associated with secondary cell wall biosynthesis and thickening, may
enhance rigidity of cells and reduce the risk of collapse during dehydration-rehydration
cycles in equatorial trees (Gall *et al.*, 2015). Moreover, equatorial plants showed lower
expression of several putative photosynthesis genes compared to subtropical plants. In

response to drought, high-light and heat stress plants minimise the photooxidative damage by

reducing the photosynthetic activity via repression of photosynthesis genes (Kimura *et al.*,

2003; Moumeni *et al.*, 2011; Wang *et al.*, 2011; Wei *et al.*, 2015). Remarkably, evidences of

selection were detected in two putative transcription factors, *RAP2.4* and *ZFN1*, which play

key roles in the regulation of osmotic stress-response (Sakamoto *et al.*, 2000; Lin *et al.*,

2008). These findings support the hypothesis that seasons marked by high VPD, caused by

the combination of high temperatures and low air humidity (McRae, 1980) in the equatorial

region, play a pivotal selective pressure for coastal trees populations.

1127

Evidences of an acquisitive resource-use strategy in the subtropical population of <u>A.</u>

<u>schaueriana</u>

Subtropical plants showed higher stomatal conductance and transpiration rates, higher

levels of green light-reflecting pigments, larger leaf area, wider leaf lamina angle and larger

xylem vessel diameter than did equatorial plants in the common garden experiment (Fig. 2-3,

Supporting information Fig. S3-S4). These characteristics enhance light energy absorbance

and carbon acquisition, at the expense of a higher cavitation risk (Stuart *et al.*, 2006; Carlson

et al., 2011). Conversely, this apparent riskier strategy may compensate for seasonal declines

in growth resulting from decreasing temperature, photoperiod and light quality in higher-

latitudes (Pecot *et al.*, 2005) (Table 1, Supporting information Fig. S1). Although low

temperatures reduce enzymatic activity (Arcus *et al.*, 2016) and, thus, plant growth, the

severity of low-temperature stress in southernmost subtropical mangrove forests of the

Americas is likely insufficient to favour the selection of freezing-tolerant adaptations, in

contrast to results reported for mangroves at their northernmost edge on the Atlantic coast of

the Americas (Cavanaugh *et al.*, 2014). At the southernmost limit of American mangrove

forests*,* minimum air temperature does not drop below 0 ▫ (Table 1, Supporting information

Fig. S1) and remains within than the expected mangrove physiological thresholds (Osland *et*

al., 2017). Additionally, the small population size of *A. schaueriana* at this location (Soares

et al., 2012) and the arrival of maladapted propagules from northerly populations likely

reduce the potential strength of selection favouring cold-adaptive traits.

Corroborating the observed differences in morphophysiological traits, we found

divergence at the molecular level that may also relate to the increasing amplitude of variation

in photoperiod and light quality towards high latitudes. Plants optimise the use of light energy

and adjust photosynthetic activity through the regulation of light-harvesting and

photosystem-component genes (Kimura *et al.*, 2003). Thus, the higher expression levels of

transcripts associated with photosynthesis in subtropical than in equatorial plants, may facilitate the absorption of light energy in subtropical plants. Although solar irradiance levels were indistinguishable between sampling sites at the time of sampling (Supporting information Table S1), transcriptomic changes in putative UV-inducible antioxidant and photodamage repair genes suggest the use of distinct strategies to respond to differential seasonality in photoperiod and light intensity between subtropical and equatorial latitudes. Notably, we observed signatures of selection in two transcripts, one showing similarity to a *UDP-GLUCOSYL TRANSFERASE*, a key enzyme in the anthocyanin biosynthesis pathway, and the other in a transcript similar to a *TPR* protein, required for chlorophyll biosynthesis and for the biogenesis of the photosynthetic apparatus (Bohne *et al.*, 2016). These results imply that light availability, in addition to VPD, may act as an environmental factor driving selection in *A. schaueriana*.

Although soil oxygen availability affects plant growth in intertidal areas, we did not focus our experiments on the relevance of hypoxia in shaping adaptive divergence in coastal wetlands (Colmer & Flowers, 2008). However, evidence of selection was detected in a transcription factor highly induced by oxygen deprivation (*HRA1*) (Giuntoli *et al.*, 2014). Additionally, the *HRA1* putative homolog also showed a 1.75-fold higher expression in subtropical leaves relative to that in equatorial leaves (Supporting information Table S1), even though sampling was conducted during the low tide at both sites. As tidal amplitude reduces with increasing latitude along the Atlantic coast of South America (Schaeffer-Novelli *et al.*, 1990) (Table 1, Fig. 4), trees are exposed to increasing anoxia conditions southwards. These findings suggest that subtropical populations may have better stress preparedness to hypoxia in comparison to equatorial populations.

Climate change and conservation implications of the results

Our results provide compelling evidences that adaptations to contrasting seasonality in light availability and VPD are involved in the organisation of diversity of the coastal foundation tree, *A. schaueriana*. The functional divergence described herein might differentially affect the sensitivity of northerly and southerly populations to a rapidly changing climate. It has been suggested that southernmost mangrove forests of the Americas could expand polewards in the near future (Godoy & De Lacerda, 2015). We expect that the observed acquisitive resource-use characteristics may favour subtropical trees growth under an increased CO_2 concentration, temperatures and rainfall, as predicted to this region by 2100 (Pachauri & Meyer, 2014). However, a great relocation of subtropical populations will be

necessary to their persistence in the face of rising sea level (Ellison, 2015), due to the narrowing of intertidal zones towards mid-latitude areas in the Atlantic coast of South America (Table 1). Even though subtropical populations apparently show a better preparedness to hypoxia in comparison to equatorial populations, this scenario is aggravated by the dense coastal occupation by urban and agricultural areas, which may preclude the landward migration of subtropical mangroves in South America. Contrastingly, equatorial populations frequently have wider plains potentially available for expansion and lower human occupation, but might be threatened by increased VPD during more intense El Niño-Southern Oscillation events (Pachauri & Meyer, 2014). Increased temperature will likely stimulate both respiration (Heskel *et al.*, 2016) and photorespiration (Jordan & Ogren, 1984) and might offset benefits in carbon acquisition caused by increased CO_2 concentration (Drake *et al.*, 1997). The critical temperature threshold for photosynthesis may be overcome more frequently, possibly reducing carbon assimilation and productivity (Saenger & Moverly, 1985). With a conservative resource-use strategy, further limitations in net carbon assimilation could lead to biomass loss or tree mortality triggered by carbon starvation (Doughty *et al.*, 2015). This scenario could be especially severe in the semi-arid northeast South American coast, particularly in the face of intense human use of adjacent plains and decreasing trends in terrestrial freshwater availability and coastal freshwater input (Osland *et al.*, 2018; Rodell *et al.*, 2018).

Our results corroborate previous studies of marine species, which associated limited north-south population connectivity with adaptation to latitudinal environmental dissimilarity (Bradbury *et al.*, 2010; Chu *et al.*, 2014; Jeffery *et al.*, 2017). As widespread coastal trees distributed along the Atlantic coast of South America show an overlapping north-south organisation of genetic variation (Takayama *et al.*, 2008; Mori *et al.*, 2015; Francisco *et al.*, 2018), we hypothesise that they may also show divergent adaptation to heterogeneous resource availability over their distribution ranges. Conservation efforts of coastal ecosystems should focus on resilient habitat-forming species, which give shelter and act as climate rescuers for several stress-sensitive species (Bulleri *et al.*, 2018). We recommend that populations of foundation trees occurring north and south from the northeast extremity of South America should warrant attention as distinct conservation management units (Moritz, 1994) for the long-term persistence of coastal biodiversity and the ecosystem services they provide.

Conclusions

Using an integrative approach and *A. schaueriana* as a model tree, we provided ecophysiological and genomic evidences that support our hypothesis that seasonal environmental heterogeneity in light availability and VPD shape the variation of traits involved in plant resource-use in coastal wetlands. The non-neutral trait diversity identified here provides unprecedented perspective into the molecular and phenotypic scales at which natural selection in contrasting environments shape functional variation of a dominant tree, along a continental coastal range. In addition, our work demonstrates the power of combining high-throughput sequencing and ecophysiological approaches in the study of local adaptation of non-model habitat-forming species from ecosystems that are highly vulnerable to ongoing climate change. To provide more realistic predictions of how dominant coastal trees may respond to current global climate changes we encourage the development of further studies accounting for phenotypically (Moran *et al.*, 2016) and genetically (Ikeda *et al.*, 2017) informed distribution modelling. Since freshwater availability has been decreasing in coastal areas worldwide (Rodell *et al.*, 2018), regulating important effects in the productivity of coastal plants (Osland *et al.*, 2018), such studies should focus on the variation involved in balance between carbon gain and water loss.

Acknowledgements

We thank Prof. Michel G. Vincentz and Prof. Juliana S. Mayer for advising and making available their laboratory for our training and performing of RNA extraction and slide preparation for the analyses of wood anatomy. We acknowledge Ilmarina C. Menezes and Stephanie K. Bajay for assistance in the field. We thank Stephanie K. Bajay for providing assistance in the germination of the propagules used in the common garden experiment. And we thank Greg Mellers and Alessandro Alves-Pereira for critically reading the manuscript. M.V.C and G.M.M. received fellowships from the São Paulo Research Foundation (FAPESP 2013/26793-7, 2013/08086-1 and 2014/22821-9). M.V.C. received a fellowship from the Coordination for the Improvement of Higher Education Personnel (CAPES 8084/2015-07). G.M.M. received a grant from the Brazilian National Council for Scientific and Technological Development (CNPq 448286/2014-9). A.P.S. received a research fellowship from CNPq (309661/2014-5).

References

Ahrens CW, Rymer PD, Stow A, Bragg J, Dillon S, Umbers KDL, Dudaniec RY. **2018**.

The search for loci under selection: trends, biases and progress. *Molecular Ecology*.

Aitken SN, Yeaman S, Holliday JA, Wang T, Curtis-McLane S. **2008**. Adaptation, migration or extirpation: climate change outcomes for tree populations. *Evolutionary Applications* **1**: 95–111.

Albert CH, Grassein F, Schurr FM, Vieilledent G, Violle C. **2011**. When and how should intraspecific variability be considered in trait-based plant ecology? *Perspectives in Plant Ecology, Evolution and Systematics* **13**: 217–225.

Alexander DH, Novembre J, Lange K. **2009**. Fast model-based estimation of ancestry in unrelated individuals. *Genome Research* **19**: 1655–1664.

Alexandersson E, Fraysse L, Sjövall-Larsen S, Gustavsson S, Fellert M, Karlsson M, Johanson U, Kjellbom P. **2005**. Whole gene family expression and drought stress regulation of aquaporins. *Plant Molecular Biology* **59**: 469–484.

De Alvarenga AMSB, Botosso PC, Soffiatti P. **2017**. Stem growth and phenology of three subtropical mangrove tree species. *Brazilian Journal of Botany*.

Antao T, Lopes A, Lopes RJ, Beja-Pereira A, Luikart G. **2008**. LOSITAN: A workbench to detect molecular adaptation based on a F st -outlier method. *BMC Bioinformatics* **9**: 1–5.

Arcus VL, Prentice EJ, Hobbs JK, Mulholland AJ, Van Der Kamp MW, Pudney CR, Parker EJ, Schipper LA. **2016**. On the Temperature Dependence of Enzyme-Catalyzed Rates. *Biochemistry* **55**: 1681–1688.

Barbier EB. **2011**. Wetlands as natural assets. *Hydrological Sciences Journal* **56**: 1360–1373.

Barrett RDH, Hoekstra HE. **2011**. Molecular spandrels: tests of adaptation at the genetic level. *Nature Reviews Genetics* **12**: 767–780.

Berardini TZ, Reiser L, Li D, Mezheritsky Y, Muller R, Strait E, Huala E. **2015**. The Arabidopsis Information Resource : Making and Mining the " Gold Standard " Annotated Reference Plant Genome. *Genesis* **53**: 474–485.

Blake JA, Christie KR, Dolan ME, Drabkin HJ, Hill DP, Ni L, Sitnikov D, Burgess S, Buza T, Gresham C, *et al*. **2015**. Gene ontology consortium: Going forward. *Nucleic Acids Research* **43**: D1049–D1056.

Bohne A V., Schwenkert S, Grimm B, Nickelsen J. **2016**. *Roles of Tetratricopeptide Repeat Proteins in Biogenesis of the Photosynthetic Apparatus*. Elsevier Inc.

Bradbury IR, Hubert S, Higgins B, Borza T, Bowman S, Paterson IG, Snelgrove PVR, Morris CJ, Gregory RS, Hardie DC, *et al*. **2010**. Parallel adaptive evolution of Atlantic cod on both sides of the Atlantic Ocean in response to temperature. *Proceedings of the Royal*

Society B: Biological Sciences **277**: 3725–3734.

Bulleri F, Eriksson BK, Queiro A, Airoldi L, Jenkins SR, Davoult D, Guizien K, Ives L,

Wahl M, Benedetti-Cecchi L. **2018**. Harnessing positive species interactions as a tool

against climate-driven loss of coastal biodiversity. *Plos Biology* **16**.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL.

2009. BLAST+: architecture and applications. *BMC bioinformatics* **10**: 421.

Carlson JE, Holsinger KE, Prunier R. **2011**. Plant responses to climate in the cape floristic

region of South Africa: Evidence for adaptive differentiation in the proteaceae. *Evolution* **65**:

108–124.

Cavanaugh KC, Kellner JR, Forde AJ, Gruner DS, Parker JD, Rodriguez W, Feller IC.

2014. Poleward expansion of mangroves is a threshold response to decreased frequency of

extreme cold events. *Proceedings of the National Academy of Sciences of the United States of*

America **111**: 723–7.

Chu ND, Kaluziak ST, Trussell GC, Vollmer S V. 2014. Phylogenomic analyses reveal

latitudinal population structure and polymorphisms in heat stress genes in the North Atlantic

snail Nucella lapillus. *Molecular Ecology* **23**: 1863–1873.

Colmer TD, Flowers TJ. **2008**. Flooding tolerance in halophytes. *New Phytologist* **0**.

Costanza R, D'Arge R, de Groot R, Farber S, Grasso M, Hannon B, Limburg K, Naeem

S, O'Neill R V., Paruelo J, *et al.* **1998**. The value of the world's ecosystem services and

natural capital. *Nature* **387**: 253–260.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE,

Lunter G, Marth GT, Sherry ST, *et al.* **2011**. The variant call format and VCFtools.

Bioinformatics **27**: 2156–2158.

Donato DC, Kauffman JB, Murdiyarso D, Kurnianto S, Stidham M, Kanninen M. **2011**.

Mangroves among the most carbon-rich forests in the tropics. *Nature Geoscience* **4**: 293–297.

Doughty CE, Metcalfe DB, Girardin CAJ, Amézquita FF, Cabrera DG, Huasco WH,

Silva-Espejo JE, Araujo-Murakami A, Da Costa MC, Rocha W, *et al.* **2015**. Drought

impact on forest carbon dynamics and fluxes in Amazonia. *Nature* **519**: 78–82.

Drake BG, Gonzàlez-Meler MA, Long SP. **1997**. More efficient plants: a consequence of

rising atmospheric CO2? *Annual Review of Plant Physiology and Plant Molecular Biology*

48: 609–639.

Duke NC, Ball MC, Ellison JC. **1998**. Factors influencing biodiversity and distributional

gradients in mangroves. *Global Ecology and Biogeography Letters* **7**: 27–47.

Duke N, Kovacs J, Griffiths A, Preece L, J. E. Hill D, Van Oosterzee P, Mackenzie J,

Morning H, Burrows D. **2017**. *Large-scale dieback of mangroves in Australia.*

Duke NC, Meynecke J, Ellison AM, Anger K, Berger U, Cannicci S, Diele K, Ewel KC,

Field CD, Koedam N, *et al.* **2007**. A World Without Mangroves ? *Science* **317**: 41.

Ellison JC. **2015**. Vulnerability assessment of mangroves to climate change and sea-level

rise impacts. *Wetlands Ecology and Management* **23**: 115–137.

Ellison AM, Farnsworth EJ, Merkt RE. **1999**. Origins of mangrove ecosystems and the

mangrove biodiversity anomaly. *Global Ecology and Biogeography* **8**: 95–115.

Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR, Heger A,

Hetherington K, Holm L, Mistry J, *et al.* **2014**. Pfam: The protein families database.

Nucleic Acids Research **42**: 222–230.

Forsythe WC, Rykiel EJ, Stahl RS, Wu H i., Schoolfield RM. **1995**. A model comparison

for daylength as a function of latitude and day of year. *Ecological Modelling* **80**: 87–95.

Francisco PM, Mori GM, Alves FM, Tambarussi E, Souza AP De. **2018**. Population

genetic structure, introgression, and hybridization in the genus Rhizophora along the

Brazilian coast. *Ecology and Evolution*: 1–14.

Gall H Le, Philippe F, Domon J-M, Gillet F, Pelloux J, Rayon C. **2015**. Cell Wall

Metabolism in Response to Abiotic Stress. *Plants* **4**: 112–166.

Giuntoli B, Lee SC, Licausi F, Kosmacz M, Oosumi T, van Dongen JT, Bailey-Serres J,

Perata P. **2014**. A Trihelix DNA Binding Protein Counterbalances Hypoxia-Responsive

Transcriptional Activation in Arabidopsis. *PLoS Biology* **12**.

Godoy MDP, De Lacerda LD. **2015**. Mangroves Response to Climate Change: A Review of

Recent Findings on Mangrove Extension and Distribution. *Annals of the Brazilian Academy*

of Sciences **87**: 651–667.

Goulas E, Schubert M, Kieselbach T, Kleczkowski LA, Gardestro P, Schro W. **2006**.

The chloroplast lumen and stromal proteomes of Arabidopsis thaliana show differential

sensitivity to short- and long-term exposure to low temperature. *The Plant Journal* **47**: 720–

734.

Gould BA, Chen Y, Lowry DB. **2018**. Gene Regulatory Divergence Between Locally

Adapted Ecotypes in Their Native Habitats. *Molecular Ecology*.

Hartung F, Suer S, Puchta H. **2007**. Two closely related RecQ helicases have antagonistic

roles in homologous recombination and DNA repair in Arabidopsis thaliana. *Proceedings of*

the National Academy of Sciences **104**: 18836–18841.

Hereford J. **2009**. A Quantitative Survey of Local Adaptation and Fitness Trade Offs. *The*

American Naturalist **173**: 579–588.

1354 **Heskel MA, O'Sullivan OS, Reich PB, Tjoelker MG, Weerasinghe LK, Penillard A,**

1355 **Egerton JJG, Creek D, Bloomfield KJ, Xiang J, *et al.* 2016**. Convergence in the

1356 temperature response of leaf respiration across biomes and plant functional types.

1357 *Proceedings of the National Academy of Sciences* **113**: 3832–3837.

1358 **Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A**. **2005**. Very high resolution

1359 interpolated climate surfaces for global land areas. *International Journal of Climatology* **25**:

1360 1965–1978.

1361 **Hoegh-Guldberg O, Bruno JF**. **2010**. The impact of climate change on the world's marine

1362 ecosystems. *Science (New York, N.Y.)* **328**: 1523–1528.

1363 **Huang J, Lu X, Zhang W, Huang R, Chen S, Zheng Y**. **2014**. Transcriptome Sequencing

1364 and Analysis of Leaf Tissue of Avicennia marina Using the Illumina Platform. *PLoS ONE* **9**:

1365 e108785.

1366 **Ikeda DH, Max TL, Allan GJ, Lau MK, Shuster SM, Whitham TG**. **2017**. Genetically

1367 informed ecological niche models improve climate change predictions. *Global Change*

1368 *Biology* **23**: 164–176.

1369 **Jeffery NW, Stanley RRE, Brendan F, Guijarro-sabaniel J, Bourret V, Bernatchez L,**

1370 **Bentzen P, Beiko RG, Gilbey J, Clément M, *et al.* 2017**. Range-wide parallel climate-

1371 associated genomic clines in Atlantic salmon. *Royal Society Open Science*: 171394.

1372 **Jombart T, Devillard S, Balloux F**. **2010**. Discriminant analysis of principal components: a

1373 new method for the analysis of genetically structured populations. *BMC genetics* **11**: 94.

1374 **Jordan D, Ogren W**. **1984**. The CO2/O2 specificity of ribulose 1,5-bisphosphate

1375 carboxylase/oxygenase. *Planta* **161**: 308–313.

1376 **Jump AS, Peñuelas J**. **2005**. Running to stand still : adaptation and the response of plants

1377 to rapid climate change. *Ecology Letters* **8**: 1010–1020.

1378 **Jung YJ, Melencion SMB, Lee ES, Park JH, Alinapon CV, Oh HT, Yun D-J, Chi YH,**

1379 **Lee SY**. **2015**. Universal Stress Protein Exhibits a Redox-Dependent Chaperone Function in

1380 Arabidopsis and Enhances Plant Tolerance to Heat Shock and Oxidative Stress. *Frontiers in*

1381 *Plant Science* **6**: 1–11.

1382 **Kawecki TJ, Ebert D**. **2004**. Conceptual issues in local adaptation. *Ecology Letters* **7**: 1225–

1383 1241.

1384 **Kazan K, Lyons R**. **2016**. The link between flowering time and stress tolerance. *Journal of*

1385 *Experimental Botany* **67**: 47–60.

1386 **Kim YO, Kim JS, Kang H**. **2005**. Cold-inducible zinc finger-containing glycine-rich RNA-

1387 binding protein contributes to the enhancement of freezing tolerance in Arabidopsis thaliana.

1388   *Plant Journal* **42**: 890–900.

1389   **Kimura M, Yamamoto YY, Seki M, Sakurai T, Sato M, Abe T, Yoshida S, Manabe K,**

1390   **Shinozaki K, Matsui M**. **2003**. Identification of Arabidopsis Genes Regulated by High Light

1391   – Stress Using cDNA Microarray. *Photochemistry and Photobiology* **77**: 226–233.

1392   **Kjerfve B, Perillo GME, Gardner LR, Rine JM, Dias GTM, Mochel FR**. **2002**. Chapter

1393   Twenty Morphodynamics of muddy environments along the Atlantic coasts of North and

1394   South America. *Proceedings in Marine Science* **4**: 479–532.

1395   **Krishnamurthy P, Mohanty B, Wijaya E, Lee D, Lim T, Lin Q, Xu J, Loh C, Kumar**

1396   **PP**. **2017**. Transcriptomics analysis of salt stress tolerance in the roots of the mangrove

1397   Avicennia officinalis. *Nature Scientific Reports* **7**: 1–19.

1398   **Langmead B, Trapnell C, Pop M, Salzberg S**. **2009**. Ultrafast and memory-efficient

1399   alignment of short DNA sequences to the human genome. *Genome Biol.* **10**: R25.

1400   **Li W, Godzik A**. **2006**. Cd-hit: A fast program for clustering and comparing large sets of

1401   protein or nucleotide sequences. *Bioinformatics* **22**: 1658–1659.

1402   **Liepman AH, Olsen LJ**. **2003**. Alanine aminotransferase homologs catalyze the

1403   glutamate:glyoxylate aminotransferase reaction in peroxisomes of Arabidopsis. *Plant*

1404   *physiology* **131**: 215–27.

1405   **Lin RC, Park HJ, Wang HY**. **2008**. Role of Arabidopsis RAP2.4 in regulating light- and

1406   ethylene-mediated developmental processes and drought stress tolerance. *Molecular Plant* **1**:

1407   42–57.

1408   **Loarie SR, Duffy PB, Hamilton H, Asner GP, Field CB, Ackerly DD**. **2009**. The velocity

1409   of climate change. *Nature* **462**: 1052–1055.

1410   **Lovelock CE, Cahoon DR, Friess DA, Guntenspergen GR, Krauss KW, Reef R, Rogers**

1411   **K, Saunders ML, Sidik F, Swales A,** *et al.* **2015**. The vulnerability of Indo-Pacific

1412   mangrove forests to sea-level rise. *Nature* **526**: 559–563.

1413   **Luu K, Bazin E, Blum MGB**. **2016**. pcadapt: An R package to perform genome scans for

1414   selection based on principal component analysis. *Molecular Ecology Resources* **33**: 67–77.

1415   **Lyu H, Li X, Guo Z, He Z, Shi S**. **2017**. De novo assembly and annotation of the Avicennia

1416   officinalis L. transcriptome. *Marine Genomics*: 2–5.

1417   **Marbà N, Duarte CM**. **2010**. Mediterranean warming triggers seagrass (Posidonia oceanica)

1418   shoot mortality. *Global Change Biology* **16**: 2366–2375.

1419   **Marshall DJ, Monro K, Bode M, Keough MJ, Swearer S**. **2010**. Phenotype-environment

1420   mismatches reduce connectivity in the sea. *Ecology Letters* **13**: 128–140.

1421   **McRae GJ**. **1980**. A Simple Procedure for Calculating Atmospheric Water Vapor

Concentration. *Journal of the Air Pollution Control Association* **30**: 394–394.

Menezes MPM de, Berger U, Mehlig U. **2008**. Mangrove vegetation in Amazonia: a review

of studies from the coast of Pará and Maranhão States, north Brazil. *Acta Amazonica* **38**:

403–420.

Michaels SD, Ditta G, Gustafson-Brown C, Pelaz S, Yanofsky M, Amasino RM. **2003**.

AGL24 acts as a promoter of flowering in Arabidopsis and is positively regulated by

vernalization. *Plant Journal* **33**: 867–874.

Moran E V., Hartig F, Bell DM. **2016**. Intraspecific trait variation across scales:

Implications for understanding global change responses. *Global Change Biology* **22**: 137–

150.

Mori GM, Zucchi MI, Souza AP. **2015**. Multiple-Geographic-Scale Genetic Structure of

Two Mangrove Tree Species: The Roles of Mating System, Hybridization, Limited Dispersal

and Extrinsic Factors. *Plos One* **10**: e0118710.

Moritz C. **1994**. Defining 'Evolutionarily Significant Units' for conservation. *Trends in*

Ecology and Evolution **9**: 373–375.

Moumeni A, Satoh K, Kondoh H, Asano T, Hosaka A, Venuprasad R, Serraj R, Kumar

A, Leung H, Kikuchi S. **2011**. Comparative analysis of root transcriptome profiles of two

pairs of drought-tolerant and susceptible rice near-isogenic lines under different drought

stress. *BMC Plant Biology* **11**.

Myouga F, Hosoda C, Umezawa T, Iizumi H, Kuromori T, Motohashi R, Shono Y,

Nagata N, Ikeuchi M, Shinozaki K. **2008**. A Heterocomplex of Iron Superoxide Dismutases

Defends Chloroplast Nucleoids against Oxidative Stress and Is Essential for Chloroplast

Development in Arabidopsis. *the Plant Cell Online* **20**: 3148–3162.

Nishizawa A, Yabuta Y, Shigeoka S. **2008**. Galactinol and Raffinose Constitute a Novel

Function to Protect Plants from Oxidative Damage. *Plant Physiology* **147**: 1251–1263.

Normile D. **2016**. El Niño's warmth devastating reefs worldwide. *Science* **352**: 15–16.

O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, Rajput B,

Robbertse B, Smith-White B, Ako-Adjei D, *et al.* 2016. Reference sequence (RefSeq)

database at NCBI: Current status, taxonomic expansion, and functional annotation. *Nucleic*

Acids Research **44**: D733–D745.

Ogawa T, Pan L, Kawai-Yamada M, Yu L-H, Yamamura S, Koyama T, Kitajima S,

Ohme-Takagi M, Sato F, Uchimiya H. **2005**. Functional Analysis of Arabidopsis Ethylene-

Responsive Element Binding Protein Conferring Resistance to Bax. *Plant physiology* **138**:

1436–1445.

Oliveira RR, Viana AJC, Reátegui ACE, Vincentz MGA. **2015**. An efficient method for
simultaneous extraction of high-quality RNA and DNA from various plant tissues. *Genetics*
and Molecular Research **14**: 18828–18838.

Osland MJ, Feher LC, Griffith KT, Cavanaugh KC, Enwright NM, Day RH, Stagg CL,
Krauss KW, Howard RJ, Grace JB, *et al.* 2017. Climatic controls on the global
distribution, abundance, and species richness of mangrove forests. *Ecological Monographs*
87: 341–359.

Osland MJ, Gabler CA, Grace JB, Day RH, McCoy ML, McLeod JL, From AS,
Enwright NM, Feher LC, Stagg CL, *et al.* 2018. Climate and plant control on soil organic
matter in coastal wetlands. *Global Change Biology*.

Pachauri RK, Meyer LA (Eds.). **2014**. *Climate Change 2014: Synthesis Report.*
Contribution of Working Groups I, II and III to the Fifth Assessment Report of the
Intergovernmental Panel on Climate Change. Geneva, Switzerland: IPCC.

Park S-Y, Yu J-W, Park J-S, Li J, Yoo S-C, Lee N-Y, Lee S-K, Jeong S-W, Seo HS, Koh
H-J, *et al.* 2007. The Senescence-Induced Staygreen Protein Regulates Chlorophyll
Degradation. *The Plant Cell* **19**: 1649–1664.

Patel RK, Jain M. **2012**. NGS QC toolkit: A toolkit for quality control of next generation
sequencing data. *PLoS ONE* **7**.

Pecot SD, Horsley SB, Battaglia MA, Mitchell RJ. **2005**. The influence of canopy, sky
condition, and solar angle on light quality in a longleaf pine woodland. *Canadian Journal of*
Forest Research **35**: 1356–1366.

Polidoro BA, Carpenter KE, Collins L, Duke NC, Ellison AM, Ellison JC, Farnsworth
EJ, Fernando ES, Kathiresan K, Koedam NE, *et al.* 2010. The loss of species: Mangrove
extinction risk and geographic areas of global concern. *PLoS ONE* **5**.

Rae L, Lao NT, Kavanagh TA. **2011**. Regulation of multiple aquaporin genes in
Arabidopsis by a pair of recently duplicated DREB transcription factors. *Planta* **234**: 429–
444.

Reef R, Lovelock CE. **2015**. Regulation of water balance in Mangroves. *Annals of Botany*
115: 385–395.

Reef R, Winter K, Morales J, Adame MF, Reef DL, Lovelock CE. **2014**. The effect of
atmospheric carbon dioxide concentrations on the performance of the mangrove Avicennia
germinans over a range of salinities. *Physiologia Plantarum* **154**.

Robinson MD, McCarthy DJ, Smyth GK. **2010**. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**: 139–140.

Rodell M, Famiglietti JS, Wiese DN, Reager JT, Beaudoing HK, Landerer FW, Lo MH.

2018. Emerging trends in global freshwater availability. *Nature* **557**: 651–659.

Russello MA, Waterhouse MD, Etter PD, Johnson EA. 2015. From promise to practice:

pairing non-invasive sampling with genomics in conservation. *PeerJ* **3**: e1106.

Saenger P, Moverly J. 1985. Vegetative phenology along the Queensland coastline.

Proceedings of the Ecological Society of Australia **13**: 257–265.

Sakamoto H, Araki T, Meshi T, Iwabuchi M. 2000. Expression of a subset of the

Arabidopsis Cys2/His2-type zinc-finger protein gene family under water stress. *Gene* **248**:

23–32.

Savolainen O, Pyhäjärvi T, Knürr T. 2007. Gene Flow and Local Adaptation in Trees.

Annual Review of Ecology, Evolution and Systematics **38**: 595–619.

Sbrocco EJ, Barber PH. 2013. MARSPEC: ocean climate layers for marine spatial ecology.

Ecology **94**: 979–979.

Schaeffer-Novelli Y, Cintrón-Molero G, Adaime RR, de Camargo TM. 1990. Variability

of mangrove ecosystems along the Brazilian coast. *Estuaries* **13**: 204–218.

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of

image analysis. *Nature Methods* **9**: 671–675.

Scholz A, Klepsch M, Karimi Z, Jansen S. 2013. How to quantify conduits in wood?

Frontiers in Plant Science **4**: 1–11.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva E V., Zdobnov EM. 2015.

BUSCO: Assessing genome assembly and annotation completeness with single-copy

orthologs. *Bioinformatics* **31**: 3210–3212.

Soares MLG, Estrada GCD, Fernandez V, Tognella MMP. 2012. Southern limit of the

Western South Atlantic mangroves: Assessment of the potential effects of global warming

from a biogeographical perspective. *Estuarine, Coastal and Shelf Science* **101**: 44–53.

Souza-Filho PWM, Cohen MCL, Lara RJ, Lessa GC, Koch B, Behling H. 2006.

Holocene Coastal Evolution and Facies Model of the Bragança Macrotidal Flat on the

Amazon Mangrove Coast, Northern Brazil. *Journal of Coastal Research*: 306–310.

Stuart SA, Choat B, Martin KC, Holbrook NM, Ball MC. 2006. The role of freezing in

setting the latitudinal limits of mangrove forests. *New Phytologist* **173**: 576–583.

Takayama K, Tateishi Y, Murata J, Kajita T. 2008. Gene flow and population subdivision

in a pantropical plant with sea-drifted seeds Hibiscus tiliaceus and its allied species: Evidence

from microsatellite analyses. *Molecular Ecology* **17**: 2730–2742.

Tohge T, Nishiyama Y, Hirai MY, Yano M, Nakajima JI, Awazuhara M, Inoue E,

Takahashi H, Goodenowe DB, Kitayama M, *et al.* 2005. Functional genomics by

integrated analysis of metabolome and transcriptome of Arabidopsis plants over-expressing

an MYB transcription factor. *Plant Journal* **42**: 218–235.

Vestbo S, Obst M, Quevedo Fernandez FJ, Intanai I, Funch P. **2018**. Present and

Potential Future Distributions of Asian Horseshoe Crabs Determine Areas for Conservation.

Frontiers in Marine Science **5**.

Wang S, Bai G, Wang S, Yang L, Yang F, Wang Y. **2016**. Chloroplast RNA-Binding

Protein RBD1 Promotes Chilling Tolerance through 23S rRNA Processing in Arabidopsis.

PLoS Genetics **12**: 1–21.

Wang D, Pan Y, Zhao X, Zhu L, Fu B, Li Z. **2011**. Genome-wide temporal-spatial gene

expression profiling of drought responsiveness in rice. *BMC Genomics* **12**.

Wee AKS, Mori GM, Lira-Medeiros CF, Núñez-Farfán J, Takayama K, Faulks L,

Suhua S, Tsuda Y, Suyama Y, Yamamoto T, *et al.* 2018. The integration and application of

genomic information in mangrove conservation. *Conservation Biology*: 1–40.

Wei W, Li QT, Chu YN, Reiter RJ, Yu XM, Zhu DH, Zhang WK, Ma B, Lin Q, Zhang

JS, *et al.* 2015. Melatonin enhances plant growth and abiotic stress tolerance in soybean

plants. *Journal of Experimental Botany* **66**: 695–707.

Wolf JBW, Lindell J, Backstrom N. **2010**. Speciation genetics: current status and evolving

approaches. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**:

1717–1733.

Young MD, Wakefield MJ, Smyth GK, Oshlack A. **2010**. Gene ontology analysis for

RNA-seq: accounting for selection bias. *Genome Biology* **11**.

Zhang A, Ren H-M, Tan Y-Q, Qi G-N, Yao F-Y, Wu G-L, Yang L-W, Hussain J, Sun S-

J, Wang Y-F. **2016**. S-Type Anion Channels SLAC1 and SLAH3 Function as Essential

Negative Regulators of Inward K $^+$ Channels and Stomatal Opening in Arabidopsis. *The*

Plant Cell **28**: 949–965.

1550

1551 **Supporting information**

1552 **Fig. S1.** Climate characterisation of equatorial and subtropical sites.

1553

1554 **Fig. S2.** Environmental data in the glasshouse and at the source sites of propagules used in

1555 the common-garden experiment.

1556

1557 **Fig. S3.** Daily curves of gas exchange in leaves of seven-months-old *Avicennia schaueriana*
1558 seedlings grown in a common-garden.

1559

1560 **Fig. S4.** *Avicennia schaueriana* from equatorial and subtropical sites grown in common-
1561 garden.

1562

1563 **Fig. S5.** Sampling of *Avicennia schaueriana* plant material used for RNA-sequencing.

1564

1565 **Fig. S6.** Characterisation of the *de novo* assembly of the *Avicennia schaueriana* reference
1566 transcriptome.

1567

1568 **Fig. S7.** Detection of differentially expressed transcripts (DET) and Gene Ontology (GO)
1569 enrichment analysis.

1570

1571 **Fig. S8.** Validation of RNA-sequencing-based differential expression of transcripts by
1572 reverse transcription real time PCR (qRT-PCR).

1573

1574 **Fig. S9.** Population structure of *A. schaueriana,* inferred from genome-wide genotyping of
1575 single nucleotide polymorphic (SNP) loci.

1576

1577 **Table S1.** Characterisation of field conditions at the time of plant material collection and
1578 storage in RNA-stabilising solution for subsequent sequencing.

1579

1580 **Table S2.** Sampling sites for plant material used in nextRAD sequencing.

1581

1582 **Table S3.** Morphological traits analysed in *Avicennia schaueriana* saplings from equatorial
1583 and subtropical sites, grown under the same environmental conditions during seven months in
1584 a glasshouse.

1585

1586 **Table S4.** Transcriptome quality parameters assessed in this work.

1587

1588 **Table S5.** Transcriptome size characterisation.

1589

1590   **Table S6.** Putative orthologous expressed sequences between *Avicennia schaueriana*
1591   transcripts obtained from flowers, stems and leaves and publicly available transcriptomes
1592   from other *Avicennia* species.

1593

1594   **Table S7.** Enriched Gene Ontology (GO) terms among differentially expressed transcripts
1595   (DET) that presented higher expression in subtropical stem samples than in equatorial stem
1596   samples.

1597

1598   **Table S8.** Enriched Gene Ontology (GO) terms among differentially expressed transcripts
1599   (DET) that presented higher expression in equatorial stem samples than in subtropical stem
1600   samples.

1601

1602   **Table S9.** Enriched Gene Ontology (GO) terms among differentially expressed transcripts
1603   (DET) that presented higher expression in subtropical leaf samples than in equatorial leaf
1604   samples.

1605

1606   **Table S10.** Enriched Gene Ontology (GO) terms among differentially expressed transcripts
1607   (DET) that presented higher expression in equatorial leaf samples than in subtropical leaf
1608   samples.

1609

1610   **Table S11.** Examples of differentially expressed transcripts (DET) of *Avicennia schaueriana*
1611   associated to the response to contrasting climate conditions between equatorial and
1612   subtropical sites.

1613

1614   **Table S12.** Oligonucleotides used in qRT-PCR reactions in this work.

1615

1616   **Table S13.** Single Nucleotide Polymorphic loci (SNP) with signature of selection present in
1617   putative protein-coding regions of the genome of *Avicennia schaueriana.*

1618

1619   **Methods S1.** Propagules and RNA sampling sites characterization.

1620

1621   **Methods S2.** RNA-sequencing data verification through qRT-PCR.

1622

1623   **Notes S1.** Validation of RNA-Seq data through qRT-PCR.

1624

Data accessibility

Expression data and sequences that support the findings have been deposited in

GenBank with the primary accession code GSE116060. A Variant Call Format file and its

complementary file, both required for all of the genome-wide SNP diversity analyses are

available upon request.

Author contributions

A.P.S., R.S.O., G.M.M. and M.V.C. designed the study. M.V.C. and G.M.M. conducted

fieldwork. M.V.C. and C.S.M. cultivated seedlings and performed analyses of

morphophysiological data. M.V.C. prepared samples and performed RNA-sequencing.

M.V.C., M.D., D.H.O. and G.M.M. analysed RNA-Seq data. C.C.S. and M.V.C. verified

RNA-Seq data through qRT-PCR. G.M.M. prepared samples and performed genotyping of

genome-wide SNP. G.M.M. and M.V.C. analysed nextRAD results. A.P.S, M.I.Z., G.M.M.

and R.S.O. contributed material/reagents/analysis tools. M.V.C. and G.M.M. wrote the

manuscript. All authors discussed the results and contributed to the manuscript.

1639

 Figures legends

1641

Figure 1. Geographical locations of *Avicennia schaueriana* **sampling sites.** Green-shaded

area represents the geographical distribution of the species. Black triangles represent the

locations of the equatorial and subtropical sampling sites for propagules and plant tissues

used in the common garden experiment and RNA-sequencing, respectively. Coloured dots

represent sampling sites of leaves used for the nextRAD genome-wide population diversity

analyses (red: population located north of the northeast extremity of South America

(NEESA); blue: population located south of the NEESA). Arrows approximately represent

the directions of the major sea currents along the Atlantic coast of South America.

1650

Figure 2. Pair-wise comparison of morphological traits between *Avicennia schaueriana*

seedlings from equatorial and subtropical sites grown under a common garden

experiment. Violin plots represent the distribution of observations for plants from equatorial

(red) and subtropical (blue) sampling sites. Box plots represent the mean, standard error, and

maximum and minimum values. For all variables represented in the figure, the absence of

difference between groups was rejected using the unpaired Mann-Whitney-Wilcoxon U-tests,

at a significance threshold of 0.05. (a) leaf inclination angle (n = 15 leaves per group, 5 plants

per group); (b) individual leaf area (n = 250 leaves per group, 3 plants per group); (c) leaf dry

mass ratio (leaf dry biomass/plant dry biomass) (n = 15 plants per group); (d) stem dry mass

ratio (stem dry biomass/plant dry biomass) (n = 15 plants per group); (e) vessel lumen area

ratio in sapwood (n = 175 per group, 3 plants per group, observations represented by black

points); (f) vessel lumen diameter (n = 700 vessels per group, 3 plants per group); (g) vessel

lumen conductivity (n = 700 vessels per group, 3 plants per group); (h) vessel lumen

resistivity (n = 700 vessels per group, 3 plants per group).

1665

Figure 3. Light reflectance of the stem epidermis of five-month-old *Avicennia*
schaueriana **seedlings grown in a common garden experiment.** Grey lines represent the
mean reflectance, and colour-shaded areas represent the standard error for each seedling
source site (red: equatorial; blue: subtropical, n = 10 plants per group). The visible light
spectrum range is highlighted in the figure.

1671

1672

1673

Figure 4. Graphical summary of the integration of ecophysiological and molecular approaches performed in this work. Strong variation

in key oceanographic and climatic factors are observed across the broad latitudinal gradient of the Atlantic coast of South America, where

dominant habitat-forming trees share a north-south organisation of the neutral genetic diversity. Besides the influence of sea currents in

propagules dispersion, the establishing success of propagules over the latitudinal range may play a role in shaping the diversity of genotypes and

phenotypes of dominant coastal plants, possibly leading to the evolution of local adaptation. To address this issue, we chose the widespread

mangrove, *Avicennia schaueriana*, as a representative species. We combined the assessment of ecophysiological resource-use traits in plants

from contrasting latitudes in a common garden experiment, comparative transcriptomics in trees sampled under contrasting latitudes, and natural

selection detection tests based on single nucleotide polymorphisms (SNP). Plants from equatorial and subtropical latitudes showed key

divergences related to the use of water and to the acquisition of carbohydrates both in the field and in common garden conditions. In addition,

north-south genetic divergence was observed in all genotyped SNPs. We also identified signatures of differential selective pressures on specific

loci associated to the accumulation of anthocyanins (UDP-GLUCOSYLTRANSFERASE), the response to osmotic stress (RAP2.4 and ZFN1),

photosynthesis (TPR) and hypoxia (HRA1). The molecular and ecophysiologic divergences observed through three independent approaches

suggest the existence of a north-south divergence in resource-use strategies, likely driven by air water vapour and light availability. These

findings highlight the power of using multidisciplinary approaches for the study of adaptation in species for which little basic biological

information is available, such as tropical coastal trees. VPD: atmospheric vapour pressure deficit; E: transpiration rate; g_s: stomatal conductance;

R_L: xylem vessel lumen resistivity; K_H: xylem vessel conductivity.

Tables

1691

1692 **Table 1.** Characterisation of subtropical and equatorial sampling sites of propagules used in

1693 the common garden experiment and of RNA used for RNA-sequencing. SD: standard

1694 deviation of the mean.

	Subtropical	Equatorial
Köppen-Geiger climate characterisation	Temperate oceanic with hot summer, without a dry season (Cfa)	Tropical monsoon (Am)
Latitude (°)	28 S	0
Annual Average tidal amplitude (m)[†]	0.45 (Microtidal)	4.00 (Macrotidal)
Annual mean temperature (°C)[‡]	20.09	26.42
Minimum temperature of the coldest month (°C)[‡]	11.76	22.04
Maximum temperature of the warmest month (°C)[‡]	28.66	31.1
Annual precipitation (mm)[‡]	1,435	2,216
Precipitation in the driest month (mm)[‡]	88	4
Precipitation in the wettest month (mm)[‡]	162	452
Mean air VPD (KPa)[‡]	1.95	2.82
Maximum air VPD (KPa)[‡]	2.47	2.95
Minimum air VPD (KPa)[‡]	1.48	2.71
Mean irradiance (KJ m^{-2} day^{-1})[‡]	14,270	17,414
Maximum irradiance (KJ m^{-2} day^{-1})[‡]	20,802	21,671
Minimum irradiance (KJ m^{-2} day^{-1})[‡]	8,201	13,874
Mean sea surface salinity (g/kg)[§]	35.50	34.96
Sea surface salinity in the saltiest month (g/kg)[§]	36.24	36.87

Sea surface salinity in the freshest month (g/kg)[§]	33.73	32.54
Average day length (hours (± SD))[¶]	12.103 (±1.251)	12.115 (±0.033)
True mangrove species in the area	*Avicennia schaueriana* *Laguncularia racemosa*	*Avicennia germinans* *Avicennia schaueriana* *Laguncularia racemosa* *Rhizophora mangle* *Rhizophora racemosa* *Rhizophora harriisoni*

VPD: Vapour pressure deficit. [†]Source: Vestbo *et al.* (2018). [‡]Source: BioClim (Hijmans *et al.*, 2005). [§]Source: MARSPEC (Sbrocco & Barber, 2013). [¶]Source: 'daylength' function from R package 'geosphere' (Forsythe *et al.*, 1995).

Sampling sites   Key environmental variables   Differential transcript expression in the field   Divergent ecophysiological traits in common garden   Population divergence and genes with signs of natural selection