Title

Local adaptation insights from genomics and ecophysiology of a neotropical mangrove

Short title

Local adaptation of a neotropical mangrove

Authors

Mariana Vargas Cruz[1], Gustavo Maruyama Mori[2], Caroline Signori Müller[1], Carla Cristina da

Silva[1], Dong-Ha Oh[3], Maheshi Dassanayake[3], Maria Imaculada Zucchi[4], Rafael Silva

Oliveira[1] and Anete Pereira de Souza[1]†

[1] Department of Plant Biology, Institute of Biology, University of Campinas, Campinas, São

Paulo, Brazil

[2]Institute of Biosciences, São Paulo State University (Unesp), São Vicente, São Paulo, Brazil

[3]Department of Biological Sciences, Louisiana State University, Louisiana, United States

[4]São Paulo Agency for Agribusiness Technology, Piracicaba, São Paulo, Brazil

†Corresponding author Email: anete@unicamp.br

Abstract

Integrating genomic and ecological data is instrumental for understanding the mechanisms of adaptive processes in natural ecosystems. In non-model species, such studies can be particularly challenging but often yield results with implications for conservation. Here, we integrate molecular and ecophysiological approaches to assess the role of selection in the north-south organisation of genetic variation in the mangrove species *Avicennia schaueriana*, a new-world tree found in tropical to temperate coastal forests along the Atlantic coast of the Americas. We found substantial divergences between populations occurring north and south of the north-eastern extremity of South America, possibly reflecting the roles of contrasting environmental forces in shaping the genetic structure of the species. In a common garden experiment, individuals from equatorial and subtropical forests were found to be divergent in traits involved in water balance and carbon acquisition, suggesting a genetic basis of the observed differences. RNA-sequencing highlighted the molecular effects of different light, temperature and air humidity regimes on individuals under field conditions at contrasting latitudes. Additionally, genome-wide polymorphisms in trees sampled along most of the species' range showed signatures of selection in sequences associated with the biogenesis of the photosynthetic apparatus, anthocyanin biosynthesis and osmotic and hypoxia stress responses. The observed functional divergence might differentially affect sensitivities of populations to our changing climate. We emphasize the necessity of independent conservation management for the long-term persistence of the species' diversity. Moreover, we demonstrate the power of using a multidisciplinary approach in adaptation studies of non-model species.

38   **Keywords**

39   Adaptation genomics, comparative transcriptomics, plant ecophysiology, Avicenniaceae

Introduction

Adaptation to contrasting environments is an ubiquitous consequence of divergent selective forces acting on intraspecific phenotypic diversity (Hereford, 2009; Kawecki & Ebert, 2004). Phenotypic variation can be achieved through plasticity during acclimation to environmental changes or through genetic variation shaped by adaptive processes (Albert, Grassein, Schurr, Vieilledent, & Violle, 2011). Though the occurrence of intraspecific divergence in adaptive traits is well recognised, its molecular basis is not yet fully understood (Savolainen, Lascoux, & Merilä, 2013). The integration of multiple independent approaches to understand the bases of adaptive variation is desirable to minimise the potential for incorrect conclusions (Barrett & Hoekstra, 2011).

In addition to being of fundamental relevance in the field of evolutionary biology, research on the molecular mechanisms underlying adaptive variation may provide valuable information for safeguarding the persistence of populations under environmental challenges. Especially, accelerated rates of contemporary climate change call for studies on functional variation and its consequences for species responses to future climate (Moran, Hartig, & Bell, 2016). Climate change forecasts include a rise in atmospheric CO_2 concentrations up to 550-1000 ppm, leading to a 0.3-4.8 °C increase in mean air temperature, a 0.26-0.82 m sea-level rise and great changes in precipitation regimes by 2100 (Pachauri & Meyer, 2014). These changes are projected to especially affect certain ecosystems, such as mangrove forests (Loarie et al., 2009), since they are distributed in narrow intertidal environments in tropical and subtropical zones and are naturally limited by annual minimum temperatures and average rainfall (Osland et al., 2017). Recent changes have already promoted shifts in the distribution of mangroves and in the density of individuals in populations (Cavanaugh et al., 2014; Duke et al., 2017; Lovelock et al., 2015; Shearman, Bryan, & Walsh, 2013). Negative impacts are

predicted mainly in regions in which aridity is expected to increase and where adjacent areas

for expansion are unavailable or do not exist (Alongi, 2015). However, in some regions,

mangroves might persist through their ability to adjust soil elevation (Lovelock et al., 2015;

McKee, Cahoon, & Feller, 2007) and to rapidly shift their distributions to new suitable areas

(Alongi, 2015; López-Medellín et al., 2011; Lovelock et al., 2015).

Yet, these predictions do not account for extant intraspecific variability across the

geographical distribution of a given species. For instance, the genetic diversity of all

mangrove species studied to date is structured into two populations occurring north and south

of the northeast extremity of South America (NEESA) (Francisco, Mori, Alves, Tambarussi,

& Souza, 2018; Mori, Zucchi, & Souza, 2015). In the NEESA region, the "South Equatorial

Current" bifurcates into "Guiana Current" and "Brazil Current", dispersing mangrove

propagules in opposite directions, likely reducing the gene flow between populations

(Francisco et al., 2018; Mori et al., 2015; Takayama, Tateishi, Murata, & Kajita, 2008) (Fig.

1). As limited gene flow is a key process determining the magnitude of local adaptation, one

could expect these populations to adapt differently to their environments (Kawecki & Ebert,

2004; Savolainen, Pyhäjärvi, & Knürr, 2007). However, the neutral nature of the molecular

markers used in previous works precludes inferences regarding environment-driven genetic

divergence.

Advances in DNA sequencing now permit the identification of neutral and putatively

adaptive genetic variation even in non-model organisms, such as mangrove trees (Andrews,

Good, Miller, Luikart, & Hohenlohe, 2016). Here, we used these tools to investigate the non-

neutral divergence between previously identified populations of the new-world mangrove tree

Avicennia schaueriana Stapf & Leechman ex Moldenke. We further explored the potential

role of environmental forces underlying such divergence along the Atlantic coast of South

America. *Avicennia schaueriana* is the most widely distributed mangrove species for which

genetic diversity information is available over its geographic range. The species is found in

the Lesser Antilles (~16 °N) and from Venezuela to the southernmost mangrove forests in the

Atlantic (~28 °S) (Soares, Estrada, Fernandez, & Tognella, 2012). To avoid generating

incorrect conclusions about the targets of selection (Barrett & Hoekstra, 2011), we integrated

three independent but complementary molecular and ecological approaches: (1) comparative

physiology of equatorial and subtropical samples grown in a common garden experiment; (2)

comparative transcriptomics of trees from equatorial and subtropical localities, sampled in

their native environments (Table 1, Supplemental Fig. 1); and (3) genome scans for signatures

of selection using high-throughput genome-wide genotyping of individuals, sampled along

almost the entire distribution of the species (Fig. 1). We discuss the implications of our results

in the context of rapidly changing climate and suggest strategies for mangrove conservation

along the Atlantic coast of South America.

Materials and Methods

Propagule sampling

Mature propagules were collected from ten *A. schaueriana* mother trees, at least 100

m apart from each other, from each of two natural populations described in a previous work

(Mori et al., 2015): (1) the southernmost range limit of American mangroves, in the

subtropical region, and (2) one equatorial site in one of the world's largest macrotidal

mangrove forests (Kjerfve et al., 2002; Souza-Filho et al., 2006), near the northernmost limit

of the species range (Fig. 1). We refer to samples collected in the former site as 'Subtropical'

and in the latter as 'Equatorial' throughout this work. A detailed characterisation of each site

can be found in Table 1 and in the Supplemental Information file (Supplemental methods S1,

Supplemental Fig. 1).

Comparative ecophysiology in a common garden experiment

Propagules of *A. schaueriana* sampled from the Equatorial and Subtropical sites were

germinated as described for *Avicennia germinans* (Reef et al., 2014). After two months, 44

similar-sized seedlings from each sampling site—with an average height of 18 cm, most with

three leaf pairs and senesced cotyledons—were transplanted to 6 L pots filled with topsoil and

sand (1:1). Seedlings were cultivated for seven months under homogenous conditions in a

glasshouse at the University of Campinas, São Paulo, Brazil (22°49' S 47°04' W), where

automatic sensors coupled with a data logger (Onset Computer Corp.) measured the

atmospheric humidity and temperature every 15 minutes. Seedlings were automatically

irrigated (daily at 10 a.m. and 5 p.m.) with a 3-minute fresh water spray. Twice a week,

nutrients were added to the soil using 600 mL of 0.4X Hoagland solution with 15.0 g L^{-1}

NaCl per pot. Pots were rotated weekly to reduce the effects of environmental heterogeneity.

Because the environmental conditions in the glasshouse differed markedly from those at each

sampling site (Supplemental Fig. 2), none of the individuals benefitted from conditions that

corresponded to those of their origins.

The light reflectance of stems was measured in ten plants from each sampling site

using a USB4000 spectrophotometer (OceanOptics, Inc.) coupled to a deuterium-halogen

light source (DH-2000; OceanOptics), using a light emission range from 200-900 nm.

Photosynthesis, stomatal conductance and transpiration rates were measured every 2.0-2.5

hours in five six-month-old individuals from each sampling site on two different days using a

Li-Cor 6400 XT (Li-Cor Corp.).

After harvest, three plants without flowers or flower buds from each sampling site

were split into leaves, stems and roots, washed with distilled water, dried for 7 days at 70 °C

and weighed. The individual leaf area, total leaf area and leaf lamina angle per plant were

measured through photographic analyses using ImageJ (Schneider, Rasband, & Eliceiri,

2012). The specific leaf area (SLA, cm^2 leaf area kg^{-1} leaf dry biomass) was also calculated

for these samples. Stems were fixed in FAA (Formaldehyde Alcohol Acetic acid), stored in

70% alcohol for wood anatomy analysis and cut into 30 µm thick transverse sections.

Sections were stained with a mixture of 1% Astra Blue and 50% alcohol (1:1) followed by

1% Safranin O. Micrographs were taken from slides using an Olympus BX51 microscope

coupled to an Olympus DP71 camera (Olympus Corp.). The following wood traits were

quantified using ImageJ and R v.4.0.0: vessel lumen area (A), vessel density in xylem

(number of vessels/xylem area), proportion of solitary vessels (number of solitary

vessels/total number of vessels), vessel grouping index (mean number of vessels per vessel

grouping), vessel lumen area ratio in xylem (vessel lumen area/xylem area) and vessel lumen

area in sapwood (vessel lumen area/sapwood area). The vessel arithmetic diameter (D), vessel

hydraulic conductivity (K_H) and lumen resistivity (R_L) were estimated according to Scholz et

al. (Scholz, Klepsch, Karimi, & Jansen, 2013).

Statistical comparisons between Equatorial and Subtropical samples were performed

in R 4.0.0 using the Mann-Whitney-Wilcoxon unpaired test for non-parametric distributions

and unpaired Student's t-test for parametric distributions with 5% significance level.

Multiple-group comparisons were conducted using one-way analysis of variance (ANOVA)

with post hoc Tukey honest significant difference (HSD) tests.

Plant material for RNA extraction and RNA-sequencing

Plant material used for RNA-sequencing (RNA-Seq) was collected in the sites

described in the "Propagule sampling" section. Leaves, stems and flowers from three adult

trees at least 100 m apart were collected in each site from July-August of 2014, corresponding

to the end of winter at the Subtropical site and the beginning of the dry season at the

Equatorial site. A detailed description of environmental conditions at the time of sampling is

available in Supplemental Table 1. Sampling occurred from 11:00 am to 4:00 pm during the

low tide at different altitudes in the intertidal zone. Plant material was washed with sodium

hypochlorite solution (0.2%) and immediately stored in RNAlater (Ambion Inc.).

We extracted RNA according to Oliveira et al. (Oliveira, Viana, Reátegui, &

Vincentz, 2015) and evaluated its integrity and purity using agarose gel electrophoresis and a

NanoVue spectrophotometer (GE Healthcare Life Sciences). lllumina TruSeq RNA Sample

Preparation kits (Illumina Inc.) were used to construct libraries. cDNA quality was assessed

using the Agilent 2100 Bioanalyzer (Agilent Technologies) and concentrations were

quantified by qPCR using the Sequencing Library qPCR Quantification kit (Illumina Inc.)

followed by sequencing using two 36-cycle TruSeq SBS Paired End kits (Illumina Inc.) and a

Genome Analyzer IIx sequencer (Illumina Inc.).

***Assembly and characterisation of the transcriptome of* A. schaueriana**

Adapter sequences were trimmed, and 72 bp paired-end reads were filtered by quality

(Phred score $\geq$ 20 for at least 70% of read length) using the NGS QC Toolkit v.2.3(Patel &

Jain, 2012). High-quality reads were used for subsequent transcriptome assembly in the CLC

Genomics Workbench (https://www.qiagenbioinformatics.com/). We used the default

settings, except for the distance between read pairs (300-500 bp) and k-mer size (45 bp).

Reads were mapped to the assembled transcripts using bowtie1 (Langmead, Trapnell,

Pop, & Salzberg, 2009) in the single-read mode using the default parameters. Transcripts

without read-mapping support were removed. Functional annotation was performed using

blastx v.2.2.31 (Camacho et al., 2009) with an e-value $< 1^{-10}$. The NCBI RefSeq (O'Leary et

al., 2016), The Arabidopsis Information Resource (TAIR) (Berardini et al., 2015) and the

NCBI non-redundant (nr) databases were used as references. To minimize contaminants, we

excluded all transcripts that were exclusively similar to non-plant sequences. Protein family

domains were identified using HMMER3 (Finn et al., 2014), which iteratively searched all

assembled sequences against the Pfam database. To assign Gene Ontology (GO) terms to

transcripts, we used the *Arabidopsis thaliana* gene association file from the Gene Ontology

Consortium (Blake et al., 2015) and retrieved the information for transcripts with similar

coding sequences in the genome of *A. thaliana*. Redundant transcripts were clustered using

CD-HIT-EST v.4.6.1 (Li & Godzik, 2006) using the local alignment mode with 95% identity

and 70% coverage of the shortest sequence thresholds. Open reading frames (ORF) in

putative protein-coding transcripts were identified using Transdecoder

(http://transdecoder.sf.net). We reduced the redundancy of transcripts in the final assembly by

retaining for each CD-HIT-EST cluster either the sequence with the longest ORF or, in the

absence of sequences containing ORF, the longest sequence.

The completeness of the final transcriptome was assessed using BUSCO (Simão,

Waterhouse, Ioannidis, Kriventseva, & Zdbnov, 2015). Additionally, a reciprocal blastn

alignment (Camacho et al., 2009) using an e-value threshold of 10^{-10} and a minimum

alignment length of 100 nucleotides with at least 70% identity was used to compare the *A.*

schaueriana transcriptome with other publicly available transcriptomes of congeneric species.

Comparative transcriptomics using RNA-sequencing

Tissue-specific count data were obtained from the number of reads uniquely mapped

to each transcript of the non-redundant transcriptome using bowtie1 (Langmead et al., 2009)

and normalised using edgeR (Robinson, McCarthy, & Smyth, 2010). Differentially expressed

transcripts (DETs) between tissue-specific samples of trees at the Equatorial and Subtropical

sites were detected using the exact test for negative binomial distributions with an adjusted P-

value < 0.05. GO term enrichment analyses of the DETs were performed using GOseq

(Young, Wakefield, Smyth, & Oshlack, 2010) with the Wallenius approximation method and

P-value < 0.05. Differential expression results were verified using reverse transcription real-

time PCR (qRT-PCR) (Supplemental methods S2).

Detection of candidate adaptive loci in **A. schaueriana**

We sampled leaves from 79 adult plants at ten locations, spanning most of the

geographic range of *A. schaueriana* (Fig. 1, Supplemental Table 2). We isolated DNA using

the DNeasy Plant Mini Kit (QIAGEN) and NucleoSpin Plant II (Macherey Nagel) following

the manufacturers' instructions. DNA quality and quantity were assessed using 1% agarose

electrophoresis and the QuantiFluor dsDNA System with the Quantus fluorometer (Promega).

Nextera-tagmented reductively-amplified DNA (nextRAD) libraries (Russello, Waterhouse,

Etter, & Johnson, 2015) were prepared and sequenced by SNPsaurus (SNPsaurus) in a HiSeq

2500 (Illumina, Inc.) with 100 bp single-end chemistry. Briefly, genomic DNA fragmentation

and short adapter ligation were performed with Nextera reagent (Illumina) followed by

amplification with one of the primers matching the adapter and extending nine arbitrary

nucleotides into the genomic DNA. Thus, the resulting amplicons were fixed at the selective

end, and their lengths were dependent on the initial Nextera fragmentation, leading to

consistent genotyping of the amplified loci. Assembly, mapping and single nucleotide

polymorphic loci (SNP) identification were performed using proprietary custom scripts

(SNPsaurus), which created a reference catalogue of abundant reads across the combined

sample set and mapped reads to this reference, allowing two mismatches and retaining

biallelic loci present in at least 10% of the samples. We further filtered markers by allowing

no more than 65% of missing data, Phred score > 30, 8x minimum coverage, only one SNP

per locus and a minor allele frequency ≥ 0.05 using vcftools v.0.1.12b (Danecek et al., 2011).

To reduce paralogy or low-quality genotype calls, we used a maximum read coverage of 56

(the average read depth times 1.5 standard deviation).

After excluding plants morphologically identified as *A. schaueriana* with genomic

signs of hybridisation with *A. germinans* (data not published), we assessed the genetic

structure in *A. schaueriana*, considering all SNPs, using discriminant analysis of principal

components (DAPC) (Jombart, Devillard, & Balloux, 2010) and ADMIXTURE v.1.3.0

(Alexander, Novembre, & Lange, 2009). For DAPC analyses, we considered the number of

groups varying from 1 to 50 and the Bayesian information criteria for inferring the number of

groups (K). Additionally, we used the optim.a.score function to avoid over-fitting during the

discrimination steps. For the ADMIXTURE analyses, we performed three separate runs for K

varying between 1 and 15, using the block-relaxation method for point estimation; computing

was terminated when estimates increased by < 0.0001, and the most likely K-value was

determined by cross-validation.

We used methods implemented with two programs to minimise false-positive signs of

natural selection: LOSITAN (Antao, Lopes, Lopes, Beja-Pereira, & Luikart, 2008), assuming

an infinite allele model of mutation, using a confidence interval of 0.99, a false-discovery rate

(FDR) of 0.1, the neutral mean FST and forcing the mean FST options; and pcadapt 3.0.4

(Luu, Bazin, & Blum, 2016), which simultaneously identifies the population structure and the

loci excessively related to this structure, using FDR < 0.1.

The sequences presenting SNP with putative evidence of selection were identified

simultaneously by pcadapt and five independent runs of LOSITAN and were searched within

expressed regions of the reference transcriptome. A reciprocal alignment between the short

sequences obtained through nextRAD (75 bp) and the longer expressed sequences assembled

from RNA-Seq data (≈ 300-11600 bp) was performed. The alignment was conducted using

blastn v.2.2.31 (Camacho et al., 2009), with a threshold of at least 50 aligned nucleotides, a

maximum of one mismatch and no gaps.

Results

Comparative physiology in a common garden experiment

To identify functional divergence between individuals from Equatorial and

Subtropical sites, we conducted a common garden experiment in a glasshouse under a

homogeneous, non-climate-controlled environment. We observed differences in key

morphophysiological traits between seedlings from contrasting latitudes (Fig. 2 and 3).

Subtropical plants showed higher transpiration rates and stomatal conductance than did

Equatorial plants (Supplemental Fig. 3). Additionally, the inclination angle of the leaf lamina

was smaller and the average size of individual leaves was smaller in Equatorial than in

Subtropical plants, but the total leaf area and specific leaf area did not significantly differ

between the groups (Fig. 2, Supplemental Table 3, Supplemental Fig. 4). Subtropical plants

accumulated more biomass in all vegetative organs (leaves, stems and roots) than did

Equatorial plants. However, the stem dry mass ratio (stem dry biomass/plant dry biomass)

was greater in Equatorial plants, whereas the leaf dry mass ratio (leaf dry biomass/plant dry

biomass) was greater in Subtropical plants. The root dry mass ratio (root dry biomass/plant

dry biomass) was indistinguishable between groups (Supplemental Table 3). Unexpectedly,

63% of the Equatorial plants flowered after six months of growth (Supplemental Fig. 4g).

Since this was not observed in any Subtropical plant, flowering plants were not used in the

biomass allocation analysis.

Stems from Subtropical samples showed wider vessel diameter than did those from

Equatorial samples, but vessel density was not significantly different between the groups (Fig.

2, Supplemental Fig. 4h-i). The sapwood space dedicated to vessel lumen area was greater in

Subtropical than in Equatorial plants. Vessels of Subtropical plants showed higher

conductivity, at a detriment of hydraulic safety, than did those of Equatorial plants (Fig. 2).

The total conductivity of the stems was not significantly different between sample groups.

Plants from the contrasting origins exhibited different stem epidermis pigmentation, with

Equatorial seedlings reflecting more red light of long wavelengths (635-700 nm) and less

green light of medium wavelengths (520-560 nm) than did Subtropical seedlings (Fig. 3).

Characterisation of the first transcriptome of **A. schaueriana**

In the absence of a reference genome, we used RNA-Seq (Z. Wang, Gerstein, &

Snyder, 2009) to obtain a *de novo* assembled transcriptome for *A. schaueriana* from leaves,

stems and flowers of six adult individuals from Equatorial and Subtropical sites

(Supplemental Fig. 5). Over 209 million paired-end 72 bp reads showing high quality were

assembled into a reference, non-redundant transcriptome containing 49,490 sequences, of

which 30,227 (61%) were putative protein-coding sequences. Over 91.9% of these reads were

mapped to a single transcript, indicating minimum redundancy and a wide representation of

sequenced data (Supplemental Table 4). Moreover, searching for universal plant orthologous

genes revealed that 91.8% of the conserved sequences in plants were present in the reference transcriptome (Supplemental Table 4). Sequences with complete ORF represented approximately 42% (12,796) of all putative protein-coding transcripts, with an average length of 1,324 nucleotides (Supplemental Table 5, Supplemental Fig. 6). Most of these putative protein-coding sequences (94.33%) showed significant similarity to proteins in the Plant RefSeq and TAIR databases (Supplemental Fig. 6c). More than 80% of these protein-coding sequences matched proteins from *Sesamum indicum* or *Erythranthe guttata*, which, as does *A. schaueriana*, belong to the order Lamiales (Supplemental Fig. 6d). We identified 27,658, 18,325 and 13,273 putative orthologs between the *A. schaueriana* reference transcriptome and transcriptomes derived from *A. marina* leaves (Huang et al., 2014), *A. officinalis* leaves (Lyu, Li, Guo, He, & Shi, 2017) and roots (Krishnamurthy et al., 2017), respectively (Supplemental Table 6).

Comparative transcriptomics between trees from the Equatorial and Subtropical sites

To identify environmental forces associated with variations in gene expression between source sites, we sampled leaves, stems and flowers from trees under uncontrolled field conditions at the end of winter at the Subtropical site and at the beginning of the dry season at the Equatorial site (Supplemental Table 1). We observed a consistency in transcript expression levels from leaves and stems among plants from the same sampling site (Supplemental Fig. 7a and 7b). However, transcript expression levels in flowers were not concordant among samples from the same origin (Supplemental Fig. 7c), leading to the identification of only one DET; thus, we did not include flowers in the subsequent analyses (Supplemental Fig. 7f). Contrastingly, we identified 1,837 and 904 transcripts showing significantly different (FDR < 0.05) relative abundance in leaves and stems, respectively,

between Equatorial and Subtropical samples (Supplemental Fig. 7d and 7e). Among the total

2,741 DETs, 1,150 (41.91%) were putative protein-coding transcripts.

The assignment of transcripts to GO terms was possible for 25,184 (83.31%) of

30,227 putative protein-coding sequences, allowing GO enrichment analyses of the DETs.

GO enrichment analyses were focused on biological processes potentially regulating the

responses of *A. schaueriana* to the contrasting climatic variables in the Equatorial and

Subtropical sites (Table 1, Supplemental Fig. 1). Analyses were conducted separately for

leaves and stems and for each of two sets of DETs: one showed higher expression levels in

Equatorial than in Subtropical samples (which we refer to these as DET-Eq) and the other

showed higher abundance in Subtropical than in Equatorial samples (which are referred as

DET-St). The enriched processes among the sets of DET included photosynthesis; plant

responses to UV, temperature stimulus and water stress; cell wall biosynthesis and cellular

respiration (Supplemental Tables 7-11, Supplemental Fig. 7i-l).

Photosynthesis: Among the DET-St, we observed various putative genes participating

in the biosynthesis of the photosynthetic apparatus, chlorophyll and photoreceptors; the

function of electron transporters and chloroplast movement coordination. Contrastingly, the

DET-Eq set showed enrichment in transcripts similar to proteins required for disassembling

the light-harvesting complex of photosystem II in thylakoid membranes and for triggering

chlorophyll degradation (Park et al., 2007) (Supplemental Table 11).

Response to UV: Both the DET-St and DET-Eq sets showed enrichment in functions

related to the response to UV-light, however, the transcript annotations differed between these

sets. The DET-St set included putative UV-B protectants and genes involved in UV-B-

induced antioxidants biosynthesis, such as plastid copper/zinc superoxide dismutases,

photosystem II repair proteins, and L-ascorbic acid. In contrast, the DET-Eq set showed

enrichment of transcripts associated with photo-oxidative damage reduction and the positive

regulation of anthocyanin biosynthesis in response to UV. Antioxidants induced by UV

irradiation (Myouga et al., 2008), such as putative iron superoxide dismutases and pyridoxine

biosynthesis genes, were among the DET-Eq (Supplemental Table 11).

Response to temperature: In the DET-St set, we observed putative genes presenting

critical roles in chloroplast protein translation during cold acclimation and that provide

tolerance to chilling temperatures (Goulas et al., 2006; S. Wang et al., 2016). For instance,

transcripts similar to the *GLYCINE-RICH RNA-BINDING PROTEIN* (*RZ1A*), which has a

chaperone activity during cold acclimation (Kim, Kim, & Kang, 2005), and to the cold-

inducible ATP-dependent *DNA HELICASE* ATSGS1, required for DNA damage-repair

(Hartung, Suer, & Puchta, 2007). Interestingly, DET-St included a putative *AGAMOUS-LIKE*

24 (*AGL24*) transcription factor, which is involved in vernalisation-induced floral transition

(Michaels et al., 2003). Contrastingly, various transcripts similar to heat shock-inducible

chaperones and to *ADENINE NUCLEOTIDE ALPHA HYDROLASE-LIKE PROTEIN*

(*AT3G53990*), involved in chaperone-mediated protein folding (Jung et al., 2015), were

among the DET-Eq set, potentially enhancing tolerance to heat in equatorial plants.

Additionally, a transcript similar to the *ETHYLENE-RESPONSIVE ELEMENT BINDING*

PROTEIN (*RAP2.3*), which confers resistance to heat and hydrogen peroxide (Ogawa et al.,

2005), was observed in this group (Supplemental Table 11).

Response to water stress: Transcripts associated with the response and tolerance to

water deficits and with cellular ion homeostasis and osmotic adjustment were enriched among

DET-Eq. For instance, a transcript similar to the *ETHYLENE-RESPONSIVE*

TRANSCRIPTION FACTOR (*RAP2.4*), which regulates the expression of several drought-

responsive genes, including aquaporins (Lin, Park, & Wang, 2008; Rae, Lao, & Kavanagh,

2011), was identified in DET-Eq. Accordingly, a putative aquaporin *PLASMA MEMBRANE*

INTRINSIC PROTEIN (PIP1;4) (Alexandersson et al., 2005) was also found in this set. We

observed in DET-Eq putative genes participating in the synthesis of raffinose, an

osmoprotective soluble trisaccharide (Nishizawa, Yabuta, & Shigeoka, 2008), and also

transcripts similar to osmosensitive ion channels belonging to the *EARLY-RESPONSIVE TO*

DEHYDRATION STRESS FAMILY. Correspondingly, we observed an ion channel protein,

SLAC1 HOMOLOGUE 3 (*SLAH3*), required for stomatal closure induced by drought stress

(A. Zhang et al., 2016), and the putative *NINE-CIS-EPOXYCAROTENOID DIOXYGENASE 3*

(*NCED3*), which increases plant resistance to water deficit through the accumulation of

abscisic acid (ABA), leading to stomatal closure. Possibly as a consequence of decreased

stomatal conductance, a transcript similar to the *ALANINE-GLYOXYLATE*

AMINOTRANSFERASE 2 HOMOLOG 3 (*AT3G08860*), which plays a central role in

photorespiration (Liepman & Olsen, 2003), showed higher expression in Equatorial than in

Subtropical samples (Supplemental Table 11).

Cell wall biosynthesis: Transcripts similar to 33 distinct proteins and transcription

factors that play central roles in the biosynthesis and deposition of cell wall components, such

as cellulose, hemicellulose, lignin and pectin, were identified among DET-Eq (Supplemental

Table 10).

Cellular respiration: DET-Eq included one putative enzyme from the tricarboxylic

acid cycle encoded in the nuclear genome, *ACONITASE 3* (*ACO3*), which converts citrate to

isocitrate, and several mitochondrial genes encoding subunits of NADH dehydrogenases,

ATP-synthases and cytochrome C oxidases (Supplemental Table 11).

We confirmed the results obtained in the computational analyses of RNA-Seq data by

qRT-PCR of ten DET detected across all leaf samples (Supplemental Fig. 8, Supplemental

Table 12, Supplemental Results).

Detection of SNP with signs of selection

To complement the analyses of differential gene expression, which could result from

plasticity and adaptive selection (Wolf, Lindell, & Backstrom, 2010), we searched for gene

sequence variation among trees sampled along the Atlantic coast of South America (Fig. 1,

Supplemental Table 2). After quality filtering of the sequenced data, we selected 77

individuals without evidence of interspecific hybridisation with *A. germinans* for downstream

analyses. We identified a set of 6,170 high-quality unlinked biallelic SNP loci with a minor

allele frequency $\geq$ 0.05 and $\geq$ 8x coverage. The overall genetic structure corroborated a

previous study based on putatively neutral microsatellite loci (Mori et al., 2015), dividing the

samples into two main groups: north and south of the NEESA (Supplemental Fig. 9).

We observed 122 loci showing significant departures from neutral expectations of

interpopulational divergence, as conservatively detected (Ahrens et al., 2018) by both pcadapt

and LOSITAN. Fifteen of these loci with putative signs of selection were aligned to *A.*

schaueriana transcripts that were similar to gene models in *A. thaliana* and *S. indicum*

(Supplemental Table 13), enabling screening for their potential functional roles. However,

five of the reference proteins did not have informative functions described for the model

plant, hindering inferences regarding their function. Conversely, among the remaining

annotated candidates, we found five putative genes involved in processes related to the

contrasting equatorial and subtropical environments (Fig. 4). One candidate locus was

detected in the putative transcription factor *RAP2.4,* which is induced in response to water and

salt stress and regulates developmental processes mediated by light intensity (Lin et al., 2008)

and the expression of aquaporins (Rae et al., 2011), which plays a role in plant water

homeostasis. Two other candidates showed similarity with the transcription factors *ZINC-*

FINGER PROTEIN 1 (*ZFN1*), involved in the regulation of the expression of several water

stress genes (Sakamoto, Araki, Meshi, & Iwabuchi, 2000)*, and *HYPOXIA RESPONSE*

ATTENUATOR (*HRA1*), strongly induced in response to low oxygen levels (Giuntoli et al.,

2014). A putative *UDP GLUCOSYL TRANSFERASE*, an enzyme that catalyses the final step

of anthocyanin biosynthesis wherein pigmentation is produced (Tohge et al., 2005), also

showed evidence of positive selection. Additionally, one candidate locus was found in a

transcript similar to a *TETRATRICOPEPTIDE REPEAT PROTEIN* (*AT2G20710*, *TPR*),

which might play a role in the biogenesis of the photosynthetic apparatus (Bohne,

Schwenkert, Grimm, & Nickelsen, 2016).

Discussion

We used complementary ecological and molecular approaches to study non-neutral

phenotypic and genetic divergences as well as the contribution of contrasting environmental

forces between two populations of the mangrove species *Avicennia schaueriana*. The genetic

structure previously detected with a few microsatellite loci between forests occurring north

and south of the NEESA (Mori et al., 2015) was confirmed using thousands of genome-wide

SNP (Supplemental Fig. 9). Equatorial plants showed morphophysiological and

transcriptomic signals that appear to minimise the effects of drought, high light and heat,

whereas traits observed in Subtropical plants suggest a maximisation of carbon assimilation,

which may be beneficial under low temperature and reduced light regime (Fig. 2-3,

Supplemental Table 11). Additionally, putative signs of selection were identified in

transcripts associated with contrasting climate variables between equatorial and subtropical

latitudes (Fig. 4).

Traits exhibited by Equatorial plants relative to Subtropical plants in the common

garden experiment, such as reduced leaf size, smaller leaf angle, higher levels of red light-

reflecting pigments, narrower vessels and lower rates of stomatal conductance, limit carbon

acquisition (Reef & Lovelock, 2015) and may have imposed constraints to carbon gain in

Equatorial plants, which also accumulated less biomass (Fig. 2-3, Supplemental Fig. 3-4). In

contrast, such characteristics allow plants to maintain suitable leaf temperature for

photosynthesis while reducing both UV exposure and water loss through the minimisation of

evaporative cooling (Reef & Lovelock, 2015; Steyn, Wand, Holcroft, & Jacobs, 2002). We

argue that the prevalence of these traits among Equatorial samples may be advantageous in

their natural environment, especially during the dry season (from August to December),

which presents high light intensity, frequently combined with high temperature (> 30 °C) and

air humidity below 70% (Table 1, Supplemental Fig. 1). Under high evaporative demand and

saline intertidal soils, water acquisition has an elevated energetic cost, representing strong

pressure in favour of water-saving adaptations (Reef & Lovelock, 2015). Accordingly,

Equatorial plants also showed lower transpiration rates than did Subtropical plants in the

common garden (Supplemental Fig. 4). In addition, 63% of the six-month-old Equatorial

plants started flowering from July-August (Supplemental Fig. 4g), which is consistent with

phenological observations reported for *A. schaueriana* in equatorial sites (Menezes, Berger, &

Mehlig, 2008). However, we found no previous records of six-month-old flowering plants in

the literature. Although a flowering peak is observed in August in southern subtropical forests

(De Alvarenga, Botosso, & Soffiatti, 2017), Subtropical plants did not flower during the

glasshouse experiment. Early flowering is a phenotype with complex genetic components and

is rarely studied in non-model organisms; however it is renowned as an adaptive mechanism

for maximising the chance of reproduction under stress (Kazan & Lyons, 2016).

Contrastingly, Subtropical plants showed higher stomatal conductance and

transpiration rates, higher levels of green light-reflecting pigments, larger leaf area, wider leaf

lamina angle and larger xylem vessel diameter than did Equatorial plants in the common

garden experiment (Fig. 2-3, Supplemental Fig. 3-4). These characteristics enhance light

energy absorbance and carbon acquisition at the expense of greater water loss and higher

cavitation risk (Carlson, Holsinger, & Prunier, 2011; Stuart, Choat, Martin, Holbrook, & Ball,

2006). These traits may compensate for declines in net primary production in higher-latitude

environments (Saenger & Snedaker, 1993) that result from restrictions in temperature and

solar irradiance (Table 1, Supplemental Fig. 1). We argue that the intensity of cold events in

southern subtropical populations of *A. schaueriana* is likely insufficient to favour the

selection of freezing-tolerant individuals, in contrast to results reported for *A. germinans* at its

northernmost distribution limit on the Atlantic coast of North America (Cavanaugh et al.,

2014). At the southernmost range edge of *A. schaueriana,* the minimum air temperature does

not drop below 0 □ (Table 1, Supplemental Fig. 1), which is higher than the expected

mangrove physiological threshold (Osland et al., 2017). Additionally, the small population

size of *A. schaueriana* at this location (Soares et al., 2012) and the arrival of maladapted

propagules from northerly populations likely reduce the potential strength of selection

favouring cold-adapted individuals.

Functional interpopulation divergence at the molecular level was evident under field

conditions. Comparative transcriptomics of trees sampled in their native habitats corroborated

the suggested effects of environmental variation in light availability, temperature and water

stress on the phenotypic divergence observed in the common garden experiment. The

transcriptomic profiles obtained at the beginning of the dry season in the Equatorial site and at

the end of winter in the Subtropical site (Supplemental Table 1) showed an enrichment of

DETs involved in photosynthesis, cellular respiration, cell wall biosynthesis and plant

responses to water stress, temperature and UV light (Supplemental Fig. 7i-l). The adaptive

relevance of these biological processes in the field was highlighted through the identification

and functional annotation of SNPs putatively under natural selection from populations along

the *A. schaueriana* geographic distribution (Supplemental Table 13). In the following

subsections, we integrate information derived from three independent approaches explored in

this work.

***Water stress as a key selective pressure in equatorial populations of* A. schaueriana**

The increased levels of transcripts similar to heat-shock proteins, to drought-induced

ion transporters, and genes that enhance heat tolerance and play central roles in stomatal

closure and photorespiration provided multiple lines of evidence of water stress in Equatorial

samples. Additionally, Equatorial plants exhibited higher expression of aquaporins and genes

involved in the accumulation of organic solutes than did Subtropical plants. These

investments improve tolerance to drought (Nishizawa et al., 2008) by lowering cell water

potential and actively transporting water through proteins in the cell membrane rather than

using passive apoplastic transport (Krishnamurthy et al., 2014; Reef & Lovelock, 2015).

Enhanced rigidity of cells reduces risks of damage during dehydration and rehydration,

thereby improving resistance to high extra-cellular osmotic pressure (Gall et al., 2015). Thus,

we argue that the higher expression of several transcripts associated with cell wall

biosynthesis and cell wall thickening in the Equatorial samples may indicate plant responses

to water stress. Further evidences of the relevance of water stress in plants at the Equatorial

site were highlighted by the identification of sequence divergence between northerly and

southerly populations in two putative osmotic stress-responsive regulatory transcription

factors (*RAP2.4* and *ZFN1*). These findings corroborate with the divergence in traits involved

in water balance between plants from Equatorial and Subtropical sites, such as vessel

diameter, leaf size, leaf angle and transpiration and stomatal conductance rates (Fig. 2,

Supplemental Fig. 3-4).

***Latitudinal variation in light quality and intensity may shape functional diversity in* A.**

schaueriana

Contrasting seasonal fluctuations in photoperiod and in light quality between low and

high latitudes (Pecot, Horsley, Battaglia, & Mitchell, 2005) (Table 1) likely influence the

differential expression observed in putative UV-inducible antioxidant and photodamage repair

genes. The adaptive relevance of these findings is supported by the sign of natural selection

found in a transcript similar to *UDP-GLUCOSYL TRANSFERASE*, a key enzyme to

anthocyanin biosynthesis, which confers protection to UV-B. Divergent morphological traits

between Equatorial and Subtropical plants grown in the common garden experiment,

including leaf inclination angle and stem light reflectance (Fig. 2 and 3) provide additional

insights into *A. schaueriana* light-related adaptations.

Low water and low light availability presumably affect photosynthesis and cellular

respiration

In response to abiotic stress conditions such as drought, heat and high light, plants

optimise the use of light energy and minimise photooxidative damage by reducing the

photosynthetic activity via the repression of light-harvesting and photosystem-component

genes (Bray, 2004; Han et al., 2009; Kimura et al., 2003; Moumeni et al., 2011; D. Wang et

al., 2011; C. Zhang et al., 2015). We argue that the lower expression of photosynthesis genes

in Equatorial than in Subtropical samples likely is further indicative of the role of water stress

in shaping divergent phenotypes in the field, but it may also result in the enhanced absorption

of light energy in Subtropical plants. Accordingly, we identified increased expression of

transcripts associated with chlorophyll biosynthesis among the DET-St set members and with

chlorophyll breakdown and leaf senescence among DET-Eq. Chlorophyll degradation

followed by leaf senescence allows the remobilisation of nutrients and reduces the water loss

through transpiration, contributing to water balance and drought tolerance (Munné-Bosch &

Alegre, 2004). Additionally, we detected a putative signature of selection in a transcript

similar to a *TPR* protein, required for chlorophyll synthesis and for the biogenesis of the

photosynthetic apparatus (Bohne et al., 2016). We suggest that differential seasonality in light

and water availability between subtropical and equatorial latitudes may be involved in the

divergence of non-neutral variability in the species. Corroborating the results obtained from

the genomic approaches, we observed functional trait divergence related to water use and

light energy absorbance in plants from contrasting latitudes cultivated under the same

conditions (Fig. 2-3, Supplemental Fig. 3-4).

Mitochondrial activity is strongly connected to photosynthesis and chloroplasts

function since it generates ATP for carbohydrate synthesis, plays a role in protecting

chloroplasts against photoinhibition, participates in dissipating reducing equivalents and

exchanges metabolites with chloroplasts during photorespiration (Millar, Whelan, Soole, &

Day, 2011; Raghavendra & Padmasree, 2003). Cellular respiration also provides ATP and

carbon compounds for secondary metabolism, playing fundamental roles in responses to

abiotic stresses, including drought (Atkin & Macherel, 2009; Millar et al., 2011). We suggest

that higher expression levels of cellular respiration genes in Equatorial than in Subtropical

samples may be a consequence of the reduced expression of photosynthesis genes and the

enhanced energetic demand caused by water stress.

Tidal amplitude variation with latitude as a diverging force along the Atlantic coast of

South America

Tidal amplitude is markedly reduced with increasing latitude along the Atlantic coast

of South America, ranging from greater than 4 m at low latitudes to less than 1 m at high

latitudes (Schaeffer-Novelli, Cintrón-Molero, Adaime, & de Camargo, 1990). The wide

variation in tidal amplitude exposes trees to varied durations of hypoxia. The identification of

a candidate SNP locus with a putative sign of selection in a transcription factor induced by

oxygen deprivation (*HRA1*) may indicate that differential tidal variation acts as a diverging

selective force between northerly equatorial and southerly subtropical populations of *A.*

schaueriana. The *HRA1* putative homolog also showed 1.75-fold higher expression in

Subtropical leaves relative to that in Equatorial leaves under field conditions. However, we

did not detect evidences at the phenotypic level in the common garden experiment suggesting

the relevance of this environmental variable.

Climate change and conservation implications of the results

The functional divergence described herein might differentially affect the sensitivity

of *A. schaueriana* populations to a rapidly changing climate. Although there is no evidence of

mangrove expansion at its southernmost range limit in the Americas (Soares et al., 2012),

researchers suggest that subtropical populations could expand polewards in the near future as

a result from the increased air and ocean temperatures and CO_2 concentrations (Godoy & De

Lacerda, 2015). We expect that the more acquisitive traits, in terms of gene expression in the

field and in morphophysiology in the common garden experiment, exhibited by plants from

the subtropical range edge may indeed favour growth under increased temperatures and

rainfall (Pachauri & Meyer, 2014). However, due to the low tidal variation (< 1 m) observed

at the southernmost mangrove distribution edge along the Atlantic coast of South America, a

greater relocation of forests will be required for species to keep pace with the rising sea level

(Ellison, 2015). In this context, dense human coastal occupations combined with the narrow

intertidal zones, characteristic of this region, make mangroves at higher latitudes more

vulnerable to habitat loss than in the equatorial region of the Atlantic coast of South America.

We expect that in contrast, despite having wider coastal plains potentially available for

expansion, we expect that Equatorial populations of *A. schaueriana* might be threatened by

the increased mean temperatures and decreased precipitation during the El Niño-Southern

Oscillation events (Pachauri & Meyer, 2014). Increased leaf temperature stimulates

respiration (Heskel et al., 2016) and photorespiration (Jordan & Ogren, 1984) and might

offset the benefits in carbon acquisition caused by increased CO_2 concentration (Drake,

Gonzàlez-Meler, & Long, 1997). The critical temperature threshold for photosynthesis is

likely to be overcome more frequently in the near future, possibly reducing carbon

assimilation and productivity (Saenger & Moverly, 1985) and, in extreme cases, causing

biomass loss triggered by cavitation or carbon starvation (Doughty et al., 2015; Rowland et

al., 2015).

For the definition of short-term mangrove conservation plans, such as the reforestation

or restoration of degraded areas on the Atlantic coast of South America, we recommend that

populations occurring north and south of the NEESA should warrant attention as distinct

conservation management units(Moritz, 1994).

Conclusions

Based on the combined analysis of our results, we argue that carbon acquisition in plants in equatorial region may be limited by the longer exposure to hypoxia, the higher vapour pressure deficit (VPD) and the higher solar irradiance, especially during the hot and dry season. Furthermore, we argue that in subtropical regions, limitations in carbon gain may result from the lower solar irradiance levels, the lower temperature and the shorter photoperiod during winter (Fig. 4). These environmental differences between equatorial and subtropical latitudes presumably control gene expression in the field, and remarkably, they may have shaped both allele frequencies in genes responding to these variables and morphophysiological traits observed in *A. schaueriana* individuals (Fig. 4). The emergence of this divergence is facilitated by the limited gene flow identified between populations north and south of the NEESA, possibly driven by the movements of the major ocean currents that are active along the Atlantic coast of South America (Fig. 1). Because a similar north-south structure of neutral diversity is also observed for other mangrove and mangrove-associated species on the Atlantic coast of South America (Francisco et al., 2018; Mori et al., 2015; Takayama et al., 2008), it is plausible that the environmental drivers of adaptive divergence in *A. schaueriana* play roles in the divergence of other species.

In addition to revealing that northerly and southerly populations of *A. schaueriana* are genetically and functionally distinct units (Wee et al., 2018), we have provided an in-depth empirical evaluation of the intraspecific variation of a long-living, non-model tree. These results should allow clearer predictions of how *A. schaueriana* and potentially other coastal plants may respond to current global climate changes by accounting for phenotypically

(Moran et al., 2016) and genetically (Ikeda et al., 2017) informed mangrove distribution

modelling.

Acknowledgements

We thank Professors Dr. Michel A. G. Vincentz and Dr. Juliana L. S. Mayer for

advising and making available their laboratory for our training and performing of RNA

extraction and slide preparation for the analyses of wood anatomy. We acknowledge Ilmarina

C. de Menezes and Stephanie K. Bajay for assistance in the field. We thank Stephanie K.

Bajay for providing assistance in the germination of the propagules used in the common

garden experiment. And we thank Greg Mellers and Alessandro Alves-Pereira for critically

reading the manuscript. M.V.C and G.M.M. received fellowships from the São Paulo

Research Foundation (FAPESP 2013/26793-7, 2013/08086-1 and 2014/22821-9). M.V.C.

received a fellowship from the Coordination for the Improvement of Higher Education

Personnel (CAPES 8084/2015-07). G.M.M. received a grant from the Brazilian National

Council for Scientific and Technological Development (CNPq 448286/2014-9). A.P.S.

received a research fellowship from CNPq (309661/2014-5).

References
Ahrens, C. W., Rymer, P. D., Stow, A., Bragg, J., Dillon, S., Umbers, K. D. L., & Dudaniec,
R. Y. (2018). The search for loci under selection: trends, biases and progress. *Molecular*
Ecology. doi:doi.org/10.1111/mec.14549
Albert, C. H., Grassein, F., Schurr, F. M., Vieilledent, G., & Violle, C. (2011). When and how
should intraspecific variability be considered in trait-based plant ecology? *Perspectives*
in Plant Ecology, Evolution and Systematics, *13*(3), 217–225.
doi:10.1016/j.ppees.2011.04.003
Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry
in unrelated individuals. *Genome Research*, *19*(9), 1655–1664.
doi:10.1101/gr.094052.109
Alexandersson, E., Fraysse, L., Sjövall-Larsen, S., Gustavsson, S., Fellert, M., Karlsson, M.,
… Kjellbom, P. (2005). Whole gene family expression and drought stress regulation of
aquaporins. *Plant Molecular Biology*, *59*(3), 469–484. doi:10.1007/s11103-005-0352-1
Alongi, D. M. (2015). The Impact of Climate Change on Mangrove Forests. *Curr. Clim.*

Change Rep., *4*, 516. doi:10.1007/s40641-015-0002-x

Alvares, C. A., Stape, J. L., Sentelhas, P. C., Gonçalves, J. L. de M., & Sparovek, G. (2013).
Köppen's climate classification map for Brazil. *Meteorologische Zeitschrift*, *22*(6), 711–
728. doi:10.1127/0941-2948/2013/0507

Andrews, K. R., Good, J. M., Miller, M. R., Luikart, G., & Hohenlohe, P. A. (2016).
Harnessing the power of RADseq for ecological and evolutionary genomics. *Nature*
Reviews Genetics, *17*(2), 81–92. doi:10.1038/nrg.2015.28

Antao, T., Lopes, A., Lopes, R. J., Beja-Pereira, A., & Luikart, G. (2008). LOSITAN: A
workbench to detect molecular adaptation based on a F st -outlier method. *BMC*
Bioinformatics, *9*(1), 1–5. doi:10.1186/1471-2105-9-323

Atkin, O. K., & Macherel, D. (2009). The crucial role of plant mitochondria in orchestrating
drought tolerance. *Annals of Botany*, *103*(4), 581–597. doi:10.1093/aob/mcn094

Barrett, R. D. H., & Hoekstra, H. E. (2011). Molecular spandrels: tests of adaptation at the
genetic level. *Nature Reviews Genetics*, *12*(11), 767–780. doi:10.1038/nrg3015

Berardini, T. Z., Reiser, L., Li, D., Mezheritsky, Y., Muller, R., Strait, E., & Huala, E. (2015).
The Arabidopsis Information Resource □: Making and Mining the " Gold Standard "
Annotated Reference Plant Genome. *Genesis*, *53*, 474–485. doi:10.1002/dvg.22877

Blake, J. A., Christie, K. R., Dolan, M. E., Drabkin, H. J., Hill, D. P., Ni, L., … Westerfeld,
M. (2015). Gene ontology consortium: Going forward. *Nucleic Acids Research*, *43*,
D1049–D1056. doi:10.1093/nar/gku1179

Bohne, A. V., Schwenkert, S., Grimm, B., & Nickelsen, J. (2016). *Roles of Tetratricopeptide*
Repeat Proteins in Biogenesis of the Photosynthetic Apparatus. *International Review of*
Cell and Molecular Biology (Vol. 324). Elsevier Inc. doi:10.1016/bs.ircmb.2016.01.005

Bray, E. A. (2004). Genes commonly regulated by water-deficit stress in Arabidopsis
thaliana. *Journal of Experimental Botany*, *55*(407), 2331–2341. doi:10.1093/jxb/erh270

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden,
T. L. (2009). BLAST+: architecture and applications. *BMC Bioinformatics*, *10*, 421.
doi:10.1186/1471-2105-10-421

Carlson, J. E., Holsinger, K. E., & Prunier, R. (2011). Plant responses to climate in the cape
floristic region of South Africa: Evidence for adaptive differentiation in the proteaceae.
Evolution, *65*(1), 108–124. doi:10.1111/j.1558-5646.2010.01131.x

Cavanaugh, K. C., Kellner, J. R., Forde, A. J., Gruner, D. S., Parker, J. D., Rodriguez, W., &
Feller, I. C. (2014). Poleward expansion of mangroves is a threshold response to
decreased frequency of extreme cold events. *Proceedings of the National Academy of*
Sciences of the United States of America, *111*(2), 723–7. doi:10.1073/pnas.1315800111

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., … Durbin,
R. (2011). The variant call format and VCFtools. *Bioinformatics*, *27*(15), 2156–2158.
doi:10.1093/bioinformatics/btr330

De Alvarenga, A. M. S. B., Botosso, P. C., & Soffiatti, P. (2017). Stem growth and phenology
of three subtropical mangrove tree species. *Brazilian Journal of Botany*.
doi:10.1007/s40415-017-0397-9

Doughty, C. E., Metcalfe, D. B., Girardin, C. A. J., Amézquita, F. F., Cabrera, D. G., Huasco,
W. H., … Malhi, Y. (2015). Drought impact on forest carbon dynamics and fluxes in
Amazonia. *Nature*, *519*(7541), 78–82. doi:10.1038/nature14213

Drake, B. G., Gonzàlez-Meler, M. A., & Long, S. P. (1997). More efficient plants: a
consequence of rising atmospheric CO2? *Annual Review of Plant Physiology and Plant*
Molecular Biology, *48*(1), 609–639. doi:10.1146/annurev.arplant.48.1.609

Duke, N., Kovacs, J., Griffiths, A., Preece, L., J. E. Hill, D., Van Oosterzee, P., … Burrows,

D. (2017). *Large-scale dieback of mangroves in Australia. Marine and Freshwater Research.* doi:10.1071/MF16322

Ellison, J. C. (2015). Vulnerability assessment of mangroves to climate change and sea-level rise impacts. *Wetlands Ecology and Management*, *23*(2), 115–137. doi:10.1007/s11273-014-9397-8

Finn, R. D., Bateman, A., Clements, J., Coggill, P., Eberhardt, R. Y., Eddy, S. R., … Punta, M. (2014). Pfam: The protein families database. *Nucleic Acids Research*, *42*(D1), 222–230. doi:10.1093/nar/gkt1223

Forsythe, W. C., Rykiel, E. J., Stahl, R. S., Wu, H. i., & Schoolfield, R. M. (1995). A model comparison for daylength as a function of latitude and day of year. *Ecological Modelling*, *80*(1), 87–95. doi:10.1016/0304-3800(94)00034-F

Francisco, P. M., Mori, G. M., Alves, F. M., Tambarussi, E., & Souza, A. P. De. (2018). Population genetic structure, introgression, and hybridization in the genus Rhizophora along the Brazilian coast. *Ecology and Evolution*, (January), 1–14. doi:10.1002/ece3.3900

Gall, H. Le, Philippe, F., Domon, J.-M., Gillet, F., Pelloux, J., & Rayon, C. (2015). Cell Wall Metabolism in Response to Abiotic Stress. *Plants*, *4*(1), 112–166. doi:10.3390/plants4010112

Giuntoli, B., Lee, S. C., Licausi, F., Kosmacz, M., Oosumi, T., van Dongen, J. T., … Perata, P. (2014). A Trihelix DNA Binding Protein Counterbalances Hypoxia-Responsive Transcriptional Activation in Arabidopsis. *PLoS Biology*, *12*(9). doi:10.1371/journal.pbio.1001950

Godoy, M. D. P., & De Lacerda, L. D. (2015). Mangroves Response to Climate Change: A Review of Recent Findings on Mangrove Extension and Distribution. *Annals of the Brazilian Academy of Sciences*, *87*(2), 651–667. doi:10.1590/0001-3765201520150055

Goulas, E., Schubert, M., Kieselbach, T., Kleczkowski, L. A., Gardestro, P., & Schro, W. (2006). The chloroplast lumen and stromal proteomes of Arabidopsis thaliana show differential sensitivity to short- and long-term exposure to low temperature. *The Plant Journal*, *47*, 720–734. doi:10.1111/j.1365-313X.2006.02821.x

Han, F., Chen, H., Li, X. J., Yang, M. F., Liu, G. S., & Shen, S. H. (2009). A comparative proteomic analysis of rice seedlings under various high-temperature stresses. *Biochimica et Biophysica Acta*, *1794*(11), 1625–1634. doi:10.1016/j.bbapap.2009.07.013

Hartung, F., Suer, S., & Puchta, H. (2007). Two closely related RecQ helicases have antagonistic roles in homologous recombination and DNA repair in Arabidopsis thaliana. *Proceedings of the National Academy of Sciences*, *104*(47), 18836–18841. doi:10.1073/pnas.0705998104

Hereford, J. (2009). A Quantitative Survey of Local Adaptation and Fitness Trade☐Offs. *The American Naturalist*, *173*(5), 579–588. doi:10.1086/597611

Heskel, M. A., O'Sullivan, O. S., Reich, P. B., Tjoelker, M. G., Weerasinghe, L. K., Penillard, A., … Atkin, O. K. (2016). Convergence in the temperature response of leaf respiration across biomes and plant functional types. *Proceedings of the National Academy of Sciences*, *113*(14), 3832–3837. doi:10.1073/pnas.1520282113

Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, *25*(15), 1965–1978. doi:10.1002/joc.1276

Huang, J., Lu, X., Zhang, W., Huang, R., Chen, S., & Zheng, Y. (2014). Transcriptome Sequencing and Analysis of Leaf Tissue of Avicennia marina Using the Illumina Platform. *PLoS ONE*, *9*(9), e108785. doi:10.1371/journal.pone.0108785

Ikeda, D. H., Max, T. L., Allan, G. J., Lau, M. K., Shuster, S. M., & Whitham, T. G. (2017).
Genetically informed ecological niche models improve climate change predictions.
Global Change Biology, *23*(1), 164–176. doi:10.1111/gcb.13470
Jombart, T., Devillard, S., & Balloux, F. (2010). Discriminant analysis of principal
components: a new method for the analysis of genetically structured populations. *BMC*
Genetics, *11*(1), 94. doi:10.1186/1471-2156-11-94
Jordan, D., & Ogren, W. (1984). The CO2/O2 specificity of ribulose 1,5-bisphosphate
carboxylase/oxygenase. *Planta*, *161*(4), 308–313. doi:10.1007/BF00398720
Jung, Y. J., Melencion, S. M. B., Lee, E. S., Park, J. H., Alinapon, C. V., Oh, H. T., … Lee, S.
Y. (2015). Universal Stress Protein Exhibits a Redox-Dependent Chaperone Function in
Arabidopsis and Enhances Plant Tolerance to Heat Shock and Oxidative Stress.
Frontiers in Plant Science, *6*, 1–11. doi:10.3389/fpls.2015.01141
Kawecki, T. J., & Ebert, D. (2004). Conceptual issues in local adaptation. *Ecology Letters*,
7(12), 1225–1241. doi:10.1111/j.1461-0248.2004.00684.x
Kazan, K., & Lyons, R. (2016). The link between flowering time and stress tolerance. *Journal*
of Experimental Botany, *67*(1), 47–60. doi:10.1093/jxb/erv441
Kim, Y. O., Kim, J. S., & Kang, H. (2005). Cold-inducible zinc finger-containing glycine-rich
RNA-binding protein contributes to the enhancement of freezing tolerance in
Arabidopsis thaliana. *Plant Journal*, *42*(6), 890–900. doi:10.1111/j.1365-
313X.2005.02420.x
Kimura, M., Yamamoto, Y. Y., Seki, M., Sakurai, T., Sato, M., Abe, T., … Matsui, M.
(2003). Identification of Arabidopsis Genes Regulated by High Light – Stress Using
cDNA Microarray. *Photochemistry and Photobiology*, *77*(2), 226–233.
doi:https://doi.org/10.1562/0031-8655
Kjerfve, B., Perillo, G. M. E., Gardner, L. R., Rine, J. M., Dias, G. T. M., & Mochel, F. R.
(2002). Chapter Twenty Morphodynamics of muddy environments along the Atlantic
coasts of North and South America. *Proceedings in Marine Science*, *4*(C), 479–532.
doi:10.1016/S1568-2692(02)80094-8
Krishnamurthy, P., Jyothi-Prakash, P. A., Qin, L., He, J., Lin, Q., Loh, C. S., & Kumar, P. P.
(2014). Role of root hydrophobic barriers in salt exclusion of a mangrove plant
Avicennia officinalis. *Plant, Cell and Environment*, *37*(7), 1656–1671.
doi:10.1111/pce.12272
Krishnamurthy, P., Mohanty, B., Wijaya, E., Lee, D., Lim, T., Lin, Q., … Kumar, P. P.
(2017). Transcriptomics analysis of salt stress tolerance in the roots of the mangrove
Avicennia officinalis. *Nature Scientific Reports*, *7*, 1–19. doi:10.1038/s41598-017-
10730-2
Langmead, B., Trapnell, C., Pop, M., & Salzberg, S. (2009). Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. *Genome Biol.*, *10*(3), R25.
doi:10.1186/gb-2009-10-3-r25
Li, W., & Godzik, A. (2006). Cd-hit: A fast program for clustering and comparing large sets
of protein or nucleotide sequences. *Bioinformatics*, *22*(13), 1658–1659.
doi:10.1093/bioinformatics/btl158
Liepman, A. H., & Olsen, L. J. (2003). Alanine aminotransferase homologs catalyze the
glutamate:glyoxylate aminotransferase reaction in peroxisomes of Arabidopsis. *Plant*
Physiology, *131*(January), 215–27. doi:10.1104/pp.011460
Lin, R. C., Park, H. J., & Wang, H. Y. (2008). Role of Arabidopsis RAP2.4 in regulating
light- and ethylene-mediated developmental processes and drought stress tolerance.
Molecular Plant, *1*(1), 42–57. doi:10.1093/mp/ssm004

Loarie, S. R., Duffy, P. B., Hamilton, H., Asner, G. P., Field, C. B., & Ackerly, D. D. (2009).
The velocity of climate change. *Nature*, *462*(7276), 1052–1055.
doi:10.1038/nature08649

López-Medellín, X., Ezcurra, E., González-Abraham, C., Hak, J., Santiago, L. S., & Sickman,
J. O. (2011). Oceanographic anomalies and sea-level rise drive mangroves inland in the
Pacific coast of Mexico. *Journal of Vegetation Science*, *22*, 143–151.
doi:10.1111/j.1654-1103.2010.01232.x

Lovelock, C. E., Cahoon, D. R., Friess, D. A., Guntenspergen, G. R., Krauss, K. W., Reef, R.,
… Triet, T. (2015). The vulnerability of Indo-Pacific mangrove forests to sea-level rise.
Nature, *526*(7574), 559–563. doi:10.1038/nature15538

Luu, K., Bazin, E., & Blum, M. G. B. (2016). pcadapt: An R package to perform genome
scans for selection based on principal component analysis. *Molecular Ecology*
Resources, *33*, 67–77. doi:10.1111/1755-0998.12592

Lyu, H., Li, X., Guo, Z., He, Z., & Shi, S. (2017). De novo assembly and annotation of the
Avicennia officinalis L. transcriptome. *Marine Genomics*, 2–5.
doi:10.1016/j.margen.2017.07.002

McKee, K. L., Cahoon, D. R., & Feller, I. C. (2007). Caribbean mangroves adjust to rising
sea level through biotic controls on change in soil elevation. *Global Ecology and*
Biogeography, *16*(5), 545–556. doi:10.1111/j.1466-8238.2007.00317.x

McRae, G. J. (1980). A Simple Procedure for Calculating Atmospheric Water Vapor
Concentration. *Journal of the Air Pollution Control Association*, *30*(4), 394–394.
doi:10.1080/00022470.1980.10464362

Menezes, M. P. M. de, Berger, U., & Mehlig, U. (2008). Mangrove vegetation in Amazonia: a
review of studies from the coast of Pará and Maranhão States, north Brazil. *Acta*
Amazonica, *38*(3), 403–420. doi:10.1590/S0044-59672008000300004

Michaels, S. D., Ditta, G., Gustafson-Brown, C., Pelaz, S., Yanofsky, M., & Amasino, R. M.
(2003). AGL24 acts as a promoter of flowering in Arabidopsis and is positively
regulated by vernalization. *Plant Journal*, *33*, 867–874. doi:10.1046/j.1365-
313X.2003.01671.x

Millar, A. H., Whelan, J., Soole, K. L., & Day, D. A. (2011). Organization and Regulation of
Mitochondrial Respiration in Plants. *Annual Review of Plant Biology*, *62*(1), 79–104.
doi:10.1146/annurev-arplant-042110-103857

Moran, E. V., Hartig, F., & Bell, D. M. (2016). Intraspecific trait variation across scales:
Implications for understanding global change responses. *Global Change Biology*, *22*(1),
137–150. doi:10.1111/gcb.13000

Mori, G. M., Zucchi, M. I., & Souza, A. P. (2015). Multiple-Geographic-Scale Genetic
Structure of Two Mangrove Tree Species: The Roles of Mating System, Hybridization,
Limited Dispersal and Extrinsic Factors. *Plos One*, *10*, e0118710.
doi:10.1371/journal.pone.0118710

Moritz, C. (1994). Defining "Evolutionarily Significant Units" for conservation. *Trends in*
Ecology and Evolution, *9*(10), 373–375. doi:10.1016/0169-5347(94)90057-4

Moumeni, A., Satoh, K., Kondoh, H., Asano, T., Hosaka, A., Venuprasad, R., … Kikuchi, S.
(2011). Comparative analysis of root transcriptome profiles of two pairs of drought-
tolerant and susceptible rice near-isogenic lines under different drought stress. *BMC*
Plant Biology, *11*. doi:10.1186/1471-2229-11-174

Munné-Bosch, S., & Alegre, L. (2004). Die and let live: Leaf senescence contributes to plant
survival under drought stress. *Functional Plant Biology*, *31*(3), 203–216.
doi:10.1071/FP03236

Myouga, F., Hosoda, C., Umezawa, T., Iizumi, H., Kuromori, T., Motohashi, R., …
Shinozaki, K. (2008). A Heterocomplex of Iron Superoxide Dismutases Defends
Chloroplast Nucleoids against Oxidative Stress and Is Essential for Chloroplast
Development in Arabidopsis. *The Plant Cell Online*, *20*(11), 3148–3162.
doi:10.1105/tpc.108.061341
Nishizawa, A., Yabuta, Y., & Shigeoka, S. (2008). Galactinol and Raffinose Constitute a
Novel Function to Protect Plants from Oxidative Damage. *Plant Physiology*, *147*(3),
1251–1263. doi:10.1104/pp.108.122465
O'Leary, N. A., Wright, M. W., Brister, J. R., Ciufo, S., Haddad, D., McVeigh, R., … Pruitt,
K. D. (2016). Reference sequence (RefSeq) database at NCBI: Current status, taxonomic
expansion, and functional annotation. *Nucleic Acids Research*, *44*, D733–D745.
doi:10.1093/nar/gkv1189
Ogawa, T., Pan, L., Kawai-Yamada, M., Yu, L.-H., Yamamura, S., Koyama, T., …
Uchimiya, H. (2005). Functional Analysis of Arabidopsis Ethylene-Responsive Element
Binding Protein Conferring Resistance to Bax. *Plant Physiology*, *138*, 1436–1445.
doi:10.1104/pp.105.063586.1436
Oliveira, R. R., Viana, A. J. C., Reátegui, A. C. E., & Vincentz, M. G. A. (2015). An efficient
method for simultaneous extraction of high-quality RNA and DNA from various plant
tissues. *Genetics and Molecular Research*, *14*(4), 18828–18838.
doi:10.4238/2015.December.28.32
Osland, M. J., Feher, L. C., Griffith, K. T., Cavanaugh, K. C., Enwright, N. M., Day, R. H.,
… Rogers, K. (2017). Climatic controls on the global distribution, abundance, and
species richness of mangrove forests. *Ecological Monographs*, *87*(2), 341–359.
doi:10.1002/ecm.1248
Pachauri, R. K., & Meyer, L. A. (Eds.). (2014). *Climate Change 2014: Synthesis Report.*
Contribution of Working Groups I, II and III to the Fifth Assessment Report of the
Intergovernmental Panel on Climate Change. *IPCC*. Geneva, Switzerland: IPCC.
Park, S.-Y., Yu, J.-W., Park, J.-S., Li, J., Yoo, S.-C., Lee, N.-Y., … Paek, N.-C. (2007). The
Senescence-Induced Staygreen Protein Regulates Chlorophyll Degradation. *The Plant*
Cell, *19*(5), 1649–1664. doi:10.1105/tpc.106.044891
Patel, R. K., & Jain, M. (2012). NGS QC toolkit: A toolkit for quality control of next
generation sequencing data. *PLoS ONE*, *7*(2). doi:10.1371/journal.pone.0030619
Pecot, S. D., Horsley, S. B., Battaglia, M. A., & Mitchell, R. J. (2005). The influence of
canopy, sky condition, and solar angle on light quality in a longleaf pine woodland.
Canadian Journal of Forest Research, *35*(6), 1356–1366. doi:10.1139/x05-069
Rae, L., Lao, N. T., & Kavanagh, T. A. (2011). Regulation of multiple aquaporin genes in
Arabidopsis by a pair of recently duplicated DREB transcription factors. *Planta*, *234*(3),
429–444. doi:10.1007/s00425-011-1414-z
Raghavendra, A. S., & Padmasree, K. (2003). Beneficial interactions of mitochondrial
metabolism with photosynthetic carbon assimilation. *Trends in Plant Science*, *8*(11),
546–553. doi:10.1016/j.tplants.2003.09.015
Reef, R., & Lovelock, C. E. (2015). Regulation of water balance in Mangroves. *Annals of*
Botany, *115*(3), 385–395. doi:10.1093/aob/mcu174
Reef, R., Winter, K., Morales, J., Adame, M. F., Reef, D. L., & Lovelock, C. E. (2014). The
effect of atmospheric carbon dioxide concentrations on the performance of the mangrove
Avicennia germinans over a range of salinities. *Physiologia Plantarum*, *154*(358–368).
doi:10.1111/ppl.12289
Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2010). edgeR: a Bioconductor package

for differential expression analysis of digital gene expression data. *Bioinformatics*, *26*(1), 139–140. doi:10.1093/bioinformatics/btp616

Rowland, L., da Costa, A. C. L., Galbraith, D. R., Oliveira, R. S., Binks, O. J., Oliveira, A. A. R., … Meir, P. (2015). Death from drought in tropical forests is triggered by hydraulics not carbon starvation. *Nature*, *528*(7580), 119–122. doi:https://doi.org/10.1038/nature15539

Russello, M. A., Waterhouse, M. D., Etter, P. D., & Johnson, E. A. (2015). From promise to practice: pairing non-invasive sampling with genomics in conservation. *PeerJ*, *3*, e1106. doi:10.7717/peerj.1106

Saenger, P., & Moverly, J. (1985). Vegetative phenology along the Queensland coastline. *Proceedings of the Ecological Society of Australia*, *13*(January 1985), 257–265.

Saenger, P., & Snedaker, S. C. (1993). Pantropical trends in mangrove above-ground biomass and annual litterfall. *Oecologia*, *96*(3), 293–299. doi:10.1007/BF00317496

Sakamoto, H., Araki, T., Meshi, T., & Iwabuchi, M. (2000). Expression of a subset of the Arabidopsis Cys2/His2-type zinc-finger protein gene family under water stress. *Gene*, *248*(1–2), 23–32. doi:10.1016/S0378-1119(00)00133-5

Savolainen, O., Lascoux, M., & Merilä, J. (2013). Ecological genomics of local adaptation. *Nature Reviews Genetics*, *14*(11), 807–20. doi:10.1038/nrg3522

Savolainen, O., Pyhäjärvi, T., & Knürr, T. (2007). Gene Flow and Local Adaptation in Trees. *Annual Review of Ecology, Evolution and Systematics*, *38*, 595–619. doi:10.1146/annurev.ecolsys.38.091206.095646

Sbrocco, E. J., & Barber, P. H. (2013). MARSPEC: ocean climate layers for marine spatial ecology. *Ecology*, *94*(4), 979–979. doi:10.1890/12-1358.1

Schaeffer-Novelli, Y., Cintrón-Molero, G., Adaime, R. R., & de Camargo, T. M. (1990). Variability of mangrove ecosystems along the Brazilian coast. *Estuaries*, *13*(2), 204–218. doi:10.1007/BF02689854

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nature Methods*, *9*(7), 671–675. doi:10.1038/nmeth.2089

Scholz, A., Klepsch, M., Karimi, Z., & Jansen, S. (2013). How to quantify conduits in wood? *Frontiers in Plant Science*, *4*(March), 1–11. doi:10.3389/fpls.2013.00056

Shearman, P., Bryan, J., & Walsh, J. P. (2013). Trends in Deltaic Change over Three Decades in the Asia-Pacific Region. *Journal of Coastal Research*, *29*(5), 1169–1183. doi:10.2112/JCOASTRES-D-12-00120.1

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015). BUSCO: Assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*, *31*(19), 3210–3212. doi:10.1093/bioinformatics/btv351

Soares, M. L. G., Estrada, G. C. D., Fernandez, V., & Tognella, M. M. P. (2012). Southern limit of the Western South Atlantic mangroves: Assessment of the potential effects of global warming from a biogeographical perspective. *Estuarine, Coastal and Shelf Science*, *101*, 44–53. doi:10.1016/j.ecss.2012.02.018

Souza-Filho, P. W. M., Cohen, M. C. L., Lara, R. J., Lessa, G. C., Koch, B., & Behling, H. (2006). Holocene Coastal Evolution and Facies Model of the Bragança Macrotidal Flat on the Amazon Mangrove Coast, Northern Brazil. *Journal of Coastal Research*, (39), 306–310.

Steyn, W. J., Wand, S. J. E., Holcroft, D. M., & Jacobs, G. (2002). Anthocyanins in vegetative tissues: a proposed unified function in photoprotection. *New Phytologist*, *155*(3), 349–361. doi:10.1046/j.1469-8137.2002.00482.x

Stuart, S. A., Choat, B., Martin, K. C., Holbrook, N. M., & Ball, M. C. (2006). The role of

freezing in setting the latitudinal limits of mangrove forests. *New Phytologist*, *173*(3),
576–583. doi:10.1111/j.1469-8137.2006.01938.x
Takayama, K., Tateishi, Y., Murata, J., & Kajita, T. (2008). Gene flow and population
subdivision in a pantropical plant with sea-drifted seeds Hibiscus tiliaceus and its allied
species: Evidence from microsatellite analyses. *Molecular Ecology*, *17*(11), 2730–2742.
doi:10.1111/j.1365-294X.2008.03799.x
Tohge, T., Nishiyama, Y., Hirai, M. Y., Yano, M., Nakajima, J. I., Awazuhara, M., … Saito,
K. (2005). Functional genomics by integrated analysis of metabolome and transcriptome
of Arabidopsis plants over-expressing an MYB transcription factor. *Plant Journal*, *42*(2),
218–235. doi:10.1111/j.1365-313X.2005.02371.x
Wang, D., Pan, Y., Zhao, X., Zhu, L., Fu, B., & Li, Z. (2011). Genome-wide temporal-spatial
gene expression profiling of drought responsiveness in rice. *BMC Genomics*, *12*.
doi:10.1186/1471-2164-12-149
Wang, S., Bai, G., Wang, S., Yang, L., Yang, F., & Wang, Y. (2016). Chloroplast RNA-
Binding Protein RBD1 Promotes Chilling Tolerance through 23S rRNA Processing in
Arabidopsis. *PLoS Genetics*, *12*(5), 1–21. doi:10.1371/journal.pgen.1006027
Wang, Z., Gerstein, M., & Snyder, M. (2009). RNA-Seq: a revolutionary tool for
transcriptomics. *Nature Reviews. Genetics*, *10*(1), 57–63.
Wee, A. K. S., Mori, G. M., Lira-Medeiros, C. F., Núñez-Farfán, J., Takayama, K., Faulks,
L., … Kajita, T. (2018). The integration and application of genomic information in
mangrove conservation. *Conservation Biology*, 1–40. doi:10.1111/cobi.13140
Wolf, J. B. W., Lindell, J., & Backstrom, N. (2010). Speciation genetics: current status and
evolving approaches. *Philosophical Transactions of the Royal Society B: Biological
Sciences*, *365*(1547), 1717–1733. doi:10.1098/rstb.2010.0023
Young, M. D., Wakefield, M. J., Smyth, G. K., & Oshlack, A. (2010). Gene ontology analysis
for RNA-seq: accounting for selection bias. *Genome Biology*, *11*(2). doi:10.1186/gb-
2010-11-2-r14
Zhang, A., Ren, H.-M., Tan, Y.-Q., Qi, G.-N., Yao, F.-Y., Wu, G.-L., … Wang, Y.-F. (2016).
S-Type Anion Channels SLAC1 and SLAH3 Function as Essential Negative Regulators
of Inward K $^{+}$ Channels and Stomatal Opening in Arabidopsis. *The Plant Cell*, *28*(4),
949–965. doi:10.1105/tpc.15.01050
Zhang, C., Zhang, L., Zhang, S., Zhu, S., Wu, P., Chen, Y., … Wu, G. (2015). Global
analysis of gene expression profiles in physic nut (Jatropha curcas L.) seedlings exposed
to drought stress. *BMC Plant Biology*, *15*(1), 1–14. doi:10.1186/s12870-014-0397-x
984

985   **Data accessibility**

986   Expression data and sequences that support the findings have been deposited in

987   GenBank with the primary accession code GSE116060. A Variant Call Format file and its

988   complementary file, both required for all of the genome-wide SNP diversity analyses are

989   available in the supporting material. Other data are available upon request.

Author contributions

A.P.S., R.S.O., G.M.M. and M.V.C. designed the study. M.V.C. and G.M.M. conducted fieldwork. M.V.C. and C.S.M. cultivated seedlings and performed analyses of morphophysiological data. M.V.C. prepared samples and performed RNA-sequencing. M.V.C., M.D., D.H.O. and G.M.M. analysed RNA-Seq data. C.C.S. and M.V.C. verified RNA-Seq data through qRT-PCR. G.M.M. prepared samples and performed genotyping of genome-wide SNP. G.M.M. and M.V.C. analysed nextRAD results. A.P.S, M.I.Z., G.M.M. and R.S.O. contributed material/reagents/analysis tools. M.V.C. and G.M.M. wrote the manuscript. All authors discussed the results and contributed to the manuscript.

 Figures and Tables

1001

Figure 1. Geographical locations of *Avicennia schaueriana* sampling sites. Green-shaded
area represents the geographical distribution of the species. Black triangles represent the
locations of the Equatorial and Subtropical sampling sites for propagules and plant tissues
used in the common garden experiment and RNA-sequencing, respectively. Coloured dots
represent sampling sites of leaves used for the nextRAD genome-wide population diversity
analyses (red: population located north of the northeast extremity of South America
(NEESA); blue: population located south of the NEESA). Arrows approximately represent
the directions of the major sea currents along the Atlantic coast of South America.

1010

Figure 2. Morphological divergence observed in seedlings of *Avicennia schaueriana* collected from Equatorial and Subtropical sampling sites and grown in a common garden experiment. Violin plots represent the distribution of observations for plants from Equatorial (red) and Subtropical (blue) sampling sites. Box plots represent the mean, standard error, and maximum and minimum values. Two group comparisons were performed using the non-parametric unpaired Mann-Whitney Wilcoxon U-tests. For all variables represented in the figure, the absence of a difference between groups was rejected at a significance threshold of 0.05. (a) leaf inclination angle (n = 15 leaves per group, 5 plants per group); (b) individual leaf area (n = 250 leaves per group, 3 plants per group); (c) leaf dry mass ratio (leaf dry biomass/plant dry biomass) (n = 15 plants per group); (d) stem dry mass ratio (stem dry biomass/plant dry biomass) (n = 15 plants per group); (e) vessel lumen area ratio in sapwood (n = 175 per group, 3 plants per group, observations represented by black points); (f) vessel lumen diameter (n = 700 vessels per group, 3 plants per group); (g) vessel lumen conductivity (n = 700 vessels per group, 3 plants per group); (h) vessel lumen resistivity (n = 700 vessels per group, 3 plants per group).

1027

Figure 3. Light reflectance of the stem epidermis of five-month-old *Avicennia schaueriana* **seedlings grown in a common garden experiment.** Grey lines represent the mean reflectance, and colour-shaded areas represent the standard error for each seedling source site (red: Equatorial; blue: Subtropical, n = 10 plants per group). The visible light spectrum range is highlighted in the figure.

1033

Figure 4. Graphical summary of the integration of ecological and molecular approaches performed in this work.
Key oceanographic and climatic factors differ markedly between equatorial and subtropical latitudinal distribution extremes of the distribution of the neotropical mangrove tree, *Avicennia schaueriana*, and possibly shape the diversity of genotypes and phenotypes in the species. To address this issue, we examined the effects of the contrasting environments on overall gene expression, its morphophysiological effects in a common garden experiment and its genomic effects through natural selection detection tests based on single nucleotide polymorphism (SNP). Plants from equatorial and subtropical latitudes showed key divergences related to the use of water and to the acquisition of carbohydrates both in the field and in common garden conditions. In addition, north-south genetic divergence was observed in all genotyped SNPs. We also identified signatures of differential selective pressures on specific loci possibly related to the accumulation of anthocyanin (*UDP-GLUCOSYLTRANSFERASE*), the response to osmotic stress (*RAP2.4* and *ZFN1*), photosynthesis (*TPR*) and hypoxia (*HRA1*). The molecular and ecologic divergences observed through three independent approaches may be related to environmental factors that strongly differ between contrasting latitudes at which the species are found. These findings highlight the power of using multidisciplinary approaches for the study of adaptation in species for which little basic biological information is available, such as tropical trees. VPD: atmospheric vapour pressure deficit; E: transpiration rate; g_s: stomatal conductance; R_L: xylem vessel lumen resistivity; K_H: xylem vessel conductivity.

1048 **Table 1.** Characterisation of Subtropical and Equatorial sampling sites of propagules used in
1049 the common garden experiment and of RNA used for RNA-sequencing. SD: standard
1050 deviation of the mean.

	Subtropical	Equatorial
Köppen-Geiger climate characterisation[†]	Temperate oceanic with hot summer, without a dry season (Cfa)	Tropical monsoon (Am)
Latitude (°)	28 S	0
Tidal amplitude	Microtidal (< 1 m)	Macrotidal (> 4 m)
Annual mean temperature (°C)[‡]	20.09	26.42
Minimum temperature of the coldest month (°C)[‡]	11.76	22.04
Maximum temperature of the warmest month (°C)[‡]	28.66	31.1
Annual precipitation (mm)[‡]	1,435	2,216
Precipitation in the driest month (mm)[‡]	88	4
Precipitation in the wettest month (mm)[‡]	162	452
Mean air VPD (KPa)[‡]	1.95	2.82
Maximum air VPD (KPa)[‡]	2.47	2.95
Minimum air VPD (KPa)[‡]	1.48	2.71
Mean irradiance (KJ m^{-2} day^{-1})[‡]	14,270	17,414
Maximum irradiance (KJ m^{-2} day^{-1})[‡]	20,802	21,671
Minimum irradiance (KJ m^{-2} day^{-1})[‡]	8,201	13,874
Mean sea surface salinity (g/kg)[§]	35.50	34.96
Sea surface salinity in the saltiest month (g/kg)[§]	36.24	36.87
Sea surface salinity in the freshest month (g/kg)[§]	33.73	32.54
Average day length (hours (± SD))[¶]	12.103 (±1.251)	12.115 (±0.033)
True mangrove species in the area	*Avicennia schaueriana* *Laguncularia racemosa*	*Avicennia germinans* *Avicennia schaueriana* *Laguncularia racemosa* *Rhizophora mangle* *Rhizophora racemosa* *Rhizophora harriisoni*

1051   VPD: Vapour pressure deficit. [†]According to Alvares et al. (Alvares, Stape, Sentelhas,
1052   Gonçalves, & Sparovek, 2013). [‡]Source: BioClim (Hijmans, Cameron, Parra, Jones, & Jarvis,
1053   2005). [§]Source: MARSPEC (Sbrocco & Barber, 2013). [¶]Source: 'daylength' function from R
1054   package 'geosphere' (Forsythe, Rykiel, Stahl, Wu, & Schoolfield, 1995).