Full title

Environment-driven genomic and trait divergence in a neotropical mangrove highlight

differential sensitivities to climate change

Short title

Environment-driven genomic and trait divergence in a non-model species

Authors

Mariana Vargas Cruz[1], Gustavo Maruyama Mori[2], Caroline Signori Müller[1], Carla Cristina da

Silva[1], Dong-Ha Oh[3], Maheshi Dassanayake[3], Maria Imaculada Zucchi[4], Rafael Silva

Oliveira[1] and Anete Pereira de Souza[1]*.

Affiliations

[1] Department of Plant Biology, Institute of Biology, University of Campinas, Campinas, São

Paulo, Brazil.

[2]Institute of Biosciences, São Paulo State University (Unesp), São Vicente, São Paulo, Brazil.

[3]Department of Biological Sciences, Louisiana State University, Louisiana, United States.

[4]São Paulo Agency for Agribusiness Technology, Piracicaba, São Paulo, Brazil.

*Corresponding author. Email: anete@unicamp.br

Abstract

Integrating genomic and ecological data are instrumental for understanding the mechanisms of adaptive processes in natural ecosystems. In non-model species, such studies can be particularly challenging, but often brings to light results with implications for conservation. Here, we integrate molecular and ecophysiological approaches to assess the role of selection in the north-south organization of the genetic variation in the mangrove species *Avicennia schaueriana*, a new-world tree found from tropical to temperate coastal forests in the Atlantic coast of the Americas. In a common garden experiment, individuals from equatorial and subtropical forests diverged in traits involved in water balance and carbon acquisition, suggesting a genetic basis of the observed differences. RNA-sequencing highlighted the molecular effects of different light, temperature and air humidity in individuals under field conditions at contrasting latitudes. Additionally, genetic polymorphisms in trees sampled along most of the species range showed signatures of selection in sequences associated with the biogenesis of the photosynthetic apparatus, anthocyanin biosynthesis and osmotic and hypoxia stress responses. We found substantial divergences between populations occurring north and south of the north-eastern extremity of South America, underpinning roles of contrasting environmental forces in shaping the genetic structure of the species. The observed functional divergence might differentially affect sensitivities of populations to our changing climate, as discussed here. We indicate the necessity of independent conservation management for the long-term persistence of the species diversity. Moreover, we demonstrate the power of using a multidisciplinary approach in adaptation studies of non-model species.

1.   Introduction

Adaptation to contrasting environments is an ubiquitous consequence of divergent selective forces acting on phenotypic diversity within species[1–3]. Phenotypic variation can be achieved through plasticity during acclimation to environmental changes, or ultimately through genetic variation shaped by adaptive processes[4]. Though the occurrence of intraspecific divergence in adaptive traits is well recognised, its molecular basis is not yet fully understood[5]. The integration of multiple independent approaches to understand the bases of adaptive trait variation is desirable to minimise the potential for incorrect conclusions[6].

In addition to its fundamental relevance in the field of evolutionary biology, research on the molecular mechanisms underlying adaptive variation has the potential to provide valuable information for ensuring species diversity persists under environmental challenges. Especially, accelerated rates of contemporary climate change call for studies on patterns of functional variation and its consequences for species responses to future climate conditions[7]. Climate change forecasts include a rise in sea-level from 0.26 to 0.82 m, an increase in mean atmospheric temperature from 0.3 to 4.8 °C and changes in precipitation regimes by the end of this century[8]. These changes are projected to especially affect certain ecosystems, as mangrove forests[9], since they are distributed in narrow intertidal environments in tropical and subtropical zones and are naturally limited by annual minimum temperatures and average rainfall[10]. Recent changes in sea level, in atmospheric temperature and in rainfall have already promoted shifts in the distribution of these ecosystems and in the density of individuals in forests[11–17]. Negative impacts are predicted mainly in regions in which aridity is expected to increase and where adjacent areas for population expansion are unavailable or do not exist[18]. However, in some regions,

mangroves are expected to persist through their ability to adjust soil elevation[11,19] and to

rapidly shift their distribution to new suitable areas[11,18,20,21].

Yet, these predictions do not account for extant intraspecific variability across the

geographical distribution of a given mangrove species. For instance, the genetic diversity of

all mangrove species studied to date is structured into two populations occurring north and

south of the northeast extremity of South America (NEESA)[22–24]. In the NEESA region, a

bifurcating subdivision of the "South Equatorial Current" into the coastal "Guiana Current"

and "Brazil Current" is observed, dispersing mangrove propagules in opposite directions and

likely reducing the gene flow between populations[22,23,25] (Fig 1). Limited gene flow is a

key process determining the magnitude of local adaptation and, when this condition is met,

one may expect populations to adapt differently to their environments[2,26]. However, the

neutral nature of the molecular markers used in previous works[22,23,25] precluded

inferences on environment-driven genetic divergence.

Fig 1. Geographical location of *Avicennia schaueriana* sampling sites. The green shaded area represents the geographical distribution of the species. Black triangles represent the location of the Equatorial and Subtropical sampling sites for propagules and plant tissues used in the common garden experiment and in RNA-sequencing, respectively. Coloured dots represent sampling sites of leaves used for extraction of genomic DNA for nextRAD sequencing (red: population located north of the northeast extremity of South America (NEESA); blue: population located south of the NEESA). Arrows approximately represent the direction of the major sea currents that act along the Atlantic coast of South America.

Advances in DNA sequencing now permit the undertaking of genome-wide studies to identify neutral and putatively adaptive genetic variation even in non-model organisms, such as mangrove trees[27]. We used these tools in the present work to investigate the non-neutral divergence between previously identified populations of the new-world mangrove tree *Avicennia schaueriana* Stapf & Leechman ex Moldenke. We further explored the potential role of environmental forces underlying such divergence along the Atlantic coast of South America. *Avicennia schaueriana* is the most widely distributed mangrove species with genetic diversity information available over this geographic scale. The species is found in the

Lesser Antilles (~16 °N) and from Venezuela to the southernmost mangrove forests in the

Atlantic (~28 °S)[28]. To avoid incorrect conclusions about the targets of selection[6], we

integrated three independent but complementary molecular and ecological approaches: (1)

comparative physiology of contrasting Equatorial and Subtropical samples grown in a

common garden experiment; (2) comparative transcriptomics of trees from Equatorial and

Subtropical localities, sampled in their native environments (Table 1, S1 Fig) and (3) genome

scans for signatures of selection using high-throughput genome-wide genotyping of

individuals, sampled along almost the entire distribution of the species (Fig 1). We further

discuss the implications of our results in the face of predicted changes in climate as well as

potential strategies for mangrove conservation along the Atlantic coast of South America.

Table 1. Characterisation of Subtropical and Equatorial sampling sites of propagules used in the
common garden experiment and of RNA used for RNA-sequencing.

Environmental features	Subtropical	Equatorial
Köppen-Geiger climate characterisation[†]	Temperate oceanic with hot summer, without a dry season (Cfa)	Tropical monsoon (Am)
Latitude	28 S	0
Tidal amplitude	Microtidal (< 1 m)	Macrotidal (> 4 m)
Annual mean temperature (°C)[‡]	20.09	26.42
Minimum temperature of the coldest month (°C)[‡]	11.76	22.04
Maximum temperature of the warmest month (°C)[‡]	28.66	31.1
Annual precipitation (mm)[‡]	1,435	2,216
Precipitation in the driest month (mm)[‡]	88	4
Precipitation in the wettest month (mm)[‡]	162	452
Mean air VPD (KPa)[‡]	1.95	2.82
Maximum air VPD (KPa)[‡]	2.47	2.95

Minimum air VPD (KPa)[‡]	1.48	2.71
Mean irradiance (KJ m^{-2} day^{-1})[‡]	14,270	17,414
Maximum irradiance (KJ m^{-2} day^{-1})[‡]	20,802	21,671
Minimum irradiance (KJ m^{-2} day^{-1})[‡]	8,201	13,874
Mean sea surface salinity (g/kg)[§]	35.50	34.96
Sea surface salinity in the saltiest month (g/kg)[§]	36.24	36.87
Sea surface salinity in the freshest month (g/kg)[§]	33.73	32.54
Average day length (hours (± sd))[¶]	12.103 (±1.251)	12.115 (±0.033)
True mangrove species at this area	*Avicennia schaueriana Laguncularia racemosa*	*Avicennia germinans Avicennia schaueriana Laguncularia racemosa Rhizophora mangle Rhizophora racemosa Rhizophora harriisoni*

VPD: Vapour pressure deficit. [†]According to Alvares *et al.* [114]. [‡]Source: BioClim[31]. [§]Source: MARSPEC[32]. [¶]Source: 'daylength' function from R package 'geosphere'[115]. sd: standard deviation of the mean.

2. Results

2.1 Comparative physiology in a common garden experiment

To identify functional divergence between individuals from Equatorial and Subtropical sites, we conducted a common garden experiment in a glasshouse, under a homogeneous, non-climate-controlled environment. We observed differences in key morphophysiological traits between seedlings from contrasting latitudes grown in this experiment (Fig 2 and 3). Subtropical plants showed higher transpiration rates and stomatal conductance than Equatorial plants (S2 Fig). Additionally, plants from the two origins were morphologically different from each other. The inclination angle of the leaf lamina was steeper in Equatorial than Subtropical plants, and the average size of individual leaves was smaller in Equatorial seedlings, even though the total leaf area and specific leaf area did not significantly differ between the groups (Fig 2, S1 Table, S3 Fig, S1 Data). Subtropical plants

accumulated more biomass in all vegetative organs (leaves, stems and roots) in comparison to

Equatorial plants under the experimental conditions. However, the biomass fraction allocated

to stems in relation to other organs (stem dry mass ratio, SMR = stem dry biomass/plant dry

biomass) was greater in Equatorial plants, whereas allocation to leaves (leaf dry mass ratio,

LMR = leaf dry biomass/plant dry biomass) was greater in Subtropical plants. Biomass

allocation to roots (root dry mass ratio, RMR = root dry biomass/plant dry biomass) did not

differ between groups (S1 Table). Unexpectedly, 63% of the Equatorial plants flowered after

the sixth month of growth (S3G Fig). Since this was not observed in any Subtropical plant,

Equatorial plants with reproductive branches were not used in the biomass allocation analysis.

Fig 2. Morphological divergence observed in seedlings of *Avicennia schaueriana* collected from
an Equatorial and a Subtropical sampling sites and grown in a glasshouse conducted common
garden experiment. Violin plots represent the distribution of observations for plants from Equatorial
(red) and Subtropical (blue) sampling sites. Box plots represent the mean, standard error, maximum
and minimum values. Two group comparisons were performed using the non-parametric unpaired
Mann-Whitney Wilcoxon U-tests. For all variables represented in the figure, the absence of a
difference between groups was rejected at a significance threshold of 0.05. (A) Leaf inclination angle

(n = 15 leaves per group, 5 plants per group); (B) individual leaf area (n = 250 leaves per group, 3
plants per group); (C) leaf dry mass ratio (Leaf dry biomass/Plant dry biomass) (n = 15 plants per
group); (D) stem dry mass ratio (Stem dry biomass/Plant dry biomass) (n = 15 plants per group); (E)
vessel lumen area ratio in sapwood (n = 175 per group, 3 plants per group, observations represented by
black points); (F) vessel lumen diameter (n = 700 vessels per group, 3 plants per group); (G) vessel
lumen conductivity (n = 700 vessels per group, 3 plants per group); (H) vessel lumen resistivity (n =
700 vessels per group, 3 plants per group).

Fig 3. Light reflectance of the stem epidermis of five-month-old *Avicennia schaueriana* seedlings
grown in a common garden experiment. Grey lines represent the mean reflectance and colour-
shaded areas represent the standard error for each seedling source site (red: Equatorial; blue:
Subtropical, n = 10 plants per group). The visible light spectrum range is highlighted in the figure.
149

Stems from Subtropical samples showed wider vessel lumen diameter than those from

Equatorial samples, but the vessel density was not significantly different between the groups

(Fig 2, S3H-S3I Fig). The sapwood space dedicated to vessel lumen area was also greater in

Subtropical than in Equatorial plants. Vessels of Subtropical plants, showed higher

conductivity at a detriment of hydraulic safety, compared to Equatorial plants (Fig 2). The

total conductivity of the stems was not significantly different between the sampling groups.

Additionally, plants from these contrasting origins exhibited different pigmentation in the

stem epidermis. Stems from Equatorial seedlings reflected more red light of long wavelengths

(635-700 nm) and less green light of medium wavelengths (520-560 nm) than Subtropical

seedlings (Fig 3, S2 Data).

2.2   Comparative transcriptomics using an RNA-sequencing experiment

In the absence of a reference genome, we used RNA-sequencing (RNA-Seq)[55] to

obtain a *de novo* assembled transcriptome for *A. schaueriana* from leaves, stems and flowers

of six adult individuals, from Equatorial and Subtropical sampling sites (S4 Fig). We obtained

over 240 million raw, paired-end, 72 bp reads and over 209 million high quality reads after

quality control (Fig 1, Table 2). High quality reads were assembled into 49,490 transcripts of

the species, after the removal of misassembled, contaminant and redundant transcripts.

Putative protein-coding sequences comprised 61% of the transcriptome (30,227 sequences).

Over 91.9% of the high-quality reads were mapped to a single transcript, providing evidence

of minimum redundancy and a wide representation of reads obtained via sequencing (S2

Table). Moreover, searching for universal plant orthologous genes using the BUSCO

pipeline[45] revealed that 91.8% of the conserved sequences in the plant database were

present in the reference transcriptome (S2 Table). Assembled sequences with complete ORF

represented approximately 42% (12,796) of all putative protein-coding transcripts, with an

average length of 1,324 nucleotides (S3 Table, S5 Fig). The majority (94.33%) of these

putative protein-coding sequences showed significant similarity to proteins in the Plant

RefSeq[40] and TAIR databases. Only a minor fraction (2.85%) of the putative protein-

coding transcripts could not be annotated using publicly available databases (S5C Fig). More

than 80% of transcripts matched proteins from *Sesamum indicum* L. or *Erythranthe guttata*

(Fisch. ex DC.) G.L. Nesom, which, together with *A. schaueriana* are grouped in the order

Lamiales (S5D Fig). We identified putative orthologous genes between *A. schaueriana* and

other closely related *Avicennia* L. (Avicenniaceae) species. This analysis revealed the

presence of 27,658, 18,325 and 13,273 putative orthologues between the *A. schaueriana*

reference transcriptome and the transcriptomes derived, respectively, from *A. marina*[56]

leaves and *A. officinalis* leaves[57] and roots[58] (S4 Table).

Table 2. Characterisation of field conditions at the moment of plant material collection and storage in
RNA-stabilizing solution for subsequent sequencing.

Sample ID	Sampling site (Climate[†])	GPS Point	Date (d/m)	Time (h:m)	TL (m)	DL (h)	Temp (°C)[‡]	RH (%)[‡]	VPD (kPa)[§]	SI (KJ/m^2)[‡]
St1	Subtropical (Cfa)	28.48 S; 48.88 W	23/08	15:15	0.5	11.3	19	84	0.351	2033.89
St2	Subtropical (Cfa)	28.36 S; 48.80 W	24/08	10:35	0.5	11.3	18.3	86	0.294	1804.15
St4	Subtropical (Cfa)	28.48 S; 48.83 W	25/08	13:11	0.5	11.3	20.9	82	0.445	1863.14
Eq4	Equatorial (Am)	00.64 S; 47.26 W	28/07	11:57	2.7	12.1	28.5	49	1.983	1940.73
Eq5	Equatorial (Am)	00.65 S; 47.26 W	29/07	12:22	2.7	12.1	28.3	51	1.883	2724.54
Eq6	Equatorial (Am)	00.64 S; 47.26 W	29/07	16:04	0.7	12.1	30.1	47	2.260	1433.68

TL: Tidal level; DL: day length; Temp: atmospheric air temperature; RH: atmospheric relative
humidity; VPD: air vapour pressure deficit; SI: solar irradiance.
[†]Köppen-Geiger Climate classification according to Alvares *et al.*[114].
[‡]Source: Brazilian National Institute of Meteorology (INMET).
[§]Estimated according to McRae[116].

To identify environmental forces associated with variations in gene expression

between source sites, we detected differentially expressed transcripts (DET) between samples

using the EdgeR[46]. Sampling was conducted at the end of winter at the Subtropical site and

at the beginning of the dry season at the Equatorial site (Table 2). Although plant material

was collected under uncontrolled field conditions, we observed a consistency in transcripts

expression levels from leaves and stems samples from plants from the same sampling site

(S6A-S6B Fig). Levels of transcript expression in flowers, however, did not present a

consistent pattern within sampling sites (S6C Fig), leading to the identification of only one

DET between flowers from Equatorial and Subtropical sites. Therefore, we did not include

flowers in the subsequent analyses (S6F Fig). Conversely, we identified 1,837 and 904

transcripts showing significantly different (FDR < 0.05) relative abundance, respectively in

leaves and stems, between samples at the Equatorial and at the Subtropical sites (S6D-S6E

Fig). Among the 2,741 DET, 1,150 (41.91%) were putative protein-coding transcripts.

The assignment of transcripts to Gene Ontology (GO) terms was possible for 25,184

(83.31%) out of 30,227 putative protein-coding sequences in the transcriptome and enabled

the performing of GO enrichment analyses of the DET. GO analyses were focused on

biological processes potentially regulating the responses of *A. schaueriana* trees to the

contrasting climatic variables in the equatorial and subtropical regions (Table 1, S1 Fig).

Analyses were conducted separately for leaves and stems and for each of the two sets of DET,

one showing increased expression levels in Equatorial than in Subtropical samples (we refer

to these as DET-Eq) and the other set showing higher abundance in Subtropical compared to

Equatorial samples (these are referred as DET-St). The enriched processes among the sets of

DET included photosynthesis, plant responses to UV, to temperature stimulus, to water stress,

to cell wall biosynthesis and to cellular respiration (S5-S9 Tables, S6I-S6-L Fig).

2.2.1   Photosynthesis

Among the DET-St, we observed the enrichment of categories related to

photosynthesis, including various putative genes participating in the biosynthesis of the

photosynthetic apparatus, in the biosynthesis of chlorophyll and photoreceptors, in the

function of electron transporters and in the coordination of chloroplasts movement. In

contrast, the DET-Eq set showed enrichment in transcripts similar to proteins required for

disassembling the light-harvesting complex of photosystem II in thylakoid membranes and for

triggering chlorophyll degradation[59] (S6I-S6J Fig, S9 Table).

2.2.2　Response to UV

Both DET-St and DET-Eq sets showed enrichment in functions related to the response

to UV-light, however, the sets of UV-responsive genes differed between these groups. The

DET-St included UV-B protectants and genes involved in the biosynthesis of antioxidants

induced by UV-B, such as plastid copper/zinc superoxide dismutases, photosystem II repair

proteins, and transcripts involved in the synthesis of the antioxidant L-ascorbic acid (vitamin

C). Conversely, the set of DET-Eq showed enrichment of putative genes involved in photo-

oxidative damage reduction and in the positive regulation of anthocyanin biosynthesis in

response to UV light. Antioxidants induced by UV irradiation[60], such as iron superoxide

dismutases and transcripts involved in the biosynthesis of the pyridoxine (vitamin B6), were

also among the DET-Eq (S9 Table 9).

2.2.3　Response to temperature

In the DET-St set, we observed putative genes presenting critical roles in chloroplast

protein translation during cold acclimation and which provide tolerance to chilling

temperatures[61,62]. For instance, transcripts similar to the *GLYCINE-RICH RNA-BINDING*

PROTEIN (*RZ1A*), which has a chaperone activity during cold acclimation[63], and to the

ATP-dependent *DNA HELICASE* ATSGS1, a cold-inducible protein required for DNA

damage-repair[64]. Interestingly, we also identified in the DET-St set a putative *AGAMOUS-*

LIKE 24 (*AGL24*) transcription factor, involved in floral transition induced by

vernalisation[65]. Contrastingly, various transcripts similar to heat shock-inducible

chaperones and an *ADENINE NUCLEOTIDE ALPHA HYDROLASE-LIKE PROTEIN*

(*AT3G53990*), involved in chaperone-mediated protein folding[66], were detected in the set

of DET-Eq, potentially enhancing tolerance to heat in the Equatorial plants. Additionally, a

transcript similar to the *ETHYLENE-RESPONSIVE ELEMENT BINDING PROTEIN*

(*RAP2.3*), which confers resistance to heat and to hydrogen peroxide[67], was observed in

this group (S9 Table).

2.2.4 Response to water stress

Transcripts associated with the response to water deprivation, to cellular ion

homeostasis and to osmotic adjustment were enriched in the DET-Eq. Dehydration-induced

transcriptional regulators and genes that improve tolerance to water deficits were among these

transcripts. For instance, a transcript similar to the *ETHYLENE-RESPONSIVE*

TRANSCRIPTION FACTOR (*RAP2.4*), induced by water stress and that regulates the

expression of several drought-responsive genes, including aquaporins[68,69] was identified in

the DET-Eq. Accordingly, a putative aquaporin *PLASMA MEMBRANE INTRINSIC*

PROTEIN (PIP1;4)[70] was also in this set. We also observed in the DET-Eq, putative genes

participating in the synthesis and accumulation of raffinose, an osmoprotective soluble

trisaccharide[71], and also transcripts similar to osmosensitive ion channels belonging to the

EARLY-RESPONSIVE TO DEHYDRATION STRESS FAMILY PROTEIN. In line with this,

we observed among the DET-Eq an ion channel protein, *SLAC1 HOMOLOGUE 3* (*SLAH3*),

required for stomatal closure induced by drought stress[72] and also the putative *NINE-CIS-*

EPOXYCAROTENOID DIOXYGENASE 3 (*NCED3*), which increases plant resistance to

water deficit, through the accumulation of abscisic acid (ABA) leading to stomatal closure.

Possibly as a consequence from decreased stomatal conductance, a transcript similar to the

ALANINE-GLYOXYLATE AMINOTRANSFERASE 2 HOMOLOG 3 (*AT3G08860*), which

plays a central role in photorespiration[73], showed higher expression in Equatorial than in

Subtropical samples (S9 Table).

2.2.5   Cell wall biosynthesis

Transcripts similar to 33 distinct proteins and transcription factors that play central

roles in the biosynthesis and deposition of distinct cell wall components, such as cellulose,

hemicellulose, lignin and pectin, were identified among the DET-Eq (S9 Table).

2.2.6   Cellular respiration

The DET-Eq were also enriched in putative cellular respiration genes, which included

one enzyme encoded in the nuclear genome, the *ACONITASE 3* (*ACO3*), which converts

citrate to isocitrate in the tricarboxylic acid cycle (TCA cycle), and several other genes from

the mitochondrial genome, encoding subunits of the NADH dehydrogenase, of the ATP-

synthase and of the cytochrome C oxidase (S9 Table).

We verified and confirmed the results obtained in the computational analyses of RNA-

Seq data by qRT-PCR of 10 DET detected across all leaf samples (S7 Fig, S10 Table, S1

Result).

2.3   Detection of genetic variants with a signature of positive selection

To complement the analyses of differential gene expression, which could result from

both plasticity and adaptive selection[74], we searched for gene sequence variation in trees

sampled along the Atlantic coast of South America (Fig 1, S11 Table). After quality filtering

of sequenced data, we selected 77 individuals, which did not show signs of interspecific

hybridization with *Avicennia germinans*, for downstream analyses. These data were used in

population genetics approaches to describe genetic structure and to perform genome-scans for

loci with signatures of positive selection. We identified a set of 6,170 high-quality unlinked

biallelic single nucleotide polymorphic loci (SNP) with a minor allele frequency ≥ 0.05 and $\geq$

8x coverage. The overall genetic structure corroborated a previous study based on putatively

neutral microsatellite loci[22], dividing the sampling sites into two main groups: north and

south of the NEESA (S8 Fig).

To investigate the role of non-neutral evolutionary forces driving the observed

population divergence, we used two distinct methods that allowed the detection of loci that

showed significant departures from a neutral expectation of interpopulation divergence. To

minimise false-positive signatures of selection, we considered 122 loci that were

conservatively detected by both pcadapt[54] and LOSITAN[75]. These loci were aligned to

transcriptome characterised herein, enabling the screening for their potential functional roles.

We found 16 candidate loci with putative evidence of selection within expressed sequences,

15 of which showed similarity to gene models in *A. thaliana* and *S. indicum* (S12 Table).

However, five of these reference proteins did not have informative function described for the

model plant, hindering inferences about their putative function in *A. schaueriana*. Conversely,

among the remaining ten annotated candidate loci, we found five genes involved in processes

related to the contrasting climate regimes of equatorial and subtropical environments. One of

the candidate SNP loci was detected in the putative transcription factor *ETHYLENE-*

RESPONSIVE TRANSCRIPTION ACTIVATOR (*RAP2.4),* which is induced in response to

water and salt stress. This gene also regulates developmental processes mediated by light

intensity[68] and the expression of aquaporins in *A. thaliana*[69], playing a role in plant water

homeostasis. Two other candidate loci with putative signature of selection were associated

with transcription factors, the *ZINC-FINGER PROTEIN 1* (*ZFN1*), involved in the regulation

of the expression of several water stress genes[76] and the *HYPOXIA RESPONSE*

ATTENUATOR (*HRA1*)[77], which is strongly induced in *A. thaliana* in response to low

oxygen levels. Additionally, the putative enzyme *UDP GLUCOSYL TRANSFERASE*, which

catalyses the final step of anthocyanin biosynthesis, wherein pigmentation is produced[78],

also showed a SNP variation with putative sign of positive selection. And one candidate locus

was present in a transcript similar to the *AT2G20710* gene model in *A. thaliana*, a

TETRATRICOPEPTIDE REPEAT (*TPR*) protein, which might play a role in the biogenesis of

the photosynthetic apparatus[79].

2.4 Data availability statement

All relevant morphophysiological data assessed from plants grown in the common

garden experiment are available in the Supplementary information files (S1-S2 Data).

Expression data and sequences that support the findings have been deposited in GenBank

with the primary accession code GSE116060. Variant Call Format (VCF) file and its

complementary file, both required for all genome-wide SNP diversity analyses are available

in the Supplementary information files (respectively S3 Data and S4 Data).

3. Discussion

In the present work, we used complementary ecological and molecular approaches to

unveil the nature of non-neutral phenotypic and genetic divergences as well as the role played

by contrasting environmental forces between two populations of the mangrove species,

Avicennia schaueriana. The overall genetic structure previously detected with a few

microsatellite loci between populations occurring north and south of the northeast extremity

of South America (NEESA)[22] was confirmed here using thousands of genome-wide SNP

loci (S8 Fig). Additionally, a common garden experiment revealed morphophysiological

differences in traits that are key to the regulation of water and carbon balances between

seedlings originated from the Equatorial and Subtropical sampling sites (Fig 2-3).

Comparatively, the Equatorial plants showed morphophysiological and transcriptomic signals

that appear to minimise effects of low water, high light and heat stresses, whereas traits

observed in Subtropical plants suggest a maximisation of carbon assimilation, beneficial

especially under low temperature and reduced light regime.

Traits exhibited by Equatorial plants compared to Subtropical plants in the common

garden experiment, such as reduced leaf size, the steeper leaf angle, higher levels of red light-

reflecting pigments, narrower vessels and lower rates of stomatal conductance, limit carbon

acquisition[80] and may have imposed constraints to carbon gain in Equatorial plants, which

also accumulated less biomass (Fig 2-3, S2-S3 Fig). Conversely, such characteristics allow

plants to maintain leaf temperature at suitable levels for photosynthesis, while reducing both

UV-exposure, through protective pigmentation and leaf inclination, and water loss, through

minimisation of evaporative cooling[80,81]. We argue that the prevalence of these traits

among Equatorial individuals may be advantageous in their natural environment, especially

during the dry season (from August to December), when high light intensity is frequently

combined with high temperature (> 30 °C) and average air humidity below 70% (Table 1, S1

Fig). In the presence of high atmospheric evaporative demand and highly saline intertidal

soils, water acquisition has an elevated energetic cost, representing a strong selective pressure

in favour of water-saving adaptations[80]. Accordingly, Equatorial plants also showed lower

transpiration rates than Subtropical plants in the common garden (S3 Fig). In addition, 63% of

the Equatorial plants showed an early transition from vegetative to reproductive stage, starting

flowering after only sixth months of growth, in July and August (S3G Fig). The period in

which flowering occurred is concordant with observations reported in literature for the species

in tropical sites, from June-August [82], however we did not find records of six-months-old

flowering plants. Even though in southern subtropical forests, a flowering peak is also

observed in August[83], Subtropical plants did not flower during the experiment. Early

flowering is a phenotype with complex genetic components, rarely studied, especially in non-

model organisms, but is an important adaptive mechanism for maximizing the chances of

reproduction under stress[84].

In contrast, Subtropical plants showed higher stomatal conductance and transpiration

rates, higher levels of green light-reflecting pigments, larger leaf area, wider leaf lamina angle

and larger xylem vessel diameter than Equatorial plants in the common garden experiment

(Fig 2-3, S2-S3 Fig). These characteristics enhance light energy absorbance and carbon

acquisition at the expense of greater water loss and higher cavitation risk[85,86]. These traits

may also compensate for declines in net primary production in higher-latitude

environments[87], resulting from compressed environmental windows of opportunity, mainly

in terms of temperature and solar irradiance (Table 1, S1 Fig). We argue that the intensity of

cold events in southern Subtropical populations of *A. schaueriana* is likely not sufficiently

expressive to favour the selection of freezing-tolerant individuals, in contrast to results

reported for the congeneric *A. germinans* at its northernmost distribution limit, on the Atlantic

coast of North America[14]. At the southern range edge of *A. schaueriana,* the annual

minimum air temperature does not drop below 0 □ (Table 1, S1 Fig), thus being higher than

the expected mangrove physiological threshold[10]. Additionally, the relatively small

population size of *A. schaueriana* at this location[28] and the arrival of maladapted

propagules from northerly populations might reduce the potential strength of natural selection

in favour of cold-adapted individuals at this Subtropical site.

Functional interpopulation divergence was also evident under field conditions at the

molecular level. Comparative transcriptomics of trees sampled in field conditions corroborate

with the suggested effects of the environmental variation in light availability, temperature and

water stress on the phenotypic divergence observed in the common garden experiment.

Transcriptomic profiles were obtained at the beginning of the dry season in the Equatorial

sampling site and at the end of winter in the Subtropical sampling site (Table 2) and showed

an enrichment of DET involved in photosynthesis, cellular respiration, cell wall biosynthesis

and plant responses to water stress, temperature stimulus and UV light (S6I-S6L Fig).

Additionally, the adaptive relevance of these biological processes in the field was highlighted

through the identification and functional annotation of SNP loci putatively under natural

selection from populations along the *A. schaueriana* geographic distribution (S12 Table). In

the following subsections, we integrate information derived from these different methods.

3.1   Water stress as a relevant selective pressure in equatorial populations

The increased levels of transcripts similar to heat-shock proteins, drought-induced ion

transporters, genes that enhance tolerance to heat, that are involved in stomatal closure and

that play central roles in photorespiration, provided multiple lines of evidence of water stress

in Equatorial samples. Additionally, compared to Subtropical plants, Equatorial plants

exhibited higher basal levels of expression of aquaporins and genes involved in the

accumulation of organic solutes. These investments improve tolerance to drought[71] by

lowering the cell water potential and rather actively transporting water through proteins in the

cell membrane than using passive apoplastic water transport[80,88,89]. The enhanced rigidity

of cells reduces the risk of cell damage during the dehydration and rehydration process and,

thus, improves resistance to the high extra-cellular osmotic pressure[90]. Therefore, we argue

that the higher expression of several transcripts similar to key secondary cell wall

biosynthesis and cell wall thickening genes in Equatorial samples also suggests plant

responses to water stress. Corroborating evidences of the relevance of water stress in the

Equatorial environmental conditions, we identified gene sequence divergence between

populations north and south of the NEESA in two putative osmotic stress-responsive

regulatory transcription factors (*RAP2.4* and *ZFN1*), one of which also induces the expression

of aquaporins (*RAP2.4*). These findings are additionally supported by the divergence in traits

related to water balance between plants from Equatorial and Subtropical sampling sites, such

as leaf size, vessel diameter, leaf angle and transpiration and stomatal conductance rates (Fig

2, S2 Fig).

3.2   Latitudinal variation in light quality and intensity may shape non-neutral diversity

Contrasting seasonal fluctuations in the photoperiod and in the light quality

throughout the year between low and high latitudes[91] (Table 1) likely influence the

differential expression of putative UV-inducible antioxidant and photodamage repair genes.

The adaptive relevance of this is supported by the natural selection sign found in a transcript

similar to a key gene to the biosynthesis of anthocyanin, which confers protection to UV-B

irradiation (*UDP-GLUCOSYL TRANSFERASE*). Divergent morphological traits between

Equatorial and Subtropical plants grown in the common garden experiment, including leaf

inclination angle and stem light reflectance, corroborate with the observed differential gene

expression and genomic signatures for positive selection (Fig 2 and 3).

3.3   Low water and low light availability may affect photosynthesis and cellular

respiration

In response to abiotic stress conditions such as drought, heat and high light, plants

optimise the use of light energy and minimise photooxidative damage, reducing the

photosynthetic activity by repressing light-harvesting and photosystem-component genes[92–

100]. We argue that the lower expression of photosynthesis genes in Equatorial than in

Subtropical samples likely reinforces the relevance of water stress in shaping divergent

phenotypes in the field, but also may result in an enhanced absorption of light energy in

Subtropical plants. Accordingly, we identified an increased expression of transcripts

associated with chlorophyll biosynthesis in the DET-St set and with chlorophyll breakdown

and leaf senescence in the DET-Eq set. The degradation of chlorophyll followed by leaf

senescence allows the remobilisation of nutrients and reduces the water loss through leaf

transpiration, contributing to water balance and to drought tolerance[101]. In addition, we

detected a putative genetic signature of selection in a transcript similar to a *TPR* protein,

required for chlorophyll synthesis and for the biogenesis of the photosynthetic apparatus[79].

Thus, we suggest that differential seasonality in light and water availability at subtropical and

equatorial latitudes may be involved in the divergence of non-neutral variability in the

species. Corroborating with the genomic approaches, we observed divergence of functional

traits related to water use and to the absorbance of light energy in plants from contrasting

latitudes cultivated under the same environmental conditions (Fig 2-3, S2-S3 Fig).

The mitochondrial activity is deeply connected to the photosynthesis and to

chloroplasts function since it generates ATP for carbohydrate synthesis, plays a role in the

protection of chloroplasts against photoinhibition, participates in the dissipation of reducing

equivalents and exchanges metabolites with chloroplasts during photorespiration[102,103].

Cellular respiration also provides ATP and carbon compounds for secondary metabolism,

playing a fundamental role in responses to abiotic stresses, including drought[102,104]. We

suggest that the higher levels of cellular respiration genes in Equatorial than in Subtropical

samples, may be a consequence of the reduced expression of photosynthesis genes and

enhanced energetic demand caused by water stress.

3.4 Tidal amplitude variation with latitude may act as a diverging force along the

Atlantic coast of South America

Tidal amplitude is markedly reduced with increasing latitude along the Atlantic coast

of South America, ranging from greater than 4 m at low latitudes to less than 1 m at high

latitudes[105]. The wide variation in tidal amplitude exposes trees to contrasting durations of

hypoxia. The identification of a candidate SNP locus with putative sign of positive selection

in a transcription factor induced by oxygen deprivation (*HRA1*) may indicate that differential

tidal variation act as a diverging selective force between northerly Equatorial and southerly

Subtropical populations of *A. schaueriana*. The *HRA1* putative homolog also showed a 1.75-

fold higher expression in Subtropical leaves relative to Equatorial leaves under field

conditions. However, we did not detect evidences at the phenotypic level in the common

garden experiment corroborating with the suggested relevance of this environmental variable.

3.5 Climate change and conservation implications of the results

The functional divergence observed in this work might differentially affect the

sensitivity of populations of *A. schaueriana* to a rapidly changing climate. Although there is

no evidence of mangrove expansion at its southernmost range limit on the Atlantic coast of

South America[28], researchers suggest that populations could expand polewards in the near

future, benefitting from the increased air and ocean temperatures and greenhouse gas

concentrations[106]. We expect that the more acquisitive phenotypic traits exhibited in the

common garden experiment by *A. schaueriana* seedlings from the subtropical range edge may

indeed be favourable for growth under increased temperatures and rainfall[8]. However, due

to the low tidal variation (< 1 m) observed in the southernmost mangrove distribution edge in

the Atlantic coast of South America, a greater relocation of forests is required for trees to

keep pace with the rising sea-level[107]. In this context, dense human coastal occupations

combined with the narrow intertidal zones, characteristic of this region, make mangroves at

higher latitudes more vulnerable to habitat loss than in the equatorial region of the Atlantic

coast of South America. Conversely, despite having wider coastal plains potentially available

for expansion, we expect that populations of *A. schaueriana* occurring north of the NEESA

will be threatened by the increased mean temperatures and the decreased precipitation during

the El-Niño Southern Oscillation events[8]. The critical temperature threshold is likely to be

reached more frequently, intensifying drought events in this region, possibly reducing carbon

assimilation and productivity[108] and, in extreme cases, causing biomass loss triggered by

cavitation or carbon starvation[109,110]. For the definition of short-term mangrove

conservation plans, such as reforestation or restoration of degraded areas on the Atlantic coast

of South America, we recommend that populations occurring north and south of the NEESA

should warrant attention as distinct conservation management units[111].

3.6   Concluding remarks

Based on the combined analysis of our results, we argue that carbon acquisition in the

equatorial region may be limited by the combination of a longer exposure to hypoxia, a higher

vapour pressure deficit (VPD) and a higher solar irradiance, especially during the hot and dry

season, whereas in the subtropical region, limitations in carbon gain may result from lower

solar irradiance levels, lower temperature and shorter photoperiod during winter (Fig 4).

These environmental differences between equatorial and subtropical latitudes seem to be

shaping gene expression in the field, and remarkably, may have shaped both allele

frequencies in genes responding to these variables and morphophysiological traits observed in

individuals of *A. schaueriana* (Fig 4). The emergence of this adaptive divergence is facilitated

by the limited gene flow identified between populations north and south of the NEESA,

possibly driven by the movement of the major ocean currents that are active on the Atlantic

coast of South America (Fig 1). Because a similar north-south divergence of neutral

molecular markers is also observed for other mangrove and mangrove-associated species on

the Atlantic coast of South America[22,23,25], it is also plausible that the environmental

drivers of divergence in *A. schaueriana* play a role in the divergence of other species. In

addition to revealing that these populations are genetically and functionally distinct

units[112], we provide an in-depth empirical evaluation of the intraspecific variation of a

long-living non-model tree. We foresee that these results should allow a clearer prediction of

how *A. schaueriana* and potentially other mangrove species may respond to the current global

climate changes by accounting for phenotypically[7] and genetically[113] informed mangrove

distribution modelling.

Fig 4. Graphical summary of the integration of ecological and high-throughput genomics data performed in this work.
Key oceanographic and climatic factors differ markedly between equatorial and subtropical latitudinal distribution extremes of the distribution of the neotropical mangrove tree, *Avicennia schaueriana*, possibly shaping the diversity of genotypes and phenotypes of the species. To address this issue, we examined the effects of these contrasting environments on the overall gene expression, its morphophysiological effects in a common garden experiment and its genomic effects through natural selection detection tests based on single nucleotide polymophic (SNP) loci. Plants from equatorial and subtropical latitudes showed key divergences related to the use of water and to the acquisition of carbohydrates both in the field and in common garden conditions. In addition, a north-south genetic divergence was observed in all genotyped SNP loci. We also identified signatures of differential selective pressures on specific loci possibly related to the accumulation of anthocyanin (*UDP-GLUCOSYLTRANSFERASE*), in response to osmotic stress (*RAP2.4* and *ZFN1*), photosynthesis (*TPR*) and hypoxia (*HRA1*). The molecular and ecologic divergences observed through three independent approaches may relate to the environmental factors that strongly differ between contrasting latitudes in which the species are found. These findings highlight the power of using multidisciplinary approaches to the study of adaptation in species with little basic biological information such as tropical trees. VPD: Atmospheric vapour pressure deficit; E: transpiration rate; g_s: stomatal conductance; R_L: xylem vessel lumen resistivity; K_H: xylem vessel conductivity.

4.   Materials and Methods

4.1   Propagules sampling sites

Ten mature propagules were collected from ten *A. schaueriana* mother trees belonging

to each of two genetically divergent natural populations described in a previous work[22]: one

north and one south of the NEESA (Fig 1). Trees were at least 100 m apart from each other at

each of the two contrasting sampling points: (1) the southernmost limit of the distribution of

American mangroves, in the subtropical region, and (2) one equatorial site in one of the

largest macrotidal mangrove forests in the world[29,30], closer to the northernmost limit of

the species range (Fig 1). We refer to the former as the 'Subtropical site' and to the latter as

the 'Equatorial site' throughout this work. A characterization of each of these sites can be

found in Table 1. Further details about these areas can be found in the Supplementary

Information file (S1 Method, S1 Fig).

To better characterise the climate from the Equatorial and Subtropical sampling sites

we have downloaded temperature, relative humidity, rainfall and solar irradiance datasets

(2008-2017) from the INMET database (the Meteorological Institute of Brazil) (S1 Fig) and

the environmental data from BioClim[31] and MARSPEC[32].

4.2   Seedlings germination and growth in a common garden experiment

Propagules of *A. schaueriana* sampled from the Equatorial and Subtropical sites

germinated on germination trays filled with mangrove soil, under shaded conditions, in open

air being irrigated daily with 1:10 (v:v) sea water:tap water solution[33]. After two months,

44 similar-sized seedlings from each sampling site — with an average height of 18 cm, most

with three leaf pairs and senesced cotyledons — were transplanted to 6 L pots filled with a

mixture of topsoil and sand (1:1 ratio). Seedlings were cultivated for seven months under the

same environmental conditions in a glasshouse at the University of Campinas, São Paulo,

Brazil (22°49' S 47°04' W). Automatic sensors coupled with a data logger (Onset Computer

Corp., Bourne, MA, USA) measured the atmospheric humidity and temperature inside the

glasshouse every 15 minutes. Seedlings were irrigated by an automated system twice per day

(10 a.m. and 5 p.m.), with a 3-minute fresh water spray. Twice a week nutrients were

manually added to the soil using 600 mL of 0.4X Hoagland solution with 15.0 g L^{-1} NaCl per

pot. All pots were rotated to reduce the effects of eventual environmental heterogeneity. The

monitored environmental conditions in the glasshouse differed markedly from both sites of

origin (S9 Fig 9). Thus, none of the individuals benefitted from environmental conditions that

corresponded to their original sampling site.

4.3   Comparative ecophysiology in a common garden experiment

The light reflectance of stems was measured in 10 plants from each sampling site

using a USB4000 spectrophotometer (OceanOptics, Inc., Dunedin, FL, USA) coupled to a

deuterium-halogen light source (DH-2000; OceanOptics, Inc., Ostfildern, Germany), using a

light emission range from 200-900 nm. To generate a daily curve of photosynthesis, stomatal

conductance and transpiration rates, five six-month-old individuals from each sampling site

were used for measurements every two or two and a half hours, in two different days, using a

Li-Cor 6400 XT (Li-Cor Corp., Lincoln, NE, USA).

After harvest, plants that did not present flowers or flower buds were split into three

parts: leaves, stems and roots. All plant parts were washed with distilled water, dried for 7

days at 70 °C and weighted. A quantitative analysis of key morphological traits was

performed using three plants from each site. The individual leaf area, total leaf area and leaf

lamina angle were measured per plant through photographs analyses using the ImageJ

software[34]. The specific leaf area (SLA, cm^2 leaf area kg^{-1} leaf dry biomass) was also

calculated for these same samples. Stems chunks were fixed in FAA (Formaldehyde Alcohol

Acetic acid) and stored in 70% alcohol for use in a wood anatomy analysis. Stems were cut

into transverse sections with a thickness of 30 μm. Sections were stained with a mixture of

1% Astra Blue and 50% alcohol (1:1) followed by 1% Safranin O (diluted in distilled water).

Micrographs of the slides were taken using an Olympus BX51 microscope coupled to an

Olympus DP71 camera (Olympus Corp., Tokyo, Japan). The following wood traits were

quantified using ImageJ[34] and R 4.0: vessel lumen area (A), vessel density in the xylem

(number of vessels/xylem area), proportion of solitary vessels (number of solitary

vessels/total number of vessels), vessel grouping index (mean number of vessels/vessel

grouping), proportion of vessel lumen area in xylem (vessel lumen area/xylem area) and

vessel lumen area in sapwood (vessel lumen area/sapwood area). The vessel arithmetic

diameter (D), vessel hydraulic conductivity (K_H) and lumen resistivity (R_L) were estimated

according to Scholz *et al.*[35].

Statistical analyses were performed in R 4.0, with a significance level of 5%, and two-

group comparisons were made using the Mann-Whitney-Wilcoxon unpaired test for non-

parametric distributions and Student's T-test for parametric distributions. Multiple-group

comparisons were conducted using one-way analysis of variance (ANOVA) with post hoc

Tukey honest significant difference (HSD) tests.

4.4   Collection of plant material for RNA extraction and RNA-sequencing

Plant material for RNA-Seq was collected in the "Propagules sampling sites"

subsection. Three adult trees were sampled in each sampling site from July-August of 2014,

at the end of winter in the Subtropical sampling site and at the beginning of the dry season in

the Equatorial sampling site (Table 2, S6 Fig). At each site, we sampled leaves, stems and

flowers of trees located at least 100 m apart from each other, from 11:00 am to 4:00 pm,

during the low tide, at different altitudes in the intertidal zone (Table 2). Plant material was

washed with sodium hypochlorite solution (0.2%), cut with a sterile slide and immediately

stored in RNAlater (Ambion Inc., Austin, TX, USA).

Total RNA extraction was performed using a protocol developed by Oliveira et

al.[36]. RNA integrity and purity were verified by agarose gel electrophoresis and using a

NanoVue spectrophotometer (GE Healthcare Life Sciences, Chalfont, Buckinghamshire, UK).

lllumina TruSeq RNA Sample Preparation protocol kits (Illumina Inc., San Diego, CA, USA)

were used to construct enriched complimentary DNA (cDNA) libraries. The quality of all

libraries was assessed using the Agilent DNA 1000 chip and Agilent 2100 Bioanalyzer

(Agilent Technologies, Santa Clara, USA). cDNA libraries concentrations were quantified by

qPCR using the Sequencing Library qPCR Quantification kit (Illumina Inc.), followed by

sequencing using TruSeq SBS Paired End kits (Illumina Inc.) and a Genome Analyzer IIx

sequencer (Illumina Inc.).

4.5   Assembly and characterization of the first reference transcriptome of **A.**

schaueriana

Adapter sequences were trimmed and 72 bp paired end reads were filtered by quality

(phred score ≥ 20 for at least 70% of read length) using the NGS QC Toolkit 2.3[37]. High-

quality reads were used in the subsequent transcriptome assembly in the CLC Genomics

Workbench (https://www.qiagenbioinformatics.com/). We used the default settings and only

set the distance between read pairs (300-500 bp), and the k-mer size to 45 bp.

Reads were mapped to the assembled transcripts using bowtie 1[38] in the single-read

mode using default parameters. We then removed transcripts that did not present read-

mapping support. Functional annotation was performed using the blastx algorithm, available

in blast+ v. 2.2.31[39], with an e-value threshold of 1^{-10}. The NCBI RefSeq[40], The

Arabidopsis Information Resource (TAIR)[41] and the NCBI non-redundant (nr) databases

were used as references. To remove contaminant transcripts, we excluded all transcripts that

were exclusively similar to non-plant sequences. Protein family domains were identified

using HMMER3[42], which iteratively searched all assembled sequences against the Pfam

database. To assign Gene Ontology (GO) terms to transcripts we used the *Arabidopsis*

thaliana gene association file from the Gene Ontology Consortium[43] and retrieved the

information for transcripts with a similar coding sequence in the genome of *A. thaliana*.

Redundant transcripts were clustered using CD-HIT-EST v. 4.6.1[44], using the local

alignment mode, with 95% identity and 70% coverage of the shortest sequence as thresholds.

Open reading frames (ORF) in putative protein coding transcripts were identified using the

Transdecoder tool (http://transdecoder.sf.net). We reduced redundancy of transcripts in the

final assembly by keeping for each CD-HIT-EST cluster either the sequence with the longest

ORF or simply the longest sequence, in the absence of sequences containing ORF.

The completeness of the final transcriptome was assessed using BUSCO[45].

Additionally, a reciprocal blastn alignment[39] using an e-value threshold of 10^{-10} and a

minimum alignment length of 100 nucleotides with at least 70% identity was used to compare

the *A. schaueriana* and other publicly available transcriptomes of congeneric species.

4.6 ***Comparative transcriptomics using RNA-sequencing data***

Tissue-specific count data was obtained from the number of reads uniquely mapped to

each transcript of the non-redundant transcriptome using bowtie 1[38]. Count data were

normalised using the EdgeR Bioconductor package[46]. Normalised data was used to detect

transcripts with significantly different counts between tissue-specific samples of trees at the

Equatorial and Subtopical sampling sites, using the exact test for negative binomial

distribution, with an adjusted P-value threshold of 0.05. GO term enrichment analyses of the

differentially expressed transcripts (DET) were performed using the GOseq package[47],

which accounts for length bias. We chose the default Wallenius approximation method and a

P-value threshold of 0.05.

Differential expression results were verified using reverse transcription real-time PCR

(qRT-PCR) (see S2 Methods).

4.7 ***Detection of* A. schaueriana *candidate loci under natural selection using genome-***

wide SNP data

We sampled adult plants at 10 locations, spanning most of *A. schaueriana* geographic

range, complementing previous population genetic studies[22] by adding the southernmost

range limit[28] (Fig 1, S1 Table). We isolated DNA from leaves sampled from 79 trees using

the DNeasy Plant Mini Kit (QIAGEN) and NucleoSpin Plant II (Macherey Nagel) following

the manufacturer's instructions. DNA quality and quantity were assessed using 1% agarose

electrophoresis and the QuantiFluor® dsDNA System in a Quantus™ fluorometer (Promega).

Nextera-tagmented, reductively-amplified DNA (nextRAD) libraries were constructed by

SNPsaurus (SNPsaurus, LLC), allowing the genotyping of SNP loci across the genome in a

consistent manner by using selective primer sequences[48]. Genomic DNA fragmentation and

short adapter ligation were performed with Nextera reagent (Illumina, Inc), followed by

amplification, which utilised one of the primers matching the adapter and extending nine

arbitrary nucleotides into the genomic DNA. Thus, the resulting amplicons were fixed at the

selective end and their lengths depended on the initial Nextera fragmentation, leading to a

consistent genotyping of the amplified loci. NextRAD libraries were then sequenced in a

HiSeq 2500 (Illumina, Inc.) with 100-bp single-end chemistry following the manufacturer's

protocol. Genotyping-by-sequencing used custom scripts (SNPsaurus, LLC), which created a

reference catalogue of abundant reads across the combined set of samples. Read mapping to

this reference allowed two mismatches. Subsequently, loci presenting sequence variation were

filtered for biallelic loci and presence in at least 10% of the total samples. Following

assembly, mapping and SNP identification, a second filtering step was performed by allowing

no more than 65% of missing data, a Phred score > 30, 8x minimum coverage, only one SNP

per loci and a minor allele frequency ≥ 0.05 using vcftools v0.1.12b[49]. To reduce false SNP

inclusion rates due to paralogy or low-quality genotype calls, we used a maximum read

coverage of 56, which resulted from the product of the average read depth and 1.5 standard

deviation.

After excluding plants morphologically identified as *A. schaueriana* with genomic

signs of hybridization with *A. germinans*, we described the overall genetic structure in *A.*

schaueriana considering both neutral and non-neutral loci in a multivariate model-free

method, the discriminant analysis of principal components (DAPC)[50], and in the

ADMIXTURE 1.3.0 program[51]. For DAPC analyses, we considered the number of groups

varying from 1 to 50 and the Bayesian information criteria for inferring the number of groups

(K), in addition to using the optim.a.score function to avoid over-fitting during the

discrimination steps. For the ADMIXTURE analyses, we performed three separate runs for K

varying between 1 and 15 and using the block-relaxation method for point estimation;

computing was terminated when estimates increased by < 0.0001. The K-value that best

described our data was determined by cross-validation.

To detect signs of natural selection, we used two methods to minimise false-positive

results. LOSITAN[52], which uses the FDIST2 method[53], assuming an infinite allele model

of mutation, using a confidence interval of 0.99, a false-discovery rate (FDR) of 0.1, the

neutral mean FST and forcing the mean FST options, and pcadapt 3.0.4[54], which

simultaneously identifies the population structure and the loci excessively related to this

structure, using an FDR threshold of 0.1.

The sequences in which loci with putative evidence of selection were identified

simultaneously by pcadapt and by five independent runs of LOSITAN were searched within

expressed regions of the reference transcriptome. A reciprocal alignment between the short

sequences obtained through nextRAD (75 bp) and the longer expressed sequences assembled

from RNA-Seq data ($\approx$ 300-11600 bp) was performed. The alignment was conducted using

the blastn algorithm from the blast+ software v. 2.2.31[39], with a threshold of at least 50

aligned nucleotides, a maximum of one mismatch and no gaps.

5.   Acknowledgements

We thank Professors Dr. Michel A. G. Vincentz and Dr. Juliana L. S. Mayer for

advising and making available their laboratory for our training and performing of,

respectively, RNA extraction procedures and slides preparation used in the analyses of wood

anatomy. We acknowledge Ilmarina C. de Menezes and Stephanie K. Bajay for assistance in

the field. We thank Stephanie K. Bajay for also giving assistance in the germination of the

propagules used in the common garden experiment. And we thank Greg Mellers and

Alessandro A. Pereira for critically reading the manuscript. M.V.C and G.M.M. received

fellowships from the São Paulo Research Foundation (FAPESP 2013/26793-7, 2013/08086-1

and 2014/22821-9). M.V.C. received fellowship from the Coordination for the Improvement

of Higher Education Personnel (CAPES 8084/2015-07, Biocomputational Biology Program).

G.M.M. received grant from the Brazilian National Council for Scientific and Technological

Development (CNPq 448286/2014-9). A.P.S. was supported by grant from CAPES

(Biocomputational Biology Program), and a received a research fellowship from CNPq

(309661/2014-5).

6.   References

1.   Hereford J. A Quantitative Survey of Local Adaptation and Fitness Trade☐Offs. Am

Nat. 2009;173: 579–588. doi:10.1086/597611

2.   Kawecki TJ, Ebert D. Conceptual issues in local adaptation. Ecol Lett. 2004;7: 1225–

1241. doi:10.1111/j.1461-0248.2004.00684.x

3.   Leimu R, Fischer M. A meta-analysis of local adaptation in plants. PLoS One. 2008;3:

1–8. doi:10.1371/journal.pone.0004010

4.   Albert CH, Grassein F, Schurr FM, Vieilledent G, Violle C. When and how should

intraspecific variability be considered in trait-based plant ecology? Perspect Plant Ecol

Evol Syst. 2011;13: 217–225. doi:10.1016/j.ppees.2011.04.003

5.   Savolainen O, Lascoux M, Merilä J. Ecological genomics of local adaptation. Nat Rev

Genet. Nature Publishing Group; 2013;14: 807–20. doi:10.1038/nrg3522

6.   Barrett RDH, Hoekstra HE. Molecular spandrels: tests of adaptation at the genetic

level. Nat   Rev   Genet.   Nature   Publishing   Group;   2011;12:   767–780.

doi:10.1038/nrg3015

7.   Moran E V., Hartig F, Bell DM. Intraspecific trait variation across scales: Implications

for understanding global change responses. Glob Chang Biol. 2016;22: 137–150.

doi:10.1111/gcb.13000

8.   Pachauri RK, Meyer LA, editors. Climate Change 2014: Synthesis Report.

Contribution of Working Groups I, II and III to the Fifth Assessment Report of the

Intergovernmental Panel on Climate Change. IPCC. Geneva, Switzerland: IPCC; 2014.

9.   Loarie SR, Duffy PB, Hamilton H, Asner GP, Field CB, Ackerly DD. The velocity of

climate change. Nature. Nature Publishing Group; 2009;462: 1052–1055.

doi:10.1038/nature08649

10.   Osland MJ, Feher LC, Griffith KT, Cavanaugh KC, Enwright NM, Day RH, et al.

Climatic controls on the global distribution, abundance, and species richness of

mangrove forests. Ecol Monogr. 2017;87: 341–359. doi:10.1002/ecm.1248

11.   Lovelock CE, Cahoon DR, Friess DA, Guntenspergen GR, Krauss KW, Reef R, et al.

The vulnerability of Indo-Pacific mangrove forests to sea-level rise. Nature. 2015;526:

559–563. doi:10.1038/nature15538

12.   Nascimento Jr. WR, Souza-Filho PWM, Proisy C, Lucas RM, Rosenqvist A. Mapping

changes in the largest continuous Amazonian mangrove belt using object-based

classification of multisensor satellite imagery. Estuar Coast Shelf Sci. 2013;117: 83–

93. doi:10.1016/j.ecss.2012.10.005

13.   Shearman P, Bryan J, Walsh JP. Trends in Deltaic Change over Three Decades in the

Asia-Pacific Region. J Coast Res. 2013;29: 1169–1183. doi:10.2112/JCOASTRES-D-

12-00120.1

14.   Cavanaugh KC, Kellner JR, Forde AJ, Gruner DS, Parker JD, Rodriguez W, et al.

Poleward expansion of mangroves is a threshold response to decreased frequency of

extreme cold events. Proc Natl Acad Sci U S A. 2014;111: 723–7.

doi:10.1073/pnas.1315800111

15.   Saintilan N, Wilson NC, Rogers K, Rajkaran A, Krauss KW. Mangrove expansion and

salt marsh decline at mangrove poleward limits. Glob Chang Biol. 2014;20: 147–157.

doi:10.1111/gcb.12341

16.   Duke N, Kovacs J, Griffiths A, Preece L, J. E. Hill D, Van Oosterzee P, et al. Large-

scale dieback of mangroves in Australia. Marine and Freshwater Research. 2017.

doi:10.1071/MF16322

17.   Eslami-Andargoli L, Dale P, Sipe N, Chaseling J. Mangrove expansion and rainfall

patterns in Moreton Bay, Southeast Queensland, Australia. Estuar Coast Shelf Sci.

Elsevier Ltd; 2009;85: 292–298. doi:10.1016/j.ecss.2009.08.011

18.   Alongi DM. The Impact of Climate Change on Mangrove Forests. Curr clim Chang

Rep. 2015;4: 516. doi:10.1007/s40641-015-0002-x

19.   McKee KL, Cahoon DR, Feller IC. Caribbean mangroves adjust to rising sea level

through biotic controls on change in soil elevation. Glob Ecol Biogeogr. 2007;16: 545–

556. doi:10.1111/j.1466-8238.2007.00317.x

20.   López-Medellín X, Ezcurra E, González-Abraham C, Hak J, Santiago LS, Sickman JO.

Oceanographic anomalies and sea-level rise drive mangroves inland in the Pacific coast

of Mexico. J Veg Sci. 2011;22: 143–151. doi:10.1111/j.1654-1103.2010.01232.x

21.   Doyle TW, Krauss KW, Conner WH, From AS. Forest Ecology and Management

Predicting the retreat and migration of tidal forests along the northern Gulf of Mexico

under sea-level rise. For Ecol Manage. 2010;259: 770–777.

doi:10.1016/j.foreco.2009.10.023

22.   Mori GM, Zucchi MI, Souza AP. Multiple-Geographic-Scale Genetic Structure of Two

Mangrove Tree Species: The Roles of Mating System, Hybridization, Limited
Dispersal and Extrinsic Factors. PLoS One. 2015;10: e0118710.
doi:10.1371/journal.pone.0118710

23. Francisco PM, Mori GM, Alves FM, Tambarussi E, Souza AP De. Population genetic
structure, introgression, and hybridization in the genus Rhizophora along the Brazilian
coast. Ecol Evol. 2018; 1–14. doi:10.1002/ece3.3900

24. Pil MW, Boeger MRT, Muschner VC, Pie MR, Ostrensky A, Boeger W a. Postglacial
north-south expansion of populations of Rhizophora mangle (Rhizophoraceae) along
the Brazilian coast revealed by microsatellite analysis. Am J Bot. 2011;98: 1031–1039.
doi:10.3732/ajb.1000392

25. Takayama K, Tateishi Y, Murata J, Kajita T. Gene flow and population subdivision in
a pantropical plant with sea-drifted seeds Hibiscus tiliaceus and its allied species:
Evidence from microsatellite analyses. Mol Ecol. 2008;17: 2730–2742.
doi:10.1111/j.1365-294X.2008.03799.x

26. Savolainen O, Pyhäjärvi T, Knürr T. Gene Flow and Local Adaptation in Trees. Annu
Rev Ecol Evol Syst. 2007;38: 595–619.
doi:10.1146/annurev.ecolsys.38.091206.095646

27. Andrews KR, Good JM, Miller MR, Luikart G, Hohenlohe PA. Harnessing the power
of RADseq for ecological and evolutionary genomics. Nat Rev Genet. Nature
Publishing Group; 2016;17: 81–92. doi:10.1038/nrg.2015.28

28. Soares MLG, Estrada GCD, Fernandez V, Tognella MMP. Southern limit of the
Western South Atlantic mangroves: Assessment of the potential effects of global
warming from a biogeographical perspective. Estuar Coast Shelf Sci. Elsevier Ltd;
2012;101: 44–53. doi:10.1016/j.ecss.2012.02.018

29.   Souza-Filho PWM, Cohen MCL, Lara RJ, Lessa GC, Koch B, Behling H. Holocene

831   Coastal Evolution and Facies Model of the Bragança Macrotidal Flat on the Amazon

832   Mangrove Coast, Northern Brazil. J Coast Res. 2006; 306–310.

833   30.   Kjerfve B, Perillo GME, Gardner LR, Rine JM, Dias GTM, Mochel FR. Chapter

834   Twenty Morphodynamics of muddy environments along the Atlantic coasts of North

835   and South America. Proc Mar Sci. Elsevier; 2002;4: 479–532. doi:10.1016/S1568-

836   2692(02)80094-8

837   31.   Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. Very high resolution

838   interpolated climate surfaces for global land areas. Int J Climatol. 2005;25: 1965–1978.

839   doi:10.1002/joc.1276

840   32.   Sbrocco EJ, Barber PH. MARSPEC: ocean climate layers for marine spatial ecology.

841   Ecology. 2013;94: 979–979. doi:10.1890/12-1358.1

842   33.   Reef R, Winter K, Morales J, Adame MF, Reef DL, Lovelock CE. The effect of

843   atmospheric carbon dioxide concentrations on the performance of the mangrove

844   Avicennia germinans over a range of salinities. Physiol Plant. 2014;154.

845   doi:10.1111/ppl.12289

846   34.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image

847   analysis.   Nat   Methods.   Nature   Publishing   Group;   2012;9:   671–675.

848   doi:10.1038/nmeth.2089

849   35.   Scholz A, Klepsch M, Karimi Z, Jansen S. How to quantify conduits in wood? Front

850   Plant Sci. 2013;4: 1–11. doi:10.3389/fpls.2013.00056

851   36.   Oliveira RR, Viana AJC, Reátegui ACE, Vincentz MGA. An efficient method for

852   simultaneous extraction of high-quality RNA and DNA from various plant tissues.

853   Genet Mol Res. 2015;14: 18828–18838. doi:10.4238/2015.December.28.32

854   37.   Patel RK, Jain M. NGS QC toolkit: A toolkit for quality control of next generation

855   sequencing data. PLoS One. 2012;7. doi:10.1371/journal.pone.0030619

856   38.   Langmead B, Trapnell C, Pop M, Salzberg S. Ultrafast and memory-efficient

857   alignment of short DNA sequences to the human genome. Genome Biol. 2009;10: R25.

858   doi:10.1186/gb-2009-10-3-r25

859   39.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al.

860   BLAST+: architecture and applications. BMC Bioinformatics. 2009;10: 421.

861   doi:10.1186/1471-2105-10-421

862   40.   O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, et al.

863   Reference sequence (RefSeq) database at NCBI: Current status, taxonomic expansion,

864   and functional annotation. Nucleic Acids Res. 2016;44: D733–D745.

865   doi:10.1093/nar/gkv1189

866   41.   Berardini TZ, Reiser L, Li D, Mezheritsky Y, Muller R, Strait E, et al. The Arabidopsis

867   Information Resource□: Making and Mining the " Gold Standard " Annotated

868   Reference Plant Genome. Genesis. 2015;53: 474–485. doi:10.1002/dvg.22877

869   42.   Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR, et al. Pfam: The

870   protein families database. Nucleic Acids Res. 2014;42: 222–230.

871   doi:10.1093/nar/gkt1223

872   43.   Blake JA, Christie KR, Dolan ME, Drabkin HJ, Hill DP, Ni L, et al. Gene ontology

873   consortium: Going forward. Nucleic Acids Res. 2015;43: D1049–D1056.

874   doi:10.1093/nar/gku1179

875   44.   Li W, Godzik A. Cd-hit: A fast program for clustering and comparing large sets of

876   protein or nucleotide sequences. Bioinformatics. 2006;22: 1658–1659.

877   doi:10.1093/bioinformatics/btl158

878   45.   Simão FA, Waterhouse RM, Ioannidis P, Kriventseva E V., Zdobnov EM. BUSCO:

879   Assessing genome assembly and annotation completeness with single-copy orthologs.

880   Bioinformatics. 2015;31: 3210–3212. doi:10.1093/bioinformatics/btv351

881   46.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for

882   differential expression analysis of digital gene expression data. Bioinformatics.

883   2010;26: 139–140. doi:10.1093/bioinformatics/btp616

884   47.   Young MD, Wakefield MJ, Smyth GK, Oshlack A. Gene ontology analysis for RNA-

885   seq: accounting for selection bias. Genome Biol. 2010;11. doi:10.1186/gb-2010-11-2-

886   r14

887   48.   Russello MA, Waterhouse MD, Etter PD, Johnson EA. From promise to practice:

888   pairing non-invasive sampling with genomics in conservation. PeerJ. 2015;3: e1106.

889   doi:10.7717/peerj.1106

890   49.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The

891   variant   call   format   and   VCFtools.   Bioinformatics.   2011;27:   2156–2158.

892   doi:10.1093/bioinformatics/btr330

893   50.   Jombart T, Devillard S, Balloux F. Discriminant analysis of principal components: a

894   new method for the analysis of genetically structured populations. BMC Genet.

895   BioMed Central Ltd; 2010;11: 94. doi:10.1186/1471-2156-11-94

896   51.   Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in

897   unrelated individuals. Genome Res. 2009;19: 1655–1664. doi:10.1101/gr.094052.109

898   52.   Antao T, Lopes A, Lopes RJ, Beja-Pereira A, Luikart G. LOSITAN: A workbench to

899   detect molecular adaptation based on a F_{st}-outlier method. BMC Bioinformatics.

900   2008;9: 1–5. doi:10.1186/1471-2105-9-323

53.   Beaumont MA, Nichols RA. Evaluating Loci for Use in the Genetic Analysis of

Population Structure. Proc R Soc B Biol Sci. 1996;263: 1619–1626.

doi:10.1098/rspb.1996.0237

54.   Luu K, Bazin E, Blum MGB. pcadapt: An R package to perform genome scans for

selection based on principal component analysis. Mol Ecol Resour. 2016;33: 67–77.

doi:10.1111/1755-0998.12592

55.   Wang Z, Gerstein M, Snyder M. RNA-Seq: a revolutionary tool for transcriptomics.

Nat Rev Genet. 2009;10: 57–63.

56.   Huang J, Lu X, Zhang W, Huang R, Chen S, Zheng Y. Transcriptome Sequencing and

Analysis of Leaf Tissue of Avicennia marina Using the Illumina Platform. PLoS One.

2014;9: e108785. doi:10.1371/journal.pone.0108785

57.   Lyu H, Li X, Guo Z, He Z, Shi S. De novo assembly and annotation of the Avicennia

officinalis L. transcriptome. Mar Genomics. Elsevier B.V.; 2017; 2–5.

doi:10.1016/j.margen.2017.07.002

58.   Krishnamurthy P, Mohanty B, Wijaya E, Lee D, Lim T, Lin Q, et al. Transcriptomics

analysis of salt stress tolerance in the roots of the mangrove Avicennia officinalis. Nat

Sci Reports. 2017;7: 1–19. doi:10.1038/s41598-017-10730-2

59.   Park S-Y, Yu J-W, Park J-S, Li J, Yoo S-C, Lee N-Y, et al. The Senescence-Induced

Staygreen Protein Regulates Chlorophyll Degradation. Plant Cell. 2007;19: 1649–

1664. doi:10.1105/tpc.106.044891

60.   Myouga F, Hosoda C, Umezawa T, Iizumi H, Kuromori T, Motohashi R, et al. A

Heterocomplex of Iron Superoxide Dismutases Defends Chloroplast Nucleoids against

Oxidative Stress and Is Essential for Chloroplast Development in Arabidopsis. Plant

Cell Online. 2008;20: 3148–3162. doi:10.1105/tpc.108.061341

61.   Wang S, Bai G, Wang S, Yang L, Yang F, Wang Y. Chloroplast RNA-Binding Protein

RBD1 Promotes Chilling Tolerance through 23S rRNA Processing in Arabidopsis.

PLoS Genet. 2016;12: 1–21. doi:10.1371/journal.pgen.1006027

62.   Goulas E, Schubert M, Kieselbach T, Kleczkowski LA, Gardestro P, Schro W. The

chloroplast lumen and stromal proteomes of Arabidopsis thaliana show differential

sensitivity to short- and long-term exposure to low temperature. Plant J. 2006;47: 720–

734. doi:10.1111/j.1365-313X.2006.02821.x

63.   Kim YO, Kim JS, Kang H. Cold-inducible zinc finger-containing glycine-rich RNA-

binding protein contributes to the enhancement of freezing tolerance in Arabidopsis

thaliana. Plant J. 2005;42: 890–900. doi:10.1111/j.1365-313X.2005.02420.x

64.   Hartung F, Suer S, Puchta H. Two closely related RecQ helicases have antagonistic

roles in homologous recombination and DNA repair in Arabidopsis thaliana. Proc Natl

Acad Sci. 2007;104: 18836–18841. doi:10.1073/pnas.0705998104

65.   Michaels SD, Ditta G, Gustafson-Brown C, Pelaz S, Yanofsky M, Amasino RM.

AGL24 acts as a promoter of flowering in Arabidopsis and is positively regulated by

vernalization. Plant J. 2003;33: 867–874. doi:10.1046/j.1365-313X.2003.01671.x

66.   Jung YJ, Melencion SMB, Lee ES, Park JH, Alinapon CV, Oh HT, et al. Universal

Stress Protein Exhibits a Redox-Dependent Chaperone Function in Arabidopsis and

Enhances Plant Tolerance to Heat Shock and Oxidative Stress. Front Plant Sci. 2015;6:

1–11. doi:10.3389/fpls.2015.01141

67.   Ogawa T, Pan L, Kawai-Yamada M, Yu L-H, Yamamura S, Koyama T, et al.

Functional Analysis of Arabidopsis Ethylene-Responsive Element Binding Protein

Conferring Resistance to Bax. Plant Physiol. 2005;138: 1436–1445.

doi:10.1104/pp.105.063586.1436

68.   Lin RC, Park HJ, Wang HY. Role of Arabidopsis RAP2.4 in regulating light- and

ethylene-mediated developmental processes and drought stress tolerance. Mol Plant.

The Authors 2007. All rights reserved.; 2008;1: 42–57. doi:10.1093/mp/ssm004

69.   Rae L, Lao NT, Kavanagh TA. Regulation of multiple aquaporin genes in Arabidopsis

by a pair of recently duplicated DREB transcription factors. Planta. 2011;234: 429–

444. doi:10.1007/s00425-011-1414-z

70.   Alexandersson E, Fraysse L, Sjövall-Larsen S, Gustavsson S, Fellert M, Karlsson M, et

al. Whole gene family expression and drought stress regulation of aquaporins. Plant

Mol Biol. 2005;59: 469–484. doi:10.1007/s11103-005-0352-1

71.   Nishizawa A, Yabuta Y, Shigeoka S. Galactinol and Raffinose Constitute a Novel

Function to Protect Plants from Oxidative Damage. Plant Physiol. 2008;147: 1251–

1263. doi:10.1104/pp.108.122465

72.   Zhang A, Ren H-M, Tan Y-Q, Qi G-N, Yao F-Y, Wu G-L, et al. S-Type Anion

Channels SLAC1 and SLAH3 Function as Essential Negative Regulators of Inward K$^+$

Channels and Stomatal Opening in Arabidopsis. Plant Cell. 2016;28: 949–965.

doi:10.1105/tpc.15.01050

73.   Liepman AH, Olsen LJ. Alanine aminotransferase homologs catalyze the

glutamate:glyoxylate aminotransferase reaction in peroxisomes of Arabidopsis. Plant

Physiol. 2003;131: 215–27. doi:10.1104/pp.011460

74.   Wolf JBW, Lindell J, Backstrom N. Speciation genetics: current status and evolving

approaches. Philos Trans R Soc B Biol Sci. 2010;365: 1717–1733.

doi:10.1098/rstb.2010.0023

75.   Ahrens CW, Rymer PD, Stow A, Bragg J, Dillon S, Umbers KDL, et al. The search for

loci under selection: trends, biases and progress. Mol Ecol. 2018;

doi:doi.org/10.1111/mec.14549

76.   Sakamoto H, Araki T, Meshi T, Iwabuchi M. Expression of a subset of the Arabidopsis

975   Cys2/His2-type zinc-finger protein gene family under water stress. Gene. 2000;248:

976   23–32. doi:10.1016/S0378-1119(00)00133-5

977   77.   Giuntoli B, Lee SC, Licausi F, Kosmacz M, Oosumi T, van Dongen JT, et al. A

978   Trihelix DNA Binding Protein Counterbalances Hypoxia-Responsive Transcriptional

979   Activation in Arabidopsis. PLoS Biol. 2014;12. doi:10.1371/journal.pbio.1001950

980   78.   Tohge T, Nishiyama Y, Hirai MY, Yano M, Nakajima JI, Awazuhara M, et al.

981   Functional genomics by integrated analysis of metabolome and transcriptome of

982   Arabidopsis plants over-expressing an MYB transcription factor. Plant J. 2005;42:

983   218–235. doi:10.1111/j.1365-313X.2005.02371.x

984   79.   Bohne A V., Schwenkert S, Grimm B, Nickelsen J. Roles of Tetratricopeptide Repeat

985   Proteins in Biogenesis of the Photosynthetic Apparatus [Internet]. International Review

986   of Cell and Molecular Biology. Elsevier Inc.; 2016. doi:10.1016/bs.ircmb.2016.01.005

987   80.   Reef R, Lovelock CE. Regulation of water balance in Mangroves. Ann Bot. 2015;115:

988   385–395. doi:10.1093/aob/mcu174

989   81.   Steyn WJ, Wand SJE, Holcroft DM, Jacobs G. Anthocyanins in vegetative tissues: a

990   proposed unified function in photoprotection. New Phytol. 2002;155: 349–361.

991   doi:10.1046/j.1469-8137.2002.00482.x

992   82.   Menezes MPM de, Berger U, Mehlig U. Mangrove vegetation in Amazonia: a review

993   of studies from the coast of Pará and Maranhão States, north Brazil. Acta Amaz.

994   2008;38: 403–420. doi:10.1590/S0044-59672008000300004

995   83.   De Alvarenga AMSB, Botosso PC, Soffiatti P. Stem growth and phenology of three

996   subtropical mangrove tree species. Brazilian J Bot. 2017; doi:10.1007/s40415-017-

997   0397-9

998   84.   Kazan K, Lyons R. The link between flowering time and stress tolerance. J Exp Bot.
999   2016;67: 47–60. doi:10.1093/jxb/erv441

85.   Stuart SA, Choat B, Martin KC, Holbrook NM, Ball MC. The role of freezing in
setting the latitudinal limits of mangrove forests. New Phytol. 2006;173: 576–583.
doi:10.1111/j.1469-8137.2006.01938.x

86.   Carlson JE, Holsinger KE, Prunier R. Plant responses to climate in the cape floristic
region of South Africa: Evidence for adaptive differentiation in the proteaceae.
Evolution (N Y). 2011;65: 108–124. doi:10.1111/j.1558-5646.2010.01131.x

87.   Saenger P, Snedaker SC. Pantropical trends in mangrove above-ground biomass and
annual litterfall. Oecologia. 1993;96: 293–299. doi:10.1007/BF00317496

88.   Krishnamurthy P, Jyothi-Prakash PA, Qin L, He J, Lin Q, Loh CS, et al. Role of root
hydrophobic barriers in salt exclusion of a mangrove plant Avicennia officinalis. Plant,
Cell Environ. 2014;37: 1656–1671. doi:10.1111/pce.12272

89.   Maurel C, Verdoucq L, Luu D-T, Santoni V. Plant Aquaporins: Membrane Channels
with Multiple Integrated Functions. Annu Rev Plant Biol. 2008;59: 595–624.
doi:10.1146/annurev.arplant.59.032607.092734

90.   Gall H Le, Philippe F, Domon J-M, Gillet F, Pelloux J, Rayon C. Cell Wall
Metabolism   in   Response   to   Abiotic   Stress.   Plants.   2015;4:   112–166.
doi:10.3390/plants4010112

91.   Pecot SD, Horsley SB, Battaglia MA, Mitchell RJ. The influence of canopy, sky
condition, and solar angle on light quality in a longleaf pine woodland. Can J For Res.
2005;35: 1356–1366. doi:10.1139/x05-069

92.   Kimura M, Yamamoto YY, Seki M, Sakurai T, Sato M, Abe T, et al. Identification of
Arabidopsis Genes Regulated by High Light – Stress Using cDNA Microarray.

Photochem Photobiol. 2003;77: 226–233. doi:https://doi.org/10.1562/0031-8655

93. Murchie EH, Hubbart S, Peng S, Horton P. Acclimation of photosynthesis to high

irradiance in rice: Gene expression and interactions with leaf development. J Exp Bot.

2005;56: 449–460. doi:10.1093/jxb/eri100

94. Wang D, Pan Y, Zhao X, Zhu L, Fu B, Li Z. Genome-wide temporal-spatial gene

expression profiling of drought responsiveness in rice. BMC Genomics. 2011;12.

doi:10.1186/1471-2164-12-149

95. Moumeni A, Satoh K, Kondoh H, Asano T, Hosaka A, Venuprasad R, et al.

Comparative analysis of root transcriptome profiles of two pairs of drought-tolerant

and susceptible rice near-isogenic lines under different drought stress. BMC Plant Biol.

2011;11. doi:10.1186/1471-2229-11-174

96. Bray EA. Genes commonly regulated by water-deficit stress in Arabidopsis thaliana. J

Exp Bot. 2004;55: 2331–2341. doi:10.1093/jxb/erh270

97. Zhang C, Zhang L, Zhang S, Zhu S, Wu P, Chen Y, et al. Global analysis of gene

expression profiles in physic nut (Jatropha curcas L.) seedlings exposed to drought

stress. BMC Plant Biol. 2015;15: 1–14. doi:10.1186/s12870-014-0397-x

98. Hasanuzzaman M, Nahar K, Alam MM, Roychowdhury R, Fujita M. Physiological,

biochemical, and molecular mechanisms of heat stress tolerance in plants. Int J Mol

Sci. 2013;14: 9643–9684. doi:10.3390/ijms14059643

99. Han F, Chen H, Li XJ, Yang MF, Liu GS, Shen SH. A comparative proteomic analysis

of rice seedlings under various high-temperature stresses. Biochim Biophys Acta.

Elsevier B.V.; 2009;1794: 1625–1634. doi:10.1016/j.bbapap.2009.07.013

100. Wong CE, Li Y, Labbe A, Guevara D, Nuin P, Whitty B, et al. Transcriptional

profiling implicates novel interactions between abiotic stress and hormonal responses

in Thellungiella, a close relative of Arabidopsis. Plant Physiol. 2006;140: 1437–1450.

doi:10.1104/pp.105.070508

101. Munné-Bosch S, Alegre L. Die and let live: Leaf senescence contributes to plant

survival under drought stress. Funct Plant Biol. 2004;31: 203–216.

doi:10.1071/FP03236

102. Millar AH, Whelan J, Soole KL, Day DA. Organization and Regulation of

Mitochondrial Respiration in Plants. Annu Rev Plant Biol. 2011;62: 79–104.

doi:10.1146/annurev-arplant-042110-103857

103. Raghavendra AS, Padmasree K. Beneficial interactions of mitochondrial metabolism

with photosynthetic carbon assimilation. Trends Plant Sci. 2003;8: 546–553.

doi:10.1016/j.tplants.2003.09.015

104. Atkin OK, Macherel D. The crucial role of plant mitochondria in orchestrating drought

tolerance. Ann Bot. 2009;103: 581–597. doi:10.1093/aob/mcn094

105. Schaeffer-Novelli Y, Cintrón-Molero G, Adaime RR, de Camargo TM. Variability of

mangrove ecosystems along the Brazilian coast. Estuaries. 1990;13: 204–218.

doi:10.1007/BF02689854

106. Godoy MDP, De Lacerda LD. Mangroves Response to Climate Change: A Review of

Recent Findings on Mangrove Extension and Distribution. Ann Brazilian Acad Sci.

2015;87: 651–667. doi:10.1590/0001-3765201520150055

107. Ellison JC. Vulnerability assessment of mangroves to climate change and sea-level rise

impacts. Wetl Ecol Manag. 2015;23: 115–137. doi:10.1007/s11273-014-9397-8

108. Saenger P, Moverly J. Vegetative phenology along the Queensland coastline. Proc Ecol

Soc Aust. 1985;13: 257–265.

109. Doughty CE, Metcalfe DB, Girardin CAJ, Amézquita FF, Cabrera DG, Huasco WH, et

al. Drought impact on forest carbon dynamics and fluxes in Amazonia. Nature.

2015;519: 78–82. doi:10.1038/nature14213

110. Rowland L, da Costa ACL, Galbraith DR, Oliveira RS, Binks OJ, Oliveira AAR, et al.

Death from drought in tropical forests is triggered by hydraulics not carbon starvation.

Nature. 2015;528: 119–122. doi:https://doi.org/10.1038/nature15539

111. Moritz C. Defining "Evolutionarily Significant Units" for conservation. Trends Ecol

Evol. 1994;9: 373–375. doi:10.1016/0169-5347(94)90057-4

112. Wee AKS, Mori GM, Lira-Medeiros CF, Núñez-Farfán J, Takayama K, Faulks L, et al.

The integration and application of genomic information in mangrove conservation.

Conserv Biol. 2018; 1–40. doi:10.1111/cobi.13140

113. Ikeda DH, Max TL, Allan GJ, Lau MK, Shuster SM, Whitham TG. Genetically

informed ecological niche models improve climate change predictions. Glob Chang

Biol. 2017;23: 164–176. doi:10.1111/gcb.13470

114. Alvares CA, Stape JL, Sentelhas PC, Gonçalves JL de M, Sparovek G. Köppen's

climate classification map for Brazil. Meteorol Zeitschrift. 2013;22: 711–728.

doi:10.1127/0941-2948/2013/0507

115. Forsythe WC, Rykiel EJ, Stahl RS, Wu H i., Schoolfield RM. A model comparison for

daylength as a function of latitude and day of year. Ecol Modell. 1995;80: 87–95.

doi:10.1016/0304-3800(94)00034-F

116. McRae GJ. A Simple Procedure for Calculating Atmospheric Water Vapor

Concentration. J Air Pollut Control Assoc. 1980;30: 394–394.

doi:10.1080/00022470.1980.10464362

1092

6. Supporting information captions

S1 Fig. Climate characterisation of Equatorial and Subtropical sampling sites. Points represent the monthly means of climate variables (2008-2017). Red line: Equatorial site (Tracuateua automatic station); blue line: Subtropical site (Santa Marta automatic station). Source: Brazilian National Institute of Meteorology (INMET). (A) Average temperature; (B) average maximum temperature; (C) average minimum temperature; (D) average solar irradiance; (E) total precipitation; (F) average air humidity; (G) average air vapour pressure deficit (VPD), estimated according to McRae(McRae, 1980).

S2 Fig. Daily curves of gas exchange in leaves of seven-months-old *Avicennia schaueriana* seedlings grown in a common-garden. Points represent the means ± standard error bars. Stars (*) represent rejection of the null hypothesis of the absence of a difference based on the unpaired Student's t-test at a significance threshold of 0.05 (n = 10 plants per group). Red line: equatorial origin; blue line: Subtropical origin. (A) Transpiration rate; (B) stomatal conductance; (C) rate of net CO_2 assimilation.

S3 Fig. *Avicennia schaueriana* from Equatorial and Subtropical sites grown in common-garden. (A) Geographical distribution of the species (green shaded zone) and location of the Equatorial (red dot) and the Subtropical (blue dot) sampling sites of propagules used in the experiment conducted in a green house (black square); (B) propagules germinating in trays filled with mangrove soil; (C-D) seedlings from the Equatorial and from the Subtropical sampling sites on the 80th day of the experiment; (E-F) seedlings from the Equatorial and Subtropical sampling sites on the 225th day of the experiment; (G) seven-months-old

1117   *Avicennia schaueriana* from the Equatorial sampling site presenting flower buds and flowers;

1118   (H-I) micrographs of stem transverse sections from pith to the outermost growth section of

1119   one representative seedling from the Equatorial and Subtropical sampling sites

1120   (magnification = 10x); stem sections were stained with Astra Blue and Safranin O; V: xylem

1121   vessels; Ph: phloem; CCC: vascular cambium.

1122

1123   **S4 Fig. Sampling of *Avicennia schaueriana* plant material used for RNA-Sequencing.**

1124   (A) Geographical distribution of the species (green zone) and locations of the Equatorial (red

1125   dot) and Subtropical (blue dot) sampling sites; (B) sampling in the native environment (top)

1126   and plant organs used for RNA extraction and sequencing (bottom).

1127

S5 Fig. Characterisation of the *de novo* assembly of the *Avicennia schaueriana* reference

transcriptome. (A) Histogram of the length of putative protein-coding and putative non-

coding transcripts; (B) classification of all predicted open reading frames (ORF) in the

reference transcriptome: complete, partial 3' (missing stop codon), partial 5' (missing start

codon), internal (missing both start and stop codons); (C) annotation of transcripts via blastx

using The *Arabidopsis* Information Resource (TAIR) protein database and The National

Center for Biotechnology Information (NCBI) RefSeq Plant protein database; (D) blastx top-

hit species distribution of transcripts annotations using NCBI RefSeq Plant protein database

as reference.

1137

1138   **S6 Fig. Detection of differentially expressed transcripts (DET) and Gene Ontology (GO)**

1139   **enrichment analysis.** (A-C) Two-dimensional scatterplot of distances among transcript

1140   expression profiles in leaf, stem and flower samples (left to right) of *Avicennia schaueriana*

1141   trees under field conditions; points represent individual samples, and distances are

approximately the log2-transformed fold changes between samples; red points represent samples from the Equatorial site and blue points represent samples from the Subtropical site. (D-F) MA-plots of transcript expression in leaves, stems and flowers; red points represent DET overexpressed in Equatorial samples; blue points represent DET overexpressed in Subtropical samples; black points represent non-differentially expressed transcripts; grey horizontal lines represent ± 1 log fold-change threshold. (G-H) Proportion of DET used in the GO enrichment analysis represented in blue and red bars. Annotation of putative coding genes was obtained using the blastx algorithm with *Arabidopsis thaliana* proteins as a reference; grey bars represent putative non-coding transcripts that are differentially expressed, and black bars represent putative coding transcripts that are differentially expressed but not annotated. (I-L) Examples of Gene Ontology (GO) terms enriched among overexpressed transcripts (DET) in Subtropical leaf samples; in Equatorial leaf samples; in Subtropical stem samples and in Equatorial stem samples. GO terms are ordered from the top down based on the P- value of enrichment (low to high, all < 0.05). Coloured bars represent the percentage of all DET that belong to each of the listed categories. Black bars represent the percentage of all transcripts in the reference transcriptome belonging to each category. Plant material of adult trees was sampled under field conditions, during the end of winter in the Subtropical site and in the beginning of the dry season in the Equatorial site. The complete lists of enriched GO terms are presented in S5-S8 Tables.

S7 Fig. Validation of RNA-Sequencing-based differential expression of transcripts by reverse transcription real time PCR (qRT-PCR). Each bar represents the relative expression of one sequenced sample. Error bars represent the standard deviation of the mean from three technical replicates of the same individual sample. Red bars represent samples from the Equatorial sampling site and blue bars, samples from the Subtropical sampling site.

1167   Non-parametric unpaired Mann-Whitney Wilcoxon U-tests were used to compare the

1168   distributions. The differential expression detected *in silico* was confirmed *in vitro* for all

1169   putative genes, at a significance threshold of 0.05. Putative genes selected for the validation

1170   of the RNA-Seq data and primers sequences are described in S10 Table.

1171

S8 Fig. Population structure of *A. schaueriana,* inferred from genome-wide genotyping

of single nucleotide polymorphic (SNP) loci. (A) Density of individuals on the discriminant

function, calculated using a multivariate method, the discriminant analysis of principal

components (DAPC); different colours represent different inferred populations; (B) map

showing the distribution of the species (green shaded area), the geographical location of

sampling sites (points) and barplots in which each colour denotes an ancestral cluster and

each line represents an individual as inferred by the method implemented in Admixture 1.3.0.

1179

S9 Fig. Environmental data in the glasshouse and at the source sites of propagules used

in the common-garden experiment. Green line: glasshouse; red line: Equatorial site

(Tracuateua automatic station, source: Brazilian National Institute of Meteorology

(INMET)); blue line: Subtropical site (Santa Marta automatic station, source: INMET).

Multiple comparisons of periodic data of each environmental variable measured during the

experiment were performed using ANOVA and post-hoc Tukey HSD tests with a

significance threshold of 0.05. * represents significant difference and hyphen (-) represents

an absence of a significant difference. (A) Average temperature; (B) average maximum

temperature; (C) average minimum temperature; (D) average air humidity; (E) average air

vapour pressure deficit (VPD).

1190

S1 Table. Morphological traits analysed in *Avicennia schaueriana* **saplings from Equatorial and Subtropical sites, grown under the same environmental conditions in a glasshouse.**

S2 Table. Transcriptome quality parameters assessed in this work.

S3 Table. Transcriptome size characterisation.

S4 Table. Putative orthologous expressed sequences between the *Avicennia schaueriana* **transcriptome sequenced in this work and publicly available transcriptomes from other congeneric**.

S5 Table. Enriched Gene Ontology (GO) terms among differentially expressed transcripts (DET) that presented higher expression in Subtropical stem samples than in Equatorial stem samples.

S6 Table. Enriched Gene Ontology (GO) terms among differentially expressed transcripts (DET) that presented higher expression in Equatorial stem samples than in Subtropical stem samples.

S7 Table. Enriched Gene Ontology (GO) terms among differentially expressed transcripts (DET) that presented higher expression in Subtropical leaf samples than in Equatorial leaf samples.

S8 Table. Enriched Gene Ontology (GO) terms among differentially expressed transcripts (DET) that presented higher expression in Equatorial leaf samples than in Subtropical leaf samples.

S9 Table. Differentially expressed transcripts of *Avicennia schaueriana* involved in the response to contrasting climate conditions between Equatorial and Subtropical sites.

S10 Table. Oligonucleotides used in qRT-PCR reactions in this work.

S11 Table. Sampling sites for plant material used in nextRAD sequencing.

S12 Table. Single Nucleotide Polymorphic loci (SNP) with signature of selection present in putative protein-coding regions of the genome of *Avicennia schaueriana*.

S1 Method. Propagules and RNA sampling sites characterization.

S2 Method. RNA-sequencing data verification through qRT-PCR.

S1 Result. Validation of RNA-Seq data through qRT-PCR.

S1 Data. Morphological traits analysed in *Avicennia schaueriana* saplings from Equatorial and Subtropical sites, grown under the same environmental conditions in a glasshouse.

S2 Data. Stem light reflectance data.

1240

1241   **S3 Data.** Variant Call Format (VCF) file required for all genome-wide SNP diversity

1242   analyses.

1243

1244   **S4 Data.** Text file complementary to S3 Data file required for all genome-wide SNP diversity

1245   analyses.