SymSim: simulating multi-faceted variability in single cell RNA sequencing

Xiuwei Zhang[1,2,*], Chenling Xu[1,*], Nir Yosef [1,2,3, ‖]

[1] Department of Electrical Engineering and Computer Sciences, Center for Computational Biology, UC Berkeley, Berkeley, CA, USA

[2] Ragon Institute of Massachusetts General Hospital, MIT and Harvard, Cambridge, MA, USA

[3] Chan-Zuckerberg Biohub Investigator

[*] These authors contributed equally

[‖] Corresponding author. Email: niryosef@berkeley.edu

The abundance of new computational methods for processing and interpreting transcriptomes at a single cell level raises the need for *in-silico* platforms for evaluation and validation. Simulated datasets which resemble the properties of real datasets can aid in method development and prioritization as well as in questions in experimental design by providing an objective ground truth. Here, we present SymSim, a simulator software that explicitly models the processes that give rise to data observed in single cell RNA-Seq experiments. The components of the SymSim pipeline pertain to the three primary sources of variation in single cell RNA-Seq data: noise intrinsic to the process of transcription, extrinsic variation that is indicative of different cell states (both discrete and continuous), and technical variation due to low sensitivity and measurement noise and bias. Unlike other simulators, the parameters that govern the simulation process directly represent meaningful properties such as mRNA capture rate, the number of PCR cycles, sequencing depth, or the use of unique molecular identifiers. We demonstrate how SymSim can be used for benchmarking methods for clustering, differential expression and trajectory inference, and for examining the effects of various parameters on their performance. We also show how SymSim can be used to evaluate the number of cells required to detect a rare population and how this number deviates from the theoretical lower bound as the quality of the data decreases. SymSim is publicly available as an R package and allows users to simulate datasets with desired properties or matched with experimental data.

The advent of single cell RNA sequencing has led to a surge of computational and statistical methods for a range of analysis tasks. Some of the methods or the tasks that they perform have originated from bulk sequencing analysis, while others address opportunities (e.g., identification of new cell states [1,2]) or technical limitations (e.g., limited sensitivity [3,4]) that are idiosyncratic to single cell genomics [5,6]. While these computational methods are often based on reasonable assumptions it is difficult to compare them to each other and assess their performance without gold standards. One approach to address this is through simulations [7–12].

Existing simulation strategies (summarized by Zappia *et al* [13]) rely primarily on fitting distributional models to observed data and then drawing from these distributions. While the resulting models provide a good fit to observed data, their parameters are often abstract and do not directly correspond to the actual processes that gave rise to the observations. This leaves an important unaddressed problem in designing and using a simulator: the need to modulate and then study the effects of specific aspects of the underlying physical processes, such as the efficiency of mRNA capture, the extent of amplification bias (e.g., by changing the number of PCR cycles, or by using unique molecular identifiers [UMI]), and the extent of transcriptional bursting. To address this, we present SymSim (**Sy**nthetic model of **m**ultiple variability factors for **Sim**ulation), a software for simulation of single cell RNA-Seq data. SymSim explicitly models three of the main sources of variation that govern single cell expression patterns [2]: allele intrinsic variation, extrinsic variation, and technical factors (Figures 1 and S1). SymSim provides the users with "knobs" to control various parameters at these three levels. First, we generate "true" numbers of molecules using a kinetic model which allows us to adjust allele intrinsic

variation and the extent of burst effect; second, we provide an intuitive interface to simulate a subpopulation structure, either discrete or along a continuum, through specification of cluster-trees, which define a low dimensional manifold from which the transcriptional kinetics is determined for every gene and every cell; third, we simulate the main stages of the library preparation process and let users control the amount of variation stemming from these steps, such as capture efficiency, amplification bias, varying sequencing depth and batch effect. Importantly, through this modeling scheme, SymSim recapitulates properties of the data (e.g., high abundance of zeros or increased noise in non-UMI protocols) without the need to explicitly force them as factors in a distributional model.

We demonstrate the utility of SymSim in two types of applications. In the first example, we use it to evaluate the performance of algorithms. We focus on the tasks of clustering and differential expression and test a number of methods under different simulation settings of biological separability and technical noise. In the second example, we use SymSim for the purpose of experimental design, focusing on the question of how many cells should one sequence to identify a certain subpopulation. The SymSim R package and an accompanying vignette are available at https://github.com/YosefLab/SymSim.

Results

The first knob: allele intrinsic variation

The first knob for controlling the simulation allows us to adjust the extent to which the infrequency of bursts of transcription adds variability to an otherwise homogenous population of cells. We use the widely accepted two-state kinetic model, in which the promoter switches

between an *on* and an *off* states with certain probabilities [14,15]. We use the notation k_{on} to represent the rate at which a gene becomes active, k_{off} the rate of the gene becoming inactive, *s* the transcription rate, and *d* the mRNA degradation rate. For simplicity, and following previous work, we fix *d* to constant value of 1 [14,16] and consider the other three parameters relative to *d* (thus becoming independent of time). Since RNA sequencing provides a single snapshot of the transcriptional process, we resort to assuming that the cells are at a steady state, and thus that the resulting single-cell measurements are drawn from the stationary distribution of the two-state kinetic model. Since *d* is fixed, we are able to express the stationary distribution for each gene analytically using a Beta-Poisson mixture [17] (Methods).

The values of the kinetic parameters (k_{on}, k_{off} and *s*) for each gene in each cell are first calculated using a product of cell-specific and gene-specific factors, then adjusted by the parameter distributions estimated from experimental data (Figure 2a, Methods). Specifically, each cell is assigned with three low-dimensional vectors (in this section, we used dimension 10; different values can be set by the user), one for each kinetic parameter. Similarly, each gene is associated with three low-dimensional vectors of the same dimension, which we term *gene effect vectors*. The value of each parameter is determined by the dot product of the two respective vectors (Figure 2a).

The coordinates of a cell's vectors represent factors of cell to cell variability that are extrinsic to the noise generated intrinsically by the process of transcription (which we model by drawing from the stationary distribution above). These values, which we term *extrinsic variability factors* (EVF) represent a low dimension manifold on which the cells lie and can be interpreted as concentrations of key proteins, morphological properties, microenvironment and more. When

simulating a homogeneous population, the EVFs of the cells are drawn from a normal distribution with a fixed mean of 1 and a standard deviation σ. σ is the *within-population variability* parameter and can be set by the user (for the results in this section σ is set to 0.5) .

The coordinates of the gene effect vectors can be interpreted as the dependence of its kinetics on the levels of EVFs. For instance, a positive value means that higher concentration of the corresponding EVF can give rise to a higher *on* rate of a certain promoter (if the EVF and gene effect vectors are both for parameter k_{on}). The gene effect values are first drawn independently from a standard normal distribution. We then replace each gene effect with a value of zero with probability η, thus ensuring that every gene is only affected by a small subset of EVFs. The *sparseness parameter* η can be set by the user; in this paper we set it to a fixed value of 0.7.

The values of kinetic parameters calculated as dot product of the EVF and gene effect vectors may not fall into realistic ranges. To map these values to realistic ranges, we first estimate the distribution of kinetic parameters of genes from real data by fitting a beta poisson model (Methods). To gain a robust estimation of the distribution of kinetic parameters to be used by SymSim, we performed the estimation multiple times with 1) different subpopulations of a dataset; 2) different imputation methods (scVI [4] and MAGIC [18]) to reduce technical variation in real data. Then we obtain aggregated distributions from the results of all the settings we considered (Figure 2b). As our reference, we used a UMI based dataset of 3,005 cortex cells by Zeisel *et al* [19] and a non-UMI based dataset of 130 IL17-expressing T helper cells (Th17) by Gaublomme *et al* [20] (See Methods for further details on the experimental data). To mitigate the effects of low sensitivity, the UMI based data was imputed using scVI [4] and MAGIC [18], and the

non-UMI data was imputed using MAGIC [18] (as scVI is only applicable to large datasets). To reduce the effects of extrinsic variation, we performed the parameter estimation separately in each of the three largest clusters in the cortex dataset (each cluster is assumed to represent a relatively homogenous subpopulation), and on the entire T cell data (a single condition, which did not contain obvious clusters) and obtained similar distribution ranges (Figures S2b). These ranges are also in line with observations from other experiments, using smFISH [21-28] or transcription inhibition based [28] methods to measure kinetic parameters (Methods and Supplementary Material Section 1). With the aggregated distributions (Figures 2b and S3a) in hand, SymSim applies a quantile approach to map the simulated parameter values that resulted from the dot product of the EVF and gene effect vectors to the range of the estimated parameters (Figure 2a, Methods).

Notably, the goal of performing kinetic parameters estimation from real data is not to estimate the true kinetic parameter values for every gene in the reference datasets (which may not be identifiable), but rather to identify the range of plausible parameter values to scale the dot products. Recent work by Larsson et al [29] reported distributions of kinetic parameters with ranges similar to our aggregated distributions (Figure S3a-b). Through simulations we show that with our procedure, we can gain comparable accuracy in estimating the distributions of parameters using imputed counts as using true counts, and both imputed and true counts yield similar ranges of the parameters to the true parameters (Figure S3c, Methods).

Finally, we account for the possibility of outlier genes with unusually high expression level, commonly observed in real data. These outlier genes are hard to model with distributional methods, and require additional parametrization [13]. This phenomena is more pronounced in

datasets from certain protocols (for example, 10x Chromium [30]) than others (for example, Smart-seq2 [31]), possibly due to selection bias which can be exacerbated by low capture rate. In SymSim, we model the high-expression outlier genes by designating a small subset of genes (whose proportion is determined by the parameter *prop_hge*) as constitutively transcribed, and adjusting their transcription rate *s* by a factor determined by the parameter *mean_hge* (>1; see Methods).

An intriguing question in the analysis of single cell RNA-seq data is the extent to which the conclusion drawn from the data (e.g., stratification into subpopulations) may be confounded by transcriptional bursting and transcriptional noise. SymSim provides a way to explore this. We first note that modality [15,17] and extent of the intrinsic noise [15] in the expression of a gene in a homogenous population of cells (*i.e.*, cells with similar EVFs) can vary for the different ranges of k_{on}, k_{off} and *s*. Specifically, one can distinguish the following three types of gene-expression distributions by the number of inflection points in the smoothed density function: unimodal with highest frequency at 0 (no inflection point), unimodal with highest frequency at non-zero value (one inflection point), and bimodal (two inflection points). Figure 2c shows the number of inflection points for different configurations of k_{on} and k_{off} with given *s*=10. This gives a clear correspondence between kinetic parameter configurations and types of gene-expression distributions. For example, when *s* is relatively large, we obtain bimodal distributions when k_{on} and k_{off} are smaller than 1.

These results thus guide us in tuning kinetic parameters to obtain desired gene-expression distributions to simulate. Specifically, we focus on adjustment of the bimodality of the distribution, which can lead to large, yet transient fluctuations in mRNA concentration at the

same cell over time, thus potentially misleading methods for cell state annotation and differential expression. To increase the overall extent of bimodality in the data, we divide (decrease) all k_{on} and k_{off} values by 10^{Bimod} (Figure 2c, yellow arrow). The parameter *Bimod* can take value from 0 to 1. This way, other properties such as burst frequency ($k_{on}/(k_{on}+k_{off})$) and synthesis rate (*s*) remain the same. In Figure 2d we show the effect of varying the *Bimod* parameter on gene-expression distribution in a simulated homogenous population. Expectedly, as *Bimod* increases, so does the number of bimodal genes, as well as the average Fano factor (Figure S2a).

The second knob: extrinsic variation via extrinsic variability factors (EVFs)

While the first knob focuses on variation within a homogeneous set of cells, the second knob allows the user to simulate multiple, different cell states. This added complexity is achieved by setting different EVF values for different cells, in a way that allows users to control cellular heterogeneity and generate discrete sub-populations or continuous trajectories. To this end, SymSim represents the desired structure of cell states using a tree (which can be specified by the user), where every subpopulation (in the discrete mode) or every cell (in the continuous mode) is assigned with a position along the tree. Different positions in the tree correspond to different expected EVF values, and the expected absolute difference between the value of an EVF of any two cells is linearly proportional to the square root of their distance in the tree (Supplementary Material Section 2).

When SymSim is applied in a discrete mode, the cells are sampled from the leaves of the tree. The set of cells that are assigned to the same leaf in the tree form a subpopulation, and their EVF values are drawn from the same distribution. As above, we draw these EVF from a

normal distribution, where the mean is determined by the position in the tree and the standard deviation is defined by the parameter σ. When SymSim is applied in a continuous mode, the cells are positioned along the edges of the tree with a small step size (which is determined by branch lengths and number of cells; Methods). The EVF values are then drawn from a normal distribution where the mean is determined by the position in the tree, and the standard deviation is defined by σ (Figure 3a).

To facilitate the correspondence between EVF values and distances in the tree we use a Brownian motion procedure as described in [32] (Methods; Figure 3a). Specifically, for each EVF we set the mean value at the root of the tree to a fixed number (default set root node to 1) and then perform Brownian motion along the branch. Figure 3a illustrates this process using populations 2 and 3 in the tree as an example. Notably, in the continuous mode, this formulation can give rise to a rich set of patterns of changes in gene expression from root ('progenitor cells') to leaves ('target cells'), including the commonly observed impulse profile [33,34] (Figure S3d-e). As an alternative, we also implemented a mode for simulating continuous data by which gene expression from root to leaves is determined explicitly by an impulse function. This might be preferable if the user would like to generate smoother changes in gene expression, or specific temporal patterns. In the following analyzes we use the Brownian motion model.

Notably, SymSim only generates a subset of EVFs from the tree, while the remaining ones are drawn from the same distribution for all subpopulations (Figure 3a). The tree-sampled subset, which we term *Diff-EVFs* (Differential EVFs) represents the conditions or factors which are different between sub-populations, and they usually account for a small proportion of all the

EVFs. The number of Diff-EVFs can be set by the user. The results in this section were produced with 60 EVFs, 20% of them are Diff-EVFs.

With this formulation, users can control the extent of between-population variation by setting the branch lengths of the input tree, and combine it with a desired level of within-population variation by setting the parameter σ. Notably, both σ and the square root of branch lengths in the tree are in units of EVF values. It is therefore the case that for any two positions in the tree, the ratio of square root tree distance to σ determines the separability between the respective distributions of the values assigned to any given Diff-EVF (Supplementary Material Section 2). As illustration, Figure 3 depicts the tSNE plots of cells from the same input tree with different σ in either a discrete (Figure 3b) or continuous (Figure 3c) mode. Notably, both panels show that the tSNE plots reflect the structure of the input tree well.

The third knob: technical variation

A large part of the variation observed in scRNA-seq datasets stems from technical sources [35–37]. The technical confounders reflect noise, reduced sensitivity and bias that are introduced during sample processing steps such as mRNA capture, reverse transcription, PCR amplification, RNA fragmentation, and sequencing. In order to introduce realistic technical variation into our model, we explicitly simulate the major steps in the experimental procedures. We implemented two library preparation protocols: (1) full length mRNAs profiling without the use of UMIs (e.g., with a standard SmartSeq2 [31]); and (2) Profiling only the end of the mRNA molecule with addition of UMIs (e.g., 10x Chromium [30]). The former protocol is usually applied for a small number of cells and with a large number of reads per cell, providing full information on transcript structure [38].

The latter is normally applied for many cells with shallower sequencing, and it is affected less by amplification and gene length biases [35].

The workflow of these steps are shown in Figure 4a (Methods). Starting from the simulated true mRNA content of a given cell (namely, number of transcripts per gene, sampled from the stationary distribution of the promoter kinetic model), the first step is mRNA capture, where every molecule is retained with probability $\hat{\alpha}$. The value of the capture efficiency $\hat{\alpha}$ associated with each cell is drawn from a normal distribution with a mean α and standard deviation β, which can be set by the user. The second step is amplification, where in every cycle SymSim selects each available molecule with a certain probability and duplicates it. The expected amplification efficiency and the number of PCR cycles can be set by the user. As an optional step, SymSim provides the option of linear amplification (e.g., as in CEL-Seq [39]). We do not apply this option in this manuscript. In the third step each amplified molecule is broken down into fragments, in preparation for further amplification, size selection and sequencing. The lengths of the simulated transcripts are obtained from the human reference genome, and the fragmentation is calibrated so that the average fragment length is 400bp, which is typical for RNA sequencing (Supplementary Material Section 3). Resulting fragments that are within an acceptable size range (100 to 1000 bp) are then carried on to the fourth and last step of sequencing.

The number of reads per cell (namely, the number of sequenced fragments) is drawn from a normal distribution whose mean is determined by the parameter *Depth,* which, along with the respective standard deviation (*Depth_sd*) can be provided by the user. To derive the final "observed" expression values we do not account for sequencing errors, and assume that every

sequenced fragment is assigned to the correct gene it originates from. For the non-UMI option, we define the raw measurement of expression as the number of reads per gene. If UMIs are used, SymSim counts every original mRNA molecule only once by collapsing all reads that originated from the same molecule. Notably, for certain depth values, the resulting distribution of number of reads per UMI is similar to the one observed in a dataset of murine cortex cells [19] (Figure S4a-b).

It has been previously shown that estimation of gene-expression levels from full length mRNA sequencing protocols has amplification biases related to sequence-specific properties like gene length and GC-content [35,40], whereas the use of UMIs can correct these biases [40,41]. In particular, we have observed a negative correlation between gene length and length-normalized gene-expression in our reference non-UMI dataset (murine Th17 cells from Gaublomme et al [20]; Figure 4b), and the same trend is reported by Phipson *et al* [40]. To account for that, we parametrize the efficiency of the PCR amplification step using a linear model that represents gene length bias (Methods). As a result, our simulated data with a non-UMI protocol shows a similar dependence of gene-expression on gene length as in experimental data (Figure 4b, real data is from [20]). In cases where UMIs are used, gene length effects are also modeled during amplification, but these effects are mitigated since each molecule is counted at most once. We therefore do not observe gene length bias in the UMI-based simulated data, similarly to the experimental data (Figure S4c, real data is from [19]). Finally, we model batch effects with multiplicative factors that are gene- and batch-specific. In Figure S4d, we show the same population of cells are separated by batches. To simplify the discussion at the remainder of this paper, we assume that the data comes from a single batch.

In Figure 4c, we show the comparison between the simulated true mRNA content of one cell and the simulated observed counts obtained with or without UMI. We consider two scenarios: the first scenario represents a study with a low technical confounding and the second one represents a highly confounded dataset. Parameters which differ between these "good" and "bad" cases in this example include capture efficiency (α), extent of amplification bias (*MaxAmpBias*) and sequencing depth (*Depth*). Using "bad" technical parameters introduce more noise to true counts, and compared to the non-UMI simulation the UMIs reduce technical noise. The histograms of true counts and four versions of simulated counts are shown in Figure S4e. Using quantile-quantile plots (Q-Q plots; Figure S4f) further demonstrates that UMIs help in maintaining a better representation of the true counts in the observed data.

The total computation time to simulate a dataset consists of time to generate true counts and time to generate observed counts from true counts. We show the runtime and memory usage for different parameter configurations in Methods.

Fitting parameters to real data

For a given real data set, SymSim can produce observed (read or UMI) counts that have similar statistical properties to the real data (Figures 4d-e), by searching in a database of simulations obtained from a range of parameter configurations (Methods). This procedure focuses on within-population variability (Similarly to Splatter [13]) and sets the values of ten parameters from both the first and third knobs (Methods). We test this function with the non-UMI Th17 dataset [20] (using all cells), the UMI cortex dataset [19] (using a subpopulation of 948 CA1 pyramidal neuron cells) and two UMI datasets from 10x Genomics, denoted by "UMI 10x t4k" and "UMI 10x

pbmc8k" (sources of experimental data are described in Methods). See Supplementary Material Section 4 for the values of the fitted parameters.

Side by side inspection of the histograms of true mRNA levels (simulated) and observed counts (simulated and experimental), indicates that SymSim can transform the simulated ground truth (Figure 4d, left) into simulated observations (Figure 4d, middle) that match the real observed data for both UMI and non-UMI protocols (Figure 4d, right). For a more quantitative analysis, we generated Q-Q plots of the distributions of mean (after adding one to all values), percent-non-zero and standard deviation (SD) of genes between simulated and experimental data (Figure 4e, Figure S5b for the UMI 10x pbmc8k dataset). Notably, we observe a certain level of inaccuracy in matching the SD at the lower ends for the non-UMI data, which can be due to lowly expressed genes. Indeed, when we exclude lowly expressed genes from the real data, the matching of SD improve substantially (Q-Q plots shown in Figure S5a). Furthemore, we conducted a similar analysis by training Splatter [13] and powSimR [12] with the same experimental datasets as input, and found that SymSim matches this data significantly better (Figure S5c). Q-Q plots of the distributions of coefficient of variation (CV) and mean (without adding one) of genes between experimental data and data simulated respectively by SymSim, Splatter and powSimR also show that SymSim provides an overall better fit than the other two simulators (Figure S5d-e).

As additional comparisons of the simulators, we plot the mean *vs* variance relationship (Figure S6a) and mean *vs* percent-nonzero relationship (Figure S6b) of genes for the experimental datasets and simulated datasets with each simulator. For both types of plots, the datasets simulated by SymSim are the most similar to the corresponding experimental datasets.

Finally, we inspected the distributions of number of expressed genes per cell (Figure S7a), distributions of number of UMIs per cell (Figure S7b) and number of UMIs per cell sorted in decreasing order (Figure S7c). SymSim and Splatter are comparable in terms of these measures and fit experimental data better than powSimR. The difference between the distributions of powsimR and those of real data in Figures S7a-b can be due to that powsimR does not attempt to reproduce unfiltered count matrices, but instead outputs data after filtering of genes and cells. Notably, we limit our parameters to the values that are available in our pre-computed grid, which explains the possible differences between observed quantities (e.g., sequencing depth per cell) in SymSim and in the real-data.

Using SymSim to evaluate computational methods for single cell RNA-seq data

SymSim can be used to benchmark methods for single cell RNA-Seq data analysis as it provides both observed counts and a reference ground truth. In this section we demonstrate the utility of SymSim as tool for benchmarking methods for clustering, differential expression and trajectory inference in a sample consisting of multiple subpopulations, using the structure depicted in Figure 3a. The design of SymSim allows us to evaluate the effect of various biological and technical confounders on the accuracy of downstream analysis. Here, we investigate the effect of total number of cells (N), within population variability (σ), mRNA capture rate (α) and sequencing depth (*Depth*). We also test the effect of the proportion of cells associated with the smallest sub-population of cells (*Prop*), using population 2 in the tree as our designated "rare" sub-population.

Using SymSim to evaluate clustering methods

We begin by inspecting the impact of each parameter on the performance of clustering methods. To this end, we simulated observed counts using the UMI option, and traversed a grid

of values for the five parameters with 18 simulation runs per configuration. The values of the remaining parameters are largely determined according to the cortex dataset [19] and specified in Supplementary Material Section 5.2. We tested three clustering methods: k-means based on euclidean distance of the first 10 principle components, k-means based on Euclidean distance in a nonlinear latent space learned by scVI [4] and SIMLR [42]. In all cases we set the expected number of clusters to the ground truth value (k=5). The accuracy of the methods is evaluated using the adjusted Rand index (ARI; higher values indicate better performance). To inspect the effects of the various parameters on clustering performance, we performed multiple linear regression between the parameters and the ARI. The regression coefficients are shown in Figure 5a. Overall, σ appears to be the most dominant factor, and the proportion of the rare population (*Prop*) is clearly positively associated with better performance. Among the technical parameters, while α plays a role on the performance especially for the rare population, the impact of *Depth* is minor.

We then focus on varying the dominant factors (except N, which we discuss in the next section), namely, the within-population variation parameter σ and mRNA capture efficiency α, for the benchmarking analysis henceforth. In particular, the range of values for σ is set such that we cover various data characteristics -- from well separated populations, to almost entirely mixed ones (Supplementary Material Section 5). We compare the performance of four clustering strategies: SIMLR [42], dimensionality reduction with scVI [4] followed by k-means, and dimensionality reduction with PCA followed by Louvain clustering [43] (implemented in Seurat [44]) or k-means. We observe better accuracy as the quality of the data increases or the within-population variation decreases (Figure 5b-c). Interestingly, comparing $\sigma=0.6$, $\sigma=0.8$ and $\sigma=1$, we can tell that when σ is high enough to make the clustering challenging, further

increasing σ does not yield obvious changes (Figure 5b). We observe a similar trend of saturation, inspecting increasing levels of capture efficiency (α), especially with scVI. Comparing the methods to each other, we see that scVI/k-means has the highest ARI in most cases; when the rare population accounts for 5% of all cells, Seurat is the second best and PCA/k-means and SIMLR are comparable; when the rare population accounts for 10% of all cells, SIMLR performs slightly worse than Seurat and PCA/k-means.

To investigate the effect of bimodality on the performance of clustering algorithms, we increased the parameter *bimod* from 0 to 1 for the dataset used in the above analysis and performed clustering. The comparison of clustering results between different methods and different values of *bimod* is shown in Figure S8a. We see that in most cases there is a decrease of performance for the same method with increase of *bimod*. We then aggregate all the values of adjusted Rand index for all methods and all the parameters of α and σ, but only group them by the *bimod* value, and performed Wilcoxon test between the two groups of values. Figure S8b shows that the difference between these two groups is small but significant (p-value $< e^{-12}$).

Using SymSim to evaluate differential expression methods

Our mechanism for simulating multiple populations automatically generates differentially expressed (DE) genes between populations (in the discrete setting; Figure 3b) or along pseudotime (in the continuous setting; Figure 3c). In the following, we use SymSim to benchmark methods for detecting DE genes, focusing on the discrete setting. We use two criteria to define the ground truth set of DE genes. The first criterion is that the number of Diff-EVFs that are associated with a non-zero gene effect value (which we denote as *nDiff-EVFgene;* Figure 6a) should be larger than zero. This criterion is motivated by our model

of transcription regulation: the kinetic parameters of a gene are affected by extrinsic factors, and changes to extrinsic factors might therefore lead to changes in the number of transcripts. Indeed, when we compare the true simulated gene expression values between subpopulations (i.e., before introducing technical confounders with the third knob), we get a uniform (random) distribution of p-values for genes with no Diff-EVFs, and an increasing skew as *nDiff-EVFgene* increases (Figure 6b, using Wilcoxon test); Figure S9a shows that the log fold change of gene-expression between subpopulations increases with *nDiff-EVFgene*. Nevertheless, in some cases, the actual expression values of genes with *nDiff-EVFgene>0* might not differ since the effects of different Diff-EVFs or the effects of modifying different kinetic parameters may cancel out. Differential expression might also be blurred by a high within-population variability. We therefore added an additional constraint and require that all DE genes must have a sufficiently large fold change in their simulated true simulated expression levels (threshold of absolute log2 fold change ranges from 0.6 to 1, details in Figure legends; Methods).

An important distinguishing feature of SymSim is that it provides an intuitive way for generating case studies for DE analysis that consist of multiple subpopulations with a predefined structure of similarity. To illustrate this, consider populations 1, 2, and 4 (Figure 6c), which form a hierarchy (2 and 4 are closer to each other and similarly distant from 1). This user-defined structure is reflected in the sizes of the sets of DE genes, obtained respectively from populations 1 vs 2 (1229 genes), 1 vs 4 (1412 genes) and 2 vs 4 (815 genes). Consistent with the hierarchy, the first two gene sets are overlapping and larger than the third one.

As an example for a benchmark study, we used four methods to detect DE genes: edgeR [45], DESeq2 [46], Wilcoxon rank-sum test and t-test on observed counts generated by various

parameter settings (Methods, Supplementary Material Section 5.3). We tested the effect of the total number of cells (N) and mRNA capture rate (α) with 10 simulation runs per parameter configuration. We use two accuracy measures: a) AUROC (area under receiver operating characteristic curve), obtained by treating the p-values output from each method as a predictor (Figure 6d, Methods); b) negative of Spearman correlation between the p-values of each detection method and the log fold difference of the true expression levels (Figure 6e, Methods).

From Figures 6d-e, one can observe that when the numbers of cells are small (30 in each population), edgeR has the best performance while the other three methods are comparable to each other. When the numbers of cells of both populations increase to 300, the two naive methods Wilcoxon test and t-test improve in their relative performance, compared to edgeR and DESeq2. The case where the number of cells are 30 and 300 appear to have performance between those of the 30 *vs* 30 case and the 300 *vs* 300 case. When increasing capture efficiency, all methods gain performance except for the case of AUROC with 300 cells. In that case, the drop in AUROC for some methods is caused by inflation in p-values as α increases, which results in lower specificity (Figure S9b). Notably, we noticed that the adjusted p-values from DESeq2 can have many missing entries (NAs), especially when *α* is low (and thus counts are low), and therefore we used its unadjusted p-values in Figure 6d-e. However, this assignment of NAs in practice filters out genes which do not pass a certain threshold of absolute magnitude (explained in DESeq2 vignette [47]). To make use of this filtering, we conducted an additional analysis where we used the adjusted p-values for DESeq2 and compare it to all other methods using only the non-filtered (non NA) genes (Figure S9c). As expected, the performance of all methods (and specifically DESeq2) improves when considering only this set of genes, and converges to high values already at lower capture efficiency rates.

To summarize, we find that edgeR has the best overall performance, with the t-test rank second followed by Wilcoxon test. We note that the aim of this section is to demonstrate the use of SymSim for methods benchmarking instead of performing a comprehensive comparisons of methods. Nevertheless, our ranking is consistent with results from a recent paper which evaluated 36 methods for DE analysis with single cell RNA-Seq data [48].

We then perform all DE methods on the datasets with *bimod*=1 and compare the results with the original dataset with *bimod*=0 (Figure S10a-c). From Figure S10a-c we can see a drop in performance when increasing *bimod*, especially when the number of cells is small (Figure S10a-b). The drop is less prominent when the number of cells in the two populations are respectively 300 and 300 (Figure S10c). Notably, the drop in the performance of clustering and DE performance cannot be simply attributed to a global decrease in gene expression levels, since increasing *bimod* does not change this statistic (Figure S10d and Methods).

Using SymSim to evaluate trajectory inference methods

The ability of SymSim to generate a continuum of cell states makes it a convenient choice to benchmark trajectory inference methods. We compare three methods including Monocle [49,50], Slingshot [51], and a minimum spanning tree (MST) algorithm implemented in the package dynverse [52] (Methods). We generate datasets with different values of σ and α with the input tree shown in Figure 3a. For each parameter configuration, we repeat the simulation 10 times. To evaluate the trajectory inference methods, we use two measures: 1) Spearman correlation between true cell order and inferred cell order. We consider cells on each lineage (a path from root to a leaf) separately and take the average of correlation on all five lineages. 2) *k*-nearest

neighbor purity (knn purity) of cells, that is, for each cell, we calculate the Jaccard Index between its *k*-nearest neighbors in the true trajectory and that in the inferred trajectory. Results are shown in Figure 7. In these plots, *k* is set to 100. In Figure 7a, we vary σ and fix α as 0.1. Both the correlation and knn purity decrease when σ increases. In Figure 7b, we vary α and fix σ as 0.6. All methods show an overall increasing trend along with α with both measures. Consistent with a recent benchmark study [52], we observe that overall Slingshot clearly outperforms the other two methods.

Experimental design

Deciding how many cells to sequence is a decision many researchers face when designing an experiment, and the optimal number of cells to sequence highly depends on the nature of the biological system under investigation and the respective technical hurdles. A previous approach to this problem [53] assumes that the goal of the experiment is to identify subpopulations of cells and provides a theoretical lower bound for the problem. This bound considers the aspect of counting cells (namely, sequencing enough representative cells from each subpopulation), but it does not account for the identifiability of each subpopulation, which may be hampered by both technical and biological factors as well as the performance of clustering algorithms.

In the following we demonstrate how SymSim can be used to shed more light on this important problem. Importantly, in its current form SymSim does not use real data to model between-population variability. We therefore interpret the results in a relative manner --- how do different variability factors shift the required number of cells, compared to each other and to the theoretical lower bound. Our example focuses on a case of one rare subset, represented by cells from population 2 (using the same tree in Figure 6c; note that one can easily generalize

this procedure to multiple rare subpopulations). We simulate observed counts with numbers of cells (N) ranging from 600 to 7000. These simulations were based on the parameters fit to the cortex dataset [19] with varying levels of σ and α (250 simulations per parameter configuration).

We applied the same four clustering methods as described in the previous section (*k*-means with scVI or PCA, Louvain clustering with PCA (Seurat) and SIMLR). We say that a given algorithm was successful in "detecting the rare population" if at least 50 cells from this set are assigned to the same cluster, and form at least 70% of the cells in that cluster. We use these labels to compute an empirical success probability P for each algorithm and each parameter configuration. Out of the 250 simulations for each parameter configuration, we randomly sample 100 randomizations 20 times, and for each 100 randomizations we can calculate a value of P. We then plot the mean and standard deviation as error bars of the 20 values of P for each N under each configuration (Figure 8a-d, Figure S11). To get an upper bound on performance that better reflects the data (rather than the choice of algorithm), we take the best P out of all algorithms, and apply cubic spline smoothing (gray curves, Figure 8a-d). In each plot we also include the theoretical limit which only requires the presence of at least 50 cells from the rare subpopulation (Methods). The theoretical curve (which is independent of all parameters except N) reaches almost 1 at N=1400. Conversely, the empirical curves vary dramatically, based on parameter values. For an easy case of low within-population variability (σ=0.4) and high capture efficiency (α=0.1) the empirical upper bound curve is close to the theoretical one (Figure 8a). This curve decreases when increasing the effect of either nuisance factor (Figure 8b,c). The reduction is substantially more dramatic for most of the methods when both nuisance factors increase, while Seurat remains robust to this change, potentially due to

that the graph-based clustering method is advantageous in reducing false positives for the rare population compared to k-means (Figure 8d).

To understand the implications on the number of cells required in a given setting, we calculated how many cells are required, in each configuration, to achieve a success rate of respectively 0.6, 0.7, 0.8 and 0.9 (Figure 8e). As expected, the resulting numbers can be much higher than the theoretical lower bound. For example, to achieve a success rate of 0.9, when the within-population variability increases ($\sigma=0.7$), we need at least 3838 cells (corresponding to $\alpha=0.01$), while with the theoretical curve, we need only less than 1200 cells. In general, the number of cells needed increase when the desired success rate and σ increase. Very low capture efficiency ($\alpha=0.001$, $\alpha=0.005$) tend to require high number of cells. Considering only the binomial sampling of cells may therefore underestimate the number of cells needed for a realistic scenario, and considerations of biological and technical variations with simulators like SymSim is merited.

Discussion

SymSim has the following features which are advantageous over existing simulators: (i) We simulate true transcript counts from a kinetic model that can be interpreted in terms of transcript synthesis rate, promoter activation and deactivation. (ii) When generating multiple discrete or continuous populations, instead of generating biological differences through directly altering the true transcript count distribution, we set Diff-EVFs, which can be interpreted as biological conditions that cause the differences between subpopulations of cells. This is a more natural and realistic way to simulate biological transcriptional differences. (iii) The EVF formulation provides an intuitive way to specify and simulate complex structures of cell-cell similarity,

without the need for manual specifications of the numbers of DE genes [13]. (iv) When generating observed counts, we simulate key steps in real experimental protocols, which automatically gives us dropout events, length bias, and distribution of library sizes. We also provide choices to use UMI based protocols or non-UMI full length mRNA protocols, as the properties of data output from these two categories can be very different.

The main input parameters to SymSim are self-explanatory with their own biological or technical meanings, which users can adjust to match an experimental dataset of interest. While the procedure of parameter fitting was developed in order to generate simulated datasets with similar properties, it may also provide additional insight, as the parameters are biologically or technically interpretable. For instance, comparing the parameters fit to the the UMI and non-UMI datasets in this study we note that the capture efficiency inferred to the latter is much higher (Supplementary Material Section 4). The modular nature of SymSim provides possibilities to generalize its application. For example, the generation of true counts with EVFs and transcription kinetics can be replaced by learning a generative model from real data, with methods such as scVI [4]. This type of extension will facilitate simulation of between-subpopulation diversity that better mimics experimental observations, albeit at the cost of using parameters that are less interpretable biologically. Another extended application of interest is to use different tree structures for different Diff-EVFs when generating multiple populations of cells, such that every tree represents a different aspect of variability between cells. For instance, using this approach, one tree can represent a differentiation process and the other can represent variability due to the physical location of the cell.

As the number and extent of biological applications of single cell genomics continues to grow, so does the extent of analytical questions one can tackle, which go beyond standard `bulk era' analysis steps (e.g., trajectory analysis, mRNA velocity [54], and more). The need for robust analytical methods therefore increases, and so does the means for proper evaluation of these methods. SymSim provides a starting point to address this challenge of flexible and feature-rich simulation for method evaluation, as it aims to directly mimic the key mechanistic properties of single cell RNA sequencing.

Methods

Simulating gene expression with the kinetic model

As shown in Figure 2a, the kinetic model of gene expression considers that a gene can be either *on* or *off* and the probabilities to transit between the two states are k_{on} and k_{off}. When the gene is *on* it is transcribed with transcription rate s. The transcripts degrade with rate d. For a given gene, based on these parameters one can simulate the number of its transcript molecules over time. The theoretical probability distribution can be calculated via the Master Equation [15,17], which is the steady state solution for the kinetic model. Alternatively, the kinetic model can be represented by a Beta-Poisson model [17], which we use in our implementation to sample expression values for a gene.

Calculating parameters for the kinetic model in SymSim simulation

For a gene in a cell, the parameters for the kinetic model k_{on}, k_{off}, and s are calculated from the cell-specific EVF vectors of this cell and the gene effect vectors of the gene (Figure 2a). To

allow independent control of the three parameters, we use one EVF vector and one gene effect vector for each parameter. Take k_{on} as an example: denoting the EVF vector as $(e_1^{kon}, e_2^{kon}, \ldots, e_p^{kon})$, and the gene effect vector for k_{on} as $(g_1^{kon}, g_2^{kon}, \ldots, g_p^{kon})$, the cell-gene specific value for k_{on} is the dot product of these two vectors. We then map these k_{on} values to the distribution of k_{on} estimated from experimental data, to obtain the matched parameters. We do so by sorting the k_{on} values (from dot products) for all genes in all cells, sampling the same number of values from the experimental k_{on} distribution (the number of values would be $m*n$, where m is the number of genes and n is the number of cells), and updating the k_{on} values to the ones sampled from the experimental distribution with the same rank. The values of k_{off} and s are calculated in the same way.

Finally, to better model the high expression outlier genes which usually account for a very small proportion of all genes, we introduce two parameters (*prop_hge* and *mean_hge*) to SymSim for these outlier genes. First, we use parameter *prop_hge* to represent the proportion of these genes. For randomly selected genes according to this proportion, we consider that they are always "on" and their gene-expression levels are sampled from a Poisson distribution with parameter s (the transcription rate). To reflect their high gene-expression level, we increase their transcription rate s calculated from EVFs and gene effects with an inflation factor. The inflation factor for each gene is the exponential (base 2) of (*mean_hge*-1+*rank*/*n*), where *rank* is the rank of mean s of a gene among all outlier genes (smaller s corresponds to smaller rank values), n is the total number of outlier genes and *mean_hge* specifies the extent of inflation to original s.

Estimating kinetic parameters from real data

We estimated kinetic parameters from experimental data using an MCMC approach. For each gene, its expression X depends on p, the proportion of time it is *on*, and the mRNA synthesis rate s. The parameter p itself is a random variable determined by the kinetic parameters k_{on} and k_{off}. We model p as a Beta distributed variable with shape parameters k_{on} and k_{off}. We model X as a Poisson distributed variable with parameter $p*s$. The distribution of X is then identical to the distribution calculated using the Master Equation [17]. The downsampling effect is modeled as a Binomial sampling with X being the number of trials, and f being the probability that a transcript is sampled for sequencing.

We fit this model to the experimental data using the Gibbs sampler implemented in RJAGS. The number of iterations is set to 2000. At every iteration, we sample each parameter from its marginal posterior conditional on the value of all other parameters. To meet the assumption that all cells share the same kinetic parameters we divide cells by clustering that is performed in the original study and fit the model to counts in a single cluster of cells at a time. We also use imputed read counts, rather than the raw read counts. We use scVI [4] and MAGIC [18] for the imputation. We used multiple starting points to diagnose convergence. For each cell cluster and each imputation method, we fit the model independently three times. We observe that different runs from the same data have very similar distributions (Figure S2c, UMI cortex data, subpopulation pyramidal CA1, imputed with scVI). We also investigate the autocorrelation and plotted the typical lag *vs* correlation plots for these chains (Figure S2d). The correlation drops as the lag increases, which is a sign of sufficient mixing of samples. We then merge all the acceptable chains to obtain the distributions for the combination of cluster and imputation

method. Finally, results from all combinations are merged to obtain the final distribution of kinetic parameters (Figure 2b).

We then perform simulations to test how well this procedure of kinetic parameter estimation reconstruct the true parameters. We simulate observed UMI counts for five discrete populations whose relationships are defined by the tree in Figure 3a, with the complete workflow of SymSim. Then we perform imputation on the observed UMI counts with both scVI and MAGIC. We then apply the kinetic parameter estimation procedure described above on each subpopulation imputed by each imputation method separately. Finally we aggregate the parameters estimated from all populations and both imputation methods, and gain the distributions estimated from imputed counts. We also apply the kinetic parameter estimation procedure on the true counts which are available.

Ranges of kinetic parameters from literature

We look into literature for the ranges of kinetic parameters k_{on}, k_{off}, and s which are experimentally measured [21–28]. The range of burst size, or s, from these studies ranges from 2-4000. And the k_{on} and k_{off} values ranges from 0.0001 to 1 per minute, and the half-life of mRNA varies from 1-10 hours, which correspond to 0.001 to 0.01 per minute. This means that k_{on}/d and k_{off}/d could take values from 10^{-2} to 10^3. The specific parameter values reported by these studies are in Supplementary Material Section 1.

Simulation of discrete and continuous populations

The structure of populations can be represented by a tree and the user can input the tree in Newick format in a text file. The differences between populations are realized through Diff-EVFs, which usually account for a small proportion of all EVFs. There are two different

modes of simulation the Diff-EVFs, Continuous and Discrete. Both modes can be modeled by Brownian motion along the tree from root to leaves, where one starts with a given value at the root (default is 1), and at each time point t, y(t) is calculated as $y(t) = y(t-1) + N(0, \Delta t)$, where $N()$ represents a Gaussian function, and Δt is the step size. The values at internal nodes of the tree are shared by all branches connecting this node. In the continuous mode, the step size between two consecutive cells on a given branch is obtained by randomly sampling n_b positions on a branch b of length l_b. In the discrete mode, the step size is the corresponding branch length and the number steps is the depth of the tips. For a given tip and a given EVF, the value we sample at the tip is used as the mean of a Gaussian distribution to sample the values for that EVF for all cells in that population with standard deviation σ (Figure 3a).

For the continuous mode we also provide an alternative option to the Brownian motion model: using impulse functions for modeling the path-specific variation. When impulse function is used, for cells sampled from branches that are not on the root-tip path for the specific EVF, they are sampled from a univariate normal with mean equal to the EVF value at their most recent common ancestor with the varying path, and standard deviation σ.

Simulation of technical steps from mRNA capturing to sequencing

We simulate two categories of library preparation protocols, one does not use UMIs (unique molecular identifiers)[30] and sequences full length mRNAs (using procedures in Smart-seq2 [31] as template), and the other uses UMIs and sequences only the 3' end of the mRNA (using the Chromium chemistry by 10x Genomics as template). In the pre-amplification step, we provide option of using linear amplification to mimic the CEL-seq protocol. As shown in Figure 4a, we take one transcript with 16 molecules as an example. To implement the UMIs, each original molecule has a variable to its count at each step. The technical steps include the following:

1) Capturing step: molecules are captured from the cell with probability α.

2) Pre-amplification step: if using non-CEL-Seq protocols, this step involves N rounds of PCR amplifications. We introduce sequence-specific biases during amplification, which includes transcript length bias and other bias assigned randomly. Parameter *lenslope* can be used to control the amount of length bias, and *MaxAmpBias* is used to tune the total amount of amplification bias. If using CEL-Seq protocol this step is the *in vitro* transcription (IVT) linear amplification.

3) Fragmentation step: the mRNAs are chopped into fragments for sequencing. If sequencing full-length mRNA, all fragments with acceptable length are kept for sequencing. If sequencing only the 3' end for UMI protocols, only fragments on the 3' end are kept for sequencing. For each transcript length, we calculate a distribution of number of fragments given expected fragment length, and use this distribution to generate the number of fragments during our simulation of the fragmentation step. The distributions are different for non-UMI and UMI protocols; the details of calculating the distributions are in Supplementary Material Section 2.

4) Amplification step: fragments go through another k rounds of PCR amplifications for all protocols, including CEL-Seq and non-CEL-Seq protocols.

5) Sequencing step: amplified fragments from the previous step are randomly selected according to a given value of sequencing depth, which is the total number of reads (fragments) to sequence.

6) After the sequencing step (assuming all reads are correctly sequenced and mapped to their original gene), we can get the UMI counts for UMI protocols and read counts for non-UMI protocols.

Note that for simplicity, this pipeline omits several steps, including reverse transcription, and library cleaning up.

The runtime to simulate a dataset is dominated by the simulation of these technical steps. The total time cost varies with mRNA capture efficiency and protocol. With one CPU core of an Intel(R) Xeon(R) CPU E5-2690 v4 @ 2.60GHz, the runtime and memory consumed to simulate a dataset with 500 cells and 10000 genes for different parameters is as following:

Capture efficiency	Protocol	Sequencing depth	Runtime (min)	Memory
0.05	UMI	100000	21.8	1.8G
0.2	UMI	100000	76.9	2.3G
0.05	nonUMI	100000	8.6	1.8G
0.2	nonUMI	100000	16.5	1.7G

Simulation of amplification biases

During PCR amplification of the full length cDNAs, the PCR amplification rate (namely, probability to be amplified) can vary for different transcripts. As a result, some transcripts are over- and some are under-amplified. This causes the unwanted amplification bias. To simulate this, for each gene, its PCR amplification rate is set to a sum of a basal amplification rate (the input parameter *rate_2PCR*, which equals to the average amplification rate across all genes) plus a bias term *B*. The bias term *B* ranges from -*MaxAmpBias* to *MaxAmpBias*, where *MaxAmpBias* is a user-specified parameter to represent the total amount of amplification bias in our system.

B is composed of two categories of biases: biases related to transcript length (referred to as gene length bias, denoted by B_{length}) and biases caused by other factors (denoted by B_{rand}). We use a linear function to model the gene length bias: we first bin all gene lengths into *nbins* bins, and get the average length in each bin: $L = (l_{bin(1)}, l_{bin(2)}, ..., l_{bin(nbins)})$. The length bias term associated with a gene in bin *i* is set to: $B_{length}(i) = lenslope * median(L) - lenslope * L(i)$

The parameter *lenslope* controls the extent of gene length bias. To ensure that B_{length} does not exceed *MaxAmpBias*, the parameter *lenslope* should be smaller than

$2 \times MaxAmpBias/(nbins - 1)$. We then set the second term B_{rand} to a random value in the

residual range [$- MaxRandBias$, $MaxRandBias$] where

$MaxRandBias = MaxAmpBias - max(B_{length})$. Namely, $B_{rand} = N(0, MaxRandBias)$, where N() is a

Gaussian.

Therefore, for a given gene with length l, its PCR amplification rate is:

$$rate\_2PCR + B_{length}(bin(l)) + B_{rand}$$

This rate is used in all rounds of PCRs in the pre-amplification step. The biases then get amplified as more PCR cycles are performed, where transcripts with higher amplification rate will likely get more molecules. Assigning a UMI to each molecule before amplification allows us to collapse all molecules with the same UMI after amplification, so different amplification rates will not affect the final molecule counts. For Figure 4b, $lenslope$ is set to 0.023, $MaxAmpBias$ is set to 0.3, $nbins$ is set to 20, and *rate_2PCR* is set to 0.7.

Fitting simulation parameters to real data

To find the best matching parameters to a real dataset, we simulate a database of datasets with a grid of parameters over a wide range. For each simulated dataset, we calculate the following statistics: mean, percent-non-zero, standard deviation of genes over all cells. Then given a real dataset, we find the simulated dataset which have the most similar distributions of the statistics to the real data, and return the corresponding parameter configurations.

The parameters and their ranges for simulating the two databases are as following:

Parameters	Values for the non-UMI database	Values for the UMI database

Sigma (σ): within-population heterogeneity	0.1, 0.2, 0.6	0.2, 0.6
Gene_effects_sd: standard deviation for generating gene effect vectors	1, 2	2
scale_s: cell size parameter, use small values for cell types known to be small	0.3, 0.5, 0.6, 1	0.1, 0.2, 0.3, 0.4, 0.5, 0.7, 1
prop_hge: proportion of extreme high expression outlier genes	0.015, 0.02, 0.025, 0.03	0.01, 0.015, 0.02, 0.025, 0.03
mean_hge: scale factor for high expression outlier genes	3, 4, 5, 6	3, 4, 5, 6
Alpha_mean (α): mean mRNA capture efficiency	0.05, 0.1, 0.15, 0.2, 0.25	0.007, 0.01, 0.02, 0.03, 0.04, 0.05, 0.07, 0.1
Alpha_sd (β): standard deviation of mRNA capture efficiency across cells	0.005, 0.01, 0.015, 0.02, 0.03, 0.045, 0.06, 0.075	7e-04, 0.001, 0.0014, 0.002, 0.003, 0.0035, 0.004, 0.005, 0.006, 0.007, 0.008, 0.009, 0.01, 0.012, 0.015, 0.02, 0.025, 0.03, 0.035, 0.05
depth_mean (*Depth*): mean sequencing depth for a set of cells	1e+05, 5e+05, 1e+06, 2e+06	45000, 70000, 95000, 150000, 3e+05, 5e+05
Depth_sd: standard deviation of sequencing depth for a set of cells	10000 ,30000, 50000, 1e+05, 150000, 2e+05, 3e+05, 6e+05	4500, 7e3, 9e3, 9.5e3, 13500, 14e3, 15e3, 19e3, 21e3, 22500, 28500, 3e4, 35e3, 4.5e4, 47500, 5e4, 7.5e4, 9e4, 15e4, 25e4
nPCR1: number of PCRs in pre-amplification phase	14, 18	10, 14

Applying dimensionality reduction and clustering methods

We apply three different dimensionality reduction methods to cluster cells simulated from multiple discrete populations: PCA, scVI and SIMLR. PCA is the naive baseline method that is also the most commonly seen in single-cell RNA-seq analysis. scVI is a more recent method that uses a zero-inflated negative binomial variational auto-encoder model to infer latent space

for each single cell. For both the first two methods, cluster identities are then assigned using k-means clustering. The third method, SIMLR, performs dimensionality reduction and cluster identity iteratively to maximize cluster separation. The fourth method, implemented in Seurat, uses PCA for dimensionality reduction and the Louvain clustering.

Simulation of differentially expressed (DE) genes

Diff-EVFs give rise to differences between populations as well as DE genes between populations. DE genes by design are the ones with non-zero gene effect values corresponding to the Diff-EVFs (Figure 6a), as the gene effect vectors are sparse with a majority of values being 0s. However, the actual expression values of these genes might not differ if the changes in two EVFs cancel out, or if the effect of change one of the kinetic parameter is canceled out by the change in another kinetic parameter. Thus we also use the log2 fold change (LFC) of mean gene-expression from the two populations as another criteria. The mean expression can be calculated based on simulated true counts, which is subject to gene-expression intrinsic noise, or based on the kinetic parameters themselves, directly from the theoretical gene-expression distribution. If the kinetic parameters of a gene in a cell is k_{on}, k_{off} and s, the expected gene-expression of this gene in this cell is $s*k_{on}/(k_{on}+k_{off})$. We use multiple thresholds ranging from 0.6 to 1 on the |LFC| to define a gene is DE, in order to avoid being biased with one single artificial threshold.

Detection of differentially expressed (DE) genes

DE genes in observed counts are detected respectively with edgeR, DESeq2, Wilcoxon test and Student t-test. For edgeR, we used the quasi-likelihood approach (QLF) with cellular detection rate (the fraction of genes that are detected with nonzero counts in each cell) as covariate. For

DESeq2, we use "local" for the fittype parameter, and we evaluate its performance respectively based on the p-values and adjusted p-values which serve as filtering of genes.

The output from each DE method is a p-value for each gene, with smaller values meaning the gene is more likely to be a DE gene. We use two metrics to evaluate the performance of a DE method: a) AUROC (area under receiver operating characteristic curve), where we apply different thresholds on the p-values to obtain different sets of predicted DE genes, and we can then plot ROC curves with different combinations of 1-*specificity* and *sensitivity*, thus calculate the area under the ROC curve. b) negative of Spearman correlation between the p-values of each detection method and the log fold difference of the true expression levels. Genes with high log fold change in true transcript counts should correspond to low p-value if the DE method works well. As the inferred p-values and log fold change in true counts are expected to be anti-correlated, we take the negative of this correlation, such that higher value correspond to better performance.

Applying trajectory inference methods

We use the R packages dynwrap (https://github.com/dynverse/dynwrap, version 0.1.0) and dynmethods (https://github.com/dynverse/dynmethods, version 0.1.0) to run the three trajectory inference methods compared in this manuscript: Monocle (version 2.6.4), Slingshot (version 0.99.12), and MST (a basic method implemented in dynmethods). All methods were run with default parameters. Both dynwrap and dynmethods are under the collection of R packages dynverse used in the manuscript by Saelens *et al* [52].

Effect of parameter *bimod* on gene expression levels

To investigate if increasing *bimod* will cause decrease in overall gene-expression levels of genes thus lead to the decrease in performance of both clustering and DE methods, we calculate the percentage change of total number of transcripts of genes from *bimod*=0 to *bimod*=1. That is, for each gene, we calculate $(\sum_{j=1}^{m} x_j - \sum_{j=1}^{m} x'_j)/\sum_{j=1}^{m} x_j$, where x_j is the number of transcripts of this gene in cell *j* when *bimod*=0, x'_j is the number of transcripts of this gene in cell *j* when *bimod*=1, and *m* is the number of cells. From Figure S10d we see that there is no consistent increase or decrease of total number of transcripts for the genes when changing *bimod*. Therefore, we conjecture that the drop in the performance of clustering and DE is rather caused by change in the distribution of gene-expression levels of genes instead of overall gene-expression levels.

Binomial model to calculate the probability of detecting a population

Assuming all sequenced cells are correctly assigned to its original population, the probability that at least *x* cells are detected from a population only depends on the binomial sampling. Denote the total number of cells by *N* and the proportion of the cells in the given population by *r*, the probability that at least *x* cells are detected for the population is:

$$1 - \sum_{k=0}^{x-1} \binom{N}{k} r^k (1-r)^{(N-k)}$$

This formula is used to generate the black curves in Figure 8a-d.

During our simulation to estimate the number of cells needed to detect a rare population, we simulate the random sampling process as follows: we start with a total of 10000 cells for all five populations with 2000 cells for each population. We set probability vector of a cell belonging

to each population as (0.25, 0.05, 0.25, 0.25, 0.2), where Population 2 is the rare population with smallest probability. For each randomization and given total number of cells N (N <= 7000), we randomly sample N cells from the pool of 10000 cells according to the probability vector.

Sources of experimental data

We use four experimental datasets throughout this paper, one is from [20] which does not use UMIs and the other three use UMIs. The non-UMI datasets profiles Th17 cells under various conditions and in our paper we use a subpopulation of 130 TGF-β1+IL-6 cells. We refer to this dataset as the non-UMI Th17 dataset. The first UMI dataset profiles 3006 cerebral cortex cells [19]. The authors found nine classes in these cells. In this paper, to get distributions of kinetic parameters to map our simulated parameters to the same distribution, we perform parameter estimation respectively on 1) 628 cells sampled from the oligodendrocyte class; 2) 715 cells sampled from the CA1 pyramidal neurons; 3) 296 cells sampled from the S1 pyramidal neurons. To verify that SymSim can simulate data with similar statistics with given experimental dataset, we use all 948 oligodendrocyte cells in the cortex dataset. The other two UMI datasets are downloaded from the 10x Genomics website: one has around 4538 Pan T Cells (denoted as the "UMI 10x t4k" dataset, https://support.10xgenomics.com/single-cell-gene-expression/datasets/2.0.1/t_4k) and the other has 8381 PBMC cells (denoted as "UMI 10x pbmc8k", data available at https://support.10xgenomics.com/single-cell-gene-expression/datasets/2.1.0/pbmc8k). For both 10x datasets, we use cluster 1 (the largest cluster) identified at their respective analysis page.

Acknowledgements

XZ was supported by grant #220558 from the Ragon Institute of MGH, MIT and Harvard. CX and NY were supported by NIH/NHLBI grant U19 AI-090023-09.

Author contributions

XZ, CX and NY conceived the study and wrote the paper. XZ and CX wrote the software and conducted the analysis.

Figure legends

Figure 1 Overview of SymSim. The *true transcript counts*, which are the number of molecules for each transcript in each cell at the time of analysis, are generated through the classical promoter kinetic model with parameters: promoter *on* rate (k_{on}), *off* rate (k_{off}) and RNA synthesis rate (*s*). The values of the kinetic parameters are determined by the product of gene-specific coefficients (termed *gene effects*) and cell-specific coefficients. The latter set of coefficients is termed extrinsic variability factors (EVF), and it is indicative of the cell state. The expected value of each EVFs is determined in accordance to the position of the cell in a user-defined tree structure. The tree dictates the structure of the resulting cell-cell similarity map (which can be either discrete or continuous) since the distance between any two cells in the tree is proportional to the expected distance between their EVF values. For homogenous populations (represented by a single location in the tree), the EVFs are drawn *iid* from a distribution whose mean is the expected EVF value and variance is provided by the user. From the true transcript counts we explicitly simulate the key experimental steps of library preparation and sequencing, and obtain *observed counts*, which are read counts for full length mRNA sequencing protocols, and UMI counts, otherwise.

Figure 2 Intrinsic variation. (**a**) A diagram of how gene and cell specific kinetic parameters are simulated from cell-specific EVF and gene-specific gene effect vectors, and how the kinetic parameters are used in a model of transcription. Each cell has a separate EVF vector for K_{on}, K_{off} and *S*. Each parameter is generated through two steps: first, for each gene in each cell, we take the dot product of the corresponding EVF and gene effect vectors. Second, the dot product values are mapped to distributions of parameters estimated from experimental data. The

"matched parameters" are used to generate true transcript counts (see Methods). (**b**) The distributions of k_{on}, k_{off} and s that are used in SymSim for simulations. These distributions are aggregated from inferred results of three subpopulations of the UMI cortex dataset (oligodendrocytes, pyramidal CA1 and pyramidal S1) after imputation by scVI and MAGIC. (**c**) A heatmap showing the effect of parameter k_{on} and k_{off} on the number of modes in transcript counts. The value of s is fixed to 10 in this plot. The red area with low k_{on} and k_{off} have one zero mode and one non-zero mode. The gray area with low k_{on} and high k_{off} has only one zero mode, and the blue area with high k_{on} and low k_{off} have one non-zero mode. The yellow arrow shows how the parameter *Bimod* can modify the amount of bimodality in the transcript count distribution. (**d**) Histogram heatmaps of transcript count distribution of the true simulated counts with varying values of *Bimod*, showing that increasing *Bimod* increases the zero-components of transcript counts and the number of bimodal genes. In these heatmaps, each row corresponds to a gene, each column corresponds to a level of expression, and the color intensity is proportional to the number of cells that express the respective gene at the respective expression level.

Figure 3 Extrinsic variation. (**a**) Illustration of generating a diverse set of cell states with SymSim. The tree represents the relationship between cells. The numbers on the edges are branch lengths; the node numbers indicate the ID of the respective subpopulation (each subpopulation is represented by a single position [leaf] in the tree). The matrix to the right depicts the derivation of EVF values. Each row corresponds to an EVF (only two are Diff-EVF), each column corresponds to a position in the tree, and the content specifies the distribution from which the EVF values are drawn. We use the notation $y_a(b)$ to represent the expected value of EVF *b* in position *a* in the tree. The rightmost plot depicts the derivation of these expected

values with Brownian motion. We use subpopulations 2 and 3 as examples for both discrete cases (sampling only cells within the subpopulations) or continuous (sampling cells along the trajectories from the root `progenitor' state [node 6] to the two `target' subpopulations [nodes 2 and 3]). (**b**) tSNE plots of five discrete populations generated from the tree structure shown in (a). Different values of σ give rise to different heterogeneity of each population. (**c**) tSNE plots of continuous populations generated from the same tree. The colors corresponds to the colors on branches in the tree shown in (a). When increasing σ, cells are more scattered around the main paths which follow the tree structure.

Figure 4 Technical variation. (**a**) A diagram showing the workflow of adding technical variation to true simulated counts. Each gray or orange square represents a molecule of the same transcript in one cell. We implement the following steps: mRNA capturing, pre-amplification (PCR or linear amplification of the cDNAs), fragmentation, amplification after fragmentation, sequencing, and calculation of UMI counts or read counts. Details of these steps can be found in Methods. (**b**) Gene length bias in both simulated and experimental data for the non-UMI protocol. (**c**) Scatter plots comparing true counts and observed counts obtained through: 1. non-UMI, "good" parameters ($α$=0.2, *MaxAmpBias*=0.1, *Depth*=1e6); 2. UMI, "good" parameters ($α$=0.2, *MaxAmpBias*=0.1, *Depth*=5e5); 3. non-UMI, "bad" parameters ($α$=0.05, *MaxAmpBias*=0.2, *Depth*=1e6); 4. UMI, "bad" parameters ($α$=0.04, *MaxAmpBias*=0.2, *Depth*=5e5). (**d**) 2D transcript counts histogram heatmap of UMI and non-UMI simulated true counts and simulated observed counts, generated with parameters which best match the input experimental counts, and histogram heatmaps of the respective experimental counts (non-UMI Th17, UMI cortex and UMI 10x t4k datasets). (**e**) Q-Q plots comparing the mean, percent non-zero and standard deviation in experimental counts and SymSim simulated observed

counts respectively for the non-UMI, UMI cortex and UMI 10x t4k datasets. A good match is indicated by most of the dots falling close to the red line.

Figure 5. Benchmarking of clustering methods. (**a**) Coefficients of various parameters from multiple linear regression between parameters and the adjusted Rand index (ARI). In the left plot the ARI are averaged over all populations, and in the right plot the ARI is only for the rare population (population 2, with 5% of all cells). (**b**) ARI of the rare populations using the three clustering methods when changing σ (α=0.04). Left plot: the rare population accounts for 5% of all the cells; right plot: the rare population accounts for 10% of all the cells. (**c**) ARI of the rare populations using the three clustering methods when changing α (σ=0.8). Left plot: the rare population accounts for 5% of all the cells; right plot: the rare population accounts for 10% of all the cells.

Figure 6. Benchmarking of DE detection methods. (**a**) Illustration of how DE genes are generated through the Diff-EVFs. Red squares in the gene effect matrix correspond to non-zero values. The two genes indicated by the arrows are DE genes by number of Diff-EVFs they have (respectively 2 and 1). (**b**) Q-Q plot comparing the p-value obtained from differential expression analysis between subpopulations 2 and 4 (using Wilcoxon test on the true simulated counts) to a uniform distribution. Genes are grouped by the number of Diff-EVFs they use and different groups are plotted in different colors. The numbers of Diff-EVFs used by genes can be thought of as the degree of DE-ness. Genes with more Diff-EVFs have p-values further diverged from uniform distribution. (**c**) Venn diagram showing that closely related populations have less DE genes between them compared to distantly related populations. We use populations 1, 2, 4 as examples: there are much more DE genes from comparison of "1 vs 2" and "1 vs 4" than " 2 vs

4", and DE genes from "1 vs 2" and "1 vs 4" have a big overlap. The DE genes are determined by log2 fold change (LFC) of true counts with criterion |LFC| > 0.8. (**d**) The AUROC (area under receiver operating characteristic curve) of detecting DE genes using four different methods from observed counts with changing capture efficiency α (σ=0.6). The populations under comparison are 2 and 4. Three sets of criteria were used to define the true DE genes and the final performance was the average performance from the three sets: (1) *nDiff-EVFgene*>0 and |LFC|>0.6; (2) *nDiff-EVFgene*>0 and |LFC|>0.8; (3) *nDiff-EVFgene*>0 and |LFC|>1. LFC was calculated with theoretical means from the kinetic parameters. (**e**) The negative of correlation between log2 fold change on theoretical mean of gene-expression and p-values obtained by a DE detection method, which changing capture efficiency α (σ=0.6). The populations under comparison are 2 and 4.

Figure 7. Benchmark trajectory inference methods. (**a**) Pseudotime correlation and knn purity of all methods when varying σ (α=0.1). (**b**) Pseudotime correlation and knn purity of all methods when varying α (σ=0.6).

Figure 8. The number of cells needed to detect a rare population. We generate five populations according to the tree structure shown in Figure 3 and set population 2 as the rare population which accounts for 5% of the cells. Other populations share 95% of the cells evenly. The criteria of "detecting the rare population" is that at least 50 cells from this population are correctly detected and the precision (positive predicted value) is at least 70%. (**a-d**) The probability of detecting the rare population with when sequencing N (x axis) cells under different σ and α configurations, with different clustering methods. The black curve represents the theoretical probability from the binomial model, assuming that all cells sequenced are assigned

correctly to the original population. The gray curve with transparency takes the maximum value at each data point from all three clustering methods with smoothing. (**e**) The heatmap shows the number of cells needed to sequence under different configurations of σ and α to detect the rare population with success rates 0.6, 0.7, 0.8, 0.9, always using the best clustering method,.

Figure S1 A flow chart of the process of SymSim generating the observed counts, with parameters involved in each step described.

Figure S2 (**a**) The Fano factor of genes across cells with different values for *Bimod*. (**b**) Distribution of kinetic parameters estimated from different experimental data. (**c**) Distribution of kinetic parameters estimated for the same data (UMI cortex dataset, population pyramidal CA1, imputed with scVI) using different starting values to obtain multiple MCMC chains. (**d**) Auto-correlation diagnose plots for the three chains in (c). The correlation values are average over all genes. For each lag value k, correlation is calculated between x_t and x_{t+k}, where x_t is a sample from MCMC at step t.

Figure S3 (**a**) distributions of kinetic parameters shown in Figure 2b in forms of burst frequency (k_{on}) and burst size (s/k_{off}). (**b**) distributions of burst frequency (k_{on}) and burst size (s/k_{off}) reported by Larsson *et al* [29] in mouse fibroblasts. (**c**) Comparison of distributions of true kinetic parameters (left column), parameters estimated from true counts (middle column), and parameters estimated from imputed counts (right column). True counts and observed counts of five populations are obtained through running SymSim with the tree shown in Figure 3a. (**d**) The gene expression of two DE genes along the pseudotime in continuous populations. The structure of populations is represented by the tree in Figure 3a. We plot gene-expression of the

lineage from the root to population 2. The number of EVFs is 20 for each of the three kinetic parameters, and 12 EVFs of parameter *s* are Diff-EVFs. σ is set to 0.4. (**e**) Heatmap of expression of genes with at least 4 Diff-EVFs along the lineage from the root to population 2.

Figure S4 (**a**) The distribution of number of reads per UMI sequenced in the cortex dataset. This plot comes from the supplementary material of the original paper [19]. (**b**) The distribution of number of reads per UMI sequenced in our simulated data, when using the UMI protocol, with 100k reads per cell. (**c**) The gene length bias in observed counts with UMI protocol, respectively from experimental and simulated data. The experimental data is the cortex dataset from paper [19]. (**d**) TSNE plot of cells simulated for one homogeneous population in two batches. (**e**) The histogram heatmap of gene expression of true simulated counts, and observed simulated counts under "good" and "bad" parameter settings. The parameters are the same as described in Figure 4c. In these heatmaps, each row corresponds to a gene, each column corresponds to a level of expression, and the color intensity is proportional to the number of cells that express the respective gene at the respective expression level. (**f**) Q-Q plots of gene expression of true simulated counts and observed simulated counts under "good" and "bad" parameter settings. The data is the same as that plotted in Figure 4c.

Figure S5 (**a**) Q-Q plots of standard deviation for genes between experimental data and SymSim simulated data for the non-UMI protocols after removing a proportion of lowly expressed genes. (**b**) Q-Q plots of mean, percent-nonzero, and standard deviation between a second 10x dataset (pbmc8k) and SymSim simulated dataset. (**c**) Q-Q plots comparing the mean, percent non-zero and standard deviation in experimental counts and simulated observed counts respectively for the non-UMI, UMI cortex and UMI 10x t4k datasets. The simulation is

performed by Splatter (left three columns) and powSimR (right three columns). (**d**) Q-Q plots comparing the coefficient of variation (CV) in experimental counts and simulated observed counts respectively for the non-UMI, UMI cortex and UMI 10x t4k datasets. Simulation is performed by SymSim (left column), Splatter (middle column) and powSimR (right column). (**e**) Q-Q plots comparing the log10(mean) in experimental counts and simulated observed counts respectively for the non-UMI, UMI cortex and UMI 10x t4k datasets. Simulation is performed by SymSim (left column), Splatter (middle column) and powSimR (right column).

Figure S6 (**a**) The mean-variance relationship of the three experimental datasets (non-UMI Th17 data, top row; UMI cortex data, middle row; UMI 10x t4k data, bottom row) and the corresponding simulated datasets by SymSim (2nd column), Splatter (3rd column) and powSimR (4th column). (**b**) The mean *vs* percent-non-zero plots of the three experimental datasets (non-UMI Th17 data, top row; UMI cortex data, middle row, UMI 10x t4k data, bottom row) and the corresponding simulated datasets by SymSim (2nd column), Splatter (3rd column) and powSimR (4th column). Red dots are the 0%, 0.2%, 0.4%, …, 100% percentiles of the mean and percent-non-zero values of all genes.

Figure S7 (**a**) Distribution of number of expressed genes per cell of the three experimental datasets (non-UMI Th17 data, left; UMI cortex data, middle; UMI 10x t4k data, right) and the corresponding simulated data by SymSim, Splatter and powSimR. (**b**) Distribution of number of UMIs per cell for the two UMI datasets and the corresponding simulated data by SymSim, Splatter and powSimR. (**c**) Number of UMIs per cell of the two UMI datasets and the corresponding simulated datasets, sorted in decreasing order.

Figure S8 Effects of *bimod* parameter on the performance of clustering methods. (**a**) Top row: ARI of the rare populations using the three clustering methods (PCA, scVI and SIMLR) with different values for *bimod* (*bimod*=0 and *bimod*=1) when changing σ (α=0.04). Left plot: the rare population accounts for 5% of all the cells; right plot: the rare population accounts for 10% of all the cells. Bottom row: ARI of the rare populations using the three clustering methods with different values for *bimod* when changing α (σ=0.8). Left plot: the rare population accounts for 5% of all the cells; right plot: the rare population accounts for 10% of all the cells. (**b**) Boxplots of aggregated ARI values, separated only by different values of *bimod*. Boxplots are plotted using the boxplot() function in R with default parameters, where the bottom and top of the box are the lower and upper quartiles respectively, and the band near the middle of the box is the median.

Figure S9 (**a**) The log fold change of genes with different number of Diff-EVFs. (**b**) The distribution of p-values of four different DE methods, with different values of capture efficiency α (noted as the title of each plot), with population size 300 vs 300. (**c**) The AUROC (top row) and negative of correlation (bottom row) measures of the four different methods for DE gene detection. In these plots the adjusted p-values from DESeq2 are used instead of the p-values, and the genes with NAs in the adjusted p-values are removed for all methods to calculate the accuracy measures. The numbers of cells in both populations for the plots on the left column are both 30, on the middle column the number of cells are respectively 30 and 300, and on the right column the numbers of cells are both 300.

Figure S10 Effects of *bimod* parameter on the performance of DE methods. In each panel, we show the AUROC (area under receiver operating characteristic curve, top row) and Negative of Correlation (bottom row) of detecting DE genes using four different methods from observed

counts with changing capture efficiency α (σ=0.6). Left column corresponds to *bimod*=0 and right column corresponds to *bimod*=1. (**a**) Number of cells in the two populations (populations 2 and 4) are 30 and 30. (**b**) Number of cells in the two populations are 30 and 300. (**c**) Number of cells in the two populations are 300 and 300. (**d**) Percentage of change of total transcript for each gene from *bimod*=0 to *bimod*=1.

Figure S11 The probability of detecting the rare population (population 2 in the tree shown in Figure 3) under a wide range of configurations of σ (Sigma) and α (Alpha). The criteria of detecting the population is that at least 50 cells are detected (true positive >=50) with precision at least 70%.

Reference

1. Amir, E.-A. D. *et al.* viSNE enables visualization of high dimensional single-cell data and reveals phenotypic heterogeneity of leukemia. *Nat. Biotechnol.* **31**, 545–552 (2013).

2. Wagner, A., Regev, A. & Yosef, N. Revealing the vectors of cellular identity with single-cell genomics. *Nat. Biotechnol.* **34**, 1145–1160 (2016).

3. Pierson, E. & Yau, C. ZIFA: Dimensionality reduction for zero-inflated single-cell gene expression analysis. *Genome Biol.* **16**, 241 (2015).

4. Lopez, R., Regier, J., Cole, M. B., Jordan, M. I. & Yosef, N. Deep generative modeling for single-cell transcriptomics. *Nat. Methods* **15**, 1053–1058 (2018).

5. Bacher, R. & Kendziorski, C. Design and computational analysis of single-cell RNA-sequencing experiments. *Genome Biol.* **17**, 63 (2016).

6. Stegle, O., Teichmann, S. A. & Marioni, J. C. Computational and analytical challenges in single-cell transcriptomics. *Nat. Rev. Genet.* **16**, 133–145 (2015).

7. L. Lun, A. T., Bach, K. & Marioni, J. C. Pooling across cells to normalize single-cell RNA sequencing data with many zero counts. *Genome Biol.* **17**, 75 (2016).

8. Lun, A. T. L. & Marioni, J. C. Overcoming confounding plate effects in differential expression analyses of single-cell RNA-seq data. *Biostatistics* **18**, 451–464 (2017).

9. Vallejos, C. A., Marioni, J. C. & Richardson, S. BASiCS: Bayesian Analysis of Single-Cell Sequencing Data. *PLoS Comput. Biol.* **11**, e1004333 (2015).

10. Korthauer, K. D. *et al.* A statistical approach for identifying differential distributions in single-cell RNA-seq experiments. *Genome Biol.* **17**, 222 (2016).

11. Ziegenhain, C. *et al.* Comparative Analysis of Single-Cell RNA Sequencing Methods. *Mol. Cell* **65**, 631–643.e4 (2017).

12. Vieth, B., Ziegenhain, C., Parekh, S., Enard, W. & Hellmann, I. powsimR: power analysis for bulk and single cell RNA-seq experiments. *Bioinformatics* **33**, 3486–3488 (2017).

13. Zappia, L., Phipson, B. & Oshlack, A. Splatter: simulation of single-cell RNA sequencing data. *Genome Biol.* **18**, 174 (2017).

14. Peccoud, J. & Ycart, B. Markovian Modeling of Gene-Product Synthesis. *Theor. Popul. Biol.* **48**, 222–234 (1995).

15. Munsky, B., Neuert, G. & van Oudenaarden, A. Using Gene Expression Noise to Understand Gene Regulation. *Science* **336**, 183–187 (2012).

16. Larson, D. R. What do expression dynamics tell us about the mechanism of transcription? *Curr. Opin. Genet. Dev.* **21**, 591–599 (2011).

17. Kim, J. K. & Marioni, J. C. Inferring the kinetics of stochastic gene expression from single-cell RNA-sequencing data. *Genome Biol.* **14**, R7 (2013).

18. van Dijk, D. *et al.* Recovering Gene Interactions from Single-Cell Data Using Data Diffusion. *Cell* **174**, 716–729.e27 (2018).

19. Zeisel, A. *et al.* Brain structure. Cell types in the mouse cortex and hippocampus revealed by single-cell RNA-seq. *Science* **347**, 1138–1142 (2015).

20. Gaublomme, J. T. *et al.* Single-Cell Genomics Unveils Critical Regulators of Th17 Cell Pathogenicity. *Cell* **163**, 1400–1412 (2015).

21. Padovan-Merhar, O. *et al.* Single mammalian cells compensate for differences in cellular volume and DNA copy number through independent global transcriptional mechanisms. *Mol. Cell* **58**, 339–352 (2015).

22. Bahar Halpern, K. *et al.* Bursty Gene Expression in the Intact Mammalian Liver. *Mol. Cell* **58**, 147–156 (2015).

23. Skinner, S. O. *et al.* Single-cell analysis of transcription kinetics across the cell cycle. *Elife*

5, e12175 (2016).

24. Dar, R. D. *et al.* Transcriptional Bursting Explains the Noise–Versus–Mean Relationship in mRNA and Protein Levels. *PLoS One* **11**, e0158298 (2016).

25. Xu, H., Sepúlveda, L. A., Figard, L., Sokac, A. M. & Golding, I. Combining protein and mRNA quantification to decipher transcriptional regulation. *Nat. Methods* **12**, 739–742 (2015).

26. Suter, D. M. *et al.* Mammalian genes are transcribed with widely different bursting kinetics. *Science* **332**, 472–474 (2011).

27. Bartman, C. R., Hsu, S. C., Hsiung, C. C.-S., Raj, A. & Blobel, G. A. Enhancer Regulation of Transcriptional Bursting Parameters Revealed by Forced Chromatin Looping. *Mol. Cell* **62**, 237–247 (2016).

28. Rabani, M., Pieper, L., Chew, G.-L. & Schier, A. F. A Massively Parallel Reporter Assay of 3′ UTR Sequences Identifies In Vivo Rules for mRNA Degradation. *Mol. Cell* **68**, 1083–1094.e5 (2017).

29. Larsson, A. J. M. *et al.* Genomic encoding of transcriptional burst kinetics. *Nature* **565**, 251–254 (2019).

30. Zheng, G. X. Y. *et al.* Massively parallel digital transcriptional profiling of single cells. *Nat. Commun.* **8**, 14049 (2017).

31. Picelli, S. *et al.* Smart-seq2 for sensitive full-length transcriptome profiling in single cells. *Nat. Methods* **10**, 1096–1098 (2013).

32. Felsenstein, J. *Inferring phylogenies*. (Sinauer Associates, 2004).

33. Sander, J., Schultze, J. L. & Yosef, N. ImpulseDE: detection of differentially expressed genes in time series data using impulse models. *Bioinformatics* **33**, 757–759 (2017).

34. Fischer, D. S., Theis, F. J. & Yosef, N. Impulse model-based differential expression

analysis of time course sequencing data. *bioRxiv* 113548 (2017). doi:10.1101/113548

35. Vallejos, C. A., Risso, D., Scialdone, A., Dudoit, S. & Marioni, J. C. Normalizing single-cell RNA sequencing data: challenges and opportunities. *Nat. Methods* **14**, 565–571 (2017).

36. Jaitin, D. A. *et al.* Massively parallel single-cell RNA-seq for marker-free decomposition of tissues into cell types. *Science* **343**, 776–779 (2014).

37. Kar, G. *et al.* Flipping between Polycomb repressed and active transcriptional states introduces noise in gene expression. *Nat. Commun.* **8**, 36 (2017).

38. Afik, S. *et al.* Targeted reconstruction of T cell receptor sequence from single cell RNA-seq links CDR3 length to T cell differentiation state. *Nucleic Acids Res.* (2017). doi:10.1093/nar/gkx615

39. Hashimshony, T., Wagner, F., Sher, N. & Yanai, I. CEL-Seq: single-cell RNA-Seq by multiplexed linear amplification. *Cell Rep.* **2**, 666–673 (2012).

40. Phipson, B., Zappia, L. & Oshlack, A. Gene length and detection bias in single cell RNA sequencing protocols. *F1000Res.* **6**, 595 (2017).

41. Islam, S. *et al.* Quantitative single-cell RNA-seq with unique molecular identifiers. *Nat. Methods* **11**, 163–166 (2014).

42. Wang, B., Zhu, J., Pierson, E., Ramazzotti, D. & Batzoglou, S. Visualization and analysis of single-cell RNA-seq data by kernel-based similarity learning. *Nat. Methods* **14**, 414–416 (2017).

43. Blondel, V. D., Guillaume, J.-L., Lambiotte, R. & Lefebvre, E. Fast unfolding of communities in large networks. *J. Stat. Mech.* **2008**, P10008 (2008).

44. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat. Biotechnol.* **36**, 411–420 (2018).

45. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140 (2010).

46. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

47. Love, M. I., Anders, S. & Huber, W. Analyzing RNA-seq data with DESeq2. (2018). Available at: http://bioconductor.org/packages/devel/bioc/vignettes/DESeq2/inst/doc/DESeq2.html. (Accessed: 29th June 2018)

48. Soneson, C. & Robinson, M. D. Bias, robustness and scalability in single-cell differential expression analysis. *Nat. Methods* (2018). doi:10.1038/nmeth.4612

49. Qiu, X. *et al.* Reversed graph embedding resolves complex single-cell trajectories. *Nat. Methods* **14**, 979–982 (2017).

50. Qiu, X. *et al.* Single-cell mRNA quantification and differential analysis with Census. *Nat. Methods* **14**, 309–315 (2017).

51. Street, K. *et al.* Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics. *BMC Genomics* **19**, 477 (2018).

52. Saelens, W., Cannoodt, R., Todorov, H. & Saeys, Y. A comparison of single-cell trajectory inference methods: towards more accurate and robust tools. *bioRxiv* 276907 (2018). doi:10.1101/276907

53. How Many Cells. Available at: https://satijalab.org/howmanycells. (Accessed: 19th May 2018)

54. La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494–498 (2018).

Figure 1

Figure 2

Figure 3

a

	pop2		pop3		pop4		pop5		pop1	
Non Diff-EVFs	$N(1, \sigma^2)$		$N(1, \sigma^2)$							
	$N(1, \sigma^2)$		$N(1, \sigma^2)$							
	$N(1, \sigma^2)$		$N(1, \sigma^2)$							
Diff-EVFs	$N(y_2(1), \sigma^2)$		$N(y_3(1), \sigma^2)$							
	$N(y_2(2), \sigma^2)$		$N(y_3(2), \sigma^2)$							

b

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure S1

External Variability Factors (EVF)
nEVFs: number of EVFs, representing dimension of the latent space of cell identify (length of EVF vectors)
Values of EVFs are sampled from $N(\mu, \sigma^2)$ for each subpopulation, where default value of μ is 1, and σ determines the within-population variation. For multiple populations, users input a tree *phyla*, and the μ for each population is generated along the tree

Gene effects
Probability of a value being non-zero is determined by *gene_effect_prob*, the non-zero values are generated by N(*geffect_mean*, *gene_effects_sd*)

K_{on} K_{off} S
k_{on} and k_{off} can be adjusted by *bimod* to increase bimodality.
A proportion *prop_hge* of genes are set to be high expression outliers and they are considered always "on" with synthesis rate *s'*, *s'* are obtained from their original *s* with multiplier determined by *mean_hge*.

Kinetic model
scale_s can scale the size of the cell, to account for that different cell types can have different size

True Transcript Counts

Library preparation and sequencing
protocol: UMI or nonUMI;
*mRNA capture efficiency α f*or all cells are sampled from a truncated Normal N(*alpha_mean*, *alpha_sd*) with minimum value 0.0005;
sequencing depth for all cells are sampled from truncated Normal N(*depth_mean*, *depth_sd*) with minimum value 200;
Other major parameters are: *lenslope* (amount of transcript length bias), *MaxAmpBias* (amount of total amplification bias), *rate_2PCR* (PCR amplification rate), *nPCR1* (number of PCR cycles in the 1st phase), *nPCR2* (number of PCR cycles in the 2nd phase), *LinearAmp* (use linear amplification of not), *nbatch* (number of batches), *gene_len* (length for all transcripts, sampled from a pool of gene lengths of human genes), *LinearAmp_coef* (coefficient for linear amplification).

Observed Transcript Counts

Figure S2

Figure S3

Figure S4

a

b

depth=1e5

c

UMI

Experimental

Simulated

Log2(mean(UMI counts))

Log2(gene length)

d

1 population, 2 batches

e

True counts

nonUMI, good quality

nonUMI, bad quality

UMI, good quality

UMI, Bad quality

genes

log10(Count) bins

freq

f

nonUMI_good
nonUMI_bad
UMI_good
UMI_bad

Simulated Observed Expression

Simulated True Expression

Figure S5

Figure S5

d

Comparison of log10(cv) between simulated and real data

SymSim   Splatter   powsimR

e

Comparison of log10(mean) between simulated and real data

SymSim   Splatter   powsimR

Figure S6

a

b

Figure S7

Figure S8

a

b

Figure S9

Figure S10

Figure S11