TITLE PAGE

How do alterations in root cap structure and integrity impact root responses to an increase in

medium strength?

J. Roué, H. Chauvet, E. Badel, F. Bizet, B. Moulia, and V. Legué

Université Clermont Auvergne, INRA, PIAF, F-63000 Clermont–Ferrand, France

Juliette Roué - juliette.roue@uca.fr

Hugo Chauvet - hugo.chauvet-thiry@inra.fr

Eric Badel - eric.badel@inra.fr

François Bizet - fr_bizet@yahoo.fr

Bruno Moulia - bruno.moulia@inra.fr

Corresponding author: valerie.legue@uca.fr, (+33) 0 473405273

Date of submission: 07/27/2018

Number of tables: 1

Number of figures: 7

Word count: 6588

Supplementary data:

Number of tables: 0

Number of figures: 2

Videos: 6

How do alterations in root cap structure and integrity impact root responses to an increase in

medium strength?

J. Roué, H. Chauvet, E. Badel, F. Bizet, B. Moulia, and V. Legué

Running title: root responses to an increase in growth medium strength

Highlight: The analysis of growth and orientation of *Arabidopsis thaliana* mutant roots affected

in root cap structure showed that intact root cap layers are required to trigger the root

responses to medium strength.

Abstract

During root progression in soil, root cap cells are the first to encounter obstacles. The root cap

is known to sense environmental cues, making it a relevant candidate for a mechanosensing

site. An original two-layer medium was developed in order to study root responses to growth

medium strength and the importance of the root cap in the establishment of these responses.

Root growth and trajectory of primary roots of *Arabidopsis thaliana* seedlings were investigated

using *in vivo* image analysis. After contact with the harder layer, the root either penetrated it or

underwent rapid curvature, allowing the reorientation of the root primary growth. The role of

the root cap in penetration and reorientation was investigated by analyzing mechanically the

responses of *Arabidopsis* mutant roots with altered caps. The primary root of *fez-2* mutant

lines, which has less root cap cell layers than wild-type roots, showed impaired penetration

ability. Conversely, *smb-3* roots of mutant lines, which display a higher number of root cap

cells, showed enhanced penetration abilities. This work highlights that alterations in root cap

structure affect the root responses to medium strength.

Key words: curvature, medium strength, Phytagel medium, root growth, root biomechanics,

root buckling, root cap

Abbreviations: COL, columella; LRC, Lateral Root Cap; SI, Sharpness Index; SMB, SOMBRERO

Introduction

Roots grow in a complex soil environment, which exhibits spatial and temporal heterogeneities, arising from sand particles, stones or hardpans of soil that constitute physical obstacles. The ability of roots to penetrate the soil and to overcome physical obstacles depends either on their ability to find a path of least resistance, or on their ability to generate sufficient force to go through the obstacle (Bengough *et al.*, 2009; Vollsnes *et al.*, 2010; Colombi *et al.*, 2017*a*). The force needed to penetrate a soil, provided by root growth, is determined by the soil strength. In natural conditions, soil strength ⌐ commonly estimated by penetrometer measurements ⌐ is in the range of 1MPa (Jin *et al.*, 2013; Kolb *et al.*, 2017). Soil heterogeneities may be overcome with soil compaction or soil drying, which results in an increase of soil strength.

In a few studies carried out in laboratories, changes in soil strength have been mimicked with simplified models appropriate to studying root responses. The elaboration of a two-layer Phytagel-based growth medium, containing two layers of distinct Phytagel concentrations, enables the roots to experience variations in medium strength when penetrating from one medium layer to another (Yamamoto *et al.*, 2008; Yan *et al.*, 2017). It has been shown that adjusting the Phytagel concentration regulated the strength of the growth medium (Schiavi *et al.*, 2016). In two-layer media experiments, where roots faced a compliant layer and a stiff layer consecutively, root tips were submitted to an increasing axial stress as they encountered the interface between the two layers. Roots which did not manage to generate sufficient force to penetrate the lower layer underwent buckling, dramatically changing the root growth trajectory and the root straightness (Massa and Gilroy, 2003; Silverberg *et al.*, 2012; Bizet *et al.*, 2016). The maximal axial force a root is able to exert before buckling, given by Euler's law, depends on the rigidity, the length of the growth zone, and the diameter of the root to the 4[th] power (Kolb *et al.*, 2017). In accordance with Euler's law, species with a higher increase in root diameter and/or decrease in root elongation in response to medium strength are more efficient in penetrating strong layers (Materechera *et al.*, 1992). Besides root diameter and growth zone length, it has been shown that root tip geometry was also an important feature modulating root response to medium strength (Ruiz *et al.*, 2016; Colombi *et al.*, 2017*b*). In compacted soils,

root tips with an acute opening angle grow faster than root tips with a blunted geometry (Iijima

et al., 2003*a*,*b*; Colombi *et al.*, 2017*b*). Colombi et *al.* (2017) showed that the root tip shape

influenced the penetration stress experienced by roots, by modulating the distribution of the

local compaction concentrated around the tip.

The shape of root tips arises from the shape of the root cap that covers and protects the root

apical meristem. The root cap is composed of the Columella (COL) located at the center of the

root cap and the Lateral Root Cap (LRC) (Arnaud *et al.*, 2010; Kumpf and Nowack, 2015). In

Arabidopsis thaliana, the mature COL and LRC are organized in 3 to 5 cell layers of increasing

age. Every formative division of root cap stem cells generates a new differentiating cell layer

which pushes the older root cap layers towards the root periphery. COL mature cells are known

to sense gravity, thanks to the amyloplasts they contain, conferring to the root cap the ability

to trigger tropisms in the root (Sato *et al.*, 2015). LRC mature cells principally act as secretory

cells which synthesize and export high molecular weight polysaccharide mucilage (Kumpf and

Nowack, 2015). Once mature, COL and LRC cell layers reach the periphery of the root cap

where they end up being sloughed off, allowing the root cap to keep a determinate, constant

size (Barlow, 2003). The balance between cell production and cell sloughing controls the root

tip shape. The study of root cap formation allowed the identification of the molecular actors

involved in cell production, differentiation and sloughing. Among these actors, FEZ and

SOMBRERO (SMB) transcription factors of the NAC family [NAM (NO APICAL MERISTEM), ATAF

(*ARABIDOPSIS* TRANSCRIPTION ACTIVATION FACTOR) and CUC (CUP-SHAPED COTYLEDON)]

were identified. FEZ specifically promotes periclinal divisions of COL and LRC stem cells

(Willemsen *et al.*, 2008). The *fez-2* mutant, resulting from the loss of function of *FEZ*, exhibits a

lower rate of periclinal division of the root cap stem cells which leads to the reduction of COL

and LRC cell layers compared to wild-type roots (Willemsen *et al.*, 2008; Bennett *et al.*, 2014).

SMB prevents the division of the differentiating root cap daughter cells and promotes their

differentiation. Additional COL and LRC cell layers are observed in the *smb-3* loss-of-function

mutant and the outer LRC cell layer reaches the end of the elongation zone (Willemsen *et al.*,

2008; Bennett *et al.*, 2010).

The objective of our study is to understand whether the structure of the root cap, controlled by
root cap integrity, is required in the establishment of root response to medium strength. We
hypothesized that the shape of the root cap could alter root responses in relation to an
increase in medium strength, with pointed caps facilitating root penetration and blunted or
domed caps complicating root penetration. We chose to study the root response to medium
strength through the *fez-2* and *smb-3 Arabidopsis thaliana* mutant lines. We characterized the
root tip geometry of *fez-2* and *smb-3* mutants precisely to understand whether the loss or the
gain of root cap cell layers impacted root tip shape. Then, the mutants were studied using a
novel experimental set-up which allows the quantitative imaging of the growth and the
orientation of the root. We cultivated Col-0, *fez-2* and *smb-3* roots in a two-layer Phytagel
medium of increasing mechanical resistance. Using a spatiotemporal analysis, we characterized
the penetration ratio, the root growth rate, and the root curvature of wild-type and mutant
roots from the upper layer to the lower layer. This simple experimental method allowed us to
describe three distinct root reorientations in response to increases in medium strength. We
discovered that the penetration ability and the reorientation in *fez-2* and *smb-3* were affected
in exactly the opposite way to what was hypothesized. Our results suggest that shape does not
seem to be the key feature influencing root response to medium strength.

Material and Methods

Plant growth medium

For all of the experiments, we used Phytagel as a substitute for Agar in the *in vitro* growth
medium. Phytagel is preferentially chosen for its higher temperature stability, higher
mechanical strength and better optical clarity (Schiavi et al., 2016). The growth medium
consisted of half-strength Murashige and Skoog (MS/2) medium adjusted to pH 5.7 and
containing Phytagel at different concentrations. The two-layer media consisted in the
superposition of an upper layer of growth medium containing 0.2% (w/v) of Phytagel and a
lower medium layer containing 0.2 % (w/v), 0.3% (w/v) or 0.5 % (w/v), called 0.2-0.2, 0.2-0.3
and 0.2-0.5 respectively (Fig. S1). The two-layer media were poured into specific culture
chambers which consisted in Polycarbonate boxes of 100 X 20 mm, and 60 mm height which

allowed root seedlings to grow vertically inside the growth medium. The two-layer media were

prepared as follows: 60 mL of sterilized MS/2 medium containing 0.2 % (w/v), 0.3% (w/v) or 0.5

% (w/v) of Phytagel were poured into the culture chambers under aseptic conditions, then,

these were rapidly closed and placed on ice. After 10 min, 20 mL of a sterilized MS/2 medium

containing 0.2 % (w/v) of Phytagel were poured over the lower medium layer.

Plant material

The wild-type strain of *Arabidopsis thaliana* (ecotype Columbia, Col-0) and *fez-2*, *smb-3*

Arabidopsis thaliana null-mutants used for the experiments were provided by B. Scheres's

laboratory (Willemsen *et al.*, 2008; Bennett *et al.*, 2010, 2014). The seeds were surface

sterilized with 30% (v/v) sodium hypochlorite solution, added to 2% (v/v) Triton X-100 for 10

min. After 5 washes in sterile distilled water, 10 seeds were gently placed on the surface of the

upper layer of the MS/2 two-layer medium. After sowing, the culture chambers were sealed

and kept at 4 °C in the dark for 2 days. Then, they were placed at 23 °C under a 16 h-

photoperiod.

Mechanical properties of Phytagel growth medium

To measure the mechanical properties of each concentration of the Phytagel medium,

compressive and penetration tests were performed, allowing the determination of the elastic

modulus and the mechanical penetration resistance of the media, respectively.

Determination of elastic modulus

Growth media containing from 0.2% to 1.2% (w/v) of Phytagel were poured into Petri dishes

having a thickness of 10 mm. 20x20x10 mm^3 Phytagel cubes were then cut and placed in the

Instron device (Tensile testing machine, Instron 5565). An uniaxial compression was applied on

the surface of the gels by the displacement of an upper plate with a constant loading rate of 0.5

mm.min^{-1}. A 100 N load cell recorded the force needed to deform the samples. Real-time load-

displacement data of each sample were exported by Console Bluehill 2 software and provided

characteristic stress-strain curves (Fig. 1A, B). After a typical contact step, the loading force

increased with the compression rate of the sample, before decreasing notably with the slipping

or the rupture of the sample. In the linear portion of the curve, *i.e.* when stress is proportional

to strain, the deformation is considered to be elastic (Schiavi *et al.*, 2016). The elastic modulus

(E) values of the Phytagel samples were determined from the ratio between the incremental

stress Δσ and the incremental strain rate Δε during elastic deformation by applying the Hooke's

law:

162
$$E = \frac{\Delta\sigma}{\Delta\varepsilon},$$

163 Where

$$\sigma = \frac{F}{A},$$
$$\varepsilon = \frac{l0}{\Delta l},$$

and F is the compression force (N), A is the surface area of the cross section of the sample (m^2)

on which the force is applied, l_0 is the initial thickness of the sample (m), Δl is the variation of

this thickness, *i.e.* the displacement of the plate (m). Compressive tests were conducted in 3

samples for each Phytagel concentration.

Determination of penetration resistance

Penetration tests were conducted in 0.2-0.2, 0.2-0.3 and 0.2-0.5 two-layer media. The samples

of the Phytagel medium were prepared as described above. A 2.5 mm diameter needle was

attached to a load cell (Instron, 100 N max) and driven downward through the growth medium

at a rate of 1 mm.min^{-1}. The real-time data were exported by Console Bluehill 2. The maximal

force required for the rupture of the surface of the lower layer was measured.

Time lapse photography and image processing

Root growth and orientation were time-lapsed 5 days after sowing. Two Nikon D7100 cameras

equipped with macro lenses (AF-S micro NIKKOR 60mm f/2.8G ED) and controlled by

DigiCamControl software, were placed on a rack in front of the culture chambers. Images of the

roots growing in the two-layer medium were acquired every 5 min during the period of the

experiment. All photographs were pre-treated using Rawtherapee software (v4.2.1) and were

then analyzed using a software developed by our laboratory, named RootStem Extractor. The
principle and procedures of the software were described in Chauvet *et al.* (2016). Briefly, the
software allowed the extraction of the topological skeleton of each root by screening pictures
perpendicularly to the root axis with a step of 0.3 pixel. The length of the root on each picture
and the local inclination as a function of the curvilinear abscissa can be computed from the
coordinates of the skeleton. The root length, as a function of time, allowed us to calculate the
root growth rate by the means of linear fit. By deriving the local inclination as a function of time
and space, curvature intensity along the root could be obtained. For each root, the curvature
intensity data are presented in a graph composed of 3 dimensions: a vertical axis corresponding
to root length from the beginning of the measurement, a horizontal axis corresponding to time
and a color bar representing the intensity of curvature.

Root tip geometry and determination of Sharpness Index

The root tips of Col-0, *fez-2* and *smb-3 Arabidopsis thaliana* lines were stained with a pseudo-
Schiff propidium iodide staining technique, as described in Truernit *et al.* (2008). They were
deposited between the slide and the cover slip and observed with a Zeiss LSM 800 laser
scanning microscope (Carl Zeiss Microscopy) coupled to a Apochromat X20 objective. From
longitudinal root observations, pictures of the longitudinal median section from the quiescent
center to the extremity of the root cap were extracted for each genotype. The area of the radial
section of the root tip was calculated using ImageJ software (v1.52d, http://imagej.nih.gov/ij/).
The vertical axis L_R and the radius of the root at the quiescent center R_R were measured (Fig.
2D1). We evaluated the Sharpness Index (SI) of the root tip according to the following equation
(Fig. 2D2):

$$SI = \frac{A_R - (L_R * 2R_R)/2}{L_R * 2R_R - (L_R * 2R_R)/2}$$

Where, A_R is the area of the root. The closer *SI* is to 0, the more the root tip is pointed.
However, the closer *SI* is to 1, the more the root tip has a rectangular shape.

Maximal root growth forces

The maximal forces roots were able to exert before buckling were estimated for Col-0, *fez-2*

and *smb-3* roots grown in 0.2-0.3 medium. After contact between the root tip and the lower

layer surface, the root tips deformed the lower medium until root deflection or root tip

deviation. The shape of the lower layer surface just before buckling was traced using ImageJ

software and the displacement (*d*) of the surface towards its initial position was measured (Fig.

4A). The shape of the root tip extremity was fitted by a circle of radius *R*. The maximal force

before root buckling F_{crit} applied on the surface by the root tip was then estimated using the

theory of contact between a sphere and a half-space (Hanaor *et al.*, 2015),

$$F_{crit} = \frac{4}{3} E^* R^{\frac{1}{2}} d^{\frac{3}{2}}$$

214   where

$$\frac{1}{E^*} = \frac{1 - v_1^2}{E_1} + \frac{1 - v_2^2}{E_2}$$

and *E1, E2* are the elastic moduli of the lower medium layer and of the root tip respectively,

and *v1, v2* are the Poisson's ratios of the lower layer and the root tip respectively. To calculate

E*, we assumed that E_2 was very close to E_1, allowing us to neglect the second part of the

formula. Moreover, according to the literature, *v1* is estimated at 0.5 (Ahearne *et al.*, 2005).

Statistical analysis

All data were statistically analyzed using R software (Rstudio, R.3.3.4, https://www.r-

project.org/). The Shapiro-Wilk test was used to test for normal distribution of the data and the

Bartlett test was used to test for homoscedasticity. One-way Analysis of Variance (ANOVA) or

two-way ANOVA were applied on the data, to determine whether the effect of each factor was

significant. When significant differences were obtained, post-hoc analyses based on the Tukey

test were applied on the data.

Results

Phytagel concentration in the medium impacts both medium rigidity and penetration resistance

The elastic modulus E, obtained from the stress-strain measurements, was calculated for the Phytagel concentrations ranging from 0.2 to 1.2% (Fig. 1A, B). The curve showed that E was highly dependent on the concentration of Phytagel and that this dependence was nonlinear (Fig. 1C). The 0.2%, 0.3%, 0.5% and 1.2% Phytagel growth medium showed E values of 3.07 ± 0.13 kPa, 7.60 ± 0.79 kPa, 31.93 ± 3.81 kPa and 158.30 ± 10.59 kPa, respectively (Fig. 1C).

Penetrometer assays were realized in 0.2-0.2, 0.2-0.3 and 0.2-0.5 two-layer media (Fig. 1D, E). The curves of the compression force as a function of the needle displacement revealed the same pattern for all of the growth media analyzed. The compression force increased markedly as the needle deformed the surface of the lower medium layer until a breaking point after which the loading force again became weak (Fig. 1E). The breaking point indicates the rupture of the lower layer surface and the penetration of the needle into the lower layer. The curves illustrate the presence of an interface between the two medium layers, whose rupture requires a specific force depending on the Phytagel concentration (Fig. 1E). The rupture of the 0.2% lower layer surface was induced by a force of 0.032 ± 0.004 N (Fig. 1F). The force required to break the 0.3% lower layer surface was of 0.054 ± 0.008 N and the force required to break the 0.5% lower layer surface was of 0.11 ± 0.009 N (Fig. 1F). The data obtained from the penetration tests showed that the strength of the interface increases significantly with the Phytagel concentration of the lower medium layer (Fig. 1F). The 0.2% layer surface was considered as the most brittle, the 0.5% surface as the strongest and the 0.3% surface as the middle-strength medium. The mechanical measurements illustrated a great regularity and repeatability for each Phytagel concentration, suggesting a stable behavior and good homogeneity of the Phytagel media.

In conclusion, the determination of Elastic modulus and penetration resistance allowed the characterization of the mechanical properties of the Phytagel media used for our experiments. An increase in the concentration of Phytagel in the medium generated increases in both medium rigidity and penetration resistance.

***fez-2* and *smb-3* *Arabidopsis thaliana* mutants exhibit an altered root cap shape**

Alterations in root cap cell layer number, cell division and differentiation in *fez-2* and *smb-3*
loss-of-function mutants have been precisely detailed (Willemsen *et al.*, 2008; Bennett *et al.*,
2010, 2014) and seem to have consequences on root tip shape. In this regard, it was important
to estimate this impact on each of the mutant lines studied. The reduction of COL and LRC cell
layers associated with the loss of function of *FEZ*, led to a shorter root cap narrower at the
base, compared to the wild-type (Fig. 2A, B, E). The estimation of the Sharpness Index (SI), of
0.30 ± 0.03 in *fez-2* roots and of 0.41 ± 0.03 in wild-type roots, showed that *fez-2* roots tips
were significantly more pointed than those of the wild-type ones (Fig. 2F). On the contrary, the
additional COL and LRC cell layers observed in the *smb-3* mutant, led to an increase in root
diameter at the quiescent center (Fig. 2C, E). The SI of the *smb-3* root tips was around 0.5 ±
0.02, illustrating their dome shape (Fig. 2F). The mean root diameter, measured at 500 μm from
the root tip, was not significantly different between *fez-2, smb-3* and wild-type roots,
suggesting that the mutations only impact the cap size (Fig. 2E). The analysis of the root growth
in the 0.2% one-layer Phytagel growth medium showed that the mean growth rate of wild-type
($3.10 ± 0.21$ μm.min^{-1}) and *fez-2* ($2.97 ± 0.18$ μm.min^{-1}) roots was not significantly different (Fig.
2G), whereas *smb-3* roots grew much more slowly, with a mean rate of $1.65 ± 0.08$ μm.min^{-1}
(Fig. 2G). Our characterization corroborates the analyses made by Bennett et al. (2010, 2014)
and the distinct phenotypes of the two mutant lines will allow the study of the role of root cap
shape in root responses to medium strength.

The medium strength impacts both the penetration ability and the growth rate of *Arabidopsis*
***thaliana* primary roots**

The penetration ability of *Arabidopsis thaliana* wild-type roots was observed in the 0.2-0.2, 0.2-
0.3 and 0.2-0.5 two-layer media (Table 1, Fig. S1). In the 0.2-0.2 medium, 100% of wild-type
roots progressed in the 0.2% upper layer, and crossed the interface to penetrate the 0.2%
lower layer. The analysis of root growth rate showed that only a short-term decrease in root
growth rate was observed 2 hours after contact (Fig. 3A). This slight decrease was followed by
the recovery of the initial root growth rate (Fig. 3A and Fig. 2G, 3μm.min^{-1}). The moment of

growth rate recovery corresponded to the rupture of the lower layer surface. One

representative example of *Arabidopsis thaliana* wild-type root penetration into the 0.2% lower

layer is presented in 3D graph (Fig. 5D), illustrating the impact of root penetration on root

growth rate and root straightness. On the graph, the increase in root length, as a function of

time, was linear before and after contact with the lower layer surface and confirmed the

transient impact of the root progression across the interface on root growth. The curvature

intensity was illustrated by the colors with the green indicating straight zones of the root. The

curvature intensity, along the penetrating root, lied between -1 and 1 mm^{-1}, illustrating root tip

oscillations around the vertical position during its growth (Fig. 5D). Right after contact, a zone

of slight curvature with an intensity below 2 mm^{-1} was initiated, represented by the horizontal

yellow strip on the graph. In the 0.2-0.3 two-layer medium, 60% of the *Arabidopsis thaliana*

primary roots penetrated the lower layer (Table 1). In parallel, the growth rate of penetrating

roots decreased by 30% within the first 30 minutes after contact with the lower layer surface

and recovered the initial growth rate within 300 minutes (Fig. 3D). The recovery of the initial

growth rate occurred after surface rupture and root penetration into the lower medium layer.

For some of the *Arabidopsis thaliana* penetrating roots, the decrease in the growth rate was

accompanied by the induction of a slight zone of curvature (data not shown). In the 0.2-0.5

two-layer medium, none of the *Arabidopsis thaliana* roots penetrated the lower layer (Table 1).

In conclusion, our results illustrated that the strength of the lower layer surface impacts the

penetration ability of *Arabidopsis thaliana* roots and affects their growth rate.

The root penetration ability is impaired in *fez-2* roots and enhanced in *smb-3* roots

To understand the role of root cap shape in response to medium strength, the penetration

abilities of *fez-2* and *smb-3* roots were compared to those of wild-type roots in 0.2-0.2, 0.2-0.3

and 0.2-0.5 two-layer media. All of the *smb-3* roots penetrated the 0.2% lower layer whereas

83% of the *fez-2* roots penetrated it (Table 1). The root growth rate of the mutant lines, 3.2

µm.min^{-1} for *fez-2* and 1.5 µm.min^{-1} for *smb-3*, was not significantly affected during root

penetration into the lower medium layer and corresponded to the initial root growth rate

observed in the 0.2% one-layer medium (Fig. 2G and Fig. 3B-C). Root penetration into the 0.3%

Phytagel lower layer was observed for 28% of the *fez-2* roots and 80% of the *smb-3* roots (Table

1). The root growth rate of *fez-2* and *smb-3* decreased by approximately 10% after root

penetration into the lower layer (Fig. 3E-F). In the 0.2-0.5 medium, only the *smb-3* roots

managed to penetrate the lower layer, with a penetration ratio of 18% (Table 1).

Thus, the *fez-2* roots exhibited an impaired penetration ability in response to mechanical

strength, whereas the *smb-3* roots exhibited an enhanced penetration ability. The root

penetration ratios revealed a variable ability of roots of a same genotype to face a given

mechanical resistance. To go further, the critical force that each root was able to exert before

reorienting was estimated.

Critical force of wild-type, *fez-2* and *smb-3* roots facing an increase in medium strength

To measure the critical force of wild-type (Col-0), *fez-2* and *smb-3* roots, the shape of the 0.3%

lower layer surface of elastic modulus E = 7.6 kPa was traced for each root just before root

curvature initiation (Fig. 4A). Wild-type roots displaced the surface at a mean maximal depth *d*

of 0.37 ± 0.03 mm, the *fez-2* roots displaced it at a mean *d* of 0.27 ± 0.01 mm and the *smb-3*

roots displaced it at a mean *d* of 0.49 ± 0.03 mm (Fig. 4B). The critical force was estimated

thanks to the values of *d*, of the radius of root tips and of the rigidity of the lower medium

layer. The mean critical force was estimated at 1.12 ± 0.14 10^{-9} N for wild-type roots, 0.66 ±

0.07 10^{-9} for *fez-2* N roots and 1.77 ± 0.18 10^{-9} N for *smb-3* roots, confirming that *smb-3* roots

were able to exert the highest force before curving while *fez-2* roots curved sooner (Fig. 4C).

_Arabidopsis thalian_a primary roots exhibit distinct curvature establishments in response to

an increase in medium strength

Representative examples of wild-type roots facing 0.3% and 0.5% lower layers are presented on

a spatiotemporal graph (Fig. 5, Fig. S3, Fig. S4). Following the contact with the 0.3% lower layer

surface, non-penetrating roots firstly grew downward against the surface and deformed it

without breaking it (Fig. 5B). A significant decrease in the root growth rate was observed (Fig.

5E). Within 2 h to 4 h after contact, a zone of curvature, illustrated by the red patch, was

observed at 500 μm to 1 mm from the root tip extremity. As roots continued to grow

downward, a second curvature zone, represented by the blue patch, appeared 3 h to 5 h after

the first one. The two curvatures exhibited a maximal intensity of 5 mm^{-1} (Fig. 3C). After their

formation, these zones of curvature were fixed in the root. Around 20 h after contact, the root

underwent a third curvature zone. This resulted in the deviation of the root tip axis allowing the

root tip to reorient its growth (Fig.5B). Root visualization in 2D did not allow the detailed

analysis of the third curvature. From now on, the term "curly phenotype" will be used to

describe roots drawing curls in response to lower layer surface strength.

In the 0.2-0.5 medium, the roots grew vertically against the stiff lower layer surface for 30 min

to 2 h after contact (Fig. 5C, F). The root growth rate decreased significantly within the first few

minutes following the contact with the lower layer surface (Fig. 5F). A first curvature was

established at approximately 400 μm from the root tip within 2 h following contact (Fig. S4).

The first curvature induced the deviation of the root tip, allowing the root to grow horizontally

along the surface of the lower layer. This first curvature exhibited a maximal intensity of 6 mm^{-1}

and was confined to its zone of formation. While the root tip reoriented its growth, a second

zone of curvature was formed at 300 μm from the root tip around one hour after the

establishment of the first curvature with a maximal intensity corresponding to 4 mm^{-1}(Fig. 5F).

Although the first curvature was fixed after its formation, the second one moved to stay at the

same distance from the root apex during root growth along the surface. During root

reorientation, no recovery of the initial growth rate was observed (Fig.S2). From now on, the

term "step-like phenotype" will be used to describe roots responding to the change in medium

strength in this way (Fig. S4).

The *fez-2* and *smb-3* mutant roots exhibit altered curvature establishment

The 20% of *fez-2* roots which did not penetrate the 0.2% lower layer exhibited the curly

phenotype (Fig. 6A, Fig.S5). The first curvature was initiated at 500 μm from the root apex and

exhibited a maximal intensity of 5 mm^{-1} (Fig. 6B). Root growth rate decreased rapidly and

notably after the contact with the lower layer surface. The *fez-2* root reorientation in the 0.2-

0.2 medium was very similar to wild-type root reorientation in the 0.2-0.3 medium (Fig. 5E and

Fig. 6B). The spatiotemporal analysis of deviation of *fez-2* root tips in the 0.2-0.3 and 0.2-0.5

media showed that *fez-2* non-penetrating roots had a step-like phenotype (Fig. 6C-F, Fig. S6, Fig. S7). In both media, the first curvature was formed within 1 h following contact at around 500 μm from the root apex and the second curvature was formed right after, very close to the root apex. The maximal intensity of the two curvatures was equal to 4 mm^{-1} in the 0.2-0.3 medium and to 2 mm^{-1} in the 0.2-0.5 medium, suggesting that the curvature intensity of *fez-2* roots depends on the strength of the lower layer (Fig. 6D and F). The second curvature intensity was very weak in step-like shaped *fez-2* roots compared to step-like wild-type roots. The root growth rate of the step-like shaped *fez-2* roots was linear after contact and during root growth reorientation (Fig. 6D and F).

In the 0.2-0.3 medium, *smb-3* non-penetrating roots exhibited the curly phenotype as observed for wild-type roots (Fig. 7A). As *smb-3* roots grow much more slowly than other genotypes, the curls were only initiated 6 h after the contact with the lower layer surface (Fig. 7B). The maximal intensity of curvatures was close to 5 mm^{-1} (Fig. 7B). In the 0.2-0.5 medium, the *smb-3* non-penetrating roots reoriented either by undergoing several deflections (Fig. 7C, Fig. S9), or by establishing two curvatures (Fig. 7E). In the first case, the roots exhibited large waves with maximal curvature intensities reaching 8 mm^{-1} (Fig. 7D). Moreover, the root growth rate of the *smb-3* roots was unchanged during the formation of the waves (Fig. 7D). In the second case, the two curvatures exhibited a similar maximal intensity of 5 mm^{-1}. The first and second curvatures were initiated at 300 μm from the root apex (Fig. 7F). The second curvature intensity was significantly higher than in wild-type roots. For these roots, the growth rate decreased slightly after the contact with the lower layer surface (Fig. 7F, Fig. S2). As in wild-type roots, no recovery of the initial root growth rate was observed after complete root reorientation (Fig. S2).

Discussion

Our experimental set-up has proved insightful for understanding root responses to medium strength. The precise characterization of media mechanical properties showed a strong correlation between medium rigidity and penetration resistance. Moreover, the precise spatiotemporal analysis of root growth rate and orientation allowed us to delve further into the

394   discrimination of passive and active mechanisms involved in root reorientation. The achieved

395   characterization of root reorientation led to a better understanding of the role of the root cap

396   in root responses to mechanical stresses.

397   Root reorientation involves both passive and active mechanisms

In our experimental system, roots grew through an upper medium layer of compliant strength

until encountering a lower layer, where penetration requires an increase in root growth

pressure. The contact between the wild-type root tip and the middle-strength and the

strongest lower layer surfaces led to a rapid decrease in root growth rate. This decrease,

observed for both penetrating and non-penetrating roots, was comparable to that observed in

poplar roots submitted to a physical obstacle, and may arise from the dedication of turgor

pressure to root penetration rather than to cell elongation (Bizet *et al.*, 2016). The root growth

against the lower layer surface creates a progressively increasing compression force on the root

(Kolb *et al.*, 2017). If the compression force reaches a critical value while the root is still in

contact with the obstacle, Euler's buckling can occur (Bizet *et al.*, 2016). We estimated that

non-penetrating wild-type roots supported a critical compression force of $1.12 \ 10^{-9}$ N before

buckling. These forces are 10^6 times less important than the critical buckling force measured on

Poplar adventitious roots (Bizet *et al.*, 2016). This difference may be explained by the difference

in root diameter between the two species and by the differences in root rigidity.

Buckling was observed for wild-type, *fez-2* and *smb-3* non-penetrating roots and led to

three distinct morphologies depending on the genotype and on the strength of the lower layer

surface. When encountering an intermediate medium strength, non-penetrating roots mostly

exhibited a curly-shape. This shape was already observed in *Medicago truncatula* roots growing

in a two-layer Phytagel media (Silverberg *et al.*, 2012). From a mathematical model based on

the theory of buckling rods, the authors assigned the curly morphology of M*edicago* roots to a

combination of helical buckling and simultaneous twisting near the root tip. Lateral

containment by the upper medium layer seemed to be a key feature modulating root helical

buckling (Silverberg *et al.*, 2012). Poplar roots grown in non-constraining hydroponic solution

and submitted to a physical obstacle buckled with a long wavelength curve shape (Bizet *et al.*,

2016). In our experimental system, some *smb-3* non-penetrating roots adopted this shape that

we called the wavy phenotype. In two-layer media, as all of the upper layers had the same

mechanical strength, the various shapes adopted by roots cannot be attributed to their

constriction level but more probably to their variability of anchorage or diameter. For most of

the curly-shaped and wavy-shaped roots, the first curvature was initiated and confined in a

zone between 400 and 500 μm from the root tip, corresponding to a zone of mechanical

weakness (Bizet *et al.*, 2016). All these observations suggest that curly and wavy phenotypes

are mainly driven by the mechanical response of the root acting as a "growing" rod blocked by

the obstacle.

The step-like phenotype was adopted by roots mostly in response to the highest increase in

medium strength. In this case, the first curvature induced the deviation of the root tip, which

implied a change in mechanical forces applied on the root tips from axial to asymmetric radial

forces. The rapidity and position of curvature initiation suggest that it may be interpreted as a

buckling. The first curvature was followed by the initiation of a second one, closer to the root

apex, allowing the root tip to regain the vertical position. This second curvature provided the

root tip with a steady form from changing cells, and was previously described as the result of

the interaction between gravitropic and thigmotropic responses (Massa and Gilroy, 2003). The

root tip deviation was preceded by a reduction in root growth rate which did not recover after

relaxation of axial mechanical stresses. The initiation of the second curvature and the

maintenance of the reduced root growth rate suggest that, even if the first curvature was

initially provoked by a buckling, the further root growth direction may be reoriented in an

active way.

The different patterns of curvature establishment suggest that the mechanical strength of the

lower layer surface determines the ensuing root response (buckling or bending). A middle-

strength medium surface allows the root tips to sink into it, preventing their deviation after

root buckling. On the contrary, a strong medium surface is barely deformed by the root tips,

which may be free to deviate and may trigger an active root reorientation.

Root cap shape is not the key feature impacting root responses to an increase in medium strength

Based on previous studies (Vollsnes *et al.*, 2010; Ruiz *et al.*, 2016; Colombi *et al.*, 2017*b*), we hypothesized that root cap shape, by changing the distribution of axial mechanical stresses around the root tip, influenced root penetration ability and root response to medium strength. It has been shown that a pointed root tip increased the concentration of axial stresses at the tip forefront rather than around the tip, resulting in enhanced crack initiation and reduced penetration stress. Contrary to what was expected, *fez-2* roots, with an acute tip, exhibited reduced penetration abilities whereas *smb-3* roots, with a dome-shaped tip, exhibited enhanced penetration abilities compared to wild-type roots. These conflicting results suggest that root cap shape may not be the key feature determining root response to medium strength. To confirm this, attention should be paid to the cap rigidity of wild-type and mutant roots. The loss or the accumulation of root cap cell layers could alter not only the cap shape but also the rigidity. A softer cap would deform more easily under an increase in medium strength, resulting in the modification of its shape.

The difference in penetration abilities between mutant and wild-type roots may be assigned to their different critical buckling forces. The *fez-2* non-penetrating roots mostly exhibited a step-like shape even when subjected to a middle-strength medium, indicating that the low resistance to buckling of *fez-2* roots did not allow them to deform the lower layer surface sufficiently to prevent the root tip deviation. On the contrary, the higher buckling resistance of *smb-3* roots allowed them to exert more axial stress on the lower layer surface and thus to penetrate the strong layers more easily. When *smb-3* roots did not penetrate, they mostly exhibited a curly phenotype, illustrating the fact that *smb-3* root tips deformed the lower layer surface more than wild-type roots. The differences in the root critical buckling force between the three *Arabidopsis* lines may be explained either by the differences in diameter or rigidity at the buckling root zone, or by the differences in growth zone length (Kolb *et al.*, 2017). Altered root cap shape or rigidity could not have influenced the resistance to buckling of the mutant roots. Our observations of the mutant root phenotypes did not reveal any alterations in root diameter, growth zone length or root hair initiation, suggesting that the root rigidity may be

altered negatively in *fez-2* and positively in *smb-3* mutant lines. We may hypothesize that the

increase in medium strength is perceived by root cap cells and that signal transduction in the

elongation zone triggers mechanisms leading to the regulation of tissue rigidity. For example,

root rigidity might be increased by auxin-dependent cell wall strengthening (Bengough *et al.*,

2011; Muratov and Baulin, 2015; Tsugama *et al.*, 2016). In *fez-2* roots, the loss of cap cell layers

may alter the mechanosensitivity and thus the regulation of cell wall extensibility in the

elongation zone in response to an increase in medium strength. The fact that the growth rate of

both *fez-2* penetrating and reorienting roots was not affected by medium strength consolidates

this hypothesis. Moreover, we observed that the second curvature of step-like shaped roots,

established through an active process, was of a lesser intensity in *fez-2* roots and of a greater

intensity in *smb-3* roots compared to wild-type roots. As *fez-2* and *smb-3* roots appear to have

an effective gravitropic response, the alteration in the curvature establishment may be the

result of a functional alteration in mutant root caps. The presence of a intact root cap with a

precise number of LRC and COL cell layers may be required to trigger the signal perception and

transduction processes within the root and thus to allow the root to respond to an increase in

medium strength.

Conclusion

During their progression in a soil, roots frequently face physical heterogeneities that they have

to displace by generating a sufficient mechanical force or circumvent by reorienting their

growth. The experimental approach presented here provides new insights into the role of the

root cap on root penetration and reorientation in response to medium strength. In two-layer

media, the roots adopted three main responses to an increase in medium strength: root

penetration in a brittle medium, root buckling without tip deviation leading to a curly

phenotype in a middle-strength medium, and root buckling followed by tip deviation leading to

a step-like phenotype in the strongest medium. The responses to medium strength were

altered in *fez-2* and *smb-3* mutant roots, which exhibit less and additional cap cell layers,

respectively. The alteration in the penetration ability, reduced in *fez-2* roots and enhanced in

smb-3 roots was the result of altered resistance to buckling rather than of altered root cap

geometries. From these results arises the hypothesis that intact root cap layers are required to trigger the root responses to medium strength.

Supplementary data

Fig. S1. The two-layer Phytagel media 0.2-0.2, 0.2-0.3 and 0.2-0.5 were used to study the responses of *Arabidopsis thaliana* roots to an increase in medium strength.

Fig. S2. Growth rate of reorienting roots of wild-type, *fez-2*, *smb-3* *Arabidopsis thaliana* lines in 0.2-0.5 two-layer medium

Fig. S3. Time-lapse movie of a reorienting wild-type root in the 0.2-0.3 two-layer medium

Fig. S4. Time-lapse movie of a reorienting wild-type root in the 0.2-0.5 two-layer medium

Fig. S5. Time-lapse movie of a reorienting *fez-2* root in the 0.2-0.2 two-layer medium

Fig. S6. Time-lapse movie of a reorienting *fez-2* root in the 0.2-0.3 two-layer medium

Fig. S7. Time-lapse movie of a reorienting *fez-2* root in the 0.2-0.5 two-layer medium

Fig. S8. Time-lapse movie of a reorienting *smb-3* root in the 0.2-0.5 two-layer medium

Acknowledgements

We would like to thank Ben Scheres and Viola Willemsen (University of Utrecht, The Netherlands) for the kind gifts of mutant lines, Nicole Brunel-Michac (Université Clermont Auvergne, INRA, PIAF) for her help and advice on microscopy, Patricia Mabrut for English editing (Université Clermont Auvergne). This research received financial support from the French Space Agency (CNES) and by the recipient of INRA programs. J.R. has a doctoral grant of the French National Ministry of Education and Research. H.C is supported by a Postdoctoral grant from the Auvergne-Rhône Alpes Council with cofunding from FEDER. F. B. was supported by a Postdoctoral grant from the CNES.

References

Ahearne M, Yang Y, El Haj AJ, Then KY, Liu KK. 2005. Characterizing the viscoelastic properties

of thin hydrogel-based constructs for tissue engineering applications. Journal of the Royal

Society Interface **2**, 455–463.

Arnaud C, Bonnot C, Desnos T, Nussaume L. 2010. The root cap at the forefront. Comptes

Rendus - Biologies **333**, 335–343.

Barlow PW. 2003. The Root Cap: Cell Dynamics , Cell Differentiation and Cap Function. Journal

of Plant Growth Regulation, 261–286.

Bengough AG, Hans J, Bransby MF, Valentine TA. 2009. PIV as a method for quantifying root

cell growth and particle displacement in confocal images. Microscopy Research and Technique

73, N27-36.

Bengough AG, McKenzie BM, Hallett PD, Valentine TA. 2011. Root elongation, water stress,

and mechanical impedance: a review of limiting stresses and beneficial root tip traits. Journal of

Experimental Botany **62**, 59–68.

Bennett T, Sanchez-perez GF, Campilho A, Willemsen V, Snel B, Scheres B. 2010. SOMBRERO ,

BEARSKIN1 , and BEARSKIN2 Regulate Root Cap Maturation in Arabidopsis. The Plant Cell **22**,

640–654.

Bennett T, van den Toorn A, Willemsen V, Scheres B. 2014. Precise control of plant stem cell

activity through parallel regulatory inputs. Development **141**, 4055–4064.

Bizet F, Bengough AG, Hummel I, Bogeat-Triboulot M-B, Dupuy LX. 2016. 3D deformation field

in growing plant roots reveals both mechanical and biological responses to axial mechanical

forces. Journal of Experimental Botany **67**, 5605–5614.

Chauvet H, Pouliquen O, Forterre Y, Legué V, Moulia B. 2016. Inclination not force is sensed by

plants during shoot gravitropism. Scientific Reports **6**, 1–8.

Colombi T, Braun S, Keller T, Walter A. 2017*a*. Artificial macropores attract crop roots and

enhance plant productivity on compacted soils. Science of the Total Environment **574**, 1283–1293.

Colombi T, Kirchgessner N, Walter A, Keller T. 2017*b*. Root Tip Shape Governs Root Elongation Rate under increased soil strength. Plant Physiology **174**, 2289–2301.

Hanaor DAH, Gan Y, Einav I. 2015. Contact mechanics of fractal surfaces by spline assisted discretisation. International Journal of Solids and Structures **59**, 121–131.

Iijima M, Barlow PW, Bengough AG. 2003*a*. Root cap structure and cell production rates of maize (Zea mays) roots in compacted sand. New Phytologist **160**, 127–134.

Iijima M, Higuchi T, Barlow PW, Bengough AG. 2003*b*. Root cap removal increases root penetration resistance in maize (Zea mays L .). Journal of Experimental Botany **54**, 2105–2109.

Jin K, Shen J, Ashton RW, Dodd IC, Parry MAJ, Whalley WR. 2013. How do roots elongate in a structured soil? Journal of Experimental Botany **64**, 4761–4777.

Kolb E, Legué V, Bogeat-Triboulot M-B. 2017. Physical root – soil interactions. Physical Biology **14**.

Kumpf RP, Nowack MK. 2015. The root cap: a short story of life and death. Journal of Experimental Botany **66**, 5651–5662.

Massa GD, Gilroy S. 2003. Touch modulates gravity sensing to regulate the growth of primary roots of Arabidopsis thaliana. Plant Journal **33**, 435–445.

Materechera SA, Alston AM, Kirby JM, Dexter AR. 1992. Influence of root diameter on the penetration of seminal roots into a compacted subsoil. Plant and Soil **144**, 297–303.

Muratov A, Baulin VA. 2015. Biophysical Chemistry Mechanism of dynamic reorientation of cortical microtubules due to mechanical stress. Biophysical Chemistry **207**, 82–89.

Ruiz S, Straub I, Schymanski SJ. 2016. Experimental Evaluation of Earthworm and Plant Root Soil Penetration – Cavity Expansion Models Using Cone Penetrometer Analogs. Vadose Zone

Journal.

Sato EM, Hijazi H, Bennett MJ, Vissenberg K, Swarup R. 2015. New insights into root

gravitropic signalling. Journal of Experimental Botany **66**, 2155–2165.

Schiavi A, Cuccaro R, Troia A. 2016. Strain-rate and temperature dependent material

properties of Agar and Gellan Gum used in biomedical applications. Journal of the Mechanical

Behavior of Biomedical Materials **53**, 119–130.

Silverberg JL, Noar RD, Packer MS, Harrison MJ, Henley CL, Cohen I, Gerbode SJ. 2012. 3D

imaging and mechanical modeling of helical buckling in Medicago truncatula plant roots.

Proceedings of the National Academy of Sciences **109**, 16794–16799.

Tsugama D, Liu S, Takano T. 2016. The bZIP Protein VIP1 Is Involved in Touch Responses in.

Plant Physiology **171**, 1355–1365.

Vollsnes a. V., Futsaether CM, Bengough a. G. 2010. Quantifying rhizosphere particle

movement around mutant maize roots using time-lapse imaging and particle image

velocimetry. European Journal of Soil Science **61**, 926–939.

Willemsen V, Bauch M, Bennett T, Campilho A, Wolkenfelt H, Xu J, Haseloff J, Scheres B. 2008.

Article The NAC Domain Transcription Factors FEZ and SOMBRERO Control the Orientation of

Cell Division Plane in Arabidopsis Root Stem Cells. Developmental Cell **15**, 913–922.

Yamamoto C, Sakata Y, Taji T, Baba T, Tanaka S. 2008. Unique ethylene-regulated touch

responses of Arabidopsis thaliana roots to physical hardness. Journal of Plant Research **121**,

509–519.

Yan J, Wang B, Zhou Y. 2017. A root penetration model of Arabidopsis thaliana in phytagel

medium with different strength. Journal of Plant Research **130**, 941–950.

Tables

Table 1. Penetration ability of Arabidopsis wild-type (Col-O), *fez-2*, and *smb-3* loss-of-function

mutant primary roots in two-layer media. Penetration ability is represented by the percentage

of roots that were able to penetrate 0.2%, 0.3% and 0.5% lower layers. For each genotype and

condition, 6 replicates of 10 roots were observed 8 days after germination. The values indicate

the average percentage over the six replicates, + or − SD. Letters indicate means significantly

different with P<0.05 (two-way ANOVA with Tukey test)

Figure legends

Fig. 1. Mechanical properties of Phytagel growth media illustrated by the determination of

Elastic modulus (A-C) and by penetrometer capacity (D-F). (A) Representation of Phytagel cube

placed between the two plates of the testing machine (Instron). (B) Stress-strain curves

resulting from compressive tests performed for medium cubes of Phytagel concentration

ranging from 0.2% to 1.2%. (C) The Elastic modulus E was estimated using the slope of the

linear portion of the curve. The circles are real data obtained for each Phytagel concentration,

squares are means, n=4. (D-F) Penetrometer assays to determine mechanical resistance of the

lower layer surface in two-layer media. (D) Representation of the needle progression in a 0.2-

0.5 two-layer medium in the mechanical testing machine. (E) Measurement of force during the

penetration of the needle in the lower medium layer of 0.2%, 0.3% and 0.5% Phytagel. The

displacement 0 corresponds to the first contact between the needle tip and the lower layer

surface (F) Penetrometer resistance of the lower layer surface of 0.2%, 0.3% or 0.5 % Phytagel.

Circles are real data, squares are means, n=4.

Fig. 2. Root shape of wild-type, *fez-2* and *smb-3 Arabidopsis thaliana* root tip

(A-D) Representative Confocal microscope images of Ps-PI stained Col-0 (A), *fez-2* (B) and *smb-3*

(C) 5 day-old roots. Asterisks indicate the quiescent center. White bar = 100 μm. (D1)

Representation of the root tip section used to measure root radius at the quiescent center (R_R)

and root length from the quiescent center to the root tip extremity (L_R). (D2) The root tip area

measured in D1 was compared to the areas of a triangle and a rectangle of the same

dimensions (L_R and R_R) to obtain the Sharpness Index (SI) of the root tip. (E) Diameter of Col-0,

fez-2 and *smb*-3 roots at the quiescent center (filled markers) and at 500 µm from the root tip

extremity, in the elongation zone (empty markers). Circles are real values and squares are

means, n=8 (F) SI of Col-0, *fez-2, smb-3* root tips. The SI lies between 0 - value corresponding to

a triangle - and 1- value corresponding to a rectangle. Circles are real values and squares are

means, n=8. (G) Root growth rate of Col-0, *fez-2, smb-3* roots growing in compliant one-layer

0.2% Phytagel-medium. Circles are real values and squares are means, n=12. Letters indicate

means significantly different with P<0.05 (ANOVA with Tukey test)

Fig. 3. Growth rate of penetrating roots of Col-0, *fez-2, smb-3* Arabidopsis thaliana lines in

two-layer media

(A-C) Root growth rate of Col-0 (A), *fez-2* (B), *smb-3* (C) roots which penetrated a compliant

lower medium layer (0.2% Phytagel). (D-F) Root growth rate of Col-0 (D), *fez-2* (E), *smb-3* (F)

roots which penetrated a middle strength lower medium layer (0.3% Phytagel). For all graphs,

circles are real values and lines are mean, 6<n<10.

Fig. 4. Critical buckling force of Col-0, *fez-2* and *smb-3* roots

(A) Schematic example of a root tip growing against a 0.3% lower layer surface of elastic

modulus E= 7.6 kPa. The gray curve represents the shape of the lower layer surface just before

root curvature initiation. d is the maximal interface displacement. R is the radius of the root tip

in contact with the surface.

(B) Shapes of the lower layer surface of Elastic modulus E = 7.6 kPa deformed by Col-0, *fez-2*

and *smb-3* root tips just before their curvature initiation. Lines are means, n=10. Dashed lines

correspond to the initial shape of the lower layer surface.

(C) Evaluation of the critical force Fcrit that the Col-0, *fez*-2 and *smb-3* roots were able to exert

on the lower layer surface before reorienting. Fcrit was computed according to the equation

depending on root tip radius R, on the maximal displacement d and on the elastic modulus of

the lower layer E. (n=10). Letters indicate means significantly different with P<0.05 (ANOVA
with Tukey test)

Fig. 5. Spatiotemporal characterization of *Arabidopsis* Col-0 root reorientation in two-layer
media. (A) Representative Col-0 root penetration into a compliant lower layer (0.2% Phytagel).
Images represent the root 2 h before contact with the lower layer surface and 0 h, 3 h and 6 h
after contact. (A1) Skeleton of the root during the penetration. (B) Representative Col-0 root
response to contact with a 0.3% lower layer at 0 h, 5 h, 10 h and 15 h after contact. This
response is called curly phenotype. (B1) Skeleton of the root during the beginning of its
reorientation, from 4 h before contact to 6 h after contact. (C) Representative wild-type root
response to contact with a 0.5% lower layer at 0 h, 3 h, 8 h and 15 h after contact. (C1) Skeleton
of the root during its reorientation, from 4 h before the contact to 14 h after the contact. White
bars correspond to 500 µm. Red arrows indicate lower layer surface (D-F) Spatiotemporal
analyses of a root penetrating a compliant lower layer (D), a root reorienting in response to
middle strength lower layer (E) and a root reorienting in response to stiff lower layer (F). For 3D
graphs, root length as a function of time is presented. The colors represent curvature intensity:
negative for right-handed curvatures and positive for left-handed curvatures. The green color
corresponds to the straight zones of the roots. The dashed lines indicate the contact.

Fig. 6. Spatiotemporal characterization of *fez-2* root reorientation in two-layer media

(A-B) Squeletton (A) and spatio-temporal analysis (B) of a root reorienting after contact with a
0.2% lower layer. (C-D) Skeleton (C) and spatio-temporal analysis (D) of a root reorienting after
contact with 0.3% lower layer. (E-F) Squeletton (E) and spatio-temporal analysis (F) of a root
reorienting after contact with a 0.5% lower layer. Dashed lines indicate the contact between
root tip and lower layer surface.

Fig. 7. Spatiotemporal characterization of *smb-3* root reorientation in two-layer media

(A-B) Squeletton (A) and spatio-temporal analysis (B) of a root reorienting after contact with a
0.3% lower layer. (C-D) Skeleton (C) and spatio-temporal analysis (D) of a root reorienting
through waves formation after contact with a 0.5% lower layer. (E-F) Squeletton (E) and spatio-

temporal analysis (F) of a root reorienting through a step-like phenotype after contact with a

0.5% lower layer. Dashed lines indicate the contact between root tip and lower layer surface.

Genotype	Two-layer medium		
Concentrations of the two layer media (% phytagel)	0.2-0.2	0.2-0.3	0.2-0.5
Rupture Force (N)	0.032 ± 0.004	0.054 ± 0.008	0.11 ± 0.009
Col-0	100.00 ± 0.00 [a]	60.00 ± 3.65 [c]	1.67 ± 1.67 [e]
fez-2	83.33 ± 4.94 [b]	28.33 ± 3.07 [d]	0.00 ± 0.00 [e]
smb-3	100.00 ± 0.00 [a]	80.00 ± 5.16 [b]	18.33 ± 4.77 [d]

Table 1. Penetration ability of Arabidopsis wild-type (Col-O), *fez-2*, and *smb-3* loss-of-function mutant primary roots in two-layer media.

Fig. 1. Mechanical properties of Phytagel media.

Fig. 2. Phenotypes of Col-0, *fez-2* and *smb-3* roots.

Fig. 3. Root growth rate of Col-0, *fez-2*, *smb-3* Arabidopsis lines in two-layer media

Fig. 4. Critical buckling force of Col-0, *fez-2* and *smb-3* roots.

Fig. 5. Spatiotemporal characterization of *Arabidopsis* Col-0 root reorientation in two-layer media.

Fig. 6. Spatiotemporal characterization of *fez-2* root reorientation in two-layer media.

Fig. 7. Spatiotemporal characterization of *smb-3* root reorientation in two-layer media.