Prospects for enhancing leaf photosynthetic capacity by manipulating mesophyll cell morphology

Running title: Photosynthetic capacity and leaf cell morphology

Tao Ren[1,2], **Sarathi M Weraduwage**[3], **Thomas D. Sharkey**[2,3, *]

[1]College of Resources and Environment, Huazhong Agricultural University, Wuhan 430070, China

[2]Department of Biochemistry and Molecular Biology, Michigan State University, East Lansing, Michigan 48824, USA

[3]Department of Energy Plant Research Laboratory and Plant Resiience Institute, Michigan State University, East Lansing, Michigan 48824, USA

Email addresses

Tao Ren 任涛 <rentao@mail.hzau.edu.cn>

Sarathi Weraduwage <sarathi@msu.edu>

*To whom correspondence should be addressed.

Thomas D. Sharkey <tsharkey@msu.edu>

Phone +1 (517) 353-3257

Number of words 6381

Number of Figures 3 (all in color)

Number of tables 2

Highlight: Cell morphology in leaves affects photosynthesis by controlling CO_2 diffusion and light distribution. Recent work has uncovered genes that control cell size, shape, and number paving the way improved photosynthesis.

Abstract

Leaves are beautifully specialized organs designed to maximize the use of light and

CO_2 for photosynthesis. Engineering leaf anatomy therefore brings great potential to

enhance photosynthetic capacity. Here we review the effect of the dominant leaf

anatomical traits on leaf photosynthesis and confirm that a high chloroplast surface

area exposed to intercellular airspace per unit leaf area (S_c) is critical for efficient

photosynthesis. The possibility of improving S_c through appropriately increasing

mesophyll cell density is further analyzed. The potential influences of modifying

mesophyll cell morphology on CO_2 diffusion, light distribution within the leaf, and

other physiological processes are also discussed. Some potential target genes

regulating leaf mesophyll cell proliferation and expansion are explored. Indeed, more

comprehensive research is needed to understand how manipulating mesophyll cell

morphology through editing the potential target genes impact leaf photosynthetic

capacity and related physiological processes. This will pinpoint the targets for

engineering leaf anatomy to maximize photosynthetic capacity.

Key words: chloroplast surface area exposed to intercellular airspace, light

distribution, mesophyll cell density, mesophyll conductance, photosynthesis, target

genes

Abbreviations: A_n, photosynthesis rate; f_{ias}, fraction of intercellular airspace; g_m,

mesophyll conductance; g_s, stomatal conductance; L_{chl}, chloroplast length; LMA, leaf

mass per unit leaf area; S_c, chloroplast surface area exposed to intercellular airspace

per unit leaf area; S_{mes}, mesophyll surface area exposed to intercellular airspace; T_{chl},

chloroplast thickness T_{cw}, mesophyll cell wall thickness; T_L, leaf thickness; T_{mes},

mesophyll layer thickness.

Introduction

Increasing the conversion efficiency of intercepted radiation into biomass through

enhancing photosynthesis has emerged as a promising focus to enhance crop

production for meeting needs of the increasing population and for plant resilience in

the face of the changing global climate (Long *et al.*, 2015*b*; Ort *et al.*, 2015; Souza *et*

al., 2017). Some current approaches to improving photosynthesis in C_3 crops, include

conversion to C_4 metabolism, engineering a photorespiratory bypass, optimizing

regeneration of RuBP etc. It is estimated that these could increase the light conversion

efficiency by 5%-60% compared with today's best cultivars (Long *et al.*, 2015*b*).

Using the rice GRCROS model, Yin and Struik (2017) simulated nine genetic

engineering strategies for enhancing C_3 leaf photosynthesis to improve crop

production under the present climate and the climate predicted for 2050. Although the

knowledge of photosynthesis has exploded recently, there is no doubt that it is still a

major challenge to understand the mechanisms, vulnerabilities, and potentials for

improvement of photosynthesis (Niinemets *et al.*, 2017).

Leaves are beautifully specialized organs evolved for efficient photosynthesis. Their

morphological differences can explain substantial variations in photosynthetic

properties among different species or the same species under different environmental

constraints (Oguchi *et al.*, 2005; Terashima *et al.*, 2011; Peguero-Pina *et al.*, 2017;

Tsukaya, 2018). For example, alterations in photosynthetic performance of sun and

shade leaves of the same plant are intimately connected with leaf anatomy (Terashima

et al., 2001, 2006; Oguchi *et al.*, 2005; Théroux-Rancourt and Gilbert, 2017). The

inseparable relationship between leaf architecture and photosynthesis thus brings

great potential to enhance photosynthetic efficiency by modifying leaf anatomy.

Tholen *et al.* (2012) described how altering leaf anatomy could significantly influence

light distribution and CO_2 diffusion in the leaf. Nonetheless, precise targets for

engineering leaf anatomy are still unknown because of poor understanding of the

molecular mechanisms controlling leaf anatomy.

Leaf mass per unit leaf area (LMA), one important leaf morphological trait that

affects photosynthesis, is also related to environmental stress and longevity of plants

(Niinemets, 2001; Poorter *et al.*, 2009; de la Riva *et al.*, 2016). There exists

considerable variation regarding the correlation between LMA and photosynthesis,.

Negative relationships between LMA and photosynthesis are associated with higher

investment in nonphotosynthetic tissue, such as cell walls (Onoda *et al.*, 2004, 2017;

Hassiotou *et al.*, 2010) and lower mesophyll conductance (Flexas *et al.*, 2008;

Niinemets *et al.*, 2009; Weraduwage *et al.*, 2016). Positive relationships are associated

with improvement of chloroplast exposure to intercellular airspace (Peguero-Pina *et*

al., 2017) and chloroplast CO_2 concentration, which ultimately enhances the

carboxylation rate (Cai *et al.*, 2014; Flexas *et al.*, 2014) under higher LMA owing to an

increase in leaf thickness. The influence of LMA on photosynthesis depends in part on

how LMA impacts mesophyll conductance. An increase in LMA related to increased

leaf density might cause mesophyll cells to be densely packed, which will inevitably

increase mesophyll CO_2 resistance (Flexas *et al.*, 2008; Niinemets *et al.*, 2009;

Weraduwage *et al.*, 2016). However, if the mesophyll cell space is large enough to

avoid excessive cell-to cell contact, it is possible to increase mesophyll and

chloroplast surface area exposed to intercellular airspace per unit leaf area and

improve mesophyll conductance (Peguero-Pina *et al.*, 2017). Recently, John *et al.*

(2017) wrote that LMA is determined by leaf cell numbers, dimensions and mass

densities. Optimizing LMA through regulating mesophyll cell morphological features

thus might be a good target to enhance leaf photosynthetic capacity.

Early in the 1960s and 1970s there were many attempts to correlate leaf anatomical

characteristics with photosynthesis. Mesophyll cell size was found to be an important

factor (Jellings and Leech, 1984). Wilson and Cooper (1970) reported that selection

for a smaller cell size could be used to improve photosynthesis and yield in ryegrass

(*Lolium perenne*). But at that time, it was not entirely known how mesophyll cell size

impacts leaf photosynthesis rate; possible factors include cell volume, cell surface

area, or the effect of cell shape on cellular packing within the leaf (Jellings and Leech,

1984). In the last two-decades, extensive investigations of leaf anatomical traits and

photosynthesis among different species or same species under various environmental

conditions have reemerged (Kogami *et al.*, 2001; Hanba *et al.*, 2002; Oguchi *et al.*,

2005; Peguero-Pina *et al.*, 2012, 2016; Sáez *et al.*, 2017). Although the typical

characteristics of leaves with high photosynthetic efficiency among different species

or environmental factors are not always consistent, a higher proportion of chloroplast

surface area exposed to intercellular airspace per unit leaf area (S_c) is generally

associated with high rates of photosynthesis. High S_c reduces the CO_2 diffusion

resistance in the leaf and improves CO_2 concentration at the sites of carboxylation.

Higher S_c values associated with more and smaller mesophyll cells could lead to

selection for smaller cells. Achieving higher S_c values through manipulating

mesophyll cell morphological traits might be a crucial spotlight of engineering leaf

anatomy to enhance leaf photosynthetic efficiency.

Furthermore, increasing explorations of the key target genes regulating cell

proliferation and expansion provides more possibilities for engineering leaf

architecture to improve photosynthetic efficiency. Takai *et al.* (2013) revealed the

gene *GREEN FOR PHOTOSYNTHESIS* (*GPS*) increases mesophyll cell number

between vascular bundles, which leads to enhanced photosynthesis rates. Lehmeier *et*

al. (2017) manipulated mesophyll cell density through promoting or repressing

KIP-RELATED PROTEIN1 (*KRP1*) and *RETINOBLASTOMA RELATED PROTEIN 1*

(*RBR1*) gene to increase photosynthetic capacity. Our recent work also found that the

leaves of *ROOT HAIR DEFECTIVE 3* (*RHD3*) defective mutants have more and

smaller mesophyll cells, and their photosynthetic rate was much higher than wild type.

All these indicate that regulation of mesophyll cell division, expansion and

arrangement in leaves is a potential target of engineering leaf anatomy by genetic

approaches to enhance photosynthetic efficiency. Therefore, this manuscript reviews

the effects of the dominant leaf anatomical traits on leaf photosynthesis, analyzes the

possibility of improving S_c through re-arranging mesophyll cells, and further

discusses the potential influences of modifying leaf anatomy on CO_2 diffusion, light

utilization and other physiological processes. Finally, some potential target genes are

explored.

2. Enhancing leaf photosynthetic capacity by altering leaf anatomy

2.1 Effects of leaf anatomical characteristics on photosynthesis

A typical C_3 leaf cross-section, comprising two epidermal layers surrounding a tight

array of columnar cells (palisades) and irregularly shaped cells (spongy), provides

powerful tool to explore various morphological features associated with leaf

physiological functions. Here we summarize 43 published research papers and

extracted the major leaf anatomical and photosynthetic parameters (Table S1),

including leaf thickness (T_L), mesophyll layer thickness (T_{mes}), fraction of intercellular

airspace (f_{ias}), *LMA*, mesophyll cell wall thickness (T_{cw}), chloroplast length (L_{chl}),

chloroplast thickness (T_{chl}), mesophyll surface area exposed to intercellular airspace

(S_{mes}), S_c, stomatal conductance (g_s), mesophyll conductance (g_m) and photosynthesis

rate (A_n) (Fig. 1). There are huge variations in leaf morphological characteristics of

different species in different environments, and the coefficient of variance (c.v.)

values vary from 22% to 79%. The smallest coefficient of variation is the length of

the chloroplast. The average chloroplast length is 4.81 μm, varying from 1.58 μm to

6.86 μm. The coefficient of variation of LMA is highest, averaging 76 g m^{-2} and

ranging from 11 g m^{-2} to 285 g m^{-2}.

162   *(Fig. 1)*

There is no significant correlation between *LMA* and A_n (Table 1), as reported in

previous studies (Hassiotou *et al.*, 2010; Veromann-Jürgenson *et al.*, 2017). Moreover,

the long distance between these two parameters indicates wide variability between

LMA and A_n (Fig. 2). As decribed in many reports (Poorter *et al.*, 2009; de la Riva *et al.*,

2016; Veromann-Jürgenson *et al.*, 2017; John *et al.*, 2017), leaf thickness, mesophyll

layer thickness, mesophyll cell wall thickness and intercellular airspace are the

dominant factors influencing *LMA*. When the *LMA* value is less than 100 g m^{-2}, an

increase in *LMA* significantly promotes S_{mes} and S_c values. However, when it exceeds

100 g m^{-2}, higher *LMA* value plays only a minor role in the improvement of the S_{mes}

and S_c value. On the contrary, higher *LMA* reduces the f_{ias} and S_c/S_{mes} values, possibly

because of densely packed mesophyll cells, further decreasing g_m (Table 2).

Despite different species and environmental conditions, significantly positive

correlations between S_c and A_n and g_m are observed; S_c is the closest positive

parameter to A_n and g_m of all summarized leaf morphological parameters. From our

results, the average S_c value is 11.1 m^2 m^{-2}, ranging from 1.2 m^2 m^{-2} of *Deschampsia*

antarctica Desv in Lagotellerie island (Sáez *et al.*, 2017) to 32.3 m^2 m^{-2} of *Abies*

pinsapo needles in the Mediterranean area (Peguero-Pina *et al.*, 2012). High variations

and significant positive correlation between S_c and A_n confirm that S_c is a potential

target of engineering leaf anatomy to enhance leaf photosynthetic capacity.

182   *(Fig.2)*

183   *(Table 1)*

184   *(Table 2)*

2.2 Increase S_c by manipulating mesophyll cell morphology

S_c is the most important leaf structural parameter in photosynthetic studies during the

last decades. von Caemmerer and Evans (1991) showed that higher S_c could explain

the relatively large CO_2 transfer conductance of rice. Almost all subsequent studies

have confirmed the importance of having a high S_c value for maximizing

photosynthesis (Terashima *et al.*, 2011). The S_c value is influenced by chloroplast

number, distribution in the mesophyll and mesophyll surface area exposed to

intercellular airspace. An increase in the number of chloroplasts in mesophyll cells is

beneficial to improve S_c (Miyazawa and Terashima, 2001). Additionally, changes in

light intensity and quality can induce movement and re-distribution of chloroplasts

within cells, further affecting S_c (Hanba *et al.*, 2002; Oguchi *et al.*, 2003). The

maximum value of S_c is determined by S_{mes}, which is the closest positive parameter to

S_c (Fig. 2). In our analysis the average S_c/S_{mes} value is 0.71, implying that nearly 3/4

of mesophyll cell surface area facing intercellular airspace is occupied by chloroplast.

For *Castanopsis sieboldii*, the S_c/S_{mes} ratio could be 1.0 after 20 days of full leaf area

expansion (Hanba *et al.*, 2002). Oguchi *et al.* (2003) described that transferring the

mature leaf from low to high light significantly increased photosynthesis rate,

associated with improving S_c. The photosynthesis rate in the low-light leaf was

significantly lower than the leaf exposed to high light because of restricted S_{mes} value.

Greater S_{mes} is a prerequisite for a larger S_c. Therefore, simultaneously raising the S_{mes}

and S_c is the key to enhance leaf photosynthetic efficiency.

Nobel *et al.* (1975) first reported in that the mesophyll cell surface area per unit leaf

area (A_{mes}/A in their notation), calculated assuming that mesophyll cells were

cylindrical with hemispheres at each end, was correlated with CO_2 assimilation rate.

Evans *et al.* (1994) optimized the calculation depending on the shape of palisade and

spongy tissue. According to our literature review, the average S_{mes} value is 15.9 m^2

m^{-2}, ranging from 2.8 m^2 m^{-2} of *Arabidopsis cgr2/3* mutant leaf (Weraduwage *et al.*,

2016) to 41.6 m^2 m^{-2} of sun *Camellia japonica* leaf (Terashima *et al.*, 2006). To

facilitate CO_2 diffusion in the leaf, most of the surface area of mesophyll cells are

exposed to intercellular airspaces (Nobel, 2009). Greater total length of mesophyll

cells exposed to intercellular airspaces thus is prerequisite to obtain the greater values

of S_{mes}. Terashima *et al.* (2011) illustrated four strategies of improving mesophyll

surface area, including (1) cell elongation, (2) cell elongation accompanied by cell

division, (3) decrease in cell size, (4) armed cells of grass species having lobes. For

lobed cells or armed cells, high surface area exposure of mesophyll cells and

chloroplasts to intercellular spaces and enhanced mesophyll conductance was found in

Oryza spp and *Viburnum* spp (Sage and Sage, 2009; Chatelet *et al.*, 2013). Adachi *et al.*

(2013) and He *et al.* (2017) showed two rice lines named BTK-a and BTK-b to have

20-50% higher photosynthetic rate than the parents because of highly developed lobes

of mesophyll cells. Although there are slight differences in the strategies to improve

mesophyll surface area, the key of the other three approaches is the trade-off between

cell size and cell density because the mesophyll cell size is negatively correlated with

cell number when the leaf thickness is constrained (Pyankov *et al.*, 1999).

To understand S_{mes} easily, here some geometrical idealization is adopted. Let us

assume that all mesophyll cells in a cross-section are spherical; cells are tightly

packed and all cell walls are exposed to the intercellular airspaces (Fig. 3). Thus, S_{mes}

is equal to the total circumference of the spherical cells divided by the width of the

cross-section according to the 2D model. If there is only a single uniform sphere in

cross-section, S_{mes} is π. If cell diameter is reduced to one-third, three layers with nine

uniform spherical cells would produce a S_{mes} of 3π. Increasing cell density would

significantly improve S_{mes} when the leaf thickness remains unchanged. Indeed,

mesophyll cell density cannot be increased indefinitely. Smaller cell volume would

result in over-lapping chloroplasts in mesophyll cells because of the strong correlation

between the number of chloroplasts in a cell and the mesophyll cell area (Pyke and

Leech, 1991, 1992). Therefore, the optimum is the minimum volume of mesophyll

cell in which all the chloroplasts are tightly attached to cell wall and avoid

over-lapping of chloroplasts. We have not rigorously examined the published

photomicrographs of leaf cross-section, however, the number of chloroplasts in

individual mesophyll cells seen in 2D cross-section is usually about 10-30. Compared

with other leaf anatomical features, the variation in length and thickness of

chloroplasts is smaller, with an average of 4.81 μm and 2.46 μm, respectively. The

typical spherical mesophyll cell area containing 10-30 chloroplasts occupying all

mesophyll surface area facing intercellular airspace is between 241-1780 μm^2. If the

mesophyll surface area is less than that value, chloroplasts might overlap in the

mesophyll cell, which would result in the decrease of S_c. In addition, adequate

mesophyll cell space is a prerequisite to achieve an increase in cell density. From Fig.

3, it could be found that f_{ias} is still 0.215, despite many more mesophyll cells. From

the literature, the average f_{ias} is 0.241 regardless of species and environment, varying

from 0.059 of *Oryza sativa* cslf6-2 mutant (Ellsworth *et al.*, 2018) to 0.675 of

Asplenium scolopendrium (Carriquí *et al.*, 2015). Great variations in f_{ias} produce the

possibility to utilize internal volume of the leaf to assemble more mesophyll cells to

improve S_{mes} and S_c. Authentic mesophyll cells are not uniformly spherical and are

more complex, nevertheless, for most species the intercellular space is sufficient to be

equipped with more mesophyll cells. Appropriate manipulation of mesophyll cell

density to increase mesophyll and chloroplast surface area exposed to intercellular air

space is a potential approach for enhancing leaf photosynthetic capacity by altering

leaf anatomy.

262   3. Potential changes of photosynthetic capacity by manipulating
263   mesophyll cell morphology

264   3.1 The mechanism of enhancing photosynthesis

265 CO_2 diffusion

After arriving at substomatal cavities, atmospheric CO_2 diffuses through intercellular

airspaces and then passes through the cell wall, cytosol, chloroplast envelope, finally

reaching the sites of carboxylation in the chloroplast. The total CO_2 diffusion

resistance between substomotal cavities and carboxylation sites is termed mesophyll

resistance and its reciprocal is defined as mesophyll conductance (g_m), which is

sufficiently small to restrict photosynthesis (Warren, 2008; Flexas *et al.*, 2008, 2012;

Evans *et al.*, 2009). Mesophyll resistance can be divided into the gas-phase and

liquid-phase resistance, in which liquid-phase resistance could account more than 90%

of mesophyll resistance (Parkhurst, 1994; Tosens *et al.*, 2012; Lu *et al.*, 2016). In the

liquid-phase, many CO_2 diffusion parameters have not been accurately quantified

(Evans *et al.*, 2009; Xiao and Zhu, 2017), however, cell wall thickness and S_c are

considered the most important anatomical parameters influencing g_m (Syvertsen *et al.*,

1995; Tomás *et al.*, 2013). Summarized data confirmed the significantly negative

correlation between g_m and cell wall thickness (Fig. 2). By contrast, g_m is positively

correlated with S_c because a large chloroplast surface area adjacent to the cell wall is

beneficial for fast diffusion through the cytosol and minimizes the total diffusion

resistance. He *et al.* (2017) reported an increase in the number of lobed mesophyll

cells induced much higher total surface area of mesophyll cells in the BTK lines,

further resulting in greater mesophyll conductance. As shown in Fig.3, without

increasing leaf thickness, increasing mesophyll cell number improves the mesophyll

surface area, thereby increasing the number of chloroplasts per unit leaf area and S_c.

Higher S_c improves CO_2 mesophyll conductance, increases CO_2 concentration at

carboxylation site and enhances photosynthetic capacity.

Light interception and distribution in leaves

Leaf anatomy is well designed to capture solar and maximize light absorption; leaves

often absorb more than 85% of the photosynthetically active radiation that hits them

(Evans and Poorter, 2001). Lens-shaped epidermal cells can concentrate incident light

severalfold, columnar palisade cells facilitate the penetration of light into the leaf, and

randomly arranged spherical spongy cells and intercellular airspaces intensely scatter

and reflect light, which increases the probability of light absorption for photosynthesis

(Vogelmann *et al.*, 1996*a,b*; Ustin *et al.*, 2001; Smith *et al.*, 2004). With depth into a

leaf, the light and chlorophyll fluorescence intensity, especially for red and blue light,

decline remarkably due to absorption and scattering, depending upon pigmentation as

well as the mesophyll structure (Vogelmann, 1993; Vogelmann and Evans, 2002).

Cui *et al.* (1991) showed more and elongate palisade cells within sun leaves facilitated

the light transmission in leaves and spread the light through the depth of the leaf while

less compact palisade cells within shade leaves helps increase light absorption

through greater scattering. Thus, altering leaf anatomy through regulating mesophyll

cell geometry and packing would greatly influence the distribution of internal light.

More and slightly-stacked mesophyll cell layers could efficiently facilitate the

penetration of light into the leaf reducing light absorption in the top most layer that

might be saturated by light allowing light to penetrate to the lower layers that may not

be saturated by light. On the other hand, smaller intercellular airspace with small

pockets could scatter light more (Smith *et al.*, 2004), increase the light pathlength and

enhance light absorption. Besides reducing CO_2 diffusion resistance, higher S_c

accompanied by more chloroplasts exposed to intercellular airspace aids efficient light

absorbtion and helps satisfy the electron requirement for CO_2 assimilation associated

with high CO_2 conductance (Terashima *et al.*, 2009). The transgenic line *ATML1$_{pro}$*:

KRP1 leaves exhibited an increase in palisade cell density improving air channel

circularity and density, promoting the proportion of light energy used and operational

election transport rate (Lehmeier *et al.*, 2017).

3.2 Other potential changes

Leaf growth is determined by the amount of photosynthetic C partitioned between leaf

area growth and *LMA* (Weraduwage *et al.*, 2015). An alteration in C investment to leaf

area or *LMA* could influence plant growth. Weraduwage *et al.* (2016) showed that

overexpression of the *Cotton Golgi-related* (*CGR*) genes *CGR2* and *CGR3,* which

mediate pectin methylesterification, increased C partitioned to leaf area growth and

improved whole-plant photosynthesis and plant growth. Greater numbers of

mesophyll cells would result in higher *LMA* (Pyankov *et al.*, 1999; Poorter *et al.*, 2009;

John *et al.*, 2017), which requires more C invested in *LMA* and reduces the available C

for leaf area growth. Additionally, mesophyll cells make more than 90% of total leaf

respiration owing to more mitochondria in mesophyll cells (Long *et al.*, 2015*a*). More

mesophyll cells mean more mitochondria and higher leaf respiration, reducing

photosynthetic C available for plant growth (Weraduwage *et al.*, 2016). Manipulating

mesophyll cell density could be used to enhance photosynthetic capacity, nonetheless,

it might change the allocation of photosynthetic C, further affecting plant growth.

Previous studies have described strong correlations between leaf hydraulic

conductance and CO_2 diffusion and photosynthesis rate (Flexas *et al.*, 2013; Scoffoni

et al., 2016; Barbour, 2017). Both leaf water transport and CO_2 delivery have

similarities in leaf anatomical traits. The leaf hydraulic system is composed of the

major and minor veins, bundle sheaths, and the mesophyll cells outside the vein

xylem. Mesophyll cell characteristics, such as mesophyll thickness, mesophyll cell

porosity and connectivity, mesophyll cell surface area for evaporation, which

significantly affect CO_2 diffusion in leaf, are key parameters determining hydraulic

conductance (Sack *et al.*, 2015). Greater porosity of mesophyll cells contributes more

to a high leaf hydraulic conductance, and broken connections between cells and fewer

pathways of water flow would cause a lower aquaporin activity (Scoffoni *et al.*, 2014).

Small leaves with smaller cells tend to have greater minor vein length per unit leaf

area, conferring high leaf hydraulic conductance (Scoffoni *et al.*, 2011). Changes in

gas exchange traits associated with altering mesophyll cell morphology would

influence leaf hydraulic conductance and transpiration. Theoretically, increasing

mesophyll cell density would not affect intercellular airspace (Fig. 3), nevertheless, an

increase in mesophyll cell density inescapably decreases intercellular air space

(Lehmeier *et al.*, 2017). Hanba *et al.* (2004) documented that the leaves of transgenic

rice Tr6322-H had higher mesophyll cell density and more developed collenchyma

coupled to a higher mesophyll conductance, however, its leaf development might be

influenced by water deficit because of increased water loss. The regulation of

mesophyll cell morphology has potential to enhance photosynthetic capacity, however,

it is worth studying whether it will increase hydraulic conductance simultaneously or

change the composition of hydraulic conductivity.

Terashima *et al.* (2006) have discussed the importance of having sufficient S_c by

decreasing cell size for efficient photosynthesis and compared the merits and demerits

of different mesophyll cell sizes. Different species can have considerably distinct

mesophyll cell sizes as a result of natural selection in various environments. Annual

herbaceous plants often have large mesophyll cells, while ever-green trees tend to

have smaller mesophyll cells. The leaves with small mesophyll cells have strong

mechanical strength, long longevity, small heat capacity, and lower expansion rate.

Additionally, they exhibit higher information cost (nuclear N- and P and chloroplast N

and P). From this perspective, plants in natural environments may not benefit from

very small mesophyll cells.

However, for agroecosystems, there might be more advantages to select small

mesophyll cells. High mechanical strength would be beneficial to enhance crop pest

resistance. The maintenance of leaf morphology depends on the cell wall, which is

helpful to increase resistance to drought stress. For intensive agricultural production,

N and P inputs are always sufficient or even excessive. If manipulating mesophyll cell

morphology could improve photosynthetic capacity and increase crop yield without

additional inputs, then N and P utilization efficiency would be higher. Selecting for

smaller mesophyll cells does not mean completely changing the shape and size of the

mesophyll cells. Instead, depending on the current size and arrangement of mesophyll

cells, it may be possible to reduce mesophyll cell size appropriately by altering cell

proliferation and expansion to improve S_c and reduce CO_2 diffusion resistance,

thereby enhancing photosynthetic efficiency. Indeed, comprehensive studies

estimating the potential for regulating mesophyll cell morphology and the related

changes on physiological and biochemical should be made.

4. Target genes could be used to regulate mesophyll cell morphology

Leaf growth has been divided into three phases: (1) leaf primordia occur at the shoot

apical meristem, (2) cells proliferate and form specific structures during primary

morphogenesis, (3) leaf cells begin to expand at secondary morphogenesis (Donnelly

et al., 1999; Andriankaja *et al.*, 2012). In the transition from primary to secondary

morphogenesis, a retrograde chloroplast signal induces the onset of cell expansion and

the leaf becomes photosyntheticaly active at the same time (Andriankaja *et al.*, 2012),

indicating that leaf growth as well as leaf morphology is inseparably correlated with

photosynthesis. Although leaf morphology is regulated in response to different growth

stages and environments, genetic control is fundamental to produce various leaves in

nature. Thus, leaf morphogenesis and the related genes are hotspots in plant biology

research. Numerous Arabidopsis mutants with altered leaf size and/or shape have been

isolated (Horiguchi *et al.*, 2006) and a key aspect of genes affecting leaf

morphogenesis also have been identified during the past years. Several articles have

reviewed in detail the critical genes regulating leaf shape, size, cell proliferation and

expansion (Beemster *et al.*, 2003; Micol, 2009; Horiguchi and Tsukaya, 2011;

Gonzalez *et al.*, 2012; Rodriguez *et al.*, 2014, 2016; Hepworth and Lenhard, 2014).

However, leaf development is a complex biological process and is determined by

many genes and pathways. The molecular mechanisms underlying leaf size and shape

are only starting to be unraveled. Although extensive research is still needed, key

genes can provide some mechanisms for regulating leaf morphology.

Cell number, which plays a more important role in determining the final size of plant

organs rather than cell size (Gázquez and Beemster, 2017), mainly depends on cell

proliferation. Based on the existing research results, we divide the principal genetic

factors regulating cell proliferation into three groups:

(1) **Direct regulators**. Overexpression or loss of these genes can directly regulate cell

proliferation. such as GROWTH-REGULATING FACTORs (*GRF*) family, which has

been widely studied (Omidbakhshfard *et al.*, 2015; Tsukaya, 2016). There are nine

members in the *Arabidopsis thaliana* GRF family (*AtGRF*), of which AtGRF1, 2, 3, 4,

5, and 9 are involved in the regulation of cell proliferation. However, while other

family members have partly overlapping roles, the function of *AtGRF5* is to regulate

cell proliferation. When *AtGRF5* and *AN3* (the transcription coactivator) were

overexpressed, plants developed larger leaves as a result of increasing cell number

rather than cell size (Horiguchi *et al.*, 2005). For AtGRFs, expression is regulated by

miRNA miR396 and ectopic expression of miR396 repressed the AtGRF activity and

resulted in smaller leaves with reduced cell number (Rodriguez *et al.*, 2010).

Additionally, overexpressing other genes, such as *AINTEGUMENTA* (*ANT*)

(Mizukami and Fischer, 2000), *Arabidopsis SKP1-LIKE1* (*ASK1*) (Zhao *et al.*, 1999),

Arabidopsis thaliana NGATHA (*AtNGA*) (Lee *et al.*, 2015), *G-PATCH DOMAIN*

PROTEIN1 (*gdp1*) (Kojima *et al.*, 2018), *KLU* (Anastasiou *et al.*, 2007),

OLIGOCELLULA1, 4, 6 (*oli1, 4, 6*) (Fujikura *et al.*, 2009), *POINTED FIRST LEAD 2*

(*PFL2*) (Ito *et al.*, 2000), *ROTUNDIFOLIA4* (*ROT4*) (Narita *et al.*, 2004),

STRUWWELPETER (SWP) (Autran *et al.*, 2002) and *TCP4* (Schommer *et al.*, 2014)

could prolong cell proliferation and produce large leaves with more cells.

(2) **Genes related to hormone signaling pathways**. Numerous hormones, such as

auxins, brassinosteroids, cytokinin, gibberellins, jasmonic acid, etc, could act as

signals for cell proliferation and expansion at the cellular level. The genes regulating

hormone signals influence cell proliferation and expansion. For example, ectopic

expression of *ARGOS* transduces auxin signals downstream of *AXR1*, which prolongs

ANT expression and regulates cell proliferation, displaying enlarged leaves attributed

to increase in cell number (Hu *et al.*, 2003). The *cro* mutants *det2* and *dwf1*, affected

in brassinosteroid biosynthesis revealed fewer and smaller cells (Nakaya *et al.*, 2002).

AHK-cytokinin receptor in *Arabidopsis thaliana* (Nishimura *et al.*, 2004),

PHYB-involved in gibberellin biosynthesis (Chaiwanon *et al.*, 2016), and *aos* (*allene*

oxide synthase) and *coil-16B* (*coronatine insensitive1*)-jasmonate receptor, could

influence cell division and expansion by regulating related hormone signals.

(3) **Genes related to pathways synthesizing structural component**. Plant cells are

encircled by a rigid cell wall and the biosynthesis and modification of cell wall would

significantly influence cell expansion and proliferation. Cellulose, hemicellulose, and

pectins are the dominant components of the plant cell (Ochoa-Villarreal *et al.*, 2012;

Tenhaken, 2015) and the genes code for enzymes catalyzing cellulose, hemicellulose,

and pectin synthesis could regulate cell expansion and proliferation. For example,

overexpression of the *fras5*, a mutant allele of *AtCesA7* gene involved in catalyzing

cellulose synthesis, reduced cellulose content, cell wall thickness, and cell elongation

(Zhong *et al.*, 2003). The *ccr1* mutant has high ferulic acid content, an intermediate in

lignin biosynthesis, and increased cell proliferation (Xue *et al.*, 2015). A knockout of

CGR2 and *CGR3* genes showed reduced cell expansion and overall plant growth

associated with reducing pectin methylesterification (Kim *et al.*, 2015).

The directly regulated genes, which might be involved in signaling, structural, and

other aspects regulating cell and leaf shape and still require further research, mainly

regulate cell proliferation. In most instances, overexpressing these genes results in

larger leaves owing to increased cell number rather than cell size. However, genes

related to hormone signaling and structural component synthesis always influence

both cell proliferation and cell expansion. Generally, an increase in the number of

cells leads to a reduction in the volume of the cells, which is called "compensatory

effect". Horiguchi and Tsukaya (2011) defined compensation as when "a decrease in

cell number in a leaf caused by genetic defect leads to an enhanced cell expansion".

They thought altered cell proliferation triggered compensation rather than cell

expansion. Moreover, compensation is induced only when cell proliferation is beyond

a threshold level (Horiguchi *et al.*, 2006). It indicates increasing cell number within a

certain range would not cause the changes in cell size, which provides the theoretical

support of regulating mesophyll cell density to improve the S_{mes} and S_c values.

Besides manipulating mesophyll cell density, improving chloroplast density is helpful

to increase the S_c values. Thus, the genes regulating chloroplast division will also be

reviewed. The *ARC* (ACCUMULATION and REPLICATION of CHLOROPLASTS)

mutants are an important genetic resource in the control of chloroplast division (Pyke

and Leech, 1994; Gao *et al.*, 2003; Cho *et al.*, 2012; Zhou *et al.*, 2015). Gao *et al.* (2003)

revealed that mutatants of *ARC5* exhibit defects in chloroplast constriction and have

enlarged dumbbell shaped chloroplasts. *GC1* (GIANT CHLOROPLAST) is another

gene involved chloroplast division, which plays a positive role at an early stage of the

division process, and *GC1* deficiency resulted in mesophyll cells harboring one to two

giant chloroplasts (Maple *et al.*, 2004). Hymus *et al*. (2013) showed overexpression of

ATHB17 increases chloroplast numbers per unit cell size due to an increase in the

number of proplastids per meristematic cell. As one of the critical genes regulating

cell proliferation, *AtGRF5* stimulates chloroplast division, resulting in a high

chloroplast number per cell and a higher photosynthesis rate (Vercruyssen *et al.*,

2015). Additionally, cytokinin promotes chloroplast ultrastructure and chlorophyll

synthesis. Cortleven and Schmulling (2015) summarized cytokinin signaling

pathways of Arabidopsis regulating chloroplast development and function through

transcription factors. As the cytokinin receptors, *AHK2* and *AHK3* genes not only

trigger cell divisions (Nishimura *et al.*, 2004), but also regulate chloroplast

development and chloroplast division. Recently, Larkin *et al.* (2016) described how

the *REC* gene helps to establish the proportion of cellular volume devoted to

chloroplasts; the *rec1* mutant has reduced size of the chloroplast compartment.

Altered chloroplast coverage in cells might induce changes in S_c and g_m.

5. Prospects to manipulate mesophyll cell morphology for enhancing photosynthesis

Regulation is disconnected from limitation (Sharkey and Weise, 2012). Light and CO_2

are the primary external constraints (limitations) on photosynthesis, therefore,

regulating leaf morphology to take maximum advantage of light and CO_2 is key to

maximize photosynthesis. During the past decades, considerable research has been

conducted to study the influence of leaf morphology on photosynthetic capacity in

different species and environments. These studies help interpret how plants achieve

high photosynthetic efficiency by optimizing leaf anatomy under different

environments. Nonetheless, for any given species, the leaf architectural traits with the

highest photosynthetic capacity is not always clear. Small differences in leaf

phenotypes and the large number of genes that affect leaf architecture make research

in this area difficult. As the most widely used species in plant biology research,

Arabidopsis thaliana has been screened and numerous mutants with different leaf size

and/or shape have been isolated (Horiguchi *et al.*, 2006; Micol, 2009). A large number

of the mutated genes determining cell proliferation, expansion and differentiation

have been identified. However, these mutations have not typically been studied in the

context of the effects on photosynthesis. Therefore, the best targets for engineering

leaf anatomy are unknown. More, and in-depth, studies should focus on (1) the

differences in photosynthetic capacity under different leaf shape and size as well as

leaf anatomy, (2) how to influence leaf photosynthesis through the potential target

genes altering leaf morphology. Such research will pinpoint the targets for altering

leaf architecture to maximize photosynthetic capacity.

Although light fluctuations significantly influence leaf photosynthesis (Kaiser *et al.*,

2018; Slattery *et al.*, 2018), CO_2 concentration is the dominant factor restricting

photosynthesis and yield in crops under current conditions (Ainsworth and Long, 2005;

Ellsworth *et al.*, 2012). In C_4 plants, the specialized Kranz anatomy helps concentrate

CO_2 concentration and enhance photosynthetic efficiency. C_3 plants do not possess

Kranz anatomy and CO_2 diffusion resistance in mesophyll cells significantly reduces

the CO_2 concentration at the sites of carboxylation. If mesophyll resistance could be

eliminated, photosynthesis rates would be improved by up to 20% (Zhu *et al.*, 2010).

Increasing chloroplast area exposed to intercellular airspace as much as possible could

reduce the CO_2 diffusion resistance and enhance mesophyll conductance and

photosynthetic capacity. Terashima *et al.* (2001, 2006, 2011) have elaborated the

importance of having sufficient S_c for efficient photosynthesis and have discussed the

various strategies to increase the S_{mes} and S_c values by manipulating cell proliferation

and expansion. They thought decreasing mesophyll cell size was an effective

approach to increase S_c and decrease CO_2 diffusion resistance. Nevertheless, most

plants do not tend to form small mesophyll cells as a result of natural selection. As

early as the 1970s, selection for a smaller cell size have been considered as critical

parameters to breed new variety with high photosynthesis and yield. Nevertheless,

owing to poor understanding of molecular mechanisms controlling leaf anatomy as

well as cell proliferation and expansion, the investigations into the influence of

mesophyll cell size on photosynthesis have not been fully developed and the selection

of more and smaller mesophyll cells is not currently considered a potential approach

to enhance photosynthesis capacity. Recent research shows increasing mesophyll cell

number through regulating genes related to the cell cycle are beneficial for improving

photosynthetic capacity (Takai *et al.*, 2013; Lehmeier *et al.*, 2017). It can be foreseen

that engineering leaf anatomy by manipulating cell proliferation and expansion using

gene regulation or gene editing technology will be a potential approach to enhance

plant photosynthetic capacity in future.

Indeed, many challenges, including photosynthesis and molecular mechanisms

controlling leaf anatomy, need to be solved. What is the potential for regulating cell

density to maximize photosynthesis? As mentioned in the compensation effect,

increasing in mesophyll cell number beyond a threshold value will trigger a decrease

in cell size (Horiguchi *et al.*, 2006). Smaller mesophyll cell size will restrict the

number and division of chloroplasts. Although the S_{mes} value is improved by

increasing mesophyll cell number, the S_c value might not be increased because of

over-lapping chloroplasts, which is unfavorable for enhancing photosynthesis. It is

therefore essential to explore the relationship between mesophyll cell traits and

mesophyll conductance and photosynthetic capacity. Additionally, an increase in leaf

mesophyll cell density would improve leaf mass density and *LMA*, which could

modify photosynthetic C partitioning to leaf area growth and growth in terms of *LMA*,

further influencing plant growth. Weraduwage *et al*. (2016) revealed that in the *cgr2/3*

mutant greater C partitioning to *LMA*, associated with an increase in leaf cell density,

resulted in smaller leaf area and reduced whole-plant photosynthesis. A great number

of cells and organelles require more energy and hence incur higher maintenance

respiratory costs (Weraduwage *et al.*, 2016). What's more, alteration in mesophyll cell

number and size also would impact the interception of light, water transport and CO_2

assimilation. Comprehensive studies to analyze the effects of manipulating mesophyll

cell morphology on other physiological processes in leaves would be useful to

optimize the approach for enhancing photosynthetic capacity.

The selection of target genes for altering leaf anatomy also requires more accurate and

extensive research. As mentioned in Lehmeier *et al.* (2017), the suppression of *RBR1*

removes some restrictions on cell division, leading to more and smaller mesophyll

cells; however, the transgenic lines with higher photosynthetic capacity grew smaller

than the wild type because of higher *LMA*. Although photosynthetic capacity is

improved by regulating these genes, such plants do not achieve higher plant biomass

and crop yield as result of the decrease of photosynthetic area. The genes coordinating

the improvement of photosynthetic capacity and plant growth can meet the demand of

modern agriculture for high yield.

Altering leaf anatomy to enhance leaf photosynthetic capacity must be achieved

without influencing leaf area. From this perspective, the genes directly regulating cell

proliferation might be considered first. For example, overexpression *AtGRF5* and *AN3*

in *Arabidopsis thaliana* results in larger leaves with an increased cell number

(Horiguchi *et al.*, 2005). Additionally, the genes controlling cell proliferation and

expansion might change other physiological processes. For example, the *cgr2/3*

mutant, double knockout of *CGR2* and *CGR3* genes, showed a reduced cell expansion

and increased mesophyll cell density as result of a reduction in the degree of pectin

methylation; however, it induced hardening of cell walls, and increased CO_2

resistance of the cell wall and reduced photosynthetic efficiency (Weraduwage *et al.*,

2016). Altering expression of ANT genes modified cell proliferation and regulated the

number of higher order veins and minor tertiary veins (Kang *et al.*, 2007), which are

closely related to leaf photosynthetic capacity (Sack *et al.*, 2013; Feldman *et al.*, 2017).

It is therefore essential to understand the various functions of the target genes for

accurate editing the genes to alter leaf morphology and enhance photosynthetic

efficiency.

582 6. Conclusion

Natural selection allows for a range of leaf architectures to effectively capture light

and exchange gases, resulting in high photosynthetic efficiency to grow and survive.

Despite the varitety of species and environmental conditions, S_c is the most important

indicator of leaf photosynthetic capacity. Having a sufficiently high S_c is a

prerequisite of efficient photosynthesis. Chloroplast number and distribution in

mesophyll cells and S_{mes} significantly influence S_c. Without overlapping mesophyll

cells, increasing mesophyll cell density is an important strategy to improve S_{mes} and S_c.

Depending on the manipulation of target genes involved in cell proliferation and

expansion, such as *AtGRF* family, it should be possible to produce more mesophyll

cells without influencing leaf area, further inducing higher S_c values and

photosynthetic capacity. It is foreseen that the chain "gene regulation-leaf

anatomy-photosynthetic capacity" will be an important strategy for enhancing

photosynthetic capacity and crop yield. Indeed, there are many challenges that need to

be thoroughly studied, such as the selection of target gene regulating cell proliferation

and expansion, the potential of adjusting mesophyll cell number and size, and the

effects of leaf anatomical traits on other physiological processes, including light

utilization, water transport etc.

Acknowledgements

This research was funded by U.S. Department of Energy Grant DE-FG02-91ER2002.

Partial salary support for T.D.S. came from Michigan AgBioResearch. T.R

acknowledges the fellowship supported by China Scholarship Council.

References

Adachi S, Nakae T, Uchida M, *et al.* 2013. The mesophyll anatomy enhancing CO_2 diffusion is a key trait for improving rice photosynthesis. Journal of Experimental Botany **64**, 1061–1072.

Ainsworth EA, Long SP. 2005. What have we learned from 15 years of free-air CO_2 enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO_2. New Phytologist **165**, 351–372.

Anastasiou E, Kenz S, Gerstung M, MacLean D, Timmer J, Fleck C, Lenhard M. 2007. Control of plant organ size by *KLUH/CYP78A5*-dependent intercellular

signaling. Developmental Cell **13**, 843–856.

Andriankaja M, Dhondt S, De Bodt S, *et al.* 2012. Exit from proliferation during leaf development in *Arabidopsis thaliana*: A not-so-gradual process. Developmental Cell **22**, 64–78.

Autran D, Jonak C, Belcram K, Beemster GTS, Kronenberger J, Grandjean O, Inzé D, Traas J. 2002. Cell numbers and leaf development in Arabidopsis: a functional analysis of the *STRUWWELPETER* gene. The EMBO Journal **21**, 6036–6049.

Barbour MM. 2017. Understanding regulation of leaf internal carbon and water transport using online stable isotope techniques. New Phytologist **213**, 83–88.

Beemster GTS, Fiorani F, Inzé D. 2003. Cell cycle: the key to plant growth control? Trends in Plant Science **8**, 154–158.

Cai Y-F, Li S-F, Li S-F, Xie W-J, Song J. 2014. How do leaf anatomies and photosynthesis of three *Rhododendron* species relate to their natural environments? Botanical Studies **55**, 36.

Carriquí M, Cabrera HM, Conesa M À., *et al.* 2015. Diffusional limitations explain the lower photosynthetic capacity of ferns as compared with angiosperms in a common garden study. Plant, Cell & Environment **38**, 448–460.

Chaiwanon J, Wang W, Zhu J-Y, Oh E, Wang Z-Y. 2016. Information integration and communication in plant growth regulation. Cell **164**, 1257–1268.

Chatelet DS, Clement WL, Sack L, Donoghue MJ, Edwards EJ. 2013. The Evolution of photosynthetic anatomy in *Viburnum* (Adoxaceae). International Journal of Plant Sciences **174**, 1277–1291.

Cho Y-H, Kim G-D, Yoo S-D. 2012. Giant chloroplast development in ethylene response1-1 is caused by a second mutation in *ACCUMULATION AND REPLICATION OF CHLOROPLAST3* in *Arabidopsis*. Molecules and Cells **33**, 99–103.

Cortleven A, Schmülling T. 2015. Regulation of chloroplast development and function by cytokinin. Journal of Experimental Botany **66**, 4999–5013.

Cui M, Vogelmann TC, Smith WK. 1991. Chlorophyll and light gradients in sun and shade leaves of *Spinacia oleracea*. Plant, Cell & Environment **14**, 493–500.

Donnelly PM, Bonetta D, Tsukaya H, Dengler RE, Dengler NG. 1999. Cell cycling and cell enlargement in developing leaves of *Arabidopsis*. Developmental Biology **215**, 407–419.

Ellsworth PV, Ellsworth PZ, Koteyeva NK, Cousins AB. 2018. Cell wall properties in *Oryza sativa* influence mesophyll CO_2 conductance. New Phytologist **219**, 66-76.

Ellsworth DS, Thomas R, Crous KY, Palmroth S, Ward E, Maier C, DeLucia E, Oren R. 2012. Elevated CO_2 affects photosynthetic responses in canopy pine and subcanopy deciduous trees over 10 years: a synthesis from Duke FACE. Global Change Biology **18**, 223–242.

Evans JR, Caemmerer S von, Setchell BA, Hudson GS. 1994. The relationship between CO_2 transfer conductance and leaf anatomy in transgenic tobacco with a reduced content of Rubisco. Australian Journal of Plant Physiology **21**, 475–95.

Evans JR, Kaldenhoff R, Genty B, Terashima I. 2009. Resistances along the CO_2 diffusion pathway inside leaves. Journal of Experimental Botany **60**, 2235–2248.

Evans JR, Poorter H. 2001. Photosynthetic acclimation of plants to growth irradiance: the relative importance of specific leaf area and nitrogen partitioning in maximizing carbon gain. Plant, Cell & Environment **24**, 755–767.

Feldman AB, Leung H, Baraoidan M, Elmido-Mabilangan A, Canicosa I, Quick WP, Sheehy J, Murchie EH. 2017. Increasing leaf vein density via mutagenesis in rice results in an enhanced rate of photosynthesis, smaller cell sizes and can reduce interveinal mesophyll cell number. Frontiers in Plant Science **8**, 1883.

Flexas J, Barbour MM, Brendel O, *et al.* 2012. Mesophyll diffusion conductance to CO_2: An unappreciated central player in photosynthesis. Plant Science **193**, 70–84.

Flexas J, Diaz-Espejo A, Gago J, Gallé A, Galmés J, Gulías J, Medrano H. 2014. Photosynthetic limitations in Mediterranean plants: A review. Environmental and Experimental Botany **103**, 12–23.

Flexas J, Ribas-Carbó M, Diaz-Espejo A, Galmés J, Medrano H. 2008. Mesophyll conductance to CO_2 : current knowledge and future prospects. Plant, Cell & Environment **31**, 602–621.

Flexas J, Scoffoni C, Gago J, Sack L. 2013. Leaf mesophyll conductance and leaf hydraulic conductance: an introduction to their measurement and coordination. Journal of Experimental Botany **64**, 3965–3981.

Fujikura U, Horiguchi G, Ponce MR, Micol JL, Tsukaya H. 2009. Coordination of cell proliferation and cell expansion mediated by ribosome-related processes in the leaves of *Arabidopsis thaliana*. The Plant Journal **59**, 499–508.

Gao H, Kadirjan-Kalbach D, Froehlich JE, Osteryoung KW. 2003. ARC5, a cytosolic dynamin-like protein from plants, is part of the chloroplast division machinery. Proceedings of the National Academy of Sciences **100**, 4328–4333.

Gázquez A, Beemster GTS. 2017. What determines organ size differences between species? A meta-analysis of the cellular basis. New Phytologist **215**, 299–308.

Gonzalez N, Vanhaeren H, Inzé D. 2012. Leaf size control: complex coordination of cell division and expansion. Trends in Plant Science **17**, 332–340.

Hanba YT, Kogami H, Terashima I. 2002. The effect of growth irradiance on leaf anatomy and photosynthesis in *Acer* species differing in light demand. Plant, Cell & Environment **25**, 1021–1030.

Hanba YT, Shibasaka M, Hayashi Y, Hayakawa T, Kasamo K, Terashima I, Katsuhara M. 2004. Overexpression of the barley aquaporin HvPIP2;1 increases internal CO_2 conductance and CO_2 assimilation in the leaves of transgenic rice plants. Plant and Cell Physiology **45**, 521–529.

Hassiotou F, Renton M, Ludwig M, Evans JR, Veneklaas EJ. 2010. Photosynthesis at an extreme end of the leaf trait spectrum: how does it relate to high leaf dry mass per area and associated structural parameters? Journal of Experimental Botany **61**, 3015–3028.

He W, Adachi S, Sage RF, Ookawa T, Hirasawa T. 2017. Leaf photosynthetic rate

and mesophyll cell anatomy changes during ontogenesis in backcrossed *indica* × *japonica* rice inbred lines. Photosynthesis Research **134**, 27–38.

Hepworth J, Lenhard M. 2014. Regulation of plant lateral-organ growth by modulating cell number and size. Current Opinion in Plant Biology **17**, 36–42.

Horiguchi G, Ferjani A, Fujikura U, Tsukaya H. 2006. Coordination of cell proliferation and cell expansion in the control of leaf size in *Arabidopsis thaliana*. Journal of Plant Research **119**, 37–42.

Horiguchi G, Kim G-T, Tsukaya H. 2005. The transcription factor AtGRF5 and the transcription coactivator AN3 regulate cell proliferation in leaf primordia of *Arabidopsis thaliana*. The Plant Journal **43**, 68–78.

Horiguchi G, Tsukaya H. 2011. Organ size regulation in plants: Insights from compensation. Frontiers in Plant Science **2**, 24.

Hu Y, Xie Q, Chua N. 2003. The Arabidopsis auxin-inducible gene *ARGOS* controls lateral organ size. The Plant Cell **15**, 1951–1961.

Hymus GJ, Cai S, Kohl EA, *et al.* 2013. Application of HB17, an *Arabidopsis* class II homeodomain-leucine zipper transcription factor, to regulate chloroplast number and photosynthetic capacity. Journal of Experimental Botany **64**, 4479–4490.

Ito T, Kim G-T, Shinozaki K. 2000. Disruption of an *Arabidopsis* cytoplasmic ribosomal protein S13-homologous gene by transposon-mediated mutagenesis causes aberrant growth and development. The Plant Journal **22**, 257–264.

Jellings AJ, Leech RM. 1984. Anatomical variation in first leaves of nine *Triticum* genotypes, and its relationship to photosynthetic capacity. New Phytologist **96**, 371–382.

John GP, Scoffoni C, Buckley TN, Villar R, Poorter H, Sack L. 2017. The anatomical and compositional basis of leaf mass per area. Ecology Letters **20**, 412–425.

Kaiser E, Morales A, Harbinson J. 2018. Fluctuating light takes crop photosynthesis on a Rollercoaster Ride. Plant Physiology **176**, 977–989.

Kang J, Mizukami Y, Wang H, Fowke L, Dengler NG. 2007. Modification of cell

proliferation patterns alters leaf vein architecture in *Arabidopsis thaliana*. Planta **226**, 1207–1218.

Kim S-J, Held MA, Zemelis S, Wilkerson C, Brandizzi F. 2015. CGR2 and CGR3 have critical overlapping roles in pectin methylesterification and plant growth in *Arabidopsis thaliana*. The Plant Journal **82**, 208–220.

Kogami H, Hanba YT, Kibe T, Terashima I, Masuzawa T. 2001. CO_2 transfer conductance, leaf structure and carbon isotope composition of *Polygonum cuspidatum* leaves from low and high altitudes. Plant, Cell & Environment **24**, 529–538.

Kojima K, Tamura J, Chiba H, Fukada K, Tsukaya H, Horiguchi G. 2018. Two nucleolar proteins, GDP1 and OLI2, function as ribosome biogenesis factors and are preferentially involved in promotion of leaf cell proliferation without strongly affecting leaf adaxial–abaxial patterning in *Arabidopsis thaliana*. Frontiers in Plant Science **8**, 2240.

Larkin RM, Stefano G, Ruckle ME, Stavoe AK, Sinkler CA, Brandizzi F, Malmstrom CM, Osteryoung KW. 2016. *REDUCED CHLOROPLAST COVERAGE* genes from *Arabidopsis thaliana* help to establish the size of the chloroplast compartment. Proceedings of the National Academy of Sciences **113**, E1116–E1125.

Lee BH, Kwon SH, Lee S-J, Park SK, Song JT, Lee S, Lee MM, Hwang Y, Kim JH. 2015. The *Arabidopsis thaliana* NGATHA transcription factors negatively regulate cell proliferation of lateral organs. Plant Molecular Biology **89**, 529–538.

Lehmeier C, Pajor R, Lundgren MR, *et al.* 2017. Cell density and airspace patterning in the leaf can be manipulated to increase leaf photosynthetic capacity. The Plant Journal **92, 981–994.

Long BM, Bahar NHA, Atkin OK. 2015*a*. Contributions of photosynthetic and non-photosynthetic cell types to leaf respiration in *Vicia faba* L. and their responses to growth temperature. Plant, Cell & Environment **38**, 2263–2276.

Long SP, Marshall-Colon A, Zhu X-G. 2015*b*. Meeting the global food demand of the future by engineering crop photosynthesis and yield potential. Cell **161**, 56–66.

Lu Z, Lu J, Pan Y, Lu P, Li X, Cong R, Ren T. 2016. Anatomical variation of mesophyll conductance under potassium deficiency has a vital role in determining leaf photosynthesis: Structural controls on g_m under K deficiency. Plant, Cell & Environment **39**, 2428–2439.

Maple J, Fujiwara MT, Kitahata N, Lawson T, Baker NR, Yoshida S, Møller SG. 2004. GIANT CHLOROPLAST 1 is essential for correct plastid division in *Arabidopsis*. Current Biology **14**, 776–781.

Micol JL. 2009. Leaf development: time to turn over a new leaf? Current Opinion in Plant Biology **12**, 9–16.

Miyazawa S-I, Terashima I. 2001. Slow development of leaf photosynthesis in an evergreen broad-leaved tree, *Castanopsis sieboldii*: relationships between leaf anatomical characteristics and photosynthetic rate. Plant, Cell & Environment **24**, 279–291.

Mizukami Y, Fischer RL. 2000. Plant organ size control: *AINTEGUMENTA* regulates growth and cell numbers during organogenesis. Proceedings of the National Academy of Sciences of the United States of America **97**, 942–947.

Nakaya M, Tsukaya H, Murakami N, Kato M. 2002. Brassinosteroids control the proliferation of leaf cells of *Arabidopsis thaliana*. Plant and Cell Physiology **43**, 239–244.

Narita NN, Moore S, Horiguchi G, Kubo M, Demura T, Fukuda H, Goodrich J, Tsukaya H. 2004. Overexpression of a novel small peptide ROTUNDIFOLIA4 decreases cell proliferation and alters leaf shape in *Arabidopsis thaliana*. The Plant Journal **38**, 699–713.

Niinemets Ü. 2001. Global-scale climatic controls of leaf dry mass per area, density, and thickness in trees and shrubs. Ecology **82**, 453–469.

Niinemets Ü, Berry JA, Caemmerer S von, Ort DR, Parry MAJ, Poorter H. 2017. Photosynthesis: ancient, essential, complex, diverse … and in need of improvement in a changing world. New Phytologist **213**, 43–47.

Niinemets Ü, Wright IJ, Evans JR. 2009. Leaf mesophyll diffusion conductance in 35 Australian sclerophylls covering a broad range of foliage structural and physiological variation. Journal of Experimental Botany **60**, 2433–2449.

Nishimura C, Ohashi Y, Sato S, Kato T, Tabata S, Ueguchi C. 2004. Histidine kinase homologs that act as cytokinin receptors possess overlapping functions in the regulation of shoot and root growth in *Arabidopsis*. The Plant Cell **16**, 1365–1377.

Nobel PS. 2009. Chapter 8 - Leaves and Fluxes. Physicochemical and Environmental Plant Physiology (Fourth Edition). San Diego: Academic Press, 364–437.

Nobel PS, Zaragoza LJ, Smith WK. 1975. Relation between mesophyll surface area, photosynthetic rate, and illumination level during development for leaves of *Plectranthus parviflorus* Henckel 1. Plant Physiology **55**, 1067–1070.

Ochoa-Villarreal M, Aispuro-Hernndez E, Vargas-Arispuro I, ngel M. 2012. Plant cell wall polymers: function, structure and biological activity of their Derivatives. In: De Souza Gomes A, ed. Polymerization. InTech, .

Oguchi R, Hikosaka K, Hirose T. 2003. Does the photosynthetic light-acclimation need change in leaf anatomy? Plant, Cell & Environment **26**, 505–512.

Oguchi R, Hikosaka K, Hirose T. 2005. Leaf anatomy as a constraint for photosynthetic acclimation: differential responses in leaf anatomy to increasing growth irradiance among three deciduous trees. Plant, Cell & Environment **28**, 916–927.

Omidbakhshfard MA, Proost S, Fujikura U, Mueller-Roeber B. 2015. Growth-Regulating Factors (GRFs): A small transcription factor family with important functions in plant biology. Molecular Plant **8**, 998–1010.

Onoda Y, Hikosaka K, Hirose T. 2004. Allocation of nitrogen to cell walls decreases photosynthetic nitrogen-use efficiency. Functional Ecology **18**, 419–425.

Onoda Y, Wright IJ, Evans JR, Hikosaka K, Kitajima K, Niinemets Ü, Poorter H, Tosens T, Westoby M. 2017. Physiological and structural tradeoffs underlying the leaf economics spectrum. New Phytologist **214**, 1447–1463.

Ort DR, Merchant SS, Alric J, *et al.* 2015. Redesigning photosynthesis to sustainably

meet global food and bioenergy demand. Proceedings of the National Academy of Sciences **112**, 8529–8536.

Parkhurst DF. 1994. Diffusion of CO_2 and other gases inside leaves. New Phytologist **126**, 449–479.

Peguero-Pina JJ, Flexas J, Galmés J, Niinemets Ü, Sancho-Knapik D, Barredo G, Villarroya D, Gil-Pelegrín E. 2012. Leaf anatomical properties in relation to differences in mesophyll conductance to CO_2 and photosynthesis in two related Mediterranean *Abies* species. Plant, Cell & Environment **35**, 2121–2129.

Peguero-Pina JJ, Sisó S, Flexas J, Galmés J, García-Nogales A, Niinemets Ü, Sancho-Knapik D, Saz MÁ, Gil-Pelegrín E. 2017. Cell-level anatomical characteristics explain high mesophyll conductance and photosynthetic capacity in sclerophyllous Mediterranean oaks. New Phytologist **214**, 585–596.

Peguero-Pina JJ, Sisó S, Sancho-Knapik D, Díaz-Espejo A, Flexas J, Galmés J, Gil-Pelegrín E. 2016. Leaf morphological and physiological adaptations of a deciduous oak (*Quercus faginea* Lam.) to the Mediterranean climate: a comparison with a closely related temperate species (*Quercus robur* L.). Tree Physiology **36**, 287–299.

Poorter H, Niinemets Ü, Poorter L, Wright IJ, Villar R. 2009. Causes and consequences of variation in leaf mass per area (LMA): a meta-analysis. New Phytologist **182**, 565–588.

Pyankov VI, Kondratchuk AV, Shipley B. 1999. Leaf structure and specific leaf mass: the alpine desert plants of the Eastern Pamirs, Tadjikistan. New Phytologist **143**, 131–142.

Pyke KA, Leech RM. 1991. Rapid image analysis screening procedure for identifying chloroplast number mutants in mesophyll cells of *Arabidopsis thaliana* (L.) Heynh. Plant Physiology **96**, 1193–1195.

Pyke KA, Leech RM. 1992. Chloroplast division and expansion is radically altered by nuclear mutations in *Arabidopsis thaliana*. Plant Physiology **99**, 1005–1008.

Pyke KA, Leech RM. 1994. A Genetic analysis of chloroplast division and expansion in *Arabidopsis thaliana*. Plant Physiology **104**, 201–207.

de la Riva EG, Olmo M, Poorter H, Ubera JL, Villar R. 2016. Leaf mass per area (LMA) and its relationship with leaf structure and anatomy in 34 Mediterranean woody species along a water availability gradient. PLOS ONE **11**, e0148788.

Rodriguez RE, Debernardi JM, Palatnik JF. 2014. Morphogenesis of simple leaves: regulation of leaf size and shape. Wiley Interdisciplinary Reviews: Developmental Biology **3**, 41–57.

Rodriguez RE, Mecchia MA, Debernardi JM, Schommer C, Weigel D, Palatnik JF. 2010. Control of cell proliferation in *Arabidopsis thaliana* by microRNA miR396. Development **137**, 103–112.

Rodriguez RE, Schommer C, Palatnik JF. 2016. Control of cell proliferation by microRNAs in plants. Current Opinion in Plant Biology **34**, 68–76.

Sack L, Scoffoni C, John GP, Poorter H, Mason CM, Mendez-Alonzo R, Donovan LA. 2013. How do leaf veins influence the worldwide leaf economic spectrum? Review and synthesis. Journal of Experimental Botany **64**, 4053–4080.

Sack L, Scoffoni C, Johnson DM, Buckley TN, Brodribb TJ. 2015. The anatomical determinants of leaf hydraulic function. In: Hacke U, ed. Functional and Ecological Xylem Anatomy. Cham: Springer International Publishing, 255–271.

Sáez PL, Bravo LA, Cavieres LA, Vallejos V, Sanhueza C, Font-Carrascosa M, Gil-Pelegrín E, Javier Peguero-Pina J, Galmés J. 2017. Photosynthetic limitations in two Antarctic vascular plants: importance of leaf anatomical traits and Rubisco kinetic parameters. Journal of Experimental Botany **68**, 2871–2883.

Sage TL, Sage RF. 2009. The Functional anatomy of rice leaves: Implications for refixation of photorespiratory CO_2 and efforts to engineer C_4 photosynthesis into rice. Plant and Cell Physiology **50**, 756–772.

Schommer C, Debernardi JM, Bresso EG, Rodriguez RE, Palatnik JF. 2014. Repression of cell proliferation by miR319-regulated TCP4. Molecular Plant **7**,

1533–1544.

Scoffoni C, Chatelet DS, Pasquet-kok J, Rawls M, Donoghue MJ, Edwards EJ, Sack L. 2016. Hydraulic basis for the evolution of photosynthetic productivity. Nature Plants **2**, 16072.

Scoffoni C, Rawls M, McKown A, Cochard H, Sack L. 2011. Decline of leaf hydraulic conductance with dehydration: Relationship to leaf size and venation architecture. Plant Physiology **156**, 832–843.

Scoffoni C, Vuong C, Diep S, Cochard H, Sack L. 2014. Leaf shrinkage with dehydration: coordination with hydraulic vulnerability and drought tolerance. Plant Physiology **164**, 1772-1788.

Sharkey TD, Weise SE. 2012. Autotrophic carbon dioxide fixation. In: Eaton-Rye J, Tripathy BC, Sharkey TD, eds. Photosynthesis. Dordrecht: Springer Netherlands, 651–674.

Slattery RA, Walker BJ, Weber APM, Ort DR. 2018. The impacts of fluctuating light on crop performance. Plant Physiology **176**, 990–1003.

Smith WK, Vogelmann TC, Critchley C (Eds.). 2004. Photosynthetic Adaptation: Chloroplast to Landscape. New York: Springer.

Souza APD, Massenburg LN, Jaiswal D, Cheng S, Shekar R, Long SP. 2017. Rooting for cassava: insights into photosynthesis and associated physiology as a route to improve yield potential. New Phytologist **213**, 50–65.

Syvertsen JP, Lloyd J, McConchie C, Kriedemann PE, Farquhar GD. 1995. On the relationship between leaf anatomy and CO_2 diffusion through the mesophyll of hypostomatous leaves. Plant, Cell & Environment **18**, 149–157.

Takai T, Adachi S, Taguchi-Shiobara F, *et al.* 2013. A natural variant of *NAL1*, selected in high-yield rice breeding programs, pleiotropically increases photosynthesis rate. Scientific Reports **3**, 2149

Tenhaken R. 2015. Cell wall remodeling under abiotic stress. Frontiers in Plant Science **5**, 771.

Terashima I, Fujita T, Inoue T, Chow WS, Oguchi R. 2009. Green light drives leaf photosynthesis more efficiently than red light in strong white light: Revisiting the enigmatic question of why leaves are green. Plant and Cell Physiology **50**, 684–697.

Terashima I, Hanba YT, Tazoe Y, Vyas P, Yano S. 2006. Irradiance and phenotype: comparative eco-development of sun and shade leaves in relation to photosynthetic CO_2 diffusion. Journal of Experimental Botany **57**, 343–354.

Terashima I, Hanba YT, Tholen D, Niinemets U. 2011. Leaf functional anatomy in relation to photosynthesis. Plant Physiology **155**, 108–116.

Terashima I, Miyazawa S-I, Hanba YT. 2001. Why are sun leaves thicker than shade leaves? — Consideration based on analyses of CO_2 diffusion in the leaf. Journal of Plant Research **114**, 93–105.

Théroux-Rancourt G, Gilbert ME. 2017. The light response of mesophyll conductance is controlled by structure across leaf profiles. Plant, Cell & Environment **40**, 726–740.

Tholen D, Boom C, Zhu X-G. 2012. Opinion: Prospects for improving photosynthesis by altering leaf anatomy. Plant Science **197**, 92–101.

Tomás M, Flexas J, Copolovici L, Galmés J, Hallik L, Medrano H, Ribas-Carbó M, Tosens T, Vislap V, Niinemets Ü. 2013. Importance of leaf anatomy in determining mesophyll diffusion conductance to CO_2 across species: quantitative limitations and scaling up by models. Journal of Experimental Botany **64**, 2269–2281.

Tosens T, Niinemets Ü, Vislap V, Eichelmann H, Castro Díez P. 2012. Developmental changes in mesophyll diffusion conductance and photosynthetic capacity under different light and water availabilities in *Populus tremula*: how structure constrains function. Plant, Cell & Environment **35**, 839–856.

Tsukaya H. 2016. Yield increase: GRFs provide the key. Nature Plants **2**, 15210.

Tsukaya H. 2018. Leaf shape evolution in the context of the primary function of the leaf as a photosynthetic orga. In: Adams III W, Terashima I, eds. The Leaf: A Platform for Performing Photosynthesis. Dordrecht: Springer Netherlands, In press.

Ustin SL, Jacquemoud S, Govaerts Y. 2001. Simulation of photon transport in a three-dimensional leaf: implications for photosynthesis. Plant, Cell & Environment **24**, 1095–1103.

Vercruyssen L, Tognetti VB, Gonzalez N, Van Dingenen J, De Milde L, Bielach A, De Rycke R, Van Breusegem F, Inzé D. 2015. GROWTH REGULATING FACTOR5 stimulates Arabidopsis chloroplast division, photosynthesis, and leaf longevity. Plant Physiology **167**, 817–832.

Veromann-Jürgenson L-L, Tosens T, Laanisto L, Niinemets Ü. 2017. Extremely thick cell walls and low mesophyll conductance: welcome to the world of ancient living! Journal of Experimental Botany **68**, 1639–1653.

Vogelmann TC. 1993. Plant tissue optics. Annual Review of Plant Physiology and Plant Molecular Biology **44**, 231–251.

Vogelmann TC, Bornman JF, Yates DJ. 1996*a*. Focusing of light by leaf epidermal cells. Physiologia Plantarum **98**, 43–56.

Vogelmann TC, Evans JR. 2002. Profiles of light absorption and chlorophyll within spinach leaves from chlorophyll fluorescence. Plant, Cell & Environment **25**, 1313–1323.

Vogelmann TC, Nishio JN, Smith WK. 1996*b*. Leaves and light capture: light propagation and gradients of carbon fixation within leaves. Trends in Plant Science **1**, 65–70.

Von Caemmerer S, Evans J. 1991. Determination of the average partial pressure of CO_2 in chloroplasts from leaves of several C_3 plants. Australian Journal of Plant Physiology **18**, 287.

Warren CR. 2008. Stand aside stomata, another actor deserves centre stage: the forgotten role of the internal conductance to CO_2 transfer. Journal of Experimental Botany **59**, 1475–1487.

Weraduwage SM, Chen J, Anozie FC, Morales A, Weise SE, Sharkey TD. 2015. The relationship between leaf area growth and biomass accumulation in *Arabidopsis*

thaliana. Frontiers in Plant Science **6**, 167.

Weraduwage SM, Kim S-J, Renna L, Anozie FC, Sharkey TD, Brandizzi F. 2016. Pectin methylesterification impacts the relationship between photosynthesis and plant growth in *Arabidopsis thaliana*. Plant Physiology, pp.00173.2016.

Wilson D, Cooper JP. 1970. Effect of selection for mesophyll cell size on growth and assimilation in *Lolium perenne* L. New Phytologist **69**, 233–245.

Xiao Y, Zhu X-G. 2017. Components of mesophyll resistance and their environmental responses: A theoretical modelling analysis. Plant, Cell & Environment **40**, 2729–2742.

Xue J, Luo D, Xu D, Zeng M, Cui X, Li L, Huang H. 2015. CCR1, an enzyme required for lignin biosynthesis in Arabidopsis, mediates cell proliferation exit for leaf development. The Plant Journal **83**, 375–387.

Yin X, Struik PC. 2017. Can increased leaf photosynthesis be converted into higher crop mass production? A simulation study for rice using the crop model GECROS. Journal of Experimental Botany **68**, 2345–2360.

Zhao D, Yang M, Solava J, Ma H. 1999. The *ASK1* gene regulates development and interacts with the *UFO* gene to control floral organ identity in Arabidopsis. Developmental Genetics **25**, 209–223.

Zhong R, Morrison III WH, Freshour GD, Hahn MG, Ye Z-H. 2003. Expression of a mutant form of cellulose synthase AtCesA7 causes dominant negative effect on cellulose biosynthesis. Plant Physiology **132**, 786–795.

Zhou Q, Wang S, Hu B, Chen H, Zhang Z, Huang S. 2015. An *ACCUMULATION AND REPLICATION OF CHLOROPLASTS 5* gene mutation confers light green peel in cucumber. Journal of Integrative Plant Biology **57**, 936–942.

Zhu X-G, Long SP, Ort DR. 2010. Improving photosynthetic efficiency for greater yield. Annual Review of Plant Biology **61**, 235–261.

Figure captions

Fig. 1. Scatterplot matrix of critical leaf morphology and photosynthetic parameters based on literature data. Note: The matrix was conducted by using the "car" package in R software. Diagonal panels show the distribution of each variable. Lower panels show scatterplot. *Tl*, leaf thickness; *Tmes*, mesophyll thickness; *fias*, fraction of intercellular air space; *LMA*, leaf dry mass per unit area; *Tcw*, mesophyll cell wall thickness; *Lchl*, chloroplast length; *Tchl*, chloroplast thickness; *Smes*, mesophyll surface area exposed to intercellular air space; *Sc*, chloroplast surface area exposed to intercellular air space; *Sc.Smes*, the ratio of Sc/Smes; *gm*, mesophyll conductance; *gs*, stomatal conductance; *An*, photosynthesis rate.

Fig. 2. Correlation between leaf morphology and photosynthetic parameters based on literature data. Note: The correlation was conducted by using the "corrr" package in R software. The closer each variable is to another represents the higher relationship while the opposite is for widely spaced variables. The line color represents the direction of the correlation. The blue line is positive correlation and the red line is negative correlation. The line shade and thickness represent the strength of the relationship. The minimum correlation coefficient required to display a line between variables is 0.3.

Fig. 3. Representations of geometrical idealization of mesophyll cells showing how geometry affects the S_{mes} and f_{ias} value. *Smes*, mesophyll surface area exposed to intercellular air space; *fias*, fraction of intercellular air space.

Figures

Fig. 1.

Fig. 2.

S_{mes}	π	2π	3π
f_{ias}	0.215	0.215	0.215

Fig. 3.

Tables

Table 1 Matrix of correlation coefficient between leaf anatomical and photosynthetic parameters

	T_L	T_{mes}	f_{ias}	LMA	T_{cw}	L_{chl}	T_{chl}	S_{mes}	S_c	S_c/S_{mes}	g_m	g_s	A_n
T_L	1.00	0.97***	0.11	0.55***	0.60***	0.36**	-0.062	0.24**	0.085	-0.38***	-0.26**	-0.22*	-0.22**
T_{mes}		1.00	0.039	0.63***	0.52***	0.37**	0.15	0.37***	0.36***	-0.18*	-0.20*	-0.30**	-0.12
f_{ias}			1.00	-0.37***	0.033	0.42**	0.52***	0.19*	-0.021	-0.31***	-0.065	-0.22*	-0.23*
LMA				1.00	0.33***	0.19	-0.20*	0.33***	0.27***	-0.12	-0.17*	-0.13	-0.018
T_{cw}					1.00	0.06	0.35***	0.071	-0.27***	-0.48***	-0.44***	-0.45***	-0.52***
L_{chl}						1.00	0.50***	0.081	0.27*	0.32**	0.32*	0.16	0.26*
T_{chl}							1.00	-0.18	-0.26**	0.017	-0.24*	-0.28**	-0.34***
S_{mes}								1.00	0.80***	-0.099	0.22**	0.03	0.18*
S_c									1.00	0.46***	0.49***	0.24**	0.47***
S_c/S_{mes}										1.00	0.44***	0.41***	0.55***
g_m											1.00	0.75***	0.76***
g_s												1.00	0.78***
A_n													1.00

Note: ***, **, and * represents significant at $P<0.001$, $P<0.01$ and $P<0.05$, respectively.

Table 2 Changes of leaf anatomical and photosynthetic parameters under different LMA

LMA (g m^{-2})	<50	50-100	>100	P value
T_L (μm)	145±61c	187±67b	320±172a	<0.001***
T_{mes} (μm)	127±54b	161±68b	306±151a	<0.001***
f_{ias}	0.276±0.165a	0.298±0.108a	0.186±0.128b	0.0076**
T_{cw} (μm)	0.242±0.135c	0.318±0.182b	0.409±0.171a	<0.001***
L_{chl} (μm)	4.38±1.53a	4.79±0.60a	4.91±0.88a	0.26
T_{chl} (μm)	2.38±0.88a	2.60±0.95a	2.09±0.78a	0.081
S_{mes} (m^2 m^{-2})	12.0±5.4b	19.6±4.5a	19.2±9.6a	<0.001***
S_c (m^2 m^{-2})	7.9±4.1b	12.6±5.1a	11.3±5.7a	<0.001***
S_c/S_{mes}	0.68±0.15a	0.72±0.18a	0.61±0.16b	0.013*
g_m (mol m^{-2} s^{-1})	0.119±0.104a	0.131±0.148a	0.062±0.042b	0.029*
g_s (mol m^{-2} s^{-1})	0.177±0.147a	0.123±0.080a	0.134±0.118a	0.073
A_n (μmol m^{-2} s^{-1})	10.9±8.3a	9.6±7.6a	9.8±5.5a	0.54

Note: ***, **, and * represents significant at $P<0.001$, $P<0.01$ and $P<0.05$, respectively.

Figures

Fig. 1.

Fig. 2.

S_{mes}	π	2π	3π
f_{ias}	0.215	0.215	0.215

Fig. 3.