Neurons of the inferior olive respond to broad classes of sensory input while subject to homeostatic control

Homeostatic complex spike firing

Chiheng Ju[1,3], Laurens W.J. Bosman[1,3*], Tycho M. Hoogland[1,2,3*],

Arthiha Velauthapillai[1], Pavithra Murugesan[1], Pascal Warnaar[1], Mario Negrello[1]

and Chris I. De Zeeuw[1,2]

[1] Department of Neuroscience, Erasmus MC, 3015 GD Rotterdam, The Netherlands

[2] Netherlands Institute for Neuroscience, Royal Netherlands Academy of Arts and Sciences, 1105 BE Amsterdam, The Netherlands

[3] These authors contributed equally

* Correspondence to Dr. Laurens Bosman, PO Box 2040, 3000 CA Rotterdam, The Netherlands, l.bosman@erasmusmc.nl; or to Dr. Tycho Hoogland, PO Box 2040, 3000 CA Rotterdam, The Netherlands, t.hoogland@erasmusmc.nl

Abstract

Cerebellar Purkinje cells integrate sensory information with motor efference copies to adapt movements to behavioral and environmental requirements. They produce complex spikes that are triggered by the activity of climbing fibers originating in neurons of the inferior olive. These complex spikes can shape the onset, amplitude and direction of movements and the adaptation of such movements to sensory feedback. Clusters of nearby inferior olivary neurons project to "microzones" which are parasagitally aligned stripes of Purkinje cells. Although it is known that sensory input preferentially recruits coherent climbing fiber activity within one or more microzones, it is still unclear whether individual Purkinje cells within a single microzone integrate climbing fiber input from multiple distinct sources, or how sensory-evoked responses depend on the stimulus strength and recent history of climbing fiber activity. By imaging complex spike responses in cerebellar lobule crus 1 to various types of sensory stimulation in awake mice we find that different sensory modalities and receptive fields have a mild, but consistent, tendency to converge on individual Purkinje cells. Purkinje cells encoding the same stimulus show increased complex spike coherence and tend to lie close together. Moreover, whereas complex spike firing is only mildly affected by variations in stimulus strength, it strongly depends on the recent history of climbing fiber activity. Our data point towards a mechanism in the olivocerebellar system that regulates complex spike firing during mono- or multisensory stimulation around a relatively low set-point, highlighting an integrative coding scheme of complex spike firing under homeostatic control.

Introduction

The olivocerebellar system is paramount for sensorimotor integration during motor behavior. The climbing fibers that originate in the inferior olive and cause complex spike firing in cerebellar Purkinje cells encode both unexpected and expected sensory events and affect initiation, execution as well as adaptation of movements (1-12). Complex spike firing frequency is typically sustained around 1 Hz and not substantially affected by the behavioral state, although short-lived increases or decreases in firing occur (2, 13-17). To date, the paradox between the persistence of complex spike firing and the behavioral relevance of individual complex spikes is still largely unresolved.

Whereas the afferents of Purkinje cells, including not only the climbing fibers but also the parallel fibers and axons of interneurons, all diverge, their efferents strongly converge upon the cerebellar nuclei, ultimately integrating many different inputs from the brain (18-20). The parallel fibers are oriented in a transverse direction along the lobular axes, while the climbing fibers and axons of the interneurons are running perpendicularly to them in line with the sagittal orientation of the dendritic trees of Purkinje cells (10, 21-28). Interestingly, the intrinsic biochemical nature as well as electrophysiological profile of individual Purkinje cells follows the organization of the climbing fibers so that Purkinje cells located in the same sagittal module receive climbing fiber input from the same olivary subnucleus and have similar identity properties, setting them apart from Purkinje cells in neighboring modules (13, 21, 29-32). Accordingly, Purkinje cell responses following electrical stimulation of major nerves of cat limbs largely adhere to the parasagittal organization of the climbing fiber zones (33, 34), which in turn can be further differentiated into smaller microzones based upon their response pattern to tactile stimulation of a particular spot on the body (35-38). Possibly, differential sensory maps even occur at the submicrozonal and individual Purkinje cell level,

but this has to our knowledge not been investigated yet. In particular, it remains unclear to

what extent different types of sensory inputs can drive complex spikes within the same

individual Purkinje cells and/or their direct neighbors and how the strength as well as the

history of these inputs influences the distribution of climbing fiber activity.

Here, we studied the impact of minimal stimuli of distinct sensory modalities on

complex spike firing of Purkinje cells in crus 1 of awake mice using *in vivo* two-photon Ca^{2+}

imaging. We found that different sensory streams appear to converge on individual inferior

olivary neurons and thereby Purkinje cells, that sensory stimulation primarily affects the

timing of the complex spikes rather than their rate, that the strength of complex spike

responses varied seamlessly from non-responsive to highly responsive, that a recent history of

high activity leads to a future of low activity, and that Purkinje cells that respond to the same

stimulus tend to be located in each other's vicinity, increasing the coherence of their

responses. Together, our data indicate that subtle and local sensory inputs can recruit mosaic

ensembles of Purkinje cells, employing population coding in a spatially and temporally

dynamic way that is in line with homeostatic control.

Results

Climbing fiber responses to tactile, auditory and/or visual input

Little is known about the distribution and convergence of different types of climbing fiber-

mediated sensory input at the level of individual, nearby Purkinje cells. Here, we performed

two-photon Ca^{2+} imaging of Purkinje cells in crus 1 of awake mice to record complex spikes

responses to various types of sensory stimulation related to the face of the mice (Fig. 1A) (39-

45). The configurations and positions of individual Purkinje cell dendrites were detected using

independent component analysis (Fig. 1B). Ca^{2+} transients were isolated per dendrite by a

template-matching procedure that reliably detects complex spikes (46-48) (Fig. 1C-E).

Figure 1 – Sensory pathways carrying facial input to the cerebellar cortex

(A) Scheme of the main routes conveying facial tactile input via the climbing fiber pathway to cerebellar

Purkinje cells (PC). Climbing fibers, which cause complex spike firing in Purkinje cells, exclusively originate from

the inferior olive. The inferior olive, in turn, is directly innervated by neurons from the trigeminal nuclei as well

as indirectly via thalamo-cortical pathways that project to the inferior olive mainly via the nuclei of the

mesodiencephalic junction (MDJ). The MDJ itself also receives direct input from the trigeminal nuclei. See the

main text for references. **(B)** *In vivo* two-photon Ca^{2+} imaging was performed to characterize Purkinje cell

complex spike responses to sensory stimulation in the medial part of crus 1. Purkinje cells were detected using

independent component analysis and the position of a Purkinje cell dendrite (yellow area on the right) within a

field of view is shown in the inset. At the end of each recording session, the brain was removed and the

location of the dye injection in medial crus 1 was confirmed through *ex vivo* epifluorescent imaging (black

circle). The white rectangle indicates the approximate recording location. **(C)** Complex spikes that were

triggered by climbing fiber activity were retrieved from fluorescent traces of individual Purkinje cells. A

representative trace obtained from the Purkinje cell dendrite illustrated in **B** is shown together with the

detected complex spikes (grey lines). The light blue episode is enlarged in **D**. Complex spikes were detected by

the combination of a threshold and a template matching algorithm. Only events with a sharp rising phase were

accepted as complex spikes. In the 60 s interval shown in **C**, there was one event with a slower rise time (see

arrow in **D**), as indicated at a larger time scale in **E**. The events in **E** are scaled to peak.

We started our study of sensory responses by using air puff stimulation of the large facial

whiskers, which is a relatively strong stimulus. In line with previous studies (2, 4, 10, 49, 50),

we found that air puff stimulation to the whiskers evoked complex spike responses in many

Purkinje cells (e.g., in 19 out of 19 Purkinje cells in the example illustrated in Fig. 2). In total,

102 out of the 118 (86%) cells analyzed were considered to be responsive to such a stimulus,

implying that the peak responses of these cells exceeded the threshold of the average + 3 s.d.

of the pre-stimulus period.

Figure 2 - Whisker stimulation evokes complex spike responses in cerebellar crus 1

(A) Complex spikes elicit large increases in the Ca^{2+} concentration within Purkinje cell dendrites that can be

resolved using *in vivo* two-photon microscopy in combination with a fluorescent Ca^{2+} indicator. An example of a

field of view with 19 identified Purkinje cell dendrites located in the medial part of crus 1 is shown with each

individual dendrite denoted by a number and a unique color. This recording was made in an awake mouse. **(B)**

The fluorescent traces of each of these dendrites show distinct Ca^{2+} transient events, which are to a large

extent associated with air puff stimulation of the facial whiskers (times of stimulation indicated by the vertical

lines). The boxed area is enlarged in Fig. S1. **(C)** Summed fluorescence trace composed of all 19 individual traces

emphasizing the participation of many Purkinje cells to the stimulus-triggered responses. **(D)** After complex

spike extraction, a clear relation between stimulus and activity was observed as illustrated by summing, for

each frame, the number of complex spikes observed over all dendrites. The vertical scale bar corresponds to

the simultaneous activity of 5 Purkinje cells. **(E)** Peri-stimulus time histogram of a Purkinje cell dendrite

(marked as number 7 in **A** and **B**) in response to air puff stimulation to the ipsilateral whiskers (154 trials). The

bin size (40 ms) corresponds to the acquisition frame rate (25 Hz). **(F)** Normalized stacked line graph of the

Purkinje cells in this field of view showing that every cell contributed to the overall response. The Purkinje cells

are ranked by their maximal response and the data are normalized so that the top line reflects the average

frequency per bin. Cell no. 7 (dashed line) had a relatively poor signal-to-noise ratio during later parts of the

recording (see Fig. S1), but it had nevertheless a complex spike response profile that was indistinguishable from

the other cells. The colors match those in the panels **A**-**B**. Inset: In total, 102 out of 118 cells analyzed (86%)

were responsive to whisker air puff stimulation (peak response exceeded average + 3 s.d. of pre-stimulus

interval). **(G)** Fluorescent traces of the trials with (red) and without (black) complex spikes fired during the first

200 ms after air puff onset. In the absence of complex spike firing, only a very small increase in fluorescence

was observed, indicating that the majority of change in fluorescence was associated with complex spike firing.

Note the longer time scale than in **E** and **F**. The lines indicate the medians and the shaded areas the inter-

quartile ranges of the 19 Purkinje cells in this field of view.

To address whether localized and more subtle stimuli recruited smaller groups of Purkinje

cells, maybe even subsets of microzones, we subsequently applied gentle tactile stimulation at

four facial locations: the whisker pad, the cheek posterior to the whisker pad, the upper lip

and the lower lip (Fig. 3A). The stimuli were given using a Von Frey filament (target force =

0.686 mN) attached to a piezo actuator. The stimulus strength was carefully calibrated to

avoid inducing responses from neighboring skin areas or nociceptive responses (see Materials

and Methods). When we evaluated the strength of the Purkinje cell responses to any of the

four tactile stimuli (998 stimulus conditions in 282 Purkinje cells), a skewed, but continuous

distribution was found (Fig. 3B). Moreover, also for individual stimulus locations skewed, but

continuous distributions of response strengths were found, with the upper lip being the least

sensitive area (Fig. S2). In other words, our data set contained Purkinje cells that did not

respond at all as well as cells that responded strongly, with all gradients in between. Thus, a

distinction between "responsive" and "non-responsive" Purkinje cells involved a somewhat

arbitrary distinction, prompting us to present most of the subsequent analyses using the entire

dataset.

Figure 3 – Purkinje cells in crus 1 respond to various types of sensory stimulation

(A) Tactile stimuli were presented to four facial regions in awake mice: the whisker pad (WP), the upper lip (UL), the lower lip (LL) and the cheek (Ch). Each stimulus was delivered with a piezo-actuator that made a muted, yet audible sound which was also delivered without touch ("sound only (SO)"). A blue light flash generated by an LED was used as visual stimulus. These experiments were performed in awake mice. (B) To avoid interference of adjacent areas, we applied gentle touches (0.686 mN). The complex spike response ratio was much reduced relative to the strong air puff stimulation to all ipsilateral whiskers illustrated in Fig. 2. A histogram of the peak responses (expressed as Z value) of all responses to either of the four tactile stimuli demonstrates that the response strength is a continuum, showing the lack of a clear separation between "responsive" and "non-responsive" Purkinje cells (998 stimulus conditions in 282 Purkinje cells). We considered Purkinje cells that showed a peak response above $Z = 3$ as "significantly responsive" (represented with black bars), but we provide most of the analyses also for the population as a whole (e.g., Fig. S2). (C) The peri-stimulus time histograms (PSTHs) of a representative Purkinje cell. The shades of grey indicate 1, 2 and 3 s.d. around the average. Each stimulus was repeated 154 times at 1 Hz. (D) For every stimulus condition, we averaged the PSTHs for all Purkinje cells that were significantly responsive to that particular stimulus (colored lines; averages ± SEM). These were contrasted to the averaged PSTH of the other Purkinje cells (black lines; averages ± SEM). The pie charts represent the fraction of Purkinje cells significantly responsive to a particular stimulus. See also Table 1. (E) The peak responses of the significantly responding Purkinje cells were the lowest for sound only and for upper lip stimulation. * $p < 0.05$; ** $p < 0.01$ (post-hoc tests after Kruskal-Wallis test) (F) As expected for complex spike responses to weak stimulation, the latencies were relatively long and variable, but consistent across types of stimulation. Only visual stimulation (LED) had a remarkably longer latency time. *** $p < 0.001$ (post-hoc tests after Kruskal-Wallis test for LED vs. whisker pad, upper lip, lower lip and cheek and $p < 0.05$ compared to sound only)

Given that the touches were delivered by a piezo-actuator that made a weak but audible sound, we also tested whether Purkinje cells responded to this sound in the absence of touch. This was the case, and, as expected, the "sound-only" stimulus evoked the weakest responses of all stimuli ($p < 0.001$; Kruskal-Wallis test; Fig. S2B). For comparison, we also included a visual stimulation, consisting of a brief flash of a blue LED. This stimulus evoked responses with a similar strength as the whisker pad and lower lip stimulation, but with a latency that was remarkably long (Fig. S2A).

To facilitate a quantitative comparison between the stimulus conditions, we subsequently focused on the subset of obvious responses, defined as having a peak amplitude exceeding our threshold for significance set at the average + 3 s.d. of the pre-stimulus period. Among these "significantly responding" Purkinje cells, we found trends as in the entire population: the lower lip recruited the strongest responses, directly followed by the whisker pad and visual stimulation, while upper lip and sound-only stimulations were less effective (Fig. 3C-E; Table 1). We also confirmed the remarkably long latency, typically more than 100 ms, for the visually evoked responses ($p < 0.001$ compared to the tactile stimuli; Kruskal Wallis test; Fig. 3F; Table 1).

Table 1

Stimulation	Responsive cells		Response parameters		Prevalence (p) $\chi^2 = 25.758$; df = 5; $p = 0.001$				
	No.	**%**	**Peak (Hz)**	**Latency (ms)**	**UL**	**LL**	**Ch**	**SO**	**LED**
Whisker pad	111 (282)	39	2.12 ± 0.82	84±42	**0.010**	0.661	**0.003**	**0.000**	0.828
Upper lip	71 (249)	29	1.89 ± 0.67	84±84		**0.031**	0.597	0.157	0.064
Lower lip	99 (264)	38	2.23 ± 0.97	84±42			**0.010**	**0.001**	1.000
Cheek	53 (203)	26	2.30 ± 1.67	84±42				0.415	**0.028**
Sound only	44 (197)	22	1.74 ± 0.65	84±84					**0.003**
LED	49 (129)	34	2.13 ± 1.18	126±0					

Table 1 – Purkinje cell responses to sensory stimulation in awake mice

Purkinje cells respond with complex firing to sensory stimulation (Fig. 3). For each type of stimulation, the

number and percentage of statistically significantly responsive cells (peak response > average + 3 s.d. of

baseline firing) is indicated (in brackets: total number of cells tested). The response peak and response latency

are indicated as medians ± inter-quartile ranges. The relative prevalence of responses was not equal for all

stimulus types, as confirmed by a 6x2 χ^2 test. Differences in relative prevalence were tested using pair-wise

Fisher's exact tests (as the χ^2 test was significant, no further correction for multiple comparisons was applied).

Bold values indicate significantly different prevalences. UL = upper lip; LL = lower lip; CH = cheek; SO = sound

only.

Convergence of sensory inputs

Next, we addressed the question whether Purkinje cells have a preference to respond to one or

multiple types of stimulation. To this end, we made for each combination of two stimuli a

scatter plot of the response strengths of each individual Purkinje cell. For each and every

combination, linear regression analysis revealed a positive correlation, implying that a

stronger response to one stimulus typically implied also a stronger response to the other (Fig.

4A). These correlations, although weak, were statistically significant for all combinations

except in two cases (i.e., upper lip vs. sound only and upper lip vs. visual stimulation; Table

2). The regression lines deviated from the 45° line, suggesting that, although there is a

tendency at the population level to combine inputs, individual Purkinje cells can display

specificity for a given stimulus. The Venn diagrams in Fig. 4B-C highlight the degree of

overlap for all combinations of two and three stimuli. For all combinations we found some but

no complete overlap. Remarkably, when comparing the observed overlap with the expected

overlap based upon a random distribution, all combinations occurred more often than

predicted. We performed a bootstrap analysis (see Materials and methods) to infer statistical

significance and indeed, most combinations were observed significantly more often than

expected from a random distribution of inputs (Fig. 4B-C, Tables 3 and 4). This also included

the visual stimulation, so that those Purkinje cells responding to a tactile stimulus typically

were responsive to visual stimulation as well.

Figure 4 – Convergence of sensory input on Purkinje cells

(A) In order to test whether sensory inputs converge on individual Purkinje cells in awake mice, we made pair-

wise comparisons of the response amplitudes to two different stimuli per Purkinje cell (scatter plots). For all

possible combinations, we found a positive slope of the linear regression analysis. For the majority of

combinations, the correlation between response strengths was highly significant: * $p < 0.05$, ** $p < 0.01$ and

*** $p < 0.001$ (Pearson correlation with Benjamini-Hochberg correction for multiple comparisons). Only upper

lip vs. sound only and upper lip vs. visual stimulation were not significantly correlated. For this analysis, we

included all Purkinje cells, whether they had a statistically significant response or not. The red dotted lines

indicate a Z-score of 3, which we set as the threshold for significance (cf. Fig. 3B). They grey arrows indicate the

fraction of observations above and below the unity line (grey dotted line). The relative strengths of every

stimulus combination were compared in a pairwise fashion (Wilcoxon tests with Benjamini-Hochberg

correction for multiple comparisons): & $p < 0.05$; && $p < 0.01$; &&& $p < 0.001$. **(B)** We performed a similar

analysis focusing only on statistically significant responses (Venn diagrams). Again, all combinations had a

positive Z score (as evaluated by a bootstrap method; see Materials and Methods), indicating more than

expected convergence. The diameter of each circle indicates the fraction of Purkinje cells showing a significant

response to that particular, color coded stimulus. The size of the bar represents the Z score of the overlapping

fraction. **(C)** The same for the combinations of three tactile stimuli. Overall, sensory streams tended to

converge, rather than diverge, on Purkinje cells. # $p < 0.10$; * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$ (Z test).

Table 2

Stimulation	Pearson correlation (R)					Significance of Pearson correlation (*p*)				
	UL	**LL**	**Ch**	**SO**	**LED**	**UL**	**LL**	**Ch**	**SO**	**LED**
Whisker pad	0.272	0.253	0.166	0.257	0.358	**<0.001**	**<0.001**	**0.018**	**0.001**	**<0.001**
Upper lip		0.192	0.189	0.130	0.129		**0.002**	**0.001**	0.126	0.179
Lower lip			0.387	0.513	0.339			**<0.001**	**<0.001**	**<0.001**
Cheek				0.485	0.362				**<0.001**	**<0.001**
Sound only					0.523					**<0.001**

Table 2 – Correlations between response probabilities

For each Purkinje cell, we made pair-wise correlations of the Z scores of the amplitudes of the responses to

different stimuli. The Pearson correlation coefficient (R) and the accompanying *p* value are indicated in this

Table. The *p* values indicated in bold reflect significant values (after Benajmini-Hochberg correction for multiple

comparisons). These data are graphically represented in Fig. 4A. WP = whisker pad; UL = upper lip; LL = lower

lip; Ch = cheek; SO = sound only.

Table 3

Stimulation	Rate of occurrence (*Z*)					Chance of occurrence (*p*)				
	UL	**LL**	**Ch**	**SO**	**LED**	**UL**	**LL**	**Ch**	**SO**	**LED**
Whisker pad	1.21	1.25	0.85	1.34	**2.75**	0.225	0.213	0.394	0.181	**0.006**
Upper lip		1.23	**2.00**	1.34	1.94		0.220	**0.045**	0.181	0.052
Lower lip			**2.39**	**2.70**	**2.62**			**0.017**	**0.007**	**0.009**
Cheek				**3.31**	**3.10**				**0.001**	**0.002**
Sound only					**2.28**					**0.022**

245 **Table 3 – Convergence of different sensory streams on individual Purkinje cells**

246 For each type of stimulus, we compared the observed rate of convergence on individual Purkinje cells to

247 chance level, using a bootstrap method based on the relative prevalence of each stimulus (cf. Fig. 4). Indicated

248 are the *Z* and the *p* values. All combinations occurred more often than expected (*Z* > 0). Bold values indicate

249 statistical significance. WP = whisker pad; UL = upper lip; LL = lower lip; Ch = cheek; SO = sound only.

Table 4

Stimulation	Rate of occurrence (*Z*)	Chance of occurrence (*p*)
WP + UL + LL	1.89	0.059
WP + LL + Ch	1.32	0.186
UL + LL + Ch	3.22	**0.001**
WP + UL + LL + Ch	3.07	**0.002**

Table 4 – Convergence of different sensory streams on individual Purkinje cells

For each type of stimulus, we compared the observed rate of convergence on individual Purkinje cells to chance level, using a bootstrap method based on the relative prevalence of each stimulus (cf. Fig. 4). Indicated are the *Z* and the *p* values. All combinations occurred more often than expected (*Z* > 0). Bold values indicate statistical significance. WP = whisker pad; UL = upper lip; LL = lower lip; Ch = cheek.

With an apparent lack of input-specificity, one might wonder whether Purkinje cells in crus 1 are encoding specific sensory events or, alternatively, respond indifferently to any external trigger. In this respect, especially the response pattern to the sound only stimulus is noticeable. A weak, but audible sound was generated by the piezo actuator used to deliver the tactile stimuli. Hence, all tactile stimuli also involved sound. Nevertheless, the Venn diagrams in Fig. 4A show that there are Purkinje cells that responded statistically significantly to sound only, but not to sound and touch delivered simultaneously. This could be taken as an argument against the input-specificity of Purkinje cells. However, one should keep in mind that the separation between "responsive" and "non-responsive" Purkinje cells is arbitrarily (cf. Fig. 3B), and to be able to draw such a conclusion, there should be a consistent absence of preferred responses. The sound only stimulus was found to be the weakest stimulus (Figs. 3E and S2B) and pair-wise comparisons of the response strengths had a strong bias towards the stimuli involving touch (Fig. 4A). This is further illustrated in a single experiment, directly comparing the responses evoked by whisker pad touch and sound only stimulation. Four randomly selected Purkinje cells from this experiment failed to show a response to sound only

stimuli in the absence of touch responses (Fig. S3A-E). This was confirmed by the group-wise

analysis of all Purkinje cells in this experiment, demonstrating a clear preference for the

whisker pad stimulation over the sound-only stimulus (Fig. S3F-G). Overall we conclude that,

although the observed responses to mild sensory stimulation are typically weak and

convergent, a certain degree of input-specificity was observed in Purkinje cells of crus 1.

Stimulus strength has limited impact on complex spike responses

To avoid recruiting responses from adjacent areas, we used weak stimulation strengths (Figs.

3 and 4). Under anesthesia, the strength of a stimulus affects the complex spike response

probability (2, 51). This finding has been reproduced in awake mice using variations in

duration and strength of peri-ocular air puff stimulation (52). As a consequence, our approach

using weak stimuli could have led to an underestimation of the number and spatial extent of

sensory climbing fiber responses. To study whether the stimulus strength could also have

affected our results, we performed an experiment in which we stimulated all whiskers

mechanically with three different strengths, the largest of which was identical to the maximal

stimulus we previously applied under anesthesia (2). The chosen stimulus intensities

maximized the variation in kinetic energy, which was the most salient feature for barrel cortex

neurons (53). Stimuli were randomly intermingled (Fig. 5A-B). Raw traces indicate that, even

at the ensemble level, (relatively) weak stimuli do not trigger responses at every trial and

spontaneous complex spike firing may occasionally appear as peaks prior to stimulus onset

(Figs. 2C, 5C and 5D).

 Figure 5 – Stimulus strength has only a minor impact on complex spike responsiveness

(A) Movements of all large facial whiskers were performed using a piezo-actuator at three different speeds

(weak: 1 mm displacement in 62 ms; moderate: 2 mm displacement in 31 ms; strong: 4 mm displacement in 16

ms). The stimulus sequence was randomly permuted. The recordings were made in awake mice. **(B)** Field of

view with 24 identified and color-coded Purkinje cells (left) and their corresponding fluorescent traces (right).

Stimuli were presented every 2 s and in between trials the laser illumination was briefly blocked to avoid

photobleaching. Note that the periods without laser illumination are not drawn to scale. The vertical shaded

areas indicate stimulus duration (which was inverse with the stimulus strength). **(C)** Summed fluorescence

trace composed of all 24 individual traces showing that not all trials evoked ensemble-wide responses. Some

spontaneous, intertrial activity was also observed. **(D)** The average number of complex spikes per frame (of 40

ms) per trial (shaded areas: ± SEM) for the three stimulus strengths show little difference for the weak and

moderate stimulation. The time course and amplitude (1-4 mm) of the three stimuli is schematized at the

bottom of the graph. Strong stimulation elicited about 30% more complex spikes, as evident from the peak

responses for each stimulus intensity. **(E)** Plotting for each of Purkinje cell the average peak response for each

stimulus condition results in many overlapping lines. Only a few cells stand out in that they show a strong

response that consistently increases with stimulus strength (red lines on top). The Purkinje cells that showed

increased responsiveness to stronger stimuli are depicted in red and the others in blue (see also Fig. S4A). **(F)**

The same data summarized with box plots for all significantly responsive cells. Data are from 209 significantly

responsive Purkinje cells (out of 340 Purkinje cells that were measured in this way). *** $p < 0.001$ (Friedman's

test).

Of the 340 Purkinje cells tested in awake mice, 209 (61%) responded significantly to at least

one stimulus strength. In these 209 Purkinje cells, we compared the complex spike responses

across three intensities. The weak and the moderate intensities (1 mm displacement reached in

62 ms and 2 mm displacement reached in 31 ms, respectively) showed a comparable number

of responses and only the strong stimulus intensity (4 mm reached in 16 ms) evoked

significantly more responses ($F(2) = 57.160$, $p < 0.001$, Friedman's test) (Figs. 5D-F, S4A).

Despite being 16 times faster (250 mm/s instead of 16 mm/s), the strongest stimulus recruited

only 28% (measured as peak response) or 34% (measured as integral of whole response

period) more complex spikes than the weak stimulus (Fig. 5E-F). The same analysis on the

whole population of Purkinje cells, including those that did not show a statistically significant

response, revealed even less of an impact of stimulus strength (Fig. S4B). We therefore

conclude that the complex spike response poorly encodes velocity of whisker displacement in

awake mice and that – within boundaries – using stronger stimuli does not necessarily lead to

qualitatively different results.

Functionally equivalent Purkinje cells tend to group together

So far, we mainly found an abundance of weak and not very specific complex spike responses

to mild sensory stimulation. One way in which such weak responses at the cellular level could

still have considerable effects on the network level would be when functionally equivalent

Purkinje cells would lie together in microzones. Indeed, as the Purkinje cells of each

microzone project to a group of adjacent neurons in the cerebellar nuclei (40, 54), population

encoding of sensory events may be a form of functional signaling in the olivocerebellar

system. Fig. 6 illustrates to what extent adjacent Purkinje cells have similar stimulus response

probabilities. In this example, Purkinje cells reacting strongly to whisker pad stimulation are

grouped together, but such a spatial clustering does not seem to be perfect, as also a few

strongly responsive cells are located in between less responsive Purkinje cells (Fig. 6A) and

as upper lip-responsive Purkinje cells are sparsely distributed across the same area (Fig. 6B).

We reasoned that spatial clustering should imply that two neighboring cells have more similar

response probabilities than randomly selected cells from the same field of view. This turned

out to be the case, but only if we confined our focus to Purkinje cells with statistically

significant responses ($Z > 3$) (Figs. 6C and 6D). Thus, especially the Purkinje cells with

stronger responses to a given stimulus type were found to be located more closely together

than could be expected from a random distribution.

Figure 6 - Purkinje cells encoding the same stimulus have a tendency to be spatially grouped

Schematic drawing of a field of view with 26 Purkinje cells organized in the medio-lateral direction of crus 1 in

an awake mouse. The color of each Purkinje cell corresponds to the maximal response to whisker pad **(A)** or

upper lip **(B)** stimulation. Purkinje cells with a filled soma had a peak response with a Z score > 3 and were

considered to be statistically significant, in contrast to those with an open soma. Responsive and non-

responsive cells are generally intermingled, but a group of "strong responders" can be observed for whisker

pad stimulation (red rectangle). **(C)** The anecdotal data in **A** suggest the presence of clusters of Purkinje cells

encoding specific stimuli. For this to be the case, one would expect that neighboring Purkinje cells have roughly

similar response strengths. We found that this assumption does not hold as the response strengths of

neighbors could not be discriminated from randomly selected cells in the same recording if all Purkinje cells are

considered (compared with bootstrap analysis based upon randomly chosen cell pairs within each field of view:

all $p > 0.8$; Z test). Data are represented in violin plots, with the grey lines indicating the 10^{th}, 25^{th}, 50^{th}, 75^{th} and

90^{th} percentiles. **(D)** When considering only the Purkinje cells with statistically significant responses, spatial

grouping does occur. For each stimulus type, the black portion of the left bar indicates the fraction of Purkinje

cells showing a significant response to that stimulus. The filled portion of the right bar indicates the fraction of

the neighbors (always on the medial side) of these significantly responsive Purkinje cells that were also

significantly responsive. As can be seen, this fraction is always substantially larger than the fraction of

significantly responsive Purkinje cells, indicating a tendency of similar Purkinje cells to group together.

Statistical significance was tested by comparing the fraction of Purkinje cells with statistically significant

responses and the fraction of neighbors of Purkinje cells with statistically significant responses that showed

statistically significant responses as well (after correction for border effects) using Fisher's exact test and after

Benjamini-Hochberg correction for multiple comparisons: * $p < 0.05$; *** $p < 0.001$.

Single whisker responses in Purkinje cells

Using our tactile stimuli we demonstrated a tendency of nearby Purkinje cells to encode the same stimulus. To study whether this would hold true for even smaller receptive fields we turned to single whisker stimulation. Using single-unit electrophysiological recordings we have previously shown that stimulation of a single whisker is sufficient to evoke complex spike responses (2). Individual whiskers can be reliably identified across mice and as such they can be qualified as minimal reproducible receptive fields. We repeated our previous single-whisker stimulation experiments now using two-photon Ca^{2+} imaging focusing on five whiskers that were stimulated individually in a random sequence. To prevent spontaneous whisking and thereby interference by other whiskers, these experiments were (unlike all other experiments in this study) performed under anesthesia. In general, the responses were specific, as many Purkinje cells responded to a particular whisker, but not to its neighboring whiskers (Fig. 7A). Overall, of the 148 Purkinje cells tested, 31 (21%) responded significantly to only one of the five whiskers tested, 14 (10%) to two and 5 (3%) to three whiskers (Fig. 7B). Not all whiskers were equally effective in recruiting Purkinje cell responses: 23 cells (15%) responded significantly to C3 stimulation, but only 4 (3%) to C1 stimulation. Likewise, pairs of more anterior whiskers had higher chances to be encoded by the same Purkinje cell (Fig. 7C; Table 5). Thus, also for single whisker stimulation, there was a balance between sensory integration and specificity.

Figure 7 – Purkinje cell responses to single whisker stimulation show weak clustering

(A) To investigate smaller receptive fields, we sequentially stimulated five of the large facial whiskers. To avoid interference with other whiskers during active movement, we performed these experiments under ketamine/xylazine anesthesia. Most Purkinje cells, if responsive to single-whisker stimulation, responded only to one of the five whiskers **(B)**. This is illustrated by five peri-stimulus time histograms (PSTHs) from a single, representative Purkinje cell. This particular cell was sensitive to stimulation of the C3 whisker only. The average

and 3 s.d. of the baseline firing are indicated (dashed line and grey area). **(C)** Purkinje cells that responded to
more than one whisker were typically responsive to the more anterior whiskers (see also Table 5). The widths
of the lines indicate the Z value of the occurrence of multiple responses per cell. **(D)** Two recording spots, in
close proximity in crus 1 of the same animal, with the identified Purkinje cell dendritic trees. For each dendrite,
the color indicates the whisker(s) to which it was responsive (see legend below with grey denoting the absence
of a statistically significant response). **(E)** For each of the two recording sites, the average of the responsive and
the non-responsive Purkinje cells is indicated (to the C3 whisker in the left panel and to the C2 whisker in the
right panel). Note that only a single cell was responsive to C2 stimulation in recording spot 2. The firing patterns
of the non-responsive cells were also averaged (black lines with shades indicating S.E.M.). **(F)** Linear regression
revealed that Purkinje cells that were surrounded by other Purkinje cells responsive to the same whisker (same
color code as in **A**) had a tendency to show stronger responses to stimulation of that whisker than Purkinje cells
that were more isolated. The x-axis represents the fraction of Purkinje cells responsive to the particular whisker
within the respective field of view. R = 0.521; $p < 0.001$.

Table 5

Whisker	Responsive cells		Rate of occurrence (Z)				Prevalence (p)			
	No.	**%**	**C1**	**C2**	**C3**	**D2**	**C1**	**C2**	**C3**	**D2**
B2	15 (148)	10	0.630	0.227	**5.130**	**2.052**	0.264	0.410	**<0.001**	0.020
C1	4 (148)	3		0.803	0.782	0.630		0.211	0.217	0.264
C2	17 (148)	11			0.222	0.208			0.412	0.418
C3	23 (148)	16				**3.100**				**0.001**
D2	15 (148)	10								

Table 5 – Purkinje cell responses to single-whisker stimulation in anesthetized mice

Purkinje cells respond with complex firing to single-whisker stimulation (cf. Figure 4). For each whisker, the
number and percentage of responsive cells is indicated (in brackets: total number of cells tested). For each
whisker, we compared the observed rate of convergence on individual Purkinje cells to chance level, using a
bootstrap methods based on the relative prevalence of each stimulus. Indicated are the Z and the p values. All
combinations occurred more often than expected ($Z > 0$). Bold values indicate statistical significance after
Benjamini-Hochberg correction for multiple comparisons.

Next, we examined the spatial distribution of responsive Purkinje cells. In Fig. 7D two nearby

recording spots from the same mouse are shown. In recording spot 1, from which the example

in Fig. 7A originates, all Purkinje cells responded to stimulation of at least one whisker.

However, in the second recording spot, only two Purkinje cells responded: both to a single,

but different whisker. For each recording spot we compared responsive vs. non-responsive

Purkinje cells, taking the average + 3 s.d. of the baseline as threshold for responsiveness. This

yielded a clear separation for the Purkinje cells in the first, but a rather poor one in the second

recording spot (Fig. 7E). In terms of number of responsive Purkinje cells and response

amplitudes, these two recordings, although made from the same lobule in the same animal,

form relatively extreme examples in our dataset. When plotting the response strength versus

the fraction of responsive Purkinje cells per field of view, we found a positive correlation

(linear regression: R = 0.5208; $p < 0.001$; Fig. 7F), implying that Purkinje cells with stronger

responses to a certain whisker tended to be surrounded by other Purkinje cells encoding the

same whisker – in line with the much stronger responses in the first than in the second

recording spot illustrated in Fig. 7D-E. Taken together, stimulating single whiskers revealed a

similar organization as did the less specific stimuli (see Fig. 6), with a clear tendency of

Purkinje cells with the same receptive field to be located close together. However, the spatial

clustering was incomplete and especially weakly responsive Purkinje cells were found to be

interspersed with completely unresponsive Purkinje cells.

Functionally equivalent Purkinje cells fire coherently

In addition to spatial clustering, a second requirement for population encoding is coherence of

complex spike firing. As coherence can be affected by anesthesia, we applied our coherence

analysis on spontaneous and modulation data obtained from Purkinje cells with significant

responses in awake mice. We defined coherence as firing within the same frame of 40 ms,

which still falls well within the subthreshold oscillation cycle of inferior olivary neurons *in*

vivo (55). Pairs of Purkinje cells with significant responses upon whisker pad stimulation

showed a significantly increased level of coherence ($p < 0.001$; two-dimensional

Kolmogorov-Smirnov test) compared to that of heterogeneous pairs (i.e. pairs of one

responsive and one non-responsive Purkinje cell) (Fig. 8A-B). When we examined the firing

pattern of the same pairs in the absence of sensory stimulation, we found similar results (Fig,

8C), indicating that it is not the sensory input per se that directs coherence. These findings

were confirmed in the whole population, taking also the other tactile and visual stimuli into

account (Fig. 8D-E). Only stimulation of upper lip, the area least represented among the

recorded Purkinje cells (Fig. 3), revealed less of a discrimination between significantly

responsive and heterogeneous pairs. Hence, we conclude that not only spatial clustering but

also coherence patterning of functionally equivalent Purkinje cells may facilitate population

encoding.

Figure 8 – Purkinje cells encoding the same response group together

(A) Representative field of view with Purkinje cell dendrites. Purkinje cells showing statistically significant

responses to whisker pad stimulation are depicted in shades of red and the other cells in shades of blue. **(B)** For

each pair of Purkinje cells we calculated the Pearson correlation coefficient (R) during 1 Hz whisker pad

stimulation. The pairs of two Purkinje cells that were both statistically significantly responsive to whisker pad

stimulation (red symbols) had on average a higher level of synchrony than the pairs connecting a responsive

and a non-responsive Purkinje cell (grey symbols; $p < 0.001$; two-dimensional Kolmogorov-Smirnov test). The

pairs consisting of two non-responsive Purkinje cells were excluded from this analysis. **(C)** Interestingly, even in

the absence of sensory stimulation, the pairs of Purkinje cells that were both responsive to whisker pad

stimulation maintained a higher level of synchrony than "heterogeneous pairs". Thus, Purkinje cells with the

same receptive field tended to fire more synchronously, even in the absence of stimulation. This analysis was

expanded in the presence **(D)** and absence **(E)** of sensory stimulation for six different types of stimulation and

illustrated as the median R value per distance category (six bin values of equal distance at a log scale). The

 shaded areas represent the inter-quartile ranges. * $p < 0.05$; *** $p < 0.001$ (two-dimensional Kolmogorov-

Smirnov tests).

Population responses

Although we found increased coherent firing in functionally equivalent Purkinje cells, the

level of coherence was still relatively low, with a regression coefficient seldomly exceeding

0.3 (Fig. 8). However, in this analysis we did not discriminate between complex spikes firing

in response to sensory stimulation and complex spikes occurring spontaneously. Therefore,

we subsequently quantified the distribution of complex spikes over time. This is illustrated for

a representative field of view in Fig. 9A. For each frame, we summed all complex spikes of

the 17 Purkinje cells in this field of view and subsequently made an aggregate peri-stimulus

time histogram of all these Purkinje cells, whereby we color-coded the number of complex

spike recorded per frame. The darker the color, the more complex spikes occurred

simultaneously. Clearly, the darker colors – and thus the stronger coherence – were observed

during the stimulus response period. This occurrence of coherent firing over time was

compared with a random redistribution of the spikes per Purkinje cell (based upon a Poisson

distribution). Note that an equal distribution would imply on average less than two complex

spikes being fired during each frame, indicating the highly patterned distribution found during

the experiments. The grey bars in Fig. 9B indicate the level of coherence that could be

expected by chance, while the red ones indicate highly unlikely values. This shows that

especially the higher levels of coherence are task-related, while spatially isolated firing occurs

irrespective of stimulation.

Figure 9 – Purkinje cells encode strong and weak sensory stimulation via synchronous firing

(A) Aggregate peri-stimulus time histograms (PSTHs) show that coherent firing of complex spikes

predominantly occurs following sensory stimulation. For each field of view, we calculated the number of

complex spikes occurring per frame, summing those of all Purkinje cells in that field of view. Subsequently, we made aggregate PSTHs where the color of each bin refers to the number of dendrites simultaneously active. In this field of view, 17 Purkinje cells were measured. Of these, 17 (100%) reacted to air puff, 12 (71%) to whisker pad, 3 (18%) to upper lip, 4 (24%) to lower lip and 1 (6%) to cheek stimulation. **(B)** Based upon a Poisson distribution of complex spikes over all dendrites and bins, one would expect between 0 and 3 simultaneously active dendrites (grey bars). The red bars indicate events involving more dendrites simultaneously than expected from a random distribution. Thus, the sparse firing as expected by chance is relatively constant throughout the trials, but the simultaneous activity of multiple dendrites is strongly enhanced following sensory stimulation. **(C)** A direct overlay of the aggregate PSTHs in response to air puff and lower lip stimulation shows that the strong response found after air puff stimulation comes at the expense of intertrial complex spikes (152 trials per condition). **(D)** For equally long recordings in the presence of different types of stimulation, equal complex spike frequencies were observed as during spontaneous activity (F(2.544, 20.348) = 2.561, *p* = 0.091, repeated measures ANOVA), indicating that sensory stimulation results in a temporal re-ordering of complex spikes, rather than to the production of more complex spikes.

Examination of the aggregate PSTHs confirms what could already be seen in Fig. 3D, in that there is a trend of reduced intertrial firing for those stimuli with a relatively strong response (Fig. 9). In this experiment, we plotted the responses to air puff stimulation, which recruited statistically significant responses in 17 out of 17 Purkinje cells, on top of those to lower lip stimulation, which recruited only 1 of the 17 Purkinje cells, illustrating the reduced baseline firing during air puff stimulation (Fig. 9C). When we calculated the average complex spike firing for each stimulus condition over the whole population of recorded Purkinje cells, we found that to be remarkably constant. Even 1 Hz air puff stimulation, able to recruit strong complex spike responses, did not result in increased complex spike firing as compared to an epoch without any form of stimulation (Fig. 9D), pointing towards a homeostatic mechanism within the inferior olive that balances out complex spike firing over longer time intervals.

 Homeostasis of complex spike firing

To further study the impact of complex spike homeostasis, we averaged the PSTHs of all

Purkinje cells with statistically significantly responses to air puff stimulation and compared

these to the firing rate in the absence of stimulation (using pseudo-triggers generated at the

same 1 Hz rate). This pair-wise comparison confirmed that the increase in complex spike

firing during the sensory-induced responses comes at the expense of intertrial firing (Fig.

10A). The same was true for the milder whisker pad stimulation. However, because the

responses were weaker, the effect on the intertrial firing was less than that following strong

air puff stimulation (Fig. 10B). The homeostatic effect was also observed following visual

stimulation, although this type of stimulation induced an oscillatory response, making the

effect less visible (Fig. 10C).

Figure 10 – Sensory stimulation results in a temporal re-ordering of complex spikes

The temporal distribution of complex spikes was compared between periods with sensory stimulation and

periods without. For this analysis, we included only Purkinje cells that displayed a significant response to the

stimulus involved (n = 117 for air puff **(A)**, n = 45 for whisker pad **(B)** and n = 27 for visual stimulation **(C)**).

Irrespective of the type of stimulation, a reduction in intertrial activity was found. This was quantified by during

the period between 500 and 250 ms prior to stimulus onset using paired t tests; * and *** indicate $p < 0.05$ and

$p < 0.001$, respectively.

Discussion

Complex spike firing may appear notoriously unpredictable as its spontaneous frequency is

low, yet sustained, and its response rate is at best moderate, with large jitters and an

ambiguous relation to stimulus strength. This raises the question as to how the inferior olive

relays its signals over time. Possibly, coherent complex spike firing by neighboring Purkinje

cells might jointly represent a stimulus, together covering the required signaling for a

particular temporal domain (37, 56-59). Furthermore, the spatial relation between somatotopic

patches and parasagittally oriented microzones has not been clarified in terms of complex

spike signaling, and it is not understood to what extent this relation also depends on the

temporal context in which the signals are generated.

Here, we investigated at the level of individual Purkinje cells whether encoding of

somatosensory input from different facial areas in lobule crus 1 occurs in striped microzones

or instead follows a more fractured arrangement. Mild touches at localized facial areas

revealed a loose version of fractured somatotopy. Purkinje cells with the same receptive field

tended to be located in each other's neighborhood, but the spatial organization was not very

strict, as highly responsive Purkinje cells were sometimes observed amidst non-responsive

ones. The functionally equivalent, adjacent Purkinje cells showed increased coherence in their

complex spike firing, in particular in response to sensory stimulation. Homeostatic

mechanisms are engaged, ensuring that over longer periods complex spike firing rates are

constant, making the short-lived coherent responses more salient.

The functional role of climbing fiber activity

Clinical manifestations of inferior olivary dysfunction range from ataxia and tremor to autism

spectrum disorders (45, 60-64). Climbing fiber-evoked complex spikes are also essential for

the proper timing, size and direction of movements as well as for the encoding of expected

and unexpected deviations from planned movements (6-8, 65-67). In addition, climbing fiber

activity is crucial for cerebellar learning by controlling synaptic plasticity at a wide variety of

synapses in the molecular layer of the cerebellar cortex (68-71). Despite these many

functions, complex spike firing is remarkably stable over longer intervals, suggesting that the

impact of complex spikes is strongly context-dependent.

This context is provided by the synaptic inputs to the inferior olive that relay

information from excitatory ascending and descending pathways as well as inhibitory

projections from the hindbrain (72). Both types converge on each individual spine present on

the dendrites of the inferior olivary neurons (44, 73). The inhibitory input comes

predominantly from the cerebellar nuclei, implying that part of the context is mediated by one

of the target regions of the climbing fibers themselves. This feedback is engaged in a closed

loop, as individual climbing fibers of each olivary subnucleus project to the Purkinje cells

located within a specific parasagittal zone (20) that converge on the neurons in the cerebellar

nuclei that project back to the same olivary subnucleus where the loop started (10, 33, 54).

Each module can be further subdivided into microzones within which complex spike

coherence is clearly enhanced upon strong stimulation (31, 37), a phenomenon that is

enhanced by strong electrotonic coupling within olivary glomeruli (48, 74-76). Strong stimuli

can also increase the coherence of Purkinje cells in adjacent bands (31) while our present data

show that weak stimuli can trigger responses of loosely grouped cells not encompassing a full

microzone. Thus, although anatomical and functional data support the microzonal

organization, the activity patterns are best understood as produced dynamically depending on

the behavioral context that is transmitted via cerebellar and extra-cerebellar synaptic input.

Receptive fields of Purkinje cells and population coding

Several existing maps of somatosensory representations of complex spike activity suggest a

fractured somatotopy (77-79). These maps were typically created by establishing, for each

recording position, the strongest input region, disregarding information on convergence of

multiple inputs. These studies made clear, however, that climbing fibers could have receptive

fields of widely different sizes, expanding to structures as large as a whole limb (80). Part of

this can be explained by the sagittally fanning climbing projections onto multiple lobules (20).

Here we show with cellular resolution within cerebellar microzones that climbing fibers

indeed do convey somatosensory input from different areas, with Purkinje cells having the

same receptive fields being located preferably in each other's proximity.

Our data confirm that increasing stimulus strength promotes complex spike responses,

but its impact might be lower than predicted by other studies (2, 51, 52), showing again the

relevance of the behavioral and historical context for complex spike firing. More salient than

the stimulus strength might be the combination of sensory inputs, in line with the trend of

their convergence on single Purkinje cells. Analyzing the response rates of Purkinje cell

ensembles revealed that the density of responsive cells is crucial for shaping the population

response. We propose that the weak spatial clustering of Purkinje cells encoding the same

stimulus, being interspersed with Purkinje cells receiving input from other sources, and the

tendency of coherent firing of Purkinje cells encoding the same stimulus both contribute to

the creation of a heterogeneous map of Purkinje cells, where each area encodes a particular

functional set of inputs. Each of the properties of the complex spikes seems rather

insignificant in isolation, but in combination may result in specific and robust population

encoding.

Non-tactile inputs

Light or sound can act as conditional stimulus to recruit increasingly more climbing fiber

activity during learning, highlighting the flexibility of this pathway (11, 81). We show here

that, also in naïve animals, visual and auditory input can converge on the same Purkinje cells

that encode somatosensory input, further strengthening the notion that climbing fiber activity

can act as an integrator of contextual input. With visual stimulation the latency was

significantly longer than expected. Probably, the presented visual stimulus is not directly

relayed from the retina, but a descending input from the visual cortex or other higher brain

regions such as the mesodiencephalic junction (72).

Homeostasis of complex spike frequency

The overall firing rate of complex spikes was stable across conditions, with response peaks

being compensated by reduced intertrial firing (see also (16)). The stronger the response peak,

the less intertrial firing, leading to homeostasis of complex spike firing over longer time

periods. Intriguingly, the complex spikes during the intertrial intervals were predominantly

fired by a few, dispersed Purkinje cells and the response peak was largely due to increased

coherence. It seems therefore that Purkinje cells display a basic complex spike firing rate,

which in the absence of functional behavior is not coherent with adjacent Purkinje cells. In

view of the strong impact of complex spikes on synaptic plasticity (68-71), this could

subserve homeostatic functions. Salient stimuli are largely encoded by coherent firing,

making them different from non-stimulus related activity.

Several mechanisms may contribute to the homeostatic control of inferior olivary

spiking, including the previously mentioned olivo-cerebellar loops. Disrupted Purkinje cell

activity can affect climbing fiber activity (82) and reduced inferior olivary activity leads to

enhanced simple spike activity (83), which in turn dampens activity in the GABAergic

neurons of the cerebellar nuclei that controls the inferior olive (84, 85). The interplay between

glutamatergic input to the neurons of the inferior olive, intracellular signaling and electrotonic

coupling may also lead to homeostatic control of complex spike firing via PKA and ßCaMKII

(86, 87). Overall, the inferior olive functions at the cross road of well-defined and rigid

anatomical structures and highly dynamic synaptic input, while subject to homeostatic

control, the sources of which still are partly to be determined.

Materials and Methods

Animals and surgery

All experimental procedures involving animals were in agreement with Dutch and European legislation and guidelines and approved in advance by an independent ethical committee (DEC Consult, Soest, The Netherlands) as required by Dutch law. Mice were group housed until the day of the experiment and kept under a regime with 12 h light and 12 h dark with food and water available *ad libitum*. The mice had not been used for other experiments prior to the ones described here.

For the experiments performed in awake mice, we recorded from in total 66 field of views located in cerebellar lobule crus 1 of 29 male C57Bl/6J mice of 4-12 weeks of age (Charles Rivers, Leiden, The Netherlands). Prior to surgery, mice were anesthetized using isoflurane (initial concentration: 4% V/V in O_2, maintenance concentration: ca. 2% V/V) and received Carprofen (Rimadyl, 5 mg/ml subcutaneously) to reduce post-surgical pain. To prevent dehydration, mice received 1 ml of saline s.c. injection before the surgeries commenced. Eyes were protected using eye ointment (Duratears, Alcon, Fort Worth, TX, USA). Body temperature was maintained using a heating pad in combination with a rectal thermometer. During surgery, we attached a metal head plate to the skull with dental cement (Superbond C&B, Sun Medical Co., Moriyama City, Japan) and made a craniotomy with a diameter of approximately 2 mm centered on the medial part of crus 1 ipsilateral to the side of somatosensory stimulation. The dura mater was preserved and the surface of the cerebellar cortex was cleaned with extracellular solution composed of (in mM) 150 NaCl, 2.5 KCl, 2 $CaCl_2$, 1 $MgCl_2$ and 10 HEPES (pH 7.4, adjusted with NaOH). After the surgery, the mice were allowed to recover from anesthesia for at least 30 minutes. Subsequently, the mice were head-fixed in the recording setup and they received a bolus-loading of the Ca^{2+} indicator Cal-

520 (0.2 mM; AAT Bioquest, Sunnyvale, CA) (88). The dye was first dissolved with 10% w/V Pluronic F-127 in DMSO (Invitrogen, Thermo Fisher Scientific, Waltham, MA, USA) and then diluted 20 times in the extracellular solution. The dye solution was pressure injected into the molecular layer (50–80 μm below the surface) at 0.35 bar for 5 min. Finally, the brain surface was covered with 2% agarose dissolved in saline (0.9% NaCl) in order to reduce motion artifacts and prevent dehydration.

For the experiments on single-whisker stimulation we made recordings under anesthesia on 17 male C57BL/6J mice of 4-12 weeks of age. The procedure was largely the same as described above, but instead of isoflurane we used ketamine/xylazine (i.p. injection via butterfly needle, initial dose: 100 mg/kg and 10 mg/kg, respectively; maintenance dose: approximately 60 mg/kg/h and 3 mg/kg/h, respectively). The mice remained under anesthesia until the end of the recording. A subset of these experiments was performed with 0.2 mM Oregon Green BAPTA-1 AM dye (Invitrogen) as this dye has been more widely used than Cal-520 (e.g., (17, 37, 56, 89). Oregon Green BAPTA-1 AM was dissolved and applied in the same way as Cal-520. We found that Cal-520 had a superior signal-to-noise ratio under our experimental conditions. As a consequence, the observed event rate was lower in the experiments using OGB-1 (OGB-1: 0.45 ± 0.26 Hz; $n = 172$ cells; Cal-520: 0.72 ± 0.40 Hz; $n = 43$ cells; median $\pm$ IQR; $U = 1719.0$, $p < 0.001$, Mann-Whitney test). The observed frequency range using Cal-520 was comparable to that found using *in vivo* single-unit recordings under ketamine/xylazine anesthesia: 0.6 ± 0.1 Hz (2). Despite an underestimation of the complex spike rate using OGB-1, we found that the ratios of Purkinje cells that responded to single whisker stimulation were similar for both dyes (OGB-1: 89 out of 373 cells (24%); Cal-520: 35 out of 152 cells (23%); $p = 0.910$; Fisher's exact test). For this reason, we combined the data from both dyes. For the analysis presented in Fig. 7, we

included only those cells that could be recorded during all stimulus conditions. All

experiments using awake data were obtained with Cal-520.

In vivo *two-photon Ca^{2+} imaging*

Starting at least 30 min after dye injection, *in vivo* two-photon Ca^{2+} imaging was performed of

the molecular layer of crus 1 using a setup consisting of a Ti:Sapphire laser (Chameleon

Ultra, Coherent, Santa Clara, CA, USA), a TriM Scope II system (LaVisionBioTec, Bielefeld,

Germany) mounted on a BX51 microscope with a 20x 1.0 NA water immersion objective

(Olympus, Tokyo, Japan) and GaAsP photomultiplier detectors (Hamamatsu, Iwata City,

Japan). A typical recording sampled a field of view of 40 x 200 μm with a frame rate of

approximately 25 Hz.

Sensory stimulation

Cutaneous stimuli were delivered to four defined regions on the left side of the face,

ipsilateral to side of the craniotomy. These regions were the whisker pad, the cheek posterior

to the whisker pad, the upper lip and the lower lip. Stimuli were applied using a Von Frey

filament (Touch Test Sensory Evaluator 2.83, Stoelting Co., IL, USA) attached to a piezo

linear drive (M-663, Physik Instrumente, Karlsruhe, Germany). Prior to the set of experiments

described here, we tested a series of 8 Von Frey filaments with a stiffness range from 0.02 g

to 1.4 g in awake head-fixed mice to select the optimal force for these experiments. We

selected the 0.07 g (0.686 mN) filament because this filament induced a mild reaction in the

mouse, but no signs of a nociceptive response (cf. Chaplan et al. (90)). The touch time was

fixed at 100 ms. As a control, we moved the stimulator without touching the face ("sound

only" condition). Visual stimuli were delivered as 10 ms pulses using a 460 nm LED (L-

7104QBC-D, Kingbright, CA, USA). The stimulation frequency was fixed at 1 Hz and the

different stimuli were applied in a random order. Video-recordings of the eye, made under

infrared illumination, revealed that the mice did not make eye movements in response to the

LED flash, but they did show reflexive pupil constriction (*data not shown*).

Single whiskers were stimulated using a piezo linear drive (M-663, Physik Instrumente)

while using a deflection of 6°. It took the stimulator approximately 30 ms to reach this

position. In our analyses, we used the estimated moment of touch as the stimulus onset time.

At the extreme position, the stimulator was paused for 150 ms before returning to the neutral

position. The stimulator was designed to minimize contact with other whiskers. Each

stimulation experiment consisted of five sessions in random order. During each block one of

the ipsilateral whiskers B2, C1, C2, C3 and D2 was stimulated. Each session consisted of 150

stimuli at 2 or 3 Hz.

Complex spike detection

Image analysis was performed offline using custom made software as described and validated

previously (46, 48, 91). In short, independent component analysis was applied to the image

stack to discover masks describing the locations of individual Purkinje cell dendrites (e.g., see

Fig. 2A). For each field of view the mask was generated only once, so that the same Purkinje

cells were analyzed for subsequent recordings of different stimulus conditions enabling paired

comparisons. Experiments during which spatial drift occurred were discarded from

subsequent analysis. The fluorescence values of all pixels in each mask were averaged per

frame. An 8% rolling baseline from a time window of 0.5 ms was subtracted from the average

fluorescence per mask (Ozden et al. 2012), after which Ca^{2+} transient events were detected

using template matching.

Statistical analysis

In general, we first tested whether parameter values were distributed normally using one-

sample Kolmogorov-Smirnov tests. If not, non-parametric tests were applied. When making

multiple comparisons, we used ANOVA tests (or the non-parametric equivalents) or used

multiple tests. In the latter case, the method of correction for multiple comparisons is

mentioned in the figure legends. For each experiment, stimuli were given in a random

sequence. Identification of Purkinje cell dendrites and event extraction were performed by a

researcher who was blind to the type of stimulus. For the experiments characterizing response

characteristics of individual Purkinje cells, we compared the fraction of responsive Purkinje

cells, the response latency and the response peak. After extracting the Ca^{2+} transient event

times, peri-stimulus time histograms (PSTHs) were constructed using the inter-frame time

(approx. 40 ms) as bin size. Stacked line plots of Purkinje cell PSTHs were sorted by weak to

strong peak responses. The data were normalized such that the top represents the average

responses of all Purkinje cells. Statistical significance of responses occurring within 200 ms

from stimulus onset was evaluated using a threshold of the mean + 3 s.d. of the firing rate

during the 500 ms prior to stimulus onset (300 ms for the single whisker stimulation

experiments, as they were performed with a higher stimulation frequency).

To calculate whether two or more inputs converged on single Purkinje cells we used a

bootstrap method. First, we determined the fraction of responsive cells for each parameter and

compared these to a randomly generated number between 0 and 1 as taken from a uniform

distribution. If for each input the randomly generated numbers were lower than the measured

fractions, we considered them as responsive to all stimuli. This procedure was repeated

10,000 times and the average and standard deviation were derived and used to calculate the *Z*

score of the experimental data. All bootstrap procedures were performed using custom-written

code in LabVIEW (National Instruments, Austin, TX, USA).

The distributions of pairs of Purkinje cells, either both responsive to a given stimulus

("responsive pairs") or one cell being responsive and the other not ("heterogeneous pairs"),

were tested using two-dimensional Kolmogorov-Smirnov tests performed in MATLAB

(MathWorks, Natick, MA, USA). Aggregate PSTHs (Fig. 9) were constructed and evaluated

as described before (4). Briefly, we calculated per individual frame the number of

simultaneously occurring events and color coded these in a PSTH combining data from all

dendrites in a field of view. Based upon the total number of complex spikes and dendrites per

recording, we calculated the expected number of simultaneous complex spikes per individual

frame based upon a Poisson distribution. The actual number of simultaneous complex spikes

was compared to this calculated distribution and a p value was derived for each number based

upon the Poisson distribution (using custom-written software in MATLAB and LabVIEW).

Unless stated otherwise, statistical tests were performed using SPSS (IBM, Armonk,

NY, USA) and data are represented in the text as average ± SEM.

Acknowledgements

The authors wish to thank A.C.H.G. IJpelaar for technical assistance and Dr. M. Schonewille for help with recording ocular movements during stimulation. Financial support was provided by the Netherlands Organization for Scientific Research (NWO-ALW; C.I.D.Z.), the Dutch Organization for Medical Sciences (ZonMW; C.I.D.Z.), Life Sciences (C.I.D.Z.), ERC-adv and ERC-POC (C.I.D.Z.) and the China Scholarship Council (No. 2010623033; C.J.). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1.   Albus JS. Theory of cerebellar function. Math Biosci. 1971;10:25-61.

2.   Bosman LWJ, Koekkoek SKE, Shapiro J, Rijken BFM, Zandstra F, van der Ende B, et al.

Encoding of whisker input by cerebellar Purkinje cells. J Physiol. 2010;588(19):3757-83.

3.   Gibson AR, Horn KM, Pong M. Activation of climbing fibers. Cerebellum. 2004;3(4):212-21.

4.   Romano V, De Propris L, Bosman LWJ, Warnaar P, Ten Brinke MM, Lindeman S, et al.

Adaptation of whisker movements requires cerebellar potentiation. bioRxiv. 2018:335489.

5.   Streng ML, Popa LS, Ebner TJ. Climbing fibers control Purkinje cell representations of

behavior. J Neurosci. 2017;37(8):1997-2009.

6.   Yang Y, Lisberger SG. Purkinje-cell plasticity and cerebellar motor learning are graded by

complex-spike duration. Nature. 2014;510(7506):529-32.

7.   Herzfeld DJ, Kojima Y, Soetedjo R, Shadmehr R. Encoding of error and learning to correct that

error by the Purkinje cells of the cerebellum. Nat Neurosci. 2018;21(5):736-43.

8.   Kitazawa S, Kimura T, Yin PB. Cerebellar complex spikes encode both destinations and errors

in arm movements. Nature. 1998;392(6675):494-7.

9.   Welsh JP, Lang EJ, Suglhara I, Llinas R. Dynamic organization of motor control within the

olivocerebellar system. Nature. 1995;374(6521):453-7.

10.   Apps R, Hawkes R, Aoki S, Bengtsson F, Brown AM, Chen G, et al. Cerebellar modules and

their role as operational cerebellar processing units. Cerebellum. 2018.

11.   Ten Brinke MM, Boele HJ, Spanke JK, Potters JW, Kornysheva K, Wulff P, et al. Evolving

models of Pavlovian conditioning: cerebellar cortical dynamics in awake behaving mice. Cell Rep.

2015;13(9):1977-88.

12.   Junker M, Endres D, Sun ZP, Dicke PW, Giese M, Thier P. Learning from the past: A

reverberation of past errors in the cerebellar climbing fiber signal. PLoS Biol. 2018;16(8):e2004344.

13.   Zhou H, Lin Z, Voges K, Ju C, Gao Z, Bosman LWJ, et al. Cerebellar modules operate at

different frequencies. eLIFE. 2014;3:e02536.

14.   Mukamel EA, Nimmerjahn A, Schnitzer MJ. Automated analysis of cellular signals from large-

scale calcium imaging data. Neuron. 2009;63(6):747-60.

15.   Bloedel JR, Ebner TJ. Rhythmic discharge of climbing fibre afferents in response to natural

peripheral stimuli in the cat. J Physiol. 1984;352:129-46.

16.   Warnaar P, Negrello M, Romano V, Owens CB, Lindeman S, Iavarona E, et al. Aperiodic

rhythms of the inferior olive. bioRxiv. 2018:408112.

17.   Hoogland TM, De Gruijl JR, Witter L, Canto CB, De Zeeuw CI. Role of synchronous activation

of cerebellar Purkinje cell ensembles in multi-joint movement control. Curr Biol. 2015;25(9):1157-65.

18.   Harvey RJ, Napper RMA. Quantitative studies on the mammalian cerebellum. Prog Neurobiol.

1991;36(6):437-63.

19.   Person AL, Raman IM. Purkinje neuron synchrony elicits time-locked spiking in the cerebellar

nuclei. Nature. 2011;481(7382):502-5.

20.   Sugihara I, Wu HS, Shinoda Y. The entire trajectories of single olivocerebellar axons in the

cerebellar cortex and their contribution to cerebellar compartmentalization. J Neurosci.

2001;21(19):7715-23.

21.   Cerminara NL, Lang EJ, Sillitoe RV, Apps R. Redefining the cerebellar cortex as an assembly of

non-uniform Purkinje cell microcircuits. Nat Rev Neurosci. 2015;16(2):79-93.

22.   Ruigrok TJH. Ins and outs of cerebellar modules. Cerebellum. 2011;10(3):464-74.

23.   Szentágothai J. The use of degeneration methods in the investigation of short neuronal

connexions. Prog Brain Res. 1965;14:1-32.

24.   Andersen P, Eccles JC, Voorhoeve PE. Postsynaptic inhibition of cerebellar Purkinje cells. J

Neurophysiol. 1964;27:1138-53.

25.   Gao W, Chen G, Reinert KC, Ebner TJ. Cerebellar cortical molecular layer inhibition is

organized in parasagittal zones. J Neurosci. 2006;26(32):8377-87.

26.   Sullivan MR, Nimmerjahn A, Sarkisov DV, Helmchen F, Wang SSH. In vivo calcium imaging of

circuit activity in cerebellar cortex. J Neurophysiol. 2005;94(2):1636-44.

27.   Sugihara I, Wu H, Shinoda Y. Morphology of single olivocerebellar axons labeled with

biotinylated dextran amine in the rat. J Comp Neurol. 1999;414(2):131-48.

28.   Sugihara I, Fujita H, Na J, Quy PN, Li BY, Ikeda D. Projection of reconstructed single Purkinje

cell axons in relation to the cortical and nuclear aldolase C compartments of the rat cerebellum. J

Comp Neurol. 2009;512(2):282-304.

29.   Xiao J, Cerminara NL, Kotsurovskyy Y, Aoki H, Burroughs A, Wise AK, et al. Systematic regional

variations in Purkinje cell spiking patterns. PLoS One. 2014;9(8):e105633.

30.   De Zeeuw CI, Ten Brinke MM. Motor learning and the cerebellum. Cold Spring Harb Perspect

Biol. 2015;7(9):a021683.

31.   Tsutsumi S, Yamazaki M, Miyazaki T, Watanabe M, Sakimura K, Kano M, et al. Structure-

function relationships between aldolase C/zebrin II expression and complex spike synchrony in the

cerebellum. J Neurosci. 2015;35(2):843-52.

32.   Suvrathan A, Payne HL, Raymond JL. Timing rules for synaptic plasticity matched to

behavioral function. Neuron. 2016;92(5):959-67.

33.   Groenewegen HJ, Voogd J, Freedman SL. The parasagittal zonation within the olivocerebellar

projection. II. Climbing fiber distribution in the intermediate and hemispheric parts of cat cerebellum.

J Comp Neurol. 1979;183(3):551-601.

34.   Oscarsson O. Termination and functional organization of the dorsal spino-olivocerebellar

path. J Physiol. 1969;200(1):129-49.

35.   Ekerot CF, Garwicz M, Schouenborg J. Topography and nociceptive receptive fields of

climbing fibres projecting to the cerebellar anterior lobe in the cat. J Physiol. 1991;441:257-74.

36.   Apps R, Garwicz M. Anatomical and physiological foundations of cerebellar information

processing. Nat Rev Neurosci. 2005;6(4):297-311.

37.   Ozden I, Sullivan MR, Lee HM, Wang SSH. Reliable coding emerges from coactivation of

climbing fibers in microbands of cerebellar Purkinje neurons. J Neurosci. 2009;29(34):10463-73.

38.   De Zeeuw CI, Hoebeek FE, Bosman LWJ, Schonewille M, Witter L, Koekkoek SK.

Spatiotemporal firing patterns in the cerebellum. Nat Rev Neurosci. 2011;12(6):327-44.

39.   Yatim N, Billig I, Compoint C, Buisseret P, Buisseret-Delmas C. Trigeminocerebellar and

trigemino-olivary projections in rats. Neurosci Res. 1996;25(3):267-83.

40.   Apps R, Hawkes R. Cerebellar cortical organization: a one-map hypothesis. Nat Rev Neurosci.

2009;10(9):670-81.

41.   Shambes GM, Gibson JM, Welker W. Fractured somatotopy in granule cell tactile areas of rat

cerebellar hemispheres revealed by micromapping. Brain Behav Evol. 1978;15(2):94-140.

42.   Bosman LWJ, Houweling AR, Owens CB, Tanke N, Shevchouk OT, Rahmati N, et al. Anatomical

pathways involved in generating and sensing rhythmic whisker movements. Front Integr Neurosci.

2011;5:53.

43.   Kubo R, Aiba A, Hashimoto K. The anatomical pathway from the mesodiencephalic junction

to the inferior olive relays perioral sensory signals to the cerebellum in the mouse. J Physiol. 2018.

44.   De Zeeuw CI, Holstege JC, Ruigrok TJH, Voogd J. Mesodiencephalic and cerebellar terminals

terminate upon the same dendritic spines in the glomeruli of the cat and rat inferior olive: an

ultrastructural study using a combination of [3H]leucine and wheat germ agglutinin coupled

horseradish peroxidase anterograde tracing. Neuroscience. 1990;34(3):645-55.

45.   De Gruijl JR, Bosman LWJ, De Zeeuw CI, De Jeu MTG. Inferior olive: All ins and outs. In: Manto

M, Schmahmann JD, Rossi F, Gruol DL, Koibuchi N, editors. Handbook of the Cerebellum and

Cerebellar Disorders. 3. Dordrecht: Springer Netherlands; 2013. p. 1013-58.

46.   Ozden I, Lee HM, Sullivan MR, Wang SSH. Identification and clustering of event patterns from

in vivo multiphoton optical recordings of neuronal ensembles. J Neurophysiol. 2008;100(1):495-503.

47.   Najafi F, Medina JF. Beyond "all-or-nothing" climbing fibers: graded representation of

teaching signals in Purkinje cells. Front Neural Circuits. 2013;7:115.

48.   De Gruijl JR, Hoogland TM, De Zeeuw CI. Behavioral correlates of complex spike synchrony in

cerebellar microzones. J Neurosci. 2014;34(27):8937-44.

49.   Brown IE, Bower JM. The influence of somatosensory cortex on climbing fiber responses in

the lateral hemispheres of the rat cerebellum after peripheral tactile stimulation. J Neurosci.

2002;22(15):6819-29.

50.   Axelrad H, Crepel F. Représentation sélective des vibrisses mystaciales au niveau des cellules

de Purkinje du cervelet par la voie des fibres grimpantes chez le rat. C R Acad Sci Hebd Seances Acad

Sci D. 1977;284(14):1321-4.

51.   Eccles JC, Sabah NH, Schmidt RF, Táboríková H. Cutaneous mechanoreceptors influencing

impulse discharges in cerebellar cortex. III. In Purkyně cells by climbing fiber input. Exp Brain Res.

1972;15(5):484-97.

52.   Najafi F, Giovannucci A, Wang SSH, Medina JF. Coding of stimulus strength via analog calcium

signals in Purkinje cell dendrites of awake mice. eLIFE. 2014;3:e03663.

53.   Arabzadeh E, Panzeri S, Diamond ME. Whisker vibration information carried by rat barrel

cortex neurons. J Neurosci. 2004;24(26):6011-20.

54.   Voogd J, Glickstein M. The anatomy of the cerebellum. Trends Neurosci. 1998;21(9):370-5.

55.   Khosrovani S, Van Der Giessen RS, De Zeeuw CI, De Jeu MTG. *In vivo* mouse inferior olive

neurons exhibit heterogeneous subthreshold oscillations and spiking patterns. Proc Natl Acad Sci U S

A. 2007;104(40):15911-6.

56.   Schultz SR, Kitamura K, Post-Uiterweer A, Krupic J, Häusser M. Spatial pattern coding of

sensory information by climbing fiber-evoked calcium signals in networks of neighboring cerebellar

Purkinje cells. J Neurosci. 2009;29(25):8005-15.

57.   Sasaki K, Bower JM, Llinás R. Multiple Purkinje cell recording in rodent cerebellar cortex. Eur J

Neurosci. 1989;1(6):572-86.

58.   Lang EJ, Sugihara I, Welsh JP, Llinás R. Patterns of spontaneous purkinje cell complex spike

activity in the awake rat. J Neurosci. 1999;19(7):2728-39.

59.   Sugihara I, Marshall SP, Lang EJ. Relationship of complex spike synchrony bands and climbing

fiber projection determined by reference to aldolase C compartments in crus IIa of the rat cerebellar

cortex. J Comp Neurol. 2007;501(1):13-29.

60.   Samuel M, Torun N, Tuite PJ, Sharpe JA, Lang AE. Progressive ataxia and palatal tremor

(PAPT): clinical and MRI assessment with review of palatal tremors. Brain. 2004;127(Pt 6):1252-68.

61.   Llinás R, Walton K, Hillman DE, Sotelo C. Inferior olive: its role in motor learing. Science.

1975;190(4220):1230-1.

62.   Bauman ML, Kemper TL. Neuroanatomic observations of the brain in autism: a review and

future directions. Int J Dev Neurosci. 2005;23(2-3):183-7.

63.   Lim CCT, Lim SA. Pendular nystagmus and palatomyoclonus from hypertrophic olivary

degeneration. New Engl J Med. 2009;360(9):e12.

64.   Welsh JP, Ahn ES, Placantonakis DG. Is autism due to brain desynchronization? Int J Dev

Neurosci. 2005;23(2-3):253-63.

65.   Ito M. Error detection and representation in the olivo-cerebellar system. Front Neural

Circuits. 2013;7:1.

66.   Simpson JI, Wylie DR, De Zeeuw CI. On climbing fiber signals and their consequence(s). Behav

Brain Sci. 1996;19(2):384-98.

67.   Wang JJ, Kim JH, Ebner TJ. Climbing fiber afferent modulation during a visually guided, multi-

joint arm movement in the monkey. Brain Res. 1987;410(2):323-9.

68.   Ito M, Kano M. Long-lasting depression of parallel fiber-Purkinje cell transmission induced by

conjunctive stimulation of parallel fibers and climbing fibers in the cerebellar cortex. Neurosci Lett.

1982;33(3):253-8.

69.   Ito M. Long-term depression. Annu Rev Neurosci. 2003;12:85-102.

70.   Coesmans M, Weber JT, De Zeeuw CI, Hansel C. Bidirectional parallel fiber plasticity in the

cerebellum under climbing fiber control. Neuron. 2004;44(4):691-700.

71. Gao Z, Van Beugen BJ, De Zeeuw CI. Distributed synergistic plasticity and cerebellar learning.

Nat Rev Neurosci. 2012;13(9):619-35.

72. De Zeeuw CI, Simpson JI, Hoogenraad CC, Galjart N, Koekkoek SKE, Ruigrok TJH.

Microcircuitry and function of the inferior olive. Trends Neurosci. 1998;21(9):391-400.

73. De Zeeuw CI, Holstege JC, Ruigrok TJ, Voogd J. Ultrastructural study of the GABAergic,

cerebellar, and mesodiencephalic innervation of the cat medial accessory olive: anterograde tracing

combined with immunocytochemistry. J Comp Neurol. 1989;284(1):12-35.

74. Sotelo C, Llinás R, Baker R. Structural study of inferior olivary nucleus of the cat:

morphological correlates of electrotonic coupling. J Neurophysiol. 1974;37(3):541-59.

75. Ruigrok TJH, de Zeeuw CI, van der Burg J, Voogd J. Intracellular labeling of neurons in the

medial accessory olive of the cat: I. Physiology and light microscopy. J Comp Neurol.

1990;300(4):462-77.

76. Devor A, Yarom Y. Electrotonic coupling in the inferior olivary nucleus revealed by

simultaneous double patch recordings. J Neurophysiol. 2002;87(6):3048-58.

77. Miles TS, Wiesendanger M. Organization of climbing fibre projections to the cerebellar cortex

from trigeminal cutaneous afferents and from the SI face area of the cerebral cortex in the cat. J

Physiol. 1975;245(2):409-24.

78. Castelfranco AM, Robertson LT, McCollum G. Detail, proportion, and foci among face

receptive fields of climbing fiber responses in the cat cerebellum. Somatosens Mot Res.

1994;11(1):27-46.

79. Rushmer DS, Woollacott MH, Robertson LT, Laxer KD. Somatotopic organization of climbing

fiber projections from low threshold cutaneous afferents to pars intermedia of cerebellar cortex in

the cat. Brain Res. 1980;181(1):17-30.

80. Thach WT, Jr. Somatosensory receptive fields of single units in cat cerebellar cortex. J

Neurophysiol. 1967;30(4):675-96.

81.   Ohmae S, Medina JF. Climbing fibers encode a temporal-difference prediction error during

cerebellar learning in mice. Nat Neurosci. 2015;18(12):1798-803.

82.   Chen X, Kovalchuk Y, Adelsberger H, Henning HA, Sausbier M, Wietzorrek G, et al. Disruption

of the olivo-cerebellar circuit by Purkinje neuron-specific ablation of BK channels. Proc Natl Acad Sci

U S A. 2010;107(27):12323-8.

83.   Montarolo PG, Palestini M, Strata P. The inhibitory effect of the olivocerebellar input on the

cerebellar Purkinje cells in the rat. J Physiol. 1982;332:187-202.

84.   Chaumont J, Guyon N, Valera AM, Dugué GP, Popa D, Marcaggi P, et al. Clusters of cerebellar

Purkinje cells control their afferent climbing fiber discharge. Proc Natl Acad Sci U S A.

2013;110(40):16223-8.

85.   Witter L, Canto CB, Hoogland TM, de Gruijl JR, De Zeeuw CI. Strength and timing of motor

responses mediated by rebound firing in the cerebellar nuclei after Purkinje cell activation. Front

Neural Circuits. 2013;7:133.

86.   Mathy A, Clark BA, Häusser M. Synaptically induced long-term modulation of electrical

coupling in the inferior olive. Neuron. 2014;81(6):1290-6.

87.   Bazzigaluppi P, Isenia SC, Haasdijk ED, Elgersma Y, De Zeeuw CI, van der Giessen RS, et al.

Modulation of murine olivary connexin 36 gap Junctions by PKA and CaMKII. Front Cell Neurosci.

2017;11:397.

88.   Tada M, Takeuchi A, Hashizume M, Kitamura K, Kano M. A highly sensitive fluorescent

indicator dye for calcium imaging of neural activity *in vitro* and *in vivo*. Eur J Neurosci.

2014;39(11):1720-8.

89.   Stosiek C, Garaschuk O, Holthoff K, Konnerth A. *In vivo* two-photon calcium imaging of

neuronal networks. Proc Natl Acad Sci U S A. 2003;100(12):7319-24.

90.   Chaplan SR, Bach FW, Pogrel JW, Chung JM, Yaksh TL. Quantitative assessment of tactile

allodynia in the rat paw. J Neurosci Methods. 1994;53(1):55-63.

91.   Ozden I, Dombeck DA, Hoogland TM, Tank DW, Wang SS. Widespread state-dependent shifts

in cerebellar activity in locomoting mice. PLoS One. 2012;7(8):e42650.

    Supporting Information - Supplementary figure legends

Figure S1 - Complex spike detection

(A) Enlarged view of the boxed area in Fig. 2B showing the time points of complex spike

firing (indicated with grey lines below each trace). **(B)** Of the same experiment, we randomly

selected four Purkinje cells of which all trials with (top) and without (bottom) a complex

spike within 200 ms of stimulus onset are shown. The signals are scaled to the maximum of

the median complex spike responses. The fat colored lines indicate the medians with the same

color code as in Fig. 2. Trials without a complex spike response can still show an increased

fluorescence, but the kinetics of the non-complex spike response differed from those of the

complex spike responses. Note that the raw trace of cell 5 had a period with increased noise

levels, making it the least reliable cell of this recording. Nevertheless, cell 5 had a complex

spike response profile that was very similar to those of the other cells (see dashed line in Fig.

2F). Cells (in other recordings) that had a worse signal-to-noise ratio than cell 5 of this

recording were excluded from analysis.

1009

Figure S2 – Purkinje cells in crus 1 respond to various types of sensory stimulation

(A) Stacked line plots illustrating the peri-stimulus time histograms of all Purkinje cells

recorded under either of the six indicated stimulus conditions (see also Fig. 3A). The cells are

ordered based upon their peak responses (calculated as Z value) during the 200 ms interval

following the stimulus onset, with the cell with the lowest response at the bottom of each

graph. The grey lines indicate cells with a peak response deemed not significant ($Z < 3$), the

colored lines indicate the significant responses ($Z > 3$). The graphs are normalized so that the

upper line depicts the averge of all cells. As shown in Fig. 3B, one cannot discriminate

between responsive and non-responsive cells in a black-and-white fashion. Instead, the cells

form a continuum from not responsive at all to highly responsive. As this way of plotting

relies on the numerical average, a skewed distribution can put emphasis on a relatively small

group of cells. As the peak amplitude distributions indeed have a skewed distribution, they are

also compared using cumulative histograms **(B)** using the same color scheme as in A. From

this representation, it is confirmed that whisker pad and lower lip stimulation yield the

strongest responses, while visual stimulation (LED) recruits a few cells with a relatively

strong response, increasing the numerical average (see **A**). Here, we tested the complete

distributions (*** $p < 0.001$; Kruskal-Wallis test). Pair-wise comparisons of all stimulus

conditions are presented in Fig. 4.

1028

Figure S3 – Sound only stimulation systematically recruited less complex spikes than

tactile stimulation

Fluorescent traces of 26 Purkinje cells in a field of view during whisker pad **(A)** and sound

only **(B)** stimulation. The moments of stimulation (at 1 Hz) are indicated by the vertical lines.

Note that the sound only stimulation involved the sound of the mechanical device delivering

tactile stimuli. Overall, whisker pad stimulation triggered more complex spike responses than

sound only as illustrated by the sum of the events **(C)**. The boxed part (10 s) is enlarged in **D**.

(E) Peri-stimulus time histograms (PSTHs) of four randomly selected Purkinje cells from the

experiment illustrated in **A** and **B**. In each, the response to whisker pad stimulation was

stronger than to sound only stimulation, which is also reflected in the median of the PSTHs of

all 26 Purkinje cells **(F)** and the median difference between whisker pad and sound only

stimulation **(G)**. The shaded areas indicate the inter-quartile range.

1041

Figure S4 – Stimulus strength has only a minor impact on complex spike responsiveness

(A) Larger view of the data represented in Fig. 5E. Shown are the average complex spike

rates during the frame with the strongest response per stimulus condition. The Purkinje cells

that showed increased responsiveness to stronger stimuli are depicted in red and the others in

blue. **(B)** The average number of complex spikes per frame (of 40 ms) per trial (shaded areas:

± SEM) for the three stimulus strengths show little difference for the weak and moderate

stimulation. The time course and amplitude (1-4 mm) of the three stimuli is schematized at

the bottom of the graph. In contrast to Fig. 6, in this plot all 340 Purkinje cells are included,

whether or not they displayed a significant response to the stimulus.

Figure 1 – Sensory pathways carrying facial input to the cerebellar cortex

(A) Scheme of the main routes conveying facial tactile input via the climbing fiber pathway to cerebellar Purkinje cells (PC). Climbing fibers, which cause complex spike firing in Purkinje cells, exclusively originate from the inferior olive. The inferior olive, in turn, is directly innervated by neurons from the trigeminal nuclei as well as indirectly via thalamo-cortical pathways that project to the inferior olive mainly via the nuclei of the mesodiencephalic junction (MDJ). The MDJ itself also receives direct input from the trigeminal nuclei. See the main text for references. **(B)** *In vivo* two-photon Ca²⁺ imaging was performed to characterize Purkinje cell complex spike responses to sensory stimulation in the medial part of crus 1. Purkinje cells were detected using independent component analysis and the position of a Purkinje cell dendrite (yellow area on the right) within a field of view is shown in the inset. At the end of each recording session, the brain was removed and the location of the dye injection in medial crus 1 was confirmed through *ex vivo* epifluorescent imaging (black circle). The white rectangle indicates the approximate recording location. **(C)** Complex spikes that were triggered by climbing fiber activity were retrieved from fluorescent traces of individual Purkinje cells. A representative trace obtained from the Purkinje cell dendrite illustrated in **B** is shown together with the detected complex spikes (grey lines). The light blue episode is enlarged in **D**. Complex spikes were detected by the combination of a threshold and a template matching algorithm. Only events with a sharp rising phase were accepted as complex spikes. In the 60 s interval shown in **C**, there was one event with a slower rise time (see arrow in **D**), as indicated at a larger time scale in **E**. The events in **E** are scaled to peak.

Figure 2 - Whisker stimulation evokes complex spike responses in cerebellar crus 1

(A) Complex spikes elicit large increases in the Ca²⁺ concentration within Purkinje cell dendrites that can be resolved using *in vivo* two-photon microscopy in combination with a fluorescent Ca²⁺ indicator. An example of a field of view with 19 identified Purkinje cell dendrites located in the medial part of crus 1 is shown with each individual dendrite denoted by a number and a unique color. This recording was made in an awake mouse. **(B)** The fluorescent traces of each of these dendrites show distinct Ca²⁺ transient events, which are to a large extent associated with air puff stimulation of the facial whiskers (times of stimulation indicated by the vertical lines). The boxed area is enlarged in Fig. S1. **(C)** Summed fluorescence trace composed of all 19 individual traces emphasizing the participation of many Purkinje cells to the stimulus-triggered responses. **(D)** After complex spike extraction, a clear relation between stimulus and activity was observed as illustrated by summing, for each frame, the number of complex spikes observed over all dendrites. The vertical scale bar corresponds to the simultaneous activity of 5 Purkinje cells. **(E)** Peri-stimulus time histogram of a Purkinje cell dendrite (marked as number 7 in **A**

and **B**) in response to air puff stimulation to the ipsilateral whiskers (154 trials). The bin size (40 ms) corresponds to the acquisition frame rate (25 Hz). **(F)** Normalized stacked line graph of the Purkinje cells in this field of view showing that every cell contributed to the overall response. The Purkinje cells are ranked by their maximal response and the data are normalized so that the top line reflects the average frequency per bin. Cell no. 7 (dashed line) had a relatively poor signal-to-noise ratio during later parts of the recording (see Fig. S1), but it had nevertheless a complex spike response profile that was indistinguishable from the other cells. The colors match those in the panels **A**-**B**. Inset: In total, 102 out of 118 cells analyzed (86%) were responsive to whisker air puff stimulation (peak response exceeded average + 3 s.d. of pre-stimulus interval). **(G)** Fluorescent traces of the trials with (red) and without (black) complex spikes fired during the first 200 ms after air puff onset. In the absence of complex spike firing, only a very small increase in fluorescence was observed, indicating that the majority of change in fluorescence was associated with complex spike firing. Note the longer time scale than in **E** and **F**. The lines indicate the medians and the shaded areas the inter-quartile ranges of the 19 Purkinje cells in this field of view.

Figure 3 – Purkinje cells in crus 1 respond to various types of sensory stimulation

(A) Tactile stimuli were presented to four facial regions in awake mice: the whisker pad (WP), the upper lip (UL), the lower lip (LL) and the cheek (Ch). Each stimulus was delivered with a piezo-actuator that made a muted, yet audible sound which was also delivered without touch ("sound only (SO)"). A blue light flash generated by an LED was used as visual stimulus. These experiments were performed in awake mice. **(B)** To avoid interference of adjacent areas, we applied gentle touches (0.686 mN). The complex spike response ratio was much reduced relative to the strong air puff stimulation to all ipsilateral whiskers illustrated in Fig. 2. A histogram of the peak responses (expressed as Z value) of all responses to either of the four tactile stimuli demonstrates that the response strength is a continuum, showing the lack of a clear separation between "responsive" and "non-responsive" Purkinje cells (998 stimulus conditions in 282 Purkinje cells). We considered Purkinje cells that showed a peak response above Z = 3 as "significantly responsive" (represented with black bars), but we provide most of the analyses also for the population as a whole

(e.g., Fig. S2). **(C)** The peri-stimulus time histograms (PSTHs) of a representative Purkinje cell. The shades of grey indicate 1, 2 and 3 s.d. around the average. Each stimulus was repeated 154 times at 1 Hz. **(D)** For every stimulus condition, we averaged the PSTHs for all Purkinje cells that were significantly responsive to that particular stimulus (colored lines; averages ± SEM). These were contrasted to the averaged PSTH of the other Purkinje cells (black lines; averages ± SEM). The pie charts represent the fraction of Purkinje cells significantly responsive to a particular stimulus. See also Table 1. **(E)** The peak responses of the significantly responding Purkinje cells were the lowest for sound only and for upper lip stimulation. * $p < 0.05$; ** $p < 0.01$ (post-hoc tests after Kruskal-Wallis test) **(F)** As expected for complex spike responses to weak stimulation, the latencies were relatively long and variable, but consistent across types of stimulation. Only visual stimulation (LED) had a remarkably longer latency time. *** $p < 0.001$ (post-hoc tests after Kruskal-Wallis test for LED vs. whisker pad, upper lip, lower lip and cheek and $p < 0.05$ compared to sound only)

Figure 4 – Convergence of sensory input on Purkinje cells

(A) In order to test whether sensory inputs converge on individual Purkinje cells in awake mice, we made pair-wise comparisons of the response amplitudes to two different stimuli per Purkinje cell (scatter plots). For all possible combinations, we found a positive slope of the linear regression analysis. For the majority of combinations, the correlation between response strengths was highly significant: * *p* < 0.05, ** *p* < 0.01 and *** *p* < 0.001 (Pearson correlation with Benjamini-Hochberg correction for multiple comparisons). Only upper lip vs. sound only and upper lip vs. visual stimulation were not significantly correlated. For this analysis, we included all Purkinje cells, whether they had a statistically significant response or not. The red dotted lines indicate a Z-score of 3, which we set as the threshold for significance (cf. Fig. 3B). They grey arrows indicate the fraction of observations above

and below the unity line (grey dotted line). The relative strengths of every stimulus combination were compared in a pairwise fashion (Wilcoxon tests with Benjamini-Hochberg correction for multiple comparisons): & *p* < 0.05; && *p* < 0.01; &&& *p* < 0.001. **(B)** We performed a similar analysis focusing only on statistically significant responses (Venn diagrams). Again, all combinations had a positive Z score (as evaluated by a bootstrap method; see Materials and Methods), indicating more than expected convergence. The diameter of each circle indicates the fraction of Purkinje cells showing a significant response to that particular, color coded stimulus. The size of the bar represents the Z score of the overlapping fraction. **(C)** The same for the combinations of three tactile stimuli. Overall, sensory streams tended to converge, rather than diverge, on Purkinje cells. # *p* < 0.10; * *p* < 0.05, ** *p* < 0.01 and *** *p* < 0.001 (Z test).

Figure 5 – Stimulus strength has only a minor impact on complex spike responsiveness

(A) Movements of all large facial whiskers were performed using a piezo-actuator at three different speeds (weak: 1 mm displacement in 62 ms; moderate: 2 mm displacement in 31 ms; strong: 4 mm displacement in 16 ms). The stimulus sequence was randomly permuted. The recordings were made in awake mice. **(B)** Field of view with 24 identified and color-coded Purkinje cells (left) and their corresponding fluorescent traces (right). Stimuli were presented every 2 s and in between trials the laser illumination was briefly blocked to avoid photobleaching. Note that the periods without laser illumination are not drawn to scale. The vertical shaded areas indicate stimulus duration (which was inverse with the stimulus strength). **(C)** Summed fluorescence trace composed of all 24 individual traces showing that not all trials evoked ensemble-wide responses. Some spontaneous, intertrial activity was also observed. **(D)** The average number of complex spikes per frame (of 40 ms) per trial (shaded areas: ± SEM) for the three stimulus strengths show little difference for the weak and moderate stimulation. The time course and amplitude (1-4 mm) of the three stimuli is schematized at the bottom of the graph. Strong stimulation elicited about 30% more complex spikes, as evident from the peak responses for each stimulus intensity. **(E)** Plotting for each of Purkinje cell the average peak response for each stimulus condition results in many overlapping lines. Only a few cells stand out in that they show a strong response that consistently increases with stimulus strength (red lines on top). The Purkinje cells that showed increased responsiveness to stronger stimuli are depicted in red and the others in blue (see also Fig. S4A). **(F)** The same data summarized with box plots for all significantly responsive cells. Data are from 209 significantly responsive Purkinje cells (out of 340 Purkinje cells that were measured in this way). *** $p < 0.001$ (Friedman's test).

A - Whisker pad

B - Upper lip

Peak amp

0   1   2   3   4   5   6   7   8   (Z)

C - All cells

| Whisker pad | Upper lip | Lower lip | Cheek | Sound | LED |

ΔPeak (Z)

n.s.   n.s.   n.s.   n.s.   n.s.   n.s.

Real Random   Real Random   Real Random   Real Random   Real Random   Real Random

D - Only significantly responsive cells

| Whisker pad | Upper lip | Lower lip | Cheek | Sound | LED |

***   ***   ***   ***   ***   *

Cells Pairs   Cells Pairs   Cells Pairs   Cells Pairs   Cells Pairs   Cells Pairs

Figure 6 - Purkinje cells encoding the same stimulus have a tendency to be spatially grouped

Schematic drawing of a field of view with 26 Purkinje cells organized in the medio-lateral direction of crus 1 in an awake mouse. The color of each Purkinje cell corresponds to the maximal response to whisker pad **(A)** or upper lip **(B)** stimulation. Purkinje cells with a filled soma had a peak response with a Z score > 3 and were considered to be statistically significant, in contrast to those with an open soma. Responsive and non-responsive cells are generally intermingled, but a group of "strong responders" can be observed for whisker pad stimulation (red rectangle). **(C)** The anecdotal data in A suggest the presence of clusters of Purkinje cells encoding specific stimuli. For this to be the case, one would expect that neighboring Purkinje cells have roughly similar response strengths. We found that this assumption does not hold as the response strengths of neighbors could not be discriminated from randomly selected cells in the same recording if all Purkinje cells are considered (compared with bootstrap analysis based upon randomly chosen cell pairs within each field of view: all $p >$ 0.8; Z test). Data are represented in violin plots, with the grey lines indicating the 10th, 25th, 50th, 75th and 90th percentiles. **(D)** When considering only the Purkinje cells with statistically significant responses, spatial grouping does occur. For each stimulus type, the black portion of the left bar indicates the fraction of Purkinje cells showing a significant response to that stimulus. The filled portion of the right bar indicates the fraction of the neighbors (always on the medial side) of these significantly responsive Purkinje cells that were also significantly responsive. As can be seen, this fraction is always substantially larger than the fraction of significantly responsive Purkinje cells, indicating a tendency of similar Purkinje cells to group together. Statistical significance was tested by comparing the fraction of Purkinje cells with statistically significant responses and the fraction of neighbors of Purkinje cells with statistically significant responses that showed statistically significant responses as well (after correction for border effects) using Fisher's exact test and after Benjamini-Hochberg correction for multiple comparisons: * $p < 0.05$; *** $p < 0.001$.

Figure 7 – Purkinje cell responses to single whisker stimulation show weak clustering
(A) To investigate smaller receptive fields, we sequentially stimulated five of the large facial whiskers. To avoid interference with other whiskers during active movement, we performed these experiments under ketamine/xylazine anesthesia. Most Purkinje cells, if responsive to single-whisker stimulation, responded only to one of the five whiskers **(B)**. This is illustrated by five peri-stimulus time histograms (PSTHs) from a single, representative Purkinje cell. This particular cell was sensitive to stimulation of the C3 whisker only. The average and 3 s.d. of the baseline firing are indicated (dashed line and grey area). **(C)** Purkinje cells that responded to more than one whisker were typically responsive to the more anterior whiskers (see also Table 5). The widths of the lines indicate the Z value of the occurrence of multiple responses per cell. **(D)** Two recording spots, in close proximity in crus 1 of the same animal, with the identified Purkinje cell dendritic trees.

For each dendrite, the color indicates the whisker(s) to which it was responsive (see legend below with grey denoting the absence of a statistically significant response). **(E)** For each of the two recording sites, the average of the responsive and the non-responsive Purkinje cells is indicated (to the C3 whisker in the left panel and to the C2 whisker in the right panel). Note that only a single cell was responsive to C2 stimulation in recording spot 2. The firing patterns of the non-responsive cells were also averaged (black lines with shades indicating S.E.M.). **(F)** Linear regression revealed that Purkinje cells that were surrounded by other Purkinje cells responsive to the same whisker (same color code as in **A**) had a tendency to show stronger responses to stimulation of that whisker than Purkinje cells that were more isolated. The x-axis represents the fraction of Purkinje cells responsive to the particular whisker within the respective field of view. R = 0.521; $p < 0.001$.

Figure 8 – Purkinje cells encoding the same response group together

(A) Representative field of view with Purkinje cell dendrites. Purkinje cells showing statistically significant responses to whisker pad stimulation are depicted in shades of red and the other cells in shades of blue. **(B)** For each pair of Purkinje cells we calculated the Pearson correlation coefficient (R) during 1 Hz whisker pad stimulation. The pairs of two Purkinje cells that were both statistically significantly responsive to whisker pad stimulation (red symbols) had on average a higher level of synchrony than the pairs connecting a responsive and a non-responsive Purkinje cell (grey symbols; $p < 0.001$; two-dimensional Kolmogorov-Smirnov test). The pairs consisting of two non-responsive Purkinje cells were excluded from this analysis. **(C)** Interestingly, even in the absence of sensory stimulation, the pairs of Purkinje cells that were both responsive to whisker pad stimulation maintained a higher level of synchrony than "heterogeneous pairs". Thus, Purkinje cells with the same receptive field tended to fire more synchronously, even in the absence of stimulation. This analysis was expanded in the presence **(D)** and absence **(E)** of sensory stimulation for six different types of stimulation and illustrated as the median R value per distance category (six bin values of equal distance at a log scale). The shaded areas represent the inter-quartile ranges. * $p < 0.05$; *** $p < 0.001$ (two-dimensional Kolmogorov-Smirnov tests).

Figure 9 – Purkinje cells encode strong and weak sensory stimulation via synchronous firing

(A) Aggregate peri-stimulus time histograms (PSTHs) show that coherent firing of complex spikes predominantly occurs following sensory stimulation. For each field of view, we calculated the number of complex spikes occurring per frame, summing those of all Purkinje cells in that field of view. Subsequently, we made aggregate PSTHs where the color of each bin refers to the number of dendrites simultaneously active. In this field of view, 17 Purkinje cells were measured. Of these, 17 (100%) reacted to air puff, 12 (71%) to whisker pad, 3 (18%) to upper lip, 4 (24%) to lower lip and 1 (6%) to cheek stimulation. **(B)** Based upon a Poisson distribution of complex spikes over all dendrites and bins, one would expect between 0 and 3 simultaneously active dendrites (grey bars). The red bars indicate events involving more dendrites simultaneously than expected from a random distribution. Thus, the sparse firing as expected by chance is relatively constant throughout the trials, but the simultaneous activity of multiple dendrites is strongly enhanced following sensory stimulation. **(C)** A direct overlay of the aggregate PSTHs in response to air puff and lower lip stimulation shows that the strong response found after air puff stimulation comes at the expense of intertrial complex spikes (152 trials per condition). **(D)** For equally long recordings in the presence of different types of stimulation, equal complex spike frequencies were observed as during spontaneous activity (F(2.544, 20.348) = 2.561, p = 0.091, repeated measures ANOVA), indicating that sensory stimulation results in a temporal re-ordering of complex spikes, rather than to the production of more complex spikes.

Figure 10 – Sensory stimulation results in a temporal re-ordering of complex spikes

The temporal distribution of complex spikes was compared between periods with sensory stimulation and periods without. For this analysis, we included only Purkinje cells that displayed a significant response to the stimulus involved (n = 117 for air puff, n = 45 for whisker pad and n = 27 for visual stimulation). Irrespective of the type of stimulation, a reduction in inter-trial activity was found. This was quantified by during the period between 500 and 250 ms prior to stimulus onset using paired t tests; * and *** indicate $p < 0.05$ and $p < 0.001$, respectively.