1

Genetic controls of mouse *Tas1r3*-independent sucrose intake

Cailu Lin[1], Michael G. Tordoff[1], Xia Li[1,2], Natalia P. Bosak[1], Masashi Inoue[1,3], Yutaka Ishiwatari[1,4], Gary K. Beauchamp[1], Alexander A. Bachmanov[1,5*], and Danielle R. Reed[1*]

[1]Monell Chemical Senses Center, Philadelphia, Pennsylvania, United States of America

[2]Current address: Sonora Quest Laboratories, Phoenix, Arizona, United States of America

[3]Current address: Tokyo University of Pharmacy and Life Science, Hachioji, Tokyo, Japan

[4]Current address: Ajinomoto Co., Inc., Tokyo, Japan

[5]Current address: GlaxoSmithKline, Collegeville, Pennsylvania, United States of America

*Corresponding author

E-mail: reed@monell.org (DRR)

Acknowledgments

We gratefully acknowledge Maria L. Theodorides, Zakiyyah Smith, Mauricio Avigdor, and Amy

Colihan for assistance with animal breeding. We also acknowledge Richard Copeland and the

consistent high-quality assistance of the animal care staff at the Monell Chemical Senses

Center and thank them for their service. Matt Kirkey assisted with genotyping the congenic mice.

Yutaka Ishiwatari assisted with genotyping markers. Longhui Chen assisted with metabolic

experiments. Anthony Sclafani made constructive comments on an earlier draft of the

manuscript.

DECLARATIONS

Conflict of interest statement. On behalf of all authors, the corresponding author states that

there are no conflicts of interest.

Funding

National Institutes of Health grants R01 DC00882, R03 DC03854 (AAB and GKB), R01

AA11028, R03 TW007429 (AAB), R03 DC03509, R01 DC04188, R01 DK55853, R01

DK094759, R01 DK058797, P30 DC011735, S10 OD018125, S10 RR025607, S10 RR026752,

and G20 OD020296 (DRR) and institutional funds from the Monell Chemical Senses Center

supported this work, including genotyping and the purchase of equipment. The Center for

Inherited Disease Research through the auspices of the National Institutes of Health provided

genotyping services.

Ethics approval: All animal study procedures were approved by the Monell Chemical Senses
Center Institutional Care and Use Committee.
Consent to participate: Not applicable.
Consent for publication: Not applicable.
Availability of data and material:
Code availability: Not applicable.
Accession IDs:

Author contributions. CL, AB and DR designed the study. CL conducted the study. CL and

DR analyzed data. XL and NB genotyped F_2 hybrids. MI conduct electrophysiological

experiment. CL and DR wrote the paper. AB, MT and GB commented and edited the paper and

all authors read and approved the manuscript.

Abstract

We have previously shown that variation in sucrose intake among inbred mouse strains is due in part to polymorphisms in the *Tas1r3* gene, which encodes a sweet taste receptor subunit and accounts for the *Sac* locus on distal Chr4. To discover other quantitative trait loci (QTLs) influencing sucrose intake, voluntary daily sucrose intake was measured in an F_2 intercross with the *Sac* locus fixed; in backcross, reciprocal consomic strains; and in single- and double-congenic strains. Chromosome mapping identified *Scon3*, located on Chr9, and epistasis of *Scon3* with *Scon4* on Chr1. Mice with different combinations of *Scon3* and *Scon4* genotypes differed more than threefold in sucrose intake. To understand how these two QTLs influenced sucrose intake, we measured resting metabolism, glucose and insulin tolerance, and peripheral taste responsiveness in congenic mice. We found that the combinations of *Scon3* and *Scon4* genotypes influenced thermogenesis and the oxidation of fat and carbohydrate. Results of glucose and insulin tolerance tests, peripheral taste tests, and gustatory nerve recordings ruled out plasma glucose homoeostasis and peripheral taste sensitivity as major contributors to the differences in voluntary sucrose consumption. Our results provide evidence that these two novel QTLs influence mouse-to-mouse variation in sucrose intake and that both likely act through a common postoral mechanism.

Keywords: sucrose consumption, metabolism, QTL, epistasis interaction, mouse

Introduction

Mice and rats avidly drink sucrose solutions, and they often become obese within a few weeks if provided with a sucrose solution in addition to food (Ramirez 1987, Sclafani 1987). Understanding the controls of sucrose consumption is thus of theoretical and practical importance for the study of motivated behavior and of diseases associated with obesity. However, the overconsumption of sucrose and ensuing obesity differ by the genetic makeup of the rodent strain (Glendinning, Breinager et al. 2010). These differences may be due in part to how much sucrose the mouse or rat drinks and in part to its metabolism. Here we focus on the genetics of sucrose consumption, which is highly preferred by nearly all rodents and is usually at least equally or perhaps more potent than some other nutritive sugars, such as glucose, in stimulating intake and weight gain (Kanarek and Orthen-Gambill 1982, Sclafani and Mann 1987).

A genetic component to sucrose consumption is supported by marked differences in the avidity for sucrose among inbred mouse and rat strains (Lush 1989, Lewis, Ahmed et al. 2005, Tordoff, Alarcon et al. 2008); moreover, rat strains have been selectively bred to drink large volumes of solutions containing sugar and other sweeteners (Dess 2000). In laboratory mice, the *Sac* (saccharin preference) locus influences intake of sweeteners, including sucrose (Fuller 1974). We and others have identified *Tas1r3*, the gene underlying the *Sac* locus, which codes for a G-protein-coupled receptor (Bachmanov, Li et al. 2001, Kitagawa, Kusakabe et al. 2001, Max, Shanker et al. 2001, Montmayeur, Liberles et al. 2001, Sainz, Korley et al. 2001) that combines with the T1R2 subunit to form the T1R2+T1R3 sweet taste receptor (Nelson, Hoon et al. 2001).

Variation in the T1R2+T1R3 sweet receptor accounts for differences in sweetener intake among several mouse strains (Reed, Li et al. 2004). Indeed, it plays such a large role in determining sweetener consumption (Inoue, Reed et al. 2004, Reed, Li et al. 2004) that it may

overshadow contribution of other genetic loci, existence of which we hypothesized based on

several pieces of evidence. First, strains of mice with the same *Tas1r3* haplotype show

substantial variation in sweetener consumption, with a large combined effect of other loci (Reed,

Li et al. 2004). Second, *Tas1r3* does not account for variation in sweetener intake in rats, and

thus other genes must be responsible for this variation (Lu, McDaniel et al. 2005). We have

confirmed this hypothesis by performing a genome scan that identified several additional

quantitative trait loci (QTLs) for sucrose consumption (see the companion paper (Lin 2020)). In

addition to the *Sac/Tas1r3* locus (named according to nomenclature as **S**ucrose **con**sumption,

QTL **2**, or *Scon2*), an especially prominent new QTL (*Scon3)* was mapped to chromosome (Chr)

9, and epistatic interaction of *Scon3* with *Scon4* (on Chr1) was detected. When we present the

data with chromosome order, *Scon4* (on Chr1) is first and then *Scon3* (on Chr9), especially for

the description of genotypes for the epistatic QTLs in Figures. We named the newly detected

sucrose QTL on Chr14 as *Scon10* (**S**ucrose **con**sumption, QTL **10**).

Although *Scon2* (*Sac/Tas1r3*), *Scon3* and *Scon4* all influence sucrose intake, the

patterns of their phenotypical effects differ: whereas the effects of *Scon2* were most apparent

when mice were offered low concentrations of sucrose solutions, the *Scon3* and *Scon4* loci

were most apparent when mice were offered high concentrations of sucrose. Thus, to isolate

the *Scon3* and *Scon4* loci, we tested mice with higher concentrations of sucrose and designed

experiments to reduce or eliminate the effects of the *Scon2* locus. To this end, we adopted a

strategy used in immunology to identify histocompatibility genes beyond the major

histocompatibility complex (MHC). The approach was to keep the MHC haplotype constant

among different strains of inbred mice while allowing other genes to vary, which revealed more

histocompatibility antigens and had important practical implications for human transplantation

biology (Almoguera, Shaked et al. 2014). Thus, we bred mice that had the same genotype at

the *Scon2* locus, which allowed us to uncover non-*Scon2* loci regulating sucrose intake.

Specifically, we bred a mouse strain that was congenic at the *Scon2* locus. We intercrossed

with an inbred strain that was identical to the congenic line at the *Scon2* locus but differed

elsewhere in the genome, and we then conducted a genome-wide screen of these intercrossed

mice. The newly discovered *Scon3* and *Scon4* loci were further pursued using consomic and

congenic mapping techniques, capturing them in fragments of the donor genome and reducing

their size over successive cycles of breeding.

To understand the underlying physiological mechanisms responsible for the different

intakes of sucrose by mice of different genotypes, we assessed the congenic mice for their

metabolism (with indirect calorimetry), gustatory nerve responses (with electrophysiology), and

blood glucose and insulin responses. Our goal was to pinpoint the regions affecting sucrose

consumption and gain information that would allow us to narrow the list of potential genes.

Methods

Animal husbandry. All animal study procedures were approved by the Monell Chemical

Senses Center Institutional Care and Use Committee. Inbred C57BL/6ByJ (B6; stock no.

001139) and 129P3/J (129; stock no. 000690) mice used as progenitor strains for the initial

stages of breeding were purchased from The Jackson Laboratory (Bar Harbor, ME, USA); we

bred all other mice in the animal vivarium at the Monell Center, located in Philadelphia,

Pennsylvania (USA). We fed all mice Rodent Diet 8604 (Harlan Teklad, Madison, WI, USA) and

gave them tap water to drink from glass bottles with stainless steel spouts (except while being

tested; see below). All mice were housed in a temperature-controlled vivarium at 23°C on a

12:12-hr light cycle, with lights off at 7 pm, barring unusual circumstances (e.g., power outages).

Pups were weaned at 21-30 days of age, and they were housed in same-sex groups.

Breeding overview. We bred and studied four types of mouse mapping populations, all

derived from B6 and 129 inbred strains (**S1 Table**): F_2 intercross, backcross, consomic, and

congenic (single and double) strains. The specific breeding strategy was different for each

population, but all were part of a large-scale program to map taste genes (Lin, Fesi et al. 2015,

Lin, Fesi et al. 2017).

F_2 intercross. The F_2 intercross was designed so that all mice had the B6 allele of the

Tas1r3 gene (Reed, Li et al. 2004). Females of the B6 inbred strain were mated with two males

from an incipient congenic strain 129P3/J.C57BL/6ByJ-*Tas1r3* (N_4F_4) (Bachmanov, Li et al.

2001, Li, Inoue et al. 2001) to produce F_1 and then F_2 generations.

Backcross and consomic. We generated serial backcross generations (**S1 Table**) by

intercrossing inbred B6 and 129 strains to produce F_1 hybrids, and then crossing F_1 (females)

with males from B6 or 129 strain to breed two reciprocal N_2 hybrids. Starting with N_3, the

selected breeders from the 129.B6 strain were mated with 129 females. We pooled four

backcross generations to form one mapping population (N_3 to N_6) used as segregating

generations for QTL mapping, and also to generate consomic mice (**S2 Table**) using marker-

assisted methods (Lin, Fesi et al. 2015, Lin, Fesi et al. 2017, Lin, Fesi et al. 2018).

Congenic strains. We branched all congenic strains from consomic strains by mating

heterozygous consomic or partially consomic mice with inbred 129 partners. This strategy

allowed us to compare phenotypes of littermates with one copy of the donor region

(heterozygous; 129/B6) to those without the donor region (homozygous; 129/129). Each

congenic mouse was potentially genetically unique (because the donor region could shorten

due to meiotic recombination), so we genotyped all congenic mice to define the donor region

breakpoints. We named congenic strains with the prefix "C": for 129.B6-*Scon4*, we named two

strains, C1 and C2; for 126.B6-*Scon3*, we named seven strains as C3-C9. All nine strains

descended from three progenitors, and the residual genomes of the congenic substrains were

tested and calculated (**S3 Table**). Specifically, we branched *Scon4* congenic mice by

backcrossing one partially consomic male from the N_8 generation (a heterozygous male from

129-Chr1[B6]) to several inbred 129 females. We branched *Scon3* congenic strains by

backcrossing several consomic mice (129.B6-Chr9) from the N_7 generation to 129 females. We

bred the double-congenic mice by mating *Scon3* heterozygous females (B6/129; N_7) with a

homozygous *Scon4* male (B6/B6; N_5F_2). Thus, for the *Scon4* locus in the double-congenic strain,

all offspring were heterozygous (129/B6). For the *Scon3* locus, half the offspring were

heterozygous (129/B6) and half were homozygous (129/129).

Limits of breeding. We had difficulty producing enough *Scon3* heterozygous or

homozygous congenic mice for in-depth phenotyping. We bred *Scon3* congenic strains with

shorter and shorter donor regions, but what few mice we could produce had to be used for

breeding. These breeding problems, similar to those encountered with the corresponding

homozygous consomics (Lin, Fesi et al. 2015), made it difficult to create a homozygous *Scon3*

congenic strain. In contrast, *Scon4* congenic mice were fertile and reproduced easily, so we did

in-depth taste tests and other phenotyping on *Scon4*, consomic, and F_2 mice only.

We had no consomic mice available for Chr14, the location of the *Scon10* locus, so we

were unable to investigate this *Scon10* locus.

Genotyping. We extracted and purified genomic DNA from tail tissue either using a

sodium hydroxide method (Truett, Heeger et al. 2000) or by proteinase K digestion followed by

high-salt precipitation (Gentra/Qiagen, Valencia, CA, USA). We measured the concentration

and purity of DNA using a small-volume spectrophotometer (Nanodrop, Wilmington DE, USA).

We genotyped these DNA samples using two methods: microsatellites and single-nucleotide

polymorphisms (SNPs). For the microsatellites, we used fluorescently labeled primers to amplify

genomic DNA by PCR and scanned the products using an ABI 3100 capillary sequencer

(Applied Biosystems, Foster City, CA, USA). For the SNPs, we genotyped using primers and

fluorescently labeled probes designed to discriminate between alleles, using an ABI Prism 7000

real-time PCR or Step One system (ABI Assay-by-Design, Applied Biosystems, Foster City, CA,

USA). All markers and their physical positions are listed in **S4 Table.**

Phenotyping. We used mice at least 8 weeks old to measure their taste solution

consumption, body composition, metabolism, glucose and insulin tolerance, and gustatory

electrophysiology. Each mouse mapping population was phenotyped separately.

Taste testing (two-bottle choice tests). We performed two types of taste tests: a basic

screen for sucrose intake and a more in-depth survey using sucrose and other taste solutions

(Tordoff 2001). For all tests, mice were 8 or more weeks old. They were individually housed for

at least 2 days before beginning these tests and they were trained to drink deionized water from

two graduated drinking tubes for at least 2 days. Daily measurements were made in the middle

of the light period by reading fluid volume to the nearest 0.1 mL. We measured body weights

before and after each test series.

For the **basic screening test**, mice were offered for 96 hr (4 days) one tube of water

and one tube of 300 mM sucrose. In later studies they were offered both tubes containing 300

mM sucrose, because we learned from the earlier tests that mice drank nearly 100% of their

daily fluid intake from the 300 mM sucrose tube; providing two tubes of sucrose ensured that

even the most avid drinkers would not drink it all in a 24-h test (each drinking tube has a

maximum volume of ~25 mL, so two tubes provided up to 50 ml sucrose solution).

For the **in-depth taste tests**, mice were offered for either 96 or 48 hr one tube of

deionized water and another tube with a taste solution (described below), switching positions of

the two drinking tubes every day to control for side bias (some mice prefer to drink from one

side regardless of the contents of the tube). Between each test series, mice usually had two

tubes of water for at least 2 days.

Mice from the F_2 intercross were tested for 96 hr (4 days) with the following taste

solutions in the order listed: 30 mM glycine, 30 mM D-phenylalanine, 1.6 and 20 mM saccharin,

50 and 300 mM sucrose, 300 mM monosodium glutamate (MSG), 3% and 10% ethanol, and 50

mM $CaCl_2$. All consomic mice, 40 of the *Scon4* homozygous congenic mice, and the inbred

control 129 and B6 mice were tested with the following: 30 mM glycine, 300 mM NaCl, 20 mM

saccharin, 300 mM sucrose, 300 mM MSG, and 3% and 10% ethanol. We purchased all taste

compounds from Sigma Chemical Co. (St. Louis, MO, USA), except the ethanol, which was

purchased from Pharmco Products (Brookfield, CT, USA).

All backcross, *Scon3* and *Scon4* heterozygous congenic, and double-congenic mice

were tested with the basic sucrose two-bottle choice test. The homozygous *Scon4* congenics

(C1) and control 129 inbred mice were tested in four batches (in addition to those 40 mice that

were tested with consomic mice) to efficiently test a large number of taste compounds. We

tested as follows, for 96 hours: **batch 1**, a concentration series of 10, 30, 100, 300, and 600 mM

sucrose (*N*=19 congenic and *N*=10 controls) for 48 hr; **batches 2 and 3**, a concentration series

of 10, 30, 100, 300, 600, and 1200 mM glucose and fructose, respectively (*N*=10 mice per

genotype group) for 48 hr; **batch 4**, 4.6% soybean oil, 6% cornstarch, and 10.5% maltodextrin

(*N*=19 congenics and 10 controls) for 96 hr.

Body composition. We measured the body composition of *Scon4* mice (C1) at 8 and 36

weeks of age using magnetic resonance (MR; Bruker minispec LF110 whole-body composition

rat and mouse analyzer; Bruker BioSpin Corp., Billerica, MA, USA). After the second MR scan,

we also performed body composition analyses by dual-energy x-ray absorptiometry (DEXA;

PIXImus II densitometer; GE software, version 2.00; Lunar Corp., Madison, WI, USA) and by

necropsy using anatomic landmarks to identify and remove the organs (Cinti 1999, Hayakawa

2001). We weighed the following organs to the nearest 0.01 g: spleen, heart, pancreas, brain,

kidney, liver, six "white" adipose depots (pericardial, inguinal, retroperitoneal, subscapular,

mesenteric, and gonadal), and a "brown" adipose depot (subscapular). We also measured body

length from the base of the teeth to the anus using electronic calipers (Fowler ProMax, Kelley

and Kelly Industrial Supply, Syracuse, NY, USA).

Metabolism, food and water intake, and activity. We examined whether *Scon4* genotype

affects metabolism, food intake, water intake, or activity, assessing the *Scon4* homozygous

congenic (C1) and a control group of inbred 129 mice. Using a TSE LabMaster (version 5.0.6;

TSE Systems, Inc., Chesterfield, MO, USA), we measured mice at 8 weeks of age and the

same mice again at 36 weeks of age. We trained the mice for 3-5 days in cages that mimicked

the experimental cages to ensure they learned to eat and drink appropriately from the

suspended food and water containers. We transferred the mice to the experimental cages and

measured oxygen consumption, carbon dioxide production, food and water intake, and physical

activity for 4 consecutive days. We estimated heat production from carbon dioxide and oxygen

consumption corrected for lean body mass.

Glucose and insulin tolerance tests. We tested homozygous *Scon4* congenic mice (C1)

and 129 inbred mice twice (at 8 weeks and again at 36 weeks of age) for changes in blood

glucose in response to exogenous glucose (glucose tolerance test; GTT) or insulin

administration (insulin tolerance test; ITT) (Murovets, Bachmanov et al. 2015). All mice were

tested during the light phase and had unrestricted access to food prior to testing.

For the GTT, we gave the mice glucose (2 g/kg, 10 ml/kg body weight; Sigma-Aldrich St.

Louis, MO, USA) either intraperitoneally (IP) (B6/B6 vs 129/129: N=20 vs 12) or by intragastric

gavage (IG) (B6/B6 vs 129/129: N=21 vs 12). For IP administration, the glucose was dissolved

in 0.9% saline; for the IG route it was dissolved in deionized water. For the ITT, we injected the

mice (B6/B6 vs 129/129: N=19 vs 11) with insulin (2 U/kg, IP; insulin, Novo Nordisk A/S,

Bagsvaerd, Denmark).

We sampled blood from the tip of the tail, with duplicate measures at each time point

(before and after the administration of glucose or insulin) using a One Touch Ultra glucometer

(LifeScan, Inc., USA). For the GTT, we sampled blood at 0, 5, 15, 30, 45, 60, 90, and 120 min;

for the ITT, we sampled at 0, 5, 30, 60, 90, and 120 min. For the GTT, animals were held in

restraint tubes during the blood draws (they were habituated to them before the tests). During

the ITT, mice were unrestrained in their home cages.

Gustatory electrophysiology. We conducted an electrophysiological experiment with

heterozygous *Scon4* (C1) and homozygous *Scon3* congenic mice (C7) and host 129 inbred

mice (5 for each group) as described elsewhere (Inoue, Li et al. 2001, Inoue, McCaughey et al.

2001, Inoue, Reed et al. 2004). Mice ranged from 26 to 50 weeks of age. Activity of the whole

chorda tympani nerve in response to lingual application of taste solutions was recorded for the

following stimuli (in mM, except for ethanol in %): 20 quinine hydrochloride; 100 NaCl; 10 HCl;

1000 glucose; 1000 maltodextrin; 1000 fructose; 100, 300, 1000 sucrose; 2, 6, 20 saccharin;

100, 300, 1000 MSG; 3, 10 ethanol. Each taste solution was washed over the tongue for 30 sec;

between taste solution presentations, the tongue was rinsed with deionized water for at least 1

min. The magnitude of the integrated response at 20 sec after stimulus onset was measured

and expressed as a proportion of the average of the previous and following responses to 100

mM NH$_4$Cl (Inoue, Reed et al. 2004).

Data analyses. We performed several steps to prepare the data for the main statistical

analysis for each mouse population. We checked the distribution of the sucrose intakes for

normality within each mapping population using the Lilliefors test (**S5 Table**), and nonnormal

data were transformed (Delignette-Muller). Male and female mice had similar sucrose intake, so

we pooled their data. For each mapping population we separately checked the correlations

between voluntary sucrose intake and (a) body weight, (b) daily habitual water intake, and (c)

age (**S1 Figure**). (Mice that habitually drank the most water also tended to drink the most

sucrose, so we used habitual water intake as a covariate in the relevant statistical analyses.) In

addition to intake, we also computed preference scores as a ratio of taste solution intake to total

fluid intake (taste solution plus water), in percent, although this preference score was an

imperfect measure because many mice drink nearly 100% of their fluid intake as 300 mM

sucrose when also offered plain water. For all statistical models we used a type 1 (sequential)

sum-of-squares model, and when testing for group differences we used Tukey's HSD tests

except as indicated. We computed statistical tests with R (version 3.3.3) and RStudio (version

1.0.136) and graphed the results using either R or Prism 6 (version 6.05; GraphPad Software,

La Jolla, CA, USA).

F$_2$ intercross. We identified genomic regions shared in common by the F$_2$ mice that

drank the most sucrose using markers from the 19 mouse autosomes and the X chromosome

(R package R/QTL, version 1.41 (Broman, Wu et al. 2003)). We estimated genotype

probabilities and genotype errors using the *calc.genoprob* function, with interval mapping by

maximum likelihood estimation (EM algorithm) for the main-effect QTLs using the *scanone*

function. We tested the significance of each marker regression against 1000 permutations of the

observed data using the n.*perm* function. We defined the confidence intervals as drops of 1

LOD (logarithm of the odds) from the peak expanded to the nearest outside markers using the

lodint function. We estimated the explained phenotypic variance for main-effect QTLs as $1 - 10^{-2\,LOD/n}$,

where n is the sample size, using the LOD score from *scanone* (Broman, Wu et al. 2003).

We investigated marker pair interactions using the *scantwo* function, evaluating significance

against the 1000 permutation tests and confirming the interaction in a general linear model

using the single marker and the interaction of marker pairs as fixed factors.

To assess whether markers identified in the analysis also affected the consumption of

other taste solutions (30 mM glycine, 300 mM NaCl, 20 mM saccharin, 300 mM sucrose, 300

mM MSG, and 3% and 10% ethanol) tested in the F$_2$ population, we analyzed solution intakes in

a general linear model using genotype of each locus as a fixed factor. We analyzed preference

scores of these solutions using one-way ANOVA followed by post hoc tests. In addition, we

computed Pearson correlations among all taste solutions using intakes (adjusted for habitual

water intakes) and preference scores separately.

Backcrosses. Using a pooled backcross population (N_3 to N_6), we conducted a general

linear model analysis of sucrose intake with genotype as a fixed factor. For each marker, we

calculated (a) the genotype means, (b) the *p*-value test statistic as the negative base 10

logarithm, and (c) the effect size using Cohen's *D* (Cohen). We report these values, including

confidence intervals (defined by 2 units of $-\log_{10}$ *p*-value drop), for the peak marker from the

mapping population. Statistical thresholds were computed with a Bonferroni correction to an α

level of 0.05 for the number of markers [*N*=67; $-\log(\alpha/N) \approx 3.13$] (Kanarek and Orthen-Gambill

1982).

Consomics. Building on the results from the F_2 intercross, which detected the main-effect

QTL on Chr9 (*Scon3*) that has epistatic interaction with the QTL on Chr1 (*Scon4*), we studied

the appropriate consomic mice for these chromosomes. For Chr9 we generated two reciprocal

consomic strains, whereas for Chr1 we were able to generate only one consomic strain. For

these three consomic strains, we analyzed sucrose intakes with a general linear model, treating

strain as a fixed factor and using the relevant host inbred strain for comparisons (129-Chr1^{B6} vs

inbred 129, 129-Chr 9^{B6} vs inbred 129, and B6-Chr 9^{129} vs inbred B6). We calculated the

genotype effect size with Cohen's *D*, and we analyzed the intakes of taste solutions other than

300 mM sucrose using the same statistical approach as we used for sucrose. We also used a

general linear model to evaluate all consomic and host inbred strains with genotype as a fixed

factor and habitual water intake as a covariate, followed by post hoc tests. We analyzed

preference scores by strain using *t*-tests.

Heterozygous congenic analysis. We analyzed the data from congenic mice using the

common segment method to find genomic regions shared by strains that have the same trait

(Shao, Sinasac et al. 2010). To this end, for all analyses of congenic mice, we used sucrose

intake as the outcome measure and the presence or absence of a donor region as a fixed factor,

using a proxy marker for the donor region (*rs3708040* for *Scon4* and *rs13480399* for *Scon3*). If

the congenic strain differed from the host inbred strain in voluntary sucrose consumption, we

concluded the associated donor region contained the causal genetic variant; if not, we

concluded otherwise.

Homozygous congenic analysis. We conducted taste tests with a concentration series

for sucrose, glucose, and fructose and with a single concentration for soybean oil, cornstarch,

and maltodextrin. For the concentration series (e.g., fructose), we conducted a two-way mixed-

design ANOVA with strain as the fixed factor (congenics vs host inbred 129) and concentration

as the repeated measure. For the single concentrations (e.g., soybean oil), we analyzed

solution intakes using strain as a fixed factor (congenics vs inbred 129). When informative (no

ceiling effect, i.e., toward 100% preference), we analyzed preference scores in the same way.

There were no differences by sex, so sex was not included in the mode.

Electrophysiology. Chorda tympani responses to individual taste solutions were not

normally distributed, so we used nonparametric Mann-Whitney *U*-tests to assess differences

between genotype groups (Inoue, Glendinning et al. 2007), and Bonferroni correction was

applied for multiple tests for series concentration of each compound.

Body composition. We evaluated the effect of *Scon4* genotype on body composition

using a two-way ANOVA (genotype x sex) followed by post hoc tests for all three measures of

body composition: MR, DEXA, and necropsy. Within each mapping population (F_2, consomics,

and 129.B6-*Scon4* congenics), we also used two-way ANOVAs (genotype x sex) to determine

whether mice with genotypes that increased voluntary sucrose intake also gained more body

weight during the 4-day sucrose intake tests.

Metabolism. Data for each measure from the TSE LabMaster were averaged over the

four 24-hr measurement periods. For each individual, parameters were corrected by lean body

mass using the algorithm in the software package provided by the manufacturer. We used *t*-

tests to compare groups (congenics B6/B6 vs host inbred 129/129).

Glucose and insulin tolerance tests. We analyzed data from GTT and ITT using two-way

mixed-design ANOVA with genotype as the fixed factor and time of blood collection as the

repeated measure.

Evaluating genes and variants in the Scon3 and Scon4 region. Using the results of the

congenic mapping, we identified specific genomic regions that likely contained the causal

variant(s) for each QTL. Within these regions, we identified the known variants and evaluated

them in several ways. First, we made a list of all genetic variants using Mouse Genomes

Project-Query SNPs, indels, or structural variations (Anonymous 2011), which contains the

results of a large-scale genome sequencing project of many inbred mouse strains, including the

129P2/OlaHsd and C57BL/6J strains (Keane, Goodstadt et al. 2011, Yalcin, Wong et al. 2011),

which are closely related to the parental inbred strains used in our study (Yang, Wang et al.

2011). From the list of genetic variants we compiled, we identified sequence variants with the

potential to cause functional changes using the Sorting Intolerant from Tolerant (SIFT) algorithm

(Anonymous 2012). Twelve genetic variants were excluded from this analysis because we could

find no unique identifier for the variant in the dbSNP database (build 138) (Anonymous 2016).

Next, using Mouse Genome Data Viewer (Anonymous 2015), we identified all genes

residing within the genomic coordinates of *Scon4* and *Scon3* defined in the congenic strains and

classified these candidate genes using the PANTHER classification system (Mi, Muruganujan et

al. 2013). Finally, using ToppGene suite (Chen, Bardes et al. 2009), we prioritized candidate

genes by comparing their functional annotations against a training set that consisted of 332

known carbohydrate metabolism mouse genes from the KEGG database (Kanehisa, Furumichi

et al. 2017). We used the Entrez identifiers of the genes (N=248) within the *Scon3* and *Scon4*

regions as input for candidate gene prioritization; therefore, only genes with an appropriate

identifier were included.

We reasoned that the causal genes from *Scon3* and *Scon4* jointly contribute to the same

pathway, so we identified gene pairs that met this criterion (i.e., one gene-pair member was

from the *Scon4* region and the other was from the *Scon3* region). We thus assessed the

protein-protein associations by searching the protein names in the STRING database (von

Mering, Jensen et al. 2005) with multiple active interaction sources that included three types of

data: (1) direct experimental evidence of protein-protein interactions; (2) less direct evidence

generated by grouping proteins of known function into a metabolic, signaling, or transcriptional

pathway; and (3) *in silico* computation techniques for detecting new protein-protein associations.

We exported gene pairs for protein-protein interactions with association scores > 0.9, which

were computed by benchmarking them against the training set (von Mering, Jensen et al. 2005),

and identified those pairs that fit our criterion.

Finally, we compared expression profiling of genes (*N*=248) within the *Scon3* and *Scon4*

critical regions at three mouse tissue sites (striatum, brown and white fat) between 129 and B6

inbred strain using publicly available GEO data (Davis and Meltzer 2007), including the closely

related strains 129S1/SvlmJ and C57Bl/6J. Grouping three samples of each strain 129P3/J and

C57BL/6J (GSE7762; (Korostynski, Piechota et al. 2007)), we conducted microarray-based

gene expression analysis for mouse striatum with a web tool GEO2R

(http://www.ncbi.nlm.nih.gov/geo/geo2r) by calculating the distribution of fluorescence intensity

values for the selected samples and then performing a LIMMA (linear model for microarray data

analysis) statistical test (Smyth 2005) after applying log transformation to the data. We

downloaded bulk RNASeq data [four samples for each of strain (129S1/SvImJ and C57Bl/6J)

for each tissue site (mouse brown and white fat); GSE91067] (Raymond E Soccio 2017), and

transformed the transcript count data. We created a matrix for the two groups' data (129 vs B6)

and fitted data with the gene-wise glmFit function of edgeR package (Robinson, McCarthy et al.

2010); we conducted likelihood ratio tests for 129 vs B6 for brown and white fat tissue

separately. The differentially expressed genes in the two types of gene profiling data sets were

reported as one $\log_2$-fold change (corresponding to a twofold change) with an associated *p*-

value (*p*<0.05) corrected for false discovery (FDR) for multiple tests (Benjamini 1995).

Results

Identification of main effect and epistatic QTLs

We mapped a highly significant main-effect QTL on Chr9 (*Scon3*) with a strong effect on 300

mM sucrose intake (explaining 37% of the trait variance) and a significant main-effect QTL on

Chr14 (*Scon10*) with a weaker effect (*7*% of trait variance); the B6 allele was dominant and

increased sucrose consumption in both cases (**Figure 1A**). The peak *Scon3* LOD score was 28

at marker *rs3023231* (106.8 Mb), with a confidence interval of 105.1-108.1 Mb (between

rs13480395 and *rs4227916*). The two-dimensional genome scan detected that *Scon3*

epistatically interacts with a new QTL for 300 mM sucrose intake, *Scon4* on Chr1 (**Figure 1B**),

with interaction LOD=1.5, and this epistatic interaction was confirmed by general linear model

analysis (**Figure 1C**). While the B6 allele of *Scon3* increased sucrose intake, the B6 allele of

Scon4 decreased sucrose intake. The phenotypic difference among *Scon4* genotypes shows up

only with 129/129 genotype of *Scon3.*

Variation at these two loci (*Scon3, Scon4*) had similar effects on the taste preferences of

the mice, strongly affecting voluntary consumption of sucrose and a few other taste compounds

(**S2 Figure**). For the *Scon4* locus, the only other compounds affected was 20 mM saccharin, but

with opposite effect direction: B6 allele of *Scon4* decrease sucrose intake but increase

saccharin intake (**S2 Figure**). For the *Scon3* locus, other compounds affected were 3% and

10% ethanol and 300 mM MSG with the same effect direction, as well as 150 mM $CaCl_2$ with

opposite effect direction (**S2 Figure**). There was no genotype effect (*Scon4* or *Scon3*) on

average daily water consumption (**S3 Figure**). Independent of genotype, mice that drank the

most 300 mM sucrose drank more MSG and ethanol but not the other taste solutions (**S4**

Figure)**.**

Confirmation of main effect and epistatic QTLs

Backcross. In the backcross population, we detected *Scon3* at the peak marker *rs302323131*;

this is the same location as in the F_2 mapping population and has an allelic effect of the same

direction and magnitude, with an –log10 (p vale) of 17.3 and a Cohen's *D* value of >0.8 (**Figure**

2A). We were not able to test for the effects of *Scon4* because these backcross mice were not

genotyped for markers on Chr1.

Consomics. The results from the consomics were consistent with those from the F_2 and

backcross populations, confirming the effects of *Scon3* and *Scon4* and their interaction. For

Scon4, mice with a B6-introgressed Chr1 drank less sucrose solution than did the host control

mice (**Figure 2C, D**), with an effect size > 1.5 (**Figure 2E**). For *Scon3,* mice with a B6-

introgressed Chr9 drank more sucrose, and mice with the 129-introgressed Chr9 drank less

sucrose, compared with host controls, with effect sizes of 1.1 and 2.7, respectively (**Figures 2C,**

D, E). Consomic mice with B6 alleles at both *Scon3* and *Scon4* drank much more sucrose than

did mice with 129 alleles at both loci (**Figures 2D, F**). We evaluated all consomic and control

mice in a single model and found that mice with different genotypes at *Scon4* and *Scon3*

differed in voluntary sucrose intake, with a marked increase in mice with certain combinations

(e.g., B6/B6 of *Scon4* and B6/B6 *Scon3*) of those alleles (**Figure 2F)**. Consomic mice also

differed in their intake of other taste solutions compared with mice from the inbred host strains

(**S5 Figure**).

129.B6-Scon4 congenics. The results from congenic mapping support the presence of

Scon4 near the centromere of Chr1 (**Figure 3**). We bred and phenotyped two *Scon4* congenic

substrains (C1 and C2; **Figure 3B**; **S6 Table**). Congenic mice from the C1 substrain (with the

donor region from 0 to 38.5 Mb) drank more sucrose than did control mice (**Figure 3C**) and

hence retained *Scon4*. However, mice from the C2 congenic strain (with a shorter donor region

spanning 24.5-38.5 Mb) drank the same volume of sucrose as did controls (**Figure 3D**) and

hence the substrain (C2) did not retain *Scon4*. Comparison of the two substrains defined the

Scon4 critical region as 0-25.4 Mb. This mapping result was supported by the outcome of the

general linear model, with the strongest association between sucrose intake and marker

genotypes in a 0-25.4 Mb region (**Figure 3A**).

In addition to heterozygous *129.B6-Scon4* congenic mice, we produced and phenotyped

homozygous congenic mice by intercrossing heterozygotes from substrain C1. *Scon4* congenic

homozygotes drank less than one-third the amount of 300 mM sucrose consumed by 129 inbred

mice (effect size of 2.6; **Figure 4**), which is consistent with *Scon4* effects in F_2 (**Figure 1C**).

129.B6-Scon3 congenics. The results from congenic mapping support the presence of

Scon3 in the same region of Chr9 (**Figure 5**) as identified in the F_2 and backcross mapping

populations. We bred and phenotyped seven *Scon3* congenic strains (C3–C9; **Figure 5B**; **S7**

Table) and found that mice from five strains (C5-C9) with donor fragments including the 1.2 Mb

critical region (105.7-106.9 Mb) drank more 300 mM sucrose than did control mice, whereas

mice from two strains (C3, C4) without this critical region did not differ from control mice (**Figure**

5C). The mapping result was supported by the outcome of the general linear model, with the

strongest relationship between sucrose intake and genotype within this 1.2 Mb region (**Figure**

5A). This region contains 6 noncoding RNA genes, 33 protein-coding genes, 6 pseudogenes,

and 9 processed transcripts (**Figure 5D**).

Scon3 and Scon4 interaction in double congenics. To confirm epistatic interaction

between *Scon3* and *Scon4* detected in F$_2$ (**Figures 1B, C)**, we intercrossed *Scon3* and *Scon4*

single congenics to produce mice with B6 donor alleles of both QTLs segregated (double

congenic, **Figure 2B**). To achieve this, we used a homozygous *Scon4* congenic male (B6/B6;

N$_5$F$_2$) mated with heterozygous *Scon3* congenic females (B6/129; N$_7$). Consequently, all

offspring were heterozygous (129/B6) for *Scon4* but had segregating (129/B6 heterozygotes

and 129/129 homozygotes) *Scon3* genotypes (**Figures 2G, H**). The double congenic mice from

this substrain that had B6 alleles at both loci (129/B6 genotype for *Scon3* and *Scon4*) drank

three times more sucrose than their littermates with genotypes 129/B6 of *Scon4* and 129/129 of

Scon3 (Cohen's *D*=5.0); mice with only one B6 allele drank much less (Cohen's *D*: *Scon4*, 0.9;

Scon3, 0.8; **Figures 2H, I**). For the genotypes for *Scon4* and *Scon3,* refer to **Figure 2G**, except

for congenic strains C2, C3 and C4, which we recoded as 129/129 because these strains did

not retain the QTLs (**Figures 3B, D,** and **Figures 5B, C**). Thus, mice with three genotypes were

included in the pooled congenic analyses: 129.B6-*Scon4* (129/B6 of *Scon4*, 129/129 of *Scon3*;

mice from substrain C1), 129.B6-*Scon3* (129/129 of *Scon4*, 129/B6 of *Scon3*; mice from

substrains C5-C9), and mice without B6 allele (129/129 of *Scon4*, 129/129 of *Scon3;* mice from

half of substrains C1, C5-C9 and all of substrains C2-C4). A single model encompassing all the

congenic data produced results very similar to those of the F$_2$ and consomics: mice differed in

voluntary sucrose intake depending on the combination of *Scon3* and *Scon4* genotypes [*Scon3*:

F(1,989)=109.7, *p*<0.00001; *Scon4*: *F*(1, 989)=8.65, *p*<0.005; *Scon3* × *Scon4*: *F*(1, 989)=40.2,

p<0.00001; **Figure 2J**].

Characterization of the *Scon3 and Scon4* QTL phenotypes

Electrophysiology. No statistically significant differences from 129 mice were found for 129.B6-

Scon3 heterozygotes or 129.B6-*Scon4* heterozygotes for any of the taste stimuli tested (100,

300 and 1000 mM sucrose; 20 mM quinine; 100 mM NaCl; 10 mM HCl; 1000 mM glycine; 1000

maltose; 1000 mM fructose; 2, 6, and 20 mM saccharin; 100, 300, and 1000 mM MSG; and 3%

and 10% ethanol; $p > 0.05$, Bonferroni correction for multiple tests; **S6 Figure**, **S8 Table**)

In the following characterization the *Scon4* QTL effect, we compared homozygous

Scon4 congenics (B6/B6) with 129 host inbred mice (129/129).

Two-bottle preference tests. The *Scon4* genotype affected the consumption of 300 and

600 mM sucrose, 600 mM glucose, and 600 mM fructose, but it had no effect at other, mostly

lower concentrations of these sugars (**Figure 6**). Likewise, *Scon4* congenic and 129 inbred mice

did not differ in their consumption of 20 mM saccharin, 30 mM glycine, or 10% ethanol (**S7A, B,**

F Figures) or of 4.6% soybean oil, 6% cornstarch, or 10.5% maltodextrin (**S8A-C Figures**).

However, *Scon4* congenic mice had lower intakes of 300 mM NaCl, 300 mM MSG, and 3%

ethanol and lower 3% ethanol preference scores compared with 129 inbred mice (**S7C-E**

Figures).

Body composition analysis. We used three methods to study body composition, MR,

DEXA, and necropsy, all of which indicated that the *Scon4* genotype had no significant effect on

body composition (MR, **S9A Figure**; DEXA, **S9B Figure**; necropsy, **S10 Figure**). From the

necropsy data, we learned that mice of different genotypes had similar organ weights, except for

the subscapular adipose depot (von Mering, Jensen et al. 2005) and heart (**S10 Figure**).

Drinking sucrose voluntarily for 4-12 days did not result in significant differences in body weight

gain among mice of different *Scon3* and *Scon4* genotypes (**S11 Figure**).

Metabolism. Homozygous *Scon4* mice differed in their metabolic gas exchange

compared with inbred host controls, especially in early adulthood. Specifically, mice from the C1

strain (B6/B6) consumed less oxygen, generated less heat (kcal/kg lean), and burned

proportionately more calories as fat than carbohydrate compared with inbred 129 mice (129/129;

Figures 7C-F). However, at both ages tested, these homozygous *Scon4* mice ate the same

amount of food (g/24 hr), drank the same amount of water (mL/24 hr), and were as active (beam

breaks/hr) as inbred 129 mice (**Figures 7A, B, G)**.

Glucose tolerance test. Homozygous *Scon4* mice did not differ from inbred 129 mice in

blood glucose or insulin in response to exogenous glucose, by either IP or IG administration,

whether tested at 8 weeks or at 36 weeks (**S12A, B Figures**).

Insulin tolerance test. After injection with insulin in 8-week-old mice, a significant

genotype × time interaction was detected, with *Scon4* congenic mice tending to have higher

insulin sensitivity (lower blood glucose) than 129 inbred mice, with post hoc tests detecting

significant difference at 90 min (**S12C Figure,** left panel). The same mice at 36 weeks showed

no effects of genotype at all and no interaction of genotype with time (**S12C Figure,** right panel).

Candidate genes

We identified and investigated genes in the *Scon3* and *Scon4* regions defined by the congenic

mapping results (*Scon3*, 9:105650529-106888925; *Scon4*, 1:1-25377067). *Scon3* harbors 55

genes and *Scon4* harbors 193 genes (**S9 Table**), of which 49 are protein-coding genes [21] and

199 are other gene types, such as long noncoding RNA (lncRNA) genes. Among 41 of the

protein-coding genes, there were 168 relevant missense and stop codon gain/loss variants (**S9**

Table). We grouped the biological processes for these genes into categories (**S13A Figure**)

and found that 31 of these known genes are involved in carbohydrate metabolism (**S9 Table**).

We identified pairs with the highest association confidence scores and containing one gene

from the *Scon4* region and another from the *Scon3* region. For example, *Rb1cc1* (RB1-inducible

coiled-coil 1) from *Scon4* and *Pik3r4* (phosphatidylinositol 3 kinase, regulatory subunit,

polypeptide 4, p150) from *Scon3* have an interaction confidence score > 0.9 (**S13B Figure**).

The prioritization of these two genes ranks first and eleventh based on the similarity to other

genes involved in carbohydrate metabolism, and one member of the pair (*Pik3r4*) has a

missense variation (**S9 Table**). Additional gene pairs also had protein-protein association

scores > 0.9: *Oprk1-Grm2* (opiate and glutamine receptors), *Grm2-Npbwr1* (glutamine and

neuropeptide receptors), and *Rpl29-Rpl7* (ribosomal proteins) (**S10 Table**).

Gene expression profiling at three mouse tissue sites (striatum, brown and white fat)

revealed a significantly differential expression of 20 candidate genes (*2610203C22Rik,*

Abhd14a, Adhfe1, Col6a4, Col6a5, Dusp7, Eya1, Glyctk, Iqcf1, Kcnq5, Mcm3, Paqr8, Parp3,

Poc1a, Prex2, Sgk3, Snhg6, Sox17, St18, Ube2w) within *Scon3* and *Scon4* (**Figure 8; S11**

Table) between 129 and B6 inbred mouse strains (twofold change with FDR<0.05).

Discussion

We identified *Scon3* and *Scon4*, two small regions of the mouse genome containing variations

that affect the voluntary intake of concentrated sucrose solution, which emerged when influential

variation in the *Tas1r3* gene was removed. We used B6 and 129 mice for these experiments

because of the marked differences in sucrose consumption between the two strains

(Bachmanov, Tordoff et al. 2001) and because we were interested in learning more about the

region on chromosome 9 identified in an earlier genome scan (see the companion paper (Lin

2020)). The utility of these mice was confirmed here by the initial *Tas1r3*-independent

intercross results, which confirmed the striking influence of a locus on Chr9 that interacts with a

separate region on Chr1.

We are unable to draw firm conclusions about the *Scon4* locus because we did not test

all taste solutions in both naive and experienced mice (**S12 Table**). The parental inbred strains

are susceptible to develop learned flavor preferences and thus prior experience with nutrients

changes their response to taste solutions (Sclafani 2006), so we cannot directly compare the

results of tests conducted with naive mice with those conducted on mice with prior experience.

Thus, while we can be sure that naive mice differ in sucrose intake by *Scon4* genotype, but we

cannot be sure that there is no effect of genotype on starch intake in naive mice because mice

drank another nonsweet but preferred solution beforehand (soybean oil).

We found that the *Scon4* genotype was related to whole-body fuel use. Under basal

conditions, mice with genotypes of *Scon4* that decreased voluntary sucrose consumption also

oxidized more carbohydrate and less fat than did mice with opposing genotypes. These genetic

effects on voluntary sucrose intake and carbohydrate oxidation do not seem to arise from the

earliest steps of intestinal absorption because there were few or no differences by genotype in

the outcomes of glucose and insulin tolerance tests. However, we cannot rule out differential

stimulation of postoral nutrient detectors that may partition nutrient oxidation and storage.

Peripheral taste sensitivity might be a contributor to the differences in voluntary sucrose

consumption, but four points make this explanation unlikely: (a) genotype affects voluntary

intake at high but not low sucrose concentrations (the ability to detect sweet taste is more

important at lower concentrations), (b) genotype effects studied here were absent for peripheral

nerve responses, (c) these small effects were in the opposite direction from those of another

genetic model of voluntary sucrose intake (Inoue, Reed et al. 2004), and (d) we found no

relationship between *Scon4* or *Scon3* in a genome scan using taste nerve responses as a trait

(see the companion paper (Lin 2020)). However, we cannot rule out a peripheral effect entirely,

given the small sample size of mice tested for peripheral neural responses to sucrose.

There are at least two ways to approach the evaluation of genes within the critical

intervals for their effect on voluntary sucrose consumption. The first method is to examine all

genes within the regions to determine if there are known connections between those genes and

voluntary sucrose consumption (narrowly) and metabolism (more broadly). No genes were

linked to voluntary sucrose consumption in the databases surveyed, but many were part of

carbohydrate and fat metabolism pathways, including *Pik3r4* (Nemazanyy, Blaauw et al. 2013),

Alas1 (Simcox, Mitchell et al. 2015)*, Acy1* (Alexandre-Gouabau, Bailly et al. 2012), *Cops5* (Tian,

Peng et al. 2010), *Atp6v1h* (Olsson, Yang et al. 2011, Baumeier, Kaiser et al. 2015), *Tlr9* (Lewis

and Cobb 2010), *Lypla1* (Hacia, Fan et al. 1999, Mootha, Bunkenborg et al. 2003), *Pkhd1*

(Kaimori, Nagasawa et al. 2007)*,* and *Gdap1* (Lopez Del Amo, Palomino-Schatzlein et al. 2017).

We further reasoned that the interaction of the *Scon3* and *Scon4* regions indicated that causal

genes might be part of the same pathway. That approach led us to identify particular gene pairs

that jointly contribute to a shared function (e.g., *Pik3r4* and *Rb1cc1*); however, their involvement

in voluntary sucrose intake awaits direct tests.

The second approach is to find genes or gene pathways that have a similar constellation

of traits, with the idea that the genes in the region could be unknown members of the known

pathway. Using this second method, we learned that many of the traits of the *Scon4* and *Scon3*

mice studied here are similar to mice with knockout or overexpression of the *FGF21* gene.

These traits include increased intake of sugar and the lack of large effects on peripheral taste

nerves and body weight or fatness (von Holstein-Rathlou, BonDurant et al. 2016). A next step

would be to evaluate the genes in the region experimentally for their roles in the *FGF21*

signaling pathway. We note that two proteins that regulate this signaling pathway, *Prdm14*

(Grabole, Tischler et al. 2013) and *Sulf1* (Buresh, Kuslak et al. 2010), are located in the *Scon4*

region. *SGLT1* and *KATP*, as an alternate potential sucrose transduction mechanism (Yee,

Sukumaran et al. 2011)*,* are located on different chromosomes from *Scon3* and *Scon4,* and it is

unlikely there is a peripheral taste effect of *Scon3* and *Scon4*, so they can be ruled out as

candidates.

Through our use of consomic mouse strains, we indirectly mapped other genes for taste-

related traits. Consomic strains contain an entire introgressed chromosome; thus, there is a

greater potential to capture more causal genes and their variants compared with the congenic

strains, which have smaller introgressed regions. As expected, our consomic mapping results

discovered new regions of interest distinct from the *Scon* loci and also confirmed many

previously reported regions that contain variation that affects intake of several other taste

compounds by mice (Phillips, Crabbe et al. 1994, Tarantino, McClearn et al. 1998, Vadasz,

Saito et al. 2000, Tordoff, Reed et al. 2008).

This study was conducted in mice, but what we learned has connections to human

health. Although the contribution of sucrose, especially in beverages, to the ills of the modern

diet remains controversial, the benefits of understanding the biological causes of excessive

sugar consumption are likely to be helpful. An earlier example of this type of benefit comes from

the discovery of a subunit of the mouse sweet receptor, T1R3 (Bachmanov, Li et al. 2001,

Kitagawa, Kusakabe et al. 2001, Max, Shanker et al. 2001, Montmayeur, Liberles et al. 2001,

Nelson, Hoon et al. 2001, Sainz, Korley et al. 2001). This discovery was useful in understanding

taste biology, but it is now apparent that this sugar-sensing receptor has many nongustatory

functions, including insulin regulation (Geraedts, Takahashi et al. 2012) and adipogenesis

(Simon, Parlee et al. 2013). Likewise, we hope this study provides insights toward

understanding the appetition for sugar and that this knowledge can be harnessed to help

prevent the negative effects of excessive sugar consumption by understanding and allaying the

human hunger for sweetness.

653　　　　　　　　　　　　　　　　　References
654

Alexandre-Gouabau, M. C., E. Bailly, T. L. Moyon, I. C. Grit, B. Coupe, G. Le Drean, H. J.
Rogniaux and P. Parnet (2012). "Postnatal growth velocity modulates alterations of proteins
involved in metabolism and neuronal plasticity in neonatal hypothalamus in rats born with
intrauterine growth restriction." J Nutr Biochem **23**(2): 140-152.
Almoguera, B., A. Shaked and B. J. Keating (2014). "Transplantation genetics: current status
and prospects." Am J Transplant **14**(4): 764-778.
Anonymous (2011). Mouse Genomes Project - Query SNPs, indels or SVs. Wellcome Trust
Sanger Institute.
Anonymous. (2012). "SIFT web server: predicting effects of amino acid substitutions on
proteins." August 15th 2015. Retrieved 10/7/2016, 2016, from
http://siftdna.org/www/SIFT_dbSNP.html.
Anonymous (2015). Mus musculus (laboratory mouse) genome view, National Center for
Biotechnology Information.
Anonymous. (2016). "dbSNP, Single Nucleotide Polymorphisms;
http://www.ncbi.nlm.nih.gov/SNP/index.html." Build 138 Retrieved 10/7/2016, 2016, from
http://www.ncbi.nlm.nih.gov/SNP/index.html.
Bachmanov, A. A., X. Li, D. R. Reed, J. D. Ohmen, S. Li, Z. Chen, M. G. Tordoff, P. J. de Jong,
C. Wu, D. B. West, A. Chatterjee, D. A. Ross and G. K. Beauchamp (2001). "Positional cloning
of the mouse saccharin preference (*Sac*) locus." Chemical Senses **26**(7): 925-933.
Bachmanov, A. A., X. Li, D. R. Reed, J. D. Ohmen, S. Li, Z. Chen, M. G. Tordoff, P. J. de Jong,
C. Wu, D. B. West, A. Chatterjee, D. A. Ross and G. K. Beauchamp (2001). "Positional cloning
of the mouse saccharin preference (Sac) locus." Chem Senses **26**(7): 925-933.
Bachmanov, A. A., M. G. Tordoff and G. K. Beauchamp (2001). "Sweetener preference of
C57BL/6ByJ and 129P3/J mice." Chemical Senses **26**: 905-913.
Baumeier, C., D. Kaiser, J. Heeren, L. Scheja, C. John, C. Weise, M. Eravci, M. Lagerpusch, G.
Schulze, H. G. Joost, R. W. Schwenk and A. Schurmann (2015). "Caloric restriction and
intermittent fasting alter hepatic lipid droplet proteome and diacylglycerol species and prevent
diabetes in NZO mice." Biochim Biophys Acta **1851**(5): 566-576.
Benjamini, Y., and Hochberg, Y (1995). "Controlling the false discovery rate: a practical and
powerful approach to multiple testing. Journal of the Royal Statistical Society Series B, 57, 289–
300."
Broman, K. W., H. Wu, S. Sen and G. A. Churchill (2003). "R/qtl: QTL mapping in experimental
crosses." Bioinformatics **19**(7): 889-890.
Buresh, R. A., S. L. Kuslak, M. A. Rusch, C. M. Vezina, S. B. Selleck and P. C. Marker (2010).
"Sulfatase 1 is an inhibitor of ductal morphogenesis with sexually dimorphic expression in the
urogenital sinus." Endocrinology **151**(7): 3420-3431.
Chen, J., E. E. Bardes, B. J. Aronow and A. G. Jegga (2009). "ToppGene Suite for gene list
enrichment analysis and candidate gene prioritization." Nucleic Acids Res **37**(Web Server issue):
W305-311.
Cinti, S. (1999). "The adipose organ." Milano: Editrice Kurtis.
Cohen, J. "Statistical power analysis for the behavioral sciences. 2nd ed. 1988, Hillsdale, N.J.: L.
Erlbaum Associates. xxi, 567 p.".
Davis, S. and P. S. Meltzer (2007). "GEOquery: a bridge between the Gene Expression
Omnibus (GEO) and BioConductor." Bioinformatics **23**(14): 1846-1847.
Delignette-Muller, M., Pouillot, R., Denis, J., and Dutang, C. "fitdistrplus: Help to Fit of a
Parametric Distribution to Non-Censored or Censored Data. R package. 2014.".
Dess, N. K. (2000). "Responses to basic taste qualities in rats selectively bred for high versus
low saccharin intake." Physiol Behav **69**(3): 247-257.
Fuller, J. L. (1974). "Single-locus control of saccharin preference in mice." J Hered **65**(1): 33-36.

Geraedts, M. C., T. Takahashi, S. Vigues, M. L. Markwardt, A. Nkobena, R. E. Cockerham, A. Hajnal, C. D. Dotson, M. A. Rizzo and S. D. Munger (2012). "Transformation of postingestive glucose responses after deletion of sweet taste receptor subunits or gastric bypass surgery." Am J Physiol Endocrinol Metab **303**(4): E464-474.

Glendinning, J. I., L. Breinager, E. Kyrillou, K. Lacuna, R. Rocha and A. Sclafani (2010). "Differential effects of sucrose and fructose on dietary obesity in four mouse strains." Physiol Behav **101**(3): 331-343.

Grabole, N., J. Tischler, J. A. Hackett, S. Kim, F. Tang, H. G. Leitch, E. Magnusdottir and M. A. Surani (2013). "Prdm14 promotes germline fate and naive pluripotency by repressing FGF signalling and DNA methylation." EMBO Rep **14**(7): 629-637.

Hacia, J. G., J. B. Fan, O. Ryder, L. Jin, K. Edgemon, G. Ghandour, R. A. Mayer, B. Sun, L. Hsie, C. M. Robbins, L. C. Brody, D. Wang, E. S. Lander, R. Lipshutz, S. P. Fodor and F. S. Collins (1999). "Determination of ancestral alleles for human single-nucleotide polymorphisms using high-density oligonucleotide arrays." Nat Genet **22**(2): 164-167.

Hayakawa, T., H. Yamasita, and T. Iwaki (2001). "A Color Atlas of Sectional Anatomy of the Mouse." Braintree Scientific Inc.

Hubbard, T. J., B. L. Aken, K. Beal, B. Ballester, M. Caccamo, Y. Chen, L. Clarke, G. Coates, F. Cunningham, T. Cutts, T. Down, S. C. Dyer, S. Fitzgerald, J. Fernandez-Banet, S. Graf, S. Haider, M. Hammond, J. Herrero, R. Holland, K. Howe, K. Howe, N. Johnson, A. Kahari, D. Keefe, F. Kokocinski, E. Kulesha, D. Lawson, I. Longden, C. Melsopp, K. Megy, P. Meidl, B. Ouverdin, A. Parker, A. Prlic, S. Rice, D. Rios, M. Schuster, I. Sealy, J. Severin, G. Slater, D. Smedley, G. Spudich, S. Trevanion, A. Vilella, J. Vogel, S. White, M. Wood, T. Cox, V. Curwen, R. Durbin, X. M. Fernandez-Suarez, P. Flicek, A. Kasprzyk, G. Proctor, S. Searle, J. Smith, A. Ureta-Vidal and E. Birney (2007). "Ensembl 2007." Nucleic Acids Res **35**(Database issue): D610-617.

Inoue, M., J. I. Glendinning, M. L. Theodorides, S. Harkness, X. Li, N. Bosak, G. K. Beauchamp and A. A. Bachmanov (2007). "Allelic variation of the Tas1r3 taste receptor gene selectively affects taste responses to sweeteners: evidence from 129.B6-Tas1r3 congenic mice." Physiol Genomics **32**(1): 82-94.

Inoue, M., X. Li, S. A. McCaughey, G. K. Beauchamp and A. A. Bachmanov (2001). "*Soa* genotype selectively affects mouse gustatory neural responses to sucrose octaacetate." Physiological Genomics **5**: 181-186.

Inoue, M., S. A. McCaughey, A. A. Bachmanov and G. K. Beauchamp (2001). "Whole-nerve chorda tympani responses to sweeteners in C57BL/6ByJ and 129P3/J mice." Chemical Senses **26**: 915-923.

Inoue, M., D. R. Reed, X. Li, M. G. Tordoff, G. K. Beauchamp and A. A. Bachmanov (2004). "Allelic variation of the *Tas1r3* taste receptor gene selectively affects behavioral and neural taste responses to sweeteners in the F_2 hybrids between C57BL/6ByJ and 129P3/J mice." J Neurosci **24**(9): 2296-2303.

Kaimori, J. Y., Y. Nagasawa, L. F. Menezes, M. A. Garcia-Gonzalez, J. Deng, E. Imai, L. F. Onuchic, L. M. Guay-Woodford and G. G. Germino (2007). "Polyductin undergoes notch-like processing and regulated release from primary cilia." Hum Mol Genet **16**(8): 942-956.

Kanarek, R. B. and N. Orthen-Gambill (1982). "Differential effects of sucrose, fructose and glucose on carbohydrate-induced obesity in rats." J Nutr **112**(8): 1546-1554.

Kanehisa, M., M. Furumichi, M. Tanabe, Y. Sato and K. Morishima (2017). "KEGG: new perspectives on genomes, pathways, diseases and drugs." Nucleic Acids Res **45**(D1): D353-D361.

Keane, T. M., L. Goodstadt, P. Danecek, M. A. White, K. Wong, B. Yalcin, A. Heger, A. Agam, G. Slater, M. Goodson, N. A. Furlotte, E. Eskin, C. Nellaker, H. Whitley, J. Cleak, D. Janowitz, P. Hernandez-Pliego, A. Edwards, T. G. Belgard, P. L. Oliver, R. E. McIntyre, A. Bhomra, J. Nicod, X. Gan, W. Yuan, L. van der Weyden, C. A. Steward, S. Bala, J. Stalker, R. Mott, R. Durbin, I. J.

Jackson, A. Czechanski, J. A. Guerra-Assuncao, L. R. Donahue, L. G. Reinholdt, B. A. Payseur, C. P. Ponting, E. Birney, J. Flint and D. J. Adams (2011). "Mouse genomic variation and its effect on phenotypes and gene regulation." Nature **477**(7364): 289-294.

Kitagawa, M., Y. Kusakabe, H. Miura, Y. Ninomiya and A. Hino (2001). "Molecular genetic identification of a candidate receptor gene for sweet taste." Biochem Biophys Res Commun **283**(1): 236-242.

Korostynski, M., M. Piechota, D. Kaminska, W. Solecki and R. Przewlocki (2007). "Morphine effects on striatal transcriptome in mice." Genome Biol **8**(6): R128.

Lewis, C. J. and B. A. Cobb (2010). "Carbohydrate oxidation acidifies endosomes, regulating antigen processing and TLR9 signaling." J Immunol **184**(7): 3789-3800.

Lewis, S. R., S. Ahmed, C. Dym, E. Khaimova, B. Kest and R. J. Bodnar (2005). "Inbred mouse strain survey of sucrose intake." Physiol Behav **85**(5): 546-556.

Li, X., M. Inoue, D. R. Reed, T. Huque, R. B. Puchalski, M. G. Tordoff, Y. Ninomiya, G. K. Beauchamp and A. A. Bachmanov (2001). "High-resolution genetic mapping of the saccharin preference locus (Sac) and the putative sweet taste receptor (T1R1) gene (Gpr70) to mouse distal Chromosome 4." Mamm Genome **12**(1): 13-16.

Lin, C., et al., (2020). "Genetics of mouse behavioral and peripheral neural responses to sucrose." In preparation.

Lin, C., B. D. Fesi, M. Marquis, N. P. Bosak, A. Lysenko, M. A. Koshnevisan, F. F. Duke, M. L. Theodorides, T. M. Nelson, A. H. McDaniel, M. Avigdor, C. J. Arayata, L. Shaw, A. A. Bachmanov and D. R. Reed (2017). "Adiposity QTL Adip20 decomposes into at least four loci when dissected using congenic strains." PLoS One **12**(12): e0188972.

Lin, C., B. D. Fesi, M. Marquis, N. P. Bosak, A. Lysenko, M. A. Koshnevisan, F. F. Duke, M. L. Theodorides, T. M. Nelson, A. H. McDaniel, M. Avigdor, C. J. Arayata, L. Shaw, A. A. Bachmanov and D. R. Reed (2018). "Burly1 is a mouse QTL for lean body mass that maps to a 0.8-Mb region of chromosome 2." Mamm Genome.

Lin, C., B. D. Fesi, M. Marquis, N. P. Bosak, M. L. Theodorides, M. Avigdor, A. H. McDaniel, F. F. Duke, A. Lysenko, A. Khoshnevisan, B. R. Gantick, C. J. Arayata, T. M. Nelson, A. A. Bachmanov and D. R. Reed (2015). "Body composition qtls identified in intercross populations are reproducible in consomic mouse strains." PLoS One **10**(11): e0141494.

Lopez Del Amo, V., M. Palomino-Schatzlein, M. Seco-Cervera, J. L. Garcia-Gimenez, F. V. Pallardo, A. Pineda-Lucena and M. I. Galindo (2017). "A Drosophila model of GDAP1 function reveals the involvement of insulin signalling in the mitochondria-dependent neuromuscular degeneration." Biochim Biophys Acta **1863**(3): 801-809.

Lu, K., A. H. McDaniel, M. G. Tordoff, X. Li, G. K. Beauchamp, A. A. Bachmanov, D. A. VanderWeele, C. D. Chapman, N. K. Dess, L. Huang, H. Wang and D. R. Reed (2005). "No relationship between sequence variation in protein coding regions of the *Tas1r3* gene and saccharin preference in rats." Chem Senses **30**(3): 231-240.

Lush, I. E. (1989). "The genetics of tasting in mice. VI. Saccharin, acesulfame, dulcin and sucrose." Genetical Research **53**: 95-99.

Max, M., Y. G. Shanker, L. Huang, M. Rong, Z. Liu, F. Campagne, H. Weinstein, S. Damak and R. F. Margolskee (2001). "*Tas1r3*, encoding a new candidate taste receptor, is allelic to the sweet responsiveness locus *Sac*." Nat Genet **28**(1): 58-63.

Mi, H., A. Muruganujan and P. D. Thomas (2013). "PANTHER in 2013: modeling the evolution of gene function, and other gene attributes, in the context of phylogenetic trees." Nucleic Acids Res **41**(Database issue): D377-386.

Montmayeur, J. P., S. D. Liberles, H. Matsunami and L. B. Buck (2001). "A candidate taste receptor gene near a sweet taste locus." Nat Neurosci **4**(5): 492-498.

Mootha, V. K., J. Bunkenborg, J. V. Olsen, M. Hjerrild, J. R. Wisniewski, E. Stahl, M. S. Bolouri, H. N. Ray, S. Sihag, M. Kamal, N. Patterson, E. S. Lander and M. Mann (2003). "Integrated

analysis of protein composition, tissue diversity, and gene regulation in mouse mitochondria."
<u>Cell</u> **115**(5): 629-640.
Murovets, V. O., A. A. Bachmanov and V. A. Zolotarev (2015). "Impaired Glucose Metabolism in
Mice Lacking the Tas1r3 Taste Receptor Gene." <u>PLoS One</u> **10**(6): e0130997.
Nelson, G., M. A. Hoon, J. Chandrashekar, Y. Zhang, N. J. Ryba and C. S. Zuker (2001).
"Mammalian sweet taste receptors." <u>Cell</u> **106**(3): 381-390.
Nemazanyy, I., B. Blaauw, C. Paolini, C. Caillaud, F. Protasi, A. Mueller, T. Proikas-Cezanne, R.
C. Russell, K. L. Guan, I. Nishino, M. Sandri, M. Pende and G. Panasyuk (2013). "Defects of
Vps15 in skeletal muscles lead to autophagic vacuolar myopathy and lysosomal disease."
<u>EMBO Mol Med</u> **5**(6): 870-890.
Olsson, A. H., B. T. Yang, E. Hall, J. Taneera, A. Salehi, M. D. Nitert and C. Ling (2011).
"Decreased expression of genes involved in oxidative phosphorylation in human pancreatic
islets from patients with type 2 diabetes." <u>Eur J Endocrinol</u> **165**(4): 589-595.
Phillips, T. J., J. C. Crabbe, P. Metten and J. K. Belknap (1994). "Localization of genes affecting
alcohol drinking in mice." <u>Alcohol Clin Exp Res</u> **18**(4): 931-941.
Ramirez, I. (1987). "When does sucrose increase appetite and adiposity?" <u>Appetite</u> **9**(1): 1-19.
Raymond E Soccio, Z. L., Eric R Chen, Yee Hoon Foong, Kiara K Benson, Joanna R Dispirito,
Shannon E Mullican, Matthew J Emmett, Erika R Briggs, Lindsey C Peed, Richard K Dzeng,
Carlos J Medina, Jennifer F Jolivert, Megan Kissig, Satyajit R Rajapurkar, Manashree Damle,
Hee-Woong Lim, Kyoung-Jae Won, Patrick Seale, David J Steger, Mitchell A Lazar (2017).
"Targeting PPARγ in the Epigenome Rescues Genetic Metabolic Defects in Mice." <u>J Clin Invest.</u>
<u>2017 Apr 3; 127(4): 1451–1462.</u>
Reed, D. R., S. Li, X. Li, L. Huang, M. G. Tordoff, R. Starling-Roney, K. Taniguchi, D. B. West, J.
D. Ohmen, G. K. Beauchamp and A. A. Bachmanov (2004). "Polymorphisms in the taste
receptor gene (*Tas1r3*) region are associated with saccharin preference in 30 mouse strains." <u>J</u>
<u>Neurosci</u> **24**(4): 938-946.
Robinson, M. D., D. J. McCarthy and G. K. Smyth (2010). "edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data." <u>Bioinformatics</u> **26**(1): 139-140.
Sainz, E., J. N. Korley, J. F. Battey and S. L. Sullivan (2001). "Identification of a novel member
of the T1R family of putative taste receptors." <u>J Neurochem</u> **77**(3): 896-903.
Sclafani, A. (1987). "Carbohydrate-induced hyperphagia and obesity in the rat: effects of
saccharide type, form, and taste." <u>Neuroscience and Biobehavioral Reviews</u> **11**(2): 155-162.
Sclafani, A. (2006). "Oral, post-oral and genetic interactions in sweet appetite." <u>Physiol Behav.</u>
Sclafani, A. and S. Mann (1987). "Carbohydrate taste preferences in rats: glucose, sucrose,
maltose, fructose and polycose compared." <u>Physiol Behav</u> **40**(5): 563-568.
Shao, H., D. S. Sinasac, L. C. Burrage, C. A. Hodges, P. J. Supelak, M. R. Palmert, C. Moreno,
A. W. Cowley, Jr., H. J. Jacob and J. H. Nadeau (2010). "Analyzing complex traits with congenic
strains." <u>Mamm Genome</u> **21**(5-6): 276-286.
Simcox, J. A., T. C. Mitchell, Y. Gao, S. F. Just, R. Cooksey, J. Cox, R. Ajioka, D. Jones, S. H.
Lee, D. King, J. Huang and D. A. McClain (2015). "Dietary iron controls circadian hepatic
glucose metabolism through heme synthesis." <u>Diabetes</u> **64**(4): 1108-1119.
Simon, B. R., S. D. Parlee, B. S. Learman, H. Mori, E. L. Scheller, W. P. Cawthorn, X. Ning, K.
Gallagher, B. Tyrberg, F. M. Assadi-Porter, C. R. Evans and O. A. MacDougald (2013).
"Artificial sweeteners stimulate adipogenesis and suppress lipolysis independently of sweet
taste receptors." <u>J Biol Chem</u> **288**(45): 32475-32489.
Smyth, G. K. (2005). "Limma: linear models for microarray data. In: Bioinformatics and
Computational Biology Solutions using R and Bioconductor, R. Gentleman, V. Carey, S. Dudoit,
R. Irizarry, W. Huber (eds.), Springer, New York, pages 397-420. ."
Tarantino, L. M., G. E. McClearn, L. A. Rodriguez and R. Plomin (1998). "Confirmation of
quantitative trait loci for alcohol preference in mice." <u>Alcohol Clin Exp Res</u> **22**(5): 1099-1105.

Tian, L., G. Peng, J. M. Parant, V. Leventaki, E. Drakos, Q. Zhang, J. Parker-Thornburg, T. J.
Shackleford, H. Dai, S. Y. Lin, G. Lozano, G. Z. Rassidakis and F. X. Claret (2010). "Essential
roles of Jab1 in cell survival, spontaneous DNA damage and DNA repair." <u>Oncogene</u> **29**(46):
6125-6137.
Tordoff, M. G., L. K. Alarcon and M. P. Lawler (2008). "Preferences of 14 rat strains for 17 taste
compounds." <u>Physiol Behav</u> **95**(3): 308-332.
Tordoff, M. G., D. R. Reed and H. Shao (2008). "Calcium taste preferences: genetic analysis
and genome screen of C57BL/6J x PWK/PhJ hybrid mice." <u>Genes Brain Behav</u> **7**(6): 618-628.
Tordoff, M. G. a. B., A.A. (2001). "http://www.monell.org/MMTPP, Monell Chemical Senses
Center. Monell Mouse Taste Phenotyping Project.".
Truett, G. E., P. Heeger, R. L. Mynatt, A. A. Truett, J. A. Walker and M. L. Warman (2000).
"Preparation of PCR-quality mouse genomic DNA with hot sodium hydroxide and tris
(HotSHOT)." <u>Biotechniques</u> **29**(1): 52, 54.
Vadasz, C., M. Saito, A. Balla, I. Kiraly, C. Vadasz, 2nd, B. Gyetvai, E. Mikics, D. Pierson, D.
Brown and J. C. Nelson (2000). "Mapping of quantitative trait loci for ethanol preference in
quasi-congenic strains." <u>Alcohol</u> **20**(2): 161-171.
von Holstein-Rathlou, S., L. D. BonDurant, L. Peltekian, M. C. Naber, T. C. Yin, K. E. Claflin, A. I.
Urizar, A. N. Madsen, C. Ratner, B. Holst, K. Karstoft, A. Vandenbeuch, C. B. Anderson, M. D.
Cassell, A. P. Thompson, T. P. Solomon, K. Rahmouni, S. C. Kinnamon, A. A. Pieper, M. P.
Gillum and M. J. Potthoff (2016). "FGF21 mediates endocrine control of simple sugar intake and
sweet taste preference by the liver." <u>Cell Metab</u> **23**(2): 335-343.
von Mering, C., L. J. Jensen, B. Snel, S. D. Hooper, M. Krupp, M. Foglierini, N. Jouffre, M. A.
Huynen and P. Bork (2005). "STRING: known and predicted protein-protein associations,
integrated and transferred across organisms." <u>Nucleic Acids Res</u> **33**(Database issue): D433-437.
Yalcin, B., K. Wong, A. Agam, M. Goodson, T. M. Keane, X. Gan, C. Nellaker, L. Goodstadt, J.
Nicod, A. Bhomra, P. Hernandez-Pliego, H. Whitley, J. Cleak, R. Dutton, D. Janowitz, R. Mott, D.
J. Adams and J. Flint (2011). "Sequence-based characterization of structural variation in the
mouse genome." <u>Nature</u> **477**(7364): 326-329.
Yang, H., J. R. Wang, J. P. Didion, R. J. Buus, T. A. Bell, C. E. Welsh, F. Bonhomme, A. H. Yu,
M. W. Nachman, J. Pialek, P. Tucker, P. Boursot, L. McMillan, G. A. Churchill and F. P. de
Villena (2011). "Subspecific origin and haplotype diversity in the laboratory mouse." <u>Nat Genet</u>
43(7): 648-655.
Yee, K. K., S. K. Sukumaran, R. Kotha, T. A. Gilbertson and R. F. Margolskee (2011). "Glucose
transporters and ATP-gated K+ (KATP) metabolic sensors are present in type 1 taste receptor 3
(T1r3)-expressing taste cells." <u>Proc Natl Acad Sci U S A</u> **108**(13): 5431-5436.

Figure 1. Identification of *Scon3* (a main-effect QTL on Chr9) and *Scon4* (an interacting QTL on Chr1) in C57BL/6ByJ x 129P3/J.C57BL/6ByJ-*Tas1r3* F$_2$ mice (N=279)

(**A**) A genome-wide scan of main-effect QTLs for 300 mM sucrose intake. The pink horizontal dashed line represents the significant linkage threshold determined by 1000 permutation tests (threshold=3.64). The black curves trace the logarithm of the odds (LOD) scores (y-axis). The genotype effect of the significant loci was confirmed in a general linear model using habitual water intake as a covariate and a post hoc comparison of least-square means (LSM) among the genotypes of these loci tested. Different letters above horizontal lines depicting LSM ± standard error indicate significant differences in daily sucrose consumption by genotype group. (**B**) Heat map for a two-dimensional genome scan with a two-QTL model. The maximum LOD score for the full model (two QTLs plus an interaction) is indicated in the lower right triangle. The maximum LOD score for the interaction model is indicated in the upper left triangle. A color-coded scale displays values for the interaction model (LOD threshold=6.5) and the full model (LOD threshold=9.1) on the left and right, respectively. A yellow circle in the upper left section shows significant interaction (LOD=13.5) between QTLs on *Scon3* and *Scon4*. (**C)** LSM ± standard error for sucrose intakes of the F$_2$ mice grouped by genotypes of the markers with the highest epistatic interaction (*D1Mit316* of *Scon4* and *rs3023231 of Scon3*). Different letters indicate statistically significant differences between genotypes (Tukey's HSD test: *p*<0.05; for details see the Methods section).

Figure 2. Confirmation of the *Scon3* and *Scon4* loci, and their interaction in backcross (A), consomic (B-E), and congenic (F-J) mice

(**A**) Backcross mice. *Scon3* was detected in the pooled backcross mice (N$_3$, N$_4$, N$_5$, and N$_6$; *N*=438). The x-axis is the location of the markers in Mb (see **S4 Table**) on mouse Chr9; the y-axis is the –log *p*-value of the statistical test by each marker genotype. The small blue bar under

the peak indicates the confidence interval (2 units of –$\log_{10}$ p-value drop from the peak). The

horizontal red line shows threshold of statistical significance, calculated using Bonferroni

correction [–log(0.05/68) ≈ 3.13]. The inset shows least-squares means (LSM) ± standard error

sucrose intake of mice grouped by peak marker genotype (*rs3023231*). (**B**) A schematic of

chromosomes of the 129.B6-double congenic strain, created by mating *Scon4* homozygous

congenic males with *Scon3* heterozygous congenic females. These double-congenic mice thus

have heterozygous genotypes (B6/129 for all mice) at the *Scon4* locus and segregating

genotypes (B6/129 or 129/129) at the *Scon3* locus. (**C-F**) Consomic mice. (**C**) A schematic of

the comparisons of mouse consomic strains and their host inbred strains, as well as the

corresponding genotypes of *Scon4* and *Scon3* in each mouse strain. (**D**) LSMs and their

standard errors for the consomic strains 129.B6-Chr1, 129.B6-Chr9, and B6.129-Chr9,

compared with their host inbred strain (129 or B6). (**E**) Effect size in Cohen's D for each group

computed by comparing homozygous consomic genotypes vs host inbred genotypes. The large

difference in effect sizes between the two reciprocal consomic strains, 129.B6-Chr9 and

B6.129-Chr9, supports the epistatic interaction of the homozygous alleles for *Scon3* and *Scon4*:

effects of *Scon3* are larger when they are combined with B6/B6 *Scon4* genotype than with

129/129 *Scon4* genotype (compare **Figure 1C**). (**F**) The interaction effect of homozygous alleles

between *Scon4* and *Scon3* was confirmed in the consomic and host inbred strains. Different

letters indicate statistically significant differences between genotypes. *Scon4*: $F_{(1,\ 225)}$=36.8,

p<0.00001; *Scon3*: $F_{(1,\ 225)}$=166.9, p<0.00001; *Scon4 × Scon3*: $F_{(1,\ 225)}$=19.4, p<0.0001. (**G-J**)

Single- and double-congenic mice. (**G**) Comparisons of littermates from the single- (top two

panels) and double- (bottom panel) congenic strains: schematic of chromosomes (left) and

corresponding *Scon4* and *Scon3* genotypes (right) in each group of mice. (**H**) LSM and their

standard errors for the congenic strains: 129.B6-*Scon4*, 129.B6-*Scon3*, and 129.B6 double

congenics, with one copy of B6 donor fragment (B6/129), compared with their littermates

without the B6-donor fragment (129/129). (**I**) Effect size in Cohen's D computed for each group

of comparisons by comparing heterozygous congenic genotype (B6/129) vs their littermates'

genotype (129/129). A large difference between the effect size observed for the double-

congenic strain and the effect sizes for 129.B6-*Scon4* or 129.B6-*Scon3* confirmed the epistatic

interaction of the heterozygous alleles between *Scon3* and *Scon4*. (**J**) The interaction effect of

heterozygous alleles between *Scon4* and *Scon3* was confirmed in congenic strains: the effects

of *Scon3* are larger when they are combined with 129/129 *Scon4* genotype (circles) than with

B6/129 *Scon4* genotype (squares; see **Figure 1C**). Different letters indicate statistically

significant differences between genotypes.

Figure 3. High-resolution mapping of the *Scon4* locus in congenic substrains

Data are from two 129.B6-*Scon4* heterozygous congenic substrains (C1 and C2) with partially

overlapping donor fragments (for markers and their positions, see **S6 Table**). (**A**) Interval

mapping of genotype-phenotype associations: using a general linear model, we computed

associations between marker genotypes and 300 mM sucrose intake in a combined group

including all 129.B6-*Scon4* heterozygous congenic mice (in this analysis, mice were grouped by

genotype at each marker rather than by substrain). The x-axis gives marker positions in Mb on

Chr1; the y-axis, $-\log_{10}$-transformed *p*-values. The horizontal red line shows significance

threshold [$-\log_{10}(0.05/N)$, where *N* is number of markers tested]; CI = confidence interval. (**B**)

Donor regions of the congenic substrains are indicated by the black bar (for the C1 strain that

retained *Scon4*) or gray bar (for the C2 strain that did not retain *Scon4*). Blue bars indicate the

regions contributed by the host (129) strain; the *Scon4* critical region (red dashed rectangle; 0-

25.4 Mb from centromere to *rs13475771*) is retained in the donor region of the C1 (*Scon4*-

positive) strain but not in C2 (*Scon4*-negative) strain. (**C and D**) Intake of 300 mM sucrose by

the *Scon4*-positive (C1; **C**) and *Scon4*-negative (C2; **D**) strains. Strain comparisons were made

using general linear model.

Figure 4. The *Scon4* homozygous allele effect on voluntary sucrose intake

The *Scon4* locus affects 300 mM sucrose consumption in homozygous 129.B6-*Scon4* congenic mice (B6/B6; C1) vs host inbred 129 mice (129/129). Mice with *Scon4* (B6/B6) and *Scon3* (129/129) have lower sucrose intakes than mice with *Scon4* (129/129) and *Scon3* (129/129). The effect size of Cohen's D is shown in right blue bar.

Figure 5. High-resolution mapping of the *Scon3* locus in congenic substrains

(**A**) For average daily sucrose intake, we computed trait-marker associations using a general linear model among all 129.B6-*Scon3* congenic mice grouped by genotype at each marker. The x-axis gives marker positions in Mb on chromosome 9 (Chr9); for other details see **Figure 3**. (**B**) Donor region of each congenic strain: black bars, strains that retained the *Scon3* locus; gray bars, strains that lost the locus; blue bars, region contributed by the host strain. (**C**) Strain comparisons with sample size (*n*) of each genotype within each strain. C3 and C4, *Scon3*-negative strains; C5–C9, *Scon3*-positive strains that share a common region (red dashed rectangle in **A and B**: 1.2-Mb donor region between 105.7 and 106.9 Mb, from *rs33653996* to *rs3023231*) that the *Scon3*-negative strains (C3 and C4) do not. (**D**) The 1.2 Mb *Scon3* critical region contains X noncoding RNA genes, Y protein-coding genes, Z pseudogenes, and N processed transcripts (Ensemble Mouse Genome Brower @GRCm38.p5 (Hubbard, Aken et al. 2007)).

Figure 6. The effects of *Scon4* are largest for the more concentrated sugar solutions

Average daily intake (left panels) and preference (right panels) for sucrose (**A**), glucose (**B**), and fructose (**C**) by homozygous 129.B6-*Scon4* congenic mice (C1; for sucrose, *N*=19; for glucose, *N*=10; for fructose, *N*=10) and host inbred 129 mice (all *N*=10) in two-bottle 48-hr preference tests. Error bars show SEM. *p<0.05, significant difference between the congenic and host inbred mice (post hoc tests). Blue, genotypes for *Scon4* and *Scon3*; black, control 129 inbred mice.

Figure 7. The *Scon4* genotype affects fuel oxidation but not food intake

Metabolic assessments in *Scon4* congenic mice (black bars; N=17; C1) and control 129 inbred

mice (gray bars; N=8) at 8 and 36 weeks of age. (**A**) Food intake: 8 weeks, $t_{(1,23)}$=1.11, p=0.282;

36 weeks, $t_{(1,23)}$=1.10, p=0.284. (**B**) Water intake: 8 weeks, $t_{(1,23)}$=0.54, p=0.597; 36 weeks,

$t_{(1,23)}$=0.96, p=0.348. (**C**) Heat production: 8 weeks, $t_{(1,23)}$=5.22, p<0.001; 36 weeks, $t_{(1,23)}$=3.49,

p<0.01. (**D**) Respiratory exchange ratio (RER): 8 weeks, $t_{(1,23)}$=5.22, p<0.001; 36 weeks,

$t_{(1,23)}$=3.49, p<0.01. (**E**) Oxygen consumption (VO$_2$): 8 weeks, $t_{(1,23)}$=4.71, p<0.001; 36 weeks,

$t_{(1,23)}$=3.16 , p<0.01. (**F**) Carbon dioxide (VCO$_2$): 8 weeks, $t_{(1,23)}$=4.03, p<0.001; 36 weeks,

$t_{(1,23)}$=0.53, p=0.598. (**G**) Activity over the entire 24-hr light/dark cycle: 8 weeks, $t_{(1,23)}$=1.85,

p=0.077; 36 weeks, $t_{(1,23)}$=0.00, p=0.999. Data are mean ± SE.

Figure 8. Gene expression profiling shows differential expression of candidate genes

within *Scon3* and *Scon4* regions between 129 and B6 inbred mouse strains

Data are from microarray-based gene expression analysis for mouse stratum (**A**) and bulk

RNASeq analyses for brown (**B**) and white (**C**) fat. Red dots show differentially expressed genes

with twofold changes between two inbred strains. The horizontal dash lines show the corrected

significance threshold (FDR=0.05).

Supplemental Figures

S1 Figure. Correlations between sucrose intake (ml/d) and phenotype within each genetic

mapping population

Data are for body weight (BW; g) by sex (M, F), average daily habitual water intake (Wat; ml/d,

M+F combined), and age (weeks, M+F combined). For sample sizes, see **S1 Table**.

S2 Figure. F$_2$ mice (*N*=279) showed genotype effects of the *Scon3 and Scon4* loci on

intake or preference for taste solutions

Group means that do not differ either have no letter subscript or share a letter subscript in

common. A=B6/B6, H=129/B6, B=129/129.

S3 Figure. No *Scon3* or *Scon4* genotype effects on average daily water intake in F_2 (A, B),

congenic (C, D), or backcross (E) mice

For sample sizes, see **S1 Table**. A=B6/B6; H=129/B6; B=129/129. Average daily water intake

was analyzed using general linear model with genotype and sex as fixed factors and body

weight as covariate. Within each panel, groups that do not share a common subscript letter

differ significantly (post hoc tests).

S4 Figure. Heat map of Pearson correlations for taste solution intakes and preference

scores of the F_2 mice (*N*=279)

S5 Figure. Chromosomes 1 and 9 harbor QTLs for the consumption of other taste

solutions.

A significant difference in intake and/or preference of a solution between a consomic strain and

host inbred 129 or B6 strain indicates presence of a QTL on this chromosome for the taste

solution ($p<0.05$). For intakes, we used a general linear model, with strain as fixed factor and

habitual water intake as a covariate, followed by post hoc test. For preference scores, we used

t-tests.

S6 Figure. Both *Scon3* and *Scon4* showed no genotype effect on the responses of the

chorda tympani gustatory nerve to lingual application of taste stimuli

We compared the heterozygous 129.B6-*Scon4* congenic mice (C1; B6/129; *N*=5) or

homozygous 129.B6-*Scon3* congenic mice (C7: B6/B6; *N*=5) and host inbred 129 mice

(129/129; *N*=5) using Mann-Whitney *U*-tests for individual solutions. We observed no significant

genotype differences ($p>0.05$) in responses of the chorda tympani gustatory nerve for the taste

stimuli (see **S8 Table**). Values are medians ± median absolute deviations.

S7 Figure. **Effects of the *Scon4* genotype on intakes and preference for other taste**

compounds

Data are for 20 mM saccharin (**A**), 30 mM glycine (**B**), 300 mM NaCl (**C**), 300 mM MSG (**D**), 3%

ethanol (**E**), and 10% ethanol (**F**) by homozygous 129.B6-*Scon4* congenics (C1: B6/B6) and

host inbred 129 mice (129/129) in two-bottle tests.

S8 Figure. No effect of the *Scon4* genotype on intakes and preferences for other caloric

compounds

Data are for 4.6% soybean oil (**A**), 6% cornstarch (**B**), or 10.5% maltodextrin (**C**) by

homozygous 129.B6-*Scon4* congenics (C1: B6/B6; *N*=19) and host inbred 129 mice (129/129;

N=10) in two-bottle tests.

S9 Figure. No effect of the *Scon4* genotype on mouse body composition at 8 or 36 weeks

of age

Mouse body composition as measured by magnetic resonance (**A**; 8 weeks: *N*=32, males=15,

females=17; 36 weeks: *N*=22, males=9, females=13) or by DEXA (*N*=26, males=12,

females=14): homozygous *Scon4* congenics (C1; B6/B6) vs host inbred 129 controls (129/129).

BMC=bone mineral content (g); BMD=bone mineral density. Groups that do not share a

common subscript letter differ by post hoc test.

S10 Figure. **Effect of *Scon4* genotype on organ and fat weight of mice dissected at 180**

days of age

No effect was observed, except for subscapular adipose depot ($F_{(1,23)}$=18.5, p<0.001) and heart

($F_{(1,23)}$=22.5, p<0.0001). Homozygous *Scon4* congenics (C1; B6/B6; males=7, females=12) vs

host inbred 129 controls (129/129; males=5, females=2). Groups that do not share a common

subscript letter differ by post hoc tests.

S11 Figure. No effect of *Scon4* genotype on body weight changes when mice were fed chow and drank sweetened water

Water was sweetened with sucrose (**A-F**), glucose (**G**), or fructose (**H**) solution. Mice drank voluntarily for 4-12 days. Data are for F$_2$ mice (**A, B**; N=279), consomic mice (**C-E**; 129.B6-Chr1 and 129 controls: N=105; 129.B6-Chr9 and 129 controls: N=87; B6.129-Chr9 and B6 controls: N=86), or congenic mice (**F-H**; sucrose: N=105; glucose: N=20; fructose: N=20). Groups that do not share a common subscript letter differ by post hoc test. A=B6/B6; H=129/B6; B=129/129.

S12 Figure. No effect of *Scon4* genotype on tolerance to glucose or insulin

Glucose was administered intraperitoneally (IP GTT; **A**) or intragastrically (IG GTT; **B**) or insulin was administered (**C**) in homozygous *Scon4* congenic mice (black squares; N=21) and the host inbred control mice (gray circles; N=12) at 8 and 36 weeks of age. Data are mean ± SEM of genotype; significance of the genotype effect was evaluated by post hoc tests. *p<0.05. The statistics are as follows:

IP, 8 weeks, genotype $F_{(1,30)}$=1.52, p=0.227; time $F_{(7,210)}$=73.5, p<0.0001, genotype × time interaction $F_{(7,210)}$=1.70, p=0.111; 36 weeks, genotype $F_{(1,24)}$=0.19, p=0.665; time $F_{(7,168)}$=107.9, p<0.0001; genotype × time interaction $F_{(7,168)}$=1.70, p=0.112.

IG, 8 weeks, genotype $F_{(1,31)}$=1.95, p=0.173; time $F_{(7, 217)}$=69.4, p<0.0001; genotype × time interaction ($F_{(7,217)}$=0.18, p=0.989); 36 weeks, genotype $F_{(1,24)}$=0.065, p=0.800; time $F_{(7,168)}$=62.2, p<0.0001; genotype × time interaction $F_{(7, 168)}$=0.53, p=0.813.

ITT, 8 weeks, genotype $F_{(1,28)}$=8.68, p=0.006; time $F_{(5, 140)}$=101.2, p<0.0001; genotype × time interaction $F_{(5,140)}$=3.70, p=0.003; 36 weeks, genotype $F_{(1,23)}$=1.012, p=0.325; time $F_{(5,115)}$=80.1, p<0.0001; genotype × time interaction $F_{(5,115)}$=1.34, p=0.253.

S13 Figure. Gene ontology analysis and protein-protein associations for genes within the *Scon3* and *Scon4* regions

(**A**) Gene ontology analysis of the biological process for genes within the *Scon4* and *Scon3*

regions. The Ensembl mouse gene identifier IDs (*N*=195; no Ensembl gene identifier IDs were

available for the 45 unknown genes) were batch-uploaded into the Panther database for *Mus*

musculus, and in total, 92 genes and 72 hits (gene IDs matched in the public database) were

processed. (**B**) Protein-protein associations for genes within *Scon4* and *Scon3* (**S10 Table**)

were identified by searching protein names using information from high-throughput experimental

data, mining of literature databases, and predictions based on genomic context analysis.

A Sucrose

B Glucose

C Fructose

Scon4 *Scon3*
B6/B6 129/129
129/129 129/129

A — Food intake, g/24h; ns; ns

B — Water intake, ml/24h; ns; ns

C — Heat, kcal/kg lean/h; $p<0.001$; $p<0.01$

D — RER; $p<0.05$; $p<0.0001$

E — VO$_2$ ml/kg lean/h; $p<0.001$; $p<0.01$

F — VCO$_2$ ml/kg lean/h; $p<0.001$; ns

G — Activity (beam breaks/h); ns; ns

8 wks   36 wks

Scon4   *Scon3*

B6/B6   *129/129*

129/129   *129/129*