Initial Characterization of the Two ClpP Paralogs of *Chlamydia*

trachomatis Suggests Unique Functionality for Each

Nicholas A. Wood[1,2], Krystal Chung[3], Nathalia Rodrigues de Almeida[4], Martin Conda-Sheridan[4],

Derek J. Fisher[3], Scot P. Ouellette[1#]

[1]Department of Pathology and Microbiology, College of Medicine, University of Nebraska Medical

Center, Omaha, NE

[2]Division of Basic Biomedical Sciences, Sanford School of Medicine, University of South Dakota,

Vermillion, SD

[3]Department of Microbiology, Southern Illinois University Carbondale, Carbondale, IL

[4]Department of Pharmaceutical Sciences, College of Pharmacy, University of Nebraska Medical

Center, Omaha, NE

Keywords: *Chlamydia*, differentiation, protein turnover, protein quality control, Clp protease

Running Title: ClpP proteases of *Chlamydia*

#Corresponding Author:

Department of Pathology and Microbiology, College of Medicine, University of Nebraska Medical

Center, 985900 Nebraska Medical Center (DRC2 5022), Omaha, NE

Tel: +1-402-559-0763 Fax: +1-402-559-5900 Email: scot.ouellette@unmc.edu

Abstract

Chlamydia is an obligate intracellular bacterium that differentiates between two distinct functional and morphological forms during its developmental cycle: elementary bodies (EBs) and reticulate bodies (RBs). EBs are non-dividing, small electron dense forms that infect host cells. RBs are larger, non-infectious replicative forms that develop within a membrane-bound vesicle, termed an inclusion. Given the unique properties of each developmental form of this bacterium, we hypothesized that the Clp protease system plays an integral role in proteomic turnover by degrading specific proteins from one developmental form or the other. *Chlamydia* has five uncharacterized *clp* genes: *clpX*, *clpC*, two *clpP* paralogs, and *clpB*. In other bacteria, ClpC and ClpX are chaperones that unfold and feed proteins into the ClpP protease to be degraded, and ClpB is a deaggregase. Here, we focused on characterizing the ClpP paralogs. Transcriptional analyses and immunoblotting determined these genes are expressed mid-cycle. Bioinformatic analyses of these proteins identified key residues important for activity. Over-expression of inactive *clpP* mutants in *Chlamydia* suggested independent function of each ClpP paralog. To further probe these differences, we determined interactions between the ClpP proteins using bacterial two-hybrid assays and Native gel analysis of recombinant proteins. Homotypic interactions of the ClpP proteins, but not heterotypic interactions between the ClpP paralogs, were detected. Interestingly, ClpP2, but not ClpP1, protease activity was detected *in vitro*. This activity was stimulated by antibiotics known to activate ClpP, which blocked chlamydial growth. Our data suggest the chlamydial ClpP paralogs likely serve distinct and critical roles in this important pathogen.

Words: 250/250

Importance

Chlamydia trachomatis is the leading cause of preventable infectious blindness and of

bacterial sexually transmitted infections worldwide. Chlamydiae are developmentally regulated,

obligate intracellular pathogens that alternate between two functional and morphologic forms with

distinct repertoires of proteins. We hypothesize that protein degradation is a critical aspect to the

developmental cycle. A key system involved in protein turnover in bacteria is the Clp protease

system. Here, we characterized the two chlamydial ClpP paralogs by examining their expression in

Chlamydia, their ability to oligomerize, and their proteolytic activity. This work will help understand

the evolutionarily diverse Clp proteases in the context of intracellular organisms, which may aid in

the study of other clinically relevant intracellular bacteria.

Words: 114/120

Introduction

Chlamydia trachomatis is one of the most prevalent human sexually transmitted infections and the leading cause of preventable infectious blindness worldwide (1). Of particular note are the negative effects associated with untreated *C. trachomatis* infections. Because of the asymptomatic nature of 60-80% of cases (2), infection by this organism can lead to complications such as pelvic inflammatory disease, which can in turn lead to infertility and ectopic pregnancies in women (3). An infant can also acquire conjunctivitis during birth should the mother be infected, which can result in irreversible blindness if left untreated.

C. trachomatis is an obligate intracellular pathogen with a complex developmental cycle (see (4) for detailed review). These pathogenic bacteria differentiate between two distinct functional and morphological forms over the course of 48 to 72 hours, depending on the strain. The electron dense elementary body (EB) is the infectious but non-dividing form that is characterized by its histone-compacted chromosome (5) and cysteine-rich, disulfide-crosslinked outer membrane (6). The reticulate body (RB) is the non-infectious and replicating form that has a relaxed chromosome structure and lipid-based cell walls that are sensitive to disruption. The typical size of each form is ~0.3 μm and ~1 μm, respectively (4). A single EB initiates infection of a host cell by attaching to the plasma membrane and inducing uptake into the cell by type III secreted effectors (7) (8). The EB remains within a host-derived vesicle, which is diverted from the endocytic pathway, and differentiates into the much larger RB. The vesicle is rapidly modified into a pathogen-specified parasitic organelle termed an inclusion (9) (10). Proliferation of this RB within the inclusion follows until such time that secondary differentiation occurs, and the RBs begin to condense their genome and crosslink their outer membrane. After a significant amount of EBs have accumulated, they exit the cell for infection of surrounding cells. Proteomic and transcriptional analyses indicate that EBs and RBs have distinct patterns of gene expression and protein repertoires (11) (12) (13) (14). Given

the striking phenotypic differences between the two developmental forms, we hypothesize that

protein turnover plays a key role in differentiation in addition to general maintenance of bacterial

homeostasis during both normal and persistent growth modes ([15](15)) ([16](16)) ([17](17)).

Even though *Chlamydiae* exhibit such a complex developmental cycle and evade the host

immune system during chronic infections, these bacteria have undergone significant reductive

evolution to eliminate unnecessary genes. This in turn suggests that most genes retained by this

organism likely serve an important function to bacterial fitness ([18](18)). Within its genome, *Chlamydia*

encodes homologs to four **c**aseino**l**ytic **p**rotease *clp* genes: *clpC*, *clpX*, *clpP* (*clpP1* and *clpP2*), and

clpB. ClpB is a suspected deaggregase, but does not appear to interact directly with other Clps for

proteomic turnover ([12](12)). ClpC and ClpX proteins are classified as AAA+ (**A**TPases **A**ssociated with

various cellular **A**ctivities) unfoldases that serve as adaptor proteins to linearize target proteins in an

ATP-dependent manner ([19](19)) ([20](20)). ATP binding, but not necessarily hydrolysis, potentiates

interaction between a homo-hexamer of these proteins and a ClpP oligomer in other organisms ([21](21))

([22](22)). ClpP is a serine protease that gives peptidase function to the resultant complex ([23](23)). ClpP

proteins have been shown to oligomerize into a tetradecameric complex of a stack of two heptamers

that can then perform proteolytic function of small peptide substrates. Upon interaction with the

adaptor protein hexamers, the N-terminal region of the ClpP complex is relaxed and stabilized,

allowing larger substrates into the proteolytic complex ([24](24)) ([25](25)).

Multiple pathogenic bacteria possess two *clpP* paralogs including *Pseudomonas aeruginosa*,

Mycobacterium tuberculosis, and *Listeria monocytogenes*,. The Baker lab characterized the dual

ClpP system of *P. aeruginosa* and demonstrated that each system may have a distinct function

contributing to virulence and fitness of the bacteria ([26](26)). While the group studying ClpP1/2 in

Mycobacterium showed heterologous interactions between these paralogs, these genes are co-

transcribed, suggesting coupled function ([27](27)). The dual ClpP proteases of *Listeria monocytogenes*

(Lm) were shown to form both hetero- and homotypic complexes ([28]) ([29]), yet the authors

demonstrated that Lm ClpP2 likely can function independently of Lm ClpP1. These studies, taken

together with the reductive evolution of the chlamydial genome, led us to hypothesize that the

chlamydial ClpPs may serve distinct and critical functions in the physiology of the organism.

To begin investigation of the role of the Clp protease system in chlamydial biology, we

initiated a series of studies to characterize the ClpP paralogs. All *clp* genes are transcribed as RB-

specific genes, and the translated product of *clpP2* was detected during the developmental cycle.

Bioinformatic and structural modeling analyses indicate that these proteins retain key residues

important for function as well as structure. Interestingly, over-expression of a mutant ClpP1 was

detrimental to chlamydial development whereas a mutant ClpP2 had no observable effect. To

explore the basis of this observation, we performed a series of *in vitro* and *in vivo* assays indicating

homo-oligomerization of each paralog. However, we did not detect hetero-oligomerization between

these paralogs. We also characterized the *in vitro* protease activity of each ClpP paralog and

observed that activity was only detected for ClpP2. Antibiotics known to activate ClpP proteases

stimulated the protease activity of ClpP2 only and blocked chlamydial growth. Combined, our data

suggest ClpP1 activity may be tightly regulated. We conclude that each chlamydial ClpP protease

serves a unique and independent role, either in differentiation or more typical physiological

processes.

Results

The *clp* genes are expressed as RB specific genes. We hypothesize, based on the arrangement of

the genes in the chromosome, that ClpC interacts with ClpP1 and that ClpX interacts with ClpP2.

This does not preclude each individual component from acting independently and is not mutually

exclusive to our hypothesis. We base our prediction of ClpP2X interaction due to the juxtaposition of

the two genes within the same operon (Fig. 1A). We then reasoned that ClpP1 would serve as the proteolytic subunit that would interact with ClpC, as both are encoded independently of each other in separate genomic contexts (Fig. 1A). Because chlamydiae are developmentally regulated bacteria, gene expression can differ from one developmental form to the other. Gene expression in *Chlamydia* can be broadly categorized into three different stages: early-, mid-, and late-developmental cycle genes (30). Early cycle genes are involved in initial differentiation from EB to RB and establishment of the nascent inclusion. Mid-cycle genes are RB specific and typically play a role in growth and division of the RBs as well as in inclusion modification. Late cycle genes function in differentiation from an RB to an EB or serve an important function in the secondary infection of a host cell. The specificity of a transcript to a particular point of the developmental cycle can give insight into the possible function of the encoded protein. To determine when during the developmental cycle the *clp* genes are transcribed, we measured the transcription of each gene over a time course of infection. Nucleic acid samples were collected for analysis at various time points of infection using wild-type *C. trachomatis* serovar L2. Our data indicate that the *clp* genes analyzed are transcribed in a pattern consistent with mid-cycle, RB specific function since the transcript levels peak after primary differentiation has occurred and a population of RBs has been established (i.e. 16 hpi; Fig. 1B). This pattern is distinct to that of canonical early or late cycle genes such as *euo* or *omcB* (31) (32) (33). The *clpX* transcript levels closely mirrored those of *clpP2* albeit with slightly lower transcript abundance, consistent with its genomic context 3' to *clpP2*.

We next sought to detect endogenous protein of the ClpP paralogs. Given the high level of homology between the *Pseudomonas aeruginosa* (Pa) ClpP1 and *C. trachomatis* (Ctr) ClpP2 (see Fig. 2), we speculated a polyclonal antibody developed against Pa_ClpP1 (kind gift of Dr. T. Baker, MIT) would detect Ctr ClpP2 by Western blot (see Suppl. Fig. S1). As seen, Ctr ClpP2 was detected from infected cell lysates starting at 16h post-infection using the polyclonal anti-Pa_ClpP1 antibody

(Fig. 1C). The polyclonal antibody developed against Pa_ClpP2 did not react with chlamydial

lysates or recombinant Ctr ClpP1 or ClpP2 (Suppl. Fig. S1). Given the relatively low levels of ClpP2

compared to the major outer membrane protein (MOMP; Fig. 1D), we cannot conclude that ClpP2 is

not present at earlier times or that an antibody specific to chlamydial ClpP2 would be more sensitive

to detect such lower levels. We are currently working towards developing antibodies against the

chlamydial Clp components. Overall, the expression data indicate that the Clp components are

expressed in the RB phase of growth and division. This does not, however, preclude these proteins

from having functions at other times during the developmental cycle.

Bioinformatics analyses identify key residues important for structure and function of each

chlamydial ClpP paralog. We hypothesized that each ClpP paralog of *C. trachomatis* serves

distinct functions in the biology and pathogenesis of this unique bacterium, despite their similarity in

expression patterns. To further test this hypothesis, we performed a series of bioinformatics analyses.

We began by looking at pairwise alignments of the chlamydial ClpP proteins against each other and

against ClpP homologs from other bacteria (Fig. 2A). Interestingly, each chlamydial ClpP shared

more homology to ClpP of *E. coli*, at 44% and 55% identity with Expect values of 4×10^{-51} and 2×10^{-83},

respectively, and to other ClpP homologs than to its paralog. This observation was supported by

the ability of the anti-Pa_ClpP1 antibody to recognize Ctr ClpP2 (Fig. 1C). The sequence identity

between Ctr ClpP1 and Ctr ClpP2 is 39% with an Expect value of 10^{-44}. Similarly, the two ClpP

paralogs of *P. aeruginosa* also share a low amount of similarity to each other compared to other

ClpP homologs via protein alignment (data not shown), and each serves a distinct function in *P.*

aeruginosa ([26](#)). Therefore, these observations support the likelihood that the chlamydial ClpP

paralogs have distinct roles.

A closer analysis of the chlamydial ClpP paralogs provides further evidence for unique

functionality of each protein. A 3-dimensional predicted structure alignment of Ctr ClpP1 and ClpP2

to *E. coli* ClpP indicates that the structural architecture would allow for homo-oligomer formation

and interaction with chaperones (Fig. 2B), though distinct differences in the oligomer interface may

prevent hetero-oligomerization (34). Additionally, the catalytic triad active site aligns well for all

ClpP homologs analyzed (Fig. 2B&C), with variation in structure being mostly limited to the N- and

C-termini. Also unique to chlamydial ClpP1 are distinct differences at residues that align to amino

acids critical to activation in other bacteria. An alanine at position 37 of Ctr ClpP1 may result in a

conformational difference in the P1 pre-activation complex that precludes P1/P2 interaction (35).

This may serve a regulatory role. Further supporting differential regulation is the lack of an aromatic

amino acid at position 57 of ClpP1 that is relatively well-conserved in other ClpP homologs,

suggesting that Ctr ClpP1 enters the "open" conformation differently than Ctr ClpP2. Ongoing

studies are investigating these residues in addition to others that are involved in complex activation.

Given the conserved catalytic residues and the structural similarity of the Ctr ClpP proteins to

homologs from a diverse set of bacteria, we conclude that the chlamydial ClpP proteins are *bona fide*

protease subunits. However, the predicted structural differences at the termini of the protein and in

other key residues suggests that the chlamydial ClpP proteins (i) do not interact with each other, (ii)

have specialized functions, and (iii) are regulated differently.

Overexpression of inactive ClpP1, but not ClpP2, negatively affects *Chlamydia*. Leveraging

recent advances in chlamydial genetics, we next determined the effect of over-expression of ClpP

proteins in *Chlamydia*. We designed and constructed plasmids encoding wild-type or inactive

mutants of each *clpP* paralog, fused to a 6xHis tag, under the control of an anhydrotetracycline

(aTc)-inducible promoter (36). Interestingly, we were unable to successfully transform *C.*

trachomatis serovar L2 with plasmids encoding the wild-type *clpP* genes. The reasons for this are

unclear (aside from the inherent difficulty in transforming this obligate intracellular bacterium) but

may indicate that leaky expression from the promoter was deleterious to the organism ([37]).

However, we successfully transformed *C. trachomatis* with plasmids encoding either ClpP1 S92A or

ClpP2 S98A active site serine mutants that are known to abolish protease activity (See Fig. 2B;

([38])). To assess the effect of over-expression of these inactive mutants, HEp2 cells were infected

with each transformant and expression was induced at 10 hours post-infection with 1 or 10 nM aTc.

Both 6- and 14-hour pulses were utilized to determine inclusion morphology at 16 and 24 hours post-

infection, respectively. At each time point and at the lower concentration of aTc inducer, the

localization of the ClpP proteins was within the bacterial cytosol. No noticeable effects were

detected after 6h of induction for the ClpP2 mutant, while the ClpP1 mutant induction appeared to

have a slight negative impact on the organisms. However, after 14h of over-expression of the mutant

ClpP1, noticeably smaller inclusion sizes were observed (Fig. 3A) whereas over-expression of the

mutant ClpP2 had no demonstrable effect (Fig. 3B). These data indicate that over-expression of

ClpP1 S92A is detrimental to chlamydiae. In support of this conclusion, we were unable to isolate a

clonal population of purified ClpP1 S92A transformants as we routinely observed bacterial

populations susceptible to antibiotic selection even after multiple passages. This observation

indicates that leaky expression may drive plasmid loss as has been observed in other chlamydial

transformation systems ([37], [39]).

The ClpP paralogs demonstrate homotypic, but not heterotypic, interactions. To better

understand the function of each ClpP paralog and why over-expression of ClpP1 S92A, but not

ClpP2 S98A, was detrimental to *Chlamydia*, we initiated a series of *in vivo* and *in vitro* biochemical

assays to characterize the properties of the ClpP proteins. Extensive evidence exists from other

bacterial systems to indicate that ClpP forms a tetradecameric complex composed of two heptameric

rings ([21-25](#), [40](#), [41](#)). To determine whether the chlamydial ClpP paralogs formed homo- and/or

heterooligomers, we first employed the bacterial adenylate cyclase-based two-hybrid (BACTH)

system to detect protein-protein interactions. This *in vivo* assay is based on the reconstitution of

adenylate cyclase activity when two functional fragments of Cya from *B. pertussis* are brought into

close proximity by interacting proteins ([42](#)). Interactions can be qualitatively determined on Xgal

plates and quantified by beta-galactosidase activity. Using the BACTH system, we observed homo-

oligomerization of each ClpP paralog, but failed to detect interactions between ClpP1 and ClpP2

(Fig. 4A&B). Importantly, the active site mutants also interacted with each other, indicating that this

mutation did not interfere with homo-oligomerization. These data are consistent with the predicted

structural and sequence differences noted (Fig. 2B&C).

As the BACTH system cannot indicate oligomerization state, we next purified recombinant

chlamydial ClpP1 and ClpP2 (22 and 23 kDa, respectively), including the active site mutants, and

analyzed their migration by native PAGE. As a positive control, we also purified *E. coli* ClpP (24

kDa). Each of the recombinant chlamydial proteins clearly formed a heptamer with a fainter, slower

migrating band indicating potential tetradecamer forms (Fig. 4C). The *E. coli* ClpP, in contrast,

migrated primarily as a tetradecamer with a less intense band at the predicted size of a heptamer. In

purifying Ctr ClpP1, we noticed the presence of a doublet band (Suppl. Fig. S2). However, the N-

terminus encodes two additional methionine residues at positions 6 and 7, thus the smaller product

may have resulted from an alternative start site when expressed in *E. coli*. Mass spectrometry

analysis of the N-terminus of each band supported this conclusion (Suppl. Fig. S2). To ensure that

only the larger, full-length product was produced, we mutated the methionine residues to leucine and

isoleucine. This recombinant ClpP1 (M6L/M7I) also migrated predominantly as a heptamer in the

Native PAGE assay (Fig. 4C). Therefore, we conclude that the smaller product did not alter the

oligomerization state of the protein.

Both our bioinformatics predictions and the BACTH data indicated that ClpP1 and ClpP2 do

not interact. To further test this, we mixed each recombinant protein in equimolar amounts and

analyzed their oligomerization state by Native PAGE. In support of our early observations, we did

not observe heteromers of each ClpP paralog as each protein ran as distinct heptamers with no

intermediate bands indicative of mixing of the monomers (Fig. 4D). However, we can neither

exclude that the *in vitro* conditions preclude an interaction between these components nor that these

paralogs interact *in vivo*. Nevertheless, given the differences in predicted structures at the termini of

each ClpP paralog (Fig. 2) and the different interaction data presented here, the parsimonious

interpretation is a model wherein each protease complex functions separately.

Chlamydial ClpP1 and ClpP2 vary in protease activity *in vitro*. To determine the functionality of

the protease complexes, we used the recombinant chlamydial ClpP proteins to assess their ability to

degrade the small, fluorogenic compound Suc-LY-AMC. The degradation of this compound by other

bacterial ClpP enzymes does not rely on ClpP activation by AAA+ chaperone complexes (43), thus

we could probe the basal protease activity of the ClpP paralogs. While ClpP2 degraded the substrate

under our given assay conditions, ClpP1 activity could not be detected (Fig. 5A). This result for

ClpP1 was not due to the presence of the alternative start site product since the M6L/M7I mutant

displayed no activity in this assay. We also observed enhanced protease activity for ClpP2 in a

sodium citrate buffer, as noted by others (44), but this buffer did not stimulate ClpP1 activity. That

ClpP1 lacks the ability to degrade Suc-LY-AMC suggests the N-terminal region of the complex (Fig.

2) may block entry of even this small reporter substrate into the active site. Importantly, the activity

of the ClpP2 protease was abolished by the active site serine mutation S98A (Fig. 5B). As a positive

control, we also tested the activity of the *E. coli* ClpP complex in our assay conditions and observed

degradation of the Suc-LY-AMP compound that was increased by sodium citrate (Fig. 5C). Thus, in

spite of the activity of chlamydial ClpP2, it displayed a significantly lower proteolytic capacity *in*

vitro compared to the *E. coli* ClpP. The apparent preference of heptameric formation by both

chlamydial ClpP1 and ClpP2 could explain the reduced activity as compared to the *E. coli* ClpP,

which preferentially formed tetradecamers *in vitro* (see Fig. 4C). These observed differences suggest

either that the assay conditions, which work for other bacterial ClpP homologs, are not optimal for

chlamydial ClpP proteases or that *Chlamydiae* tightly regulate the activity of their ClpP complexes.

ClpP-activating antibiotics stimulate protease activity of ClpP2, but not ClpP1, and block

chlamydial growth. Recently, a novel class of antibiotics that target ClpP have been developed (45).

ACP and ADEP derivatives activate ClpP by eliminating or reducing its need for chaperone binding.

The unregulated proteolysis that results has been demonstrated to eliminate persister bacteria (46).

We used these compounds to determine whether the activated chlamydial ClpP proteases could

degrade a more complex substrate, casein. As noted with the Suc-LY-AMP substrate, ClpP1 failed to

degrade casein in the presence of any of the ACP1 derivatives (Fig. 6A). By contrast, the ACP1

compounds stimulated ClpP2 to degrade casein (Fig. 6B), albeit with some detectable substrate

remaining at the end of the assay. As before, the *E. coli* ClpP was more efficient in degrading the

substrate with little to no casein detectable at the end of the assay (Fig. 6C; (47)). ClpP2 was thus

able, in the presence of these drugs that relax the N-termini of the complex, to degrade larger

substrates with intrinsically disordered regions (i.e. casein). The inability to detect proteolytic

activity of ClpP1 again suggests that additional factors may regulate its activity consistent with the

differences we observed in our bioinformatics analysis (Fig. 2). Also consistent with the difference

in activation is the lack of an aromatic amino acid at residue 57 of ClpP1. This particular residue of

the hydrophobic pocket has recently been shown to be instrumental in ADEP-induced dysregulation

of proteolytic activity, which supports a model where chlamydial ClpP1 is unaffected by activating

drugs (35). However, we cannot exclude that we have not identified the optimal conditions necessary

to demonstrate the enzyme activity of wild-type ClpP1.

As we were able to demonstrate the ability of the ACP1 compounds to activate ClpP2, we

next sought to test the effect of these compounds directly on *Chlamydia* growth and viability. To

ensure that any effect on the bacteria was not due to host cell death, we first tested cell viability of

infected, ACP-treated HEp2 cells. Indeed, we noted limited loss (~10%) in host cell viability for

both 25 and 50 μg/mL concentrations following 16 hours of drug treatment (Fig. 6D), which

reinforces that any loss of bacterial viability would not be due to a reduction in host cell viability.

The ACP drugs all showed a remarkable impact on *Chlamydia* at both concentrations tested (Fig.

6E). The lower drug concentration resulted in a 50-fold reduction in recoverable EBs when added at

8 hours post-infection whereas the higher concentration led to almost complete abrogation of

chlamydial viability. Given the effects of the ACP compounds on ClpP protease activity *in vitro*,

these data suggest that uncoupling of ClpP2 activity from its cognate ATPase strongly inhibits

chlamydial growth.

Discussion

The Clp protease system has been extensively characterized in *E. coli* and *Bacillus subtilis*,

and its importance in the growth and virulence of various pathogens (opportunistic or not) has been

examined (see (23) for review). Since the discovery of the ClpP protease (48), interaction studies

demonstrated that the ClpP proteins typically interact as a tetradecameric stack of two heptamers

(49). The serine active sites are buried within these barrel-like structures (50). ClpP hydrolysis of

large protein substrates is increasingly stimulated in the presence of AAA+ chaperone binding (51).

However, complexed ClpP alone processes small substrates of up to 30 amino acids regardless of the

presence of ATP, suggesting that active site availability and complex accessibility are the two main

mechanisms regulating ClpP activity (52). Substrates that enter this complex by either chaperone-

mediated or chaperone-independent means are hydrolyzed into short peptides of 6-8 amino acids

with some degree of amino acid specificity (44, 53).

While not unprecedented, the presence of two encoded *clpP* paralogs in a bacterial

chromosome is unusual. In *P. aeruginosa*, another bacterium encoding dual ClpP proteases, the two

isoforms have not been shown to interact in any test condition, and the different transcriptional

profiles and differential effects on virulence factor production indicate these paralogs likely have

distinct roles in *P. aeruginosa* (26). Conversely, the two ClpP proteins of *M. tuberculosis* and *L.*

monocytogenes heterologously interact as two homotypic heptamers, which facilitates an increase in

function (27, 29, 54). The activation of one homotypic ClpP heptamer activates the heptamer of the

other isoform in *M. tuberculosis*, providing a novel mechanism for activation (41). Of note is that *L.*

monocytogenes ClpP2 may also form a completely homotypic tetradecamer of one ClpP paralog

(55). Again, the Clp proteolytic subunits play an integral role in virulence factor regulation in

addition to enhancement of bacterial survival during intracellular growth (56). Taken together, these

studies demonstrate the diverse function of dual ClpP peptidase systems in the pathogenesis and

growth of various bacteria.

The work presented here to characterize the *C. trachomatis* ClpP paralogs is the first to

explore the function and role of the Clp protease system in chlamydiae. Our results suggest that the

two ClpP proteases likely serve distinct roles in chlamydial physiology and, we hypothesize, in the

complex developmental cycle of this organism. We utilized bioinformatics techniques as an initial

approach to investigate the potential function of the ClpP proteins. We observed significant

similarity of the chlamydial ClpP paralogs to ClpP homologs of more studied organisms such as *E.*

coli. In support of this, most of the hallmark conserved residues, including the active site triad, are

present in the chlamydial ClpP paralogs. Both ClpP1 and ClpP2 have hydrophobic residues aligned

with those of other bacterial ClpPs, suggesting the presence of AAA+ adaptor protein docking sites

(57). Indeed, chlamydial ClpC and ClpX encode the evolutionarily retained IGF/L loop motifs that

facilitate interaction with the ClpP tetradecameric complex (Wood and Ouellette, unpublished

observation) (41). In spite of the similarities between the ClpP paralogs, there are also notable

differences. The C-termini of both chlamydial ClpPs contain two residues that comprise the

hydrophobic pocket (47, 58), but the alignment shows the presence of charged residues (D15, K40,

K41, and D66) in ClpP1 while the conserved residues appear to be uncharged. In addition, ClpP1

lacks a highly conserved tyrosine residue (D15), a conserved glutamine (A36), and a conserved

aromatic residue (V57), all of which contribute to canonical activation and tetradecamer chamber

access for other homologs (35). Of note is the predicted alpha helix at the N-terminus of ClpP1 and

the C-terminus of ClpP2, which may contribute to some form of steric hindrance to prevent

interaction between ClpP1 and ClpP2 (59). Indeed, we could not detect an interaction between ClpP1

and ClpP2 using both *in vivo* and *in vitro* techniques. Rather, we detected only homoheptameric

oligomers. While our *in vitro* assays have failed to show P1/P2 complex formation, we cannot rule

out the possibility of protein modification *in vivo* promoting heteromers. Further studies examining

the role of the N- and C-termini will be conducted to determine whether or not they play a role in

complex formation and/or specificity.

Given the role of the N-terminal regions in regulation of the ClpP protease via reduced access

into the proteolytic complex (60), the hypothesis of a difference in interaction or regulation of the

ClpP complexes in *C. trachomatis* is reasonable based on the data presented here. We detected

ClpP2 protease activity against both a small oligopeptide substrate and the more complex casein

substrate. However, we did not detect proteolytic activity of ClpP1 *in vitro*, even in the presence of

well-characterized drugs that activate ClpP complexes in the absence of an ATPase, which is in

agreement with an important and highly regulated role for ClpP1 in *C. trachomatis*. The structural

conservation and presence of the catalytic active site makes it unlikely that ClpP1 is a catalytically

inactive protease. Rather, some form of modification or conformational change in structure, possibly

chaperone or adaptor mediated, may be necessary to promote full complex formation and activity

both *in vitro* and *in vivo*. In this model, an unknown regulatory factor may modulate ClpP1 activity.

In a manner consistent with our bioinformatics data, ClpP1 may be activated non-canonically, as the

mechanism needed to open the channel at the V57 residue is likely different than those targeting the

aromatic amino acids present in other homologs. We are investigating this and other possibilities.

Our data lead us to hypothesize that each ClpP protein in *Chlamydia* forms a separate

protease complex and, therefore, serves a different function. Whether each ClpP interacts

specifically with ClpX or ClpC remains to be determined. However, the genomic proximity of ClpP2

to ClpX suggests a likely interaction with ClpX by sequestration immediately following translation;

thus, we speculate that ClpP1 likely interacts preferentially with ClpC. Our qPCR data suggest that

the *clpX*-encoding gene follows the *clpP2*-encoding gene in an operon, further supporting coupled

function and an *in vivo* complex of ClpP2 and ClpX oligomers. Complex formation is currently

under investigation as is what role each complex has *in vivo*. It is also possible that ClpP

tetradecamers may function independently *in vivo* in a housekeeping role to degrade small

oligopeptides resulting from degradation of larger proteins and/or import from the host cell.

Chlamydiae encode a large repertoire of oligopeptide transporters, and the short peptide substrates

are an important source of nutrients, particularly early during infection ([61](#)).

One of the unexpected outcomes of these initial studies was the failure to transform *C.*

trachomatis with wild-type ClpP constructs. Tet-inducible systems have reduced leakiness compared

to other inducible expression systems ([62](#)) ([63](#)), and this leakiness has been observed in constructs

used to transform *C. trachomatis* ([37](#)). When used for the expression of essential gene products or in

situations where gene dosage effects are important, the leaky expression can lead to plasmid loss

and/or the inability to transform *Chlamydia*. Based on repeated failure to transform *C. trachomatis*

with vectors encoding the wild-type ClpP proteins, we conclude that leaky expression led to

rejection of the plasmid (see also ([37](#))). This suggests that stoichiometric ratios of functional Clp

enzymes are important to bacterial homeostasis. In spite of the failure to transform *C. trachomatis*

with wild-type ClpP expressing vectors, we were successful in transforming *C. trachomatis* with

vectors encoding catalytically inactive ClpP mutants. Interestingly, over-expression of the inactive

ClpP mutants in *Chlamydia* resulted in differential effects on chlamydial growth and morphology.

ClpP2 S98A has, even upon the highest level of induction, little negative effect on inclusion

morphology. The disconnect between the transformation of wild-type and inactive mutants could

mean that the ratio of wild-type ClpP2 to its cognate AAA+ chaperone is vital, perhaps to prevent

unregulated proteolysis, but incorporation of inactive forms of ClpP2 (activity loss confirmed *in*

vitro, see Fig. 5B) into the tetradecameric oligomer is not necessarily harmful to *C. trachomatis*. The

ability of ClpP2 to degrade a small peptide reporter (Fig. 5A) is in congruence with a model where a

chaperone-independent ClpP2 tetradecamer targets and degrades imported peptides. Given that

overexpression of inactive ClpP2 seems to have limited negative impact on chlamydial development,

an inactive ClpP2 oligomer may bind small peptides, preventing extensive targeting of other, more

vital substrates and negating any dominant negative effect. Conversely, over-expression of ClpP1

S92A and its incorporation into the endogenous ClpP1 proteolytic machinery has a clear negative

impact on *Chlamydia*. We observed that the mutant ClpP proteins interacted with the wild-type

proteins in the BACTH assay (Wood and Ouellette, data not shown), suggesting the potential of the

inactive proteins to form complexes with the endogenous proteins. Overall, these data suggest that,

should each mutant protein oligomerize with endogenous wild-type protein *in vivo*, subunit

poisoning of the ClpP1 complex is deleterious to chlamydiae whereas the ClpP2 complex is more

tolerant.

We hypothesize that proteolytic Clp subunit regulation by a chaperone protein is vital to

chlamydial survival. Based on our studies with the ACP1 antibiotics, our assays showed that ClpP2

acts in a manner similar to other well-characterized ClpP proteases when artificially activated by

these compounds (Fig. 6A&C). Consequently, treatment of *Chlamydia*-infected cells with ACP1 (or

one of its derivatives) significantly and drastically reduced chlamydial growth (Fig. 6E), which we

speculate is due to the effects on ClpP2 function. Whether the negative effect stems from

dysregulation of ClpP2 leading to uncontrolled proteolysis, which is supported by the *in vitro*

degradation of casein, or from blocking of ClpP2/AAA+ chaperone complex formation, resulting in

the inability to degrade larger substrates not accessible to ClpP2 alone, is currently under

investigation. In spite of the inability of ClpP2 S98A over-expression to significantly impact

chlamydial growth, the antibiotics data suggest that disruption of ClpP2 via dysregulation of activity

is overwhelmingly negative to chlamydial physiology. However, we cannot exclude a possible effect

of these antibiotics on ClpP1 *in vivo*. Clearly, over-expression of ClpP1 S92A was deleterious to

chlamydial growth, thus we conclude that each ClpP paralog is essential to normal growth and

development.

We have shown here an initial characterization of the chlamydial Clp protease system

focusing on the ClpP protease components. Overall, the sensitivity of the organisms to perturbations

in ClpP activity suggests a critical function for these paralogs in maintenance of chlamydial

physiology. We speculate that these systems will also play an integral role in differentiation and

possibly persistence (64) (65), which may provide a mechanism that could be leveraged to block

growth or eliminate persister cells (46). Further characterization of the chlamydial Clp system could

facilitate development of targeted therapeutics for treatment of *C. trachomatis* infection, thereby

lowering dependence on broad-spectrum antibiotics. Ongoing efforts are investigating this

hypothesis as well as the function of the AAA+ chaperones.

Materials and Methods

Strains and Cell Culture: The human epithelial cell line HEp2 was utilized in the transcriptional and

antibiotic studies and was routinely cultivated and passaged in Iscove's Modified Dulbecco's

Medium (IMDM, Gibco/ThermoFisher; Waltham, MA), and 10% FBS (Sigma; St. Louis, MO).

McCoy mouse fibroblasts were used for the purpose of chlamydial transformation, and human

epithelial HeLa cells were used for plaque purification of the resulting chlamydial transformants as

well as for protein isolation and assessment of ClpP2 expression. All of these cell lines were

passaged routinely in Dulbecco's Modified Eagle's Medium (DMEM, Gibco/ThermoFisher).

Density gradient purified *Chlamydia trachomatis* L2/434/Bu (ATCC VR902B) EBs were used for

the antibiotic studies. *C. trachomatis* serovar L2 EBs (25667R) naturally lacking the endogenous

plasmid were prepared and used for transformation [see (66)].

Transcript Analysis Using RT-qPCR:

HEp2 cells seeded in 6-well plates were infected at a multiplicity of infection of 1 with *C.*

trachomatis serovar L2. At the indicated times post-infection, total RNA and DNA were collected

from duplicate wells (65). Briefly, total RNA was collected from infected cells using Trizol reagent,

extracted with chloroform, and the aqueous phase precipitated with an equal volume of isopropanol

according to the manufacturer's instructions (ThermoFisher). DNA was removed from total RNA by

rigorous DNase-free treatment (ThermoFisher) before 1 µg was reverse transcribed with Superscript

III RT (ThermoFisher). Equal volumes of cDNA were used for qPCR. Total DNA was collected

from infected cells by trypsinizing the cells, pelleting for 5 min at 400 xg, and resuspending in PBS.

464 Samples were freeze-thawed three times before processing with the DNeasy Blood and Tissue kit

465 according to the manufacturer's instructions (Qiagen). 150 ng of total DNA were used for qPCR.

466 Transcripts and genomic DNA were quantified by qPCR in 25 μL reactions using 2x SYBR Green

467 Master Mix in an ABI7300 thermal cycler in comparison to a standard curve generated from purified

468 *C. trachomatis* L2 genomic DNA. Transcripts were normalized to genomic DNA.

470 **C. trachomatis *propagation and detection of ClpP2:*** DFCT28, a GFP-expressing *C. trachomatis*

471 434/Bu clone ([67](#)), was routinely grown in and titered (using the IFU assay) on HeLa cells ([68](#)).

472 Briefly, cells were maintained in Dulbecco's modified Eagle medium (DMEM) supplemented with

473 10% fetal bovine serum (FBS) and grown at 37 °C with 5% CO_2. For chlamydial infection

474 experiments, HeLa cells were grown until confluent in 6-well tissue culture dishes and then infected

475 with DFCT28 at an MOI of ~3 using centrifugation at 545 xg for one hour. The infected cells were

476 then incubated at 37 °C with 5% CO_2 with DMEM/FBS supplemented with 0.2 μg/mL

477 cycloheximide and 1x non-essential amino acids. At various times post infection, the medium was

478 removed, cells were washed twice with 2 ml of PBS, and cells were lysed via addition of 200 μL

479 Laemmli buffer with β-mercaptoethanol followed by heating at 90-100 °C for 5 minutes. Chlamydial

480 protein samples or purified, recombinant ClpP samples were run on 12% SDS-PAGE gels and either

481 stained for total protein with Coomassie Brilliant Blue or transferred to nitrocellulose for western

482 blotting. Blots were probed with a rabbit polyclonal anti-Pa_ClpP1 (*Pseudomonas aeruginosa*

483 designation, similar to ClpP2 from *C. trachomatis*) diluted 1:10000 in 5% milk Tris-buffered saline

484 (mTBS) or anti-Pa_Clp2 (*Pseudomonas aeruginosa* designation, similar to ClpP1 from *C.*

485 *trachomatis*) at 1:2500. The antibodies were kindly provided by Dr. T. Baker (Massachusetts

486 Institute of Technology) ([26](#)). After incubating with primary antibodies, blots were washed with

487 Tween (0.5%)-TBS (TTBS) and then probed with a goat, anti-rabbit IgG poly-HRP conjugated

secondary antibody (Thermo Scientific 32260) diluted 1:1000 in mTBS. As a control for chlamydial

protein, blots were also probed for the major outer membrane protein (MOMP) using a mouse

monoclonal anti-MOMP antibody (1:1000; Abcam, ab41193) and a goat anti-mouse IgG HRP

conjugated secondary antibody (1:2000; Millipore AP124P). After incubation with the secondary

antibodies, blots were washed with TTBS followed by TBS and then incubated with

chemiluminescent substrate (EMD Millipore Immobilon ECL) for imaging on a Bio-Rad Chemidoc

MP.

Bioinformatics Analysis: Gene maps and sequences of genes of *Chlamydia trachomatis* used were

obtained from STDGen database (http://stdgen.northwestern.edu). RefSeq protein sequences from *E.*

coli, B. subtilis, M. tuberculosis, and *P. aeruginosa* were acquired from the NCBI protein database

(https://www.ncbi.nlm.nih.gov/guide/proteins/). The ClpP1 vs. ClpP2 protein alignment to find

sequence   identity   was   performed   using   NCBI   Protein   BLAST   function

(https://blast.ncbi.nlm.nih.gov/Blast.cgi) (69). Multiple sequence alignments were performed using

Clustal Omega (70) with default settings and were presented using Jalview Version 2 (71). PDB files

for   predicted   3D   structures   were   acquired   from   the   Phyre2   website

(http://www.sbg.bio.ic.ac.uk/phyre2/html/page.cgi?id=index) (72). Protein models and model

alignments were rendered using the UCSF Chimera package from the Computer Graphics

Laboratory, University of California, San Francisco (supported by NIH P41 RR-01081) (73).

Plasmid Construction: A full list of the primers and plasmids used is included in the supplementary

material (S1). Plasmids for the Bacterial Adenylate Cyclase Two-Hybrid (BACTH) system were

cloned using the Gateway® recombination system (74). The genes were amplified from *Chlamydia*

trachomatis L2 genomic DNA with primers designed to add an *attB* recombination site on either side

of the gene. The PCR products were then incubated with a pDONR^{TM}221 entry vector (containing

attP recombination sites) in the presence of BP Clonase II (Invitrogen) that inserts the gene via the

flanking *attB* sites and removes *ccdB* endotoxin flanked by the *attP* sites encoded on the plasmid,

allowing for positive selection. The result of the BP reaction was an entry vector containing the gene

of interest flanked by *attL* sites. 2 μL were transformed into DH5α chemically competent *E. coli* and

plated onto an LB agar plate containing 50 μg/mL kanamycin. Plasmid from an individual colony

was purified and used for the LR reaction into one of three destination vectors (pST25-DEST,

pSNT25-DEST, or pUT18C-DEST). The same entry vector for any given gene was used for all three

LR reactions to insert into the destination vector with LR Clonase II. 150 ng of the entry vector was

incubated with 150ng of destination vector for 1 hour at room temperature. 2 μL were used to

transform XL1 *E. coli*, which were plated on the appropriate selection plate. Purified plasmid from

an individual colony was sequence verified prior to use the in BACTH assay (see below).

524   Constructs for chlamydial transformation were created using the HiFi Cloning (New England

Biolabs) protocol. To add the C-terminal 6xHis tag to the Clp proteins, the genes were amplified

from the genome with a primer to add the poly-histidine tag. These products served as the template

for PCR reactions to add the necessary overlap for the HiFi reaction. Primers were generated using

the NEBuilder® assembly tool available from New England BioLabs (http://nebuilder.neb.com). The

backbone used was the pTLR2 derivative of the pASK plasmid (36). The pTLR2 backbone was

digested using FastDigest BshTI and Eco52I restriction enzymes and dephosphorylated using

FastAP (ThermoFisher), and then 50ng of digested plasmid was incubated with a 2:1 ratio of insert

copy number to backbone and 2x HiFi master mix (NEB). Following a 15 minute incubation of the

reaction mix at 50° C, 2 μL of the reaction was transformed into DH5α chemically competent *E. coli*

(NEB) and plated on the appropriate antibiotic selection plate. Positive clone sequences were

verified by Eurofins Genomics. Sequence verified plasmids were transformed into *dam-/dcm- E. coli*

(New England BioLabs) in order to produce demethylated plasmid, which was sequence and digest

verified prior to transformation into *C. trachomatis* (see below).

Strains created or used in this study are listed in the supplementary material. *E. coli* strains

were maintained on LB agar plates and grown in LB medium or on LB agar plates supplemented

with 100 µg/ml ampicillin as needed. Chlamydial genomic DNA for cloning was obtained from EBs

using a phenol:chloroform extraction after extensive heat treatment in the presence of proteinase K

(75), and *E. coli* genomic DNA was isolated using sodium hydroxide lysis of colonies. The

chlamydial *clpP1* and *clpP2* along with *clpP* from *E. coli* were amplified via PCR using the primers

listed in Supplemental Information and Phusion DNA polymerase (Thermo Scientific). PCR

products were cloned into the pLATE31 expression vector from Thermo Scientific as directed by the

manufacturer to create fusion proteins with a C-terminal 6x His-tag. Plasmids were initially

transformed into *E. coli* NEB10 and selected on LB agar ampicillin plates. Transformants were

screened for inserts using colony PCR with Fermentas Master Mix (Thermo Scientific) and positive

clones were grown for plasmid isolation (GeneJet Plasmid Miniprep Kit, Thermo Scientific). DNA

inserts were sequenced by Macrogen USA and sequence-verified plasmids were then transformed

into *E. coli* BL21(DE3) bacteria for protein production.

Chlamydial Transformation: The protocol followed was a modification of the method developed by

Mueller and Fields (76). For transformation, 10^6 *C. trachomatis* serovar L2 EBs (25667R) naturally

lacking the endogenous plasmid were incubated with 2 µg of unmethylated plasmid in a volume of

50 µL $CaCl_2$ at room temperature for 30 minutes. Each reaction was sufficient for a confluent

monolayer of McCoy cells in one well of a six well plate that had been plated a day prior. The

transformants were added to a 1 mL overlay of room temperature HBSS per well, and an additional 1

mL of HBSS was then added to each well. The plate was centrifuged at 400 xg for 15 min at room

temperature, where the beginning of this step was recorded as the time of infection. Following the

spin, the plate was incubated for 15 minutes at 37° C. This infection was recorded as T_0. The

inoculum was aspirated at the end of the incubation and replaced with antibiotic-free DMEM+10%

FBS. 8 hours post-infection, the media was replaced with DMEM containing 1 μg/mL

cycloheximide, 10 μg/mL gentamicin, and 1 U/mL penicillin. Cells infected with transformants were

passaged every 48 hours until a population of penicillin resistant bacteria was established. These EBs

were then harvested and frozen in sucrose/phosphate (2SP; (66)) solution at -80° C prior to titration.

Determining the Effect of Overexpression of Clp Mutant Proteins via Immunofluorescence

Analysis: Transformed *C. trachomatis* containing a mutant *clp* gene under control of an

anhydrotetracycline (aTc) inducible promoter was used to infect a monolayer of HEp2 cells on

coverslips with penicillin as a selection agent. Samples were induced with increasing, subtoxic

amounts of aTc at 8 hours post-infection and were methanol fixed after either a 6 hour or 14 hour

pulse (16 or 24 hours post-infection). Fixed cells were incubated with an anti-Ctr serovar L2 guinea

pig primary antibody (kindly provided by Dr. Rucks, UNMC) to stain for the organism and a goat

anti-guinea pig Alexa488 conjugated secondary antibody for visualization of organisms within the

inclusion. Additionally, a mouse anti-6xHis tag was used, followed by a goat anti-mouse Alexa594

secondary antibody for confirmation of protein expression and localization. Finally, the samples

were stained with DAPI to visualize the host cell and bacterial DNA. Representative images were

taken on a Zeiss Fluoview1000 laser scanning confocal microscope with a 60x objective and 2x

digital zoom and were equally color corrected using Adobe Photoshop CC.

Determining Protein-Protein Interactions with the BACTH System: To test interaction of the Clp

proteins, the Bacterial Adenylate Cyclase Two-Hybrid (BACTH) assay was utilized. This assay

relies on reconstitution of adenylate cyclase activity in adenylate cyclase deficient (Δ*cya*) DHT1 *E.*

coli. The genes of interest are translationally fused to one of either subunit, denoted as T18 and T25,

of the *B. pertussis* adenylate cyclase toxin. Each *clpP* gene cloned into one of the pST25, pSNT25,

or pUT18C Gateway® vectors was tested for both homotypic and heterotypic interactions ([10]). One

plasmid from the T25 background and one from the T18 background were co-transformed into the

DHT1 CaCl$_2$ competent *E. coli* and were plated on a double antibiotic minimal M63 medium

selection plate supplemented with 0.5 mM IPTG for induction of the protein, 40 μg/mL Xgal to give

a visual readout upon cleavage, and 0.2% maltose as a unique carbon source. These plates also

contain 0.04% casein hydrolysate to supplement the bacteria with the branched chain amino acids as

Δ*cya* DHT1 *E. coli* cannot synthesize these amino acids. Blue colonies were indicative of positive

interaction between proteins since both the *lac* and *mal* operons require reconstituted cAMP

production from interacting T25 and T18 fragments to be expressed. Leucine zipper motifs were

used for controls in pKT25 and pUT18C backgrounds on the appropriate antibiotic selection plates

because these have been previously shown to interact ([77]). β-galactosidase activity was then

measured from 8 colonies to quantify protein interactions. Random positive colonies were used to

inoculate individual wells and grown 24 hours at 30° in M63 with 0.2% maltose and appropriate

antibiotics. These bacteria were permeabilized with 0.1% SDS and chloroform prior to addition of

0.1% o-nitrophenol-β-galactoside (ONPG). The reaction was stopped using 1 M NaHCO$_3$ after

precisely 20 minutes of incubation at room temperature. Absorbance at the 405 wavelength was

recorded and normalized to bacterial growth (OD$_{600}$), dilution factor, and time (in minutes) of

incubation prior to stopping the reaction. Totals were reported in relative units (RU) of β-

galactosidase activity.

Purification of Recombinant ClpP1 and ClpP2: His-tagged Ctr ClpP1, Ctr ClpP2, and Ec ClpP

were purified from 500 mL cultures of BL21(DE3) *E. coli* transformed with the respective plasmid

based on the protocol described in (78). Samples were induced with 0.5 mM IPTG and incubated

with shaking for 20 hours at 18°C. Cultures were pelleted and frozen at -20°C prior to purifications.

Buffers used are listed in Table S2. Samples were suspended, sonicated, bound to HisPur Cobalt

Resin (Thermo Scientific), and washed in buffer A. Proteins were eluted from the resin using buffer

B. Buffer exchange for buffer C was performed using a Millipore Amicon Ultra 15 filtration units (3

kDa cut-off). ClpP proteins were quantified using the Bio-Rad Protein assay, assessed for purity on

12% SDS-PAGE gels with Coomassie staining, and identified using anti-His-tag western blot.

Blotting was performed using a mouse monoclonal anti-6x His antibody (1:1000; Millipore HIS.H8)

and a goat anti-mouse IgG HRP conjugated secondary antibody (1:2000). Protein samples were

aliquoted and stored at -80°C.

In Vitro *Analysis of ClpP1 and ClpP2 Homo-Oligomerization:* 5 µg of purified protein was

incubated in buffer D at 37°C for 1 hour before being mixed with a 5x native sample buffer (5 mM

Tris [pH 6.8], 38 mM glycine, 0.06% bromophenol blue) and analyzed on a BioRad MiniProtean 4-

20% gradient gel for Native-PAGE. Samples were run for 90 minutes at 200V. Gels were assessed

using Coomassie staining.

Assessment of ClpP activity **in vitro.** Fluorometric peptide assay: The ClpPs (at 1 µM) were added

to 500 µM of Suc-Luc-Tyr-AMC (Boston Biochem) dissolved in buffer E (50 mM Tris-HCl [pH 8],

200 mM KCl, and 1 mM DTT) or buffer F (with 0.2 M sodium citrate) (44). Final reaction volumes

were 50 µl. Reactions were monitored over six hours at 37 °C using a BioTek Synergy HT plate

reader set at an excitation of 340/360 and an emission of 440/460 with readings taken at five-minute

intervals. Casein degradation assays: Casein (Sigma-Aldrich) was dissolved in buffer E and 1 µg was

used per assay. Samples containing casein and 1 µM of the respective ClpP were incubated at 37 °C

for 3 hours with or without the respective ACP compound (500 µM). Reactions were halted by

mixing with 2x Laemmli buffer containing β-mercaptoethanol and heating at 90-100 °C for 5

minutes. Samples were analyzed for digestion of casein using 12% SDS-PAGE gels followed by

Coomassie staining.

Effect of ACP compounds on chlamydial growth and host cell viability. Antibiotic stocks of ACP1,

ACP1a, and ACP1b were synthesized (47), resuspended at 25 mg/mL in DMSO, and frozen at -

20°C. Methods for the synthesis, purification, and analysis of these compounds is available in

Supplementary Information. For assessment of cell viability upon treatment, four wells of a 96 well

plate with a confluent monolayer of HEp2 cells were either infected with density gradient purified

wild type *C. trachomatis* serovar L2 with an MOI of 1 or left uninfected. These wells were either

treated or not with 25 or 50 µg/mL of ACP1, ACP1a, or ACP1b, with a set of DMSO only samples

used as a control. Antibiotics were added eight hours post-infection (hpi). At 24 hpi, 100 µL of 2x

Resazurin (Abcam) was added to three wells of each treatment condition, adding only DMEM to the

fourth as a background control. Following a four-hour incubation at 37º C, absorbance at the 570 nm

wavelength was recorded using a Tecan plate reader. The wells were averaged, subtracting

background absorbance from samples without dye. Absorbance was reported as percentage of the

untreated samples. To quantify the effect of the drug on *C. trachomatis*, variable treatments of each

drug (25 µg and 50 µg) were added or not eight hours post-infection. Cell lysates containing *C.*

trachomatis were collected in 2SP and frozen at -80° C prior to serial titration on a fresh cell layer.

Inclusion forming units (IFUs), a proxy for recoverable EBs from the initial infection samples, were

calculated from the average number of inclusions in 15 fields of view multiplied by the number of

fields of view within the well and corrected for the dilution factor and volume of inoculum.

Acknowledgements

We would like to thank Dr. Elizabeth Rucks of UNMC for providing us with multiple

antibodies for analysis by immunofluorescence, and for thoughtful review of the manuscript. We

thank Dr. Walid Houry (Univ. of Toronto) for providing us with an initial sample of the ACP1b

antibiotic, Dr. Tania Baker (MIT) for providing us with Pa_ClpP specific antibodies for use in our

experiments, and Dr. H. Caldwell (NIH/NIAID) for eukaryotic cell lines. We also thank Amanda

Blocker (SIUC) for providing assistance with protein purification and activity assays. Funding for

this project was supported by an NSF CAREER award (1810599) to SPO and University funds to

DJF. This project was also funded by the Department of Defense office of the Congressionally

Directed Medical Research Programs (CDMRP), 2017 Peer Reviewed Medical Research Program

(PRMRP) Discovery Award (PR172445) to MCS.

References:

1. WHO. 2017. Trachoma. http://www.who.int/mediacentre/factsheets/fs382/en/. Accessed

2. Hafner LM, Wilson DP, Timms P. 2014. Development status and future prospects for a

vaccine against *Chlamydia trachomatis* infection. Vaccine 32:1563-1571.

3. CDC. 2017. 2017 Sexually Transmitted Diseases Surveillance.

https://www.cdc.gov/std/stats16/chlamydia.htm#foot-1. Accessed

4. AbdelRahman YM, Belland RJ. 2006. The chlamydial developmental cycle. FEMS

Microbiology Reviews 29:949-959.

5. Hackstadt T, Baehr W, Ying Y. 1991. *Chlamydia trachomatis* developmentally regulated

protein is homologous to eukaryotic histone H1. Proceedings of the National Academy of

Sciences of the United States of America 88:3937-3941.

6. Hatch TP, Allan I, Pearce JH. 1984. Structural and polypeptide differences between

envelopes of infective and reproductive life cycle forms of *Chlamydia* spp. Journal of

Bacteriology 157:13-20.

7. Clifton DR, Fields KA, Grieshaber SS, Dooley CA, Fischer ER, Mead DJ, Carabeo RA,

Hackstadt T. 2004. A chlamydial type III translocated protein is tyrosine-phosphorylated at

the site of entry and associated with recruitment of actin. Proceedings of the National

Academy of Sciences of the United States of America 101:10166.

8. Lane BJ, Mutchler C, Al Khodor S, Grieshaber SS, Carabeo RA. 2008. Chlamydial Entry

Involves TARP Binding of Guanine Nucleotide Exchange Factors. PLoS Pathogens

4:e1000014.

9. Hodinka RL, Davis CH, Choong J, Wyrick PB. 1988. Ultrastructural study of endocytosis of

Chlamydia trachomatis by McCoy cells. Infection and Immunity 56:1456-1463.

10.   Moore ER, Ouellette SP. 2014. Reconceptualizing the chlamydial inclusion as a pathogen-

specified parasitic organelle: an expanded role for Inc proteins. Frontiers in Cellular and

Infection Microbiology 4:157.

11.   Belland RJ, Zhong G, Crane DD, Hogan D, Sturdevant D, Sharma J, Beatty WL, Caldwell

HD. 2003. Genomic transcriptional profiling of the developmental cycle of *Chlamydia*

trachomatis. Proceedings of the National Academy of Sciences of the United States of

America 100:8478-8483.

12.   Vandahl BBS, Birkelund S, Christiansen G. 2004. Genome and proteome analysis of

Chlamydia. PROTEOMICS 4:2831-2842.

13.   Skipp PJS, Hughes C, McKenna T, Edwards R, Langridge J, Thomson NR, Clarke IN. 2016.

Quantitative Proteomics of the Infectious and Replicative Forms of Chlamydia trachomatis.

PLoS ONE 11:e0149011.

14.   Saka HA, Thompson JW, Chen Y-S, Kumar Y, Dubois LG, Moseley MA, Valdivia RH.

2011. Quantitative proteomics reveals metabolic and pathogenic properties of Chlamydia

trachomatis developmental forms. Molecular Microbiology 82:1185-1203.

15.   Beatty WL, Byrne GI, Morrison RP. 1993. Morphologic and antigenic characterization of

interferon gamma-mediated persistent Chlamydia trachomatis infection in vitro. Proceedings

of the National Academy of Sciences of the United States of America 90:3998-4002.

16.   Raulston JE. 1997. Response of Chlamydia trachomatis serovar E to iron restriction in vitro

and evidence for iron-regulated chlamydial proteins. Infection and Immunity 65:4539-4547.

17.   Matsumoto A, Manire GP. 1970. Electron Microscopic Observations on the Effects of

Penicillin on the Morphology of Chlamydia psittaci. Journal of Bacteriology 101:278-285.

18.   Stephens RS, Kalman S, Lammel C, Fan J, Marathe R, Aravind L, Mitchell W, Olinger L,

715   Tatusov RL, Zhao Q, Koonin EV, Davis RW. 1998. Genome Sequence of an Obligate

716   Intracellular Pathogen of Humans: *Chlamydia trachomatis*. Science 282:754.

717   19.   Stinson BM, Nager AR, Glynn SE, Schmitz KR, Baker TA, Sauer RT. 2013. Nucleotide

718   binding and conformational switching in the hexameric ring of a AAA+ machine. Cell

719   153:628-639.

720   20.   Kirstein J, Schlothauer T, Dougan DA, Lilie H, Tischendorf G, Mogk A, Bukau B, Turgay K.

721   2006. Adaptor protein controlled oligomerization activates the AAA+ protein ClpC. The

722   EMBO Journal 25:1481-1491.

723   21.   Baker TA, Sauer RT. 2012. ClpXP, an ATP-powered unfolding and protein-degradation

724   machine. Biochimica et biophysica acta 1823:15-28.

725   22.   Alexopoulos JA, Guarné A, Ortega J. 2012. ClpP: A structurally dynamic protease regulated

726   by AAA+ proteins. Journal of Structural Biology 179:202-210.

727   23.   Yu AYH, Houry WA. 2007. ClpP: A distinctive family of cylindrical energy-dependent

728   serine proteases. FEBS Letters 581:3749-3757.

729   24.   Grimaud R, Kessel M, Beuron F, Steven AC, Maurizi MR. 1998. Enzymatic and Structural

730   Similarities between the *Escherichia coli* ATP-dependent Proteases, ClpXP and ClpAP.

731   Journal of Biological Chemistry 273:12476-12481.

732   25.   Bewley MC, Graziano V, Griffin K, Flanagan JM. 2006. The asymmetry in the mature

733   amino-terminus of ClpP facilitates a local symmetry match in ClpAP and ClpXP complexes.

734   Journal of structural biology 153:113-128.

735   26.   Hall BM, Breidenstein EBM, de la Fuente-Núñez C, Reffuveille F, Mawla GD, Hancock

736   REW, Baker TA. 2017. Two Isoforms of Clp Peptidase in *Pseudomonas aeruginosa* Control

737   Distinct Aspects of Cellular Physiology. Journal of Bacteriology 199:e00568-16.

27.   Akopian T, Kandror O, Raju RM, UnniKrishnan M, Rubin EJ, Goldberg AL. 2012. The

active ClpP protease from *M. tuberculosis* is a complex composed of a heptameric ClpP1 and

a ClpP2 ring. The EMBO Journal 31:1529-1541.

28.   Dahmen M, Vielberg M-T, Groll M, Sieber Stephan A. 2015. Structure and Mechanism of

the Caseinolytic Protease ClpP1/2 Heterocomplex from Listeria monocytogenes.

Angewandte Chemie International Edition 54:3598-3602.

29.   Balogh D, Dahmen M, Stahl M, Poreba M, Gersch M, Drag M, Sieber SA. 2017. Insights

into ClpXP proteolysis: heterooligomerization and partial deactivation enhance chaperone

affinity and substrate turnover in *Listeria monocytogenes*. Chemical Science 8:1592-1600.

30.   Shaw EI, Dooley CA, Fischer ER, Scidmore MA, Fields KA, Hackstadt T. 2002. Three

temporal classes of gene expression during the *Chlamydia trachomatis* developmental cycle.

Molecular Microbiology 37:913-925.

31.   Wichlan DG, Hatch TP. 1993. Identification of an early-stage gene of *Chlamydia psittaci*

6BC. Journal of Bacteriology 175:2936-2942.

32.   Douglas AL, Hatch TP. 2000. Expression of the transcripts of the sigma factors and putative

sigma factor regulators of *Chlamydia trachomatis L2*. Gene 247:209-214.

33.   Ouellette SP, Rueden KJ, Gauliard E, Persons L, de Boer PA, Ladant D. 2014. Analysis of

MreB interactors in Chlamydia reveals a RodZ homolog but fails to detect an interaction with

MraY. Frontiers in Microbiology 5:279.

34.   Martin A, Baker TA, Sauer RT. 2008. Diverse pore loops of the AAA+ ClpX machine

mediate unassisted and adaptor-dependent recognition of ssrA-tagged substrates. Molecular

cell 29:441-450.

35.   Ni T, Ye F, Liu X, Zhang J, Liu H, Li J, Zhang Y, Sun Y, Wang M, Luo C, Jiang H, Lan L,

761   Gan J, Zhang A, Zhou H, Yang C-G. 2016. Characterization of Gain-of-Function Mutant

762   Provides New Insights into ClpP Structure. ACS Chemical Biology 11:1964-1972.

763   36.   Wickstrum J, Sammons LR, Restivo KN, Hefty PS. 2013. Conditional Gene Expression in

764   *Chlamydia trachomatis* Using the Tet System. PLoS ONE 8:e76743.

765   37.   Ouellette SP. 2018. Feasibility of a Conditional Knockout System for *Chlamydia* Based on

766   CRISPR Interference. Frontiers in Cellular and Infection Microbiology 8:59.

767   38.   Feng J, Michalik S, Varming AN, Andersen JH, Albrecht D, Jelsbak L, Krieger S, Ohlsen K,

768   Hecker M, Gerth U, Ingmer H, Frees D. 2013. Trapping and Proteomic Identification of

769   Cellular Substrates of the ClpP Protease in *Staphylococcus aureus*. Journal of Proteome

770   Research 12:547-558.

771   39.   Rucks EA, Olson MG, Jorgenson LM, Srinivasan RR, Ouellette SP. 2017. Development of a

772   Proximity Labeling System to Map the Chlamydia trachomatis Inclusion Membrane.

773   Frontiers in Cellular and Infection Microbiology 7:40.

774   40.   Joshi SA, Hersch GL, Baker TA, Sauer RT. 2004. Communication between ClpX and ClpP

775   during substrate processing and degradation. Nature Structural &Amp; Molecular Biology

776   11:404.

777   41.   Schmitz KR, Carney DW, Sello JK, Sauer RT. 2014. Crystal structure of *Mycobacterium*

778   *tuberculosis* ClpP1P2 suggests a model for peptidase activation by AAA+ partner binding

779   and substrate delivery. Proceedings of the National Academy of Sciences of the United States

780   of America 111:E4587-E4595.

781   42.   Karimova G, Pidoux J, Ullmann A, Ladant D. 1998. A bacterial two-hybrid system based on

782   a reconstituted signal transduction pathway. Proceedings of the National Academy of

783   Sciences of the United States of America 95:5752-5756.

43.   Woo KM, Chung WJ, Ha DB, Goldberg AL, Chung CH. 1989. Protease Ti from Escherichia

coli requires ATP hydrolysis for protein breakdown but not for hydrolysis of small peptides.

Journal of Biological Chemistry 264:2088-2091.

44.   Gersch M, Stahl M, Poreba M, Dahmen M, Dziedzic A, Drag M, Sieber SA. 2016. Barrel-

shaped ClpP Proteases Display Attenuated Cleavage Specificities. ACS Chemical Biology

11:389-399.

45.   Brötz-Oesterhelt H, Beyer D, Kroll H-P, Endermann R, Ladel C, Schroeder W, Hinzen B,

Raddatz S, Paulsen H, Henninger K, Bandow JE, Sahl H-G, Labischinski H. 2005.

Dysregulation of bacterial proteolytic machinery by a new class of antibiotics. Nature

Medicine 11:1082.

46.   Conlon BP, Nakayasu ES, Fleck LE, LaFleur MD, Isabella VM, Coleman K, Leonard SN,

Smith RD, Adkins JN, Lewis K. 2013. Killing persister cells and eradicating a biofilm

infection by activating the ClpP protease. Nature 503:365-370.

47.   Leung E, Datti A, Cossette M, Goodreid J, McCaw Shannon E, Mah M, Nakhamchik A,

Ogata K, El Bakkouri M, Cheng Y-Q, Wodak Shoshana J, Eger Bryan T, Pai Emil F, Liu J,

Gray-Owen S, Batey Robert A, Houry Walid A. 2011. Activators of Cylindrical Proteases as

Antimicrobials: Identification and Development of Small Molecule Activators of ClpP

Protease. Chemistry & Biology 18:1167-1178.

48.   Hwang BJ, Park WJ, Chung CH, Goldberg AL. 1987. *Escherichia coli* contains a soluble

ATP-dependent protease (Ti) distinct from protease La. Proceedings of the National

Academy of Sciences of the United States of America 84:5550-5554.

49.   Flanagan JM, Wall JS, Capel MS, Schneider DK, Shanklin J. 1995. Scanning Transmission

Electron Microscopy and Small-Angle Scattering Provide Evidence that Native *Escherichia*

coli ClpP is a Tetradecamer with an Axial Pore. Biochemistry 34:10910-10917.

50.   Wang J, Hartling JA, Flanagan JM. 1997. The Structure of ClpP at 2.3 Å Resolution Suggests

a Model for ATP-Dependent Proteolysis. Cell 91:447-456.

51.   Lee ME, Baker TA, Sauer RT. 2010. Control of substrate gating and translocation into ClpP

by channel residues and ClpX binding. Journal of molecular biology 399:707-718.

52.   Thompson MW, Singh SK, Maurizi MR. 1994. Processive degradation of proteins by the

ATP-dependent Clp protease from *Escherichia coli*. Requirement for the multiple array of

active sites in ClpP but not ATP hydrolysis. Journal of Biological Chemistry 269:18209-

18215.

53.   Jennings LD, Bohon J, Chance MR, Licht S. 2008. The ClpP N-terminus coordinates

substrate access with protease active site reactivity. Biochemistry 47:11031-11040.

54.   Benaroudj N, Raynal B, Miot M, Ortiz-Lombardia M. 2011. Assembly and proteolytic

processing of mycobacterial ClpP1 and ClpP2. BMC Biochemistry 12:61-61.

55.   Dahmen M, Vielberg MT, Groll M, Sieber Stephan A. 2015. Structure and Mechanism of the

Caseinolytic Protease ClpP1/2 Heterocomplex from *Listeria monocytogenes*. Angewandte

Chemie International Edition 54:3598-3602.

56.   Gaillot O, Bregenholt S, Jaubert F, Di Santo JP, Berche P. 2001. Stress-Induced ClpP Serine

Protease of *Listeria monocytogenes* Is Essential for Induction of Listeriolysin O-Dependent

Protective Immunity. Infection and Immunity 69:4938-4943.

57.   Martin A, Baker TA, Sauer RT. 2007. Distinct static and dynamic interactions control

ATPase-peptidase communication in a AAA+ protease. Molecular cell 27:41-52.

58.   Lee B-G, Park EY, Lee K-E, Jeon H, Sung KH, Paulsen H, Rübsamen-Schaeff H, Brötz-

Oesterhelt H, Song HK. 2010. Structures of ClpP in complex with acyldepsipeptide

antibiotics reveal its activation mechanism. Nature Structural &Amp; Molecular Biology

17:471.

59.   Bewley MC, Graziano V, Griffin K, Flanagan JM. 2009. Turned on for degradation: ATPase-

833   independent degradation by ClpP. Journal of structural biology 165:118-125.

834   60.   Li DHS, Chung YS, Gloyd M, Joseph E, Ghirlando R, Wright GD, Cheng Y-Q, Maurizi MR,

835   Guarné A, Ortega J. 2010. Acyldepsipeptide Antibiotics Induces the Formation of a

836   Structured Axial Channel in ClpP: a Model for the ClpX/ClpA Bound State of ClpP.

837   Chemistry & biology 17:959-969.

838   61.   Ouellette SP, Dorsey FC, Moshiach S, Cleveland JL, Carabeo RA. 2011. Chlamydia Species-

839   Dependent Differences in the Growth Requirement for Lysosomes. PLoS ONE 6:e16783.

840   62.   Lutz R, Bujard H. 1997. Independent and tight regulation of transcriptional units in

841   Escherichia coli via the LacR/O, the TetR/O and AraC/I1-I2 regulatory elements. Nucleic

842   Acids Research 25:1203-1210.

843   63.   Wang B, Barahona M, Buck M. 2015. Amplification of small molecule-inducible gene

844   expression via tuning of intracellular receptor densities. Nucleic Acids Research 43:1955-

845   1964.

846   64.   Kane CD, Vena RM, Ouellette SP, Byrne GI. 1999. Intracellular Tryptophan Pool Sizes May

847   Account for Differences in Gamma Interferon-Mediated Inhibition and Persistence of

848   Chlamydial Growth in Polarized and Nonpolarized Cells. Infection and Immunity 67:1666-

849   1671.

850   65.   Ouellette SP, Rueden KJ, Rucks EA. 2016. Tryptophan Codon-Dependent Transcription in

851   Chlamydia pneumoniae during Gamma Interferon-Mediated Tryptophan Limitation.

852   Infection and Immunity 84:2703-2713.

853   66.   Wang Y, Kahane S, Cutcliffe LT, Skilton RJ, Lambden PR, Clarke IN. 2011. Development

854   of a Transformation System for *Chlamydia trachomatis*: Restoration of Glycogen

855   Biosynthesis by Acquisition of a Plasmid Shuttle Vector. PLoS Pathogens 7:e1002258.

856   67.   Illingworth M, Hooppaw AJ, Ruan L, Fisher DJ, Chen L. 2017. Biochemical and Genetic

857   Analysis of the Chlamydia GroEL Chaperonins. Journal of Bacteriology 199:e00844-16.

858   68.   Scidmore MA. 2006. Cultivation and Laboratory Maintenance of Chlamydia trachomatis.

859   Current Protocols in Microbiology 00:11A.1.1-11A.1.25.

860   69.   Altschul SF, Wootton JC, Gertz EM, Agarwala R, Morgulis A, Schäffer AA, Yu Y-K. 2005.

861   Protein Database Searches Using Compositionally Adjusted Substitution Matrices. The FEBS

862   journal 272:5101-5109.

863   70.   Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H,

864   Remmert M, Söding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation of high-

865   quality protein multiple sequence alignments using Clustal Omega. Molecular Systems

866   Biology 7:539-539.

867   71.   Troshin PV, Procter JB, Barton GJ. 2011. Java bioinformatics analysis web services for

868   multiple sequence alignment—JABAWS:MSA. Bioinformatics 27:2001-2002.

869   72.   Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. 2015. The Phyre2 web portal

870   for protein modelling, prediction and analysis. Nature protocols 10:845-858.

871   73.   Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE.

872   2004. UCSF Chimera—A visualization system for exploratory research and analysis. Journal

873   of Computational Chemistry 25:1605-1612.

874   74.   Ouellette SP, Gauliard E, Antosová Z, Ladant D. 2014. A Gateway®-compatible bacterial

875   adenylate cyclase-based two-hybrid system. Environmental Microbiology Reports 6:259-267.

876   75.   Ouellette SP, Hatch TP, AbdelRahman YM, Rose LA, Belland RJ, Byrne GI. 2006. Global

877   transcriptional upregulation in the absence of increased translation in Chlamydia during

878   IFNγ-mediated host cell tryptophan starvation. Molecular Microbiology 62:1387-1401.

76.   Mueller KE, Wolf K, Fields KA. 2017. Chlamydia trachomatis Transformation and Allelic

Exchange Mutagenesis. Current Protocols in Microbiology 45:11A.3.1-11A.3.15.

77.   Robichon C, Karimova G, Beckwith J, Ladant D. 2011. Role of Leucine Zipper Motifs in

Association of the *Escherichia coli* Cell Division Proteins FtsL and FtsB. Journal of

Bacteriology 193:4988-4992.

78.   Trentini DB, Suskiewicz MJ, Heuck A, Kurzbauer R, Deszcz L, Mechtler K, Clausen T.

2016. Arginine phosphorylation marks proteins for degradation by a Clp protease. Nature

539:48.

 Figure Legends

 Figure 1: The *clp* genes are transcribed, and ClpP2 is produced, during RB growth. (A) Gene

 maps of the *clp* genes in the *Chlamydia trachomatis* genome. Gene numbers are in numerical order

 from left to right and reflect the serovar D numbering scheme of Stephens et al. ([18](#)). The *clp* genes

 are circled in red. The *clpP2* gene is *ct706*. **(B) RT-qPCR** analysis of the *clp* genes from two

 independent time course experiments of a *C. trachomatis* serovar L2 infection of HEp2 cells. Total

 RNA and DNA were collected at the indicated times post infection and processed as described in

 Materials and Methods. Equivalent amounts of cDNA were used for each assay and analyzed in

 triplicate. Results were reported as a ratio of cDNA to genomic DNA. Standard deviations for each

 were typically less than 10% of the sample. Note that some transcripts were not detectable at 1 hpi.

 Temporal expression of **(C)** ClpP2, using a *Pseudomonas aeruginosa* ClpP1-specific antibody, and

 (D) the chlamydial major outer membrane protein (MOMP) was assessed. Whole cell lysates of total

 protein from a time course of *Chlamydia*-infected cells **(E)** were separated by SDS-PAGE and

 transferred to a PVDF membrane for blotting.

 Figure 2: Bioinformatics analyses suggest both chlamydial ClpP paralogs are functional

 proteases. (A) Pairwise alignments performed using NCBI-BLAST (default settings) and presented

 as %Identity/%Similarity/%Coverage. Multiple sequence alignment performed using Clustal Omega

 default settings and presented using Jalview Version 2. Organisms included are *C. trachomatis* (Ctr),

 Escherichia coli (Ec), *Bacillus subtilis* (Bs), *Mycobacterium tuberculosis* (Mtb), and *Pseudomonas*

 aeruginosa (Pa). Conserved residues highlighted in varying shades of blue depending on

 conservation strength across species, with the catalytic triad in red, the oligomer interface in yellow,

 and the hydrophobic docking region in green. Residues in which ClpP1 has a different substitution

compared to other ClpP proteins are denoted by Ω. Residues involved in activation conformational

changes denoted by ψ ([35]) **(B)** Predicted 3D structures of Ctr ClpP1, Ctr ClpP2, and Ec ClpP

(peptide colors as shown in multiple sequence alignment) acquired from Phyre2 and aligned and

colored in Chimera (UCSF). Active site residues are circled in red. **(C) Space filling model** of

predicted 3D structures as in (B). "Rear" is a 180° rotation around the Y axis to show the surface of

the protein not shown in the "Front" image. The active site is again circled in red.

Figure 3: Over-expression of inactive ClpP1, but not ClpP2, inhibits chlamydial inclusion

growth. *C. trachomatis* serovar L2 was transformed with anhydrotetracycline (aTc)-inducible shuttle

vectors encoding active site mutants of each ClpP paralog with a 6xHis tag at the C-terminus. HEp2

cells were infected with each transformant, and expression of the inactive ClpP_6xHis was induced

at 10h post infection (hpi). **(A) ClpP1(S92A) overexpression** assay. Overexpression of inactive

ClpP1 has a negative impact on inclusion growth. **(B) ClpP2(S98A) overexpression** assay.

Parameters same as described above. Overexpression does not appear to negatively affect

Chlamydia. For (A) and (B), samples were stained for major outer membrane protein (CtrL2; green),

6xHis tagged ClpP protein of interest (red), and DNA (blue) with a composite image of all channels

merged (Merge). Representative images of three independent experiments are presented. Scale bars

are equal to 10 μm. Images were acquired on a Fluoview1000 laser scanning confocal microscope

with a 60x objective and a 2x digital magnification.

Figure 4: The chlamydial ClpP paralogs self-associate but do not form hetero-oligomeric

complexes. (A) Bacterial adenylate cyclase-based two-hybrid assay of homotypic interactions of

C. trachomatis (Ctr) ClpP1 and ClpP2. The indicated constructs encoding each *clpP* paralog fused to

the T18 or T25 fragment of the Cya toxin of *B. pertussis* were co-transformed into Δ*cya E. coli*

936 DHT1 and plated on minimal medium containing X-gal. Interactions between proteins results in the

937 formation of blue colonies. Representative images of results from at least three independent

938 experiments are shown along with positive (Zip/Zip) and negative (T25 empty vector versus T18-

939 Zip) controls. Photos set to grayscale. **(B) β-Galactosidase assay** results from experiments

940 performed as described in Panel (A). Y-axis is a measurement of relative units of beta-galactosidase

941 activity. X-axis indicates the test conditions. Interactions are considered positive when at least five

942 times the activity of the negative control is measured. Only homotypic interactions were positive in

943 these assays. Note that the inactive mutants of each ClpP also interact homotypically. **(C) Test of**

944 **ClpP Oligomerization by Native-PAGE**. 5 µg samples were run on 4-20% native-PAGE gels and

945 stained with Coomassie for protein detection. Representative results from at least three experiments

946 with independent protein purifications are shown. The Ctr ClpP1 methionine mutant (M6L/M7I) was

947 run in a separate lane and the contrast was uniformly adjusted to help with band visualization. Native

948 molecular weight markers are shown to the left of the gel. *E. coli* ClpP (Ec ClpP) is included as a

949 positive control on the far right of the gel. **(D) Test of Hetero-oligomerization between Ctr ClpP1**

950 **and Ctr ClpP2.** Each recombinant protein was incubated together prior to electrophoresis. The gel

951 has been cropped and enlarged to aid in detecting P1 and P2 hetero-oligomers, which appear to be

952 absent consistent with panel (B) results. The predicted heptamer/tetradecamer sizes in kDa are: Ctr

953 ClpP1, 154/308; Ctr ClpP2 161/322; and Ec ClpP 168/336.

955 **Figure 5: Chlamydial ClpP2, but not ClpP1, has chaperone-independent protease activity.** *In*

956 *vitro* protease activity of the *C. trachomatis* (Ctr) ClpP proteins versus a peptide substrate. ClpP

957 samples (1 µM) were incubated with the fluorometric peptide Suc-Luc-Tyr-AMC, and fluorescence

958 was analyzed over a time. **(A) Relative activity** is shown for each protein assayed with or without

959 the sodium citrate buffer. Use of sodium citrate buffer is indicated as "Sod-Cit". **(B) Loss of activity**

is observed for the Ctr ClpP2 catalytic serine mutant. **(C)** *E. coli* **ClpP** (Ec ClpP) was used as a

positive control. Experiments were run a minimum of three times and baseline assay values obtained

with buffer alone were subtracted from each sample. Average values are reported with standard

error.

Figure 6: ClpP2, but not ClpP1, activity is stimulated by the antibiotic ACP1, which is

detrimental to chlamydial growth. (A-C) Casein (1 μg) was incubated with 1 μM of the respective

ClpP at 37 °C for 3 hours with or without the ACP compounds (used at 500 μM). Reactions were

halted by mixing samples with Laemmli followed by boiling. Samples were run on 12% SDS-PAGE

and stained with Coomassie. Molecular weight markers are shown to the left of each gel.

Representative gels are shown; at least three experiments were performed for each protein. Ctr = *C.*

trachomatis. Ec = *E. coli* **(D) Cell viability** of ACP treated cells were analyzed with resazurin

assays. Values are reported as a percentage of the vehicle control and are representative of three

independent experiments. **(E) Reinfection models** of ACP drug-treated cells reported on a Log_{10}

scale. A student's two-tailed T test was used to compare each parameter to the vehicle control (***=

P<0.0001). For (D) and (E), HEp2 cells were infected with *C. trachomatis* serovar L2, and the ACP1

compounds or DMSO only was added to the infected cultures at 8 hours post-infection (hpi). At 24

hpi, cell viability was assessed for (D), or infected cells were collected to re-infect fresh monolayers

to determine the recovery of inclusion forming units (IFUs) as described in Materials and Methods

for (E).

Fig. 1

A.

	Ctr ClpP1	Ctr ClpP2	Ec ClpP	Bs ClpP	Mtb ClpP1	Mtb ClpP2	Pa ClpP1	Pa ClpP2
Ctr ClpP1	-	39%/58%/85%	44%/63%/84%	41%/61%/91%	38%/55%/87%	38%/58%/79%	42%/62%/80%	36%/62%/86%
Ctr ClpP2	39%/58%/91%	-	55%/78%/78%	58%/77%/100%	52%/74%/87%	44%/68%/90%	58%/78%/91%	45%/66%/88%

```
                                          Ψ                              ΨΩ      ΩΩ
Ctr P1   1 -------------------MPEGEMMHKLQDVIDRKLLDSRRIFFSEPVTEKSAAEAIKKLWYLELTNPGQ  52
Ctr P2   1 ---------------MTLVPYVVEDTGRGERAMDIYSRLLKDRIVMIGQEITEPLANTVIAQLLFLMSEDPTK  58
Ec P     1 ---MSYSGERDNFA-PHMALVPMVIEQTSRGERSFDIYSRLLKERVIFLTGQVEDHMANLIVAQMLFLEAENPEK  71
Bs P     1 ------------------MNLIPTVIEQTNRGERAYDIYSRLLKDRIIMLGSAIDDNVANSIVSQLLFLAAEDPEK  58
Mtb P1   1 ------------------MSQVTDMRSNSQGLSLTDSVYERLLSERIIFLGSEVNDEIANRLCAQILLLAAEDASK  58
Mtb P2   1 -----MNSQNSQIQPQARYILPSFIEHSSFGVKESNPYNKLFEERIIFLGVQVDDASANDIMAQLLVLESLDPDR  70
Pa P1    1 MSRNSFIPHVPDIQ-AAGGLVPMVEQSARGERAYDIYSRLLKERVIFLVGQVEDYMANLVVAQLLFLEAENPEK  74
Pa P2    1 ----MK----------TDDKDREGGDSHGAIGAKLMEYALKVRKVFVTGGVDEKMAKDVVQQLHILASI-SDD  58

             Ψ              Ω              Ω
Ctr P1  53 PIVFVINSPGGSVDAGFAVWDQIKMISSPLTTVVTGLAASMGSVLSLCAVPGRRFATPHARIMIHQPSIGGTITG 127
Ctr P2  59 DIQIFINSPGGYITAGLAIYDTIRFLGCDVNTYCIGQAASMGALLLSAGTKGKRYALPHSRMMIHQPS--GGIIG 131
Ec P    72 DIYLYINSPGGVITAGMSIYDTMQFIKPDVSTICMGQAASMGAFLLTAGAKGKRFCLPNSRVMIHQPL--GGYQG 144
Bs P    59 EISLYINSPGGSITAGMAIYDTMQFIKPKVSTICIGMAASMGAFLLAAGEKGKRYALPNSEVMIHQPL--GGAQG 131
Mtb P1  59 DISLYINSPGGSISAGMAIYDTMVLAPCDIATYAMGMAASMGEFLLAAGTKGKRYALPHARILMHQPL--GGVTG 131
Mtb P2  71 DITMYINSPGGFTSLMAIYDTMQYVRADIQTVCLGQAASAAAVLLAAGTPGKRMALPNARVLIHQPSLSGVIQC 145
Pa P1   75 DIHLYINSPGGSVTAGMSIYDTMQFIKPNVSTTCIGQACSMGALLLAGGAAGKRYCLPHSRMMIHQPL--GGFQG 147
Pa P2   59 PIYMFVNSPGGHVESGDMIFDAIRFITPKVIMIGSGSVASAGALIYAAADKENRYSLPNTRFLLHQPS--GGIQG 131

                                            Ω
Ctr P1 128 QATDLDIHAREILKTKARIIDVYVEATGQSPEVIEKAIDRDMWMSANEAMEFGLLDGILFSFNDL------- 192
Ctr P2 132 TSADIQLQAAEILTLKKHLSNILAECTGQSVEKIIEDSERDFFMGAEEAIAYGLIDKVISSAKETKDKSIAS 203
Ec P   145 QATDIEIHAREILKVKGRMNELMALHTGQSLEQIERDTERDRFLSAPEAVEYGLVDSILTHRN--------- 207
Bs P   132 QATEIEIAAKRILLKDKLNKVLAERTGQPLEVIERDTDRDNFKSAEEALEYGLIDKILTHTEDKK------- 197
Mtb P1 132 SAADIAIQAAEQFAVIKKEMFRLNAEFTGQPIERIEADSDRDRWFTAAEALEYGFVDHIITRAHVNGEAQ--- 200
Mtb P2 146 QFSDLEIQAAEIERMRTLMETTLARHTGKDAGVIRKDTDRDKILTAEEAKDYGIIDTVLEYRKLSAQTA--- 214
Pa P1  148 QASDIEIHAKEILFIKERLNQILAHHTGQPLDVIARDTDRDRFMSGDEAVKYGLIDKVMTQRDLAV------ 213
Pa P2  132 PASNIEIYRREIVRMKERLDRIFAEATGQTPEKISADTERDFWLNAEEAVQYGLVNKIIVSEREITLPGQ-- 201
```

B.

Oligomer Interface	(yellow)
Catalytic Triad	(red)
Hydrophobic Docking Site	(green)

C.

Front   Rear

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6