Functional Testing of Thousands of Osteoarthritis-Associated Variants for Regulatory Activity

Jason Chesler Klein[1]*#, Aidan Keith[1]*, Sarah J. Rice[2], Colin Shepherd[2], John Loughlin[2], Jay Shendure[1,3,4#]

*authors contributed equally

1. Department of Genome Sciences, University of Washington, Seattle, WA 98195
2. Skeletal Research Group, Institute of Genetic Medicine, Newcastle University, International Centre for Life, Newcastle-upon-Tyne, NE1 3BZ, UK
3. Brotman Baty Institute for Precision Medicine, Seattle, WA 98195
4. Howard Hughes Medical Institute, University of Washington, Seattle, WA 98195

Correspondence to jcklein@uw.edu and shendure@uw.edu

Abstract

To date, genome-wide association studies have implicated at least 35 loci in osteoarthritis, but due to linkage disequilibrium, we have yet to pinpoint the specific variants that underlie these associations, nor the mechanisms by which they contribute to disease risk. Here we functionally tested 1,605 single nucleotide variants associated with osteoarthritis for regulatory activity using a massively parallel reporter assay. We identified six single nucleotide polymorphisms (SNPs) with differential regulatory activity between the major and minor alleles. We show that our most significant hit, rs4730222, drives increased expression of an alternative isoform of *HBP1* in a heterozygote chondrosarcoma cell line, a CRISPR-edited osteosarcoma cell line, and in chondrocytes derived from osteoarthritis patients.

Main text

Genome-wide association studies (GWAS) have successfully implicated thousands of genetic loci in common human diseases. Most of the underlying signal is believed to derive from variation in non-coding regulatory sequences. However, because of linkage disequilibrium (LD), it has been extraordinarily challenging for the field to identify the variants that causally underlie each association.

Over the past decade, we and others developed massively parallel reporter assays (MPRAs) to increase the throughput at which regulatory sequences can be tested for functional potential[1–4]. An MPRA involves cloning thousands of candidate regulatory sequences to a single reporter gene, transfecting them to a cell line *en masse*, and performing deep sequencing of the resulting transcripts to quantify the degree of transcriptional activation mediated by each candidate

41 regulatory sequence. MPRAs have previously been applied to characterize variants underlying
42 eQTLs (in LCLs)[5], red blood cell traits (in K562 and human erythroid progenitors/precursors)[6],
43 cancer-associated common variants (in HEK293 cells)[7], and adiposity-associated common
44 variants (in HepG2 cells[8]).
45
Here we sought to apply an MPRA (specifically, STARR-seq[4]) to quantify the relative regulatory
potential of SNPs residing on haplotypes implicated in osteoarthritis (OA), with the aim of
pinpointing causal variants. We compiled a list of 35 lead SNPs associated with OA in
Europeans via GWAS, with minor allele frequencies over 5%[9–26]. Each SNP represents an
independent signal with $p<5e-8$ (genome-wide significant; n=20) or $p<5e-5$ (genome-wide
suggestive; n=15) (**Table S1**). We identified all SNPs in LD with an $r^2>0.8$ in Europeans using
rAggr (**Fig. 1A**), resulting in a list of 1,605 candidate SNPs. For the major and minor allele of
each SNP, we synthesized 196 nt of genomic sequence, centered on the SNP and flanked by
adaptor sequences, on a microarray (230 nt oligos; **Fig. 1B**).
55
During PCR amplification of array-derived oligos, we appended 5 nt degenerate barcodes (**Fig.
1C**), such that each allele would be represented by multiple independent measurements in the
subsequent experiment. We then cloned these barcoded oligos into the human STARR-seq
vector and transfected Saos-2 cells, an osteosarcoma cell line. In STARR-seq, the candidate
regulatory sequences are located within the transcript itself (**Fig. 1D**). From the transfected Saos-
2 cells, we extracted, amplified and sequenced both DNA and RNA corresponding to the cloned
region, and then calculated activity scores as the normalized log2 (ratio of RNA reads / ratio of
DNA reads) for each barcode-allele combination (**Fig. 1E-F**). For all alleles with greater than
five independent measurements over three biological replicates (independent transfections of the
same library), we averaged the allele activity scores to a single value. Due to bottlenecking
during library construction, some alleles were under sampled and therefore excluded from
further analysis. Altogether, we obtained activity scores for 1,953 of 3,210 alleles (61%), and
activity scores for both alleles of 752 of 1,605 SNPs (47%).
69
We first asked whether these STARR-seq-based activity scores correlated with biochemical
marks for putative enhancers. For this analysis, activity scores corresponding to two alleles of the
same SNP were collapsed. We ranked and split the resulting 1,203 activity scores for distinct
genomic sequences into quintiles, and then intersected these with datasets of biochemical marks
of putative enhancers in cartilage and bone (H3K27ac in bone marrow-derived chondrocytes,
H3K27ac in human embryonic limb buds, and ATAC-seq in knee OA cartilage)[27,28]
(**Supplementary Fig. 1**). The highest scoring quintile was significantly enriched for overlap with
H3K27ac ChIP-seq peaks in embryonic limb bud from E41 (2.1-fold, Bonferroni-corrected chi-
square p=0.0096), E44 (1.8-fold, p=0.044), and E47 (2.0-fold, p=0.0072), but not with knee OA
cartilage ATAC-seq peaks nor bone marrow-derived chondrocyte H3K27ac peaks. These results
are in line with our use of an osteosarcoma rather than a cartilage-derived cell line. The highest

scoring quintile includes 240 genomic regions, 67 of which overlap putative enhancers from at
least one dataset. In contrast, the least active quintile includes 239 genomic regions, only 37 of
which overlap putative enhancers from at least one dataset (1.8-fold difference, chi-square
p=9.7e-4). Altogether, these enrichments demonstrate that at least a subset of the 1,605 genomic
regions tested here correspond to enhancers in OA-relevant tissues. All activity scores are
included in **Table S2.**

We next sought to ask whether any alleles are differentially active, focusing on the 752 SNPs for
which we successfully measured activity scores for both alleles (**Table S3**). Overall, activity
scores for two alleles of a given SNP were highly correlated, with an overall Spearman
correlation of 0.96 (**Fig. 1G**). This was reassuring, given that each pair of alleles was separately
synthesized and cloned, and therefore at non-identical abundances in the STARR-seq library.
After correcting for multiple testing with Benjamini-Hochberg (BH) at a 5% FDR, we identified
6 SNPs whose alleles demonstrated significantly differential functional activity in Saos-2 cells
(**Fig. 1G, Table S3**). The most significant SNP, rs4730222, is located in the 5' UTR of several
non-canonical isoforms of HMG-Box Transcription Factor 1 (*HBP1).* Other significant SNPs
include rs80095766 (intronic to *COG5*), rs2286798 (intronic to *ITIH1*), rs11745630 (downstream
of *PIK3R1*), rs6976 (3' UTR of *GLT8D1*), and rs1563351 (intronic to *LOC102723886*). Two
pairs of the six significant SNPs are at the same loci (the ones near *HBP1* and *COG5*, both at
chromosome 7q22.3; and the ones near *ITIH1* and *GLT8D1*, both at chromosome 3p21.1).

We chose to further characterize rs4730222 for several reasons. First, this SNP was the most
significant hit from our reporter assay, with a substantial difference in activity between the two
alleles (2.5-fold increased expression of the minor, disease-associated allele, BH adjusted p-
value=2.4e-6). Second, it overlaps several marks for active regulatory elements. Third, we have
previously observed reduced expression of the canonical *HBP1* transcript in relevant tissues from
carriers of the OA-associated allele[29], interestingly opposite the effect observed here. HBP1 is a
transcriptional repressor that regulates the Wnt-beta-catenin pathway as well as superoxide
production, both of which have been implicated in OA development and progression[30,31,32].

rs4730222 overlaps with several marks associated with active regulatory DNA: H3K27ac (mark
for active enhancers and promoters) from bone marrow-derived chondrocytes[28], human
embryonic limb bud at E33, E41, E44 and E47[27], and ENCODE layered data, H3K4me3 (mark
for active promoters) in ENCODE layered data, and ATAC-seq (mark for open chromatin) peaks
from articular knee cartilage of OA patients[33] (**Fig. 2A**). *HBP1* contains several transcript
isoforms, with three probable alternative promoters identified from cap-selected clones and nine
validated alternative polyadenylation sites[34]. One of the three probable alternative promoters
contains rs4730222 at position +80 relative to the alternative TSS. We therefore hypothesized
that the variant may alter expression of this or another *HBP1* isoform. To test this, we first
confirmed that the alternative TSS overlapping rs4730222 is utilized in Saos2 and SW1353 cells

(a chondrosarcoma cell line) with qRT-PCR with primers contained within this 5' UTR as well
as spanning to the following exon. However, despite multiple attempts, we failed to successfully
amplify the isoform from the alternative TSS to the canonical stop, suggesting that the
alternative TSS may belong to a truncated isoform of the *HBP1* transcript. Based on PCR and
Sanger sequencing, the truncated isoform is most likely ENST00000497535, which contains the
alternative 5' UTR and first two exons (**Fig. 2A, Supplementary Fig. 2**).
127
After confirming that rs4730222 is transcribed as part of a *HBP1* isoform in osteogenic and
chondrogenic cells (albeit not to the canonical stop), we genotyped rs4730222 in several cell
lines (SW1353, Tc28a/2, Saos-2, chondrogenic progenitor cells) in hopes of identifying a
heterozygous line. As SW1353 was heterozygous for rs4730222, we tested it for allelic
expression imbalance (AEI) of the transcribed SNP. In all three biological replicates, we
observed a significant allelic imbalance (Fisher's exact test, $p < 1e-5$), with the minor allele
showing a 1.3 to 1.4-fold relative enrichment in RNA/DNA compared to the major allele. This
was more modest but directionally concordant with 2.5-fold enrichment of the minor allele in the
reporter assay (**Fig. 2B**).
137
The OA-implicated haplotype at 7q22.3 is a ~500 kb region of high linkage disequilibrium, and
consequently accounts for 283 of the 1,605 SNPs tested here. The observed AEI of the *HBP1*
alternative isoform in SW1353 cells, which are likely heterozygous for the entire OA-associated
haplotype, could be consequent to any or several of these SNPs, or to other forms of genetic
variation. To test whether rs4730222 causally underlies the allelic imbalance of the *HBP1*
alternative transcript, we introduced the minor allele of rs4730222 into Saos-2 cells, which are
homozygous for the major allele, through CRISPR-mediated homology directed repair (HDR).
We generated four biological replicates (*i.e.* independently edited cell populations), and
quantified the RNA-DNA ratio of indel-free reads derived from the major vs. minor allele.
Similar to the SW1353 AEI (1.3 to 1.4-fold), we identified a 1.4 to 1.6-fold relative enrichment
in the RNA/DNA ratio for the minor allele compared to the major allele (Fisher's exact test,
$p < 1e-5$; **Fig. 2C**). This confirms that the minor allele at rs4730222 causally underlies
upregulation of the *HBP1* alternative isoform.
151
We next sought to test whether rs4730222 drives increased transcription of the alternative TSS in
osteoarthritic tissue. To do so, we tested for AEI in chondrocytes derived from OA patients. For
each of nine patients heterozygous for rs4730222, we extracted RNA and DNA and amplified
and sequenced the 5'UTR containing the SNP (three technical replicates per patient) from DNA
and cDNA. Despite low cDNA concentrations and low expression of the alternative TSS, we
observed an overall trend in AEI in accordance with our reporter assay and cell models (median
AEI=1.12, Mann Whitney U-test on 27 DNA ratios vs. 27 RNA ratios, p=0.003). Interestingly,
one patient appeared to exhibit AEI in the opposite direction[35,36] (**Supplementary Fig. 3**).
160

In summary, we set out to functionally test the regulatory effects of 1,605 SNPs that potentially
underlie 35 GWAS signals for osteoarthritis. We succeeded in generating reproducible
measurements of regulatory activity for about two-thirds of the regions tested, and of differential
regulatory activity for about half of the regions tested. The most highly active regions in our
assay were ~2-fold enriched for biochemical marks associated with enhancers. We furthermore
identified six SNPs, which each drove differential expression at an FDR of 5%. The most
significant of these, rs4730222, resides in the 5' UTR of multiple isoforms of *HBP1*, a
transcriptional repressor. The minor, OA-associated risk allele of rs4730222 increases
transcription of alternative isoform(s) of *HBP1*. We validated this finding in SW1353 cells,
CRISPR-edited Saos2 cells, and in chondrocytes derived from knee OA patients.

We previously described reduced expression of the canonical *HBP1* transcript in OA tissue,
relative to healthy tissue[29]. Here, we do not observe AEI of the canonical *HBP1* transcript, but
rather AEI of an alternative transcript. We speculate that the impact of rs4730222, wherein the
disease-associated allele consistently increases expression of an alternative transcript,
secondarily reduces expression of canonical *HBP1*. One possibility is that the alternative isoform
encodes a non-coding RNA with an upstream open reading frame (uORF). This isoform of
HBP1 may have a *trans* effect on the canonical transcript, or have its own, yet uncharacterized,
function in the cell. Additionally, alternative TSSs have been shown to modulate translational
efficiency and tissue specificity of genes[37,38]. Through one or several of these mechanisms, the
expression levels of this isoform might modulate HBP1 levels in certain tissues. Finally, we note
that the rs4730222-overlapping isoform expressed in SW1353 and Saos2 is a truncated version
of *HBP1*. Therefore, it may disrupt endogenous activity of the gene, *e.g.* by acting as a
dominant-negative. Distinguishing between these mechanistic possibilities for rs4730222 and
HBP1, as well as for other MPRA-prioritized OA-associated SNPs, should be a high priority for
the field.

Acknowledgements
The authors thank the Shendure and Loughlin labs, particularly J. Alexander, M. Gasperini, and
S. Kim, for helpful discussions. This work was funded by grants from the National Institutes of
Health (UM1HG009408, R01CA197139, R01HG006768) to J.S. J.C.K. was supported in part by
1F30HG009479 from the NHGRI. J.S. is an investigator of the Howard Hughes Medical
Institute. S.J.R., C.S. and J.L. were supported by Arthritis Research UK (grant 20771), and by
the Medical Research Council and Arthritis Research UK as part of the MRC-Arthritis Research
UK Centre for Integrated Research into Musculoskeletal Aging (CIMA, grant references JXR
10641, MR/P020941/1 and MR/R502182/1).

Author Contributions
The project was conceived and designed by J.C.K., A.K., J.S., and J.L. S.J.R. and C.S. collated
and generated the OA patient DNA and cDNA samples. J.L. compiled OA lead variants. J.C.K.

and A.K. performed all experiments and analyses. J.C.K., A.K. and J.S. wrote the manuscript.
All authors read and approved the final version of the manuscript.
203

204 **Main Figures**
205

206
207

Figure 1. Schematic and results from massively parallel reporter assay. A) For each GWAS-
lead SNP, we identified all SNPs in LD with $r^2 > 0.8$. Colored lines indicate SNPs in the same
LD block. B) For all SNPs, we extracted 196 nt of genomic sequence centered at the SNP, and
separately synthesized the minor (hollow circle) and major alleles, flanked by common adaptor
sequences (cyan and purple). C-D) We amplified our library from the array via PCR with
primers directed at the common adaptors, in the process appending 5 nt degenerate barcodes
(black lines) and additional sequences homologous to the vector (cyan). We cloned our barcoded
library of all major and minor alleles into the STARR-seq vector. Each putative regulatory
region is cloned into the 3' UTR of a reporter gene (cyan) with a minimal promoter (dark blue).
E) We transfected our library into Saos-2 cells via electroporation. 48 hours post transfection, we
extracted RNA and DNA. F) We determined the abundance of each allele-barcode combination
in the mRNA and DNA population through sequencing. G). For each allele, we calculated one
activity score as the average log2(RNA/DNA) across all independent measurements. We
identified six SNPs with significantly different activity between the major and minor alleles.
222

Figure 2. Functional Validation of rs4730222. A) Gene model of *HBP1* ensemble isoform
ENST00000468410 (major) and ENST00000497535 (alternative). UCSC genome browser
zoomed in to the two transcriptional start sites (http://genome.ucsc.edu). Rs4730222, within the
5' UTR of the alternative transcript, is indicated by a vertical grey line traversing the annotation
tracks. The four black tracks are H3K27ac performed in human embryonic limb bud at E33, E41,
E44 and E47 respectively[27]. The green track is H3K27ac data from chondrocytes derived from
cultured bone marrow mesenchymal stem cells[28]. Layered H3K27ac is H3K27ac ChIP-seq (a
marker for active enhancers and active promoters) layered from GM12878, H1-hESC, HSMM,
HUVEC, K562, NHEK, and NHLF cells. Layered H3K4me3 is H3K4me3 ChIP-seq (marker for
active promoters) layered from the same seven ENCODE cell lines. B) Allelic expression
imbalance in SW1353, a chondrosarcoma cell line heterozygote for rs4730222. Black bars
represent the fraction of the minor allele in DNA and grey bars indicate the fraction of the minor
allele in cDNA. C) Allelic expression imbalance in Saos-2 cells with the minor allele of
rs4730222 introduced through CRISPR-mediated HDR. Bars are the same as in **Fig. 2B**.

Lead SNP	Nearest protein coding gene	Ref	Annotation	#Linked SNPs tested	Significance
rs6976	GLT8D1	9	3' UTR	280	GWS
rs12107036	TP63	9	intronic	1	GWSu
rs10948172	SUPT3H	9	intronic	213	GWSu
rs9350591	FILIP1	9	intergenic	73	GWS
rs3815148	HBP1	10	intronic	284	GWS
rs4836732	ASTN2	9	intronic	2	GWS
rs10492367	KLHDC5	9	intergenic	5	GWS
rs835487	CHST11	9	intronic	7	GWS
rs11842874	MCF2L	11	intronic	8	GWS
rs225014	DIO2	12	exonic	19	GWSu
rs945006	DIO3	13	3' UTR	1	GWSu
rs3204689	ALDH1A2	14	3' UTR	52	GWS
rs8044769	FTO	9	intronic	8	GWSu
rs12982744	DOT1L	15	intronic	21	GWSu
rs6094710	NCOA3	16	intergenic	11	GWS
rs143383	GDF5	17	5' UTR	98	GWS
rs4764133	MGP	18	intergenic	137	GWS
rs3850251	ENPP3	18	intronic	6	GWSu
rs754106	LRCH1	19	intronic	10	GWSu
rs6766414	STT3B	19	intergenic	29	GWSu
rs2862851	TGFA	20	intronic	25	GWS
rs10471753	PIK3R1	20	intergenic	29	GWS
rs2236995	SLBP	20	intronic	4	GWS
rs496547	TREH	20	intergenic	3	GWS
rs4867568	LSP1P3	21	intergenic	6	GWSu
rs788748	IGFBP3	22	intergenic	10	GWSu
rs12901499	SMAD3	23	intronic	28	GWSu
rs4907986	COL11A1	24	intronic	23	GWSu
rs1241164	COL11A1	24	intronic	29	GWSu
rs833058	VEGF	24	intergenic	1	GWSu
rs10116772	GLIS3	25	intronic	8	GWS
rs2820436	ZC3H11B	26	intronic	43	GWS
rs11335718	ANXA3	26	intronic	1	GWS
rs11780978	PLEC	26	intronic	110	GWS
rs2521349	MAP2K6	26	intronic	20	GWS

Supplementary Table 1. Lead SNPs. List of 20 genome-wide significant (GWS) and 15 genome-wide
suggestive (GWSu) variants compiled in May 2017.

Supplementary Figures

Supplementary Figure 1. Overlap between tested sequences and enhancer marks. The 1,203

SNPs were split into 5 quintiles of ~240 sequences each, based on their normalized RNA/DNA

activity score. Q1 refers to the sequences with the highest activity scores and q5 refers to the

sequences with the lowest activity scores. We then overlapped each quintile with peaks called

from OA ATAC-seq[33], BMD-chondrocytes H3K27ac ChIP-seq[28], and human embryonic limb

bud H3K27ac ChIP-seq from E33, E41, E44, and E45[27]. Y-axis is the fraction of the 240 SNPs

in each quintile overlapping peaks.

Supplementary Figure 2. Relative expression of different HBP1 isoforms. A) Y axis is the Cq value from RT-qPCR from SW1353 cells. The major allele uses a forward primer at the start codon and reverse primer at the conserved stop codon of the major isoforms. The 5' UTR primer set amplifies a short product, entirely within the alternative 5' UTR. The truncated primer set amplifies both ENST00000497535 and ESNT00000485846. The full length primer set includes a forward primer in the alternative TSS and reverse primer at the stop codon of the major isoform. We do not identify any full length product utilizing the alternative TSS. B) Gel of qPCR products. First lane is a 2-log ladder. Second lane is the 5' UTR amplification. Expected size is 156 bp. Third lane is the truncated amplification. The primer set should amplify both ENST00000497535 (expected size 548bp) and ENST00000485846 (expected size 846bp). However, ENST00000485846 contains an internal exon while ESNT00000497535 does not. We ran the PCR product on a gel and Sanger sequenced the purified PCR product, which did not include the internal exon. Fourth lane is amplifying from the alternative 5' UTR to canonical stop. There was no amplification product. Fifth lane is the major isoform (amplifying from canonical start to canonical stop. Expected size is 1,455 bp.

286 **Supplementary Figure 3.** Allelic Expression Imbalance (AEI) in cartilage from patients
287 receiving total knee replacements. Black bars are the fraction of DNA reads aligning to the minor
288 allele. Grey bars are the fraction of RNA reads aligning to the minor allele. Bars indicate the
289 minimum and maximum fraction from three technical replicates.

Methods

Identification and design of target SNPs

We selected SNPs that had a minor allele frequency >5% and had been reported as being associated with OA in European populations at a significance level that surpassed or approached the genome-wide threshold of <5e-8. The deadline date for inclusion was May 2017. In total, 35 SNPs were identified, each representing an independent association signal (Table S1). We ran rAGGr on our list of 35 candidate SNPs to identify all variants with a minimum minor allele frequency >= 0.001 in linkage disequilibrium with an r^2>0.8 in Europeans (CEU+FIN+GBR+IBS+TSI) based on 1000 Genomes, Phase 3, Oct 2014. We then filtered out any polymorphisms greater than one nucleotide, resulting in a list of 1,605 SNPs. For each variant, we extracted 196 nt of genomic sequence centered around the SNP using BEDTOOLS getfasta, and edited the SNP to create both the minor and major alleles (3,210 sequences). To each 196 nt sequence, we appended HSS_clon_F (5' - TCTAGAGCATGCACCGG - 3') to the 5' end and DO_R6 (5'- GCCGGTCAGAATGATGG -3') to the 3' end. We then ordered the 3,210 sequences in duplicate as part of an Agilent 244K 230-mer array.

Library Generation

We amplified our sequences off of the Agilent array with HSS_clon_F and R6_5N_HSSR (5'-CCGGCCGAATTCGTCGANNNNNCCATCATTCTGACCGGC -3') using KAPA HiFi HotStart ReadyMix in a 50 uL reaction with 0.75 ng DNA and SYBR Green on a MiniOpticon Real-Time PCR system (Bio-Rad) and stopped the reaction before plateauing (13 cycles). This reaction amplified our library, added a 5 nt degenerate barcode to each sequence, and added both adapters for cloning into the human STARR-seq vector. We purified the PCR product using a 1.5x AMPure cleanup following manufacturer's protocol. We then ligated 6 ng of our purified PCR into 25 ng of linearized human STARR-seq backbone using the NEBuilder HiFi DNA Assembly Cloning Kit following manufacturer's protocol. We transformed 1.2 uL of the ligation product in 50 uL of NEB C3020 cells, grew up overnight in 100 mL of LB+Amp, and extracted the library using a Zymo Research ZymoPURE Plasmid Midiprep Kit.

STARR-seq Screen

We transfected 1.5 million Saos-2 cells with 20 ug of our library in triplicate using the Thermo Fisher Scientific Neon Transfection System with resuspension buffer R at 1250V, 40ms, 1 shock, with 100 uL pipettes, in triplicate. After electroporation, we added the cells to 10cm plates with pre-warmed media (McCoy's 5A with 10% FBS and 1x Pen/Strep). 48 hours post transfection, we extracted both DNA and RNA from each replicate using the Qiagen ALLPrep DNA/RNA Mini Kit. DNA was eluted in 80 uL and RNA was eluted in 30 uL. RNA was treated with Thermo Fisher Scientific TURBO DNase following manufacturer's protocol and reverse transcribed using Thermo Fisher Scientific SuperScript III Reverse Transcriptase in a 20 uL reaction with 8 uL of RNA. For each replicate, we amplified DNA in two reactions, each with 2

ug of DNA using NEBNext High Fidelity 2X PCR Master Mix with primers HSS_NF_pu1 (5'-
CTAAATGGCTGTGAGAGAGCTCAGGTACAACTGATCTAGAGCATGCACC   -3')   and
HSS_R_pu1.(5'- ACTTTATCAATCTCGCTCCAAACCCTTATCATGTCTGCTCGAAGC -3')
and stopped before plateauing (15 cycles). After PCR, products were purified with a 1.5x
AMPure cleanup, and pooled together. For each replicate, we also amplified cDNA in two
reactions, each with 10 uL of RT product in 50 uL reactions with NEBNext High Fidelity 2X
PCR   Master   Mix   with   primers   HSS_F_pu1   (5'-
CTAAATGGCTGTGAGAGAGCTCAGGGGCCAGCTGTTGGGGTGTCCAC-3')   and
HSS_R_pu1 (5'- ACTTTATCAATCTCGCTCCAAACCCTTATCATGTCTGCTCGAAGC -3')
and stopped before plateauing (18-20 cycles). After PCR, products were purified with a 1.5x
AMPure cleanup, eluted in 50 uL each, and pooled together. For the cDNA samples, we
performed a nested reaction using KAPA HiFi HotStart ReadyMix in a 50 uL reaction with 1 uL
of   the   pooled   outer   PCR   reaction   with   HSS-NF-pu1   and   pu1R   (5'-
ACTTTATCAATCTCGCTCCAAACC -'3) and stopped before plateauing (7 cycles). Reactions
were purified with a 1.5x AMPure cleanup and eluted in 50 uL each. Flow cell adapters and
indexes were added to all DNA and cDNA reactions through an additional round of PCR using
Kapa HiFi HotStart ReadyMix in 50 uL reactions with 1 uL of the first DNA PCR or 1 uL of the
inner   cDNA   PCR   using   an   indexed   pu1_P5   primer   (5'-
AATGATACGGCGACCACCGAGATCTACACNNNNNNNNNNACGTAGGCCTAAATGGC
TGTGAGAGAGCTCAG   -3')   and   an   indexed   pu1_P7   primer   (5'-
CAAGCAGAAGACGGCATACGAGATNNNNNNNNNGACCGTCGGCACTTTATCAATCT
CGCTCCAAACC -3') and stopped before plateauing (6 cycles). The libraries were sequenced
on an Illumina NextSeq 500/550 v2 300 cycle mid-output kit.
354
Analysis of STARR-seq Screen
We aligned all sequencing reads to a reference fasta file of our variants using BWA mem and
extracted reads from error-free molecules [39]. Each variant contained several different 5 nt
barcodes added through PCR. We counted the number of reads from each replicate for each
variant-barcode combination in the DNA and cDNA pool. If there were at least 10 DNA reads
and at least 1 RNA read, we calculated an activity score as the log2(number of RNA reads from
the variant-barcode combination normalized to the total number of RNA reads, divided by the
number of DNA reads from the variant-barcode combination normalized to the total number of
DNA reads). We then combined all variant-barcode activity scores from each replicate, and for
any variant with at least five different measurements, we averaged the activity score for a final
activity score for each variant. This resulted in activity scores for 1,953 of the 3,210 alleles. 752
of the 1,605 variants contained measurements for both alleles. For each of the 752 variants with
measurements for both alleles, we tested whether the 2 alleles drove different expression by
performing a Mann-Whitney U Test for each variant using SciPy v0.19.1 with Python v2.7.3.
We then performed a Benjamini-Hochberg correction with an FDR = 0.05 to correct for multiple
testing.

371
Allelic Imbalance of rs4730222 in SW1353 cells
We first genotyped several osteogenic and chondrogenic cell lines for rs4730222 (SW1353, Tc28a/2, Saos-2, chondrogenic progenitor cells) using rs4730222_sangerF (5'- TACGCAGTTCGAATGAATGGGCTC -3') and rs4730222_sangerR (5'- AGCTACAAAAACCTGGCTGTCCAC -3'). PCR products were purified with a 1.5x AMPure cleanup and Sanger sequenced with rs4730222_sangerF.

We then tested for allelic imbalance of rs4730222 in the isoforms expressing the SNP in SW1353. We performed three independent DNA and RNA extractions using the Qiagen ALLPrep DNA/RNA Mini Kit. DNA was eluted in 80 uL and RNA was eluted in 30 uL. RNA was treated with TURBO DNase and reverse transcribed with SuperScript III Reverse Transcriptase. We then amplified the 5'UTR around rs4730222 from each DNA and cDNA sample using KAPA HiFi HotStart ReadyMix in a 50 uL reaction with 100 ng DNA or 5 uL cDNA with HBP1_5UTR_F_pu1 (5'- CTAAATGGCTGTGAGAGAGCTCAGAGTCCGGGCTGCGGTCACATGATG -3') and HBP1_5UTR_R_pu1 (5'- ACTTTATCAATCTCGCTCCAAACCAGCTACAAAAACCTGGCTGTCCAC -3') and stopped DNA reactions at 25 cycles and cDNA reactions at 32 cycles. Products were purified using a 1.5x AMPure cleanup, and flow cell adapters and indexes were added using an indexed pu1_P5 primer and an indexed pu1_P7 primer. Libraries were spiked into a Miseq v2 300 cycle run. Reads were aligned to a fasta reference file using BWA mem and the number of perfect reads coming from both alleles was quantified from both DNA and cDNA from each replicate.

CRISPR Knock-in of rs4730222 in Saos-2 cells
rs4730222 falls within a potential Cas9 PAM site (5'- ACGCGATGAATGGCGAAAGA G**G**G -3'). We therefore designed a guideRNA that would target rs4730222, so that the minor-allele donor would not be re-cut. We ordered the following oligos from IDT: rs4730222_guideF (5'- CACCGACGCGATGAATGGCGAAAGA -3') and rs4730222_guideR (5'- AAACTCTTTCGCCATTCATCGCGTC -3') and followed the Zhang lab protocol to clone them into the px458 plasmid (SpCas9-2A-EGFP and single guide RNA).

We created our donor vector in two steps. First, we amplified a 1,459 bp region around rs4730222 with the following primers, which also append 16 bp homologous sequence to puc19 onto each side of the amplicon: HBP1_puc19F (5'- TCGGTACCCGGGGATCAAGTAGGAAAGTTTCGGTTGAGGAG -3') and HBP1_puc19R (5'- TCGACTCTAGAGGATCAACTGAACAGATGACCGACTCTACC -3). We then cloned this into a linearized puc19 plasmid using Clontech's In-Fusion HD Cloning Kit following manufacturer's protocol, transformed into Stellar Competent cells, grew up a single colony and extracted plasmid using the Zymo Research ZymoPURE Plasmid Midiprep Kit. We then re-linearized the puc19-HBP1 wild-type plasmid via PCR with puc19_HBP1-linF (5'- GTGGGGGATGGACTTGGCGTG -3') and puc19-HBP1-linR (5'-

CTCCTCAACCGAAACTTTCCTACTT -3'). We also amplified a small region around rs4730222, while mutating the SNP, using mut_insF (5'-AAGTAGGAAAGTTTCGGTTGAGGAG -3') and mut_insR (5'-CCAAGTCCATCCCCCACGCTCTTTCGCCATTCATCGCG -3'). We then cloned the mutated insert into puc19-HBP1-wt using the In-Fusion HD Cloning Kit and grew up a single colony with the minor allele at rs4730222 flanked by 600-850 bp of homology on each side.

We transfected 1 million Saos-2 cells with 10 ug of our px458-rs4730222 guide and 10 ug of our donor library containing the minor allele using the Neon Transfection system as described above. 72 hours post transfection, we performed FACS on a BD FACS Aria III to isolate ~150,000 GFP+ cells (transfected with px458), which we then expanded. On day 10 post transfection, we extracted DNA and RNA, performed reverse transcription with Superscript III, and amplified the region surrounding rs4730222 from both DNA (using HBP1_5UTR_F_pu1 and HBP_DNA_Routside (5'- TAGGTGGGCAATCCTGGGAGAAGGTAC -3')), and RNA (using HBP1_5UTR_F_pu1 and HBP_RNA_Routside (5'- TGCCAGATTCTGACTCACTATTTGC -3')) in 50 uL reactions using KAPA HiFi 2x ReadyMix. We then purified the PCR reactions with a 1.5x AMPure cleanup, eluted in 50 uL, and used 1 uL in a nested reaction with pu1L (5'-CTAAATGGCTGTGAGAGAGCTCAG -3') and HBP1_5UTR_R_pu1. Reactions were purified with a 1.5x AMPure cleanup, and flow cell adapters and indexes were added using an indexed pu1_P5 primer and an indexed pu1_P7 primer. Libraries were spiked into a Miseq v2 300 cycle run. Reads were aligned to a fasta reference file using BWA mem and the number of perfect reads coming from both alleles was quantified from both DNA and cDNA from each replicate.

Allelic imbalance of rs4730222 in osteoarthritis patients' chondrocytes

Cartilage tissue samples were obtained from OA patients who had undergone joint replacement surgery at the Newcastle upon Tyne NHS Foundation Trust hospitals. The Newcastle and North Tyneside Research Ethics Committee granted ethical approval for the collection, with each donor providing verbal and written informed consent (REC reference number 14/NE/1212). Our patient ascertainment criterion

has been described in detail previously [40,41]. The cartilage was removed from the joint using a scalpel and was collected distal to the OA lesion. The tissue samples were stored frozen at -80°C and ground to a powder using a Retsch Mixermill 200 (Retsch Limited) under liquid nitrogen. Nucleic acids were then extracted from the ground tissue using TRIzol reagent (Life Technologies) according to the manufacturer's instructions, with the upper aqueous phase separated for RNA isolation, while the interphase and lower organic phase were used to isolate DNA. RNA was reverse transcribed using the SuperScript First-Strand cDNA synthesis kit (Invitrogen). Matched DNA and cDNA were amplified with KAPA HiFi 2x ReadyMix and SYBR Green and halted before plateauing. The primers sequences were as follows:

5'-CTAAATGGCTGTGAGAGAGCTCAGAGTCCGGGCTGCGGTCACATGATG-3'; and

5'-ACTTTATCAATCTCGCTCCAAACCAGCTACAAAAACCTGGCTGTCCAC-3'.

All samples were purified with a 1.5x AMPure cleanup following manufacturer's instructions
and eluted in 50 uL Qiagen Elution Buffer. 1 uL of purified product was then indexed for
Illumina sequencing using an indexed pu1_P5 primer and an indexed pu1_P7 primer. Libraries
were spiked into a Miseq v2 300 cycle run. Reads were aligned to a fasta reference file using
BWA mem and the number of aligning reads coming from both alleles was quantified from both
DNA and cDNA from each replicate. For statistical analysis, a Mann Whitney U-test was
performed comparing DNA vs. RNA abundances of the minor allele for the 54 values (27 for
DNA and 27 for RNA; nine patients x three replicates).
458
Characterization of HBP1 rs4730222-containing isoforms
We designed the following set of primers to differentiate between different HBP1 isoforms:
Major (1stExonF: 5'- GTGTGGGAAGTGAAGACAAATCAGATGC -3' and LastExonR: 5'-
CTTCCACCTGTCACCAAGGATCACAC   -3'),   5'   UTR   (UTR_qPCR_F: 5'-
CAGTCTCCGCCTTTCAACCTATG   -3'   and   UTR_qPCR_R: 5'-
ATGAACTCGAGTGTAGAGTGCACAG -3'), Truncated (UTR_qPCR_F and Exon6_R:
CCACCTCATTTTCACGGTAAGTAG -3') and Full Len (UTR_qPCR_F and LastExonR). We
performed technical triplicates for each qPCR using KAPA Robust 2x Hotstart Readymix with
cDNA from wild-type SW1353 cells, letting the reaction go for 40 cycles. We then ran products
on a gel, and differentiated between ENST00000497535 and ENST00000485846 based both on
size and Sanger sequencing.
470
471

REFERENCES

1. Patwardhan, R. P. *et al.* High-resolution analysis of DNA regulatory elements by synthetic saturation mutagenesis. *Nat. Biotechnol.* **27,** 1173–1175 (2009).

2. Patwardhan, R. P. *et al.* Massively parallel functional dissection of mammalian enhancers in vivo. *Nat. Biotechnol.* **30,** 265–270 (2012).

3. Melnikov, A. *et al.* Systematic dissection and optimization of inducible enhancers in human cells using a massively parallel reporter assay. *Nat. Biotechnol.* **30,** 271 (2012).

4. Arnold, C. D. *et al.* Genome-wide quantitative enhancer activity maps identified by STARR-seq. *Science* **339,** 1074–1077 (2013).

5. Tewhey, R. *et al.* Direct Identification of Hundreds of Expression-Modulating Variants using a Multiplexed Reporter Assay. *Cell* **172,** 1132–1134 (2018).

6. Ulirsch, J. C. *et al.* Systematic Functional Dissection of Common Genetic Variation Affecting Red Blood Cell Traits. *Cell* **165,** 1530–1545 (2016).

7. Liu, S. *et al.* Systematic identification of regulatory variants associated with cancer risk. *Genome Biol.* **18,** 194 (2017).

8. Vockley, C. M. *et al.* Massively parallel quantification of the regulatory effects of noncoding genetic variation in a human cohort. *Genome Res.* **25,** 1206–1214 (2015).

9. Consortium, A. & Others. arcOGEN Collaborators, Zeggini E, Panoutsopoulou K, Southam L, Rayner NW, et al. Identification of new susceptibility loci for osteoarthritis (arcOGEN): a genome-wide association study. *Lancet* **380,** 815–823 (2012).

10. Kerkhof, H. J. M. *et al.* A genome-wide association study identifies an osteoarthritis susceptibility locus on chromosome 7q22. *Arthritis Rheum.* **62,** 499–510 (2010).

11. Day-Williams, A. G. *et al.* A variant in MCF2L is associated with osteoarthritis. *Am. J.*

Hum. Genet. **89,** 446–450 (2011).

12. Meulenbelt, I. *et al.* Identification of DIO2 as a new susceptibility locus for symptomatic

osteoarthritis. *Hum. Mol. Genet.* **17,** 1867–1875 (2008).

13. Meulenbelt, I. *et al.* Meta-analyses of genes modulating intracellular T3 bio-availability

reveal a possible role for the DIO3 gene in osteoarthritis susceptibility. *Ann. Rheum. Dis.* **70,**

164–167 (2011).

14. Styrkarsdottir, U. *et al.* Severe osteoarthritis of the hand associates with common variants

within the ALDH1A2 gene and with rare variants at 1p31. *Nat. Genet.* **46,** 498–502 (2014).

15. Evangelou, E. *et al.* The DOT1L rs12982744 polymorphism is associated with osteoarthritis

of the hip with genome-wide statistical significance in males. *Ann. Rheum. Dis.* **72,** 1264–

1265 (2013).

16. Evangelou, E. *et al.* A meta-analysis of genome-wide association studies identifies novel

variants associated with osteoarthritis of the hip. *Ann. Rheum. Dis.* **73,** 2130–2136 (2014).

17. Miyamoto, Y. *et al.* A functional polymorphism in the 5' UTR of GDF5 is associated with

susceptibility to osteoarthritis. *Nat. Genet.* **39,** 529–533 (2007).

18. den Hollander, W. *et al.* Genome-wide association and functional studies identify a role for

matrix Gla protein in osteoarthritis of the hand. *Ann. Rheum. Dis.* **76,** 2046–2053 (2017).

19. Panoutsopoulou, K. *et al.* Radiographic endophenotyping in hip osteoarthritis improves the

precision of genetic association analysis. *Ann. Rheum. Dis.* **76,** 1199–1206 (2017).

20. Castaño-Betancourt, M. C. *et al.* Novel Genetic Variants for Cartilage Thickness and Hip

Osteoarthritis. *PLoS Genet.* **12,** e1006260 (2016).

21. Yau, M. S. *et al.* Genome-Wide Association Study of Radiographic Knee Osteoarthritis in

North American Caucasians. *Arthritis Rheumatol* **69,** 343–351 (2017).

22. Evans, D. S. *et al.* Genome-wide association and functional studies identify a role for

IGFBP3 in hip osteoarthritis. *Ann. Rheum. Dis.* **74,** 1861–1867 (2015).

23. Valdes, A. M. *et al.* Genetic variation in the SMAD3 gene is associated with hip and knee

osteoarthritis. *Arthritis Rheum.* **62,** 2347–2352 (2010).

24. Rodriguez-Fontenla, C. *et al.* Assessment of osteoarthritis candidate genes in a meta-

analysis of nine genome-wide association studies. *Arthritis Rheumatol* **66,** 940–949 (2014).

25. Casalone, E. *et al.* A novel variant in GLIS3 is associated with osteoarthritis. *Ann. Rheum.*

Dis. **77,** 620–623 (2018).

26. Zengini, E. *et al.* Genome-wide analyses using UK Biobank data provide insights into the

genetic architecture of osteoarthritis. *Nat. Genet.* **50,** 549–558 (2018).

27. Cotney, J. *et al.* The evolution of lineage-specific regulatory activities in the human

embryonic limb. *Cell* **154,** 185–196 (2013).

28. Herlofsen, S. R. *et al.* Genome-wide map of quantified epigenetic changes during in vitro

chondrogenic differentiation of primary human mesenchymal stem cells. *BMC Genomics*

14, 105 (2013).

29. Raine, E. V. A., Wreglesworth, N., Dodd, A. W., Reynard, L. N. & Loughlin, J. Gene

expression analysis reveals HBP1 as a key target for the osteoarthritis susceptibility locus

that maps to chromosome 7q22. *Ann. Rheum. Dis.* **71,** 2020–2027 (2012).

30. Luyten, F. P., Tylzanowski, P. & Lories, R. J. Wnt signaling and osteoarthritis. *Bone* **44,**

522–527 (2009).

31. Berasi, S. P., Xiu, M., Yee, A. S. & Paulson, K. E. HBP1 repression of the p47phox gene:

cell cycle regulation via the NADPH oxidase. *Mol. Cell. Biol.* **24,** 3011–3024 (2004).

32. Scott, J. L. *et al.* Superoxide dismutase downregulation in osteoarthritis progression and

end-stage disease. *Ann. Rheum. Dis.* **69,** 1502–1510 (2010).

33. Liu, Y. *et al.* Chromatin accessibility landscape of articular knee cartilage reveals aberrant

enhancer regulation in osteoarthritis. *bioRxiv* 274043 (2018). doi:10.1101/274043

34. Thierry-Mieg, D., Thierry-Mieg, J. & NCBI/NLM/NIH. AceView: Gene:HBP1, a

comprehensive annotation of human, mouse and worm genes with mRNAs or

ESTsAceView. Available at:

https://www.ncbi.nlm.nih.gov/IEB/Research/Acembly/av.cgi?db=36a&c=Gene&l=HBP1.

(Accessed: 9th June 2018)

35. Gee, F., Rushton, M. D., Loughlin, J. & Reynard, L. N. Correlation of the osteoarthritis

susceptibility variants that map to chromosome 20q13 with an expression quantitative trait

locus operating on NCOA3 and with functional variation at the polymorphism rs116855380.

Arthritis & Rheumatology **67,** 2923–2932 (2015).

36. Shepherd, C. *et al.* Functional characterisation of the osteoarthritis genetic risk residing at

ALDH1A2 identifies rs12915901 as a key target variant. *Arthritis Rheumatol* (2018).

doi:10.1002/art.40545

37. Wang, X., Hou, J., Quedenau, C. & Chen, W. Pervasive isoform☐specific translational

regulation via alternative transcription start sites in mammals. *Mol. Syst. Biol.* **12,** 875

(2016).

38. Reyes, A. & Huber, W. Alternative start and termination sites of transcription drive most

transcript isoform differences across human tissues. *Nucleic Acids Res.* **46,** 582–592 (2018).

39. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

arXiv [q-bio.GN] (2013).

40. Southam, L. *et al.* An SNP in the 5'-UTR of GDF5 is associated with osteoarthritis

susceptibility in Europeans and with in vivo differences in allelic expression in articular

cartilage. *Hum. Mol. Genet.* **16,** 2226–2232 (2007).

41. Egli, R. J. *et al.* Functional analysis of the osteoarthritis susceptibility-associated GDF5

regulatory polymorphism. *Arthritis Rheum.* **60,** 2055–2064 (2009).