Constitutively active RAS in *S. pombe* causes persistent Cdc42 signalling but only transient MAPK activation

Emma J. Kelsall[1,8], Ábel Vértesy[2,6,8], Kees Straatman[3], Mishal Tariq[1], Raquel Gadea[1,4], Chandni Parmar[1], Gabriele Schreiber[2], Shubhchintan Randhawa[1,7], Cyril Dominguez[1,5], Edda Klipp[2]* and Kayoko Tanaka[1,9]*

Department of Molecular and Cell Biology, University of Leicester

Theoretical Biophysics, Institute of Biology, Humboldt-Universität zu Berlin

Centre for Core Biotechnology Services, University of Leicester

Marco Formacion Profesional, Calle del Conde de Aranda 7-9, 50004 Zaragoza, Spain

Leicester Institute of Structural and Chemical Biology, University of Leicester

Present address: Hubrecht Institute-KNAW (Royal Netherlands Academy of Arts and Sciences) and University Medical Center, 3584 CT Utrecht, The Netherlands

Present address: Sanjay Gandhi Post Graduate Institute of Medical Sciences, Department of Molecular Medicine & Biotechnology, Lucknow, India 226014

These authors contributed equally

Lead Contact

*Correspondence: kt96@le.ac.uk (K.T.), edda.klipp@rz.hu-berlin.de (E.K.)

Highlights

(1) Constitutive Ras1.GV causes over-activation of Cdc42 in *S. pombe* pheromone signaling

(2) Ras1.GV induces an acute but only transient MAPK^{Spk1} activation

(3) The RAS effector pathways MAPK^{Spk1} and Cdc42 compete with each other for active Ras1

(4) Predictive modelling explains MAPK^{Spk1} activation dynamics in 24 signaling-mutants

eTOC Blurb

S. pombe Ras1 activates the MAPK^{Spk1} and Cdc42 pathways. Kelsall et al. report that the constitutively active Ras1.G17V mutation, which causes morphological anomalies, induces prolonged Cdc42 activation and MAPK^{Spk1} activation followed by attenuation. Mathematical modelling shows competition between the MAPK^{Spk1} and Cdc42 pathways for active Ras1.

Summary

The small GTPase RAS is a signalling hub. Oncogenic RAS mutations are assumed to over-activate all of the downstream pathways. We tested this assumption in fission yeast, where, RAS-mediated pheromone signalling (PS) activates the MAPK^{Spk1} and Cdc42 pathways. Unexpectedly, we found that constitutively active Ras1.G17V induced acute transient MAPK^{Spk1} activation, whilst Cdc42 activation persisted. Acute transient MAPK^{Spk1} activation was also seen in the deletion mutant of Cdc42-GEF^{Scd1}, a Cdc42 activator. We have built a mathematical model using PS negative-feedback circuits and competition between the two Ras1 effectors, MAPKKK^{Byr2} and Cdc42-GEF^{Scd1}. The model robustly predicted the MAPK^{Spk1} activation dynamics of an additional 21 PS mutants. Supporting the model, we show that a recombinant Cdc42-GEF^{Scd1} fragment competes with MAPKKK^{Byr2} for Ras1 binding. Our study has established a concept that constitutive RAS yields biased Cdc42/Rac over-activation, providing a strong rationale to interfere with this process to target oncogenic RAS in humans.

Key words

Ras, MAPK, Cdc42/Rac, yeast pheromone signalling

Introduction

Proto-oncogene Ras GTPase family members are widely conserved and play pivotal roles in cell growth, differentiation and apoptosis (Cox and Der, 2010). The physiological impact of Ras mutations is highlighted in the resultant tumorigenesis and developmental disorders (Prior et al., 2012; Schubbert et al., 2007). More than 99% of identified oncogenic RAS mutations occur at codons 12, 13 and 61 of human Ras isoforms (Prior et al., 2012) and impair efficient GTP hydrolysis (Trahey and McCormick, 1987). This results in accumulation of GTP-bound Ras, which is generally considered to cause constitutive activation of the downstream effector pathways, such as ERK and PI3K signalling pathways (Lin et al., 1998; Zhu et al., 1998). Interestingly, mouse embryonic fibroblasts (MEFs) derived from the *K-ras^{G12D}* mouse model show neither an increased basal level of active ERK and Akt nor constitutive activation of ERK and Akt upon growth factor stimulation even though the *K-ras^{G12D}* MEFs showed enhanced proliferation and partial transformation (Tuveson et al., 2004). This observation indicates that not all effector pathways become constitutively activated by oncogenic Ras, presumably because of an efficient negative feedback loop. Meanwhile, it is reasonable to expect that oncogenic RAS over-activates some of its effector pathways to initiate tumorigenesis. Small GTPases, including Cdc42 and Rac, may be such effector pathways, as they are required in oncogenic-RAS-driven tumorigenesis (Malliri et al., 2002; Qiu et al., 1997; Stengel and Zheng, 2012).

We wished to understand how multiple Ras effector pathways respond to the Ras-triggered signals in a physiological setting. We employed the model organism fission yeast. In this organism, a unique Ras homologue, Ras1, plays a key role in pheromone signalling to cause mating of haploid cells and sporulation in diploid cells (Yamamoto, 1996)(Fig. 1A, B). Upon nutritional starvation, cells of opposite mating types (*h$^+$* and *h$^-$*) exchange mating pheromones. Gpa1, the α-subunit of the pheromone receptor-coupled G-protein, relays the pheromone signal into the cell (Obara et al., 1991). Activated Gpa1 together with Ras1 then activate MAPK cascade consisting of Byr2 (MAPKKK), Byr1 (MAPKK) and Spk1 (MAPK) (Fukui et al., 1986; Masuda et al., 1995; Nadin-Davis and Nasim, 1988; Nadin-Davis et al., 1986a; Nadin-Davis et al., 1986b; Wang et al., 1991; Xu et al., 1994). An intriguing observation is that the *ras1.G17V* mutant, an equivalent of mammalian *ras.G12V* mutant prevalent in cancer, produces an excessively elongated shmoo, or a conjugation tube, upon exposure to the mating pheromone (Nadin-Davis et al., 1986a). This "elongated" *ras1.G17V* phenotype has been interpretted that

Ras1 might be directly responsible for amplifying the pheromone signal at an intracellular level

(Yamamoto, 1996).

Ras1 also regulates cell morphology during vegetative growth; whilst deletion of either *gpa1*,

MAPKKK^{byr2}, *MAPKK^{byr1}* or *MAPK^{spk1}* does not result in any obvious phenotypes during vegetative

cell growth (Obara et al., 1991; Sipiczki, 1988; Toda et al., 1991), *ras1Δ* cells lose the typical rod-

shape morphology of fission yeast to become rounded (Fukui et al., 1989; Nadin-Davis et al.,

1986a). Studies based on recombinant protein assays and yeast-2-hybrid analysis demonstrated

that Ras1 interacts with both MAPKKK^{Byr2} and Scd1, a GDP-GTP exchange factor (GEF) for Cdc42,

which regulates the actin cytoskeleton and cell morphology (Chang et al., 1994; Gronwald et al.,

2001; Tu et al., 1997). These observations suggest that Ras1 simultaneously regulates both the

pheromone MAPK^{Spk1} and the Cdc42 pathways at the cell membrane (Weston et al., 2013).

Indeed, a dynamic "Cdc42 zone" at the cell cortex prior to mating has been observed (Merlini et

al., 2013) and Ras1 and MAPK^{Spk1} cascade components are found there and are involved in the

process (Dudin et al., 2016; Merlini et al., 2013; Merlini et al., 2016; Merlini et al., 2018; Weston

et al., 2013). However, it is not yet understood how Ras1 interplay between the two pathways.

By establishing conditions to induce highly synchronous mating of fission yeast cells, for the first

time we were able to follow MAPK^{Spk1} activation during the physiological mating process in

wildtype and in mutants showing various mating phenotypes. This comprehensive set of

quantitative measurements allowed us to build a mathematical model of the Ras-mediated

pheromone signalling. Our model can serve as a prototype of a branched Ras-mediated

signalling pathway, demonstrating a competition of downstream pathways for a common

upstream activator, Ras1. The model also highlights the physiological importance of the

bipartite activation of MAPKKK^{Byr2}: a Ras1-dependent and a Ras1-independent mechanism, the

latter of which employs the adaptor protein Ste4 (Barr et al., 1996; Okazaki et al., 1991). The

adaptor^{Ste4} can be targeted to downregulate MAPK^{Spk1} even in the presence of *ras1.G17V*

mutation. Finally, our study reveals the crucial role played by Cdc42 in the *ras1.G17V* mutant

causing the *ras1.G17V* phenotype.

Results

(1) A highly synchronous mating assay allows the precise measurement of MAPK^{Spk1} activity

To directly measure fission yeast pheromone signalling, we quantitated MAPK^{Spk1} phosphorylation throughout the mating process (Fig. S1, Fig. 1A and B). We established a protocol to induce highly synchronous mating and employed cells where the endogenous MAPK^{Spk1} is tagged with GFP-2xFLAG. Under these conditions, homothallic h^{90} cells started to mate 7-hours after induction of mating (Fig. 1C, grey line). Phosphorylated (active) MAPK^{Spk1} (**pp**MAPK^{Spk1}) levels were quantitated as described in Materials and Methods. The **pp**MAPK^{Spk1} signal was first detected three hours after induction of mating and reached its peak at about seven hours, when cell fusion was also initially observed (Fig. 1C, blue line). The **pp**MAPK^{Spk1} then gradually decreased to a non-zero level as meiosis continued towards sporulation. The observation established that the MAPK^{Spk1}-GFP activation occurs as the mating process progresses and declines before sporulation, around 15 hours post induction. It was also noted that the total MAPK^{Spk1}-GFP was essentially not expressed during the vegetative cycle, but it was promptly induced by nitrogen starvation (Supplementary Fig. S2A and S2B). The *mapk^{spk1}* gene is a known target of the transcription factor Ste11 (Mata and Bahler, 2006), which itself is activated (phosphorylated) by MAPK^{Spk1} (Kjaerulff et al., 2005). This positive feedback loop likely facilitates a swift increase of MAPK^{Spk1} expression upon nitrogen starvation. We found that MAPK^{Spk1}-GFP localised to both the cytosol and the nucleus, with some nuclear accumulation, before it gradually disappeared as the mating process came to the end (Fig. 1D). Interestingly, transient foci of GFP signals were also found at the cell cortex, as has been reported for MAPKK^{Byr1}, the activator of MAPK^{Spk1} (Dudin et al., 2016) (Fig. 1D, yellow arrows).

(2) Constitutively active MAPKK$^{Byr1.DD}$ mutant causes constitutive activation of MAPK^{Spk1}

Activation of MAPKK family kinases is mediated by dual phosphorylation of conserved Ser/Thr residues (Zheng and Guan, 1994). These correspond to serine 214 and threonine 218 of the MAPKK^{Byr1}. A MAPKK^{Byr1} mutant termed MAPKK$^{Byr1.DD}$, which carries aspartic acid substitution at these sites, was expected to act as a constitutively active MAPKK (Ozoe et al., 2002). We introduced the *MAPKK$^{byr1.DD}$* mutation at its chromosome locus and measured the activation profile of its target, MAPK^{Spk1} during the mating process. The increase of the **pp**MAPK^{Spk1} signal in the *MAPKK$^{byr1.DD}$* mutant strain was delayed compared to the wildtype strain and the

ppMAPK^{Spk1} accumulated at a slower rate (Fig. 1E, light green line). However, the level of

ppMAPK^{Spk1} remained high after reaching its highest intensity at around 16 hours after

induction, resulting in a constitutive phosphorylation of MAPK^{Spk1} (Fig. 1E and Supplementary

Fig. S2C and S2D). This result highlights that the suggested de-phosphorylation of **pp**MAPK^{Spk1}

by phosphatases Pmp1 and Pyp1 (Didmon et al., 2002) is not efficient in downregulating the

pheromone signalling in the presence of MAPKK$^{Byr1.DD}$. The MAPKK$^{Byr1.DD}$ strain is also less

competent in finding the partner mating cell than cells with wildtype MAPKK^{Byr1}. We also

confirmed the intriguing *"fus"* (fusion deficient) phenotype of cells expressing MAPKK$^{Byr1.DD}$ from

its native promoter (Fig.1F) (Dudin et al., 2016; Ozoe et al., 2002). These cells find their partners

and pair up successfully but fail to fuse with each other, resulting in *fus* phenotype. Consistent

with the observed slower increase in **pp**MAPK^{Spk1} (Fig.1E) the nuclear localisation of MAPK^{Spk1}

was also delayed compared to Wildtype cells (Fig 1F). A strong MAPK^{Spk1}-GFP signal was then

observed in the nuclei of the paired *fus* cells and the nuclear MAPK^{Spk1}-GFP signal persists even

24 hours after the induction of mating (Fig. 1F). Interestingly, the projection tips of the paring

cells often show increased MAPK^{Spk1}-GFP signal (Fig. 1F, yellow arrows). These results conclude

that MAPK^{Spk1} activation is highly influenced by the MAPKK^{Byr1} status and phosphatases directly

regulating MAPK^{Spk1} cannot counteract MAPKK$^{Byr1.DD}$.

(3) The *ras1.G17V* mutation causes acute transient MAPK^{Spk1} activation

The fission yeast equivalent of human oncogenic *ras.G12V* is *ras1.G17V,* which induces an

excessively elongated shmoo, and the cells fail to recognize a partner and become sterile (Fukui

et al., 1986; Nadin-Davis et al., 1986a). The "elongated shmoo" phenotype was interpreted as

an excess activation of the downstream pathway(s) of Ras1, leading to a prediction that the

ras1.G17V causes over-activation of MAPK^{Spk1} (Weston et al., 2013; Yamamoto, 1996). However,

no direct evidence has been provided.

Quantitation of the **pp**MAPK^{Spk1}-GFP in the *ras1.G17V* mutant showed an acute increase of the

ppMAPK^{Spk1}-GFP upon induction of mating (Fig. 1G, Green line). However, the signal intensity

declined gradually and by 16 hours after induction the level was comparable to wildtype cells,

indicating that down-regulation of **pp**MAPK$^{Spk1-GFP}$ is effective, unlike in *MAPKK$^{byr1.DD}$* mutant.

Correspondingly, the cellular MAPK^{Spk1}-GFP signal also declined 24 hours after induction of

mating (Fig. 1H). Collectively, these observations indicate that the down-regulation mechanism

for **pp**MAPK^{Spk1} is robust and resistant to Ras1.G17V. It was also noted that the peak intensity of

the **pp**MAPK^{Spk1} was somewhat lower and the rate of signal reduction was slightly decreased

compared to wildtype cells (Fig. 1G).

(4) The elongated *ras1.G17V* shmoos develop with a minimum level of MAPK^{Spk1}: neither

amplitude nor duration of ppMAPK^{Spk1} signal influences the *ras1.G17V* phenotype

Having observed that *MAPKK$^{byr1.DD}$* and *ras1.G17V* show different MAPK^{Spk1} activation profiles

(sustained vs transient) and different morphological phenotypes ("*fus*" vs elongated), we

examined a link between the MAPK^{Spk1} activation profiles and the cell morphology. The

MAPK^{Spk1} phosphorylation profile of the cells harbouring both *ras1.G17V* and *MAPKK$^{byr1.DD}$*

mutations (*ras1.G17V MAPKK$^{byr1.DD}$* double mutant) was comparable to the cells harbouring

MAPKK$^{byr1.DD}$ mutation only; it showed a slow increase of the **pp**MAPK^{Spk1}-GFP level, which

reached the plateau at about 16 hours after induction of mating (Fig. 2A, red line). The nuclear

ppMAPK^{Spk1}-GFP signal in the *ras1.G17V MAPKK$^{byr1.DD}$* double mutant was also present 24 hours

after induction of mating, unlike the *ras1.G17V* single mutant cells, confirming that the

ras1.G17V MAPKK$^{byr1.DD}$ double mutant cells retained a high **pp**MAPK^{Spk1} level. Thus, in terms of

the activation status of MAPK^{Spk1}, *MAPKK$^{byr1.DD}$* is epistatic to *ras1.G17V*.

In terms of cell morphology, the *ras1.G17V MAPKK$^{byr1.DD}$* double mutant cells showed the

"elongated" *ras1.G17V* phenotype (Fig. 2B), but not the "paired" *MAPKK$^{byr1.DD}$* phenotype.

Therefore, *ras1.G17V* was epistatic to *MAPKK$^{byr1.DD}$* in terms of the elongated shmoo

morphology.

Interestingly, obvious shmoo formation in the *ras1.G17V MAPKK$^{byr1.DD}$* double mutant was first

noticed 16 hours after induction of mating, much later than the *ras1.G17V* single mutant, but as

a similar timing as the *MAPKK$^{byr1.DD}$* single mutant (Fig. 2B). Given the slow increase of the

ppMAPK^{Spk1} signal in mutants harbouring the *MAPKK$^{byr1.DD}$* mutation, we predicted that the

delayed appearance of the *ras1.G17V* shmoo meant that the *ras1.G17V* shmoo formation still

requires a certain level of MAPK^{Spk1} activity. Indeed, when *MAPKK^{byr1}* was deleted in the

ras1.G17V mutant, not only was the MAPK^{Spk1} activation and nuclear MAPK^{Spk1}-GFP signal

abolished (Fig. S1F), but also shmoo formation was abrogated as in the *MAPKK^{byr1}Δ* single

mutant (Fig. 2C). Based on these observations, we concluded that the cell morphology is

determined by the molecular status of Ras1, and not by the MAPK^{Spk1} activation profile. Yet, the

ras1.G17V phenotype still requires MAPK^{Spk1} activity, which determines the timing of shmoo

formation.

(5) Cdc42 is required for the shmoo formation but not for the MAPK^{Spk1} activation

During the vegetative cycle, *ras1Δ* cells show spherical cell morphology (Fukui et al., 1986; Nadin-Davis et al., 1986a) and polarised localisation of active Cdc42 is compromised (Kelly and Nurse, 2011), (Fig. 4C, D), indicating that Ras1 is involved in Cdc42 activation. The GTP-loaded Cdc42 is then predicted to activate the downstream Ste20-like kinase, Pak1/Shk1 (Endo et al., 2003; Marcus et al., 1995; Ottilie et al., 1995; Verde et al., 1995), resulting in actin reorganisation and shmoo formation under mating conditions (Bendezu and Martin, 2013; Merlini et al., 2016).

To confirm that Cdc42 acts downstream of Ras1, we generated a double mutant strain harbouring *ras1.G17V* and deletion of *scd1*, encoding a GDP-GTP exchanging factor for Cdc42 (Cdc42-GEF^{Scd1}). The *ras1.G17V* elongated shmoo phenotype was lost in the double mutant and instead, the cells showed a mating-deficient phenotype similar to the *cdc42-GEF^{scd1}Δ* single mutant (Fig 3A). The result supports the model that Cdc42 acts downstream of Ras1 to cause morphological changes.

Intriguingly, in the strains harbouring the *cdc42-GEF^{scd1}Δ* mutation, a nuclear MAPK^{Spk1}-GFP signal appeared (Fig. 3A), indicating that activation of MAPK^{Spk1} may not be impaired by lack of active Cdc42. This was unexpected because in a previous study, it was predicted that activation of Cdc42 contributes to activation of MAPKKK^{Byr2} (Tu et al., 1997). To clarify this issue, we measured **pp**MAPK^{Spk1} in the *cdc42-GEF^{scd1}Δ* mutant. Strikingly, in these cells MAPK^{Spk1} activation occurred with a reproducible advancement of the initial activation timing compared to the wildtype cells (Fig. 3B). The result shows that MAPK^{Spk1} activation does not require Cdc42 activity. Additionally, the faster activation of MAPK^{Spk1} raises the interesting possibility that two Ras1 effectors, MAPKKK^{Byr2} and Cdc42-GEF^{Scd1}, are competing with each other for activated Ras1, thus, lack of Cdc42-GEF^{Scd1} results in an advanced MAPK^{Spk1} activation (modelled in Fig. 7A).

Substantial MAPK^{Spk1} activation in the *cdc42-GEF^{scd1}Δ* mutant means that the mating deficiency of this mutant is unlikely to be the result of the lack of MAPK^{Spk1} activation. Indeed, introduction of *MAPKK$^{byr1.DD}$* to the *cdc42-GEF^{scd1}Δ* mutant did not restore the mating deficient phenotype, even though nuclear MAPK^{Spk1}-GFP highly accumulated (Fig. 3C). The result shows that active Cdc42 function is absolutely required for the mating process regardless of the MAPK^{Spk1}

activation status. Taken together with the essential role of MAPK^{Spk1}, we concluded that the

mating pheromone signalling feeds into at least two pathways, MAPK^{Spk1} and Cdc42.

(6) Ras1 activates two effector pathways, MAPK^{Spk1} and Cdc42.

In order to further clarify the role of Ras1 we examined the MAPK^{Spk1} activation status and cell

morphology in the following four strains: *ras1Δ* mutant, *ras1Δ MAPKK$^{byr1.DD}$* double mutant,

MAPKKK^{byr2}Δ mutant and *MAPKKK^{byr2}Δ MAPKK$^{byr1.DD}$* double mutant. As mentioned earlier,

deletion of *ras1* causes cells to show a round morphology (Fig. 3F), with reduced cortical signal

of CRIB-GFP, an indicator of the active GTP-bound form of Cdc42, showing that Cdc42 activation

is compromised (Fig.4C, D). *ras1* deletion also causes substantial reduction, but not complete

elimination, of the **pp**MAPK^{Spk1} (Fig. 3D, red line, and Supplementary Fig. S3A); thus, Ras1 plays

an important role in activating both Cdc42 and MAPK^{Spk1} pathways. Introduction of the

MAPKK$^{byr1.DD}$ mutation into the *ras1Δ* mutant cells induces the constitutive **pp**MAPK^{Spk1} (Fig. 3D,

green line and Supplementary Fig. S3A) but does not affect the round cell morphology and cells

remain sterile (Fig. 3F, the 2nd left panel. Note the accumulating MAPK^{Spk1}-GFP at 16 hours after

induction of mating), as was the case for the *Cdc42-GEF^{Scd1}Δ MAPKK$^{byr1.DD}$* double mutant (Fig.

3C).

In a striking contrast, the sterile phenotype of the *MAPKKK^{byr2}Δ*, associated with complete lack

of shmoo formation (Fig. 3F, the 2nd right panel), was converted to the "*fus*" phenotype, when

combined with the *MAPKK$^{byr1.DD}$* mutation (Fig. 3F, the far right panel). As expected, the

MAPKKK^{byr2}Δ MAPKK$^{byr1.DD}$ double mutant shows MAPK^{Spk1} constitutive activation (Fig. 3E and

Supplementary Fig. S3B). Thus, unlike the cases of *scd1Δ* or *ras1Δ*, lack of *MAPKKK^{byr2}* can be

bypassed by constitutive activation of MAPK^{Spk1}, indicating that the sole role of MAPKKK^{Byr2} is to

activate the MAPK^{Spk1} unlike its upstream activator, Ras1, which also activates Cdc42 pathway (a

model presented in Fig. 7A).

(7) Ras1.G17V causes accumulation of Cdc42-GTP at the cell cortex.

Having observed a relatively mild influence of Ras1.G17V towards the MAPK^{Spk1} activation, we

next examined whether the Cdc42 pathway was affected by the *ras1.G17V* mutation. We

visualized the active GTP-bound form of Cdc42 (Cdc42^{GTP}) using CRIB-GFP that specifically binds

to Cdc42^{GTP} (Tatebe et al., 2008). As previously observed, dynamic foci of CRIB-GFP appeared on

the cell cortex upon induction of mating (Bendezu and Martin, 2013)(Fig. 4A and B). In our

experimental condition, more than 80% of wildtype cells showed the cortical CRIB-GFP signal at
4.5 hours after induction of mating (Fig. 4B). The cortical CRIB-GFP foci became concentrated at
the site of mating and quickly disappeared once cells fused successfully to form zygotes (Fig. 4A
and B). In striking contrast, in the *ras1.G17V* mutant cells, the cortical CRIB-GFP signal persisted,
often at the elongated tip end of the cells, even 12.5 hours after induction of mating (Fig. 4A
and B). The signal could still be seen in about 40% of the cells 22.5 hours after induction of
mating (Fig. 4A and B). The result shows that the Cdc42 pathway is excessively activated in the
ras1.G17V mutant and the localisation pattern of Cdc42^{GTP} indicates that the signature
"elongated" *ras1.G17V* morphological phenotype is caused by deregulation of the Cdc42
pathway.
Ras1-mediated Cdc42 pathway activation has been also indicated during the vegetative growth
where the *ras1Δ* mutant shows a round cell morphology (Chang et al., 1994; Kelly and Nurse,
2011). However, unlike during the mating process, the *ras1.G17V* mutation does not cause an
obvious morphological phenotype during the vegetative growth. We predicted that, during the
vegetative growth, rigorous negative regulation occurs for Cdc42 by GTPase activation
protein(s) (GAPs), such as Rga4 (Das et al., 2007; Kelly and Nurse, 2011; Tatebe et al., 2008) to
counteract the effect of *ras1.G17V*. To examine this possibility, we compared Cdc42 activation
status of vegetatively growing wildtype, *ras1Δ*, *ras1.G17V*, *rga4Δ*, and *rga4Δ ras1.G17V* double
mutant cells (Fig. 4C and D). As previously described, CRIB-GFP showed a clearly polarized signal
at the growing cell tips in the wildtype strain (Tatebe et al., 2008)(Fig. 4C and D). In the *ras1Δ*
mutant, the cells were round and CRIB-GFP signal on the cell cortex had largely disappeared as
was seen in the *cdc42-GEF^{scd1}Δ* mutant (Kelly and Nurse, 2011). In contrast, the cortical CRIB-
GFP signal was clearly increased in the *ras1.G17V* single mutant although the cell morphology
appeared largely similar to the wildtype cells (Fig. 4C, D). These results indicate a direct
involvement of Ras1 in activating Cdc42. Meanwhile, the *rga4Δ* mutant cells showed slight
alterations to the cell morphology, accompanied with less polarized distribution of cortical CRIB-
GFP signal, as has been reported (Fig. 4C and D) (Das et al., 2007; Kelly and Nurse, 2011; Tatebe
et al., 2008). Strikingly, the *rga4Δ ras1.G17V* double mutant showed a clear morphological
alteration (big round cells) and the strongest cortical CRIB-GFP signal among all the mutants
examined (Fig. 4C and D). The result fits well with our hypothesis that Ras1.G17V is activating
Cdc42 even during vegetative growth, but the overall effect of Ras1.G17V is counteracted by the
Cdc42-GAP, Rga4.
302

(8) Ras1 and an adaptor protein Ste4 are both necessary to fully activate MAPKKK[Byr2]

Although Ras1 clearly plays the major role to activate MAPK[Spk1], a marginal, but detectable level

of **pp**MAPK[Spk1] was still induced in the *ras1Δ* mutant (Fig. 3D and Supplementary Fig. S3A),

indicating that there is a Ras1-independent mechanism to activate MAPK[Spk1]. Previous studies

proposed an adaptor protein, Ste4, to be involved in the activation of MAPKKK[Byr2](Barr et al.,

1996; Okazaki et al., 1991; Ramachander et al., 2002; Tu et al., 1997). We therefore examined

whether Ste4 is required for MAPK[Spk1] activation.

In contrast to the *ras1Δ* mutant, we detected virtually no MAPK[Spk1] phosphorylation in the

ste4Δ mutant (Fig. 5A and Supplementary Figure S3C), indicating that the adaptor[Ste4] is a

prerequisite for the MAPK[Spk1] activation and the **pp**MAPK[Spk1] signal observed in the *ras1Δ*

mutant is dependent on Ste4 function. Introduction of *ras1.G17V* mutation neither restored the

MAPK[Spk1] activation nor mating (Fig. 5A, B), thus, activation of Ras1 cannot take over Ste4

function. In a striking contrast, the *ste4Δ MAPKK[byr1.DD]* double mutant showed the *"fus"*

phenotype as the *MAPKK[byr1.DD]* single mutant cells and induced constitutive MAPK[Spk1] activation,

indicating that Ste4 is solely required for MAPK[Spk1] activation (Fig. 5A, B).

Taken together, MAPKKK[Byr2] is activated through a mechanism involving both Ras1 and Ste4, but

Ste4 only conveys the signal towards the MAPK[Spk1], while Ras1 also activates Cdc42.

(9) Ste6, a Ras1 GTP-GDP exchange factor, contributes to both the MAPK[Spk1] and the Cdc42
322

pathway activation
323

There are two GDP-GTP exchange factors (GEFs) identified for Ras1: Ste6 and Efc25 (Hughes et

al., 1990; Tratner et al., 1997). As to the functional differences, Ste6 is essential for mating but is

dispensable during the vegetative cycle whilst Efc25 is dispensable for mating but is required to

maintain the cell morphology during the vegetative growth (Hughes et al., 1990; Tratner et al.,

1997). There has been an interesting proposition that Ste6 may specifically help Ras1 to activate

the MAPK[Spk1] pathway, but not the Cdc42 pathway, whilst Efc25 specifically facilitates Ras1 to

activate the Cdc42 pathway (Papadaki et al., 2002). We examined this hypothesis by monitoring

the MAPK[Spk1] activation status and conducting genetic epistasis analysis of *ras1.G17V* and

MAPKK[byr1.DD] in the *ste6Δ* mutant.

In *ste6Δ* cells, MAPK[Spk1] phosphorylation was found somewhat reduced but occurred at a clearly

detectable level. The signal increased when the *ras1.G17V* mutation was introduced (Fig. 5C and

Supplementary Figure S3D). When *ste6Δ and MAPKK^byr1.DD* were combined, MAPK^Spk1 signalling

recapitulated the *MAPKK^byr1.DD* activation profile (Fig. 5C and Supplementary Figure S3D).

Nonetheless, the "pheromone-insensitive sterile" morphology of *ste6Δ* was only rescued by

ras1.G17V, as previously reported, exhibiting the "elongated" phenotype (Hughes et al., 1990),

but not by MAPKK^byr1.DD (Fig. 5D). The result indicates that, unlike *ste4Δ* mutant, the mating

deficiency of *ste6Δ* is not caused by mere lack of MAPK^Spk1 activation but by lack of Ras1

activation. We concluded that Ste6 functions to activate Ras1, which then activates *both* the

MAPK and Cdc42 pathways in response to pheromone signalling.

(10) Activation mutant of Gpa1 mimics the full pheromone signalling

In order to generate an integrated prototype Ras signalling model, we further investigated the

upstream signal input machinery. Previous studies showed that Gpa1 plays the primary role in

pheromone signalling (Obara et al., 1991). In agreement, in the *gpa1Δ* mutant we detected no

MAPK^Spk1 activation nor nuclear accumulation of MAPK^Spk1-GFP (Fig. 6A, B and Supplementary

Fig. S4A). Introducing the *ras1.G17V* to the *gpa1Δ* strain did not rescue the complete lack of

MAPK^Spk1 activation nor did it induce a shmoo-like morphological change (Fig. 6A, B and

Supplementary Fig. S4A), supporting our earlier observation that a Ras1-independent

mechanism, involving Ste4 is essential for MAPK^Spk1 activation. Meanwhile, introducing the

MAPKK^byr1.DD mutation caused the constitutive activation of MAPK^Spk1 (Fig. 6A and

Supplementary Fig. S4A), but cells showed no morphological change (Fig. 6B). When both

ras1.G17V and *MAPKK^byr1.DD* mutations were introduced into the *gpa1Δ* strain, MAPK^Spk1 was

activated and a shmoo-like morphological change occurred (Fig. 6A, B and Supplementary Fig.

S4A). Therefore, activation of both of these two molecules is required and sufficient to mimic

the pheromone signalling.

To further confirm that the Gpa1 is a central component of pheromone signalling, we looked

into the MAPK^Spk1 activation status in the constitutively active *gpa1.QL* mutant, which exhibits a

"shmoo-like" morphological change in the heterothallic *h⁻* strain without the mating partner

(Obara et al., 1991).

Upon nitrogen starvation, the *h⁻ gpa1.QL* mutant strain showed morphological changes and a

strong MAPK^Spk1 activation (Fig. 6C red line, Fig. 6D and Supplementary Fig. S4B). This response

was largely dependent on the Ras1 function as the *h⁻ gpa1.QL ras1Δ* double mutant exhibited a

significantly reduced level of **pp**MAPK^Spk1 and a round cell morphology, two typical features of

ras1Δ cells (Fig. 6C, D and Supplementary Fig. S4B). On the other hand, the *h⁻ ras1.G17V* single

mutant showed a very low level of MAPK^{Spk1} activation, comparable to the one observed in the

h⁻ wildtype strain, with no apparent morphological alternation, confirming that sole activation

of Ras1 does not substitute the pheromone signalling.

The *h⁻ MAPKK$^{byr1.DD}$* mutant induced a strong constitutive MAPK^{Spk1} activation, confirming that

the MAPKK$^{Byr1.DD}$ molecule can activate MAPK^{Spk1} regardless of the pheromone signal input (Fig.

6C, light-blue line and Supplementary Fig. S4B). However, cell morphology was unchanged (Fig.

6D). Collectively, these results support the model where Gpa1 acts as the central transducer of

the pheromone signalling, which can be mimicked only if both MAPK^{Spk1} and Ras1 are activated.

(11) A holistic modelling framwork of MAPK^{Spk1} activation

Based on quantitative **pp**MAPK^{Spk1} measurements in wildtype and various mutant strains (Fig.

1C, E, G, and Fig. 3B), we constructed a mathematical model of the MAPK^{Spk1} signalling

dynamics. The aim of the model is to test whether a simple competition of the MAPK and the

Cdc42 pathways for a shared pool of active Ras^{GTP}, can explain why the *scd1Δ* strain shows the

similar **pp**MAPK^{Spk1} activation profile as the *ras1.G17V* strain.

We designed a reductionist model of 6 ordinary differential equations to represent key steps of

pheromone signalling (Fig. 7A, Supplementary Fig. S5 and materials and methods). Model

simulations and parameter estimations were performed in COPASI (Hoops et al., 2006) and

details of the modelling process are described in Materials and Methods. Each biochemical

process is referred as [L1]-[L10] as depicted in Fig. 7A. The signalling components were set to

interact without delay based on the observations that signalling components are localized in

close proximity (Fig. 1 and Fig. 4)(Dudin et al., 2016; Merlini et al., 2016; Merlini et al., 2018).

The framework of the modelling process is as follows: Genes encoding pheromones, receptors,

Gpa1, Ste4 and Ste6 are all known to be under regulation of Ste11, the master transcriptional

regulator for meiotic genes (Hughes et al., 1994; Mata and Bahler, 2006; Mata et al., 2002; Mata

et al., 2007; Sugimoto et al., 1991) and these components are grouped into the Pheromone

Sensing (PS) unit. During the vegetative growth, the PS unit is set to zero. Nitrogen starvation

activates Ste11 (Kjaerulff et al., 2007; Sugimoto et al., 1991), which induces the PS unit [L1]. The

PS unit activates MAPKKK^{Byr2} in a twofold manner: Directly by Ste4 [L3] and through Ras1

[L4](Fig. 7A). Activated MAPKKK^{Byr2} then triggers activation of MAPKK^{Byr1} [L5] that activates

MAPK[Spk1] [L6]. Since activated MAPK[Spk1] further activates Ste11 (Kjaerulff et al., 2005), MAPK[Spk1]

has a positive feedback loop on its own expression via Ste11 [L2,L7] (Fig. 7A).

As the pheromone signalling was found to induce a transient **pp**MAPK[Spk1] peak (Fig. 1C),

ppMAPK[Spk1] activity is ought to be regulated by a delayed downregulation. Because the

MAPKK[byr1.DD] mutant completely lacks downregulation (Fig. 1E), downregulation occurring

downstream of MAPKK[byr1] (e.g.: Pyp1 and Pmp1) were considered physiologically insignificant.

Meanwhile, Sxa2 (a serine carboxypeptidase against a mating pheromone P-factor) and Rgs1 (a

regulator of Gpa1), both of which are induced upon successful pheromone signalling (Imai and

Yamamoto, 1992; Mata and Bahler, 2006; Pereira and Jones, 2001; Watson et al., 1999),

receptor internalization (Hirota et al., 2001) and regulation of the *mapk[spk1]* transcript or other

components by antisense RNA (Bitton et al., 2011) fit well to the criteria for the negative

feedback. We represented all these potential negative feedbacks collectively as a single circuit,

[L8]. Importantly, this downregulation [L8] works unperturbed in the presence of Ras1.G17V, by

acting through the Ste4-dependent MAPKKK[Byr2] activation process (Fig. 1G, Fig. 7A).

Ras1-GTP activates both the MAPKKK[Byr2] and the Cdc42 pathways [L9] (Chang et al., 1994). As

opposed to previous expectations that active Cdc42[GTP] is required to activate MAPKKK[Byr2] (Tu et

al., 1997), deletion of Cdc42-GEF[Scd1] does not compromise MAPK[Spk1] activation (Fig. 3B) and

rather, it makes **pp**MAPK[Spk1] dynamics remarkably similar to that of the Ras1.G17V strain (Fig.

3B and Fig. 1G, plotted together in Fig. 7B): in both cases, **pp**MAPK[Spk1] peaks earlier than the

wildtype case. At a molecular level, reactions depleting the Ras1[GTP] pool are compromised in

both strains. Therefore, to explain these observations, we hypothesized that MAPK[Spk1] and

Cdc42 pathways are competing for the common Ras1[GTP] pool. In this manner, one of the

pathways can modulate the other by changing the amount of unbound (available) Ras1[GTP].

In support of this prediction, we showed that binding of recombinant Ras1.G17V[GTP] to a

MAPKKK[Byr2] fragment was reduced by the presence of a Cdc42-GEF[Scd1] fragment *in vitro* (Fig. 7C,

D). In this assay, bacterially expressed GST tagged fragments of both MAPKKK[Byr2] (65-180) and

Cdc42-GEF[Scd1] (760-872) showed a specific binding towards the GTP-loaded Ras1.G17V (1-172)

(Fig. 7C). However, the binding of Ras1.G17V[GTP] to MAPKKK[Byr2] was substantially decreased

when the Cdc42-GEF[Scd1] fragment was added (Fig. 7D). The result likely refects the intrinsic

biochemical competitive nature of MAPKKK[Byr2] and Cdc42-GEF[Scd1] for Ras1 binding. As we show

below, this simple hypothesis successfully describes pheromone signalling mutants tested in this

study, suggesting that no unproven cross links are necessary to reproduce the *in vivo*

observations.

An intriguing common feature observed in both Cdc42-GEF^{scd1}Δ and *ras1.G17V* mutants is that

ppMAPK^{Spk1} peaks not only at an earlier time point, but also with a *lower amplitude* as

compared to the wildtype. If increased Ras1^{GTP} levels simply accelerate MAPK^{Spk1} activation, as

has been conventionally assumed, **pp**MAPK^{Spk1} production should peak earlier *and higher*

(Supplementary Fig S6B, best fit out of 1000 global fits), and the addition of Ras1^{GTP} only should

increase the amplitude, but not affect timing (Supplementary Fig S6C). We confirmed these

results in the best models from 1000 global fits (Materials and Methods). The comparison

between our experimental results and the *in silico* predictions suggests that the role of Ras1^{GTP}

is more complex than previously thought.

Strikingly, if we hypothesize that Ras1^{GTP} also contributes to the negative feedback [L10] (Fig.

7A), we recapitulate the "*earlier and lower*" peak of **pp**MAPK^{Spk1} in both the *ras1.G17V* and the

Cdc42-GEF^{scd1}Δ mutants (Fig. 7B). We currently do not have a direct experimental evidence to

support [L10]. However, considering the fact that Ras1^{GTP} likely acts as a physical signalling hub

at the cell cortex, mediating MAPKKK^{Byr2} recruitment and activation, which leads to recruitment

of MAPKK^{Byr1} and MAPK^{Spk1}, Ras1^{GTP} may work as a two-way amplifier, both assisting localized

MAPK^{Spk1} activation at the shmoo site, as well as helping the negative feedback by concentrating

the affected molecules.

The model successfully recapitulated the experimental results for wildtype, *ras1.G17V*,

MAPKK$^{Byr1.DD}$ and *cdc42-GEF^{scd1}Δ* mutants (Fig. 7B). To test the predictive capacity of the model,

we next performed an *in silico* experiment where we titrated increasing amounts of Ras1^{GTP} in

the wildtype condition before nitorgen removal. In agreement with our hypothesis, we obtained

a *ras1.G17V*-like **pp**MAPK^{Spk1} activation profile with increasing amount of Ras1^{GTP}, i.e., the

ppMAPK^{Spk1} peaks earlier with a lower peak intensity (Fig. 7E). The result further supported that

Ras1^{GTP} availability alone is sufficient to explain both the *ras1.G17V* and *Cdc42-GEF^{scd1}Δ*

phenotypes.

To further test the predictive value of the model, we asked whether it could predict **pp**MAPK^{Spk1}

dynamics in the 21 other strains, which were measured (Fig. 2-6), but not used for fitting the

model. We implemented each mutation in the wildtype model (Supplementary Table S2) and

the model accurately predicted relative **pp**MAPK^{Spk1} dynamics in 17 cases, or showed

predicitons in close proximity to the observed **pp**MAPK^{Spk1} dynamics in the 4 remaining cases

(Fig. 7F). Concluding from these results, our model likely represents the physiological framework

of fission yeast RAS-MAPK signalling.

Discussion

By quantitating the MAPK^{Spk1} and Cdc42 activation status during the mating process and

conducting epistasis analysis between numerous signalling mutants, we showed that Ras1

coordinates activation of both the MAPK^{Spk1} cascade and the Cdc42 pathway. Furthermore, we

revealed that the *ras1.G17V* mutant phenotype is caused by deregulation of Cdc42, rather than

altered activation of MAPK^{Spk1} in physiological setting. Based on the experimental data, we built

a mathematical model, which hypothesize that the MAPK^{Spk1} cascade is subject to robust

feedback regulation and two Ras1 effectors, Cdc42-GEF^{Scd1} and MAPKKK^{Byr2}, are competing for

active Ras1. This model faithfully recapitulates MAPK^{Spk1} activation profiles in the wildtype and

all mutant strains examined in this study. The model implies that targeting one of the RAS

effector pathways can potentially result in a complex outcome, rather than simply shutting

down the targeted effector pathway. We concluded that fission yeast pheromone Ras signalling

is not only defined by compartmentalisation (Onken et al., 2006) but rather a coordination of

events involving both the MAPK^{Spk1} and Cdc42 pathways.

In this study, we confirmed that Gpa1 is the central player of the pheromone signalling. It is

likely that Gpa1 is the most downstream molecule conveying the complete pheromone signal.

Considering that all the pheromone signalling components examined so far have been found at

the shmoo site (Dudin et al., 2016; Merlini et al., 2016; Merlini et al., 2018)(this study), an

attractive hypothesis is as follows: firstly, the activated pheromone receptor Map3/Mam2

locally activates Gpa1, which activates Ras1 and MAPK^{Spk1}. This then leads to a localised

activation of the Cdc42, causing shmoo formation in the direction of a mating partner (Fig. 7G).

The *ras1.G17V* mutation led to a acute activation of MAPK^{Spk1} compared with the wildtype cells.

The G17V mutantion of Ras1 is equivalent of G12V mutation of mammalian RAS, which results

in a substantial reduction of both intrinsic and GAP-mediated GTPase activities (Trahey and

McCormick, 1987). Therefore a larger fraction of Ras1 is expected to be in the GTP-bound form

in the *ras1.G17V* mutant cells. By mathematical modelling, we showed that the increased

Ras1^{GTP} pool can explain a faster acute activation of MAPK^{Spk1}.

To our surprise, the constitutive Ras1.G17V mutation did not induce over-activation of

MAPK[Spk1]. The attenuated MAPK[Spk1] activation in the presence of Ras1.G17V indicated that an

efficient feedback mechanism is in place to counteract the effect of Ras1.G17V. Strikingly, the

same trend has been reported in the mouse model of the *K-ras*[G12D] mutation integrated at the

endogenous chromosome locus (Tuveson et al., 2004). Therefore it is highly likely that the

MAPK cascade is generally robust against upstream oncogenic constitutive stimulation.

Based on our observation that the MAPK[Spk1] is constitutively activated in the *MAPKK*[byr1.DD]

mutant, we predict that the negative regulation occurs upstream of, or at the same level as,

MAPKK[Byr1], rather than phosphatases that directly regulate MAPK[Spk1]. In humans, ERK is shown

to phosphorylate RAF proteins, the prototype MAPKKKs, to contribute to ERK signal attenuation

(Brummer et al., 2003; Dougherty et al., 2005; Ritt et al., 2010). In future studies it will be

important to determine whether MAPK[Spk1] can directly downregulate MAPKKK[Byr2] in a

physiological setting.

Our results also show that an adaptor[Ste4] plays a crucial role in activating MAPKKK[Byr2], abolishing

ppMAPK[Spk1] production even in the presence of *ras1.G17V* mutation. This suggests that the

adaptor[Ste4] fits well to be one of the major targets by the negative feedback loop against

ppMAPK[Spk1]. This mechanism is shared by budding yeast, where an adaptor protein, Ste50,

modulates MAPKKK[Ste11] (Ramezani-Rad, 2003). In humans, although such an adaptor protein for

RAF proteins has yet to be identified, multiple RAF-interacting proteins, including 14-3-3

proteins, as well as formation of heterodimers between BRAF and CRAF, have been studied for

their Ras-independent mechanism to activate RAF proteins (Lavoie and Therrien, 2015).

Collectively, MAPK cascades seem to retain a general resistance to oncogenic RAS mutations in

physiological settings.

Whilst lack of Cdc42 activation does not impair MAPK[Spk1] activation, we found that MAPK[Spk1]

activity is required for shmoo formation even in the presence of Ras1.G17V (Fig. 2C). Therefore,

the two Ras1 effectors, Cdc42 and MAPK[Spk1] pathways, are not completely separable. The

situation is reminiscent of the *K-ras*[G12D] MEFs (Tuveson et al., 2004). In this system, the *K-ras*[G12D]

MEFs showed morphological anomalies. As both ERK and AKT phosphorylation levels in the *K-*

ras[G12D] MEFs resembled wildtype, these pathways unlikely caused the morphological

phenotype. Nonetheless, inhibitors against MAPK and PI3K pathways reverted the *K-ras*[G12D]–

induced abnormal morphology back to the one similar to wildtype. The observation suggests

that MAPK and PI3K pathways somehow contribute to the *K-ras*[G12D] morphological phenotype;

for example, a basal level of MAPK and PI3K pathway activation may be a prerequisite for the *K-*

ras^G12D–induced morphological anomalies.

The molecular mechanism of how MAPK^Spk1 contributes to Cdc42 activation will require further

studies. Key components in pheromone signalling are transcriptionally up-regulated upon

pheromone signalling (Xue-Franzen et al., 2006). This is driven by MAPK^Spk1, which activates the

master transcriptional regulator Ste11 (Kjaerulff et al., 2005; Mata and Bahler, 2006; Xue-

Franzen et al., 2006). Therefore, the contribution of MAPK^Spk1 to Cdc42 activation is expected to

occur, at least partly, through transcriptional activation.

In addition, localisation of signalling components may be regulated by MAPK^Spk1. In budding

yeast, Cdc24, the GEF for Cdc42, is sequestered into the nucleus by an adaptor protein Far1

(Nern and Arkowitz, 2000; Shimada et al., 2000). Upon the pheromone signalling, budding yeast

MAPK^Fus3 phosphorylates Far1, which then brings Cdc24 out to the shmoo site, leading to Cdc42

activation on the cell cortex (Hegemann et al., 2015).

MAPK^Spk1 may also directly phosphorylate to activate Cdc42 and/or its regulatory proteins such

as Cdc42-GEF^Scd1, Scd2 or GAP-Cdc42^Rga4, all of which function at the shmoo site during the

mating (Bendezu and Martin, 2013; Dudin et al., 2016)(Fig.4 A and B). In agreement with this

hypothesis, a transient MAPK^Spk1 and MAPKK^Byr1 signal on the cell cortex was observed during

the mating process (Fig. 1)(Dudin et al., 2016). Localisation of MAPK at the growing cell tips was

also observed in other fungi including *S. cerevisiae* and *N. crassa* (Chen et al., 2010; Fleissner et

al., 2009; Maeder et al., 2007; van Drogen et al., 2001). In budding yeast, MAPK^Fus3 can directly

phosphorylates Bni1, a formin that organises actin filaments, to facilitate shmoo formation

(Matheos et al., 2004).

Interestingly, during the vegetative growth when expression of MAPK^Spk1 is repressed,

Ras1.G17V is still capable of activating Cdc42 (Fig. 4C and D). Whether other MAPKs, such as

Sty1 or Pmk1, contribute to Cdc42 activation during the vegetative cell cycle will be an

important question to answer. Intriguingly, recent studies show that Sty1 inhibits, rather than

assists, establishment of the Cdc42 polarity module (Mutavchiev et al., 2016). Collectively, it is

likely that the Cdc42 polarity module is regulated in a context dependent manner by multiple

MAPKs in a range of ways.

In this study we revealed the vital contribution of Cdc42 to induce the *ras1.G17V* phenotype in

fission yeast pheromone signalling. In mouse models, small GTPases, Cdc42 and Rac, are

required for *H-ras^G12V* induced transformation (Malliri et al., 2002; Stengel and Zheng, 2012).

Therefore, oncogenic RAS-induced Cdc42/Rac misregulation may be a common basis of

oncogenicity of mutated-RAS-induced signalling. Specifically targeting this process may

therefore be an effective strategy against oncogenic RAS-driven tumourigenesis.

Acknowledgment

The authors thank Tatsuya Maeda, Thibault Mayor, Janni Petersen, Louise Fairall, John Schwabe,

David Critchley, Andrey Reviyakin, Gary Willars and Mohan Harihar for helpful suggestions,

stimulating discussions and critical reading of the manuscript. We thank PROTEX and PNACL at

University of Leicester for their technical assistance. We are grateful to Kazu Shiozaki, Keith Gull

and Yeast Genetic Resource Center for providing strains and antibodies. This work was funded

by the Wellcome Trust Institutional Strategic Support Fund WT097828/Z/11/Z,

WT097828/Z/11/B and the Deutsche Forschungsgemeinschaft (DFG) EXC 81. A.V. was supported

by German Academic Exchange Service (DAAD) A0981674.

Author Contributions

E.J.K., S.R. and K.T. generated yeast strains. E.J.K., G.S. and K.T. monitored activation status of

MAPK^{Spk1} and Cdc42. A.V. and E.K. conducted mathematical modelling. E.J.K., K.S. and K.T.

conducted image analysis. M.T., R.G., C.P. and C.D. conducted biochemical analysis of Ras1, Byr2

and Scd1. E.J.K., A.V., E.K. and K.T. designed the experiments and interpreted the data.

Declaration of Interests

The authors declare no competing interests.

References

Barr, M.M., Tu, H., Van Aelst, L., and Wigler, M. (1996). Identification of Ste4 as a potential
regulator of Byr2 in the sexual response pathway of Schizosaccharomyces pombe. Mol Cell
Biol *16*, 5597-5603.
Bendezu, F.O., and Martin, S.G. (2013). Cdc42 explores the cell periphery for mate selection
in fission yeast. Current biology : CB *23*, 42-47.
Bitton, D.A., Grallert, A., Scutt, P.J., Yates, T., Li, Y., Bradford, J.R., Hey, Y., Pepper, S.D., Hagan,
I.M., and Miller, C.J. (2011). Programmed fluctuations in sense/antisense transcript ratios
drive sexual differentiation in S. pombe. Mol Syst Biol *7*, 559.

Brummer, T., Naegele, H., Reth, M., and Misawa, Y. (2003). Identification of novel ERK-
mediated feedback phosphorylation sites at the C-terminus of B-Raf. Oncogene *22*, 8823-
8834.
Chang, E.C., Barr, M., Wang, Y., Jung, V., Xu, H.P., and Wigler, M.H. (1994). Cooperative
interaction of S. pombe proteins required for mating and morphogenesis. Cell *79*, 131-141.
Chen, R.E., Patterson, J.C., Goupil, L.S., and Thorner, J. (2010). Dynamic localization of Fus3
mitogen-activated protein kinase is necessary to evoke appropriate responses and avoid
cytotoxic effects. Mol Cell Biol *30*, 4293-4307.
Cox, A.D., and Der, C.J. (2010). Ras history: The saga continues. Small GTPases *1*, 2-27.
Das, M., Wiley, D.J., Medina, S., Vincent, H.A., Larrea, M., Oriolo, A., and Verde, F. (2007).
Regulation of cell diameter, For3p localization, and cell symmetry by fission yeast Rho-GAP
Rga4p. Molecular biology of the cell *18*, 2090-2101.
Didmon, M., Davis, K., Watson, P., Ladds, G., Broad, P., and Davey, J. (2002). Identifying
regulators of pheromone signalling in the fission yeast Schizosaccharomyces pombe.
Current genetics *41*, 241-253.
Dougherty, M.K., Muller, J., Ritt, D.A., Zhou, M., Zhou, X.Z., Copeland, T.D., Conrads, T.P.,
Veenstra, T.D., Lu, K.P., and Morrison, D.K. (2005). Regulation of Raf-1 by direct feedback
phosphorylation. Mol Cell *17*, 215-224.
Dudin, O., Merlini, L., and Martin, S.G. (2016). Spatial focalization of pheromone/MAPK
signaling triggers commitment to cell-cell fusion. Genes Dev *30*, 2226-2239.
Endo, M., Shirouzu, M., and Yokoyama, S. (2003). The Cdc42 binding and scaffolding
activities of the fission yeast adaptor protein Scd2. J Biol Chem *278*, 843-852.
Fleissner, A., Leeder, A.C., Roca, M.G., Read, N.D., and Glass, N.L. (2009). Oscillatory
recruitment of signaling proteins to cell tips promotes coordinated behavior during cell
fusion. Proceedings of the National Academy of Sciences of the United States of America
106, 19387-19392.
Fukui, Y., Kozasa, T., Kaziro, Y., Takeda, T., and Yamamoto, M. (1986). Role of a ras homolog
in the life cycle of Schizosaccharomyces pombe. Cell *44*, 329-336.
Fukui, Y., Miyake, S., Satoh, M., and Yamamoto, M. (1989). Characterization of the
Schizosaccharomyces pombe ral2 gene implicated in activation of the ras1 gene product.
Mol Cell Biol *9*, 5617-5622.
Gronwald, W., Huber, F., Grunewald, P., Sporner, M., Wohlgemuth, S., Herrmann, C., and
Kalbitzer, H.R. (2001). Solution structure of the Ras binding domain of the protein kinase
Byr2 from Schizosaccharomyces pombe. Structure *9*, 1029-1041.
Hegemann, B., Unger, M., Lee, S.S., Stoffel-Studer, I., van den Heuvel, J., Pelet, S., Koeppl, H.,
and Peter, M. (2015). A Cellular System for Spatial Signal Decoding in Chemical Gradients.
Dev Cell *35*, 458-470.
Hirota, K., Tanaka, K., Watanabe, Y., and Yamamoto, M. (2001). Functional analysis of the C-
terminal cytoplasmic region of the M-factor receptor in fission yeast. Genes Cells *6*, 201-
214.
Hoops, S., Sahle, S., Gauges, R., Lee, C., Pahle, J., Simus, N., Singhal, M., Xu, L., Mendes, P., and
Kummer, U. (2006). COPASI--a COmplex PAthway SImulator. Bioinformatics *22*, 3067-3074.
Hughes, D.A., Fukui, Y., and Yamamoto, M. (1990). Homologous activators of ras in fission
and budding yeast. Nature *344*, 355-357.
Hughes, D.A., Yabana, N., and Yamamoto, M. (1994). Transcriptional regulation of a Ras
nucleotide-exchange factor gene by extracellular signals in fission yeast. J Cell Sci *107 (Pt*
12), 3635-3642.
Imai, Y., and Yamamoto, M. (1992). Schizosaccharomyces pombe sxa1+ and sxa2+ encode
putative proteases involved in the mating response. Mol Cell Biol *12*, 1827-1834.

Kelly, F.D., and Nurse, P. (2011). Spatial control of Cdc42 activation determines cell width in
fission yeast. Molecular biology of the cell *22*, 3801-3811.
Kjaerulff, S., Andersen, N.R., Borup, M.T., and Nielsen, O. (2007). Cdk phosphorylation of the
Ste11 transcription factor constrains differentiation-specific transcription to G1. Genes Dev
21, 347-359.
Kjaerulff, S., Lautrup-Larsen, I., Truelsen, S., Pedersen, M., and Nielsen, O. (2005).
Constitutive activation of the fission yeast pheromone-responsive pathway induces ectopic
meiosis and reveals ste11 as a mitogen-activated protein kinase target. Mol Cell Biol *25*,
2045-2059.
Lavoie, H., and Therrien, M. (2015). Regulation of RAF protein kinases in ERK signalling. Nat
Rev Mol Cell Biol *16*, 281-298.
Lin, A.W., Barradas, M., Stone, J.C., van Aelst, L., Serrano, M., and Lowe, S.W. (1998).
Premature senescence involving p53 and p16 is activated in response to constitutive
MEK/MAPK mitogenic signaling. Genes Dev *12*, 3008-3019.
Maeder, C.I., Hink, M.A., Kinkhabwala, A., Mayr, R., Bastiaens, P.I., and Knop, M. (2007).
Spatial regulation of Fus3 MAP kinase activity through a reaction-diffusion mechanism in
yeast pheromone signalling. Nat Cell Biol *9*, 1319-1326.
Malliri, A., van der Kammen, R.A., Clark, K., van der Valk, M., Michiels, F., and Collard, J.G.
(2002). Mice deficient in the Rac activator Tiam1 are resistant to Ras-induced skin tumours.
Nature *417*, 867-871.
Marcus, S., Polverino, A., Chang, E., Robbins, D., Cobb, M.H., and Wigler, M.H. (1995). Shk1, a
homolog of the Saccharomyces cerevisiae Ste20 and mammalian p65PAK protein kinases, is
a component of a Ras/Cdc42 signaling module in the fission yeast Schizosaccharomyces
pombe. Proceedings of the National Academy of Sciences of the United States of America *92*,
6180-6184.
Masuda, T., Kariya, K., Shinkai, M., Okada, T., and Kataoka, T. (1995). Protein kinase Byr2 is a
target of Ras1 in the fission yeast Schizosaccharomyces pombe. J Biol Chem *270*, 1979-
1982.
Mata, J., and Bahler, J. (2006). Global roles of Ste11p, cell type, and pheromone in the control
of gene expression during early sexual differentiation in fission yeast. Proceedings of the
National Academy of Sciences of the United States of America *103*, 15517-15522.
Mata, J., Lyne, R., Burns, G., and Bahler, J. (2002). The transcriptional program of meiosis
and sporulation in fission yeast. Nature genetics *32*, 143-147.
Mata, J., Wilbrey, A., and Bahler, J. (2007). Transcriptional regulatory network for sexual
differentiation in fission yeast. Genome biology *8*, R217.
Matheos, D., Metodiev, M., Muller, E., Stone, D., and Rose, M.D. (2004). Pheromone-induced
polarization is dependent on the Fus3p MAPK acting through the formin Bni1p. J Cell Biol
165, 99-109.
Merlini, L., Dudin, O., and Martin, S.G. (2013). Mate and fuse: how yeast cells do it. Open Biol
3, 130008.
Merlini, L., Khalili, B., Bendezu, F.O., Hurwitz, D., Vincenzetti, V., Vavylonis, D., and Martin,
S.G. (2016). Local Pheromone Release from Dynamic Polarity Sites Underlies Cell-Cell
Pairing during Yeast Mating. Current biology : CB *26*, 1117-1125.
Merlini, L., Khalili, B., Dudin, O., Michon, L., Vincenzetti, V., and Martin, S.G. (2018).
Inhibition of Ras activity coordinates cell fusion with cell-cell contact during yeast mating. J
Cell Biol *217*, 1467-1483.
Mutavchiev, D.R., Leda, M., and Sawin, K.E. (2016). Remodeling of the Fission Yeast Cdc42
Cell-Polarity Module via the Sty1 p38 Stress-Activated Protein Kinase Pathway. Current
biology : CB *26*, 2921-2928.

Nadin-Davis, S.A., and Nasim, A. (1988). A gene which encodes a predicted protein kinase
can restore some functions of the ras gene in fission yeast. EMBO J *7*, 985-993.
Nadin-Davis, S.A., Nasim, A., and Beach, D. (1986a). Involvement of ras in sexual
differentiation but not in growth control in fission yeast. EMBO J *5*, 2963-2971.
Nadin-Davis, S.A., Yang, R.C., Narang, S.A., and Nasim, A. (1986b). The cloning and
characterization of a RAS gene from Schizosaccharomyces pombe. J Mol Evol *23*, 41-51.
Nern, A., and Arkowitz, R.A. (2000). Nucleocytoplasmic shuttling of the Cdc42p exchange
factor Cdc24p. J Cell Biol *148*, 1115-1122.
Obara, T., Nakafuku, M., Yamamoto, M., and Kaziro, Y. (1991). Isolation and characterization
of a gene encoding a G-protein alpha subunit from Schizosaccharomyces pombe:
involvement in mating and sporulation pathways. Proceedings of the National Academy of
Sciences of the United States of America *88*, 5877-5881.
Okazaki, N., Okazaki, K., Tanaka, K., and Okayama, H. (1991). The ste4+ gene, essential for
sexual differentiation of Schizosaccharomyces pombe, encodes a protein with a leucine
zipper motif. Nucleic acids research *19*, 7043-7047.
Onken, B., Wiener, H., Philips, M.R., and Chang, E.C. (2006). Compartmentalized signaling of
Ras in fission yeast. Proceedings of the National Academy of Sciences of the United States of
America *103*, 9045-9050.
Ottilie, S., Miller, P.J., Johnson, D.I., Creasy, C.L., Sells, M.A., Bagrodia, S., Forsburg, S.L., and
Chernoff, J. (1995). Fission yeast pak1+ encodes a protein kinase that interacts with Cdc42p
and is involved in the control of cell polarity and mating. EMBO J *14*, 5908-5919.
Ozoe, F., Kurokawa, R., Kobayashi, Y., Jeong, H.T., Tanaka, K., Sen, K., Nakagawa, T., Matsuda,
H., and Kawamukai, M. (2002). The 14-3-3 proteins Rad24 and Rad25 negatively regulate
Byr2 by affecting its localization in Schizosaccharomyces pombe. Mol Cell Biol *22*, 7105-
7119.
Papadaki, P., Pizon, V., Onken, B., and Chang, E.C. (2002). Two ras pathways in fission yeast
are differentially regulated by two ras guanine nucleotide exchange factors. Mol Cell Biol *22*,
4598-4606.
Pereira, P.S., and Jones, N.C. (2001). The RGS domain-containing fission yeast protein,
Rgs1p, regulates pheromone signalling and is required for mating. Genes Cells *6*, 789-802.
Prior, I.A., Lewis, P.D., and Mattos, C. (2012). A comprehensive survey of Ras mutations in
cancer. Cancer Res *72*, 2457-2467.
Qiu, R.G., Abo, A., McCormick, F., and Symons, M. (1997). Cdc42 regulates anchorage-
independent growth and is necessary for Ras transformation. Mol Cell Biol *17*, 3449-3458.
Ramachander, R., Kim, C.A., Phillips, M.L., Mackereth, C.D., Thanos, C.D., McIntosh, L.P., and
Bowie, J.U. (2002). Oligomerization-dependent association of the SAM domains from
Schizosaccharomyces pombe Byr2 and Ste4. J Biol Chem *277*, 39585-39593.
Ramezani-Rad, M. (2003). The role of adaptor protein Ste50-dependent regulation of the
MAPKKK Ste11 in multiple signalling pathways of yeast. Current genetics *43*, 161-170.
Ritt, D.A., Monson, D.M., Specht, S.I., and Morrison, D.K. (2010). Impact of feedback
phosphorylation and Raf heterodimerization on normal and mutant B-Raf signaling. Mol
Cell Biol *30*, 806-819.
Schubbert, S., Bollag, G., and Shannon, K. (2007). Deregulated Ras signaling in
developmental disorders: new tricks for an old dog. Curr Opin Genet Dev *17*, 15-22.
Shimada, Y., Gulli, M.-P., and Peter, M. (2000). Nuclear sequestration of the exchange factor
Cdc24 by Far1 regulates cell polarity during yeast mating. Nature Cell Biology *2*, 117.
Sipiczki, M. (1988). The role of sterility genes (ste and aff) in the initiation of sexual
development in Schizosaccharomyces pombe. Molecular & general genetics : MGG *213*, 529-
534.

Stengel, K.R., and Zheng, Y. (2012). Essential role of Cdc42 in Ras-induced transformation
revealed by gene targeting. PLoS One *7*, e37317.
Sugimoto, A., Iino, Y., Maeda, T., Watanabe, Y., and Yamamoto, M. (1991).
Schizosaccharomyces pombe ste11+ encodes a transcription factor with an HMG motif that
is a critical regulator of sexual development. Genes Dev *5*, 1990-1999.
Tatebe, H., Nakano, K., Maximo, R., and Shiozaki, K. (2008). Pom1 DYRK regulates
localization of the Rga4 GAP to ensure bipolar activation of Cdc42 in fission yeast. Current
biology : CB *18*, 322-330.
Toda, T., Shimanuki, M., and Yanagida, M. (1991). Fission yeast genes that confer resistance
to staurosporine encode an AP-1-like transcription factor and a protein kinase related to
the mammalian ERK1/MAP2 and budding yeast FUS3 and KSS1 kinases. Genes Dev *5*, 60-
73.
Trahey, M., and McCormick, F. (1987). A cytoplasmic protein stimulates normal N-ras p21
GTPase, but does not affect oncogenic mutants. Science *238*, 542-545.
Tratner, I., FourticqEsqueoute, A., Tillit, J., and Baldacci, G. (1997). Cloning and
characterization of the S-pombe gene efc25(+), a new putative guanine nucleotide exchange
factor. Gene *193*, 203-210.
Tu, H., Barr, M., Dong, D.L., and Wigler, M. (1997). Multiple regulatory domains on the Byr2
protein kinase. Mol Cell Biol *17*, 5876-5887.
Tuveson, D.A., Shaw, A.T., Willis, N.A., Silver, D.P., Jackson, E.L., Chang, S., Mercer, K.L.,
Grochow, R., Hock, H., Crowley, D*., et al.* (2004). Endogenous oncogenic K-ras(G12D)
stimulates proliferation and widespread neoplastic and developmental defects. Cancer Cell
5, 375-387.
van Drogen, F., Stucke, V.M., Jorritsma, G., and Peter, M. (2001). MAP kinase dynamics in
response to pheromones in budding yeast. Nat Cell Biol *3*, 1051-1059.
Verde, F., Mata, J., and Nurse, P. (1995). Fission yeast cell morphogenesis: identification of
new genes and analysis of their role during the cell cycle. J Cell Biol *131*, 1529-1538.
Wang, Y., Xu, H.P., Riggs, M., Rodgers, L., and Wigler, M. (1991). byr2, a Schizosaccharomyces
pombe gene encoding a protein kinase capable of partial suppression of the ras1 mutant
phenotype. Mol Cell Biol *11*, 3554-3563.
Watson, P., Davis, K., Didmon, M., Broad, P., and Davey, J. (1999). An RGS protein regulates
the pheromone response in the fission yeast Schizosaccharomyces pombe. Mol Microbiol
33, 623-634.
Weston, C., Bond, M., Croft, W., and Ladds, G. (2013). The coordination of cell growth during
fission yeast mating requires Ras1-GTP hydrolysis. PLoS One *8*, e77487.
Xu, H.P., White, M., Marcus, S., and Wigler, M. (1994). Concerted action of RAS and G
proteins in the sexual response pathways of Schizosaccharomyces pombe. Mol Cell Biol *14*,
50-58.
Xue-Franzen, Y., Kjaerulff, S., Holmberg, C., Wright, A., and Nielsen, O. (2006). Genomewide
identification of pheromone-targeted transcription in fission yeast. BMC Genomics *7*, 303.
Yamamoto, M. (1996). The molecular control mechanisms of meiosis in fission yeast.
Trends Biochem Sci *21*, 18-22.
Zheng, C.F., and Guan, K.L. (1994). Activation of MEK family kinases requires
phosphorylation of two conserved Ser/Thr residues. EMBO J *13*, 1123-1131.
Zhu, J., Woods, D., McMahon, M., and Bishop, J.M. (1998). Senescence of human fibroblasts
induced by oncogenic Raf. Genes Dev *12*, 2997-3007.
782
783

Figure legends

Figure 1. Distinct modes of MAPK^{Spk1} temporal phosphorylation profile and morphological

changes during sexual differentiation in wildtype, *MAPKK$^{byr1.DD}$* and *ras1.G17V* mutants.

(A) A pictorial representation of wildtype fission yeast sexual differentiation. (B) A list of key

signalling components of the fission yeast pheromone signalling pathway. The diagram reflects

the prediction that Gpa1 and Ras1 separately contribute to activation of MAPKKK^{Byr2} activation

although the precise mechanism is unknown (Xu et al., 1994). At the same time, Ras1 activation

is expected to be at least partly under influence of active Gpa1 because the *ste6* gene, encoding

a Ras1 activator, is strongly induced upon successful pheromone signaling (Hughes et al., 1994).

(C)-(H) Cells were induced for sexual differentiation by the plate mating assay system as

described in the materials and methods. (C), (E) and (G) Quantified **pp**MAPK^{Spk1} signal from

western blots of wildtype (KT3082) (C), *MAPKK$^{byr1.DD}$* (KT3435) (E) and *ras1.G17V* (KT3084) (G)

cells. Three biological replicates were used for quantitation (error bars are ±SEM). α-tubulin was

used as a loading control and quantitation was carried out using the Image Studio ver2.1

software (Licor Odyssey CLx Scanner). For the wildtype samples in (C), the % of cells mating is

also indicated. The wildtype **pp**MAPK^{Spk1} result (C) is also presented in (E) and (G) as a reference.

(D), (F) and (H) Cellular morphology (brightfield) and localization of MAPK^{Spk1}-GFP over a 24 hour

time-course in wildtype (D), *MAPKK$^{byr1.DD}$* (F) and *ras1.G17V* (H) cells. Time after induction of

mating in hours is indicated on the left. At each time point, a bright-field image and a GFP signal

image were taken and processed as described in materials and methods. Green asterisks in the

time 24 h in the *ras1.G17V* cell image (H) indicate auto-fluorescence signal from inviable cell

debris, which were presumably produced through cytokinesis failure or cell lysis. Yellow arrows

in panels (D) and (F) indicate transient accumulation of MAPK^{Spk1}-GFP at the shmoo tips. Scale

bars represent 10μm.

Figure 2. In the *ras1.G17V MAPKK$^{byr1.DD}$* double mutant, the MAPK^{Spk1} phosphorylation profile

follows *MAPKK$^{byr1.DD}$* single mutant phenotype whilst cell morphology mimics the *ras1.G17V*

single mutant phenotype.

(A) MAPK^{Spk1} phosphorylation status in the *ras1.G17V MAPKK$^{byr1.DD}$* double mutant cells

(KT3439). Cells were induced for mating by the plate mating assay system as described in the

materials and methods. Quantitated **pp**MAPK^{Spk1} signal (arbitrary unit) from western blots is

presented. Results of two biological replicates (each derived from three technical replicates,

error bars are ±SEM) are presented in red. (B) The terminal mating phenotype of *ras1.G17V*

MAPKK^byr1.DD double mutant is a phenocopy of *ras1.G17V* single mutant which shows the

"elongated" morphology. Images were taken of *ras1.G17V MAPKK^byr1.DD* double mutant

(KT3439) in the same way as in Fig. 1. Time after induction of mating in hours is indicated on the

left. (C) There is no morphological change in the absence of MAPK^Spk1 signalling. Cell images of

MAPKK^byr1Δ (KT4700) and *ras1.G17V MAPKK^byr1Δ* (KT5030) strains are shown. Images were

taken in the same way as in Fig. 1. Time after induction of mating in hours is indicated on the

left of each series. Scale bars represent 10μm.

Figure 3. Ras1 activates both MAPK^Spk1 and Cdc42 pathways during pheromone signalling.

(A) *scd1Δ* morphology and MAPK^Spk1-GFP signal. Images of WT (KT3082), *scd1Δ* (KT4061) and

scd1Δ ras1.G17V double mutant (KT4056) were taken in the same way as in Fig.1. Numbers on

the left represents hours after induction of mating. (B) MAPK^Spk1 phosphorylation state in *scd1Δ*

(KT4061) cells after mating induction. Results of three biological replicates (error bars are ±SEM)

are presented. The wildtype **pp**MAPK^Spk1 result presented in Fig.1 (C) is also shown in blue as a

reference. (C) Cell images of *scd1Δ MAPKK^byr1.DD* double mutant (KT4047) were taken in the

same way as in Fig.1. Numbers on the left represents hours after induction of mating. (D)

MAPK^Spk1 phosphorylation state in *ras1Δ* (KT4323), *MAPKK^byr1.DD* (KT3435) and *ras1Δ*

MAPKK^byr1.DD (KT4359) cell extracts. Original Western blotting data is presented in Fig. S3A. (E)

MAPK^Spk1 phosphorylation state in *mapkkk^byr2Δ* (KT3763), *MAPKK^byr1.DD* (KT3435) and

mapkkk^byr2Δ MAPKK^byr1.DD (KT4010) cell extracts. Original Western blotting data is presented in

Fig. S3B. For (D) and (E), quantification was carried out using the Image Studio ver2.1 (Li-cor). (F)

Cell images of the strains mentioned in (D) and (E) were taken in the same way as in Fig.1.

Numbers on the left represents hours after induction of mating. For all the images presented in

(A), (C) and (F), scale bars represent 10μm.

Figure 4. Ras1.G17V induces cortical Cdc42^GTP accumulation

(A) Cell morphology and localisation of Cdc42^GTP, indicated by CRIB-GFP signal, during the sexual

differentiation process. Wildtype (KT5077) and *ras1.G17V* (KT5082) mutant cells were induced

for mating/sexual differentiation by the plate mating assay condition (Materials and Methods)

and live cell images were taken at the indicated time after induction of mating/sexual

differentiation. Representative CRIB-GFP signal images are presented. Cells with cortical CRIB-

GFP foci are indicated by orange stars. Rapidly-disappearing CRIB-GFP signals at the fusion site

of wildtype mating cells are indicated by green arrows at time 8.5h image. Scale bar: 10 μm. (B)

Quantitation of the results presented in (A). At each time point (4.5h, 6.5h, 10.5h, 12.5h and

22.5h after induction of mating/sexual differentiation), 150 cells were examined whether they

have cortical CRIB-GFP foci. % cells with cortical CRIB-GFP foci is presented. The experiment was

repeated for three times and the mean values and SDs are plotted in the graph. (C) Cell

morphology and localisation of Cdc42^{GTP}, indicated by CRIB-GFP signal, during vegetative

growth. Representative CRIB-GFP signal images of cells of wildtype (KT5077), *ras1Δ* (5107),

ras1.G17V (KT5082), *rga4Δ* (5551) and *rga4Δ ras1.G17V* (KT5554) are presented. Scale bar: 10

μm. (D) Quantitated CRIB-GFP signals on the cell cortex of cells presented in (C). Intensity of GFP

signal on the cell cortex was measured along one of the cell tips as indicated as a magenta

dotted line in the example image on the right (Scale bar: 10 μm) as stated in the Materials and

Methods. 40 cells without septum were measured for each strain and the average curve from all

aligned traces per strain was calculated, and displayed with respective standard error of the

mean curves (dashed lines) as described in Materials and Methods.

Fig. 5. Distinct contributions of Ste4 and Ste6 to MAPK^{Spk1} phosphorylation.

(A) Ste4 is essential for MAPK^{Spk1} activation. MAPK^{Spk1} phosphorylation status in *ste4Δ* (KT4376),

ste4Δ ras1.G17V (KT5143) and *ste4Δ MAPKK$^{byr1.DD}$* (KT5136) at times-points 0, 8 ,16 and 24

hours post mating induction are presented. Original Western blotting membranes are presented

in Fig. S3C. (B) Incapability of *ste4Δ* to cause pheromone-induced morphological change is

suppressed by *MAPKK$^{byr1.DD}$* but not by *ras1.G17V*. Cell images of *ste4Δ* (KT4376), *ste4Δ*

ras1.G17V (KT5143) and *ste4Δ MAPKK$^{byr1.DD}$* (KT5136) strains were taken 24 hours after

induction of mating. Scale bar: 10 μm. (C) Lack of Ste6 does not result in the complete loss of

MAPK^{Spk1} phosphorylation. MAPK^{Spk1} phosphorylation status in *ste6Δ* (KT4333), *ste6Δ ras1.G17V*

(KT4998) and *ste6Δ MAPKK$^{byr1.DD}$* (KT5139) at times-points 0, 8 ,16 and 24 hours post mating

induction are presented. Original Western blotting membranes are presented in Fig. S3D. (D)

Incapability of *ste6Δ* to cause pheromone-induced morphological change is suppressed by

ras1.G17V mutation but not by *MAPKK$^{byr1.DD}$* mutation. Cell images of *ste6Δ* (KT4333), *ste6Δ*

ras1.G17V (KT4998) and *ste6Δ MAPKK^byr1.DD* (KT5139) strains were taken 24 hours after

induction of mating. Scale bar: 10 μm.

Fig. 6. Gpa1 transduces the pheromone signalling by activating MAPK^Spk1 and Ras1 pathways

(A) MAPK^Spk1 phosphorylation status in homothallic *gpa1Δ* (KT4335), *gpa1Δ ras1.G17V*

(KT5023), *gpa1Δ MAPKK^byr1.DD* (KT4353) and *gpa1Δ ras1.val17 MAPKK^byr1.DD* (KT5035) at times-

points 0, 8 ,12, 16 and 24 hours after mating induction. Original Western membrane is

presented in Fig. S4A. (B) Cell images of the above mentioned strains at 16 hours after mating

induction. All the cell images were taken and processed as in Figure 1. Scale bar is 10 μm. (C)

MAPK^Spk1 phosphorylation status in *h⁻* WT (KT4190), *h⁻ gpa1.QL* (KT5059), *h⁻ ras1.G17V* (KT4233),

h⁻ gpa1.QL ras1Δ (KT5070) and h⁻ *MAPKK^byr1.DD* (KT4194) at times-points 0, 8 ,12 and 24 after

mating induction. Note that while the activation induced in the *gpa1.QL* mutant was down-

regulated, *MAPKK^byr1.DD* induced a constitutive activation. Original Western membrane is

presented in Fig. S4B. (D) Cell images of the above strains at 12 h after induction of mating. All

the cell images were taken and processed as in Figure 1. Scale bar is 10 μm.

Fig. 7. Mathematical modelling of the fission yeast pheromone pathway dynamics.

(A) Components and frameworks of the mathematical model in wildtype and signalling mutants:

ras1.G17V, *Cdc42GEF^scd1Δ,* and *MAPKK^byr1.DD*. Changes corresponding to each mutant are

indicated as follows: Grey: removed components or interactions, orange: increased level of

activity. For the exact implementation of the mutants, see Materials and Method. The

measured component, **pp**MAPK^Spk1, is highlighted in green. (B) Measured and simulated

ppMAPK^Spk1 activation profiles in wildtype, *ras1.G17V*, *Cdc42GEF^scd1Δ* and *MAPKK^byr1.DD* mutants.

Dashed lines: model simulations. Diamonds: experimental data presented in Fig. 1C, E, G, and

Fig. 3B; error bars: SEM. (C) GTP-loaded Ras1.G17V (1-172) directly binds to Byr2 (65-180) and

Scd1 (760-872). *In vitro* GST pull-down assays of bacterially expressed Ras1.G17V (1-172), GST-

Byr2 (65-180) and GST-Scd1 (760-872) were conducted as described in materials and methods.

GTP-loaded Ras1.G17V (1-172) was found to bind to both GST-Byr2 (65-180) and GST-Scd1 (760-

872). (D) Two Ras1 effectors, Byr2 and Scd1, compete for GTP-loaded Ras1.G17V (1-172). In

vitro GST pull-down assays of bacterially expressed Ras1.G17V (1-172) and GST-Byr2 (65-180)

were conducted as in (A). Addition of Scd1 (760-872) fragment interfered with Ras1-Byr2

binding (the 4th lane). Quantitated signal intensities of the Ras1.G17V (1-172) band in the gel are

shown in the right panel. (E) Simulated ppMAPK^{Spk1} dynamics in the wildtype model at

increasing concentrations of Ras1^{GTP} added *in silico* to the system. Increased Ras1^{GTP}

concentration causes advanced and reduced **pp**MAPK^{Spk1} peak intensities. (F) The model fitted

to the 4 strains (as above, in red) correctly predicts ppMAPK^{Spk1} dynamics in the additional 21

signalling mutant strains measured in this study.(G) Schematic diagram of the fission yeast

pheromone signalling pathway, highlighting the branched pheromone sensing, and that

ultimately both branches are necessary for mating.

A homothalic *h90* cells (mixed population of *h+* and *h-* cells)

vegetative growth → Nitrogen starvation → G1 arrest → pheromone signal → shmoo formation → mating → karyogamy → sporulation

B RAS-MAPK signalling components in fission yeast pheromone signaling pathway

Pheromone (P-factor, M-factor) → Receptor (Mam2, Map3) → Receptor-coupled G protein (Gpa1) → Ras (Ras1) → MAPKKK (Byr2) → MAPKK (Byr1) → MAPK (Spk1)

Figure 1. Distinct modes of MAPK^{Spk1} temporal phosphorylation profile and morphological changes during sexual differentiation in wildtype, *MAPKK$^{byr1.DD}$* and *ras1.G17V* mutants.

(A) A pictorial representation of wildtype fission yeast sexual differentiation. (B) A list of key signalling components of the fission yeast pheromone signalling pathway. The diagram reflects the prediction that Gpa1 and Ras1 separately contribute to activation of MAPKKK^{Byr2} activation although the precise mechanism is unknown (Xu et al., 1994). At the same time, Ras1 activation is expected to be at least partly under influence of active Gpa1 because the ste6 gene, encoding a Ras1 activator, is strongly induced upon successful pheromone signaling (Hughes et al., 1994). (C)-(H) Cells were induced for sexual differentiation by the plate mating assay system as described in the materials and methods. (C), (E) and (G) Quantified **pp**MAPK^{Spk1} signal from western blots of wildtype (KT3082) (C), *MAPKK$^{byr1.DD}$* (KT3435) (E) and *ras1.G17V* (KT3084) (G) cells. Three biological replicates were used for quantitation (error bars are ±SEM). α-tubulin was used as a loading control and quantitation was carried out using the Image Studio ver2.1 software (Licor Odyssey CLx Scanner). For the wildtype samples in (C), the % of cells mating is also indicated. The wildtype **pp**MAPK^{Spk1} result (C) is also presented in (E) and (G) as a reference. (D), (F) and (H) Cellular morphology (brightfield) and localization of MAPK^{Spk1}-GFP over a 24 hour time-course in wildtype (D), *MAPKK$^{byr1.DD}$* (F) and *ras1.G17V* (H) cells. Time after induction of mating in hours is indicated on the left. At each time point, a bright-field image and a GFP signal image were taken and processed as described in materials and methods. Green asterisks in the time 24 h in the ras1.G17V cell image (H) indicate auto-fluorescence signal from inviable cell debris, which were presumably produced through cytokinesis failure or cell lysis. Yellow arrows in panels (D) and (F) indicate transient accumulation of MAPK^{Spk1}-GFP at the shmoo tips. Scale bars represent 10μm.

Figure 2. In the *ras1.G17V MAPKK^byr1.DD* double mutant, the MAPK^Spk1 phosphorylation profile follows *MAPKK^byr1.DD* single mutant phenotype whilst cell morphology mimics the *ras1.G17V* single mutant phenotype.

(A) MAPK^Spk1 phosphorylation status in the *ras1.G17V MAPKK^byr1.DD* double mutant cells (KT3439). Cells were induced for mating by the plate mating assay system as described in the materials and methods. Quantitated **pp**MAPK^Spk1 signal (arbitrary unit) from western blots is presented. Results of two biological replicates (each derived from three technical replicates, error bars are ±SEM) are presented in red. (B) The terminal mating phenotype of *ras1.G17V MAPKK^byr1.DD* double mutant is a phenocopy of *ras1.G17V* single mutant which shows the "elongated" morphology. Images were taken of *ras1.G17V MAPKK^byr1.DD* double mutant (KT3439) in the same way as in Fig. 1. Time after induction of mating in hours is indicated on the left. (C) There is no morphological change in the absence of MAPK^Spk1 signalling. Cell images of *MAPKK^byr1Δ* (KT4700) and *ras1.G17V MAPKK^byr1.DD* (KT5030) strains are shown. Images were taken in the same way as in Fig. 1. Time after induction of mating in hours is indicated on the left of each series. Scale bars represent 10μm.

Figure 3. Ras1 activates both MAPK^{Spk1} and Cdc42 pathways during pheromone signalling.
(A) *scd1Δ* morphology and MAPK^{Spk1}-GFP signal. Images of WT (KT3082), *scd1Δ* (KT4061) and *scd1Δ ras1.G17V* double mutant (KT4056) were taken in the same way as in Fig.1. Numbers on the left represents hours after induction of mating. (B) MAPK^{Spk1} phosphorylation state in *scd1Δ* (KT4061) cells after mating induction. Results of three biological replicates (error bars are ±SEM) are presented. The wildtype result presented in Fig.1 (C) is also shown in blue as a reference. (C) Cell images of *scd1Δ MAPKK$^{byr1.DD}$* double mutant (KT4047) were taken in the same way as in Fig.1. Numbers on the left represents hours after induction of mating. (D) MAPK^{Spk1} phosphorylation state in *ras1Δ* (KT4323), *MAPKK$^{byr1.DD}$* (KT3435) and *scd1Δ MAPKK$^{byr1.DD}$* (KT4359) cell extracts. Original Western blotting data is presented in Fig. S3A. (E) MAPK^{Spk1} phosphorylation state in *mapkkk^{byr2}Δ* (KT3763), *MAPKK$^{byr1.DD}$* (KT3435) and *mapkkk^{byr2}Δ MAPKK$^{byr1.DD}$* (KT4010) cell extracts. Original Western blotting data is presented in Fig. S3B. For (D) and (E), quantification was carried out using the Image Studio ver2.1 (Li-cor). (F) Cell images of the strains mentioned in (D) and (E) were taken in the same way as in Fig.1. Numbers on the left represents hours after induction of mating. For all the images presented in (A), (C) and (F), scale bars represent 10μm.

Figure 4. Ras1.G17V induces cortical Cdc42^{GTP} accumulation

(A) Cell morphology and localisation of Cdc42^{GTP}, indicated by CRIB-GFP signal, during the sexual differentiation process. Wildtype (KT5077) and *ras1.G17V* (KT5082) mutant cells were induced for sexual differentiation by the plate mating assay condition (Materials and Methods) and live cell images were taken at the indicated time after induction of mating/sexual differentiation. Representative CRIB-GFP signal images are presented. Cells with cortical CRIB-GFP foci are indicated by orange stars. Rapidly-disappearing CRIB-GFP signals at the fusion site of wildtype mating cells are indicated by green arrows at time 8.5h image. Scale bar: 10 μm. (B) Quantitation of the results presented in (A). At each time point (4.5h, 6.5h, 10.5h, 12.5h and 22.5h after induction of mating/sexual differentiation), 150 cells were examined whether they have cortical CRIB-GFP foci. % cells with cortical CRIB-GFP foci is presented. The experiment was repeated for three times and the mean values and SDs are plotted in the graph. (C) Cell morphology and localisation of Cdc42^{GTP}, indicated by CRIB-GFP signal, during vegetative growth. Representative CRIB-GFP signal images of cells of wildtype (KT5077), *ras1Δ* (5107), *ras1.G17V* (KT5082), *rga4Δ* (5551) and *rga4Δ ras1.G17V* (KT5554) are presented. Scale bar: 10 μm. (D) Quantitated CRIB-GFP signals on the cell cortex of cells presented in (C). Intensity of GFP signal on the cell cortex was measured along one of the cell tips as indicated as a magenta dotted line in the example image on the right (Scale bar: 10 μm) as stated in the Materials and Methods. 40 cells without septum were measured for each strain and the average curve from all aligned traces per strain was calculated, and displayed with respective standard error of the mean curves (dashed lines) as described in Materials and Methods.

Fig. 5. Distinct contributions of Ste4 and Ste6 to MAPK^Spk1 phosphorylation.
(A) Ste4 is essential for MAPK^Spk1 activation. MAPK^Spk1 phosphorylation status in *ste4Δ* (KT4376), *ste4Δ ras1.G17V* (KT5143) and *ste4Δ MAPKK^byr1.DD* (KT5136) at times-points 0, 8 ,16 and 24 hours post mating induction are presented. Original Western blotting membranes are presented in Fig. S3C. (B) Incapability of *ste4Δ* to cause pheromone-induced morphological change is suppressed by *MAPKK^byr1.DD* but not by *ras1.G17V*. Cell images of *ste4Δ* (KT4376), *ste4Δ ras1.G17V* (KT5143) and *ste4Δ MAPKK^byr1.DD* (KT5136) strains were taken 24 hours after induction of mating. Scale bar: 10 μm. (C) Lack of Ste6 does not result in the complete loss of MAPK^Spk1 phosphorylation. MAPK^Spk1 phosphorylation status in *ste6Δ* (KT4333), *ste6Δ ras1.G17V* (KT4998) and *ste6Δ MAPKK^byr1.DD* (KT5139) at times-points 0, 8 ,16 and 24 hours post mating induction are presented. Original Western blotting membranes are presented in Fig. S3D. (D) Incapability of *ste6Δ* to cause pheromone-induced morphological change is suppressed by *ras1.G17V* mutation but not by *MAPKK^byr1.DD* mutation. Cell images of *ste6Δ* (KT4333), *ste6Δ ras1.G17V* (KT4998) and *ste6Δ MAPKK^byr1.DD* (KT5139) strains were taken 24 hours after induction of mating. Scale bar: 10 μm.

Fig. 6. Gpa1 transduces the pheromone signalling by activating MAPK^{Spk1} and Ras1 pathways.
(A) MAPK^{Spk1} phosphorylation status in homothallic *gpa1Δ* (KT4335), *gpa1Δ ras1.G17V* (KT5023), *gpa1Δ MAPKK$^{byr1.DD}$* (KT4353) and *gpa1Δ ras1.val17 MAPKK$^{byr1.DD}$* (KT5035) at times-points 0, 8 ,12, 16 and 24 hours after mating induction. Original Western membrane is presented in Fig. S4A. (B) Cell images of the above mentioned strains at 16 hours after mating induction. All the cell images were taken and processed as in Figure 1. Scale bar is 10 μm. (C) MAPK^{Spk1} phosphorylation status in *h⁻* WT (KT4190), *h⁻ gpa1.QL* (KT5059), *h⁻ ras1.G17V* (KT4233), *h⁻ gpa1.QL ras1Δ* (KT5070) and *h⁻ MAPKK$^{byr1.DD}$* (KT4194) at times-points 0, 8 ,12 and 24 after mating induction. Note that while the activation induced in the gpa1.QL mutant was down-regulated, *MAPKK$^{byr1.DD}$* induced a constitutive activation. Original Western membrane is presented in Fig. S4B. (D) Cell images of the above strains at 12 h after induction of mating. All the cell images were taken and processed as in Figure 1. Scale bar is 10 μm.

Fig. 7. Mathematical modelling of the fission yeast pheromone pathway dynamics.
(A) Components and frameworks of the mathematical model in wildtype and signalling mutants: *ras1.G17V, Cdc42GEF^scd1Δ*, and
MAPKK^byr1.DD. Changes corresponding to each mutant are indicated as follows: Grey: removed components or interactions, orange:
increased level of activity. For the exact implementation of the mutants, see Materials and Method. The measured component, **pp**MAPK^Spk1,
is highlighted in green. (B) Measured and simulated **pp**MAPK^Spk1 activation profiles in wildtype, *ras1.G17V, Cdc42GEF^scd1Δ* and
MAPKK^byr1.DD mutants. Dashed lines: model simulations. Diamonds: experimental data presented in Fig. 1C, E, G, and Fig. 3B; error bars:
SEM. (C) GTP-loaded Ras1.G17V (1-172) directly binds to Byr2 (65-180) and Scd1 (760-872). *In vitro* GST pull-down assays of bacterially
expressed Ras1.G17V (1-172), GST-Byr2 (65-180) and GST-Scd1 (760-872) were conducted as described in materials and methods. GTP-
loaded Ras1.G17V (1-172) was found to bind to both GST-Byr2 (65-180) and GST-Scd1 (760-872). (D) Two Ras1 effectors, Byr2 and Scd1,
compete for GTP-loaded Ras1.G17V (1-172). *In vitro* GST pull-down assays of bacterially expressed Ras1.G17V (1-172) and GST-Byr2
(65-180) were conducted as in (A). Addition of Scd1 (760-872) fragment interfered with Ras1-Byr2 binding (the 4th lane). Quantitated signal

intensities of the Ras1.G17V (1-172) band in the gel are shown in the right panel. (E) Simulated ppMAPK^{Spk1} dynamics in the wildtype model at increasing concentrations of Ras1^{GTP} added *in silico* to the system. Increased Ras1^{GTP} concentration causes advanced and reduced **pp**MAPK^{Spk1} peak intensities. (F) The model fitted to the 4 strains (as above, in red) correctly predicts ppMAPK^{Spk1} dynamics in the additional 21 signalling mutant strains measured in this study.(G) Schematic diagram of the fission yeast pheromone signalling pathway, highlighting the branched pheromone sensing, and that ultimately both branches are necessary for mating.