Management performance mapping and the value of information for regional prioritization of management interventions

C. E. Buddenhagen[1234], J. Andrade Piedra[5], G. A. Forbes[5], P. Kromann[56], I. Navarrete[5], S. Thomas-Sharma[7], Y. Xing[123], R. A. Choudhury[1238], K. F. Andersen[1239], E. Schulte-Geldermann[5,10], K. A. Garrett[123]

[1]Plant Pathology Department, University of Florida, Gainesville, FL, USA

[2]Institute for Sustainable Food Systems, University of Florida, Gainesville, FL, USA

[3]Emerging Pathogens Institute, University of Florida, Gainesville, FL, USA

[4]AgResearch, Ltd., Ruakura, Hamilton, New Zealand

[5]International Potato Center, Lima, Peru, Quito, Ecuador, and Nairobi, Kenya

[6]Field Crops, Wageningen University & Research, Lelystad, The Netherlands

[7]Department of Plant Pathology and Crop Physiology, Louisiana State University Agricultural Center, Baton Rouge, LA, USA

[8]School of Earth, Environment, Marine Science, University of Texas Rio Grande Valley, TX, USA

[9]Department of Plant Pathology, Kansas State University, Manhattan, KS, USA

[10]Department of Agriculture, University of Applied Sciences Bingen, Bingen, Germany

Corresponding author: karengarrett@ufl.edu

ABSTRACT

Policymakers and donors often need to identify the locations and settings where technologies are most likely to have important effects, to increase the benefits from agricultural development or extension efforts. Higher quality information may help to target the high-payoff locations. The value of information (VOI) in this context is formalized by evaluating the results of decision making guided by a set of information compared to the results of acting without taking the information into account. We present a framework for management performance mapping that includes evaluating the VOI for decision making about geographic priorities in regional intervention strategies, in case studies of Andean and Kenyan potato seed systems. We illustrate use of Bayesian network models and recursive partitioning to characterize the relationship between seed health and yield responses and environmental and management predictors used in studies of seed degeneration. These analyses address the expected performance of an intervention based on geographic predictor variables. In the Andean example, positive selection of seed from asymptomatic plants was more effective at high altitudes in Ecuador. In the Kenyan example, there was the potential to target locations with higher technology adoption rates and with higher potato cropland connectivity, i.e., a likely more important role in regional epidemics. Targeting training to high performance areas would often provide more benefits than would random selection of target areas. We illustrate how assessing the VOI can help inform targeted development programs and support a culture of continuous improvement for interventions.

Additional keywords: agricultural development, disease, Ecuador, GIS, intervention ecology, Kenya, pest management, potato, seed degeneration, translational science, value of information, virus, yield gap

A central problem in applied spatial ecology is how to partition management efforts across

landscapes. Interventions by governments or development organizations are often designed to

increase regional crop yield, for example by improving disease management. International,

governmental, and non-governmental organizations that seek to reduce poverty, enhance food

security, and support ecosystem services, need strategies to geographically target interventions

after identifying priorities using participatory approaches with stakeholders. We propose

"management performance mapping" as a tool for translating experimental results to support

identification of geographic priorities by policy makers and donors. Management performance

mapping consists of scaling up models based on an often limited number of observations, to

visualize how specific interventions are likely to perform at a regional scale (Altieri and Nicholls

2008; van Bussel et al. 2015; van Wart, Kersebaum, et al. 2013; Grassini et al. 2015).

Management performance mapping can have a number of applications, such as providing a

summary of recommendations for extension programs, or evaluating which type of management

is most effective for a set of locations. In this paper, we focus on management performance

mapping to inform targeting of interventions to support a management component known to be

effective under some circumstances, where the goal is to identify the locations where it will be

most effective. This approach may be particularly useful in low-income countries where

smallholder farmers have fewer options, and there is interest in making a valuable new option

available through a system intervention. Management performance mapping can be implemented

to visualize the impact of proposed interventions, to improve decision-making and policymaking,

as a component of adaptive management in development (Fig. 1).

Digital or precision agriculture and species distribution models both address components

of spatial prioritization and are thus related to management performance mapping. The question

of how to optimize information use for decision making is addressed at the within-field scale in

precision agriculture (Tittonell and Giller 2013), allowing well-resourced farmers, and

potentially smallholder farmers (Cook et al. 2003), to collect and utilize spatially explicit data

sets (in near real-time) about crop performance. Inputs such as fertilizer, pesticides, and

irrigation are applied to areas of the field where they are most needed to maximize yields.

Species distribution models address the problem of optimal targeting indirectly, by

providing information about where invasive (or endangered) species, including pathogens, are

most likely to be found (Austin 2007; Hijmans and Graham 2006; Sheppard et al. 2014), often

grappling with problems in statistical inference (Stolar and Nielsen 2015) also relevant to

management performance mapping. Species distribution models are generally designed to draw

inference beyond the regions where data were collected, by estimating species niche parameters

based on maps of species occurrence or abundance throughout a species' native and introduced

range (Sutherst and Maywald 1985; Wang et al. 2017; Phillips et al. 2018; Bourdôt and

Lamoureaux 2019). Management performance mapping for disease management can

incorporate both information about which environments are conducive to pathogen and vector

reproduction, and which environments are conducive to effective management.

The value of information (VOI) concept is useful for evaluating the benefits of basing

strategies on management performance mapping. Assessing the VOI involves quantifying the

expected benefit of reducing uncertainty (Canessa et al. 2015), as described further below. VOI

analyses offer a means of both evaluating information and benefits, and assessing the role of

uncertainty when comparing management options (Hirshleifer and Riley 1979; Macauley 2006;

Canessa et al. 2015). VOI analyses compare outcomes from decision making with and without

particular units of information, taking into account the stakes for making good or bad decisions,

such as differences in yield or profit (Fig. 2). In studies of willingness-to-pay, such as farmer

willingness-to-pay for technologies, the utility functions for technologies are closely related to

the VOI (Breidert et al. 2006; Asante Bright Owusu et al. 2011; Hanemann 1991). Of course,

decision-maker willingness-to-act based on information is necessary for information valuation to

be meaningful. For example, overly confident decision-makers may not be influenced by new

information, or they may not reflect on the uncertainty that is inherent in the information

available. Many examples in the VOI literature focus on agriculture, such as the uncertainty risk

distribution for farm yield (Hirshleifer and Riley 1979), the value of weather forecasting for

farmers (Lave 1963), and risk assessment for crop futures (Danthine 1978). A related area of

application of VOI concepts is in invasion biology more generally and in conservation biology,

where decisions must also be made about where to prioritize efforts (Canessa et al. 2015;

Johnson et al. 2017; Wilson 2015). VOI analyses have so far seen little application in plant

pathology, crop epidemiology, or seed system development, where they have the potential to

improve research prioritization and decision making.

We present case studies of management performance mapping and the application of VOI

analysis that focus on smallholder management of "seed degeneration" in agricultural systems.

Seed degeneration is the reduction in yield or quality caused by an accumulation of pathogens

(often viruses) and pests in planting material over successive cycles of propagation, where

vegetatively-propagated crops deserve particular attention because of their higher risk of disease

transmission (Thomas-Sharma et al. 2016). Establishing improved seed systems is challenging,

especially in low-income countries, due in part to the many system components that must be

integrated for seed system success (Jaffee and Srivastava 1994; McGuire and Sperling 2016;

McQuaid et al. 2016; Sperling 2008; Gildemacher et al. 2009; Bentley and Vasques 1998;

Almekinders et al. 2019). In informal seed systems in low-income countries, farmers typically

use seed saved from the previous season for replanting, often leading to reduced yields, e.g., 5-

50% reduction (Devaux et al. 2010a), especially when farmers are unfamiliar with approaches

for selecting healthier seed from their fields with reduced pathogen risk. Despite the challenges

(Almekinders et al. 2019), seed system improvement has great potential for improving regional

agriculture, by providing healthier seed of better varieties, and has been a major focus of

agricultural development efforts funded by many agencies (e.g., national plant protection

agencies, The Bill and Melinda Gates Foundation, USAID, and FAO) (Jaffee and Srivastava

1994; McGuire and Sperling 2016; Almekinders et al. 1994).

Optimizing yield by reducing disease impacts, and improving seed quality, is a primary

goal of many seed system interventions. Governments and institutions with a strong focus on

science for development, such as CGIAR, work on a suite of factors linked to seed system

health. Farmer training efforts focus on options for disease management and optimal decision-

making. International development efforts for improved seed systems seek to increase farmer

access to disease-free, disease-resistant, high-quality seed, to improve farmer practices, to

implement "integrated seed health strategies" (Thomas-Sharma et al. 2016), and to implement

realistic phytosanitary thresholds (Choudhury et al. 2017). Despite concerted efforts, many

systems may revert to largely informal systems (sub-optimal seed sourced on-farm much of the

time) after interventions. For example, 98% of potato seed sources in the Andes were reported as

informal (Louwaars et al. 2013; Devaux et al. 2010a). Interventions are more likely to succeed if

they are affordable, and help farmers to be profitable (McGuire and Sperling 2013; Sperling et

al. 2013). As an example, positive selection is an on-farm management intervention that can

provide large yield benefits, e.g., 28-55% increases (mean 32%) (Gildemacher et al. 2012, 2011;

Schulte-Geldermann et al. 2012) and is often recommended as part of an integrated seed health

strategy (Thomas-Sharma et al 2016). Under positive selection, farmers select healthy appearing

plants and mark them for later harvesting of seed. Training farmers in the techniques of positive

selection can be an effective component of an integrated seed health strategy, and we use

positive selection as the example management in our case studies.

A challenge for management performance mapping – as for species distribution

modelling, digital agriculture, and most analyses designed to draw inference about larger

geographic areas – is to make the most of the available data while avoiding overinterpretation of

results. Often data about agricultural management performance exist, or can be collected inside

of existing intervention projects, but the data are collected at the scale of fields, farms or

individual plant performance measures. Multiple factors influence plant productivity apart from

management, generating uncertainty about the pay-off from management choices even where

data are relatively abundant. We discuss considerations for use of limited data. The Andean

case study addressed below illustrates both the challenge and potential value of management

performance mapping. Greater vector activity is often assumed in lower elevations, suggesting

that virus management in seed materials would be more important in these regions. Field

observations in Ecuador, though based on a limited number of fields, suggest that the reverse is

true for this case. We evaluate the VOI from management performance mapping to guide

selection of intervention locations if this counterintuitive observation is indeed representative for

the region.

Our objectives in this study are to (i) introduce and illustrate the concept of management

performance mapping and associated methods, (ii) introduce the use of VOI analysis in this

context, and (iii) illustrate the application of management performance mapping for potato seed

degeneration management by positive selection of seed in the Andes and in Kenya. We also

illustrate how analysis of likely management performance at individual sites can be combined

with other geographic considerations, such as cropland connectivity as a proxy for the role of

locations in epidemic spread for the region (Xing et al. 2020).

METHODS

We describe the steps involved in producing management performance maps (Fig. 1),

using the example of training farmers in positive selection to identify plants more likely to

produce healthy seed. Then we illustrate management performance mapping for a seed

degeneration data set from a potato seed study in Ecuador and a study of management adoption

in Kenya (Gildemacher et al. 2012; Kromann et al. 2017). As a step in preparing the Andean

management performance maps, we illustrate the application of Bayesian networks and recursive

partitioning for assessing the influence of disease, environmental factors, and management on

yield. We also evaluate the potential VOI for guiding the selection of locations in development

interventions for potato seed health in Ecuador and Kenya based on the estimated effects,

although we note that in these cases more data would be needed before proceeding to action in

the field based on these analyses. We illustrate steps 1 through 5 of the management

performance mapping pipeline (Fig. 1), while steps 6 through 8 would also be key to achieving

outcomes in the field in an adaptive management approach (Shea et al. 2014). To illustrate the

potential for combining management performance mapping (evaluated for each geographic pixel

independently) with other types of spatial processes that may include the potential roles of

locations in epidemic spread, we also provide an example of integration with a cropland

connectivity analysis (Xing et al. 2020), described below.

1) Formulate questions about the performance of a specific management strategy

across a geographic region. In these case studies, we evaluate the effects of positive selection

of farm-saved seed potato for virus disease management. In general, the identification of

management strategies for evaluation will likely be more successful if the process includes

participatory input from stakeholders. In the Andean case study, our questions are: Where would

training in positive selection likely produce the greatest benefit for yield in Ecuador and

Colombia? And how does the variety grown and the time since seed replacement influence the

benefit for yield? In the Kenyan case study, our question is: Where would training in positive

selection likely produce the greatest benefit for yield, choosing among three regions of Kenya?

2) Assemble data related to the performance of the management strategy. We use

two data sets as case studies. The first is from potato production in the Ecuadorian Andes, from a

study designed for parameter estimation for a seed degeneration model. This study (Kromann et

al. 2017) monitored seed degeneration in two potato cultivars, at three altitudes, and considered

the use of on-farm seed management options. The two cultivars were INIAP-Fripapa and

Superchola (perceived by farmers to be susceptible and resistant to degeneration, respectively).

The field trials were carried out during three cycles of planting at three sites representing three

altitudes (<2700 masl, 3000 masl and > 3400 masl, where the site <2700 masl was moved during

the course of the experiment). Twelve 49 m^2 plots were planted each year, two plots of each

cultivar at each altitude/site. In each whole plot, three types of seed management were carried

out in subplots: positive selection, roguing and random selection. The response variables

included (1) virus incidence (Potato virus X (PVX), Potato virus Y (PVY), Potato virus S (PVS),

Potato leaf roll virus (PLRV), Andean potato latent virus (APLV), and Andean potato mottle

virus (APMoV)) in plants at emergence, flowering and in tubers, evaluated using DAS-ELISA,

(2) incidence and severity of pest damage and diseases in tubers, and (3) tuber yield.

This Ecuadorian study was designed for parameter estimation for a seed degeneration

model (Thomas-Sharma et al. 2017). The main components of this model relate to seed health

(virus incidence, time/seasons since certified seed was last obtained), cultivar, environmental

factors (weather), management (seed propagation and selection) and yield data for samples of

individual potato plants (Kromann et al. 2017).   A single site represented each altitude in this

data set, so variability within a scenario can only be evaluated at the individual plant level. Lack

of replication at the field level is a limitation for management performance mapping, because an

analysis intended for providing recommendations for project implementation would be stronger

if multiple farms per altitude provided estimates of farm-to-farm variation in management

performance within an altitude range. We focus on yield data as the response in the management

performance mapping example, with potential predictors being farm altitude (across three

altitudes), seasons since certified seed was obtained, and the management performance of

positive selection compared to roguing or random seed selection as management strategies.

Climate variables – precipitation, humidity and temperature, from the WorldClim data base (Fick

and Hijmans 2017) – were also evaluated as potential predictors, but were not effective

predictors of either disease incidence or yield, probably at least in part because only three fields

per year were evaluated (data and analysis not shown).

The second data set was published data about seed health management, and positive

selection training and adoption rates in three counties in Kenya (Gildemacher et al. 2012). We

used this data to illustrate integrating information about the likelihood that farmers in a region

will adopt a technology (Gildemacher et al. 2012), another key component of intervention

success.

3) Identify predictor variables for management performance. There are many

potential predictor variables for performance indicators (Thomas-Sharma et al., in preparation)

and a wide range of methods can be used to identify important predictors, including regression

analysis, generalized linear models, and generalized additive models. We illustrate two types of

machine learning algorithms – classification and regression trees, and Bayesian network analysis

– to evaluate potential predictors, focusing on yield as the response used as a management

performance indicator. These two methods were used to identify predictor variables for the

effect of positive selection on yield for the Kromann et al. (2017) dataset. Simpler approaches to

identifying key predictors may also often prove useful in application of management

performance mapping.

Classification and regression trees. Classification and regression trees have been applied

in agricultural systems for land and soil classification, climate change impact assessment, risk

assessment, and evaluation of toxin levels and disease conduciveness in plants (Langemeier et al.

2016; Novak and LaDue 1999; Etter et al. 2006; Caley and Kuhnert 2006; Paul and Munkvold

2004; Tittonell and Giller 2013). The strength of the recursive partitioning method lies in its

ability to deal with non-linearity in data and to depict and support interpretation of the outputs in

a decision-tree format. A limitation of this method is that it may perform relatively poorly with

continuous variables or large numbers of unordered variables. We illustrate use of the rpart

package in R in the following two examples using the Kromann et al (2017) data set.

Effect of seed selection, time since seed renewal, and altitude on yield, evaluated with

recursive partitioning (Andes). We evaluated yield as the response variable, with predictors

being the use of positive selection (as opposed to roguing or random seed selection), the time

since seed renewal through purchase of certified seed (either three seasons or less than three

seasons), and the effect of altitude (across three altitudes). Because altitude is available as a

potential geographic predictor variable for the region, it is a candidate for extrapolating analysis

of the performance of positive selection to a wider area in Step 4 (Fig. 1).

Effect of potato cultivar and its interactions on yield, evaluated with recursive

partitioning (Andes). In a previous study of a grower cooperative in Tungurahua, Ecuador,

Superchola (one of the most important potato varieties in Ecuador) and INIAP-Fripapa were sold

and grown at a ratio of approximately 2:1 by volume (Buddenhagen et al. 2017). Our analysis of

the Kromann et al. dataset (2017) also focuses on these two varieties. We evaluate the effects of

cultivar, altitude, and management by estimating mean per-plant yields across treatment

combinations, and by using recursive partitioning in rpart.

Bayesian networks. Bayesian networks (Therneau et al. 2010) have been applied in

natural resource management systems for applications such as vegetation classification, optimal

decision making, disease management, adaptive management of wildlife habitat, and expert

elicitation (Geenen and Van Der Gaag 2005; Aguilera et al. 2011; Kristensen and Rasmussen

2002; Perez-Ariza et al. 2012; Howes et al. 2010). A Bayesian network is a directed, acyclic

graph whose nodes represent predictor variables and links represent dependencies. The

relationships between variables are quantified in conditional probability tables, where the set of

all tables together represents the full joint distribution. Important strengths of the Bayesian

network method include its ability to infer probabilistic relationships among many variables

simultaneously. The network structure can be set manually by the user or learned from the data

using a variety of algorithms. In the case of exact estimation algorithms, it is possible to set

values for any combination of nodes and produce new posterior probabilities for each variable in

the network. A limitation of this method is the cost of some of the most advanced Bayesian

network software. In addition, combinations of continuous and categorical data can be

problematic for some commonly-used Bayesian network algorithms (Aguilera et al. 2011). Tools

available for Bayesian network analysis include BI-CAMML, Hugin and Netica (Aguilera et al.

2011). R packages include bnlearn, gRain and pcalg (Nagarajan et al. 2013). We selected Netica

for this illustration because it is relatively affordable, the algorithms it uses allow for immediate

updating of conditional probabilities based on selected levels for variables, it has a powerful

graphical interface, and it is widely used in ecological and environmental analyses (Aguilera et

al. 2011).

294   ***Effect of positive selection on yield, evaluated in Bayesian networks (Andes).*** The

295   benefit of positive selection (in the third cropping cycle after certified seed purchase) was

296   evaluated in a Bayesian network in Netica. Netica's Tree-Augmented Naive Bayes (*TAN*)

297   classifier algorithm was used to estimate the conditional probability tables and the network

298   structure. From the conditional probability tables we estimated yields above (7.7 t/ha) and below

299   (3.2 t/ha) the threshold altitude identified in the analysis: 2895 m.a.s.l.

300   ***A simple analysis of regional differences in adoption of training recommendations***

301   ***(Kenya).*** In this case study, we evaluated regional differences in farmers' adoption of positive

302   selection after training (Table 1), reported by Gildemacher et al. (2012) as follows for three

303   Kenyan counties: Nakuru 46%, Nyandarua 19%, and Narok 18%.

304   **4) Estimate management performance across the geographic study region.** For

positive selection of on-farm seed in the Andes, we selected for analysis and extrapolation a

major potato growing region stretching from southern Ecuador to southern Colombia. Using

potato production geographic data layers from SPAM 2005 v3.2 Global Data (IFPRI and IIASA

2016) we focused on pixels with >200 ha potato production per pixel (where a pixel represents 5-

arc minutes, approximately 10,000 ha). Here 51% of potato production is above 2895 m (the

altitude threshold identified in the analyses above) based on SPAM estimates (You et al. 2012).

The resulting management performance map will indicate that these regions would be priorities

for targeting training in positive selection if decisions are based solely on this analysis of the data

from Kromann et al. (2017).

For positive selection of on-farm seed in Kenya, rather than extrapolating the estimates of

management performance for positive selection, we simply compare the relative performance of

the counties (Table 1). The resulting management performance map will indicate prioritization

among these counties if decisions for targeting positive selection training are based solely on the

data from Gildemacher et al. (2009).

5) Evaluate the value of information for management intervention or policy. We

assessed the value of information for decisions about where to invest management interventions,

for a scenario where the estimates from Kromann et al. (2017) do correctly represent the region.

For the purposes of this illustration, we considered cases where decision makers either have or

do not have information about the geographic differences in management performance (Fig. 2).

In the absence of information, they might select any location for management with equal

probability. An estimate of the value of information would be the difference in the benefit of

investment for locations selected based on the information ("informed location selection"), and

the benefit for locations selected randomly ("uninformed location selection"). For example,

informed site selection might direct site selection to farms above or below the altitude threshold

identified in analysis (e.g., 2895 m.a.s.l. identified in Bayesian network analysis), depending on

whether higher or lower elevations provide greater benefits. In the case where decision makers

have a prior belief that is not supported by the data, and it is in fact an incorrect belief, the value

of information would be the difference between investment outcomes based on the

misconception ("misinformed location selection") and outcomes based on informed investments.

For example, there could be a prior belief that a particular pathogen will be more prevalent at

lower elevations, due to a higher abundance of vectors, resulting in a prior belief that positive

selection would be more important at lower elevations. We evaluated uninformed, informed, and

misinformed management choices related to spatially distributed differences in yield, disease,

cultivar and the rates with which best practices are adopted.

339   ***VOI for positive selection targeting in the Andes.*** Comparison of yield improvements

due to positive selection training – with and without the information from Kromann et al. (2017)

– has as a first step determining how common each trait combination is in the landscape being

considered. Then the probability of randomly including a particular trait combination can be

estimated. The proportion of Ecuadorian farmers using certified seed was previously reported at

2% (Devaux et al. 2010b), and many farmers lack access to certified seed, though for some

organized farming groups the proportion using certified or quality-declared seed can be as high

as 46% (Buddenhagen et al. 2017). We take the frequency of farms in this landscape being

planted with certified seed ("new seed") at any given time as being approximately 2% (so that a

farm drawn at random has probability $p = 0.02$ of being planted with certified seed, although this

is an approximation because it is generally the wealthier farmers, government programs, or non-

governmental organizations who acquire certified seed). Farm altitude, based on the geographic

analysis described above for higher density potato regions, is above the altitude threshold

identified in recursive partitioning approximately 51% of the time.   For simplicity, we treat the

potato cultivar planted as 33% INIAP-Fripapa and 66% Superchola, based on estimates for the

province of Tungurahua from Buddenhagen et al. (2017).

VOI for targeting positive selection in Kenya. The average benefit of positive selection

was reported by Gildemacher et al. (2012) as 3.4 tons per ha (~$350 per ha). This translated to a

per-household benefit of $156 per season for a farm of average size for the region. Meanwhile,

the cost of training was $38 per farmer. In this case, the expected first-year benefit was $44 per

household when training occurred in a randomly selected region (without regard to adoption

rate) (Table 1). We compare this outcome to the outcome using information about frequencies of

adoption.

Integration with another criterion for selecting priority locations: cropland

connectivity (Ecuador and Colombia). The data layer of estimated management performance

is one important factor for deciding where to prioritize management efforts. The management

performance map developed up to this stage is generated pointwise, in that it treats each location

(point) as independent from other locations. However, some locations will have more important

roles in epidemics than others, due to factors such as the location's position in spatial epidemic

networks. Thus, targeting some locations will have more important effects to slow regional

epidemics, for seed degeneration pathogens such as viruses that tend to be spread from one field

to another. We also evaluated the layer of management performance estimates for positive

selection integrated with a data layer of the potato "cropland connectivity risk index", a measure

of the likely importance of locations for spatial movement through potato growing areas (Xing et

al. 2020; Margosian et al. 2009), as described below.

The potato cropland connectivity analysis was based on the potato crop harvested area

data from SPAM 2005 v3.2 Global Data (IFPRI and IIASA 2016). This data has pixel resolution

5-arc min, and those cells with harvested area greater than 200 ha were included in the cropland

connectivity risk analysis (Xing et al., 2020). As described in more detail in Xing et al. (2020),

the distance between pairs of cells was evaluated in a sensitivity analysis for both inverse power-

law models (parameters 0.5, 1, and 1.5) and negative exponential models (parameters 0.05, 0.1,

0.2, 0.3, and 1). Three network link thresholds (0.001, 0.0001, 0.00001) were applied separately

to each adjacency matrix to represent three different scenarios in the network analysis in a

sensitivity analysis. A cropland connectivity risk index (CCRI) was calculated as the scaled

weighted sum of betweenness centrality, node strength, the sum of nearest neighbours' node

degrees, and eigenvector centrality, as in Xing et al. (2020). For each realization in the sensitivity

analysis, the mean CCRI was evaluated across the 24 parameter combinations. This mean CCRI

was then mapped in combination with the map of management performance estimates, to

identify locations important both for the CCRI (indicating a potentially important epidemic role)

and for management benefits from positive selection.

RESULTS

3) Identifying predictor variables for management performance

Positive selection and yield for Andean potato.

In the recursive partitioning analysis, higher per plant yields were generally obtained

from INIAP-Fripapa (compared to Superchola) in the first two years after the certified seed was

purchased, the highest yields being obtained for altitudes over 3278 m (Fig. 3). The highest

yields for Superchola were found above 2895 m altitude. If a farmer can afford to replace seed

more frequently, and the farm is over 3200 m, INIAP-Fripapa yielded higher than Superchola

(and their value in the market was comparable in 2016 – 0.29 and 0.33 USD per kg,

respectively).

Plants three years post-certified seed purchase yielded 33% less per plant. The benefits of

positive selection allow yields to approach the average yields for recently purchased certified

seed. There were no differences with respect to cultivar three years after certified seed purchase,

suggesting positive selection was equally valuable in both varieties for seed that had gone

through more than two planting cycles.

The Bayesian network analysis indicated that high yielding plants were found more

commonly in plots where first generation certified seed was used, at higher altitudes, for the

INIAP-Fripapa cultivar, and where there was a lower minimum temperature and higher rainfall

six months after planting, as well as low levels of PVX, PLRV, and PVY (Fig. 4). Positive

selection was less likely to be the management implemented if the yield was low. All the viruses

except PVY, and PLRV for low yield plants, were more likely to be absent (frequency = 0) than

present. Each virus was relatively more likely to be absent if a plant was in the high yield

category compared to plants in the low yield category (Fig. 4). The uncertainty was high

compared to the observed values, indicating that another cycle of data collection would be

needed before implementing project plans based on this data.

Adoption of positive selection for Kenyan potato. This analysis was based on the

probability of adoption of positive selection, where higher adoption rates resulted in a higher

payoff for intervention investment. Adoption rates were 46, 19 and 18% in three counties (Table

1) (Gildemacher et al. 2012, 2011). Thus, based on this measure alone, selection of the county

with 46% adoption rate would approximately double the benefits obtained from a training

intervention focused only where there was 18% or 19% adoption.

4) Estimating management performance across the geographic study region, and

integrating with cropland connectivity estimates.

Applying models to a map of the relevant region and integrating data layers for

Andean potato. The mapped estimates of the management performance of positive selection for

Andean potato yield (based on altitude) and the locations where potato cropland connectivity risk

was highest based on the CCRI (Fig. 5) were combined to identify locations both (a)

independently likely to have the best management outcomes, and (b) likely important for

regional management of disease spread (high CCRI). Locations that meet both criteria were

observed near the border of Ecuador and Colombia, and near Ambato and Riobamba in Ecuador

(Fig. 5).

Technology adoption rates for three counties in Kenya. The county with the highest

adoption rate for positive selection, Nakuru (Table 1), was intermediate in terms of the cropland

connectivity index (Fig. 6). The cropland connectivity index was high for multiple locations in

and near Nyandarua county.

5) Evaluate the value of information for management intervention or policy.

VOI for targeted implementation of positive selection (yield as response, Bayesian

networks to identify predictors) For equivalent farm sizes above and below 2895 m

(representing 49% of the area), we estimated the benefit of training under uninformed (random)

site selection by using the weighted mean of the benefit above and below 2895 m.a.s.l.

(representing 51% of the cultivated area), which is 6.5 t/ha ($0.51 * 7.7 + 0.49 * 3.2 = 6.5$).

The estimated benefit under informed site selection, selecting locations above 2895 m, is

7.7 t/ha – a difference of 1.2 t/ha from random site selection. Under misinformed site selection,

if the assumption was that positive selection provides more benefits at low altitude (perhaps due

to greater pathogen load), then the benefit is 3.2 tons per ha, 4.5 tons per ha less than the optimal

allocation, and 3.3 tons per ha less than the uninformed (random) site selection option.

VOI for targeted implementation of positive selection (yield as response, recursive

partitioning to identify predictors) Assuming that positive selection training targeted farmers

randomly with respect to the observed frequency of the categories, the weighted mean benefit of

positive selection would be 8.7 tons per ha (Fig. 3). Preferentially targeting sites at high altitude

(but sampling randomly with respect to seed age and cultivar) provides higher benefits (9

tons/ha), otherwise targeting low altitude sites provides lower returns at (8.3 tons/ha). Targeting

farmers who plant farm-saved seed, three years since purchase of certified seed, provides little

benefit: 8.8 tons/ha compared to random targeting of farmers (under the scenario where use of

certified seed is rare at 2%). Targeting the 2% that do use new seed provides a benefit of 3.1

tons/ha, although these more successful farmers may not need interventions. By far the greatest

benefit is provided by targeting farmers that grow INIAP-Fripapa (benefit of 10.9 tons/ha) as

opposed to Superchola (benefit of 7.6 tons/ha).

Regional differences in adoption of training recommendations in Kenya. In the

example of Kenyan potato seed technology adoption, targeted selection of high adoption rate

areas for training (where the per-farmer benefit was $72) would increase the return by $28 per

farmer trained (Table 1). Realized benefits would vary depending on the farm size.

DISCUSSION

The examples given here show how management performance mapping can be used to

target sites for project interventions. We illustrate how identifying locations where positive

selection of on-farm saved seed has the highest performance for increasing yield (Andes) or the

highest adoption rates (Kenya) can provide substantial regional benefits. While the particular

examples here would require additional information to advance to the field with confidence

(steps 6 through 8 in Fig. 1), they illustrate how a management performance mapping framework

can be implemented An NGO or government extension agency with limited resources could use

such an approach to better target rural development interventions. We compared uninformed and

informed allocation of resources, for scenarios where the management performance models are

correct (i.e., scenarios where the data perfectly represent the region of interest), to assess the

value of the information used for targeting interventions. In a simple scenario, using Bayesian

network analysis to identify altitude as a management performance predictor, we found that the

benefit of positive seed selection was highest (an increase of 4.5 tons per ha) at high altitudes,

and uninformed allocation of farmer training would provide a net benefit of 1.2 tons per ha less

than targeted training. Incorrectly assuming that better outcomes for positive selection would be

obtained at lower altitudes, perhaps because aphid vector abundances were thought to be higher,

would have produced 3.3 and 4.5 tons per ha less for random site selection and optimal

allocation, respectively, for this scenario (Bertschinger et al. 2017).

Along with the magnitude of management effects on yield, adoption rates are also key to

successful interventions (Parsa et al. 2014). Based on the data about the benefits of adoption

rates of positive selection of seed in Kenya, we found that unless adoption rates were higher than

24%, the first-year benefit per household would not exceed the $38 per farmer cost of training

(although, presumably, the benefits would continue to accrue in subsequent years). Also, random

allocation of training effort would only yield a $44 benefit (over the cost of the training) per

household. Gildemacher et al. (2017) also point out that adoption rates were lower in drought

years, suggesting that prediction of adoption rates could be difficult if based on regional patterns

in a single year. Observed adoption rates may vary in predictable ways based on disease

incidence in the current or previous season, in-season weather conditions, language spoken,

literacy, cultural differences between trainer and trainee, wealth or other factors. When these

relationships are understood and spatial data are available for key predictor variables for

adoption, these variables could form a part of selection criteria for farmer training initiatives (and

the approach to the training could be altered to improve adoption rates).

Our example decision, deciding where to implement training for improved disease

management, represents a class of decisions where there is confidence that the activity will

provide a benefit. Management performance mapping is applied to guide implementation to

locations where there is some evidence that the benefit will be greater than in other locations.

For this class of decisions, the risk is often low that limited data is "worse than no data at all". In

the management performance mapping context, the null hypothesis is often that the benefit of

implementation will be the same in all locations. In evaluating where there is evidence to reject

this hypothesis, there is not a strong motivation to particularly avoid false positives or Type I

error (rejecting a null hypothesis when the null hypothesis is true), because a false negative or

Type II error (failing to reject a null hypothesis when the null hypothesis is false) is arguably just

as bad. The main risk of "bad data" would be from data with a strong bias that would lead to

misinformed decisions. The cost of "bad data" may also go up if the logistical costs (of

transport, communications, etc.) of targeting locations incorrectly identified is higher than

targeting locations at random or selecting locations based on convenience. There is the potential

for these risks to be managed in real time during project implementation by incorporating

distributed or "big" data sourced from farmer phone apps, rapid disease detection methods or

citizen science initiatives (Nakato et al. 2016; Boykin et al. 2019). In our example data from

Ecuador, our only estimate of uncertainty within a scenario was based on variability among

individual plants, while a person making decisions about regional priorities would strongly

prefer to have information about farm-to-farm variability within each scenario. One of the

potential applications of VOI analysis is to determine whether collecting more or better data

about management performance is justified (Ades et al. 2004), not just for the sake of more

statistical power in general, but because the information improves farmer decision-making under

a realistic range of conditions.

Two key factors for adoption of positive selection are market price and the varieties

grown in a region, in terms of their rates of seed degeneration. The number of seasons over

which positive selection is adopted is also an important factor helping to determine the return on

investment in training. The study of adoption rates in Kenya was performed once and may be

limited to the circumstances at the time of sampling. At the time of the training in positive

selection, there was a shift in Nyandarua from the variety Tigoni to Shangi, so there might have

been more interest in acquiring the new variety than in improving old seed stock (Okello et al.

2018; Kaguongo et al. 2008). Nyandarua also has an apparent role in spread of potato cyst

nematode in the region (Mburu et al. 2018; Mwangi et al. 2015), along with bacterial wilt

problems, which may have made positive selection less effective there. Potato farming has a

longer history in Nyandarua. In Narok, potatoes are less important and conditions are less

favorable, with the main potato variety grown being Dutch Robijn. In Nakuru, with the highest

adoption rate, more varieties are grown and potato farming is more recent, in generally good

growing conditions. These differing factors in the three counties, combined with changes over

time such as the occurrence of droughts, can modify the likelihood of adoption of positive

selection. The yield improvements from positive selection in Kenya, averaging 30% (Schulte-

Geldermann et al. 2012), make it an attractive technology for development investments. In this

system, there is the possibility of farmers actually improving seed quality over time, rather than

simply slowing decline, and understanding this potential could also support decisions.

Formulating a strategy for targeting positive selection in Kenya would be strengthened by new

data about how the differences among these and other counties influence the current likelihood

of technology adoption.

Combining data layers for evaluating optimal intervention strategies can provide more

insight, along with potential challenges due to uncertainty and different spatial resolutions

(Sutton and Armsworth 2014). Evaluating the risk of disease due to cropland connectivity (Xing

et al. 2020) in combination with independent location characteristics can position the analysis in

the larger context of disease management for the region. Cropland connectivity may change

over the course of the year, as potato is present or absent. For example, in Kenya some parts of

Nyandarua county, such as Njambini and Oljororok, have potato in the field the entire year, due

to the availability of groundwater coming from the Aberdare range during the dry season. Three

crops a year are very common, and likely affect pest and disease cycles. Consistently highly

connected locations may be more important targets for achieving impacts on regional epidemic

spread, although there is also the potential for highly connected locations experiencing high

inoculum loads to respond poorly to some types of management. A broader systems analysis –

for example, impact network analysis (Garrett et al. 2018) which integrates across management

performance, socioeconomic or innovation networks (Fritsch and Kauffeld-Monz 2010; Leeuwis

and Aarts 2011), and biophysical networks such as epidemic networks – can aid in identifying

intervention locations that prioritize across multiple goals. For farmer decision making, flexible

decision rules and reducing variability of risk may be priorities (Bert et al. 2006; Andrieu et al.

2015).

Crop and epidemic models may provide valuable data layers if they incorporate spatially

mappable variables. Estimating the effects of management strategies, such as variety

deployment, depends on understanding the yield potential, perhaps based on a combination of

weather or climate data and data about regional management practices (van Wart, van Bussel, et

al. 2013; Araya et al. 2010; Reynolds et al. 2018). Disease modeling may be used to evaluate the

likely effects of management, such as addressing the problem of seed degeneration (Thomas-

Sharma et al. 2017; Jones et al. 2010). Spatial epidemic components such as seed trade networks

(McQuaid et al. 2017; Buddenhagen et al. 2017; Andersen et al. 2019) may also be valuable

components of more refined management performance mapping.

Management performance mapping to identify target locations for interventions, and the

VOI analysis therein, is potentially useful for many problems in agriculture or intervention

ecology. There is a constellation of approaches that address related goals. Yield gap analyses that

incorporate maps can address some of the same goals as management performance mapping

(Schulthess et al. 2013; Silva et al. 2017; Lobell et al. 2015, 2009; van Ittersum et al. 2016;

Grassini et al. 2015; van Bussel et al. 2015). For example, yield gap analysis attempts to identify

the most important factors that influence yield, especially factors that are controllable. The focus

of management performance mapping for intervention targeting, however, is on providing spatial

information about the intervention impact of management options. Management performance

maps would ideally incorporate and account for interacting human dimensions (e.g., learning,

financial liquidity, capital, institutions; Arneth et al. 2014), as well.

The benefits from management performance mapping may be enhanced if maps and VOI

calculations are updated with new information sources over time in an adaptive management

scheme (Shea et al. 2014; Bennett et al. 2018). Empirical data can be supplemented with

models, expert opinion, and local knowledge (Petsakos et al. 2018; Tulloch et al. 2014) to

understand changes in factors such as pesticide resistance, new varieties, and new management

such as irrigation. Projects may expand due to new stakeholder priorities and new situations on

the ground. As new pests and pathogens enter a region (Bebber et al. 2014), they will likely

necessitate alterations in current best management practices. For example, in Ecuador potato

purple top disease has become a major problem (Caicedo et al. 2015; Castillo-Carrillo et al.

2018) since the Ecuadorian experiments reported here were performed. 'Candidatus Liberibacter

solanacearum', associated with zebra chip disease, and its vector the tomato potato psyllid,

Bactericera cockerelli, have also been reported in Ecuador (Caicedo et al. 2020; Castillo-Carrillo

et al. 2019). New strategies for potato best management practices in Ecuador will need to

address purple top and the risk of zebra chip, including uncertainty about causal agents. In other

scenarios, multiple outcomes may be important, such as a combination of benefits and

environmental costs of management (Laurance et al. 2014), pesticide effects on non-target

species in disease management, or conservation management focusing on both biodiversity

hotspots and locations with keystone species (Smith et al. 2007). Our examples addressed

management performance mapping with performance defined in terms of the mean performance

observed. Other potential criteria for selecting regions for investment might emphasize different

priorities (Table 2). For example, effective altruism concepts can be used to target stakeholders

to maximize research benefit (Garrett et al. 2020).

In summary, management performance mapping provides a process to extrapolate from

available data to make evidence-based decisions about where to invest in disease and crop

management or training initiatives. Scenario analyses to support decision making (Wiebe et al.

2018) can build on the framework developed here.

ACKNOWLEDGMENTS

This research was undertaken as part of, and funded by, the CGIAR Research Program on Roots,
Tubers and Bananas (RTB) and supported by CGIAR Trust Fund contributors
https://www.cgiar.org/funders/. We also appreciate support by the CGIAR Research Program on
Climate Change and Food Security (CCAFS), Bill and Melinda Gates Foundation grant
OPP1080975, USDA NIFA grant 2015-51181-24257, USDA APHIS grant 11-8453-1483-CA,
the USAID Feed the Future Haiti Appui à la Recherche et au Développement Agricole (AREA)
project grant AID-OAA-A-15-00039, US NSF Grant EF-0525712 as part of the joint NSF-NIH
Ecology of Infectious Disease program, US NSF Grant DEB-0516046, and the University of
Florida.
617

LITERATURE CITED

Ades, A. E., Lu, G., and Claxton, K. 2004. Expected value of sample information calculations in medical decision modeling. Med Decis Making. 24:207–227.

Aguilera, P. A., Fernández, A., Fernández, R., Rumí, R., and Salmerón, A. 2011. Bayesian networks in environmental modelling. Environmental Modelling & Software. 26:1376–1388.

Almekinders, C. J. M., Louwaars, N. P., and de Bruijn, G. H. 1994. Local seed systems and their importance for an improved seed supply in developing countries. Euphytica. 78:207–216.

Almekinders, C. J. M., Walsh, S., Jacobsen, K. S., Andrade-Piedra, J. L., McEwan, M. A., de Haan, S., et al. 2019. Why interventions in the seed systems of roots, tubers and bananas crops do not reach their full potential. Food Sec. 11:23–42.

Altieri, M. A., and Nicholls, C. I. 2008. Scaling up agroecological approaches for food sovereignty in Latin America. Development. 51:472–480.

Andersen, K. F., Buddenhagen, C. E., Rachkara, P., Gibson, R., Kalule, S., Phillips, D., et al. 2019. Modeling Epidemics in Seed Systems and Landscapes To Guide Management Strategies: The Case of Sweet Potato in Northern Uganda. Phytopathology. 109:1519–1532.

Andrieu, N., Descheemaeker, K., Sanou, T., and Chia, E. 2015. Effects of technical interventions on flexibility of farming systems in Burkina Faso: Lessons for the design of innovations in West Africa. Agricultural Systems. 136:125–137.

Araya, A., Keesstra, S. D., and Stroosnijder, L. 2010. A new agro-climatic classification for crop suitability zoning in northern semi-arid Ethiopia. Agricultural and Forest Meteorology. 150:1057–1064.

Arneth, A., Brown, C., and Rounsevell, M. D. A. 2014. Global models of human decision-making for land-based mitigation and adaptation assessment. Nature Clim Change. 4:550–557.

Asante Bright Owusu, Otoo Emmanuel, Wiredu Alexander Nana, Acheampong Patricia, Osei-Adu Jonas, and Nsiah-Frimpong Benedicta. 2011. Willingness to adopt the vine multiplication technique in seed yam production in the forest savanna transition agro-ecological zone, Ghana. Journal of Development and Agricultural Economics. 3:710–719.

Austin, M. 2007. Species distribution models and ecological theory: A critical assessment and some possible new approaches. Ecological Modelling. 200:1–19.

Bebber, D. P., Holmes, T., and Gurr, S. J. 2014. The global spread of crop pests and pathogens: The global spread of crop pests and pathogens. Global Ecology and Biogeography. 23:1398–1407.

Bennett, J. R., Maxwell, S. L., Martin, A. E., Chadès, I., Fahrig, L., and Gilbert, B. 2018. When to monitor and when to act: Value of information theory for multiple management units and limited budgets ed. Hedley Grantham. J Appl Ecol. 55:2102–2113.

Bentley, J. W., and Vasques, D. 1998. The seed potato system in Bolivia: organisational growth and missing links. Agricultural Research and Extension Network. :1–12.

Bert, F. E., Satorre, E. H., Toranzo, F. R., and Podestá, G. P. 2006. Climatic information and decision-making in maize crop production systems of the Argentinean Pampas. Agricultural Systems. 88:180–204.

Bertschinger, L., Bühler, L., Dupuis, B., Duffy, B., Gessler, C., Forbes, G. A., et al. 2017.
Incomplete infection of secondarily infected potato plants–an environment dependent
underestimated mechanism in plant virology. Frontiers in Plant Science. 8:74.
Bourdôt, G. W., and Lamoureaux, S. L. 2019. *Abutilon theophrasti* –a comparison of two climate
niche models. New Zealand Journal of Agricultural Research. :1–12.
Boykin, Sseruwagi, Alicai, Ateka, Mohammed, Stanton, et al. 2019. Tree lab: portable genomics
for early detection of plant viruses and pests in Sub-Saharan Africa. Genes. 10:632.
Breidert, C., Hahsler, M., and Reutterer, T. 2006. A review of methods for measuring
willingness-to-pay. Innovative Marketing. 2:8–32.
Buddenhagen, C. E., Hernandez Nopsa, J. F., Andersen, K. F., Andrade-Piedra, J., Forbes, G. A.,
Kromann, P., et al. 2017. Epidemic network analysis for mitigation of invasive pathogens
in seed systems: Potato in Ecuador. Phytopathology. 107:1209–1218.
van Bussel, L. G. J., Grassini, P., Van Wart, J., Wolf, J., Claessens, L., Yang, H., et al. 2015.
From field to atlas: Upscaling of location-specific yield gap estimates. Field Crops
Research. 177:98–108.
Caicedo, J., Crizón, M., Pozo, A., Cevallos, A., Simbaña, L., Rivera, L., et al. 2015. First report
of "*Candidatus* Phytoplasma aurantifolia" (16SrII) associated with potato purple top in
San Gabriel-Carchi, Ecuador. New Dis. Rep. 32:20.
Caicedo, J. D., Simbaña, L. L., Calderón, D. A., Lalangui, K. P., and Rivera-Vargas, L. I. 2020.
First report of 'Candidatus Liberibacter solanacearum' in Ecuador and in South America.
Australasian Plant Dis. Notes. 15:6.
Caley, P., and Kuhnert, P. M. 2006. Application and evaluation of classification trees for
screening unwanted plants. Austral Ecology. 31:647–655.
Canessa, S., Guillera-Arroita, G., Lahoz-Monfort, J. J., Southwell, D. M., Armstrong, D. P.,
Chadès, I., et al. 2015. When do we need more data? A primer on calculating the value of
information for applied ecologists ed. Olivier Gimenez. Methods in Ecology and
Evolution. 6:1219–1228.
Castillo-Carrillo, C., Fu, Z., and Burckhardt, D. 2019. First record of the tomato potato psyllid
Bactericera cockerelli from South America. Bulletin of Insectology. 72:85–91.
Castillo-Carrillo, C., Paltrinieri, S., Bustamante, J. B., and Bertaccini, A. 2018. Detection and
molecular characterization of a 16SrI-F phytoplasma in potato showing purple top disease
in Ecuador. Australasian Plant Pathol. 47:311–315.
Choudhury, R. A., Garrett, K. A., Klosterman, S. J., Subbarao, K. V., and McRoberts, N. 2017.
A framework for optimizing phytosanitary thresholds in seed systems. Phytopathology.
107:1219–1228.
Cook, S., O'Brien, R., Corner, R., Oberthur, T., Stafford, J., and Werner, A. 2003. Is precision
agriculture irrelevant to developing countries. Precision Agriculture. :115–120.
Danthine, J.-P. 1978. Information, futures prices, and stabilizing speculation. Journal of
Economic Theory. 17:79–98.
Devaux, A., Ordinola, M. E., Hibon, A., and Flores, F. A. 2010a. *El sector papa en la región*
andina: Diagnóstico y elementos para una visión estratégica (Bolivia, Ecuador y Perú).
International Potato Center.
Devaux, A., Ordinola, M. E., Hibon, A., and Flores, F. A. 2010b. *El sector papa en la región*
andina: Diagnóstico y elementos para una visión estratégica (Bolivia, Ecuador y Perú).
International Potato Center.

Etter, A., McAlpine, C., Wilson, K., Phinn, S., and Possingham, H. 2006. Regional patterns of
agricultural land use and deforestation in Colombia. Agriculture, Ecosystems &
Environment. 114:369–386.
Fick, S. E., and Hijmans, R. J. 2017. Worldclim 2: new 1-km spatial resolution climate surfaces
for global land areas: new climate surfaces for global land areas. Int. J. Climatol.
37:4302–4315.
Fritsch, M., and Kauffeld-Monz, M. 2010. The impact of network structure on knowledge
transfer: an application of social network analysis in the context of regional innovation
networks. The Annals of Regional Science. 44:21–38.
Garrett, K. A., Alcalá-Briseño, R. I., Andersen, K. F., Brawner, J., Choudhury, R. A., Delaquis,
E., et al. 2020. Effective altruism as an ethical lens on research priorities. Phytopathology.
110:708–722.
Garrett, K. A., Alcalá-Briseño, R. I., Andersen, K. F., Buddenhagen, C. E., Choudhury, R. A.,
Fulton, J. C., Hernandez Nopsa, J. F., Poudel, R., and Xing, Y. 2018. Network analysis:
A systems framework to address grand challenges in plant pathology. Annual Review of
Phytopathology 56:559-580.
Geenen, P. L., and Van Der Gaag, L. C. 2005. Developing a Bayesian network for clinical
diagnosis in veterinary medicine: from the individual to the herd. In *Proceedings of the*
Third Bayesian Modelling Applications Workshop; Edinburgh, Citeseer.
Gildemacher, P. R., Demo, P., Barker, I., Kaguongo, W., Woldegiorgis, G., Wagoire, W. W., et
al. 2009. A description of seed potato systems in Kenya, Uganda and Ethiopia. American
Journal of Potato Research. 86:373–382.
Gildemacher, P. R., Leeuwis, C., Demo, P., Borus, D., Schulte-Geldermann, E., Mundia, P., et al.
2012. Positive selection in seed potato production in Kenya as a case of successful
research-led innovation. International Journal of Technology Management & Sustainable
Development. 11:67–92.
Gildemacher, P. R., Schulte-Geldermann, E., Borus, D., Demo, P., Kinyae, P., Mundia, P., et al.
2011. Seed potato quality improvement through positive selection by smallholder farmers
in Kenya. Potato Research. 54:253–266.
Grassini, P., van Bussel, L. G. J., Van Wart, J., Wolf, J., Claessens, L., Yang, H., et al. 2015.
How good is good enough? Data requirements for reliable crop yield simulations and
yield-gap analysis. Field Crops Research. 177:49–63.
Hanemann, W. M. 1991. Willingness to pay and willingness to accept: how much can they differ?
The American Economic Review. 81:635–647.
Hijmans, R. J., and Graham, C. H. 2006. The ability of climate envelope models to predict the
effect of climate change on species distributions. Global Change Biology. 12:2272–2281.
Hirshleifer, J., and Riley, J. G. 1979. The analytics of uncertainty and information-an expository
survey. Journal of Economic Literature. 17:1375–1421.
Howes, A. L., Maron, M., and Mcalpine, C. A. 2010. Bayesian networks and adaptive
management of wildlife habitat: Bayesian networks and adaptive management.
Conservation Biology. 24:974–983.
IFPRI, and IIASA. 2016. *Global Spatially-Disaggregated Crop Production Statistics Data for*
2005. Version 3.2. International Food Policy Research Institute (IFPRI) and International
Institute for Applied Systems Analysis (IIASA). Available at:
https://doi.org/10.7910/DVN/DHXBJX.

van Ittersum, M. K., van Bussel, L. G. J., Wolf, J., Grassini, P., van Wart, J., Guilpart, N., et al. 2016. Can sub-Saharan Africa feed itself? Proceedings of the National Academy of Sciences. 113:14964–14969.

Jaffee, S., and Srivastava, J. 1994. The roles of the private and public sectors in enhancing the performance of seed systems. The World Bank Research Observer. 9:97–117.

Johnson, F. A., Smith, B. J., Bonneau, M., Martin, J., Romagosa, C., Mazzotti, F., et al. 2017. Expert elicitation, uncertainty, and the value of information in controlling invasive species. Ecological Economics. 137:83–90.

Jones, R. A. C., Salam, M. U., Maling, T. J., Diggle, A. J., and Thackray, D. J. 2010. Principles of predicting plant virus disease epidemics. Annu. Rev. Phytopathol. 48:179–203.

Kaguongo, W., Gildemacher, P., Demo, P., Wagoire, W., Kinyae, P., Andrade, J., et al. 2008. *Farmer practices and adoption of improved potato varieties in Kenya and Uganda.* Lima, Peru: International Potato Center.

Kristensen, K., and Rasmussen, I. A. 2002. The use of a Bayesian network in the design of a decision support system for growing malting barley without use of pesticides. Computers and Electronics in Agriculture. 33:197–217.

Kromann, P., Andrade-Piedra, J. L., Navarrete, I, Taipe, A., and Gómez, J. 2017. Dataset for: Potato seed degeneration in Ecuador. International Potato Center Dataverse, V2. Available at: http://dx.doi.org/10.21223/P3/3CT90C [Accessed December 1, 2017].

Langemeier, C. B., Robertson, A. E., Wang, D., Jackson-Ziems, T. A., and Kruger, G. R. 2016. Factors affecting the development and severity of Goss's bacterial wilt and leaf blight of corn, caused by *Clavibacter michiganensis* subsp. *nebraskensis*. Plant Disease. 101:54–61.

Laurance, W. F., Clements, G. R., Sloan, S., O'Connell, C. S., Mueller, N. D., Goosem, M., et al. 2014. A global strategy for road building. Nature. 513:229–232.

Lave, L. B. 1963. The value of better weather information to the raisin industry. Econometrica. 31:151–164.

Leeuwis, C., and Aarts, N. 2011. Rethinking Communication in Innovation Processes: Creating Space for Change in Complex Systems. The Journal of Agricultural Education and Extension. 17:21–36.

Lobell, D. B., Cassman, K. G., and Field, C. B. 2009. Crop yield gaps: their importance, magnitudes, and causes. Annual Review of Environment and Resources. 34:179–204.

Lobell, D. B., Thau, D., Seifert, C., Engle, E., and Little, B. 2015. A scalable satellite-based crop yield mapper. Remote Sensing of Environment. 164:324–333.

Louwaars, N. P., de Boef, W. S., and Edeme, J. 2013. Integrated seed sector development in Africa: a basis for seed policy and law. Journal of Crop Improvement. 27:186–214.

Macauley, M. K. 2006. The value of information: Measuring the contribution of space-derived earth science data to resource management. Space Policy. 22:274–282.

Margosian, M. L., Garrett, K. A., Hutchinson, J. S., and With, K. A. 2009. Connectivity of the American agricultural landscape: assessing the national risk of crop pest and disease spread. BioScience. 59:141–151.

Mburu, H., Cortada, L., Mwangi, G., Gitau, K., Kiriga, A., Kinyua, Z., et al. 2018. First Report of Potato Cyst Nematode Globodera pallida Infecting Potato (Solanum tuberosum) in Kenya. Plant Disease. 102:1671–1671.

McGuire, S., and Sperling, L. 2013. Making seed systems more resilient to stress. Global Environmental Change. 23:644–653.

McGuire, S., and Sperling, L. 2016. Seed systems smallholder farmers use. Food Security.
8:179–195.
McQuaid, C. F., van den Bosch, F., Szyniszewska, A., Alicai, T., Pariyo, A., Chikoti, P. C., et al.
2017. Spatial dynamics and control of a crop pathogen with mixed-mode transmission.
PLOS Computational Biology. 13:e1005654.
McQuaid, C. F., Sseruwagi, P., Pariyo, A., and van den Bosch, F. 2016. Cassava brown streak
disease and the sustainability of a clean seed system. Plant Pathology. 65:299–309.
Mwangi, J. M., Kariuki, G. M., Waceke, J. W., and Grundler, F. M. 2015. First report of
Globodera rostochiensis infesting potatoes in Kenya. New Disease Reports. 31:18–18.
Nagarajan, R., Scutari, M., and Lèbre, S. 2013. *Bayesian networks in R with applications in*
systems biology. New York: Springer-Verlag.
Nakato, G. V., Beed, F., Bouwmeester, H., Ramathani, I., Mpiira, S., Kubiriba, J., et al. 2016.
Building agricultural networks of farmers and scientists via mobile phones: case study of
banana disease surveillance in Uganda. Canadian Journal of Plant Pathology. 38:307–316.
Novak, M. P., and LaDue, E. 1999. Application of recursive partitioning to agricultural credit
scoring. Journal of Agricultural and Applied Economics. 31:109–122.
Okello, J. J., Lagerkvist, C. J., Kakuhenzire, R., Parker, M., and Schulte-Geldermann, E. 2018.
Combining means-end chain analysis and goal-priming to analyze Tanzanian farmers'
motivations to invest in quality seed of new potato varieties. British Food Journal.
120:1430–1445.
Parsa, S., Morse, S., Bonifacio, A., Chancellor, T. C. B., Condori, B., Crespo-Perez, V., et al.
2014. Obstacles to integrated pest management adoption in developing countries.
Proceedings of the National Academy of Sciences. 111:3889–3894.
Paul, P. A., and Munkvold, G. P. 2004. A model-based approach to preplanting risk assessment
for gray leaf spot of maize. Phytopathology. 94:1350–1357.
Perez-Ariza, C. B., Nicholson, A. E., and Flores, M. J. 2012. Prediction of coffee rust disease
using Bayesian networks. In *Proceedings of the Sixth European Workshop on*
Probabilistic Graphical Models, , p. 259–266.
Petsakos, A., Hareau, G., Kleinwechter, U., Wiebe, K., and Sulser, T. B. 2018. Comparing
modeling approaches for assessing priorities in international agricultural research.
Research Evaluation. 27:145–156.
Phillips, C. B., Kean, J. M., Vink, C. J., and Berry, J. A. 2018. Utility of the CLIMEX 'match
climates regional' algorithm for pest risk analysis: an evaluation with non-native ants in
New Zealand. Biol Invasions. 20:777–791.
Reynolds, M., Kropff, M., Crossa, J., Koo, J., Kruseman, G., Molero Milan, A., et al. 2018. Role
of Modelling in International Crop Research: Overview and Some Case Studies.
Agronomy. 8:291.
Schulte-Geldermann, E., Gildemacher, P. R., and Struik, P. C. 2012. Improving seed health and
seed performance by positive selection in three Kenyan potato varieties. Am. J. Potato
Res. 89:429–437.
Schulthess, U., Timsina, J., Herrera, J. M., and McDonald, A. 2013. Mapping field-scale yield
gaps for maize: An example from Bangladesh. Field Crops Research. 143:151–156.
Shea, K., Tildesley, M. J., Runge, M. C., Fonnesbeck, C. J., and Ferrari, M. J. 2014. Adaptive
management and the value of information: learning via intervention in epidemiology ed.
Andrew P. Dobson. PLoS Biol. 12:e1001970.

Sheppard, C. S., Burns, B. R., and Stanley, M. C. 2014. Predicting plant invasions under climate
change: are species distribution models validated by field trials? Glob Change Biol.
20:2800–2814.
Silva, J. V., Reidsma, P., Laborte, A. G., and van Ittersum, M. K. 2017. Explaining rice yields
and yield gaps in Central Luzon, Philippines: An application of stochastic frontier
analysis and crop modelling. European Journal of Agronomy. 82:223–241.
Smith, C. S., Howes, A. L., Price, B., and McAlpine, C. A. 2007. Using a Bayesian belief
network to predict suitable habitat of an endangered mammal – The Julia Creek dunnart
(*Sminthopsis douglasi*). Biological Conservation. 139:333–347.
Sperling, L. 2008. *When Disaster Strikes: a guide to assessing seed system security*. CIAT.
Sperling, L., Ortiz, O., and Thiele, G. 2013. *Seed systems. Conceptual frameworks for guiding*
practical interventions. ed. CGIAR. CGIAR.
Stolar, J., and Nielsen, S. E. 2015. Accounting for spatially biased sampling effort in presence-
only species distribution modelling ed. Janet Franklin. Diversity Distrib. 21:595–608.
Sutherst, R. W., and Maywald, G. F. 1985. A computerised system for matching climates in
ecology. Agriculture, Ecosystems & Environment. 13:281–299.
Sutton, N. J., and Armsworth, P. R. 2014. The grain of spatially referenced economic cost and
biodiversity benefit data and the effectiveness of a cost targeting strategy: spatial grain in
conservation planning. Conservation Biology. 28:1451–1461.
Thomas-Sharma, S., Andrade-Piedra, J., Carvajal Yepes, M., Hernandez Nopsa, J. F., Jeger, M.
J., Jones, R. A. C., et al. 2017. A risk assessment framework for seed degeneration:
informing an integrated seed health strategy for vegetatively propagated crops.
Phytopathology. 107:1123–1135.
Tittonell, P., and Giller, K. E. 2013. When yield gaps are poverty traps: The paradigm of
ecological intensification in African smallholder agriculture. Field Crops Research.
143:76–90.
Tulloch, A. I. T., Tulloch, V. J. D., Evans, M. C., and Mills, M. 2014. The value of using
feasibility models in systematic conservation planning to predict landholder management
uptake: management feasibility models. Conservation Biology. 28:1462–1473.
Wang, C., Hawthorne, D., Qin, Y., Pan, X., Li, Z., and Zhu, S. 2017. Impact of climate and host
availability on future distribution of Colorado potato beetle. Sci Rep. 7:4489.
van Wart, J., van Bussel, L. G. J., Wolf, J., Licker, R., Grassini, P., Nelson, A., et al. 2013. Use
of agro-climatic zones to upscale simulated crop yield potential. Field Crops Research.
143:44–55.
van Wart, J., Kersebaum, K. C., Peng, S., Milner, M., and Cassman, K. G. 2013. Estimating crop
yield potential at regional to national scales. Field Crops Research. 143:34–43.
Wiebe, K., Zurek, M., Lord, S., Brzezina, N., Gabrielyan, G., Libertini, J., et al. 2018. Scenario
Development and Foresight Analysis: Exploring Options to Inform Choices. Annu. Rev.
Environ. Resour. 43:545–570.
Wilson, E. C. F. 2015. A practical guide to value of information analysis. PharmacoEconomics.
33:105–121.
Xing, Y., Hernandez Nopsa, J. F., Andersen, K. F., Andrade-Piedra, J., Beed, F. D., Blomme, G.,
et al. 2020. Global cropland connectivity: a risk factor for invasion and saturation by
emerging pathogens and pests. BioScience. In press.

You, L., Crespo, S., Guo, Z., Koo, J., Ojo, W., Sebastian, K., et al. 2012. MapSpaM - Home of
the Spatial Production Allocation Model. Available at: http://mapspam.info/ [Accessed
December 15, 2017].
890

891   Table 1. Regional adoption rates after positive seed selection training in Kenya and the expected
892   benefit of training given the adoption rate (Gildemacher et al. 2011, 2012). The average benefit
893   is that expected under random allocation of training effort to the regions without regard to
894   adoption rates.
895

County	Observed adoption rate	Per household benefit $USD	Expected benefit of training in year 1
Nakuru	0.46	156	72
Nyandarua	0.19	156	30
Narok	0.18	156	28
		Average Benefit:	44

896
897

898 Table 2. Potential criteria for identifying priority sites for interventions (such as training farmers
899 to use positive selection for improved seed health).
900

Criterion	Rationale
Regions where expected absolute benefit is greatest	Greatest benefit to regional food production
Regions where expected proportional gain is greatest	Greatest benefit to regional farmers
Regions where outcomes before intervention are lowest	Benefit to regions in greatest need
Regions where outcomes before intervention are highest	Benefit to regions currently best adapted for production
Incorporating measures of uncertainty at the farm level	
Regions where the 5th percentile benefit is greatest	Consistent benefit across farmers
Regions where the trimmed mean is greatest	Greatest benefit for typical farmers

901
902

Management performance mapping steps	Andes case study	Kenya case study
1) Formulate questions about the performance of a specific management strategy across a geographic region	1) Where does positive selection of farm-saved potato seed provide the greatest benefit in Ecuador and Colombia?	1) Where does positive selection of farm-saved seed provide the greatest benenfit in three regions of Kenya?
2) Assemble data related to the performance of the management strategy	2) Per plant yield, viruses, seed age, altitude, and management (including positive selection)	2) Positive selection adoption rate and yield response to positive selection
3) Identify predictor variables for management performance	3) Altitude, variety, and time since seed replacement - identified using Bayesian networks and classification trees	3) Proportional differences in regional adoption rates
4) Estimate management performance across the geographic study region	4) Extrapolation based on altitude for major potato regions of Ecuador and Colombia	4) (No extrapolation in this case study)
5) Evaluate the value of information for management intervention or policy	5) Comparison of intervention yield benefits with and without acting on information about altitude, variety and time since seed replacement	5) Comparison of intervention yield benefits with and without acting on information about adoption rates

6) Evaluate what additional information may be needed before proceeding in participatory planning sessions
7) Design intervention based on geographic locations identified
8) Monitor project outcomes and iteratively adapt strategies based on new data and stakeholder input

904
905
Figure 1. Steps in the management performance mapping pipeline. Selected development
interventions should ideally take place in a culture of continuous improvement, based on ongoing
monitoring and evaluation with stakeholders, and incorporating experimentation to facilitate
adaptive management. Two case studies show how the steps may be implemented. Management
performance mapping operates in this context by scaling up field, farm, and plot derived
information to larger scale landscapes, regions or countries.
912
913

Environment A **High** management performance	Environment B **Low** management performance	Environment C **Low** management performance
Yield without management: 5 Yield with management: 9 Benefit: 4	Yield without management: 5 Yield with management: 6 Benefit: 1	Yield without management: 5 Yield with management: 5 Benefit: 0

Type of site selection	Probability of site selection within each of these three environments			Expected site performance
Random (uninformed)	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{3}$	$4*\frac{1}{3}+1*\frac{1}{3}+0*\frac{1}{3}=1\frac{2}{3}$
Informed	1	0	0	$1*4=4$
Misinformed	0	$\frac{1}{2}$	$\frac{1}{2}$	$1*\frac{1}{2}+0*\frac{1}{2}=\frac{1}{2}$

Value of information comparing informed site selection to random site selection: $4-1\frac{2}{3}=2\frac{1}{3}$

Value of information comparing informed site selection to misinformed site selection: $4-\frac{1}{2}=3\frac{1}{2}$

Figure 2. The of information (VOI) for data used to guide site selection for interventions can be evaluated as illustrated here for a hypothetical case. Suppose there are three types of environment, each equally common, and a measure of how well the management being evaluated performs in each environment: improvements in yield in environments A, B, and C of 4, 1, and 0 units, respectively. If sites are selected at random for intervention, without information about yield in the different environments, the average benefit from management is 1 2/3 units. If sites are selected considering the information about better management performance in environment A, and thus only environment A is targeted, then the average benefit from management is 4 units. If misinformation leads to the incorrect belief that management is more effective in environments B and C and these environments are equally targeted, then the average benefit from management is 1/2 unit. The VOI comparing informed site selection to random site selection is 2 1/3 units. The VOI comparing informed site selection to misinformed site selection is 3 1/2 units.

Figure 3. Recursive partitioning results in a decision tree format, with per plant yield (g) of
potato in Ecuador as the response variable, based on the Kromann et al. (2017) dataset. Branches
to the left are results when the logical statements at the nodes are true, and branches to the right
are results when the logical statements are false. The upper numbers in the boxes are the mean
yields for that condition, and the percent values are the proportion of the data for which the
condition applies. For "Cycle >= 2.5", "yes" indicates that the time since seed replacement with
certified seed was greater than 2 years, while 'no' indicates that it was 2 or fewer years. For
"Management = Random, Roguing", "yes" indicates that either roguing or random seed selection
was implemented, while 'no' to that option indicates that positive selection was implemented.
For "Variety = Superchola", "no" indicates that the variety was INIAP-Fripapa. (Darker colors
indicate a higher number of 'no' answers for that condition compared to other conditions.)
944

945
946
947

948
949

Figure 4. Ecuadorian potato yield and the factors associated with yield from a Bayesian network
analysis carried in Netica using the Kromann et al. (2017) dataset. The two networks indicate the
frequency distribution of a set of twelve potential predictor variables for plants with low yield
(top) and high yield (bottom). The lower text for each node gives the estimated mean and
uncertainty. Positive selection was less likely to be the management implemented for cases
where the yield was low (top network). All the viruses except *Potato virus Y*, and *Potato leaf roll*

virus for low yield plants, were more likely to be absent (frequency = 0) than present. Each virus
was relatively more likely to be absent if a plant had high yield (bottom) than if it had low yield.
958
959

961
962
963
964
Figure 5. Ecuador and southern Colombia, with potato production indicated based on SPAM
estimates. An altitude of 2895 m.a.s.l. was identified as a cut-off for management performance
for positive selection of plants for on-farm seed saving. Pixels above 2895 m elevation (51% of
the pixels) are indicated with a dot, where pixels are included if the harvested area estimate is
greater than 200 ha. The graticules are 1-degree squares. Higher values of the potato cropland
connectivity risk index estimated for Ecuador and southern Colombia are indicated by darker
colors, indicating likely more important roles in potato epidemics. Targeting sites for farmer
training in positive selection, might be based on the combination of being above the altitude cut-
off for positive selection performance, and being in high cropland connectivity locations such
that improved management would have the potential to positively influence other regions.
975
976

977
978

979
980
981
Figure 6. Cropland connectivity in the area of three counties in Kenya, where darker shading indicates a higher cropland connectivity risk index. Cropland connectivity is a measure of the likely importance of a pixel for epidemic spread through potato production. When the three counties indicated were studied to evaluate adoption rates for positive selection of plants for on-farm seed saving, Nakuru county was reported to have over twice the adoption rate. Targeting for training in positive selection methods could take into account the higher adoption rate in Nakuru and the higher cropland connectivity in Nyandarua.