Damage-induced reactive oxygen species enable regenerative signalling by the rapid repositioning of Hedgehog expressing cells.

Montserrat Garcia Romero[1§], Gareth McCathie[1§], Philip Jankun[1]
and Henry Hamilton Roehl[1♭]

[1]Bateson Centre
Department of Biomedical Sciences
The University of Sheffield
Firth Court, Western Bank
Sheffield S10 2TN
United Kingdom

[§]These authors contributed equally
[♭]Corresponding author

Abstract

Aquatic vertebrates have been studied for their remarkable ability to regenerate
limbs and tails after amputation. Previous studies indicate that reactive oxygen
species (ROS) signalling is required to initiate regeneration, but the cellular
mechanism by which this takes place is poorly understood. The developmental
signalling molecules retinoic acid, Hedgehog, Wnt and FGF, have been shown to
have pro-regenerative roles in many systems. However, whether these are playing
roles that are specific to regeneration, or are simply recapitulating their
developmental functions has remained unclear. To address these questions, we have
analysed zebrafish larval tail regeneration. We find evidence that ROS released
upon wounding mobilise notochord cells resulting in repositioning of these cells to
the damage site. These notochord cells secrete Hedgehog ligands which are required
for regeneration. Intriguingly, we find that Hedgehog signalling is not required for
normal tail development suggesting that it plays a regeneration specific role. Our
results provide a model for how ROS signalling initiates tail regeneration, and
indicate that developmental signalling pathways can play regenerative functions
that are not directly related to their known developmental roles. This understanding
may facilitate development of regenerative therapies that involve the application of
signalling molecules to damaged tissues or organs.

Introduction
21
The study of regenerative biology aims to understand the mechanisms and limitations of
endogenous regenerative capacity. During vertebrate appendage regeneration in
salamander limb, mouse digit, and *Xenopus* tadpole tail and zebrafish fin, a conserved
sequence of events takes place after tissue is removed[1,2]. Immediately following tissue
damage the wound is closed by active movement of the surrounding epithelia. Next, the
epithelial covering of the wound thickens to form what is called the wound epithelia and
cells migrate under the wound epithelia to form a tightly packed mass proliferative cells
called the blastema. During the final phase of regeneration the blastemal cells and the
wound epithelium proliferate and differentiate to give rise to the missing tissue.
31
How wounding initiates regeneration is a central question in regenerative biology. Tissue
damage results in the immediate release of signals such as calcium, ATP, and reactive
oxygen species (ROS) which act to stimulate wound closure and the immune response
thus limiting detrimental effects of injury[3]. Of these initial signals, ROS stand out as a
good candidate for the activation of regeneration. Upon wounding, calcium release results
in a rapid burst of ROS that is likely to be produced by Duox, a NADPH Oxidase[3]. Then
ROS, primarily in the relatively stable form of hydrogen peroxide, are thought to diffuse
into the neighbouring tissue to act as a paracrine signal (referred to as ROS signalling).
ROS exert their effects through the reversible oxidation of cysteine residues in key
regulatory proteins[4]. Although precisely how ROS signalling is able to confer specific
cellular responses is still poorly understood, studies focused upon the MAPK and Wnt
pathways suggest that ROS levels may serve to modulate the activity of diverse signalling
pathways[5,6]. These studies indicate that members of the thioredoxin family of redox
sensors bind signalling pathway components in a ROS dependent manner.
46
The importance of ROS signalling to vertebrate regenerative biology has only recently
become apparent. Research on zebrafish larvae has shown that ROS are required for
axonal regeneration[7] and that ROS acts via a Src Family Kinase (SFK) to promote
regeneration of the fin fold[8]. Other zebrafish studies have shown that adult heart and fin
require ROS to regrow after wounding[9,10]. ROS have also been shown to act during

Xenopus tadpole regeneration[11]. Thus ROS signalling may act more generally as signal
that serves to both coordinate the damage response and to initiate the regeneration
program.
55
Another crucial question is how the regeneration of an organ differs from the initial
development of that organ[12,13]. For example, the mesenchyme of the developing limb bud
resembles the blastema of the amputated limb in both morphology and its expression of
msx genes. Similarly, the apical ectodermal ridge of the limb bud resembles the wound
epithelium and both structures express *dlx* genes. This suggests that once these structures
have appeared, regeneration follows the previously established developmental program.
Developmental signalling pathways such as Wnt/β-Catenin, FGF, Hedgehog, Retinoic
Acid (RA), Notch and BMP have been shown to play important roles during regeneration.
We must therefore ask whether the regenerative roles of these pathways are unique to
regeneration or are simply a recapitulation of earlier developmental roles.
66
To shed light on these questions, we have analysed how developmental pathways direct
zebrafish larval tail regeneration. Here we provide evidence that Hedgehog signalling is a
key regulator of tail regeneration, acting upstream of the Wnt/β-Catenin, FGF and RA
signalling pathways. This finding is surprising given that Hedgehog signalling does not
play a role in tail development. In addition, we propose that the source of the regenerative
Hedgehog signal is notochord cells that are rapidly repositioned to the stump immediately
following wounding. Our data suggest that this movement is dependent upon release of
ROS from the wound site and requires SFK activity and microtubule polymerisation.
Together these data suggest a model that ROS signalling initiates tail regeneration by
relocating Hedgehog expressing notochord cells to the wound site.
77
78 **Results:**
79
Regeneration in zebrafish larvae has been studied in two contexts: fin fold excision and
tail excision[14,15,16]. During fin fold excision, tissue removal is limited to epithelium and fin
mesenchymal cells in the caudal region of the tail (Fig. S1a). On the other hand, tail
excision involves partial removal of neural tube, notochord, muscle, pigment cells, blood
vessels as well as the caudal fin fold (Fig. S1a). Within minutes after tail excision
notochord cells move out of the notochord sheath to give rise to a cluster of cells (the
"notochord bead)") that sit on the stump of the tail (Supplementary Movie 1). Formation
of the notochord bead appears to be caused by contraction of the anterior/posterior body
axis resulting in pressure build up in the notochord (Fig. S2). To determine the timing of
regeneration after tail excision we examined the expression markers of the different
stages during regeneration (Fig. S1b). By 24 hours post excision (hpe), *dlx5a* expression
marks the forming wound epithelium, and by 48hpe the blastema is marked by strong
expression of *msxc*. A previous study of fin fold regeneration found that the early
blastema is marked by the RA synthesis gene *raldh2* and that RA signalling is required
for regeneration[17]. Consistent with this, we find that *raldh2* is upregulated in the forming
blastema at 24hpe. Increased expression of the muscle differentiation gene *myod* is seen
between 48 and 81hpe suggesting that regrowth takes place during this interval. Although
these genes are expressed during tail development, they are not detected immediately
prior to excision (Fig. S1b). This indicates that tail excision reactivates expression of
these genes. If operated fish are raised past larval development, they appear
morphologically normal. However, skeletal visualisation reveals that the internal
structure of the tail is modified perhaps due to a defect in notochord extension (Fig. S1c).
102
To begin to understand how developmental signalling coordinates regeneration, we first
focused on the FGF pathway to identify ligands and downstream targets that are induced
by tissue removal. Similar to previous studies that identified *fgf20a* as a damage-induced

ligand in the fin fold[18], we found that *fgf10a* transcripts are upregulated at 24hpe in cells
adjacent to the extruded notochord bead (Fig. 1a). We also found that the target gene
pea3 is induced in the surrounding cells, indicating that these cells are responding to FGF
signalling at this time (Fig. 1a). Neither gene is expressed strongly in unoperated tails at
this time (Fig. S3a,b). The allele *tbvbo* encodes a null mutation in *fgf10a* that does not
have a discernable effect on normal tail development (Fig. S4)[19]. Consistent with a role for
FGF signalling in larval tail regeneration we found that *fgf10a*[-/-] larvae have reduced
regenerative capacity (Fig. 1b). To determine whether the wound epithelium and
blastema form properly, we tested levels of expression of *dlx5a* and *raldh2* in *fgf10a*[-/-]
larvae and in fish treated with the FGF receptor inhibitor SU5402[20]. Both conditions did
not show an observable difference in expression suggesting that FGF signalling does not
play a role in the initial patterning of the regenerating tail (Fig. 1c). A previous study
found that FGF signalling is required for regeneration specific proliferation in the fin
fold[15], and similarly we found that the number of proliferating cells in *fgf10a*[-/-] larvae is
reduced compared to wild-type fish (Fig. 1d). These results suggest that FGF signalling is
required for damage-induced proliferation after tail excision.
122
We next investigated the role of Wnt/β-Catenin signalling during tail regeneration by
screening for Wnt genes that are activated after excision. We found that *wnt10a* is
upregulated starting at 18hpe in cells neighbouring the notochord bead and the Wnt/β-
Catenin downstream target gene *tcf7* is upregulated also in this region (Fig. 2a). Neither
gene is expressed strongly in unoperated tails at this time (Fig. S3c,d). To further
investigate we used the Wnt/β-Catenin pathway inhibitor IWR-1[21] to turn off signalling
following excision. Treatment with IWR-1 resulted in the absence of tail regrowth
suggesting that Wnt/β-Catenin signalling is required for regeneration (Fig. 2b). To
determine whether *dlx5a* or *raldh2* expression depend upon the Wnt/β-Catenin pathway,
we used a heatshock inducible *dickkopf-1b* transgene (*hsp70l:dkk1b-GFP*)[22] to inhibit
signalling and a glycogen synthase kinase antagonist (GskXV) to activate signalling[23]. We
found that whereas inhibition abolished expression of both genes, pathway activation led
to expansion of the expression domains of *dlx5a* and *raldh2* (Fig. 2c). Since both FGF
and Wnt/β-Catenin pathways appear to act at the same time (24-48hpe), we wondered
whether they might regulate each other's activity. By manipulation of both pathways we
found that each pathway has a positive effect on the other pathway's activity (Fig. 2d,e).
These data indicate that Wnt/β-Catenin signalling patterns the early regenerating tail and
interacts with FGF signalling.
141
The third developmental pathway that we have investigated is the Hedgehog signalling
pathway. We first tested whether the Hedgehog pathway influences regeneration using
the inhibitor cyclopamine[24] (Fig. 3a,b). Cyclopamine treatment following tail excision
resulted in loss of regeneration and loss of *dlx5a* and *msxc* expression. Consistent with an
early role for Hedgehog signalling we found that treatment from -12hpe to 12hpe is
sufficient to block regeneration. To control for off-target effects of the cyclopamine
treatment, we used another Hedgehog pathway inhibitor, LDE225[25], and obtained similar
results (Fig. 3a,b). To determine how Hedgehog signalling controls regeneration, we
tested for expression of Wnt, FGF and RA genes after cyclopamine treatment and found
that their expression is abolished (Fig. 3c). Consistent with Hedgehog signalling acting
upstream of FGF signalling we found that cyclopamine treatment strongly reduced
proliferation after tail excision (Fig. 3d). Given the early timing of the Hedgehog
signalling requirement, it is possible that this pathway acts upstream of the other
developmental pathways. To test this, we performed chemical epistasis by treating
regenerating fish with cyclopamine to block the Hedgehog pathway while activating the
Wnt/β-Catenin pathway with GskXV. We found that GskXV treatment is sufficient to
restore expression of the regeneration markers *dlx5a*, *msxc* and *raldh2* indicating that the
Hedgehog and Wnt/β-Catenin pathways form a linear pathway (Fig. 3e). Together these

data present a model that the Hedgehog pathway plays a key role during regeneration by
activating the Wnt/β-Catenin, FGF and RA pathways.
162
To determine the source and timing of Hedgehog signalling, we looked at expression of
Hedgehog ligands as well as the downstream target *patched1* (*ptch1*). We found that two
Hedgehog ligands (*sonic hedgehog a*, *shha* and *indian hedgehog b*, *ihhb*) are strongly
expressed in the notochord bead (Fig. 4a and Fig. S5a). Although this expression appears
to be limited to the cells in the bead, this restricted detection is an artefact of the
wholemount in situ hybridisation method: When fixed larvae are cut along the coronal
axis or obliquely to reveal the notochord prior to hybridisation, then expression of *ihhb* is
detected in the notochord before and after excision (Fig. 4b). This artefact is likely to be
because notochord sheath cells deposit a dense extracellular matrix that may restrict
penetration of components during in situ hybridisation. Consistent with this model, we
find that there is a low level of expression of *ptch1* around the caudal tip of the notochord
prior to excision (Fig. S3f), and high level expression after the notochord bead has
formed (Fig. S5a). To further test whether Hedgehog signalling acts upstream of other
developmental pathways, we looked to see whether manipulation of any of these
pathways affects the wound-induced expression of *ptch1* and found that none of the
treatments altered its expression (Fig. S5b). Together these data suggest the model that
formation of the notochord bead provides for a novel source of Hedgehog signalling that
acts upstream of other developmental pathways.
181
In the fin fold excision model of regeneration, the notochord is not injured suggesting that
the initiation of regeneration may take place by a mechanism that does not rely upon the
notochord bead and Hedgehog signalling. To test this we treated fish with cyclopamine
after fin fold excision. We found a mild effect in that the fin fold regenerates at a slower
rate when Hedgehog signalling is blocked (Fig. S6a). Consistent with this we did not
detect upregulation of *ihhb* or *ptch1* after fin fold excision (Fig. S6b). Although these
results suggest that the Hedgehog pathway may act during fin fold regeneration, they
indicate that it is unlikely to play the same crucial role that it does during tail regeneration.
190
Given that ROS signalling has been proposed to activate regeneration in different models,
it is possible that ROS initiate larval tail regeneration upstream of Hedgehog signalling.
High levels of ROS are synthesized in cells along the edge of the stump starting
immediately after excision (Fig. S7a). To begin to test the role of ROS we blocked their
generation with the flavoenzyme inhibitor diphenylene iodonium (DPI) which inhibits
NADPH oxidases[3]. Treatment with 150μM DPI from 1 hour prior to excision until 1hpe
is sufficient to reduce regeneration by 50% (Fig. S7b,c) and strongly reduces wound-
induced ROS levels (Fig. S7a,d). DPI may affect other signalling pathways besides ROS
that involve flavoenzymes or indeed may have unrelated off-target effects[26]. To control for
these effects of the DPI treatment, we utilised the ROS scavenger MCI186[11] which also
results in reduced ROS levels and regenerated tail length (Fig. S7d,e). The effect of MCI
186 is milder that that of DPI but increasing levels of MCI 186 results in toxicity, so we
kept the levels of MCI 186 below this threshold. Consistent with ROS signalling
activating regeneration upstream of the Hedgehog pathway, we found that wound-
induced expression of *ptch1*, *raldh2*, *tcf7* and *pea3* is reduced following DPI treatment
(Fig. S7f).
207
We next sought to determine whether ROS signalling potentially regulates regeneration
by promoting the formation of the notochord bead. When we measured the size of the
notochord bead at 4hpe, we found that DPI reduced bead formation, but that MCI
treatment did not (Fig. 5a). As SFKs have been proposed to act downstream of ROS
signalling during fin fold regeneration, we tested whether the SFK inhibitor PP2[27]
influences bead size. We also used nocodazole which interferes with microtubule

polymerisation in an attempt to block bead formation by an independent mechanism. We
found that both compounds had a strong effect on bead formation (Fig. 5a). To assess
how these inhibitors affect wound-induced activation of Hedgehog pathway, we analysed
levels of *ptch1* and *ihhb* expression. We found that these compounds reduce the amount
of *ihhb* transcripts in the notochord bead (Fig. 5b,c) and reduce *ptch1* to levels similar to
those seen with DPI treatment (Fig. S8). Although nocodozole treatment would also be
predicted to affect ciliogenesis and thus Hedgehog signal transduction, the observation
that treatment blocks bead formation and Hedgehog ligand expression suggest that
nocodozole exerts its effect upstream of ciliogenesis. Intriguingly, DPI treatment can
result in the loss of *ihhb* expression even when the bead is fully extruded suggesting that
a burst of ROS production may be required for both notochord extrusion and expression
of *ihhb* in the notochord bead (Fig. S9). Together these data suggest that ROS/SFK
dependent bead formation is a necessary step in the regeneration of the tail.
227
To further control for potential unintended effects of chemical treatments we decided to
test whether we could rescue DPI and PP2 treatment by activating Wnt/β-Catenin
signalling as we have done for cyclopamine treatments (Fig. 3e). We treated fish at the
time of excision with either DPI or PP2 to inhibit bead formation, allowed the fish to
recover and then the next day activated WNT/β-Catenin signalling with GskXV. Rescue
was quantified by measuring the area of *raldh2* expression after RNA in situ analysis.
Both DPI and PP2 treated fish show a significant level of recovery of *raldh2* expression
after treatment with GskXV (Fig. S10). This experiment suggests that the effects of DPI
and PP2 are not simply due to toxic off-target effects and support the hypothesis that
ROS and SFKs act upstream of Wnt/β-Catenin signalling. In addition we attempted to
rescue formation of the notochord bead by supplementing the fish water with H_2O_2 during
DPI treatment (data not shown). These tests were unsuccessful perhaps because the
timing, concentration or distribution of endogenous ROS is difficult to mimic with
exogenously provided H_2O_2. Given these limitations, the DPI treatments should be
interpreted with caution and further analysis of ROS signalling is needed to confirm its
role in regeneration.
244
Having found that DPI and PP2 both strongly reduce notochord bead extrusion, we
decided test whether these compounds affect early cell shape changes seen directly after
tail excision (Supplementary Movie 1). Within the first few minutes after excision there
is a change in the curvature of the cell membranes which initially bow towards the
anterior, then change to bow towards the posterior (Fig. S2). This change suggests that
these cells are passively being forced towards the open end of the notochord perhaps due
to increased pressure within the notochord. To quantify this change we chose to measure
the Menger curvature which is the inverse of the radius of a circle that approximates the
curved arc of the cell membrane. A cell membrane that runs perpendicular to the
notochord sheath will result in a Menger curvature of 0, one bowed to the posterior
results in a positive value and one bowed to the anterior a negative value. We measured
changes in notochord cell curvature during the first 20 minutes after excision and found
that while control fish show a dramatic change in curvature, those treated with DPI or
PP2 resemble uncut fish (Fig. S11).
259
A second early morphological change after tail excision is contraction of the trunk along
the anterior/posterior axis (Supplementary Movie 1). Contraction along this axis could
result in increased pressure within the notochord that may result in expulsion of cells
from the open end of notochord. We measured trunk length of individual animals before
excision and then measured the same animal again at 2hpe (Fig. S12). We found that
whereas untreated fish contract by an average of 4.4%, DPI and PP2 only contract by
1.7% and 1.2% respectively. Taken together these data suggest that ROS/SFK signalling
is required for early cell movements immediately following tail excision.

268
Given the complex signalling interactions that take place during regeneration, we
wondered whether these interactions are also required during normal tail development.
Although interactions between the FGF, Wnt/β-Catenin and RA pathways are crucial for
tail formation in chick, mouse and zebrafish[28, 29, 30, 31], roles for ROS and Hedgehog signalling
have not been described. To test whether ROS act during tail development, we treated
larvae with DPI during axis elongation and found that morphologically tails are
unaffected by this treatment (Fig. 6a). Likewise, expression of Hedgehog ligands is not
affected by DPI treatment during tail development (Fig. 6b). To functionally test whether
Hedgehog signalling acts on the tail bud, we treated fish with cyclopamine and found that
blocking Hedgehog signalling does not affect *raldh2* or *tcf7* levels (Fig. 6c). Consistent
with this, we found that cyclopamine treated fish have relatively normal tail development
except for the formation of U-shaped somites due to a known function of Hedgehog
signalling in somite patterning (Fig. 6d)[24]. Furthermore, fish carrying null mutations in the
Hedgehog receptor gene *smoothened* (*smo*[b577])[32] form a tail, but lack Hedgehog signalling
(Fig 6d). *smoothened* is absolutely required for Hedgehog signalling as it has no paralogs
and there is no evidence for its redundancy with other genes. The possibility that low
levels of maternal Smoothened activity may play a role in tail development is very
unlikely as maternal-zygotic *smoothened* mutants form a tail[33]. These data indicate that
proposed regulatory interactions mediated by ROS and Hedgehog do not act during
normal tail development and have specific functions during tail regeneration.
289
290 **Discussion**
291
In the course of this study, we have uncovered a mechanism by which damage-induced
ROS signalling may initiate regeneration (Fig. 7). Our model does not exclude the
involvement of other signals such as TGFβ, EGF (Epidermal Growth Factor), hypoxia
and non-canonical WNT signalling which are also likely to play roles in larval tail
regeneration[14, 22, 34, 35]. Rather this model is intended to provide a framework for future
analysis of larval tail regeneration in fish. Surprisingly our data suggest that ROS
influence regeneration by causing the rapid repositioning of Hedgehog-expressing
notochord cells to the site of the wound. Prior to injury, notochord cells express
Hedgehog ligands but this expression has little or no direct effect on cells neighbouring
the notochord as judged by expression of *ptch1*. Once the sheath is breached and
notochord cells are extruded to form a bead, high levels of Hedgehog ligands signal to the
surrounding tissue. Another surprising result is that although Hedgehog signalling is
required for tail regeneration, it is dispensable for tail development. Thus the Hedgehog
pathway plays a regeneration-specific role in this context and acts as a relay between the
immediate damage response and the expression of signalling pathways known to
coordinate both development and redevelopment of the tail.
308
The important role played by the notochord extrusion and Hedgehog pathway activation
in zebrafish larvae may be conserved during tail regeneration in other aquatic vertebrates
such as frogs and salamanders. One study has shown that in *Xenopus* tadpoles a
notochord bead forms at the stump within the first few hours of tail excision, and that
these extruded notochord cells express Hedgehog ligands from 24-48hpe[36]. This study
went on to show that tadpoles treated with cyclopamine immediately following tail
excision have reduced regeneration. Another study has shown that ROS signalling is
required for induction of FGF and Wnt/β-Catenin signalling at 36hpe[11]. Based upon this
timing, it is possible that Hedgehog signalling from the notochord bead also acts to link
ROS signalling to redevelopment of the tadpole tail. The picture from axolotl is
intriguing, as although Hedgehog signalling is important for axolotl tail regeneration, the
expression of *Sonic hedgehog* is restricted to the floor plate of the neural tube[37]. It would
be fascinating if the Hedgehog pathway has maintained its role as a key regulator of tail

regeneration in axolotl despite its expression being limited to the floor plate. In addition
to its role in tail regeneration, Hedgehog ligands *shha*, *ihhb* and *desert hedgehog* have
been shown to be upregulated during heart regeneration where they direct epicardial
regeneration[38] and Hedgehog signalling activates Wnt/β-Catenin genes during newt limb
regeneration[39]. As these tissues also require ROS production to regenerate, it will be
interesting to test whether the Hedgehog pathway acts to link ROS signalling to
redevelopment in these contexts.
329
In our model of tail regeneration, we have placed the FGF, Wnt/β-Catenin and RA
pathways in a in a "redevelopment" module because broadly speaking these pathways
interact in a similar ways during development and regeneration (Figure 7). This makes
sense because it is unlikely that organisms would evolve entirely novel mechanisms to
regrow tissue, when a pre-existing developmental module would suffice. However, this
may be an oversimplification as there are some fundamental differences between the
origins and movement of precursor cells during development and redevelopment. Our
preliminary comparison of cross talk between these pathways also suggests that
developmental interactions are present but in an altered form during redevelopment (data
not shown).
340
In contrast, Hedgehog signalling plays a novel role during tail regeneration, raising the
question of how Hedgehog signalling has evolved to act upstream of the redevelopment
module. If in nature larvae are often losing the end of their tails, then it is reasonable to
suggest that Hedgehog regulation of the redevelopment module has evolved due to
selective pressure. Alternatively, Hedgehog signalling may interact with FGF, Wnt/β-
Catenin and RA pathways during another developmental process that is unrelated to tail
regeneration. In this case its regulatory role could have evolved and be maintained by
selection for this unrelated developmental process. This second model does not require
selective pressure to evolve Hedgehog's regenerative role, and rather suggests that an
existing developmental signalling network has been co-opted to serve during regeneration.
351
A related question is how ROS/SFK signalling has evolved to regulate notochord
extrusion. Here we propose that ROS cause contraction along the anterior/posterior axis
which results a build-up of pressure within the trunk of the fish. Given that the notochord
sheath forms a tube-like structure, compaction along the anterior/posterior axis could
build up pressure and force notochord cells rapidly out of the open end. The driving force
for contraction may be the mass movement of epithelial cells towards the stump that has
been described after fin fold excision[40]. The molecular mechanism for notochord bead
extrusion is likely to involve ROS signalling through SFKs as well as microtubule
polymerisation. Further evidence for this mechanism comes from a range of models.
Studies in *Xenopus* have shown that microtubule polymerisation is required for wound
closure[41] and ROS have been shown to play a role in wound closure in *C.elegans*[42] and
more recently in zebrafish[43]. *Yoo et al* have shown that the SFK Lyn acts as a receptor for
ROS signalling[27] and several studies have suggested the model that SFKs directly
phosphorylate microtubules to promote their polymerisation and/or stabilisation[44,45,46]. Thus,
it is possible that the notochord bead results from ROS-dependent morphological changes
that pressurise the notochord forcing cells out of the open end. Once formed, the
notochord bead then acts as a novel source of Hedgehog signalling to promote tail
redevelopment.

Figures

Figure 1: FGF signalling is required for cell proliferation during larval tail regeneration. a, Arrowheads point to expression of the FGF ligand, *fgf10a*, and the downstream target, *pea3*, which become evident by 24hpe. **b,** *fgf10a⁻* fish have reduced tail regrowth compared to wild-type fish. **c,** Expression of *dlx5a* and *raldh2* are not strongly affected when FGF signalling is reduced. SU5402 treatment was done from 26 hpe to 30hpe at 10μM concentration. This treatment is sufficient to abolish detection of the FGF downstream target *pea3*. **d,** Whereas in wild-type fish DNA mitosis increases after tail excision, in *fgf10a⁻* fish mitosis remains constant. Fish were stained with Phospho-Histone H3 (ser10) antibody in green.

Figure 2: Wnt/β-Catenin signalling is required for patterning during larval regeneration.
a, Expression of the Wnt ligand, *wnt10a*, and the downstream target, *tcf7*, are upregulated by 18hpe (arrowheads). **b,** Larvae treated with 10μM IWR-1 continuously after excision show no signs of regeneration at 96hpe. **c,** Expression domains of *dlx5a* and *raldh2* are abolished when Wnt/β-Catenin signalling is inhibited and increased when Wnt/β-Catenin signalling is up-regulated. **d,** Expression of *fgf10a* and *pea3* are down-regulated in *hs:dkk1* larvae and up-regulated in fish treated with GskXV. **d,** Expression of the Wnt/β-Catenin target gene *axin2* is reduced in fish treated with SU5402.

Figure 3: Hedgehog signalling plays a central role during larval tail regeneration.
a, Continuous inhibition with the Hedgehog pathway inhibitors cyclopamine and LDE225 blocks regeneration. Also, pulse treatment around the time of tail excision blocks regeneration at 96hpe. **b,** Larvae treated with cyclopamine and LDE225 from 0-48hpe have reduced expression of *dlx5a* and *msxc*. **c,** Wound-induced expression of markers for the Wnt/β-Catenin, FGF and RA pathways is lost after treatment with cyclopamine. Larvae in the *wnt10a* and *pea3* panels were treated with cyclopamine from 0-30hpe. *tcf7*, *fgf10a* and *raldh2* panels were treated with cyclopamine from 0-48hpe. **d,** Cell proliferation is inhibited by treatment with cyclopamine. This analysis was done as in Figure 1d. **e,** Upregulation of Wnt/β-Catenin signalling after cyclopamine treatment restores expression of *dlx5a*, *msxc* and *raldh2*. Arrowheads indicate partial rescue of marker expression.

Figure 4: The Hedgehog pathway is activated immediately after tail excision.
a, *ihhb* expression is detected within the notochord bead and continues until 30hpe. Expression of *ptch1* is upregulated by 18hpe and continues after 30hpe. Arrowheads point to expression domains. Oblique and coronal sections reveal that *ihhb* is expressed in the notochord before (**b**) and after (**c**) tail excision. Arrowheads point to expression in the notochord.

Figure 5: Comparison of the effects of DPI, MCI 186, PP2 and nocodazole on notochord bead extrusion and *ihhb* expression.
a, Bead extrusion is affected strongly by PP2 treatment and not by MCI 186 treatment. DPI and nocodozole have intermediate effects on extrusion. **b,** *ihhb* expression after different treatments. **c,** Quantification of *ihhb* stain from panel b shows that all treatments reduced the level of expression in the notochord bead. DPI treatment started at 1 hour prior to wounding and ended at 1hpe, after which larvae were washed in E3 buffer and incubated until 6hpe. MCI186 treatment started at 0hpe and ended at 3hpe. PP2 treatment started at 1 hour prior to wounding and ended at 6hpe. Nocodazole treatment started at 0hpe until 6hpe.

Figure 6: ROS and Hedgehog signalling do not have similar roles during tail development.
a, DPI treatment from 14-24hpf does not affect larval tail morphology. **b,** *shha* and *ihhb* expression in the tail is not affected by DPI treatment at from 14-18hpf. **c,** *tcf7* and *raldh2* expression in the tail bud is not affected by treatment with cyclopamine from 8-16hpf. **d,** Loss of Hedgehog signalling does not affect tail formation. Larvae were treated with cyclopamine from 8-10hpf. Homozygous *smo*[b577] mutants lack zygotic Hedgehog signalling activity.

Figure 7: A model for tail regeneration.

Figure S1: Larval tail regeneration. a, When cut in the pigment gap (zone between the two red arrows), the notochord bead (the green arrow) is extruded within 3hpe and the tail regenerates within 4-5 days. The position of fin fold excision is shown for comparison. **b,** Different stages of regeneration are marked by the expression of *dlx5a*, *raldh2*, *myod* and *msxc*. Uncut fish are shown to the right for comparison. **c,** Live images of the tail after metamorphosis is complete (left panels) and Von Kossa stain showing the failure of the notochord to extend dorsally (right panels). In unoperated fish the uroneural and hypurals 3-5 form around and are attached to the notochord which grows dorsally as the adult tail forms. After excision these bones all form on the urostyle suggesting that the dorsal flexure of the notochord has not occurred. Abbreviations: urostyle (ust), uroneural (uro) and hypural 1-5 (1-5).

Figure S2: There appears to be a change in pressure within the notochord at the start of extrusion. Close-up of notochord after tail excision to show cell membranes are bowed towards anterior at the start and then become bowed towards the posterior. Arrowhead marks a cell membrane. Images captured every 30 seconds as in Supplemental Movie 1. Scale bar = 100µm.

Figure S3: Expression of signalling genes in the larval tail.
a, *fgf10a* is expressed weakly in neuromasts (arrowheads). **b,** *pea3* is expressed in neuromasts (arrowheads) and has weak expression in the fin fold. **c,** *wnt10a* has weak expression in the ventral fin fold (arrowhead). **d,** *tcf7* has no detectable expression. **e,** *ihhb* has no detectable expression. **f,** *ptch1* has weak expression in tissue neighbouring the notochord (arrowhead).

Figure S4: *fgf10a* **does not act during tail development. a,** Arrowhead indicates *fgf10a* expression in the tail. **b,** Development of the *fgf10a⁻* larval tail is indistinguishable from wild-type. **c,** Measurement of tail growth does not show a significant difference between *fgf10a⁻* and wild-type fish.

Figure S5: *shha* **is expressed in the notochord bead and Hedgehog signalling is unaffected by manipulation of other developmental pathways.**
a, *shha* is expressed in the notochord bead (arrowheads). **b,** The activity of the Hedgehog pathway was monitored by in situ hybridisation for *ptch1*. Larvae were treated with GskXV at 12.5μM from 40-48hpe to activate the Wnt/β-Catenin pathway. *hs:dkk1b-GFP* fish were heatshocked for three hours at 39°C at 28hpe and allowed to recover for 17 hours at 28.5°C to inhibit the Wnt/β-Catenin pathway. Larvae were treated with SU5402 from 44-48hpf to inhibit the FGF pathway. Larvae were treated with DEAB from 1-48hpe to block the RA pathway. Larvae were treated with RA from 43.5-48hpf to activate the RA pathway.

Figure S6: Hedgehog signalling does not play a crucial role in fin fold regeneration.
a, Treatment with 20μM cyclopamine is sufficient to block regeneration and has a milder affect on fin fold excision. Regrowth was measured from images in Fiji/ImageJ software by placing a rectangle parallel to the body that started at the end of the notochord and finished at the caudal end of the fin fold. **b,** Neither *ptch1* nor *ihhb* are noticeably upregulated following fin fold excision. Arrowheads indicate low levels of *ptch1* expression in unoperated and fin fold excision larvae.

Figure S7: ROS activity immediately after tail excision. a, Time course showing production of ROS after tail excision is reduced in larvae treated with 150μM DPI from 1 hour prior to excision. Larvae were bathed in 10μM PFBS-F (205429, Santa Cruz) to detect ROS. **b,** Optimisation to show that DPI treatment for as little as 2hrs (1hr pre-treatment and 1hr post-treatment) is sufficient to reduce tail regrowth by >50%. **c,** Representative larval tails showing the extent of tail regrowth after DPI treatment. **d,** Quantification of PFBS-F fluorescence shows that DPI treatment has a stronger effect on ROS levels than MCI186. **e,** Comparison of the efficacy of DPI to MCI186 in regards to regenerated tail length. **f,** DPI treatment inhibits wound-induced activation of the Hedgehog, Wntβ-Catenin, RA and FGF pathways. Larvae were pre-treated with 150 μM DPI for one hour and for 6hpe in 100μM DPI. Larvae were then washed in E3 buffer and incubated until 24hpe or 48hpf in the case of *tcf7*. Arrowheads point to expression of *pea3* and *ptch1* that is found in unoperated animals (Supplementary Fig. S2).

Figure S8: Wound-induced *ptch1* expression is reduced at 24hpe after treatment with inhibitors. Samples were treated as in Figure 5c.

Figure S9: Treatment with DPI, MCI 186, PP2 or nocodazole and comparison of bead size to expression of *ihhb*. Data from panels 5a and 5c plotted together for direct comparison. Regression lines suggest that nocodazole has the strongest correlation and DPI has the weakest.

Figure S10: Treatment with DPI and PP2 is partially rescued by upregulation of the Wnt/β-Catenin pathway. Larvae were operated on at 72hpf and treated with DPI, PP2 alone or in combination with GskXV. For DPI treatments, larvae were pre-treated for one hour prior to excision at 150μM concentration, then in 100μM for 5hpe. For PP2 treatments, larvae were pre-treated for thirty minutes prior to excision at 20μM concentration, then in 20μM for 3hpe. Larvae were then treated with 10μM GskXV from 24hpe to 32hpe and stained for *raldh2* expression as a marker of the blastema.

Figure S11: Menger curvature measurements show that notochord cell shape changes are altered by treatment with DPI or PP2. Fish were pretreated for two hours with DPI or PP2 at 28.5°C. Then fish were operated and incubated at 21°C with imaging taking place at twenty minutes post excision.

Figure S12: Wound induced contraction of the trunk is reduced by treatment with DPI or PP2. Fish were pretreated for two hours with DPI or PP2. Then fish were imaged immediately prior to tail excision, incubated individually for a further two hours in DPI or PP2 and imaged again.

Supplemental Movie 1. Tissue movement following tail excision.
This movie shows the rapid repositioning of notochord cells to form the notochord bead. The white arrow marks one cell's movement (1.5μm/minute). Note that the notochord cell membranes are bowed towards the anterior at the start of the movie and rapidly change to bow posteriorly within the first five minutes. This may indicate that pressure is building up within the notochord, and that subsequent notochord cell movement is passive rather than an active migration.

Green dots mark reference points within the fin fold epithelia and show the movement of the tissue towards the posterior. Over the duration of the movie the path between these dots shrinks by 11%. Red dots mark individual mesenchymal cells, these move together by 9% during the movie. Measurement of the length of the dorsal edge of the body using melanocyte position also shows a contraction of 9%. Together these indicate that the trunk shrinks by approximately 10% along the anterior/posterior axis during the first three hours after excision. Measurements along the dorsal/ventral axis of each somite show that the trunk becomes wider by approximately 7%. Consistent with this, the angle of the somites increases by 5% on average.

REFERENCES

1.   Gemberling M, Bailey TJ, Hyde DR, Poss KD. The zebrafish as a model for complex tissue regeneration. *Trends in genetics : TIG* **29**, 611-620 (2013).

2.   Tanaka EM, Reddien PW. The cellular basis for animal regeneration. *Developmental cell* **21**, 172-185 (2011).

3.   Niethammer P, Grabher C, Look AT, Mitchison TJ. A tissue-scale gradient of hydrogen peroxide mediates rapid wound detection in zebrafish. *Nature* **459**, 996-999 (2009).

4.   Holmstrom KM, Finkel T. Cellular mechanisms and physiological consequences of redox-dependent signalling. *Nat Rev Mol Cell Biol* **15**, 411-421 (2014).

5.   Funato Y, Miki H. Redox regulation of Wnt signalling via nucleoredoxin. *Free Radic Res* **44**, 379-388 (2010).

6.   Ichijo H, *et al*. Induction of apoptosis by ASK1, a mammalian MAPKKK that activates SAPK/JNK and p38 signaling pathways. *Science* **275**, 90-94 (1997).

7.   Rieger S, Sagasti A. Hydrogen peroxide promotes injury-induced peripheral sensory axon regeneration in the zebrafish skin. *PLoS biology* **9**, e1000621 (2011).

8.   Yoo SK, Freisinger CM, LeBert DC, Huttenlocher A. Early redox, Src family kinase, and calcium signaling integrate wound responses and tissue regeneration in zebrafish. *The Journal of cell biology* **199**, 225-234 (2012).

9.   Gauron C, *et al*. Sustained production of ROS triggers compensatory proliferation and is required for regeneration to proceed. *Scientific reports* **3**, 2084 (2013).

10.   Han P, *et al*. Hydrogen peroxide primes heart regeneration with a derepression mechanism. *Cell research* **24**, 1091-1107 (2014).

11.   Love NR, *et al*. Amputation-induced reactive oxygen species are required for successful Xenopus tadpole tail regeneration. *Nature cell biology* **15**, 222-228 (2013).

12.   Nacu E, Tanaka EM. Limb regeneration: a new development? *Annual review of cell and developmental biology* **27**, 409-440 (2011).

13.   Wehner D, Weidinger G. Signaling networks organizing regenerative growth of the zebrafish fin. *Trends in genetics : TIG* **31**, 336-343 (2015).

14.   Rojas-Munoz A, *et al*. ErbB2 and ErbB3 regulate amputation-induced proliferation and migration during vertebrate regeneration. *Developmental biology* **327**, 177-190 (2009).

15.   Kawakami A, Fukazawa T, Takeda H. Early fin primordia of zebrafish larvae regenerate by a similar growth control mechanism with adult regeneration. *Dev Dyn* **231**, 693-699 (2004).

16.   Yoshinari N, Kawakami A. Mature and juvenile tissue models of regeneration in small fish species. *Biol Bull* **221**, 62-78 (2011).

17.   Mathew LK, *et al*. Comparative expression profiling reveals an essential role for raldh2 in epimorphic regeneration. *J Biol Chem* **284**, 33642-33653 (2009).

18.   Whitehead GG, Makino S, Lien CL, Keating MT. fgf20 is essential for initiating zebrafish fin regeneration. *Science* **310**, 1957-1960 (2005).

19.   Norton WH, Ledin J, Grandel H, Neumann CJ. HSPG synthesis by zebrafish Ext2 and Extl3 is required for Fgf10 signalling during limb development. *Development* **132**, 4963-4973 (2005).

20.   Roehl H, Nusslein-Volhard C. Zebrafish pea3 and erm are general targets of FGF8 signaling. *Curr Biol* **11**, 503-507 (2001).

21.   Yin A, Korzh S, Winata CL, Korzh V, Gong Z. Wnt signaling is required for early development of zebrafish swimbladder. *PLoS One* **6**, e18431 (2011).

22.   Stoick-Cooper CL, *et al*. Distinct Wnt signaling pathways have opposing roles in appendage regeneration. *Development* **134**, 479-489 (2007).

23.   Felber K, Elks PM, Lecca M, Roehl HH. Expression of osterix Is Regulated by FGF and Wnt/beta-Catenin Signalling during Osteoblast Differentiation. *PLoS One* **10**, e0144982 (2015).

24.   Wolff C, Roy S, Ingham PW. Multiple muscle cell identities induced by distinct levels and timing of hedgehog activity in the zebrafish embryo. *Curr Biol* **13**, 1169-1181 (2003).

25.   Pan S, *et al*. Discovery of NVP-LDE225, a Potent and Selective Smoothened Antagonist. *ACS medicinal chemistry letters* **1**, 130-134 (2010).

26.   Li Y, Trush MA. Diphenyleneiodonium, an NAD(P)H oxidase inhibitor, also potently inhibits mitochondrial reactive oxygen species production. *Biochem Biophys Res Commun* **253**, 295-299 (1998).

27.   Yoo SK, Starnes TW, Deng Q, Huttenlocher A. Lyn is a redox sensor that mediates leukocyte wound attraction in vivo. *Nature* **480**, 109-112 (2011).

28.   Martin BL, Kimelman D. Canonical Wnt signaling dynamically controls multiple stem cell fate decisions during vertebrate body formation. *Developmental cell* **22**, 223-232 (2012).

29.   Wilson V, Olivera-Martinez I, Storey KG. Stem cells, signals and vertebrate body axis extension. *Development* **136**, 1591-1604 (2009).

30.   Cunningham TJ, Kumar S, Yamaguchi TP, Duester G. Wnt8a and Wnt3a cooperate in the axial stem cell niche to promote mammalian body axis extension. *Dev Dyn* **244**, 797-807 (2015).

31.   Zhao X, Duester G. Effect of retinoic acid signaling on Wnt/beta-catenin and FGF signaling during body axis extension. *Gene expression patterns : GEP* **9**, 430-435 (2009).

32.   Varga ZM, *et al.* Zebrafish smoothened functions in ventral neural tube specification and axon tract formation. *Development* **128**, 3497-3509 (2001).

33.   Mich JK, Chen JK. Hedgehog and retinoic acid signaling cooperate to promote motoneurogenesis in zebrafish. *Development* **138**, 5113-5119 (2011).

34.   Heber-Katz E. Oxygen, Metabolism, and Regeneration: Lessons from Mice. *Trends Mol Med* **23**, 1024-1036 (2017).

35.   Pfefferli C, Jazwinska A. The careg element reveals a common regulation of regeneration in the zebrafish myocardium and fin. *Nat Commun* **8**, 15151 (2017).

36.   Taniguchi Y, Watanabe K, Mochii M. Notochord-derived hedgehog is essential for tail regeneration in Xenopus tadpole. *BMC developmental biology* **14**, 27 (2014).

37.   Schnapp E, Kragl M, Rubin L, Tanaka EM. Hedgehog signaling controls dorsoventral patterning, blastema cell proliferation and cartilage induction during axolotl tail regeneration. *Development* **132**, 3243-3253 (2005).

38.   Wang J, Cao J, Dickson AL, Poss KD. Epicardial regeneration is guided by cardiac outflow tract and Hedgehog signalling. *Nature* **522**, 226-230 (2015).

39.   Singh BN, Doyle MJ, Weaver CV, Koyano-Nakagawa N, Garry DJ. Hedgehog and Wnt coordinate signaling in myogenic progenitors and regulate limb regeneration. *Developmental biology* **371**, 23-34 (2012).

40.   Mateus R, *et al.* In vivo cell and tissue dynamics underlying zebrafish fin fold regeneration. *PLoS One* **7**, e51766 (2012).

41.   Bement WM, Mandato CA, Kirsch MN. Wound-induced assembly and closure of an actomyosin purse string in Xenopus oocytes. *Curr Biol* **9**, 579-587 (1999).

42.   Xu S, Chisholm AD. C. elegans epidermal wounding induces a mitochondrial ROS burst that promotes wound repair. *Developmental cell* **31**, 48-60 (2014).

43.   LeBert D, *et al.* Damage-induced reactive oxygen species regulate vimentin and dynamic collagen-based projections to mediate wound repair. *Elife* **7**, (2018).

44.   Iliev AI, *et al.* Rapid microtubule bundling and stabilization by the Streptococcus pneumoniae neurotoxin pneumolysin in a cholesterol-dependent, non-lytic and Src-kinase dependent manner inhibits intracellular trafficking. *Mol Microbiol* **71**, 461-477 (2009).

45.   Katagiri K, Katagiri T, Kajiyama K, Yamamoto T, Yoshida T. Tyrosine-phosphorylation of tubulin during monocytic differentiation of HL-60 cells. *J Immunol* **150**, 585-593 (1993).

46.   Marie-Cardine A, Kirchgessner H, Eckerskorn C, Meuer SC, Schraven B. Human T lymphocyte activation induces tyrosine phosphorylation of alpha-tubulin and its

association with the SH2 domain of the p59fyn protein tyrosine kinase. *Eur J Immunol* **25**, 3290-3297 (1995).