Depletion of limiting rDNA structural complexes triggers chromosomal

instability and replicative aging of *Saccharomyces cerevisiae*

Ryan D. Fine[1], Nazif Maqani[1], Mingguang Li[1,2], Elizabeth Franck[1], and Jeffrey S. Smith[1,*]

[1]Department of Biochemistry and Molecular Genetics, University of Virginia School of

Medicine, Charlottesville, VA 22908. [2]Department of Laboratory Medicine, Jilin Medical

University, Jilin, 132013, China

[*]Corresponding Author

Department of Biochemistry and Molecular Genetics

University of Virginia School of Medicine

Pinn Hall, Box 800733

Charlottesville, VA 22908

Phone: 434-243-5864

Fax: 434-924-5069

Email: jss5y@virginia.edu

Keywords: Sir2, cohesin, RENT, monopolin, yeast, replicative lifespan, aging, rDNA

Abstract

Sir2 is a highly conserved NAD^+-dependent histone deacetylase that functions in heterochromatin formation and promotes replicative lifespan (RLS) in the budding yeast, *Saccharomyces cerevisiae*. Within the yeast rDNA locus, Sir2 is required for efficient cohesin recruitment and maintaining stability of the tandem array. In addition to the previously reported depletion of Sir2 in replicatively aged cells, we discovered that subunits of the Sir2 containing complexes, SIR and RENT, were depleted. Several other rDNA structural protein complexes also exhibited age-related depletion, most notably the cohesin complex. We hypothesized that mitotic chromosome instability (CIN) due to cohesin depletion could be a driver of replicative aging. ChIP assays of the residual cohesin (Mcd1-13xMyc) in moderately aged cells showed strong depletion from the rDNA and initial redistribution to the point centromeres, which was then lost in older cells. Despite the shift in cohesin distribution, sister chromatid cohesion was partially attenuated in aged cells and the frequency of chromosome loss was increased. This age-induced CIN was exacerbated in strains lacking Sir2 and its paralog, Hst1, but suppressed in strains that stabilize the rDNA array due to deletion of *FOB1* or through caloric restriction (CR). Furthermore, ectopic expression of *MCD1* from a doxycycline-inducible promoter was sufficient to suppress rDNA instability in aged cells and to extend RLS. Taken together we conclude that age-induced depletion of cohesin and multiple other nucleolar chromatin factors destabilize the rDNA locus, which then results in general CIN and aneuploidy that shortens RLS.

Introduction

Budding yeast replicative lifespan (RLS) was originally described decades ago as the number of times a mother cell divides before losing viability (MORTIMER AND JOHNSTON 1959), and has been an effective model system for the identification and/or characterization of several conserved aging-related genes and pathways, including *SIR2*, AMPK (Snf1), and TOR signaling (WASKO AND KAEBERLEIN 2014). *SIR2* is possibly the most famous yeast gene associated with replicative aging and encodes the founding family member of the NAD^+-dependent histone/protein deacetylases, commonly known as sirtuins (reviewed in (BUCK *et al.* 2004)). The NAD^+ dependence of sirtuins provides a direct link between metabolism and cellular processes regulated by these enzymes. In fact, recent evidence points to depletion of cellular NAD^+ pools as a potential mechanism for aging-associated disease, which could be mediated by impairment of sirtuins or other NAD^+ consuming enzymes (GOMES *et al.* 2013). Therefore, understanding how sirtuins are impacted by aging and how they regulate age-altered cellular processes is of intense interest.

Eukaryotic genomes generally encode for several sirtuin homologs. The *Saccharomyces cerevisiae* genome, for example, encodes *SIR2* and four additional Homologs of Sir Two (*HST1-HST4*) (BRACHMANN *et al.* 1995; DERBYSHIRE *et al.* 1996). Sir2 and its fellow Silent Information Regulator proteins, Sir1, Sir3, and Sir4 were originally shown to establish and maintain silencing of the silent mating loci, *HML* and *HMR* (RINE AND HERSKOWITZ 1987). These proteins form the so-called SIR complex that is recruited to and then spreads across the *HM* loci and telomeres to form hypoacetylated heterochromatin-like domains (reviewed in (GARTENBERG AND SMITH 2016)). Sir2 is required for replicative longevity and its abundance is significantly reduced in replicatively aged yeast cells (DANG *et al.* 2009), presenting a possible mechanism for the

decline of Sir2-dependent processes during aging. including gene silencing. Indeed, the depletion

of Sir2 in aged cells causes hyperacetylated H4K16 and silencing defects at subtelomeric loci

(DANG *et al.* 2009). It has been reported that aged cells become sterile (mating-incompetent) due

to loss of silencing at *HML* and *HMR* (SMEAL *et al.* 1996), which results in co-expression of the

normally repressed α1/α2 and a1/a2 transcription factor genes encoded at these loci. In theory,

this should induce a diploid-like or "pseudodiploid" gene expression pattern and sterility, as is

observed for a silencing defective *sir2Δ* mutant (RINE AND HERSKOWITZ 1987). However, more

recent experiments point toward a silencing-independent mechanism of sterility, whereby

aggregation of the Whi3 protein in aged cells makes them insensitive to pheromones (SCHLISSEL

et al. 2017).

Alternative models for Sir2 control of RLS have focused on the rDNA tandem array

where Sir2 is important for cohesin recruitment (KOBAYASHI *et al.* 2004; GANLEY AND

KOBAYASHI 2014). Cohesin association with the rDNA also requires Tof2 and the Lrs4/Csm1

(cohibin) complex (HUANG *et al.* 2006). Sir2 silences RNA polymerase II-dependent

transcription at the rDNA locus via a nucleolar HDAC complex called RENT (BRYK *et al.* 1997;

SMITH AND BOEKE 1997), consisting of Sir2, Net1, and Cdc14 subunits (SHOU *et al.* 1999;

STRAIGHT *et al.* 1999). Specifically, RENT represses transcription of endogenous non-coding

RNAs from the intergenic spacer (IGS) regions (LI *et al.* 2006). Derepression of the bidirectional

promoter (E-pro) within IGS1 in *sir2Δ* cells displaces cohesin from the rDNA, thus destabilizing

the array by making it more susceptible to unequal sister chromatid exchange (KOBAYASHI AND

GANLEY 2005). Mild Sir2 overexpression, on the other hand, enhances silencing, suppresses

recombination between repeats, and extends RLS (SMITH *et al.* 1998; KAEBERLEIN *et al.* 1999).

Extrachromosomal rDNA circles (ERCs) derived from these unequal recombination events specifically accumulate to high levels in old mother cells (SINCLAIR AND GUARENTE 1997), where they can interfere with G1 cyclin expression (NEUROHR *et al.* 2018). Such an ERC-centric model is supported by RLS extension of *fob1Δ* strains (DEFOSSEZ *et al.* 1999). Fob1 binds to the rDNA at IGS1 to block DNA replication forks from colliding with elongating RNA polymerase I molecules (KOBAYASHI AND HORIUCHI 1996). The blocked forks can collapse, resulting in double-stranded DNA breaks that trigger unequal sister chromatid exchange (TAKEUCHI *et al.* 2003). The frequency of rDNA recombination and ERC production is reduced in a *fob1Δ* mutant due to loss of the fork block, thus extending RLS (DEFOSSEZ *et al.* 1999). More recently, this rDNA-centric model of aging has been extended to include general rDNA instability having negative effects on genome integrity, including ERC accumulation, and is also considered a critical contributor to aging (GANLEY AND KOBAYASHI 2014).

In addition to promoting cohesin recruitment to the rDNA, Sir2 is also required for establishing sister chromatid cohesion (SCC) at *HML* and *HMR* (CHANG *et al.* 2005; WU *et al.* 2011). Moreover, we previously observed significant overlap between Sir2 and cohesin at additional binding sites throughout the genome (LI *et al.* 2013). Outside heterochromatin, the cohesin loading complex (Scc2/Scc4) deposits cohesin (Mcd1, Irr1, Smc1, Smc3) onto centromeres and other cohesion associated regions (CARs) (CIOSK *et al.* 2000; KOGUT *et al.* 2009), in order to maintain SCC until anaphase, when Mcd1 is cleaved by separase to facilitate sister chromatid separation (reviewed in (MARSTON 2014)). Cohesin defects therefore result in chromosome instability (CIN) due to improper chromosome segregation (reviewed in (WOOD *et al.* 2010)). A previous study found that cells deleted for *SIR2* have a CIN phenotype related to hyperacetylation of H4K16 (CHOY *et al.* 2011), though the functional relationship to cohesin was

not considered. Given the natural depletion of Sir2 from replicatively aging yeast cells (DANG *et*

al. 2009), we hypothesized the frequency of CIN should increase with age. Here, we establish

that CIN is indeed more frequent in aged cells and is associated with SCC defects. We go on to

show that like Sir2, cohesin subunits are depleted from aged mother cells, providing a likely

reason for problems with SCC. Interestingly, despite the overall reduction in cohesin protein

levels, the chromosomal distribution of cohesin enrichment was not uniform. In moderately aged

mother cells, enrichment at the rDNA was drastically reduced, while binding at centromeres was

not, suggesting a mechanism by which SCC is preferentially maintained at centromeres to ensure

cell viability. However, this comes at the expense of chronic rDNA instability that is exacerbated

by additional age-induced reductions in the RENT and cohibin/monopolin complexes. The

defects in rDNA stability can be suppressed by overexpressing the Mcd1 subunit of cohesion,

which also extends RLS, thus making cohesin a dose dependent longevity factor. Lastly, the age

associated CIN phenotype is suppressed by deleting *FOB1* or by CR growth conditions,

suggesting a model whereby rDNA instability on chromosome XII caused by RENT, cohibin,

and cohesin depletion drives the mitotic segregation defects of other chromosomes during

replicative aging.

Methods

Yeast strains, plasmids, and media

Yeast strains were grown at 30°C in Yeast Peptone Dextrose (YPD) or Synthetic

Complete (SC) medium for strains bearing plasmids (MATECIC *et al.* 2010). *SIR2, HST1*, or

FOB1 open reading frames were disrupted with one-step PCR-mediated gene replacement using

kanMX4, natMX4, or *hphMX4* drug resistance markers, respectively. The *HMR* deletion by

replacement with *hphMX4* spans sacCer3 genome chrIII coordinates 293170-294330. All C-

terminally 13xMyc (EQKLISEEDL) tagged proteins were targeted at endogenous loci in the

Mother Enrichment Program (MEP) strains UCC5181 and UCC5179 (LINDSTROM AND

GOTTSCHLING 2009), followed by mating to generate homozygous diploids. All deletions and

fusions were confirmed by colony PCR, western blotting, or both. pRF4 was constructed by PCR

amplifying the *MCD1* open reading frame from ML1 genomic DNA and ligating into *Pst*I and

*Not*I sites of pCM252 (BELLI *et al.* 1998), a tetracycline inducible overexpression vector

(Euroscarf). pRF10 and pRF11 were constructed by removing the expression cassette by *Pvu*II

blunt end digestion of pCM252 and pRF4, respectively, and ligating it between the *Pvu*II sites of

pRS405 bearing *LEU2*, thus replacing the *TRP1* marker with *LEU2*. pRF10 and 11 were then

digested with *EcoRV* and integrated into the genome at *leu2Δ1*. pSB760 and pSB766 are

integrating and *2μ LEU2* vectors, respectively, bearing a single copy of *SIR2* (BUCK *et al.* 2002).

All strains used in this work are listed in Table S1 and all primers are listed in Table S2.

Isolation of aged yeast cells

Aged yeast cell enrichment was based on the MEP strain background (LINDSTROM AND

GOTTSCHLING 2009; LINDSTROM *et al.* 2011). For all assays, 1 μL of stationary phase culture

was inoculated into 100 mL of YPD medium and then grown into log phase. Approximately

1×10^8 cells were harvested, and centrifuged cell pellets washed 3 times with 1x phosphate

buffered saline (PBS). Cells were then resuspended in 1 mL PBS and mixed with 5 mg of Sulfo-

NHS-LC-Biotin (Pierce) per 1×10^8 cells for 30 min at room temperature. After biotin labeling,

5×10^7 cells were added to 1.5 L YPD cultures containing 1 μM estradiol, and 100 μg/mL

ampicillin to prevent bacterial contamination. These cultures were allowed to grow for 24 hr to

36 hr before processing in an assay specific manner (see below). For non-MEP strain

backgrounds, estradiol was not added to the cultures.

Western blotting

Two 1.5 L cultures were used for each western blot experiment, corresponding to

approximately 2×10^7 total aged cells after purification for each biological replicate. Cells were

pelleted using a Sorvall RC-5B Plus centrifuge with an SLA-3000 rotor at 2000 rpm, then

resuspended at a density of 6×10^8 cells/mL in RNAlater (Ambion) for 45 min in two separate

conical tubes. Following fixation, cells were pelleted and resuspended in 45 mL of cold PBS, 2

mM EDTA in 50 mL conical tubes. The mixture was incubated at 4°C for 30 min with 800 µL of

Streptavidin MicroBeads (Miltenyi Biotec), which were then purified through an autoMACS Pro

Cell Separator using the posseld2 program (UVA Flow Cytometry Core Facility). A 20 µL

aliquot of each output was used for bud scar counting using calcofluor white staining, before

combining the isolated samples into a single microfuge tube. Samples were frozen at -80°C

before protein extraction. Thawed cells were vortexed twice for 1 min in 20% TCA

(trichloroacetic acid) with ~100 µL of acid washed glass beads with a brief cooling period in

between vortexing. Beads were allowed to settle before transferring the supernatant to a fresh

microfuge tube. A 250 µL wash of 5% TCA was applied twice to the beads and pooled with the

initial lysis sample. Proteins were precipitated at 10,000 rpm in a microfuge for 5 min at 4°C.

Pellets were then resuspended in 50 µL of 1x SDS sample buffer (50 mM Tris-HCl pH 6.8, 2%

SDS, 10% Glycerol, 3.6 M 2-mercaptoethanol) and neutralized with 30 µL of 1M Tris-HCl, pH

8.0. Samples were run on a 9% (w/v) SDS-polyacrylamide gel and transferred to an Immobilon-

P membrane (Millipore). Membranes were incubated for 1 hour at room temperature in 1xTBST

+ 5% non-fat milk with primary antibodies (1:2000 α-Myc 9E10, 1:5000; α-Vma2 (Life

Technologies); 1:5000 α-Sir2 (Santa Cruz Biotechnology); 1:1000 α-Sir4 (Santa Cruz

Biotechnology); 1:1000 α-Sir3 (Santa Cruz Biotechnology). HRP-conjugated secondary

antibodies (Promega) were diluted 1:5000, and detected using chemiluminescence with HyGLO

(Denville Scientific). Quantitation was performed with ImageJ by using the rectangle tool to

outline protein bands and an equivalent sized box for background. After subtraction of

background, the signal of the aged cell band was divided by the signal of the Vma2 loading

control, and finally normalized to the quantity of the Vma2 normalized young cell band which

was arbitrarily set at 1.0.

ChIP Assays with aged and young cell populations

Two 1.5 L cultures were used for each biological replicate. After centrifugation, cells

were washed with 1xPBS then resuspended with 45 mL of PBS and incubated with 800 µL of

streptavidin microbeads, followed by sorting with the autoMACS Pro Cell Separator. Sorted

cells were immediately crosslinked with 1% formaldehyde for 20 min at room temperature, then

transferred to screw cap microcentrifuge tubes and the pellets flash frozen in liquid nitrogen.

Cells were thawed and lysed in 600 µL FA140 Lysis buffer (50 mM HEPES, 140 mM NaCl, 1%

Triton X-100, 1 mM EDTA, 0.1% SDS, 0.1 mM PMSF, 1x protease inhibitor cocktail; Sigma)

by shaking with acid-washed glass beads in a Mini-Bead beater (Biospec Products). Cell lysates

were recovered and sonicated for 30 cycles of 30 sec "on" and 30 sec "off" in a Diagenode

Bioruptor followed by centrifugation at 16,000 x g. A $1/10^{th}$ supernatant volume input was taken

for each sample and crosslinking reversed by incubating overnight at 65°C in 150 µL elution

buffer (TE, 1% SDS). The remaining supernatant was used for immunoprecipitation overnight at

4°C with 5 µg of primary antibody and 30 µL of protein G magnetic beads (Pierce), followed by

washing 1x with FA-140 buffer, 2x with FA-500 buffer (FA-140 with 500 mM NaCl), and 2x

with LiCl solution (10 mM Tris-HCl, pH 8.0, 250 mM LiCl, 0.5% NP-40, 0.5% SDS, 1 mM

EDTA). DNA was eluted twice with 75 µL of elution buffer in a 65°C water bath for 15 min.

The eluates were combined and crosslinking reversed. Input and ChIP DNA samples were

purified by an Invitrogen PureLink™ PCR purification kit. Finally, ChIP DNA was quantified by

real-time PCR and normalized to the input DNA signal. Young cells were collected flow-through

from the autoMACS cell sorter and then processed as described for the aged cells.

Sister chromatid cohesion assay

From 50 mL log phase SC cultures of strains 3349-1B, 3312-7A, 3460-2A, RF258,

RF278, and RF290, 5×10^7 cells were washed and biotinylated as described in the Isolation of

aged yeast cells section. This population was transferred into a 1.5 L SC culture and allowed to

grow for 14 hr. The original biotinylated cells were then purified by incubation with 300 µL of

streptavidin micro beads followed by gravity filtration through a Miltenyi LS column. The

column was washed twice with 5 ml of PBS and then processed as described below for young

cells.

From the original log phase culture, 5×10^7 cells were transferred to a fresh 50 mL SC

culture and arrested in mitosis with 10 µg/mL nocodazole for 1.5 hr. For the *mcd1-1* strain 3312-

7A, cells were also shifted to 37°C at this time. For bud scar staining, 5 mg of calcofluor white

was dissolved in 1 mL of PBS and any remaining aggregate removed by centrifugation. The 1

mL of soluble calcofluor was then added to the 50 mL SC culture. Non-arrested cells were

directly stained with calcofluor. Cells were then pelleted and washed in PBS. Following staining,

200 µL of 4% paraformaldehyde was added directly to the cell pellet and allowed to crosslink for

15 min at room temperature. The cell pellet was washed once with PBS and resuspended in

~100-200 µL of 0.1 MKPO$_4$/1 M sorbitol, pH 6.5. Images were captured with a Zeiss Axio

Observer z1 widefield microscope using a 64x oil objective lens.

Replicative lifespan assays

Lifespan assays were carried out as previously described (STEFFEN *et al.* 2009). Briefly,

small aliquots of log phase cultures were dripped in a straight line onto solid agar YPD with 2%

glucose. From the initial populations, a minimum of 32 virgin daughter cells were picked for

lifespan assays with daughter cells being selectively pulled away from mother cells using a

fiberoptic dissection needle and on a Nikon Eclipse 400 microscope. All virgin daughters were

required to bud at least one time to be included in the experiment and dissection was carried out

over the course of several days with temporary incubation at 4°C in between dissection periods

to stop division. Cells were considered dead when they stopped dividing for a minimum of 2

generation times (180 min). For p-values indicated in the text, a Wilcoxon rank-sum test was

conducted for respective lifespan assays using the basic wilcox.test function in R.

Mini-chromosome loss (sectoring) assay

The colony sectoring assay was performed on SC plates with adenine limited to 80 µM.

Frequency of mini-chromosome loss represents the number of ½ red/white sectored colonies

divided by the sum of sectored and white colonies. Cells were plated to an approximate density

of 500 cells/plate based on counts from a Brightline hemocytometer. Any plates bearing greater

than 1000 cells were discarded. Three biological replicates of each strain were performed, with at

least 10 plates counted per replicate. For aged cell populations, ~5×10^6 biotinylated cells were

aged in 1.5 L of YPD for 24 hr. Cells were incubated with 300 µL of streptavidin magnetic beads

(New England Biolabs) and manually washed 4 times with PBS on a magnetic stand, then plated

onto the limiting adenine SC plates such that ~500 colonies appeared. Bud scars were not

counted because the size of the beads prohibited visualization.

RT-qPCR measurement of *MCD1* overexpression

Doxycycline was added to log phase cultures at a concentration of 2 µg/mL for 4 hr in

order to induce expression of *MCD1*. Total RNA was extracted using a standard acid phenol

extraction protocol (AUSUBEL *et al.* 2000). cDNA was created from 1 µg of RNA using a Verso

cDNA synthesis kit (Thermo Fisher). *MCD1* expression levels were quantified on an Applied

Biosystems StepOne real time PCR machine with primers JS2844 and JS2949, and normalized to

actin transcript levels (primers JS1146 and JS1147).

Hi-C library construction

Log-phase cultures were cross-linked with 3% formaldehyde for 20 min and quenched

with a 2x volume of 2.5M glycine. Cell pellets were washed with dH_2O and stored at -80°C.

Thawed cells were resuspended in 5 ml of 1X NEB2 restriction enzyme buffer (New England

Biolabs) and poured into a pre-chilled mortar containing liquid N_2. Nitrogen grinding was

performed twice as previously described (BELTON AND DEKKER 2015), and the lysates were then

diluted to an OD_{600} of 12 in 1x NEB2 buffer. 500 µl of cell lysate was used for each Hi-C library

as follows. Lysates were solubilized by the addition of 50 µl 1% SDS and incubation at 65°C for

10 min. 55 µl of 10% Triton X-100 was added to quench the SDS, followed by 10 µl of 10X

NEB2 buffer and 15 µl of *Hin*dIII (New England Biolabs, 20 U/µl) to digest at 37°C for 2 hr. An

additional 10 µl of *Hin*dIII was added for digestion overnight. The remainder of the protocol was

based on previously published work with minor exceptions (BURTON *et al.* 2014). In short, ends

were filled in with dNTPs and biotinylated dCTP at 0.4 mM concentration using Klenow Exo-

(NEB) for 1 hr at 37°C. After a brief heat inactivation, blunt ends were ligated together in 3 mL

reaction volumes with T4 DNA-ligase for 6 hr at 16°C with a minimum DNA concentration of

0.5 ng/µL. Following ligation, cross-links were reversed at 70°C O/N and DNA was purified by

phenol/chloroform extraction and ethanol precipitation. Unligated-biotinylated ends were

removed using T4 DNA Polymerase (NEB). DNA was purified one final time with two

Zymogen DNA Clean and Concentrate-5 Kit columns per ligation reaction and eluted with 65

µL TE (130 µL total). Chromatin was quantitated with a Qubit fluorometer and then sheared

using a Diagenode Bioruptor. Hi-C sequencing libraries were prepared with reagents from an

Illumina Nextera Mate Pair Kit (FC-132-1001) using the standard Illumina protocol of End

Repair, A-tailing, Adapter Ligation, and 12 cycles of PCR. PCR products were size selected and

purified with AMPure XP beads before sequencing with an Illumina Miseq or Hiseq.

Hi-C data analysis

Iteratively corrected heatmaps appearing in this manuscript were produced using python

scripts from the Mirny lab hiclib library, publicly available at:

http://mirnylab.bitbucket.org/hiclib/index.html. Briefly, reads were mapped using the iterative

mapping program and then run through the fragment filtering program using the default

parameters. Raw heat maps were further filtered to remove diagonal reads and iteratively

corrected. Finally, the iteratively corrected heatmaps were normalized for read count differences

to make them comparable. The *cdc15-2* sample data was pulled from the SRA database at

SRP094582 (LAZAR-STEFANITA *et al.* 2017), and our data is available from GEO at GSE117037.

Data availability

Strains and plasmids are available upon request. Supplemental Tables S1 and S2, containing lists

of all strains and oligodeoxynucleotides used in the study, are included in the Supplemental

Information file. Supplemental Figures S1 through S7 are also included in the same file.

Results

Sir2 binding partners are depleted in aged yeast cells

Sir2 is a dosage dependent longevity factor, such that strains provided one extra *SIR2*

gene copy have an extended RLS (KAEBERLEIN *et al.* 1999). Mother cells experience a

progressive reduction of Sir2 protein during normal replicative aging that presumably contributes

to the aging process (DANG *et al.* 2009). Sir2 does not function in isolation, so we hypothesized

that protein levels of key Sir2-interacting partners could also be depleted with age. To isolate

sufficient quantities of aged cells for western blot assays we turned to <u>M</u>other <u>E</u>nrichment

<u>P</u>rogram (MEP) strains developed by the Gottschling lab (LINDSTROM AND GOTTSCHLING 2009).

The aged cell purification procedure was validated by increased average bud scar counts with

calcofluor white and the expected reduction of Sir2 protein (Figure 1A). The vacuolar protein

Vma2, used as a loading control for these western blots, does not deplete with age (LINDSTROM

et al. 2011). Since Sir2 is the catalytic subunit of both SIR and RENT (Figure 1B), it was

important to know which complexes were impacted by age. As shown in Figure 1C and D, Sir4

was strongly depleted in aged cells while Sir3 was actually enriched. Sir3 enrichment in aged

cells was also observed in an earlier proteomics screen (JANSSENS *et al.* 2015). Such a stark

difference was considered relevant because Sir2 and Sir4 interact as a heterodimer that associates

with the acetylated H4 N-terminal tail (MOAZED *et al.* 1997), while Sir3 is subsequently

recruited following H4K16 deacetylation to complete SIR holocomplex formation on

heterochromatin (OPPIKOFER *et al.* 2011). Myc-tagged Net1 (RENT complex) was also strongly

depleted from aged cells (Figure 1E), indicating that Sir2/Sir4 and the nucleolar RENT complex

are both depleted during aging. It should be noted that the Sir2 paralog, Hst1, which has the

capacity to compensate for loss of Sir2 (HICKMAN AND RUSCHE 2007; LI *et al.*), was also

partially depleted from aged cells (Figure 1F). Detailed bud scar and protein quantitations from

triplicate biological samples are reported in Table S3.

Chromosome instability (CIN) increases during replicative aging

325   Considering the observed depletion of multiple heterochromatin factors in aged yeast

cells, as well as the CIN phenotype of a *sir2Δ* mutant (CHOY *et al.* 2011), we next tested whether

aged cells had a CIN phenotype that was exacerbated by complete loss of *SIR2* and/or *HST1*.

Strains utilized for this experiment have an artificial chromosome III bearing a suppressor tRNA

gene, *SUP11* (SPENCER *et al.* 1990). Loss of the chromosome prevents suppression of an ochre

stop codon in *ade2-101*, resulting in the classic *ade2* red colony phenotype. The frequency of

nondisjunction events was measured by counting half-sectored red/white colonies from young

and aged cell populations (Figure 1G). Sectoring was elevated in young populations of *sir2Δ*

and *hst1Δ* mutants, and additively increased in a *sir2Δhst1Δ* double mutant (Figure 1H, black

bars, left side of panel). Interestingly, sectoring was significantly higher for aged populations of

each strain (Figure 1H, gray bars, left side of panel), suggesting additional age-associated factors

were involved. We next tested whether the *sir2Δ* effect on sectoring was related to a

pseudodiploid phenotype caused by derepression of the *HM* loci. This reporter strain background

was *MATα*, so we deleted *HMR* (chrIII 293170-294330) to eliminate the a1/a2 transcription

factors. Reversal of the pseudodiploid phenotype was confirmed by restoration of mating to the

sir2Δ hmrΔ strains (data not shown). Importantly, this manipulation significantly suppressed

sectoring of the young *sir2Δ* and *sir2Δ hst1Δ* mutants, but not the *hst1Δ* mutant (Figure 1H,

middle of panel), indicating there was indeed a *sir2Δ*-induced pseudodiploid effect that

contributed to mini-chromosome loss (Figure 1H, black bars, middle of panel). However, aging

still increased sectoring in each strain even when *HMR* was deleted (Figure 1H, gray bars,

middle of panel), suggesting that the aging-associated CIN factor was unrelated to mating type

control.

Because of the observed Net1 depletion (Figure 1E), we hypothesized that age-induced

mini-chromosome loss could be related to rDNA instability caused by loss of the RENT

complex. To address this idea, *FOB1* was deleted from WT, *sir2Δ*, *hst1Δ*, and *sir2Δ hst1Δ*

reporter strains to stabilize the rDNA, followed by retesting the sectoring phenotypes. As shown

in Figure 1H (right side of panel), frequency of sectoring observed for each aged *fob1Δ* strain

was generally similar to that observed with young *FOB1*[+] versions of the strains (Figure 1H,

black bars, left side of panel), suggesting that destabilization of the rDNA during aging does

contribute to the instability of other chromosomes. The *hmrΔ* and *fob1Δ* mutations were

potentially suppressing chromosome loss through independent mechanisms, thus begging the

question of whether combining them would fully suppress CIN in a *sir2Δ hst1Δ hmrΔ fob1Δ*

quadruple mutant. Remarkably, aged cells from this mutant combination lost the mini

chromosome marker at a very low rate comparable to young WT cells, with no statistical

difference (Figure 1H, 1-way ANOVA).

Cohesin levels are depleted in aged yeast cells

The above results raised the question of what factor(s) related to rDNA stability,

chromosome segregation, and Sir2, was becoming defective in aged cells. Cohesin perfectly fit

this profile and was earlier shown in mammalian oocytes to become depleted with age (reviewed

in (JESSBERGER 2012)). To test whether cohesin was another Sir2-linked factor depleted during

yeast aging, we C-terminally myc-tagged the Mcd1 or Smc1 subunits (Figure 2A) in the MEP

strain background. Western blotting demonstrated that both subunits were significantly depleted

from aged cells (Figure 2B), implying depletion of the whole cohesin complex. We also

observed depletion of the cohibin/monopolin subunit Lrs4 (Figure S2), which can function as a

cohesin clamp at the rDNA (HUANG *et al.* 2006). Furthermore, a Myc-tagged Scc2 subunit of the

Scc2/Scc4 cohesin loading complex was age-depleted (Figure 2C), though not as severely as the

cohesin complex, predicting that the remaining cohesin complex could still be loaded onto

chromatin in aged cells. ChIP assays for Mcd1-myc in MEP cells aged for 24 hr demonstrated

strong depletion from the rDNA intergenic spacer (IGS1) (Figure 2D), but was surprisingly

enriched at the centromeres of chromosomes XI and III. This effect did not appear to

significantly extend into the pericentric region of chromosome III (Figure 2E). In order to rule

out primer specificity or chromosome size effects, we tested three centromeres from other

chromosomes and observed the same trend for each of them (Figure 2F). To confirm the

enrichment was centromere specific, we tested two additional sites on chromosome IV that were

15kb and 200kb away from the centromere, and observed no increase in the aged samples

compared to young (Figure 2G). To test if the enrichment of Mcd1-myc at centromeres in aged

cells was specific to cohesin, and not a ChIP artifact, we next tested the distribution of Sir4-myc,

which was also depleted from aged cells (Figure 1C). In this case, Sir4-myc was depleted from

TELXV in aged cells (one of its normal targets) without any apparent redistribution to

centromeres (*CEN4*) or the rDNA (Figure 2H), indicating that not all age-depleted proteins

become enriched at centromeres. Based on these results, we hypothesized that as cohesin starts

depleting during replicative aging, its association with the rDNA is severely affected while a

significant portion of the remaining complex is retained and potentially redistributed to

centromeres. This is consistent with an earlier finding that cohesin preferentially associates with

pericentromeric regions instead of chromosome arms when Mcd1 expression is artificially

reduced below 30% of normal (HEIDINGER-PAULI *et al.* 2010). We next asked if the centromere

enrichment would be lost in an older cell population by extending growth of the MEP culture

time course to 36 hr. As shown in Figure 2I, the Mcd1-Myc association with *CEN4* and *CEN14*

returned to the original levels observed in young cells, while remaining depleted at the rDNA

array (Fig 2J). It is important to note that in a recent report (PAL *et al.* 2018), cohesin enrichment

at centromeres was actually reduced in very old yeast cells (>25 generations). Combined with

our findings, a model emerges whereby cohesin enrichment at centromeres is initially enhanced

during aging, but then catastrophically lost as cells approach the end of their lifespan.

Sister chromatid cohesion is compromised in aged yeast cells

Previous work found that SCC was surprisingly normal despite the forced reduction

(<30%) of Mcd1 protein levels (HEIDINGER-PAULI *et al.* 2010), leading us to ask whether

cohesion would be maintained in our moderately aged yeast cells that were also depleted for

cohesin, yet still showed enrichment at centromeres. To this end, we utilized strains with a LacO

array located approximately 10 kb away from centromere IV (*CEN4*) as a proxy for centromeric

cohesion, or at the *LYS4* locus on chromosome IV, located approximately 400 kb away from the

centromere, as a proxy for arm cohesion (UNAL *et al.* 2008; GUACCI AND KOSHLAND 2012).

Differential positioning of the array had no significant impact on RLS (Figure S3). SCC was

monitored by LacI-GFP appearing either as one dot in the case of cohesion maintenance or two

dots in the case of cohesion loss. Using an *mcd1-1* temperature sensitive mutant as a positive

control (GUACCI *et al.* 1997), we observed a large increase in two dots when cells were

synchronized in mitosis with nocodazole and shifted to 37°C (Figure 3A and B). WT cells for the

equivalent reporter strain were next biotinylated and aged for 24 hours, followed by purification

with magnetic streptavidin beads and arrest with nocodazole. Analyzing cells with >7 bud scars,

which is older than the average bud scar count of our western blot experiments, revealed a mild

loss of cohesion (Figure 3C and D). A similar analysis was then performed without the use of

nocodazole to rule out any side-effects related to triggering the mitotic spindle checkpoint. To

avoid misinterpreting anaphase events as lost SCC, we C-terminally tagged the spindle pole body

subunit Spc42 with dsRED, and only counted GFP dots from large-budded cells where the

spindle pole bodies had not separated between the mother and daughter cell. With this analysis,

the fold-change of SCC defect between young and aged cells was more pronounced (Figure 3E

and F), but the absolute frequency of loss was still much weaker than observed with the *mcd1-1*

mutant, which was not surprising given that mean lifespan of this strain background is ~16

generations (Figure S3). In very old cells (>25 bud scars) of a different strain background, an

independent study observed SCC loss on chromosome XII at a frequency that approached 50%

(PAL *et al.* 2018) suggesting that CIN may become most severe in the oldest cells where cohesin

is not only depleted from the rDNA, but also from centromeres.

RLS is modulated by Mcd1 expression levels

Since Sir2 and cohesin are both naturally depleted from replicatively aging yeast cells

(Figures 1A and 2B), and mild Sir2 overexpression extends RLS (KAEBERLEIN *et al.* 1999), we

hypothesized that manipulating cohesin expression levels would also impact CIN and RLS in a

dose-dependent manner. We initially attempted to overexpress the Mcd1 subunit from a

galactose inducible *GAL1* promoter and then measure mini-chromosome loss frequency by

counting ½ sectored colonies. However, simply growing the reporter strain in galactose-

containing media, even with an empty expression cassette, resulted in severe mini-chromosome

loss compared to glucose-containing media (Figure 4A and B). This effect was specific to

galactose, as growth with another non-preferred carbon source (raffinose) had no effect on

sectoring (Figure 4A and B). Though not useful for assaying the effects of *MCD1*

overexpression, it was still possible that the unexpectedly high CIN phenotype would correlate

with reduced RLS. We therefore measured RLS with the mini-chromosome reporter strain on

YEP plates with 2% glucose, galactose, or raffinose. As shown in Figure 4C, galactose strongly

decreased the mean RLS by ~50% compared to glucose (9.2 vs. 18.9 divisions, $p<1.0 \times 10^{-15}$),

while raffinose only had a marginal effect (15.5 divisions, $p<0.01$). To confirm the galactose

effect on RLS was not specific to the mini-chromosome strain, lifespan assays were repeated

with the commonly used strains BY4741 (*MAT*a) and BY4742 (*MAT*α). Again, a significant

decrease in mean lifespan was observed for BY4741 (17.7 divisions) and BY4742 (18.9

divisions) on galactose as compared to glucose (24.3 and 24.2 divisions, respectively) (Figure

S4, p<0.001), suggesting that galactose triggers a high rate of CIN through an unknown

mechanism that also shortens RLS.

To circumvent the use of galactose for *MCD1* overexpression we turned to an inducible

"Tet-On" promoter that is activated by doxycycline (BELLI *et al.* 1998). Strains harboring this

integrated cassette transcriptionally overexpressed *MCD1* approximately 2- to 7-fold compared

to the empty vector control (Figure S5A and B). In the mini-chromosome reporter strain, *MCD1*

overexpression significantly reduced the frequency of ½ sectored colonies in both young and

aged populations (Figure 4D), in agreement with *MCD1* being isolated as a high copy suppressor

of CIN using a different reporter system (ZHU *et al.* 2015). Next, *MCD1* was overexpressed in a

strain containing an ochre stop codon in the *MCD1* open-reading frame (*mcd1L12STOP*) that

reduces Mcd1 protein to ~30% of normal (HEIDINGER-PAULI *et al.* 2010). With an empty

pCM252 control *CEN* vector, *mcd1L12STOP* exhibited a 40% reduction in mean RLS (9.9

divisions) compared to an isogenic WT control (16.5 divisions), indicating that Mcd1 depletion

shortens RLS (Figure 4E, p<0.001). Overexpressing *MCD1* almost fully restored longevity to the

mutant (14.7 divisions) and, remarkably, also extended RLS of the control WT strain to 19.6

divisions (p<1.0×10^{-7}), which was primarily due to improved survival during the first ~15

divisions and then followed by a steeper decline in viability (Figure 4E). This biphasic pattern

was highly reproducible and suggested that the *CEN MCD1* OE plasmid was being lost around

mid-life due to increased chromosome loss, as previously seen in aged cells of our CIN assay

(Figure 1H). To account for this potential variable, we also integrated the empty and *MCD1* OE

vectors into a different FY834 strain background related to the long lived BY4741/42

background (WINSTON *et al.* 1995; BRACHMANN *et al.* 1998). Not only did *MCD1* OE extend the

mean RLS (28.0 divisions) in this background compared to the control (21.0 divisions), but the

maximum number of divisions was also increased by 25% (Figure S6, $p<1.0 \times 10^{-7}$). We therefore

conclude that similar to Sir2, Mcd1 is a strong dose-dependent longevity factor.

Sir2 functions upstream of cohesin for RLS (KOBAYASHI et al. 2004; KOBAYASHI AND

GANLEY 2005), and for SCC at HMR (WU et al. 2011), implying that Sir2 is upstream of cohesin

loading and function. However, this relationship could potentially be more complex since both

factors are depleted with age. To explore this further, we next tested if SIR2 overexpression

could rescue the short RLS of an Mcd1-depleted mcd1L12STOP strain by integrating a second

copy of SIR2 (2xSIR2) at the LEU2 locus. As shown in Figure 4F, 2xSIR2 partially rescued mean

RLS of the mcd1L12STOP strain (14.1 versus 9.9 divisions, $p<1.0 \times 10^{-4}$), and also increased

maximum RLS of the WT strain. Reciprocally, we asked if MCD1 overexpression would

suppress the short RLS of a sir2Δ or sir2Δ hst1Δ mutant. The double mutant was included to rule

out any redundancy between the two sirtuins. Mean RLS was clearly not increased by MCD1

overexpression as compared to empty vector for either the single (10.7 versus 10.2) or double

mutant (10.1 versus 9.7 divisions, Figure 4G), confirming that SIR2 was required for MCD1 in

regulating RLS. Interestingly, the sir2Δ hst1Δ fob1Δ hmrΔ quadruple mutant combination, which

suppressed CIN in aged cells (Figure 1H), fully restored RLS of the short lived sir2Δ hst1Δ

combination to WT levels (Figure 4G, p=n.s.), suggesting that CIN is a key driver of replicative

aging downstream of Sir2 and Hst1.

Considering the strong depletion of cohesin from rDNA in aged cells (Figure 2D), and

extended RLS when MCD1 was overexpressed (Figure 4E), we next tested if age-induced rDNA

instability was suppressed by MCD1 overexpression using a reporter strain harboring ADE2 in

the rDNA array (KAEBERLEIN et al. 1999). There was a large increase of red/white ½ sectoring

(marker loss) from aged cells that was suppressed upon MCD1 overexpression (Figure 4H). In

the absence of *SIR2*, ½ sectoring was high from young and aged cells when the empty vector

(pRF10) was integrated, and *MCD1* overexpression did not significantly reduce rDNA instability

in either population (Figure 4H), indicating that at least some Sir2 was required for Mcd1 to

impact rDNA stability. We conclude that loss of Sir2 and cohesin in aging cells causes rDNA

array instability that generally exacerbates CIN.

RLS extension by CR correlates with improved chromosome stability

Reducing glucose concentration in the growth medium is effective at extending RLS and

is considered a form of caloric restriction (CR) for yeast (JIANG *et al.* 2000; LIN *et al.* 2002).

There have been several hypotheses put forth for the underlying mechanisms, including

stabilization of the rDNA, as CR suppresses recombination within the rDNA (LAMMING *et al.*

2005; RIESEN AND MORGAN 2009; SMITH *et al.* 2009). Since *MCD1* overexpression suppressed

rDNA recombination and extended RLS, we hypothesized that CR may suppress the abbreviated

RLS of a cohesin-depleted *mcd1L12STOP* mutant strain. Indeed, CR extended RLS of both the

WT and *mcd1L12STOP* strains (Figure 5A, $p<0.01$ and $p<1.0 \times 10^{-7}$ respectively). However, the

suppression was apparently not due to maintenance of global cohesin levels because steady state

Mcd1-13xMyc was still depleted in glucose restricted aged cells (Figure 5B). CR also strongly

suppressed minichromosome loss in young and aged cells, even in the *sir2Δ hst1Δ* double mutant

(Figure 5C). Importantly, this CR effect also correlated with almost complete rescue of RLS for

the *sir2Δ hst1Δ* mutant (Figure 5D, p=n.s.). Taken together, the results support a mechanism for

RLS extension by CR, whereby stabilization of the rDNA locus helps maintain general mitotic

chromosome stability to protect against aneuploidy.

The rDNA array has an opportunity to interact with centromeres during anaphase

To conclude this study, we asked whether there is any mechanistic connection between

the rDNA and centromeres that could cause CIN. If rDNA instability has a direct effect on SCC

during aging, then we should observe elevated dissociation of sister chromatids in a *sir2Δ* mutant

and improved SCC in a *fob1Δ* mutant. However, as shown in Figure S7, the frequency of 2 GFP

dots in the SCC assay for these two mutants in aged populations was comparable to WT (see

Figure 3E and F). Alternatively, the rDNA could potentially affect centromere function through

direct contacts. Previous Hi-C analysis of the yeast genome and fluorescence microscopy of

nucleolar proteins positioned the rDNA off to one side of the nucleus, apparently secluded from

the rest of the genome (GOTTA *et al.* 1997; DUAN *et al.* 2010). The repetitive nature of rDNA

precludes it from appearing in Hi-C contact maps, but closer inspection of iteratively corrected

chromosome XII contact maps at 10 kb resolution indicated a clear interaction between unique

sequences flanking the centromere-proximal (left) edge of the array and *CEN12* (Figure 6A,

yellow arrow). We hypothesized that this contact was regulated by Sir2 since it is in the vicinity

of a known tRNA boundary (tQ(UUG)L) for rDNA silencing (BISWAS *et al.* 2009), but deletion

of *SIR2* had no effect on the contact (data not shown). Interestingly, all centromeres of the yeast

genome, including *CEN12*, cluster together in asynchronous cell population Hi-C data (Figure

6B, yellow arrows; (DUAN *et al.* 2010)), which potentially places them in proximity with the

rDNA given the association of *CEN12* with sequences flanking the rDNA. During anaphase, the

rDNA is thought to be separated from centromeres, but analysis of Hi-C data for cell-cycle

synchronized *cdc15-2*, which arrests cells in anaphase, revealed that the rDNA-proximal/*CEN12*

contact specifically occurs during anaphase (Figure 6C, (LAZAR-STEFANITA *et al.* 2017)), during

which time the centromeres are still clustered together by the spindle pole body (Figure 6D).

Taken together, these results suggest that the rDNA may transiently contact the centromeres

during mitosis, providing a potential window of time for a destabilized rDNA array to negatively

impact the integrity of general chromosome segregation during mitosis (Figure 6E).

Discussion

Nuclear protein depletion during replicative aging as a paradigm for aging pathologies

During this study the majority of chromatin-associated proteins that we analyzed by

western blotting were depleted in replicatively aged yeast cells. The only protein unaffected by

age that we tested, other than the vacuolar Vma2 control, was Sir3 (Figure 1D). A similar

proportion of homologous recombination proteins were depleted in an independent analysis of

aged cells, with Rad52 the only one tested that was not affected (PAL *et al.* 2018). These results

suggest there is at least some selectivity to the depletion of nuclear proteins in aged cells.

However, the large number of depleted factors also makes it likely that targeted nuclear protein

deficiency could lead to multiple age-associated phenotypes. Replicatively aging yeast cells

appear especially susceptible to this phenomenon, as even total core histone levels are depleted

(HU *et al.* 2014). Evidence also exists for histone depletion during aging of metazoan organisms,

including mammals (reviewed in (SONG AND JOHNSON 2018)). More generally, global protein

turnover is elevated in cells from prematurely aging progeria patients, which may trigger higher

translation rates (BUCHWALTER AND HETZER 2017). This is significant because reducing

translation is a means of extending lifespan in multiple organisms (MEHTA *et al.* 2010). The

mechanism(s) driving nuclear protein depletion in aged yeast mother cells or other organisms

remain unclear.

The specificity for Sir2/Sir4 depletion over Sir3 is intriguing given that Sir2 and Sir4

form a tight complex that allosterically stimulates the deacetylase activity of Sir2 (HSU *et al.*

2013), while Sir3 is a subunit of the SIR holocomplex (OPPIKOFER *et al.* 2011). Since Sir3 levels

are elevated in aged cells (Figure 1D; (JANSSENS *et al.* 2015)), it is likely to have a function

independent of the SIR complex during aging. The mechanism for Sir2/Sir4 depletion from aged

cells remains uncharacterized, though in non-aging cell populations the stability/turnover of Sir4,

but not Sir2, is mediated by the E3 ubiquitin ligase San1 (DASGUPTA *et al.* 2004), which has also

been implicated as a quality control E3 ligase for mutated/unfolded nuclear proteins (GARDNER

et al. 2005). Whether San1 controls Sir4 stability during aging remains unknown, but since Sir4

is more severely depleted than Sir2 in aged cells (Figures 1A and 2C), Sir4 could be selectively

depleted from the SIR complex, thus leaving Sir2 unprotected and subject to turnover through a

different mechanism. Consistent with this idea, we note that Sir4 is depleted independently of

Sir2 during extended G1 arrest (LARIN *et al.* 2015). Alternatively, Sir2/Sir4 could be equally

depleted as a complex from telomeres and the *HM* loci (not necessarily via San1), leaving the

nucleolar pool of Sir2/RENT as more resistant to aging. Under this scenario, protecting integrity

of the rDNA array could take precedence over other heterochromatic domains. Interestingly, the

Schizosaccharomyces pombe San1 ortholog has also been implicated in a chaperone-assisted

degradation pathway that functions in quality control of kinetochores to promote chromosome

stability (KRIEGENBURG *et al.* 2014).

Sir2 depletion in replicatively aged yeast cells is reminiscent of Sirt1 depletion in serially

passaged mouse embryonic fibroblasts (MEFs), which correlates with declining mitotic activity

(SASAKI *et al.* 2006). Sir2 and Sirt1 are both known to function in regulating DNA replication

origins (HOGGARD *et al.* 2018; UTANI AND ALADJEM 2018), and the effect of deleting *SIR2* on

early origin firing is thought to be mediated by competition for limiting factors with the repeated

rDNA origins (YOSHIDA *et al.* 2014). Furthermore, CR has been proposed to extend RLS by

reducing rDNA origin firing, which improves overall genome replication (KWAN *et al.* 2013).

This may help explain why CR can extend RLS and suppress CIN even when *SIR2* and *HST1* are

deleted. Conversely, depleted cohesin in old cells could potentially cause rDNA instability by

impacting DNA replication and repair.

593 A precarious balance between rDNA and centromeric cohesin

SCC ensures chromosomes are not segregated until the Mcd1/Scc1 cohesin subunit is

cleaved in response to a mitotic spindle checkpoint signal that all chromosomes are properly

attached to microtubules and aligned at the metaphase plate (reviewed in (MARSTON 2014)).

Cohesin is also critical during meiosis, and it is well established in mammals that SCC defects

occur in the oocytes of older mothers, causing meiotic chromosome missegregation events

during both anaphase I and II (JESSBERGER 2012). This phenomenon is believed to be a major

mechanism for increased aneuploidy risk that usually results in embryonic lethality, or in the

case of chromosome 21 trisomy, Down's syndrome. The meiotic cohesin subunit Rec8 is

depleted in the oocytes of older mice, as is Shugoshin (Sgo2), which normally protects/maintains

centromeric cohesin (LISTER *et al.* 2010). More recent experiments in *Drosophila* suggest that

oxidative stress in aged oocytes contributes to the SCC defects (PERKINS *et al.* 2016). Our results

in replicatively aging yeast cells reveal that aging-induced cohesin depletion and the resulting

chromosome missegregation can extend to mitotic cells. Though cohesin depletion or defects

have not been reported for aging mammalian somatic cells, the mitotic spindle checkpoint

protein BubR1 is depleted in dynamic somatic tissues such as spleen in aged mice (BAKER *et al.*

2004). Deficiency of this protein results in premature aging phenotypes (BAKER *et al.* 2004),

while overexpression extends lifespan (BAKER *et al.* 2013). This is similar to the effects we

observe with Mcd1 depletion and overexpression on yeast RLS. Interestingly, BubR1 is also a

deacetylation target of Sirt2, which appears to stabilize the protein and extend lifespan, thus

linking mitotic spindle checkpoint regulation to NAD^+ metabolism (NORTH *et al.* 2014). It

remains unclear if Sir2, Hst1, or other sirtuins regulate the yeast BubR1 ortholog, Mad3, or

additional checkpoint and kinetochore proteins.

SCC is the canonical function for cohesin, though the complex also functions in

establishing and regulating genome organization at the level of chromatin structure, gene

regulation, and double strand break (DSB) repair (reviewed in (UHLMANN 2016)). Among these

various processes, SCC at centromeres appears the most critical because artificial depletion of

Mcd1 to <30% of normal levels results in preferential cohesin binding to pericentromeric regions

rather than cohesin associated regions (CARs) on chromosome arms (HEIDINGER-PAULI *et al.*

2010). SCC was also well maintained in these strains at the expense of normal chromosome

condensation, DNA repair, and rDNA stability (HEIDINGER-PAULI *et al.* 2010). In aged yeast

cells, we observed relative enrichment of Mcd1-myc at centromeres as compared to loss at the

rDNA (IGS1) locus (Figures 2D and 2E) consistent with pericentromeric cohesin retention in the

artificially depleted system. Despite maintaining the cohesin complex at centromeres, SCC was

still slightly impaired in the aged cells, but only if we analyzed cells >7 generations old. We

suspect cohesin was reduced at centromeres in these older cells, which would be consistent with

the loss of centromeric Mcd1 enrichment when cells were aged longer for 36 hr, but it was also

likely that some of the centromere-localized Mcd1 was non-functional. These results are in line

with an independent study that analyzed significantly older mother cells (~25 generations) and

observed reduced cohesin enrichment at centromeres and severe loss of SCC (PAL *et al.* 2018).

Another recent study using single cell microfluidics found that chromosome loss was very

common just prior to the last cell division (NEUROHR *et al.* 2018). Collectively, the results

suggest that centromere-associated cohesin is preferentially retained during the initial stages of

replicative aging, but then eventually breaks down below a critical threshold in the oldest cells.

Numerous nuclear proteins are depleted in aged yeast cells, not just cohesin subunits, so

we hypothesize that defects in other nuclear processes mediated by such factors also contribute

to SCC defects and chromosome instability either directly or indirectly. The depleted cohesin

loading complex (Scc2/4) is an obvious candidate due to its role in loading cohesin at

centromeres and CARs. Similarly, the depleted cohibin complex (Figure S2) is proposed to act as

a cohesin clamp onto rDNA chromatin (HUANG *et al.* 2006), and also functions at centromeres to

maintain mitotic integrity (BITTO *et al.* 2015). Sir2 and Hst1 are also obvious candidates given

the earlier finding that H4K16 deacetylation at centromeres by Sir2 helps maintain chromosome

stability (CHOY *et al.* 2011). Part of this effect is apparently due to the pseudo-diploid phenotype

of a *sir2Δ* mutation, which has been previously shown to impact RLS (KAEBERLEIN *et al.* 1999).

Hst1 also binds centromeric DNA *in vitro* and *in vivo* (OHKUNI AND KITAGAWA 2011), though

the functional relevance of that association remains uncharacterized. The suppression of age-

associated mini-chromosome loss in the absence of *FOB1* clearly points to rDNA instability as

an unexpected source of general CIN. Such a relationship is reinforced by the observed depletion

of nucleolar proteins Net1 and Lrs4 in aged cells (Figures 2E and S2), both of which are required

for normal rDNA/nucleolar integrity and stable cohesin association with the rDNA (SMITH *et al.*

1999; STRAIGHT *et al.* 1999; HUANG *et al.* 2006).

How could destabilization of the rDNA locus result in general chromosome instability

and shortened RLS? As depicted in Figure 6, unique sequence flanking the rDNA contacts the

centromere of chromosome XII, thus placing it in proximity to other centromeres during

anaphase. Whether the actual rDNA genes contact centromeres remains unclear due to the

current limitations of Hi-C analysis with repetitive DNA. However, specific regions of the rDNA

were previously shown to associate with various non-rDNA chromosomal regions using an

anchored 4C approach (O'SULLIVAN *et al.* 2009). Furthermore, multiple nucleolar associated

domains have been identified in metazoan cells that copurify with nucleoli (MATHESON AND

KAUFMAN 2016). Loss of cohesin from the rDNA could potentially disrupt long-range

interactions with centromeres or non-centromeric regions of cohesin association that influence

chromosome integrity. One potential mechanism could be significant disruption of overall

chromosome condensation during mitosis, as cohesin appears to play a larger role in the DNA

looping associated with chromosome condensation in budding yeast than previously thought

(SCHALBETTER *et al.* 2017).

Interestingly, another class of nuclear factors depleted in aged yeast cells are several

DNA repair proteins (PAL *et al.* 2018). Consequently, the lack of proper DNA repair while the

rDNA becomes destabilized correlates with fragmentation of chromosome XII and the other

chromosomes (PAL *et al.* 2018). Rad52 foci also appear in aged cells indicating persistent DNA

damage (NEUROHR *et al.* 2018). It was proposed that accumulation of breaks and rearrangements

ultimately causes cell death during replicative aging. Such cells were significantly older (>25

divisions) compared to the cells in our study, which exhibited a maximum of 13 divisions after

24 hr. Alternatively, it is possible that these presumably random rearrangements disrupt normal

SCC, leading to CIN.

Aneuploidy as an aging mechanism

All 16 *S. cerevisiae* chromosomes harbor essential genes, so if a single chromosome is lost from a haploid yeast cell, then the affected mother or daughter cell should become inviable and no longer divide. Given the elevated frequency of chromosome loss during replicative aging, the chances of generating an inviable mother cell during a replicative aging assay increases after each subsequent division. Therefore, at least a portion of the replicative lifespan in haploid yeast cells is controlled by the ability to maintain all 16 chromosomes. Complete loss of a chromosome would not be an immediate viability issue for diploid cells, however, due to the chances of losing both homologs in a single mitosis being exceedingly rare. On the other hand, haploid strains that are disomic for individual chromosomes are often short lived, with longer chromosomes typically having larger effects (SUNSHINE *et al.* 2016). It was hypothesized that such strains suffer from proteotoxic stress due to inappropriate protein expression levels. Therefore, a similar mechanism could shorten RLS in a diploid strain that is trisomic for an individual chromosome, though this has not yet been tested. Aneuploidy is also a hallmark of aging in the germline (NAGAOKA *et al.* 2012), and somatic tissues of mammals (LUSHNIKOVA *et al.* 2011; BAKER *et al.* 2013), making it a conserved feature of aging from yeast to humans.

Another exciting feature of this study is the suppression of CIN by CR growth conditions that extend RLS. This effect was independent of the reduced cohesin levels in aged cells, and even improved RLS of the cohesin-depleted strains. Since SCC is normal in the cohesin-depleted strain (HEIDINGER-PAULI *et al.* 2010), we hypothesize that CR reinforces other processes that are defective due to reduced cohesin or other depleted factors that promote rDNA stability. Indeed, CR is known to suppress rDNA instability in yeast cells (RIESEN AND MORGAN 2009; SMITH *et*

al. 2009), and improve overall genome replication efficiency (KWAN *et al.* 2013). Hi-C analysis

also suggests there could be direct effects of rDNA structure on centromere function, which will

be a focus of future investigation.

Acknowledgments

We thank Dan Gottschling and all lab members for kindly providing yeast strains and initial

advice on the MEP system. Stefan Bekiranov, Job Dekker, Jon Belton, Maitreya Dunham, Ivan

Liachko, Maxim Imakev, and Anton Goloborodko all provided valuable advice on Hi-C

protocols and analysis methods. We thank Doug Koshland for providing the Mcd1 reduction and

cohesion assay strains, and Matt Kaeberlein for rDNA marker loss strains. We also thank Jef

Boeke, Marc Gartenberg, and Dan Gottschling for providing antibodies. Special thanks to Todd

Stukenberg for microscopy assistance. Lastly, thanks to David Auble for critically reading the

manuscript and providing comments prior to submission. We declare no conflicts of interest.

References

Ausubel, F. M., R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman *et al.* (Editors), 2000

Current Protocols in Molecular Biology. John Wiley & Sons, Inc., New York.

Baker, D. J., M. M. Dawlaty, T. Wijshake, K. B. Jeganathan, L. Malureanu *et al.*, 2013 Increased

expression of BubR1 protects against aneuploidy and cancer and extends healthy

lifespan. Nat Cell Biol 15**:** 96-102.

Baker, D. J., K. B. Jeganathan, J. D. Cameron, M. Thompson, S. Juneja *et al.*, 2004 BubR1

insufficiency causes early onset of aging-associated phenotypes and infertility in mice.

Nat Genet 36**:** 744-749.

Belli, G., E. Gari, L. Piedrafita, M. Aldea and E. Herrero, 1998 An activator/repressor dual

system allows tight tetracycline-regulated gene expression in budding yeast. Nucleic

Acids Res 26**:** 942-947.

Belton, J. M., and J. Dekker, 2015 Measuring chromatin structure in budding yeast. Cold Spring

Harb Protoc 2015**:** 614-618.

Biswas, M., N. Maqani, R. Rai, S. P. Kumaran, K. R. Iyer *et al.*, 2009 Limiting the extent of the

RDN1 heterochromatin domain by a silencing barrier and Sir2 protein levels in

Saccharomyces cerevisiae. Mol Cell Biol 29**:** 2889-2898.

Bitto, A., A. M. Wang, C. F. Bennett and M. Kaeberlein, 2015 Biochemical genetic pathways

that modulate aging in multiple species. Cold Spring Harb Perspect Med 5.

Brachmann, C. B., A. Davies, G. J. Cost, E. Caputo, J. Li *et al.*, 1998 Designer deletion strains

derived from *Saccharomyces cerevisiae* S288C: a useful set of strains and plasmids for

PCR-mediated gene disruption and other applications. Yeast 14**:** 115-132.

Brachmann, C. B., J. M. Sherman, S. E. Devine, E. E. Cameron, L. Pillus *et al.*, 1995 The *SIR2*

gene family, conserved from bacteria to humans, functions in silencing, cell cycle

progression, and chromosome stability. Genes Dev 9**:** 2888-2902.

Bryk, M., M. Banerjee, M. Murphy, K. E. Knudsen, D. J. Garfinkel *et al.*, 1997 Transcriptional

silencing of Ty1 elements in the *RDN1* locus of yeast. Genes Dev 11**:** 255-269.

Buchwalter, A., and M. W. Hetzer, 2017 Nucleolar expansion and elevated protein translation in

premature aging. Nat Commun 8**:** 328.

Buck, S. W., C. M. Gallo and J. S. Smith, 2004 Diversity in the Sir2 family of protein

deacetylases. J Leukoc Biol 75**:** 939-950.

Buck, S. W., J. J. Sandmeier and J. S. Smith, 2002 RNA polymerase I propagates unidirectional

spreading of rDNA silent chromatin. Cell 111**:** 1003-1014.

Burton, J. N., I. Liachko, M. J. Dunham and J. Shendure, 2014 Species-level deconvolution of

metagenome assemblies with Hi-C-based contact probability maps. G3 (Bethesda) 4**:**

1339-1346.

Chang, C. R., C. S. Wu, Y. Hom and M. R. Gartenberg, 2005 Targeting of cohesin by

transcriptionally silent chromatin. Genes Dev 19**:** 3031-3042.

Choy, J. S., R. Acuna, W. C. Au and M. A. Basrai, 2011 A role for histone H4K16

hypoacetylation in *Saccharomyces cerevisiae* kinetochore function. Genetics 189**:** 11-21.

Ciosk, R., M. Shirayama, A. Shevchenko, T. Tanaka, A. Toth *et al.*, 2000 Cohesin's binding to

chromosomes depends on a separate complex consisting of Scc2 and Scc4 proteins. Mol

Cell 5**:** 243-254.

Dang, W., K. K. Steffen, R. Perry, J. A. Dorsey, F. B. Johnson *et al.*, 2009 Histone H4 lysine 16

acetylation regulates cellular lifespan. Nature 459**:** 802-807.

Dasgupta, A., K. L. Ramsey, J. S. Smith and D. T. Auble, 2004 Sir Antagonist 1 (San1) is a

ubiquitin ligase. J Biol Chem 279**:** 26830-26838.

Defossez, P. A., R. Prusty, M. Kaeberlein, S. J. Lin, P. Ferrigno *et al.*, 1999 Elimination of

replication block protein Fob1 extends the life span of yeast mother cells. Mol Cell 3**:**

447-455.

Derbyshire, M. K., K. G. Weinstock and J. N. Strathern, 1996 *HST1*, a new member of the *SIR2*

family of genes. Yeast 12**:** 631-640.

Duan, Z., M. Andronescu, K. Schutz, S. McIlwain, Y. J. Kim *et al.*, 2010 A three-dimensional

model of the yeast genome. Nature 465**:** 363-367.

Ganley, A. R., and T. Kobayashi, 2014 Ribosomal DNA and cellular senescence: new evidence

supporting the connection between rDNA and aging. FEMS Yeast Res 14**:** 49-59.

Gardner, R. G., Z. W. Nelson and D. E. Gottschling, 2005 Degradation-mediated protein quality

control in the nucleus. Cell 120**:** 803-815.

Gartenberg, M. R., and J. S. Smith, 2016 The nuts and bolts of transcriptionally silent chromatin

in *Saccharomyces cerevisiae*. Genetics 203**:** 1563-1599.

Gomes, A. P., N. L. Price, A. J. Ling, J. J. Moslehi, M. K. Montgomery *et al.*, 2013 Declining

NAD^+ induces a pseudohypoxic state disrupting nuclear-mitochondrial communication

during aging. Cell 155**:** 1624-1638.

Gotta, M., S. Strahl-Bolsinger, H. Renauld, T. Laroche, B. K. Kennedy *et al.*, 1997 Localization

of Sir2p: the nucleolus as a compartment for silent information regulators. EMBO J 16**:**

3243-3255.

Guacci, V., and D. Koshland, 2012 Cohesin-independent segregation of sister chromatids in

budding yeast. Mol Biol Cell 23**:** 729-739.

Guacci, V., D. Koshland and A. Strunnikov, 1997 A direct link between sister chromatid

cohesion and chromosome condensation revealed through the analysis of *MCD1* in *S.*

cerevisiae. Cell 91**:** 47-57.

Heidinger-Pauli, J. M., O. Mert, C. Davenport, V. Guacci and D. Koshland, 2010 Systematic

reduction of cohesin differentially affects chromosome segregation, condensation, and

DNA repair. Curr Biol 20**:** 957-963.

Hickman, M. A., and L. N. Rusche, 2007 Substitution as a mechanism for genetic robustness: the

duplicated deacetylases Hst1p and Sir2p in *Saccharomyces cerevisiae*. PLoS Genet 3**:**

e126.

Hoggard, T. A., F. Chang, K. R. Perry, S. Subramanian, J. Kenworthy *et al.*, 2018 Yeast

heterochromatin regulators Sir2 and Sir3 act directly at euchromatic DNA replication

origins. PLoS Genet 14**:** e1007418.

Hsu, H. C., C. L. Wang, M. Wang, N. Yang, Z. Chen *et al.*, 2013 Structural basis for allosteric

stimulation of Sir2 activity by Sir4 binding. Genes Dev 27**:** 64-73.

Hu, Z., K. Chen, Z. Xia, M. Chavez, S. Pal *et al.*, 2014 Nucleosome loss leads to global

transcriptional up-regulation and genomic instability during yeast aging. Genes Dev 28**:**

396-408.

Huang, J., I. L. Brito, J. Villen, S. P. Gygi, A. Amon *et al.*, 2006 Inhibition of homologous

recombination by a cohesin-associated clamp complex recruited to the rDNA

recombination enhancer. Genes Dev 20**:** 2887-2901.

Janssens, G. E., A. C. Meinema, J. Gonzalez, J. C. Wolters, A. Schmidt *et al.*, 2015 Protein

biogenesis machinery is a driver of replicative aging in yeast. Elife 4**:** e08527.

Jessberger, R., 2012 Age-related aneuploidy through cohesion exhaustion. EMBO Rep 13**:** 539-

546.

Jiang, J. C., E. Jaruga, M. V. Repnevskaya and S. M. Jazwinski, 2000 An intervention

resembling caloric restriction prolongs life span and retards aging in yeast. FASEB J 14**:**

2135-2137.

Kaeberlein, M., M. McVey and L. Guarente, 1999 The SIR2/3/4 complex and SIR2 alone

promote longevity in *Saccharomyces cerevisiae* by two different mechanisms. Genes Dev

13**:** 2570-2580.

Kobayashi, T., and A. R. Ganley, 2005 Recombination regulation by transcription-induced

cohesin dissociation in rDNA repeats. Science 309**:** 1581-1584.

Kobayashi, T., and T. Horiuchi, 1996 A yeast gene product, Fob1 protein, required for both

replication fork blocking and recombinational hotspot activities. Genes Cells 1**:** 465-474.

Kobayashi, T., T. Horiuchi, P. Tongaonkar, L. Vu and M. Nomura, 2004 *SIR2* regulates

recombination between different rDNA repeats, but not recombination within individual

rRNA genes in yeast. Cell 117**:** 441-453.

Kogut, I., J. Wang, V. Guacci, R. K. Mistry and P. C. Megee, 2009 The Scc2/Scc4 cohesin

loader determines the distribution of cohesin on budding yeast chromosomes. Genes Dev

23**:** 2345-2357.

Kriegenburg, F., V. Jakopec, E. G. Poulsen, S. V. Nielsen, A. Roguev *et al.*, 2014 A chaperone-

assisted degradation pathway targets kinetochore proteins to ensure genome stability.

PLoS Genet 10**:** e1004140.

Kwan, E. X., E. J. Foss, S. Tsuchiyama, G. M. Alvino, L. Kruglyak *et al.*, 2013 A natural

polymorphism in rDNA replication origins links origin activation with calorie restriction

and lifespan. PLoS Genet 9**:** e1003329.

Lamming, D. W., M. Latorre-Esteves, O. Medvedik, S. N. Wong, F. A. Tsang *et al.*, 2005 *HST2*

mediates *SIR2*-independent life-span extension by calorie restriction. Science 309**:** 1861-

1864.

Larin, M. L., K. Harding, E. C. Williams, N. Lianga, C. Dore *et al.*, 2015 Competition between

heterochromatic loci allows the abundance of the silencing protein, Sir4, to regulate *de*

novo assembly of heterochromatin. PLoS Genet 11**:** e1005425.

Lazar-Stefanita, L., V. F. Scolari, G. Mercy, H. Muller, T. M. Guerin *et al.*, 2017 Cohesins and

condensins orchestrate the 4D dynamics of yeast chromosomes during the cell cycle.

EMBO J 36**:** 2684-2697.

Li, C., J. E. Mueller and M. Bryk, 2006 Sir2 represses endogenous polymerase II transcription

units in the ribosomal DNA nontranscribed spacer. Mol Biol Cell 17**:** 3848-3859.

Li, M., B. J. Petteys, J. M. McClure, V. Valsakumar, S. Bekiranov *et al.*, 2010 Thiamine

biosynthesis in *Saccharomyces cerevisiae* is regulated by the NAD$^+$-dependent histone

deacetylase Hst1. Mol Cell Biol 30**:** 3329-3341.

Li, M., V. Valsakumar, K. Poorey, S. Bekiranov and J. S. Smith, 2013 Genome-wide analysis of

functional sirtuin chromatin targets in yeast. Genome Biol 14**:** R48.

Lin, S. J., M. Kaeberlein, A. A. Andalis, L. A. Sturtz, P. A. Defossez *et al.*, 2002 Calorie

restriction extends *Saccharomyces cerevisiae* lifespan by increasing respiration. Nature

418**:** 344-348.

Lindstrom, D. L., and D. E. Gottschling, 2009 The mother enrichment program: a genetic system

for facile replicative life span analysis in *Saccharomyces cerevisiae*. Genetics 183**:** 413-

422, 411si-413si.

Lindstrom, D. L., C. K. Leverich, K. A. Henderson and D. E. Gottschling, 2011 Replicative age

induces mitotic recombination in the ribosomal RNA gene cluster of *Saccharomyces*

cerevisiae. PLoS Genet 7**:** e1002015.

Lister, L. M., A. Kouznetsova, L. A. Hyslop, D. Kalleas, S. L. Pace *et al.*, 2010 Age-related

meiotic segregation errors in mammalian oocytes are preceded by depletion of cohesin

and Sgo2. Curr Biol 20**:** 1511-1521.

Lushnikova, T., A. Bouska, J. Odvody, W. D. Dupont and C. M. Eischen, 2011 Aging mice have

increased chromosome instability that is exacerbated by elevated Mdm2 expression.

Oncogene 30**:** 4622-4631.

Marston, A. L., 2014 Chromosome segregation in budding yeast: sister chromatid cohesion and

related mechanisms. Genetics 196**:** 31-63.

Matecic, M., D. L. Smith, X. Pan, N. Maqani, S. Bekiranov *et al.*, 2010 A microarray-based

genetic screen for yeast chronological aging factors. PLoS Genet 6**:** e1000921.

Matheson, T. D., and P. D. Kaufman, 2016 Grabbing the genome by the NADs. Chromosoma

125**:** 361-371.

Mehta, R., D. Chandler-Brown, F. J. Ramos, L. S. Shamieh and M. Kaeberlein, 2010 Regulation

of mRNA translation as a conserved mechanism of longevity control. Adv Exp Med Biol

694**:** 14-29.

Moazed, D., A. Kistler, A. Axelrod, J. Rine and A. D. Johnson, 1997 Silent information regulator

protein complexes in *Saccharomyces cerevisiae*: a SIR2/SIR4 complex and evidence for

a regulatory domain in SIR4 that inhibits its interaction with SIR3. Proc Natl Acad Sci U

S A 94**:** 2186-2191.

Mortimer, R. K., and J. R. Johnston, 1959 Life span of individual yeast cells. Nature 183**:** 1751-

1752.

Nagaoka, S. I., T. J. Hassold and P. A. Hunt, 2012 Human aneuploidy: mechanisms and new

insights into an age-old problem. Nat Rev Genet 13**:** 493-504.

Neurohr, G. E., R. L. Terry, A. Sandikci, K. Zou, H. Li *et al.*, 2018 Deregulation of the G1/S-

phase transition is the proximal cause of mortality in old yeast mother cells. Genes Dev

32**:** 1075-1084.

North, B. J., M. A. Rosenberg, K. B. Jeganathan, A. V. Hafner, S. Michan *et al.*, 2014 SIRT2

induces the checkpoint kinase BubR1 to increase lifespan. Embo j 33**:** 1438-1453.

O'Sullivan, J. M., D. M. Sontam, R. Grierson and B. Jones, 2009 Repeated elements coordinate

the spatial organization of the yeast genome. Yeast 26**:** 125-138.

Ohkuni, K., and K. Kitagawa, 2011 Endogenous transcription at the centromere facilitates

centromere activity in budding yeast. Curr Biol 21**:** 1695-1703.

Oppikofer, M., S. Kueng, F. Martino, S. Soeroes, S. M. Hancock *et al.*, 2011 A dual role of

H4K16 acetylation in the establishment of yeast silent chromatin. EMBO J 30**:** 2610-

2621.

Pal, S., S. D. Postnikoff, M. Chavez and J. K. Tyler, 2018 Impaired cohesion and homologous

recombination during replicative aging in budding yeast. Sci Adv 4**:** eaaq0236.

Perkins, A. T., T. M. Das, L. C. Panzera and S. E. Bickel, 2016 Oxidative stress in oocytes

during midprophase induces premature loss of cohesion and chromosome segregation

errors. Proc Natl Acad Sci U S A 113**:** E6823-e6830.

Riesen, M., and A. Morgan, 2009 Calorie restriction reduces rDNA recombination independently

of rDNA silencing. Aging Cell 8**:** 624-632.

Rine, J., and I. Herskowitz, 1987 Four genes responsible for a position effect on expression from

HML and *HMR* in *Saccharomyces cerevisiae*. Genetics 116**:** 9-22.

Sasaki, T., B. Maier, A. Bartke and H. Scrable, 2006 Progressive loss of SIRT1 with cell cycle

withdrawal. Aging Cell 5**:** 413-422.

Schalbetter, S. A., A. Goloborodko, G. Fudenberg, J. M. Belton, C. Miles *et al.*, 2017 SMC

complexes differentially compact mitotic chromosomes according to genomic context.

Nat Cell Biol 19**:** 1071-1080.

Schlissel, G., M. K. Krzyzanowski, F. Caudron, Y. Barral and J. Rine, 2017 Aggregation of the

Whi3 protein, not loss of heterochromatin, causes sterility in old yeast cells. Science 355**:**

1184-1187.

Shou, W., J. H. Seol, A. Shevchenko, C. Baskerville, D. Moazed *et al.*, 1999 Exit from mitosis is

triggered by Tem1-dependent release of the protein phosphatase Cdc14 from nucleolar

RENT complex. Cell 97**:** 233-244.

Sinclair, D. A., and L. Guarente, 1997 Extrachromosomal rDNA circles--a cause of aging in

yeast. Cell 91**:** 1033-1042.

Smeal, T., J. Claus, B. Kennedy, F. Cole and L. Guarente, 1996 Loss of transcriptional silencing

causes sterility in old mother cells of *S. cerevisiae*. Cell 84**:** 633-642.

Smith, D. L., Jr., C. Li, M. Matecic, N. Maqani, M. Bryk *et al.*, 2009 Calorie restriction effects

on silencing and recombination at the yeast rDNA. Aging Cell 8**:** 633-642.

Smith, J. S., and J. D. Boeke, 1997 An unusual form of transcriptional silencing in yeast

ribosomal DNA. Genes Dev 11**:** 241-254.

Smith, J. S., C. B. Brachmann, L. Pillus and J. D. Boeke, 1998 Distribution of a limited Sir2

protein pool regulates the strength of yeast rDNA silencing and is modulated by Sir4p.

Genetics 149**:** 1205-1219.

Smith, J. S., E. Caputo and J. D. Boeke, 1999 A genetic screen for ribosomal DNA silencing

defects identifies multiple DNA replication and chromatin-modulating factors. Mol Cell

Biol 19**:** 3184-3197.

Song, S., and F. B. Johnson, 2018 Epigenetic mechanisms impacting aging: a focus on histone

levels and telomeres. Genes (Basel) 9: 201.

Spencer, F., S. L. Gerring, C. Connelly and P. Hieter, 1990 Mitotic chromosome transmission

fidelity mutants in *Saccharomyces cerevisiae*. Genetics 124**:** 237-249.

Steffen, K. K., B. K. Kennedy and M. Kaeberlein, 2009 Measuring replicative life span in the

budding yeast. J Vis Exp. 28: e1209.

Straight, A. F., W. Shou, G. J. Dowd, C. W. Turck, R. J. Deshaies *et al.*, 1999 Net1, a Sir2-

associated nucleolar protein required for rDNA silencing and nucleolar integrity. Cell 97**:**

245-256.

Sunshine, A. B., G. T. Ong, D. P. Nickerson, D. Carr, C. J. Murakami *et al.*, 2016 Aneuploidy

shortens replicative lifespan in *Saccharomyces cerevisiae*. Aging Cell 15**:** 317-324.

Takeuchi, Y., T. Horiuchi and T. Kobayashi, 2003 Transcription-dependent recombination and

the role of fork collision in yeast rDNA. Genes Dev 17**:** 1497-1506.

Uhlmann, F., 2016 SMC complexes: from DNA to chromosomes. Nat Rev Mol Cell Biol 17**:**

399-412.

Unal, E., J. M. Heidinger-Pauli, W. Kim, V. Guacci, I. Onn *et al.*, 2008 A molecular determinant

for the establishment of sister chromatid cohesion. Science 321**:** 566-569.

Utani, K., and M. I. Aladjem, 2018 Extra View: Sirt1 Acts as a gatekeeper of replication

initiation to preserve genomic stability. Nucleus 9**:** 261-267.

Wasko, B. M., and M. Kaeberlein, 2014 Yeast replicative aging: a paradigm for defining

conserved longevity interventions. FEMS Yeast Res 14**:** 148-159.

Winston, F., C. Dollard and S. L. Ricupero-Hovasse, 1995 Construction of a set of convenient

Saccharomyces cerevisiae strains that are isogenic to S288C. Yeast 11**:** 53-55.

Wood, A. J., A. F. Severson and B. J. Meyer, 2010 Condensin and cohesin complexity: the

expanding repertoire of functions. Nat Rev Genet 11**:** 391-404.

Wu, C. S., Y. F. Chen and M. R. Gartenberg, 2011 Targeted sister chromatid cohesion by Sir2.

PLoS Genet 7**:** e1002000.

Yoshida, K., J. Bacal, D. Desmarais, I. Padioleau, O. Tsaponina *et al.*, 2014 The histone

deacetylases Sir2 and Rpd3 act on ribosomal DNA to control the replication program in

budding yeast. Mol Cell 54**:** 691-697.

Zhu, J., D. Heinecke, W. A. Mulla, W. D. Bradford, B. Rubinstein *et al.*, 2015 Single-cell based

quantitative assay of chromosome transmission fidelity. G3 (Bethesda) 5**:** 1043-1056.

Figure Legends

Figure 1. Depletion of Sir2 complexes and elevated chromosome instability in replicatively

aging cells. (A) Western blot of Sir2 protein levels in young and aged cells. Vma2 serves as a

loading control. **(B)** Depiction of the SIR and RENT complexes, which share Sir2 as a catalytic

subunit. **(C, D)** Western blots of native Sir4 and Sir3. **(E, F)** Western blots of 13xMyc tagged

Sir2 paralog, Hst1, and the Net1 subunit of RENT. **(G)** Schematic of artificial chromosome loss

assay for ½ sectored colonies. **(H)** Quantification of sectoring frequency for young or aged cell

populations. Approximately 10,000 colonies were analyzed for each strain across several

biological replicates. *$p < 0.05$, two-tailed student t-test. Bud scar counts are an average for each

enriched population. Qty: indicates the mean western signal of each protein relative to Vma2,

with the ratio for young cells set at 1.0 (n=3 biological replicates).

Figure 2. Chromosome instability is linked to rDNA stability and cohesin redistribution

(A) Schematic of the cohesin complex subunits. **(B)** Western blot of Mcd1-13xMyc and Smc1-

13xMyc from young and aged cells. **(C)** Western blot of 13xMyc tagged Scc2. **(D, E, F, G)**

ChIP-qPCR of Mcd1-13xMyc in young cells (black bars) and aged cells (gray bars) at the

indicated loci normalized to background signal in the *STE2* ORF. **(H)** ChIP-qPCR of Sir4-

13xMyc in young and aged cells normalized to an intergenic *PDC1* site that shows SIR complex

association (LI *et al.* 2013). **(I, J)** ChIP-qPCR of Mcd1-13xMyc from a time course of young, 24

hr, and 36 hr aged cells at indicated loci normalized to *STE2* background signal. Asterisks

indicate significant differences between young and aged cells (p<0.05).

Figure 3. Sister chromatid cohesion is weakened in aged yeast cells. (A) Representative

control images of arm cohesion in an *mcd1-1* mutant at permissive (25°C) and non-permissive

(37°C) temperatures. **(B)** Quantification of cohesion maintenance (1-dot) or loss (2-dots) from

100 *mcd1-1* cells. **(C)** Representative images of young (log-phase) or aged cells arrested with

nocodazole (Nz) and monitored for centromere or arm cohesion. **(D)** Quantifying cohesion loss

(2-dots) in cells with at least 7 bud scars. *p<0.05, two-tailed t-test (n>50 cells). Quantification

of cohesion loss for young and aged cells in the absence of nocodazole **(F)** Representative

images of nocodazole untreated cells. White scale bar represents 2 microns.

Figure 4. Modulation of RLS and CIN by manipulating *MCD1* and *SIR2* expression levels.

(A) Representative images of chromosome loss (sectoring) for WT (YPH278) cells grown

continuously in 2% glucose, galactose, or raffinose. **(B)** Quantification of half-sector colonies

shown in 4A. *p<0.001, two-tailed student t-test. **(C)** RLS assay of YPH278 cells growing on

rich YEP agar plates containing either 2% glucose, galactose, or raffinose. (n=64, each

condition; mean RLS = ♦ 18.9, ■ 9.2, ▲ 15.5) **(D)** Quantification of chromosome loss (1/2

sectors) for strains with an integrated tet^{On} empty (pRF10) or *MCD1* (pRF11) construct. RLS

assays of WT (*MCD1⁺*) and *mcd1L12STOP* strains with integrated pRF10 or pRF11 constructs

(mean RLS = ♦16.2, ■ 10.3, ▲ 19.2, ✕15.6). **(F)** RLS assay of WT and *mcd1L12STOP* strains

with integrated empty, pRS306, or *SIR2* containing plasmid, pJSB186 (n=40 cells each; mean

RLS = ♦21.5 ■ 9.9 ▲ 30.4 ✕14.8). **(G)** RLS assay with *MCD1* OE or *fob1Δ hmrΔ* rescue of

sir2Δ or *sir2Δ hst1Δ* mutants (mean RLS = ♦16.5, ■ 10.1, ▲ 9.7, ✕10.7, ✳ 10.25, ● 16.7). **(H)**

rDNA recombination (marker loss, ½ sector) assay with rDNA marker loss reporter strain

W303AR bearing *ADE2* within the rDNA array. (* p<0.05, two-tailed t-test).

Figure 5. CR suppresses the CIN and RLS defects of *sir2Δ* and *mcd1* mutants. **(A)** RLS

assay of WT (JH5275b) and *mcd1L12STOP* (JH5276b) strains under normal 2% glucose and CR

(0.5% glucose) conditions (n=32 cells; mean RLS = ♦21.3, ■ 8.1, ▲ 24.1, ✕13.8). **(B)** Western

blot of Mcd1-13xMyc for cultures grown in YEP media containing 0.5% glucose. **(C)**

Chromosome loss (sectoring) assay of strains grown in media containing 0.5% glucose. Data

from Figure 1H included for reference. *p<0.05, two-tailed t-test. **(D)** RLS assay of WT and

sir2Δhst1Δ mutant strains under normal and CR conditions. (n=32 cells; mean rls = ♦19.6, ■

9.3, ▲ 23.5, ✕17.1).

Figure 6. Chromosomal instability during replicative aging is linked to the rDNA.

(A) Iteratively corrected and read-normalized heatmap of ChrXII Hi-C contact data at 10 kb

resolution in WT cells growing asynchronously, revealing an interaction between *CEN12* (bin

15) and unique sequence adjacent to the rDNA (bin 45), indicated by yellow arrow. **(B)** Contact

heatmap of chromosomes XI, XII, and XIII showing centromere clustering in asynchronously

growing WT cells. Yellow arrows indicate examples of centromere alignment with *CEN12* and

sequence adjacent to the rDNA. **(C)** Iteratively corrected Chromosome XII heatmap from *cdc15*

cells arrested in anaphase. Yellow arrow indicates the interaction between CEN12 and the

rDNA-adjacent bin. **(D)** Contact map showing interactions between centromeres in the anaphase-

arrested *cdc15* mutant. **(E)** Model for cohesin and Sir2 depletion from the rDNA of aged mother

cells. Cohesin is initially retained at centromeres (Middle Age, black arrow), and then ultimately

lost in the oldest cells. ERCs begin to form during middle age and may contribute in a self-

perpetuating cycle resulting in chronic destabilization of the rDNA array due to loss of cohesin,

RENT, and the cohibin complex (red).

A ChrIV Arm Cohesion (Nz)

mcd1-1 25C   *mcd1-1* 37C

C Young   Aged

Arm (Nz)

Cen. (Nz)

F

Cen.   Young

Aged

Cal. White   LacI-GFP   SPB   Merge

A ChrXII Asynchronous

bin 15 *CEN12*
bin 45 *tQ(UUG)L*

High

Low

B Asynchronous

High

Low

ChrX   ChrXII   ChrXIII

C ChrXII Anaphase *cdc15*

bin 15 *CEN12*
bin 45 *tQ(UUG)L*

High

Low

D Anaphase *cdc15*

High

Low

ChrX   ChrXII   ChrXIII

E

Young   Middle Age   Old

rDNA   rDNA   rDNA

Sir2   Cohesin   Centromeres   ERCs