Competition for a limited pool of cohesin between centromeres and the rDNA

triggers chromosomal instability that shortens replicative lifespan

Ryan D. Fine[1], Nazif Maqani[1], Mingguang Li[1,2], Elizabeth Franck[1], and Jeffrey S. Smith[1,*]

[1]Department of Biochemistry and Molecular Genetics, University of Virginia School of

Medicine, Charlottesville, VA 22908. [2]Department of Laboratory Medicine, Jilin Medical

University, Jilin, 132013, China

[*]Corresponding Author

Department of Biochemistry and Molecular Genetics

University of Virginia School of Medicine

Pinn Hall, Box 800733

Charlottesville, VA 22908

Phone: 434-243-5864

Fax: 434-924-5069

Email: jss5y@virginia.edu

Keywords: Sir2, Hst1, Mcd1, cohesin, Net1, RENT, Lrs4, monopolin, replicative lifespan,

chromosome instability, rDNA, aging

Abstract

Sir2 is a highly conserved NAD$^+$-dependent histone deacetylase that functions in heterochromatin formation and promotes replicative lifespan (RLS) in the budding yeast, *Saccharomyces cerevisiae*. Within the yeast rDNA locus, Sir2 is required for efficient cohesin recruitment and maintaining stability of the tandem array. In addition to the rDNA, ChIP-seq of an epitope-tagged cohesin subunit (Mcd1-13xMyc) in a *sir2Δ* mutant revealed subtle reductions of cohesin binding at all 16 centromeres. Coupled with the previously reported chromosome instability in *sir2Δ* cells and depletion of Sir2 in aged cells, we hypothesized that mitotic chromosome instability (CIN) due to Sir2 depletion could be a driver of replicative aging. In addition to Sir2, we discovered that other subunits of the Sir2-containing SIR and RENT complexes were depleted in aged cells, as were subunits of the cohesin and monopolin/cohibin complexes, implying the possibility of CIN. ChIP assays of the residual Mcd1-13xMyc in aged cells showed strong depletion from the rDNA and possible redistribution to centromeres, most likely in an attempt to maintain chromosome stability. Despite the shift in cohesin distribution, sister chromatid cohesion was partially attenuated in old cells and the frequency of chromosome loss was increased. This age-induced CIN was exacerbated in strains lacking Sir2 and its paralog, Hst1, but suppressed in strains that stabilize the rDNA array due to the deletion of *FOB1* or through caloric restriction (CR). Furthermore, ectopic expression of *MCD1* from a doxycycline-inducible promoter was sufficient to suppress to rDNA instability in aged cells and to extend RLS. Taken together we conclude that age-induced depletion of cohesin and multiple other nucleolar chromatin factors destabilize the rDNA locus, which then results in general CIN and aneuploidy that shortens RLS.

Author summary

The aging process is generally characterized by the breakdown of multiple cellular processes, including the maintenance of genome integrity. Alterations from the normal chromosome number is known as aneuploidy, and commonly occurs in oocytes (eggs) of older mothers. The risk of aneuploidy increases with age and can cause diseases such as Down's syndrome (chromosome 21 trisomy). It is therefore important to understand how aging causes the chromosome instability (CIN) that leads to aneuploidy. In this study we have investigated whether CIN is associated with aging of mitotic cells using the budding yeast, *Saccharomyces cerevisiae*, as a model system. We show that chromosomes are indeed lost during aging of these cells due to the depletion of several key proteins that function in organizing and maintaining chromosome architecture, including the cohesin complex. Restoring cohesin levels was sufficient to delay aging, and adding extra cohesin even extended normal lifespan. Lastly, caloric restriction (CR), a dietary modification known to extend lifespan of yeast and other model organisms, strongly prevented chromosome instability, suggesting a possible new mechanism for how CR delays age-associated conditions at the cellular level.

Introduction

Budding yeast replicative lifespan (RLS) was originally described decades ago as the number of times a mother cell divides before losing viability [1], and has been an effective model system for the identification and/or characterization of several conserved aging-related genes and pathways, including *SIR2*, AMPK (Snf1), and TOR signaling [2]. *SIR2* is probably the most famous yeast gene associated with replicative aging and encodes the founding family member of the NAD^+-dependent histone/protein deacetylases, commonly known as sirtuins (reviewed in [3]). The NAD^+ dependence of sirtuins provides a direct link between metabolism and cellular processes regulated by these enzymes. In fact, recent evidence points to depletion of cellular NAD^+ pools as a potential mechanism for aging-associated disease, which could be mediated by impairment of sirtuins or other NAD^+ consuming enzymes [4]. Therefore, understanding how sirtuins are impacted by aging and determining how they contribute to the regulation of age-altered cellular processes is of intense interest in the aging field.

Eukaryotic genomes generally encode for several different sirtuin homologs. The *Saccharomyces cerevisiae* genome, for example, encodes *SIR2* and four additional Homologs of Sir Two (*HST1-HST4*) [5]. Sir2 was originally shown to establish and maintain silencing of the silent mating loci, *HML* and *HMR*, along with its fellow Silent Information Regulator (SIR) proteins, Sir1, Sir3, and Sir4 [6]. Additionally, Sir2, Sir3, and Sir4 form the so-called SIR complex that is recruited to and then spreads across the *HM* loci and telomeres to form hypoacetylated heterochromatin-like domains (reviewed in [7]). Sir2 protein levels are significantly reduced in replicatively old yeast cells [8], presenting a possible mechanism for the decline of Sir2-dependent processes during aging, including gene silencing. Indeed, depletion of Sir2 in old cells causes hyperacetylated H4K16 and silencing defects at subtelomeric loci [8].

Haploid cells lacking Sir2 derepress *HML* and *HMR*, resulting in co-expression of mating type

regulatory genes that together induce a diploid-like gene expression pattern that produces a non-

mating "pseudodiploid" phenotype [6]. Old haploid mother cells are sterile, presumably due to

the pseudodiploid effect [9], but more recent experiments point toward a silencing-independent

mechanism caused by aggregation of the Whi3 protein and loss of pheromone sensitivity [10].

Alternative models for Sir2 control of RLS have focused on the rDNA tandem array. Sir2

functions in the silencing of RNA polymerase II-dependent transcription at the rDNA locus on

chromosome XII via a nucleolar complex called RENT [11, 12], which consists of Sir2, Net1,

and Cdc14 [13, 14]. RENT-dependent H3 and H4 deacetylation represses transcription of

endogenous non-coding RNAs from the intergenic spacer (IGS) regions [15]. Derepression of

the bidirectional promoter (E-pro) within IGS1 of *sir2Δ* cells displaces cohesin from the rDNA,

thus destabilizing the array by making it more susceptible to unequal sister chromatid exchange

[16]. A predominant model for *SIR2* control of replicative aging postulates that

extrachromosomal rDNA circles (ERCs) derived from these unequal recombination events

specifically accumulate to high levels in mother cells, consequently causing cell death through an

unknown mechanism [17]. Such an ERC-centric model is supported by the extended RLS

extension of *fob1Δ* strains [18]. Fob1 binds to the rDNA in IGS1 to block DNA replication forks

from colliding with RNA polymerase I transcribing in the opposite direction [19]. The blocked

replication forks can collapse and result in double-stranded DNA breaks that trigger the unequal

sister chromatid exchange [20]. The frequency of rDNA recombination and ERC accumulation is

significantly reduced in a *fob1Δ* mutant due to loss of the fork block, thus extending RLS [18].

This rDNA-centric model of replicative aging has been modified over the years such that overall

rDNA instability, rather than ERC accumulation, is now considered the critical deleterious factor

toward lifespan [21]. Strains with virtually no ERC accumulation in old cells, but a high

frequency of rDNA recombination, are short-lived [22]. Also consistent with this idea, *sir2Δ* still

reduces RLS in a *fob1Δ* strain that has limited ERCs [23]. This raises several important

questions, including 1) does Sir2 regulate RLS through mechanisms independent of the rDNA,

and 2) how does rDNA instability shorten RLS? The experiments in this current study address

both of these questions.

As mentioned above, Sir2 is required for efficient cohesin association with the rDNA,

with stabilization of the tandem array a major mechanism for maintaining replicative longevity.

The cohesin complex in *S. cerevisiae* consists of four subunits, Mcd1/Scc1, Irr1/Scc3, Smc1, and

Smc3 [24]. The cohesin loading complex (Scc2/Scc4) deposits cohesin onto centromeres and

other sites across the chromosome arms (including rDNA) during S-phase to maintain sister

chromatid cohesion (SCC) until anaphase, when the Mcd1 subunit is cleaved by separase to

facilitate sister chromatid separation (reviewed in [24]). Cohesin defects result in chromosome

instability (CIN) and aneuploidy due to improper chromosome segregation during mitosis

(reviewed in [25]. Interestingly, cells lacking *SIR2* or heavily overexpressing *SIR2* have

increased rates of chromosome loss [26, 27]. Aneuploidy has been shown to shorten yeast RLS

by inducing proteotoxicity through improper expression levels [28]. Taken together, these results

suggest that Sir2 could impact RLS by regulating mitotic chromosome segregation during aging.

In this study, we investigated a potentially novel role for Sir2-limited yeast RLS through

maintenance of accurate chromosome segregation. In the process, we uncovered chromosome

instability and SCC defects in aging yeast cells that correlated with reduced levels of cohesin and

cohesin loader subunits. Forcibly manipulating Mcd1 subunit levels was sufficient to modify

RLS, suggesting a causative relationship to aging. Interestingly, cohesin association with

chromosome arms and the rDNA was generally reduced in old cells, but maintained or even

increased at centromeres, suggesting a programmed mechanism by which SCC is preferentially

maintained at centromeres to ensure cell viability. However, this comes at the expense of chronic

rDNA instability that is exacerbated by age-induced reductions in the RENT and

cohibin/monopolin complexes. Lastly, reducing rDNA instability by deleting *FOB1* suppressed

the general CIN phenotype in aged cells, leading to a model whereby rDNA instability caused by

cohesin depletion drives the defects in mitotic segregation of other chromosomes during

replicative aging.

Results

We previously reported a statistically significant overlap in genome-wide Sir2 binding sites

measured by ChIP-seq with cohesin binding sites measured by ChIP-Chip microarray analysis

[29, 30], consistent with the idea of a functional relationship between the two. Since deleting

SIR2 reduced cohesin recruitment at individually selected loci [30], including the rDNA [31], we

decided to explore this result further using ChIP-Seq with exponentially growing WT and *sir2Δ*

strains in which the Mcd1 subunit of cohesin was C-terminally tagged with 13 copies of the myc

epitope (13xMyc). A composite plot of the sequencing data normalized to 1X read coverage

indicated both replicates of the *sir2Δ* strain exhibited dramatically reduced levels of Mcd1-myc

binding within the rDNA locus as compared to the WT strain (Fig 1A). This result was

confirmed with ChIP-qPCR using an IGS1 primer pair (Fig 1B) and supported earlier ChIP

results from the Nomura lab showing that cohesin recruitment to the rDNA requires Sir2 [31]. It

was also reported that cohesin released from the rDNA in a *sir2Δ* mutant preferentially

relocalized to pericentromeric regions as measured by fluorescence microscopy of GFP-tagged

Smc3 [32]. We attempted to recapitulate this result by creating a composite profile of all 16

centromeres but did not detect the same pericentromeric enrichment of Mcd1-myc in our *sir2Δ*

ChIP-seq datasets (Fig 1C). There was instead a trend toward reduced cohesion association.

Standard ChIP assays also did not reveal evidence of significant centromeric or pericentromeric

enrichment (Fig 1D). However, we cannot rule out the possibility that the ChIP signal was

diluted within an asynchronous cell population, compared to the microscopy data that examined

mitotic cells.

Cohesin levels are depleted in old yeast cells

Strains deleted for *SIR2* are short lived, but they likely do not fully recapitulate all

cellular changes that accompany normal replicative aging. We therefore hypothesized that more

severe effects of cohesin localization at centromeres could be uncovered by comparing

replicatively young vs. old cells in which Sir2 protein levels naturally declined. To isolate

sufficient quantities of old cells for ChIP assays we turned to Mother Enrichment Program

(MEP) strains developed by the Gottschling lab [33]. The aged cell purification procedure was

validated by increased bud scar staining with calcofluor white and by the expected reduction of

Sir2 protein (Fig 2A). The vacuolar protein Vma2, used as a loading control, does not deplete

with age [34], and detailed bud scar and protein quantitation from triplicate biological samples

are reported in S1 Fig. Since Sir2 is the catalytic subunit of both SIR and RENT (Fig 2B), it was

important to know which complexes were impacted by age. As shown in Figs 2C and 2D, Sir4

was depleted in old cells while Sir3 was not. Such a stark difference was considered highly

relevant because Sir2 and Sir4 tightly interact as a heterodimer that associates with the acetylated

H4 N-terminal tail [35]. Sir3 is subsequently recruited following H4K16 deacetylation to

complete SIR holocomplex formation on heterochromatin [36]. Myc-tagged Net1 (RENT

complex) was also depleted from old cells (Fig 2E), indicating that Sir2/Sir4 and the RENT

complex are both depleted during replicative aging, thus revealing a previously unrecognized

level of specificity in age-related Sir2 protein homeostasis. It should be noted that the Sir2

paralog, Hst1, which has the capacity to compensate for loss of Sir2, is also partially depleted

from old cells (S2A Fig).

Considering the depletion of multiple heterochromatin factors in old yeast cells and the

generally reduced Mcd1-myc ChIP signal in *sir2Δ* cells, we hypothesized that cohesin complex

levels may also drop in old cells. We attempted to individually myc-tag three different subunits

of cohesin (Mcd1, Smc1, and Irr1) in the MEP strain background. All cohesin subunits are

essential, and only two of the tagged proteins (Mcd1 and Smc1) were functional based on

recovery of viable cells and proper size on western blots. Both of these subunits were

significantly depleted in old cells (Fig 2F and S2B Fig), indicating the entire cohesion complex

was affected. Furthermore, a Myc-tagged Scc2 subunit of the Scc2/Scc4 cohesin loading

complex was age-depleted (S2C Fig), predicting that cohesion complex association with

chromatin should be generally sparse. ChIP assays for Mcd1-myc in aged cells did demonstrate

strong depletion from the rDNA intergenic spacer IGS1 as expected, but binding was

surprisingly enhanced at the centromere of chromosome IV (Fig 2G). Sir4-myc, on the other

hand, was depleted from *TELXV* in old cells (one of its normal targets) without shifting its

enrichment to centromeres (*CEN4*) or the rDNA (Fig 2H), indicating that not all age-depleted

proteins become enriched at centromeres. From these results we conclude that as cohesin is

depleted during replicative aging, a significant portion of the remaining complex is retained and

potentially redistributed to centromeres, consistent with an earlier finding that cohesin

195 preferentially associates with pericentromeric regions instead of chromosome arms when Mcd1

196 expression is artificially reduced below 30% of normal [37].

198 **Sister chromatid cohesion is compromised in old yeast cells**

199   SCC was surprisingly normal despite artificially reduced Mcd1 levels [37], leading us to

200 ask whether cohesion would be maintained in old yeast cells that were also depleted of cohesin.

201 To this end, we utilized strains with LacO arrays located approximately 10 kb away from

202 centromere IV (*CEN4*) or on the arm of chromosome IV at the *LYS4* locus located approximately

203 400 kb away from the centromere as proxy for arm cohesion [38, 39]. Differential positioning of

204 the array had no significant impact on RLS (S3 Fig). SCC was monitored by LacI-GFP

205 appearing either as one dot in the case of maintained cohesion or two dots in the case of cohesion

206 loss. Using an *mcd1-1* temperature sensitive mutant as a positive control [40], we observed a

207 significant increase in two dots when cells were synchronized in mitosis with nocodazole and

208 shifted to 37°C (Fig 3A and B). WT cells were next biotinylated and aged for 24 hours, followed

209 by purification with magnetic streptavidin beads. The maximum bud scar count was 13, which

210 was roughly half the maximum lifespan for this strain as measured by RLS assay. There was a

211 trend toward lost centromere and arm cohesion in old cells (>5 bud scars) when cultures were

212 synchronized in mitosis with nocodazole (Fig 3C, D, and E). When cells were not arrested with

213 nocodazole and only large budded (mitotic) cells were analyzed, significant cohesion loss was

214 observed in old mother cells compared to young (Fig 3E and F). Therefore, the limited amount of

215 cohesin complex remaining in old cells is functional, but has difficulty maintaining SCC when

216 microtubules are naturally allowed to exert pulling force on centromeres.

Chromosome instability (CIN) increases during replicative aging

Based on the cohesion defect in aged yeast cells, we next asked if they also display

elevated chromosome instability. Strains for this experiment have an artificial chromosome III

bearing a suppressor tRNA gene, *SUP11* [41]. Loss of the chromosome prevents suppression of

an ochre stop codon in *ADE2*, resulting in the classic *ade2* red colony phenotype. The frequency

of nondisjunction events was measured by counting half-sectored red/white colonies from young

and aged cell populations (Fig 4A). A *sir2Δ* mutant was included in the experiment because it

was previously reported to increase CIN [26], as was an *hst1Δ* mutant because of its potential for

functional overlap with Sir2 [42, 43]. Sectoring was elevated in young populations of *sir2Δ* and

hst1Δ mutants, and additively elevated in a *sir2Δ hst1Δ* double mutant (Fig 4B, left panel).

Interestingly, sectoring was even higher for aged populations of each strain, suggesting an

independent age-associated factor was involved. We next tested whether the *sir2Δ* effect on

sectoring was related to the pseudodiploid phenotype caused by derepression of the *HM* loci.

This reporter strain background was *MATα*, so we deleted *HMR* (chrIII 293170-294330) to

eliminate the a1/a2 transcription factors, and then confirmed reversal of the pseudodiploid

phenotype by restoration of mating to the *sir2Δ hmrΔ* strains (data not shown). Importantly, this

procedure significantly suppressed sectoring of the young *sir2Δ* and *sir2Δ hst1Δ* mutants, but not

the *hst1Δ* mutant (Fig 4B, middle panel), indicating there was indeed a *sir2Δ*-induced

pseudodiploid effect that suppressed CIN (Fig 4B, middle panel). Aging increased sectoring in

each strain even when *HMR* was deleted, indicating the aging-associated CIN factor was

unrelated to mating type control.

Even though SCC was unaffected by forced cohesin depletion, rDNA array condensation

and stability were significantly impaired [37]. We therefore hypothesized that age-induced

chromosome instability could be related to rDNA instability caused by natural Sir2 and cohesin

depletion. To address this idea, *FOB1* was deleted to stabilize the rDNA and the sectoring

phenotype retested. As shown in Fig 4B (right panel), the age-associated increase in

chromosome instability for each mutant was generally reduced to that observed with young

FOB1[+] cells (Fig 4B, left panel), consistent with rDNA instability driving overall CIN.

What could be the mechanistic connection between the rDNA and centromeres? Previous

Hi-C analysis of the yeast genome and fluorescence microscopy of nucleolar proteins positioned

the rDNA off to one side of the nucleus, apparently secluded from the rest of the genome [44,

45]. The repetitive nature of rDNA precludes it from appearing in Hi-C contact maps, but closer

inspection of chromosome XII at 10 kb resolution indicated a clear interaction between unique

sequences flanking the centromere-proximal (left) edge of the array and *CEN12* (Fig 4C).

Further analysis of observed/expected contacts using HOMER revealed a looped structure

between the rDNA and *CEN12* (Fig 4D), which was recently reported to occur during anaphase

[46]. Interestingly, all centromeres in the yeast genome, including *CEN12*, cluster together in Hi-

C contact maps (Fig 4E; [44]), which by default also places them in contact with the left flank of

the rDNA. Taken together, these results suggest that the rDNA comes in close contact with

centromeres during mitosis, providing a potential window of time for rDNA instability to

physically impact the integrity of general chromosome segregation.

Growth on galactose induces CIN and shortens RLS

Since Sir2 and cohesin are both naturally depleted from replicatively aging yeast cells,

and mild Sir2 overexpression extends RLS [23], we hypothesized that manipulating cohesin

expression levels would also impact CIN and RLS. We initially attempted to overexpress the

Mcd1 subunit from a galactose inducible *GAL1* promoter and then measure minichromosome

loss frequency by counting ½ sectored colonies. However, simply growing the reporter strain in

galactose-containing media, even with an empty expression cassette, resulted in severe

minichromosome loss compared to glucose-containing media (Figs 5A and B). This effect was

specific to galactose, as growth with another non-preferred carbon source (raffinose) had no

effect on sectoring (Figs 5A and 5B). We next measured RLS with the minichromosome reporter

strain on YEP plates with 2% glucose, galactose, or raffinose. As shown in Fig 5C, galactose

decreased the mean RLS by ~50% compared to glucose (9.2 vs. 18.9 divisions), while raffinose

only marginally decreased lifespan to an average of 15.5 cell divisions. To confirm the galactose

effect on RLS was not specific to the minichromosome strain, RLS assays were repeated with the

well-characterized strains BY4741 (*MAT*a) and BY4742 (*MAT*α). Again, a significant decrease

in mean lifespan was observed for BY4741 (17.7 divisions) and BY4742 (18.9 divisions) on

galactose as compared to glucose (24.3 and 24.2 divisions, respectively) (Fig 5D), suggesting

that galactose triggers high rates of CIN through an unknown mechanism that also shortens RLS.

RLS is modulated by Mcd1 expression levels

To circumvent the use of galactose for *MCD1* overexpression we turned to an inducible

"Tet-On" promoter that is activated by doxycycline [47]. Strains harboring this integrated

cassette transcriptionally overexpressed *MCD1* approximately 7-fold compared to the empty

vector control (S4 Fig). *MCD1* was overexpressed in a "WT" strain (JH5275b) and a mutant

(JH5276b) containing an ochre stop codon in the *MCD1* open-reading frame (*mcd1L12STOP*)

that reduced Mcd1 protein levels to ~30% of normal [37]. With an empty pRS405 control vector

integrated, *mcd1L12STOP* exhibited a 40% reduction in mean RLS (9.9 divisions) compared to

WT (16.5 divisions), and overexpressing *MCD1* almost fully restored mean RLS to the mutant

(14.7 divisions) (Fig 6A). Especially intriguing was the observation that *MCD1* overexpression

extended mean RLS in the WT background to 19.6 division, an increase that was primarily due

to improved survival during the first ~15 divisions, followed by a steeper decline later (Fig 6A).

This biphasic pattern was highly reproducible, as it was observed in three independent RLS

experiments. Mechanistically, there could be a secondary age-related event at the RLS mid-point

that is beyond rescue by ectopic *MCD1* expression. Alternatively, doxycycline-induction could

eventually be attenuated due to the extended incubation times of RLS assays. Either way, the

results clearly indicate that similar to Sir2, the Mcd1 subunit of cohesin is a dosage-dependent

longevity factor.

Previous studies reported that Sir2 was required for efficient cohesin association with the

rDNA due to silencing of E-pro [16, 31], implying that Sir2 was upstream of cohesin. This

functional relationship could be more complex in the context of RLS, especially since both

factors are depleted with age. To explore further, we tested whether *SIR2* overexpression could

rescue the short RLS of an Mcd1-depleted *mcd1L12STOP* strain by integrating a second copy of

SIR2 (2x*SIR2*) at the *LEU2* locus. As shown in Fig 6B, 2x*SIR2* partially rescued mean RLS of

the *mcd1L12STOP* strain (14.1 versus 9.9 divisions), and also increased maximum RLS of the

WT strain as expected. Reciprocally, we asked whether *MCD1* overexpression could suppress

the short RLS of a *sir2Δ hst1Δ* mutant. The double mutant was chosen to avoid any redundancy

between the two sirtuins. Mean RLS was clearly not increased by *MCD1* overexpression as

compared to empty vector (10.1 versus 9.7 divisions, Fig 6C), confirming that *SIR2* was

upstream of *MCD1* in regulating RLS.

Considering the strong depletion of cohesin from rDNA in old cells (Fig 2G), and

extended RLS when *MCD1* was overexpressed (Fig 6A), we next tested whether aging-induced

rDNA instability was suppressed by *MCD1* overexpression using a reporter strain harboring

ADE2 in the rDNA array [23]. There was a large increase of red/white sectoring (marker loss) in

old cells that was suppressed upon *MCD1* overexpression (Fig 6D). In the absence of *SIR2*,

however, red/white sectoring was high in both young and old cells when the empty vector

(pRF10) was integrated, and *MCD1* overexpression did not significantly reduce rDNA instability

in either population (Fig 6D), indicating that at least some Sir2 was required for Mcd1 to impact

rDNA stability. We conclude that loss of Sir2 and cohesin in aging cells causes rDNA array

instability that exacerbates CIN.

RLS extension by CR correlates with improved chromosome stability

Reducing glucose concentration in the growth medium is effective at extending RLS and

is considered a form of caloric restriction (CR) for yeast [48, 49]. There have been several

hypotheses put forth for the underlying mechanisms, including stabilization of the rDNA [21].

Since *MCD1* overexpression suppressed rDNA recombination and extended RLS, we

hypothesized that CR would suppress the shortened RLS of a cohesion-depleted *mcd1L12STOP*

mutant strain. Indeed, CR extended RLS of both the WT and *mcd1L12STOP* strains (Fig 7A).

However, the suppression was apparently not due to maintenance of global cohesion levels

because steady state Mcd1-13xMyc was still depleted in restricted old cells (Fig 7B). CR also

strongly suppressed minichromosome loss in young and aged cells, even in the *sir2Δ hst1Δ*

double mutant (Fig 7C). Importantly, this CR effect also correlated with almost complete rescue

of RLS for the *sir2Δ hst1Δ* mutant (Fig 7D). Taken together, the results reveal a new mechanism

for RLS extension by CR, whereby stabilization of the rDNA locus helps maintain general

mitotic chromosome stability to protect against aneuploidy.

Discussion

Nuclear protein depletion during replicative aging as a paradigm for aging pathologies

During this study the majority of chromatin-associated proteins analyzed by western

blotting were depleted in replicatively aged yeast cells. The only protein unaffected by age, other

than the vacuolar Vma2 control, was Sir3. A similar proportion of homologous recombination

proteins were depleted in an independent analysis of old cells, with Rad52 the only one not

affected [50]. These results indicate there is at least some selectivity to the depletion of nuclear

proteins in old cells. However, the large number of depleted factors also makes it likely that

targeted nuclear protein deficiency could lead to multiple age-associated phenotypes.

Replicatively aging yeast cells appear especially susceptible to this phenomenon, as even total

core histone levels are depleted [51]. Evidence also exists for histone depletion during aging of

metazoan organisms, including mammals (reviewed in [52]). More generally, global protein

turnover is elevated in cells from prematurely aging progeria patients, which may trigger higher

translation rates [53]. This is significant because reducing translation is a means of extending

lifespan in multiple organisms [54]. Ribosomal proteins appear to accumulate in old cells [55],

again consistent with protein synthesis being a driver of aging [56]. The mechanism(s) driving

nuclear protein depletion in old yeast mother cells remain unclear.

The specificity for Sir2/Sir4 depletion over Sir3 is intriguing given that Sir2 and Sir4

form a tight complex that allosterically stimulates the deacetylase activity of Sir2 [57]. Sir2/Sir4

also binds to H4K16-acetylated chromatin independent of Sir3, which is subsequently recruited

to complete the SIR holocomplex once H4K16 is deacetylated [36]. The mechanism for Sir2/Sir4

depletion from old cells remains uncharacterized, though in non-aging cell populations the

stability/turnover of Sir4, but not Sir2, is mediated by the E3 ubiquitin ligase San1 [58], which

has also been implicated as a quality control E3 ligase for mutated/unfolded nuclear proteins

[59]. Whether San1 controls Sir4 stability during aging remains unknown, but since Sir4 is more

severely depleted than Sir2 in old cells (Figs 2A and 2C), Sir4 could be selectively depleted from

the SIR complex, thus leaving Sir2 unprotected and then subject to turnover through a different

mechanism. Alternatively, Sir2/Sir4 could be equally depleted as a complex from telomeres and

the *HM* loci (not necessarily via San1), leaving the nucleolar pool of Sir2/RENT as more

resistant to aging. Under this scenario, protecting the integrity of the rDNA array could take

precedence over other heterochromatic domains. Interestingly, the *Schizosaccharomyces pombe*

San1 ortholog has also been implicated in a chaperone-assisted degradation pathway that

functions in quality control of kinetochores to promote chromosome stability [60].

Sir2 depletion in replicatively aged yeast cells is reminiscent of Sirt1 depletion in serially

passaged mouse embryonic fibroblasts (MEFs), which correlates with declining mitotic activity

[61]. Sir2 and Sirt1 are both known to function in regulating DNA replication origins [62, 63],

and the effect of deleting *SIR2* on early origin firing is thought to be mediated by competition for

limiting factors with the repeated rDNA origins [64]. Furthermore, CR has been proposed to

extend RLS by reducing rDNA origin firing, which improves overall genome replication [65].

Therefore, depleted cohesin in old cells could potentially cause rDNA instability by impacting

DNA replication.

A precarious balance between rDNA and centromeric cohesion

Sister chromatid cohesion (SCC) ensures chromosomes are not segregated until the

Mcd1/Scc1 cohesin subunit is cleaved in response to a mitotic spindle checkpoint signal that all

chromosomes are properly attached to microtubules and aligned at the metaphase plate (reviewed

in [24]). Cohesin is also critical during meiosis, and it is well established in mammals that SCC

defects occur in the oocytes of older mothers, causing meiotic chromosome missegregation

events during both anaphase I and II [66]. This phenomenon is believed to be a major

mechanism for increased aneuploidy risk that usually results in embryonic lethality, or in the

case of chromosome 21 trisomy, Down's syndrome in children. The meiotic cohesin subunit

Rec8 is depleted in the oocytes of older mice, as is Shugoshin (Sgo2), which normally

protects/maintains centromeric cohesin [67]. More recent experiments in *Drosophila* suggest that

oxidative stress in aged oocytes contributes to the SCC defects [68]. Our results in replicatively

aging yeast cells reveal that aging-induced cohesin depletion and the resulting chromosome

missegregation can extend to mitotic cells. Though cohesin depletion or defects have not been

reported for mammalian somatic cells, the mitotic spindle checkpoint protein BubR1 is depleted

in dynamic somatic tissues such as spleen [69]. Deficiency of this protein results in premature

aging phenotypes [69], while overexpression extends lifespan [70]. This is similar to the effects

we observe with Mcd1 depletion and overexpression on yeast RLS. Interestingly, BubR1 is also

a deacetylation target of Sirt2, which appears to stabilize the protein and extend lifespan, thus

linking mitotic spindle checkpoint regulation to NAD^+ metabolism [71]. It remains unclear if

Sir2, Hst1, or other sirtuins regulate the yeast BubR1 ortholog, Mad3, or additional checkpoint

and kinetochore proteins.

SCC is the canonical function for cohesin, though the complex also functions in

establishing and regulating genome organization at the level of chromatin structure, gene

regulation, and double strand break (DSB) repair (reviewed in [72]). Among these various

processes, SCC at centromeres appears the most critical because artificial depletion of Mcd1 to

<30% of normal levels results in preferential cohesin binding to pericentromeric regions rather

than cohesin associated regions (CARs) on chromosome arms [37]. SCC was also well

maintained in these strains at the expense of normal chromosome condensation, DNA repair, and

rDNA stability [37]. In aged yeast cells we observed relative enrichment of Mcd1-myc at

centromeres as compared to loss at the rDNA (IGS1) locus (Fig 2G), suggesting an attempt by

cells to maintain centromere cohesion at all costs, similar to what occurred in the artificially

depleted system. Despite maintaining the cohesin complex at centromeres, SCC was still

impaired in the older cells, especially when tension was maintained. In contrast, an independent

study that analyzed significantly older mother cells (~20 generations) than in our study (~6

generations) observed reduced cohesin enrichment at centromeres [50]. Collectively, the results

suggest that centromere-associated cohesin is preferentially retained during the initial stages of

replicative aging, but then eventually breaks down below a critical threshold in the oldest cells.

Numerous nuclear proteins are depleted in old yeast cells, not just cohesin subunits, so

we hypothesize that defects in other nuclear processes mediated by such factors also contribute

to SCC defects and chromosome instability either directly or indirectly. The depleted cohesin

loading complex (Scc2/4) is an obvious candidate due to its role in loading cohesin onto

chromatin. Similarly, the depleted Lrs4/Csm1(cohibin complex) is proposed to act as a cohesin

clamp onto rDNA chromatin [73], and also functions at centromeres to maintain mitotic integrity

[74]. Sir2 and Hst1 are also obvious candidates given the earlier finding that H4K16

deacetylation at centromeres by Sir2 helps maintain chromosome stability [26]. Hst1 also binds

centromeric DNA *in vitro* and *in vivo* [75], though the functional relevance of that association

remains uncharacterized. The suppression of age-associated mini-chromosome loss in the

absence of *FOB1* clearly points to rDNA instability as an unexpected source of general CIN.

Such a relationship is reinforced by the observed depletion of nucleolar proteins Net1 and Lrs4

in old cells (Figs 2E and S2D), both of which are required for normal rDNA/nucleolar integrity

and stable cohesin association with the rDNA [14, 73, 76].

How could destabilization of the rDNA locus result in general chromosome instability

and shortened RLS? As depicted in Fig 4, unique sequence flanking the rDNA on chromosome

XII contacts the centromere of each chromosome, including XII. Whether the actual rDNA genes

contact centromeres remains unclear due to the limitations of Hi-C analysis with repetitive DNA.

However, specific regions of the rDNA were previously shown to associate with various non-

rDNA chromosomal regions using an anchored 4C approach [77]. Loss of cohesin from the

rDNA could potentially disrupt long-range interactions with centromeres or non-centromeric

regions of cohesin association that influence chromosome integrity. One potential mechanism

could be significant disruption of overall chromosome condensation during mitosis, as the

cohesin appears to play the major role in chromosome condensation in budding yeast instead of

the condensin complex [46], which is different from metazoan organisms where condensin is the

primary condensation machinery.

Interestingly, another class of nuclear factors depleted in old yeast cells are several DNA

repair proteins [50]. Consequently, the lack of proper DNA repair while the rDNA becomes

destabilized correlates with fragmentation of chromosome XII and the other chromosomes, with

rDNA sequences actually being transferred into the other chromosomes. It was proposed that

accumulation of breaks and rearrangements ultimately causes cell death during replicative aging.

Such cells were significantly older (>25 divisions) compared to the cells in our study, which

averaged 5-6 divisions. Alternatively, it is possible that these presumably random

rearrangements disrupt normal SCC, leading to CIN.

Aneuploidy as an aging mechanism

All 16 *S. cerevisiae* chromosomes harbor essential genes, so if a single chromosome is

lost from a haploid yeast cell, then the affected mother or daughter cell should become inviable

and no longer divide. Given the elevated frequency of chromosome loss during replicative aging,

the chances of generating an inviable mother cell during a replicative aging assay increases after

each subsequent division. Therefore, at least a portion of the replicative lifespan in haploid yeast

cells is controlled by the ability to maintain all 16 chromosomes. Complete loss of a

chromosome would not be an immediate viability issue for diploid cells, however, as the chances

of losing both homologs in a single mitosis are exceedingly rare. On the other hand, haploid

strains that are disomic for individual chromosomes are often short lived, with longer

chromosomes typically having larger effects [28]. It was hypothesized that such strains suffer

from proteotoxic stress due to inappropriate protein expression levels. Therefore, a similar

mechanism could shorten RLS in a diploid strain that is trisomic for an individual chromosome,

though this has not yet been tested. Aneuploidy is also a hallmark of aging in the germline [78],

and somatic tissues of mammals [70, 79], making it a conserved feature of aging from yeast to

humans.

Another exciting feature of this study is the suppression of CIN by CR growth conditions

that extend RLS. This effect was independent of the reduced cohesin levels in aged cells, and

even improved RLS of the cohesin-depleted strains. Since SCC is normal in the cohesin-depleted

strain [37], we hypothesize that CR reinforces other processes that are defective due to reduced

cohesin or other depleted factors that promote rDNA stability. Indeed, CR is known to suppress

rDNA instability in yeast cells [80, 81], and improve overall genome replication efficiency [65].

Hi-C analysis also suggests there could be direct effects of rDNA structure on centromere

function, which will be the focus of future investigation.

Methods

Yeast strains, plasmids, and media

Yeast strains were primarily grown at 30°C in Yeast Peptone Dextrose (YPD) or

Synthetic Complete (SC) medium for strains bearing plasmids [82]. *SIR2, HST1*, or *FOB1* open

reading frames were disrupted with one-step PCR-mediated gene replacement using *kanMX4*,

natMX4, or *hphMX4* drug resistance markers, respectively. The *HMR* deletion by replacement

with *hphMX4* spans sacCer3 genome coordinates of chrIII 293170-294330. All C-terminally

13xMyc (EQKLISEEDL) tagged proteins were fused at the endogenous loci of haploid MEP

strains UCC5181 and UCC5179, followed by crossing to generate homozygous diploids. All

deletions and fusions were confirmed by colony PCR, western blotting, or both. pRF4 was

constructed by PCR amplifying the *MCD1* open reading frame from ML1 genomic DNA and

ligating into *Pst*I and *Not*I sites of pCM252, a tetracycline/doxycycline inducible overexpression

vector available from Euroscarf (http://www.euroscarf.de). pRF10 and pRF11 were constructed

by removing the expression cassette by *Pvu*II blunt end digestion of pCM252 and pRF4,

respectively, and ligating it between the *Pvu*II sites of pRS405 bearing the *LEU2*, thus replacing

the *TRP1* marker with *LEU2*. pSB760 and pSB766 are integrating and *2µ LEU2* vectors,

respectively, bearing a single copy of *SIR2* [83]. All strains used in this work are listed in S1

Table and all primers are listed in S2 Table.

Old yeast cell isolation

Old yeast cell enrichment was based on MEP (Mother Enrichment Program) strains [33,

34]. For all assays, 1 µL of stationary phase culture was inoculated into 100 mL of YPD medium

and then grown into log phase. Approximately $1x10^8$ cells were harvested, and centrifuged cell

pellets washed 3 times with 1x phosphate buffered saline (PBS). Cells were then resuspended in

1 mL of PBS and mixed with 5 mg of Sulfo-NHS-LC-Biotin (Pierce) per $1x10^8$ cells for 30

minutes at room temperature. After biotin labeling, $5x10^7$ cells were added to 1.5 L YPD cultures

containing 1µM estradiol, and 100 µg/mL ampicillin to prevent bacterial contamination. These

cultures were allowed to grow for 24 hours before being processed in a downstream assay

specific manner (see below). For non-MEP strain backgrounds, estradiol was not added to the

cultures.

Old cell western blots

Two large scale cultures (1.5 L) were used for each western blot experiment

corresponding to approximately $2x10^7$ total old cells after purification. Cells were pelleted using

a Sorvall RC-5B Plus centrifuge with an SLA-3000 rotor at 2000 rpm, and then resuspended at a

density of $6x10^8$ cells/mL in RNAlater (Ambion) for 45 minutes in two separate conical tubes.

Following fixation, cells were pelleted and resuspended in 45 mL of cold 1xPBS, 2 mM EDTA

in 50 mL conical tubes. The mixture was incubated at 4°C for 30 min with 800 µL of

Streptavidin MicroBeads (Miltenyi Biotec), which were then purified through an autoMACS Pro

Cell Separator using the posseld2 program (UVA flow cytometry core facility). A 20 µL aliquot

of each output was used for bud scar counting using calcofluor white staining, before combining

the isolated samples into a single microfuge tube. Samples were frozen at -80°C before protein

extraction. Thawed cells were vortexed twice for 1 min in 20% TCA (trichloroacetic acid) with

~100 µL of acid washed glass beads with a brief cooling period in between vortexing. Beads

were allowed to settle before removing the supernatant to a fresh microfuge tube. A 250 µL wash

of 5% TCA was applied twice to the beads and pooled with the initial lysis sample. Proteins were

precipitated at 10,000 rpm in a microfuge for 5 min at 4°C. The pellets were resuspended in 50

µL of 1x SDS sample buffer (50 mM Tris-HCl pH 6.8, 2% SDS, 10% Glycerol, 3.6 M 2-

mercaptoethanol) and neutralized with 30 µL of 1M Tris-HCl, pH 8.0. Samples were run on a

9% (w/v) SDS-polyacrylamide gel and transferred to an Immobilon-P membrane (Millipore).

Membranes were incubated for 1 hour at room temperature in 1xTBST + 5% non-fat milk with

primary antibodies (1:2000 α-Myc 9E10, 1:5000; α-Vma2 (Life Technologies); 1:5000 α-Sir2

(Santa Cruz Biotechnology); 1:1000 α-Sir4 (Santa Cruz Biotechnology); 1:1000 α-Sir3 (Santa

Cruz Biotechnology). HRP-conjugated secondary antibodies (Promega) were diluted 1:5000, and

detected using chemiluminescence with HyGLO (Denville Scientific).

ChIP Assays with old and young cells populations

Two 1.5 L cultures were used for each biological replicate sample. After centrifugation,

cells were washed with 1xPBS and resuspended in 45 mL of 1xPBS and incubated with 800 µL

of streptavidin microbeads, followed by sorting with the autoMACS Pro Cell Separator. Sorted

cells were immediately crosslinked with 1% formaldehyde for 20 min at room temperature, then

transferred to screw cap microcentrifuge tubes and the pellets flash frozen in liquid nitrogen.

Cells were thawed and lysed in 600 µL FA140 Lysis buffer (50 mM HEPES, 140 mM NaCl, 1%

Triton X-100, 1 mM EDTA, 0.1% SDS, 0.1 mM PMSF, 1x protease inhibitor cocktail; Sigma)

by shaking with acid-washed glass beads in a Mini-Bead beater (Biospec Products). Cell lysates

were recovered and sonicated for 30 cycles of 30 sec "on" and 30 sec "off" in a Diagenode

Bioruptor followed by centrifugation at 16,000 x g. A 1/10th supernatant volume input was taken

for each sample and crosslinking reversed by incubating overnight at 65°C in 150 µL elution

buffer (TE, 1% SDS). The remaining supernatant was used for immunoprecipitation overnight at

4°C with 5 µg of primary antibody and 30 µL of protein G magnetic beads (Pierce), followed by

washing 1x with FA-140 buffer, 2x with FA-500 buffer (FA-140 with 500 mM NaCl), and 2x

with LiCl solution (10 mM Tris-HCl, pH 8.0, 250 mM LiCl, 0.5% NP-40, 0.5% SDS, 1 mM

EDTA). DNA was eluted twice with 75 µL of elution buffer in a 65°C water bath for 15 min.

The eluates were combined and crosslinking reversed. Input and ChIP DNA samples were

purified by an Invitrogen PureLink™ PCR purification kit. Finally, ChIP DNA was quantified by

real-time PCR and normalized to the input DNA signal. Young cells were collected from flow

through out of the autoMACS cell sorter and then processed as described for the aged cells. For

ChIP assays on unsorted populations, cells were grown to log phase in 100 mL YPD before

formaldehyde cross-linking for 20 min at 30°C while shaking. Additionally, 2.5 mg of protein

was used for the IP instead of the entire lysate, and the chromatin solution was sonicated for 50

cycles rather than 30.

ChIP-Seq Library Preparation

Chromatin was prepared in a similar manner to old cell ChIP assays with the following

exceptions. Cells were grown to log phase in 200 mL YPD and fixed for 20 min with 1%

formaldehyde. Lysate volumes were adjusted to 4 mL in FA-140 lysis buffer and sonicated 5

times with a Branson 250 Digital Sonifier for 30 sec at 70% amplitude. Samples were cooled on

ice for at least 1 min between sonication periods. The lysate was split into 3 microcentrifuge

tubes and spun for 10 min at 10,000 rpm in a microfuge at 4°C. The entire supernatant from 1

tube was used for immunoprecipitation overnight with 15 μg of α-Myc 9E10 antibody and 30 μL

of protein G magnetic beads (Pierce). After washing, elution, and reverse cross-linking, 100 mg

of Proteinase K was added for 2 hr at 42°C. Chromatin was purified in a Zymogen DNA Clean

and Concentrator-5 Kit and quantified using a Qubit fluorometric device. A minimum of 20 ng

of DNA was used as input for NEBNext® library construction. ML150 libraries were sequenced

on an Illumina Nextseq and ML163 libraries on an Illumina Miseq by the UVA Genome

Analysis and Technology Core. Sequencing files are available at GEO accession number

GSE117037. Sequencing reads were mapped to the sacCer3 genome using bowtie2 with default

settings. Corresponding bam files were further processed to produce the coverage plots shown in

Fig 1 with the Deeptools package of Galaxy (https://usegalaxy.org/).

Sister chromatid cohesion assay

From 50 mL log phase SC cultures of strains 3349-1B, 3312-7A, and 3460-2A, 5×10^7

cells were washed and biotinylated as described in the Old Cell Isolation section. This population

was transferred into a 1.5 L SC culture and allowed to grow for 12 hr. The original biotinylated

population was then purified by incubation with 300 μL of streptavidin micro beads followed by

gravity filtration through a Miltenyi LS column. The column was washed twice with 5 ml of PBS

and the enriched cell/bead population was aged for an additional 12 hr in a fresh 1.5 L SC culture

before being purified for a second time with an additional 200 μL of beads. Eluted cells (still

bound to beads) were processed as described below for young cells.

From the original log phase culture, 5×10^7 cells were arrested in a fresh 50 mL SC culture

containing 10 µg/mL nocodazole for 1.5 hr. For the *mcd1-1* strain 3312-7A, cells were also

shifted to 36°C at this time. Bud scars were stained with the addition of 1 mL of supernatant

from a solution of PBS and 5 mg of Calcofluor white that was centrifuged to eliminate

aggregates. Non-arrested cells were directly stained with calcofluor. Cells were then pelleted and

washed in PBS. Following staining, 200 µL of 4% paraformaldehyde was added directly to the

cell pellet and allowed to crosslink for 15 min at room temperature. The cell pellet was washed

once with PBS and resuspended in ~100-200 µL of 0.1 MKPO$_4$/1 M sorbitol, pH 6.5. Images

were captured with a Zeiss Axio Observer z1 widefield microscope using a 64x oil objective

lens.

Replicative lifespan assays

Lifespan assays were carried out essentially the same as previously described [84].

Briefly, small aliquots of log phase cultures were dripped in a straight line onto solid agar YPD

with 2% glucose. From the initial populations, a minimum of 32 virgin daughter cells were

picked for lifespan assays with daughter cells being selectively pulled away from mother cells

using a fiberoptic dissection needle and on a Nikon Eclipse 400 microscope. All virgin daughters

were required to bud at least one time to be included in the experiment and dissection was

carried out over the course of several days with temporary incubation at 4°C in between

dissection periods to stop division. Cells were considered dead when they stopped dividing for a

minimum of 2 generation times (180 min).

Chromosome loss (sectoring) assay

The colony sectoring assay was performed on SC plates with adenine limited to 80 µM.

Frequency of mini-chromosome loss represents the number of ½ or ¼ red/white sectored

colonies divided by the sum of sectored and white colonies. Cells were plated to an approximate

density of 500 cells/plate based on counts from a Brightline hemacytometer. Any plates bearing

greater than 1000 cells were discarded. Three biological replicates of each strain were

performed, with at least 10 plates counted per replicate. For old cell populations, ~5x10^6

biotinylated cells were aged in 1.5 L of YPD for 24 hr. Cells were incubated with 300 µL of

streptavidin magnetic beads (New England Biolabs) and manually washed 4 times with PBS on a

magnetic stand, then plated onto the limiting adenine SC plates such that ~500 colonies

appeared. Bud scars were not counted because the size of the beads prohibited visualization.

RT-qPCR measurement of *MCD1* overexpression

Doxycycline was added to log phase cultures at a concentration of 2 µg/mL for 6 hr in

order to induce expression of MCD1 from pRF4. Total RNA was extracted using a standard acid

phenol extraction protocol [85]. cDNA was created from ~1 µg of RNA using a Verso cDNA

synthesis kit (Thermo Fisher). *MCD1* expression levels were quantified on an Applied

Biosystems StepOne real time PCR machine with primers JS2844 and JS2949, and normalized to

actin transcript levels (primers JS1146 and JS1147).

Hi-C analysis

Log-phase cultures were cross-linked with 3% formaldehyde for 20 min and quenched

with a 2x volume of 2.5M Glycine. Cell pellets were washed with dH$_2$O and stored at -80°C.

Thawed cells were resuspended in 5 ml of 1X NEB2 restriction enzyme buffer (New England

Biolabs) and poured into a pre-chilled mortar containing liquid N_2. Nitrogen grinding was

performed twice as previously described [86], and the lysates were then diluted to an OD_{600} of 12

in 1x NEB2 buffer. 500 µl of cell lysate was used for each Hi-C library as follows. Lysates were

solubilized by the addition of 50 µl 1% SDS and incubation at 65°C for 10 min. 55 µl of 10%

TritonX-100 was added to quench the SDS, followed by 10 µl of 10X NEB2 buffer and 15 µl of

*Hin*dIII (New England Biolabs, 20 U/µl) to digest at 37°C for 2 hr. An additional 10 µl of

*Hin*dIII was added for digestion overnight. The remainder of the protocol was based on

previously published work with minor exceptions [87], see supplemental experimental

procedures for details. Hi-C sequencing libraries were prepared with reagents from an Illumina

Nextera Mate Pair Kit (FC-132-1001) using the standard Illumina protocol of End Repair, A-

tailing, Adapter Ligation, and 12 cycles of PCR. PCR products were size selected and purified

with AMPure XP beads before sequencing with an Illumina Miseq or Hiseq.

Author Contributions

Conceptualization; R.D.F. and J.S.S.; Methodology, R.D.F., M.L. and J.S.S.; Software, R.D.F.;

Strain Creation and Validation, R.D.F., N.M., E.F., and M.L.; Plasmid Creation and Validation,

R.D.F., E.F., and M.L; Formal Analysis, R.D.F.; Data Curation, R.D.F.; Writing-Original Draft,

R.D.F. and J.S.S.; Writing Review & Editing, R.D.F., M.L., and J.S.S; Supervision, J.S.S.;

Project Administration, J.S.S.; Funding Acquisition, J.S.S.

Acknowledgments

We thank Dan Gottschling and all lab members for kindly providing yeast strains and initial

advice on using the MEP system. Stefan Bekiranov, Job Dekker, Jon Belton, Maitreya Dunham,

Ivan Liachko, Maxim Imakev, and Anton Goloborodko all provided valuable advice on Hi-C protocols and analysis methods. We thank Doug Koshland for providing the Mcd1 reduction and cohesion assay strains, and Matt Kaeberlein for rDNA marker loss strains. Special thanks to Todd Stukenberg for microscopy assistance. Lastly, thanks to David Auble for critically reading the manuscript and providing comments prior to submission. We declare no conflicts of interest.

References

1. Mortimer RK, Johnston JR. Life span of individual yeast cells. Nature. 1959;183(4677):1751-2. PMID: 13666896.

2. Wasko BM, Kaeberlein M. Yeast replicative aging: a paradigm for defining conserved longevity interventions. FEMS Yeast Res. 2014;14(1):148-59. PMID: 24119093.

3. Buck SW, Gallo CM, Smith JS. Diversity in the Sir2 family of protein deacetylases. J Leukoc Biol. 2004;75(6):939-50. PMID: 14742637.

4. Gomes AP, Price NL, Ling AJ, Moslehi JJ, Montgomery MK, Rajman L, et al. Declining NAD$^+$ induces a pseudohypoxic state disrupting nuclear-mitochondrial communication during aging. Cell. 2013;155(7):1624-38. PMID: 24360282.

5. Brachmann CB, Sherman JM, Devine SE, Cameron EE, Pillus L, Boeke JD. The *SIR2* gene family, conserved from bacteria to humans, functions in silencing, cell cycle progression, and chromosome stability. Genes Dev. 1995;9(23):2888-902. PMID: 7498786.

6. Rine J, Herskowitz I. Four genes responsible for a position effect on expression from *HML* and *HMR* in *Saccharomyces cerevisiae*. Genetics. 1987;116(1):9-22. PMID: 3297920.

7. Gartenberg MR, Smith JS. The nuts and bolts of transcriptionally silent chromatin in *Saccharomyces cerevisiae*. Genetics. 2016;203(4):1563-99. PMID: 27516616.

8.   Dang W, Steffen KK, Perry R, Dorsey JA, Johnson FB, Shilatifard A, et al. Histone H4

lysine 16 acetylation regulates cellular lifespan. Nature. 2009;459(7248):802-7. PMID:

19516333.

9.   Smeal T, Claus J, Kennedy B, Cole F, Guarente L. Loss of transcriptional silencing

causes sterility in old mother cells of *S. cerevisiae*. Cell. 1996;84(4):633-42. PMID: 8598049.

10.   Schlissel G, Krzyzanowski MK, Caudron F, Barral Y, Rine J. Aggregation of the Whi3

protein, not loss of heterochromatin, causes sterility in old yeast cells. Science.

2017;355(6330):1184-7. PMID: 28302853.

11.   Bryk M, Banerjee M, Murphy M, Knudsen KE, Garfinkel DJ, Curcio MJ. Transcriptional

silencing of Ty1 elements in the *RDN1* locus of yeast. Genes Dev. 1997;11(2):255-69. PMID:

9009207.

12.   Smith JS, Boeke JD. An unusual form of transcriptional silencing in yeast ribosomal

DNA. Genes Dev. 1997;11(2):241-54. PMID: 9009206.

13.   Shou W, Seol JH, Shevchenko A, Baskerville C, Moazed D, Chen ZW, et al. Exit from

mitosis is triggered by Tem1-dependent release of the protein phosphatase Cdc14 from nucleolar

RENT complex. Cell. 1999;97(2):233-44. PMID: 10219244.

14.   Straight AF, Shou W, Dowd GJ, Turck CW, Deshaies RJ, Johnson AD, et al. Net1, a

Sir2-associated nucleolar protein required for rDNA silencing and nucleolar integrity. Cell.

1999;97(2):245-56. PMID: 10219245.

15.   Li C, Mueller JE, Bryk M. Sir2 represses endogenous polymerase II transcription units in

the ribosomal DNA nontranscribed spacer. Mol Biol Cell. 2006;17(9):3848-59. PMID:

16807355.

16.   Kobayashi T, Ganley AR. Recombination regulation by transcription-induced cohesin

dissociation in rDNA repeats. Science. 2005;309(5740):1581-4. PMID: 16141077.

17.   Sinclair DA, Guarente L. Extrachromosomal rDNA circles--a cause of aging in yeast.

Cell. 1997;91(7):1033-42. PMID: 9428525.

18.   Defossez PA, Prusty R, Kaeberlein M, Lin SJ, Ferrigno P, Silver PA, et al. Elimination of

replication block protein Fob1 extends the life span of yeast mother cells. Mol Cell.

1999;3(4):447-55. PMID: 10230397.

19.   Kobayashi T, Horiuchi T. A yeast gene product, Fob1 protein, required for both

replication fork blocking and recombinational hotspot activities. Genes Cells. 1996;1(5):465-74.

PMID: 9078378.

20.   Takeuchi Y, Horiuchi T, Kobayashi T. Transcription-dependent recombination and the

role of fork collision in yeast rDNA. Genes Dev. 2003;17(12):1497-506. PMID: 12783853.

21.   Ganley AR, Kobayashi T. Ribosomal DNA and cellular senescence: new evidence

supporting the connection between rDNA and aging. FEMS Yeast Res. 2014;14(1):49-59.

PMID: 24373458.

22.   Ganley AR, Ide S, Saka K, Kobayashi T. The effect of replication initiation on gene

amplification in the rDNA and its relationship to aging. Mol Cell. 2009;35(5):683-93. PMID:

19748361.

23.   Kaeberlein M, McVey M, Guarente L. The SIR2/3/4 complex and SIR2 alone promote

longevity in *Saccharomyces cerevisiae* by two different mechanisms. Genes Dev.

1999;13(19):2570-80. PMID: 10521401.

24.   Marston AL. Chromosome segregation in budding yeast: sister chromatid cohesion and

related mechanisms. Genetics. 2014;196(1):31-63. PMID: 24395824.

25.   Wood AJ, Severson AF, Meyer BJ. Condensin and cohesin complexity: the expanding

repertoire of functions. Nat Rev Genet. 2010;11(6):391-404. PMID: 20442714.

26.   Choy JS, Acuna R, Au WC, Basrai MA. A role for histone H4K16 hypoacetylation in

Saccharomyces cerevisiae kinetochore function. Genetics. 2011;189(1):11-21. PMID: 21652526.

27.   Holmes SG, Rose AB, Steuerle K, Saez E, Sayegh S, Lee YM, et al. Hyperactivation of

the silencing proteins, Sir2p and Sir3p, causes chromosome loss. Genetics. 1997;145(3):605-14.

PMID: 9055071.

28.   Sunshine AB, Ong GT, Nickerson DP, Carr D, Murakami CJ, Wasko BM, et al.

Aneuploidy shortens replicative lifespan in *Saccharomyces cerevisiae*. Aging Cell.

2016;15(2):317-24. PMID: 26762766.

29.   Glynn EF, Megee PC, Yu HG, Mistrot C, Unal E, Koshland DE, et al. Genome-wide

mapping of the cohesin complex in the yeast *Saccharomyces cerevisiae*. PLoS Biol.

2004;2(9):E259. PMID: 15309048.

30.   Li M, Valsakumar V, Poorey K, Bekiranov S, Smith JS. Genome-wide analysis of

functional sirtuin chromatin targets in yeast. Genome Biol. 2013;14(5):R48. PMID: 23710766.

31.   Kobayashi T, Horiuchi T, Tongaonkar P, Vu L, Nomura M. SIR2 regulates

recombination between different rDNA repeats, but not recombination within individual rRNA

genes in yeast. Cell. 2004;117(4):441-53. PMID: 15137938.

32.   Stephens AD, Quammen CW, Chang B, Haase J, Taylor RM, 2nd, Bloom K. The spatial

segregation of pericentric cohesin and condensin in the mitotic spindle. Mol Biol Cell.

2013;24(24):3909-19. PMID: 24152737.

33.   Lindstrom DL, Gottschling DE. The mother enrichment program: a genetic system for

facile replicative life span analysis in *Saccharomyces cerevisiae*. Genetics. 2009;183(2):413-22.

PMID: 19652178.

34.   Lindstrom DL, Leverich CK, Henderson KA, Gottschling DE. Replicative age induces

mitotic recombination in the ribosomal RNA gene cluster of *Saccharomyces cerevisiae*. PLoS

Genet. 2011;7(3):e1002015. PMID: 21436897.

35.   Moazed D, Kistler A, Axelrod A, Rine J, Johnson AD. Silent information regulator

protein complexes in Saccharomyces cerevisiae: a SIR2/SIR4 complex and evidence for a

regulatory domain in SIR4 that inhibits its interaction with SIR3. Proc Natl Acad Sci USA.

1997;94(6):2186-91. PMID: 9122169.

36.   Oppikofer M, Kueng S, Martino F, Soeroes S, Hancock SM, Chin JW, et al. A dual role

of H4K16 acetylation in the establishment of yeast silent chromatin. EMBO J.

2011;30(13):2610-21. PMID: 21666601.

37.   Heidinger-Pauli JM, Mert O, Davenport C, Guacci V, Koshland D. Systematic reduction

of cohesin differentially affects chromosome segregation, condensation, and DNA repair. Curr

Biol. 2010;20(10):957-63. PMID: 20451387.

38.   Guacci V, Koshland D. Cohesin-independent segregation of sister chromatids in budding

yeast. Mol Biol Cell. 2012;23(4):729-39. PMID: 22190734.

39.   Unal E, Heidinger-Pauli JM, Kim W, Guacci V, Onn I, Gygi SP, et al. A molecular

determinant for the establishment of sister chromatid cohesion. Science. 2008;321(5888):566-9.

PMID: 18653894.

40.   Guacci V, Koshland D, Strunnikov A. A direct link between sister chromatid cohesion

and chromosome condensation revealed through the analysis of *MCD1* in *S. cerevisiae*. Cell.

1997;91(1):47-57. PMID: 9335334.

41.   Spencer F, Gerring SL, Connelly C, Hieter P. Mitotic chromosome transmission fidelity

mutants in *Saccharomyces cerevisiae*. Genetics. 1990;124(2):237-49. PMID: 2407610.

42.   Hickman MA, Rusche LN. Substitution as a mechanism for genetic robustness: the

duplicated deacetylases Hst1p and Sir2p in *Saccharomyces cerevisiae*. PLoS Genet.

2007;3(8):e126. PMID: 17676954.

43.   Li M, Petteys BJ, McClure JM, Valsakumar V, Bekiranov S, Frank EL, et al. Thiamine

biosynthesis in *Saccharomyces cerevisiae* is regulated by the NAD^+-dependent histone

deacetylase Hst1. Mol Cell Biol. 2010;30(13):3329-41. PMID: 20439498.

44.   Duan Z, Andronescu M, Schutz K, McIlwain S, Kim YJ, Lee C, et al. A three-

dimensional model of the yeast genome. Nature. 2010;465(7296):363-7. PMID: 20436457.

45.   Gotta M, Strahl-Bolsinger S, Renauld H, Laroche T, Kennedy BK, Grunstein M, et al.

Localization of Sir2p: the nucleolus as a compartment for silent information regulators. EMBO J.

1997;16(11):3243-55. PMID: 9214640.

46.   Schalbetter SA, Goloborodko A, Fudenberg G, Belton JM, Miles C, Yu M, et al. SMC

complexes differentially compact mitotic chromosomes according to genomic context. Nat Cell

Biol. 2017;19(9):1071-80. PMID: 28825700.

47.   Belli G, Gari E, Piedrafita L, Aldea M, Herrero E. An activator/repressor dual system

allows tight tetracycline-regulated gene expression in budding yeast. Nucleic Acids Res.

1998;26(4):942-7. PMID: 9461451.

48.   Jiang JC, Jaruga E, Repnevskaya MV, Jazwinski SM. An intervention resembling caloric

restriction prolongs life span and retards aging in yeast. FASEB J. 2000;14(14):2135-7. PMID:

11024000.

49.   Lin SJ, Kaeberlein M, Andalis AA, Sturtz LA, Defossez PA, Culotta VC, et al. Calorie

restriction extends *Saccharomyces cerevisiae* lifespan by increasing respiration. Nature.

2002;418(6895):344-8. PMID: 12124627.

50.   Pal S, Postnikoff SD, Chavez M, Tyler JK. Impaired cohesion and homologous

recombination during replicative aging in budding yeast. Sci Adv. 2018;4(2):eaaq0236. PMID:

29441364.

51.   Hu Z, Chen K, Xia Z, Chavez M, Pal S, Seol JH, et al. Nucleosome loss leads to global

transcriptional up-regulation and genomic instability during yeast aging. Genes Dev.

2014;28(4):396-408. PMID: 24532716.

52.   Song S, Johnson FB. Epigenetic mechanisms impacting aging: A focus on histone levels

and telomeres. Genes. 2018;9(4). PMID: 29642537.

53.   Buchwalter A, Hetzer MW. Nucleolar expansion and elevated protein translation in

premature aging. Nat Commun. 2017;8(1):328. PMID: 28855503.

54.   Mehta R, Chandler-Brown D, Ramos FJ, Shamieh LS, Kaeberlein M. Regulation of

mRNA translation as a conserved mechanism of longevity control. Adv Exp Med Biol.

2010;694:14-29. PMID: 20886753.

55.   Janssens GE, Veenhoff LM. The natural variation in lifespans of single yeast cells is

related to variation in cell size, ribosomal protein, and division time. PLoS One.

2016;11(12):e0167394. PMID: 27907085.

56.   Janssens GE, Meinema AC, Gonzalez J, Wolters JC, Schmidt A, Guryev V, et al. Protein

biogenesis machinery is a driver of replicative aging in yeast. eLife. 2015;4:e08527. PMID:

26422514.

57.   Hsu HC, Wang CL, Wang M, Yang N, Chen Z, Sternglanz R, et al. Structural basis for

allosteric stimulation of Sir2 activity by Sir4 binding. Genes Dev. 2013;27(1):64-73. PMID:

23307867.

58.   Dasgupta A, Ramsey KL, Smith JS, Auble DT. Sir Antagonist 1 (San1) is a ubiquitin

ligase. J Biol Chem. 2004;279(26):26830-8. PMID: 15078868.

59.   Gardner RG, Nelson ZW, Gottschling DE. Degradation-mediated protein quality control

in the nucleus. Cell. 2005;120(6):803-15. PMID: 15797381.

60.   Kriegenburg F, Jakopec V, Poulsen EG, Nielsen SV, Roguev A, Krogan N, et al. A

chaperone-assisted degradation pathway targets kinetochore proteins to ensure genome stability.

PLoS Genet. 2014;10(1):e1004140. PMID: 24497846.

61.   Sasaki T, Maier B, Bartke A, Scrable H. Progressive loss of SIRT1 with cell cycle

withdrawal. Aging Cell. 2006;5(5):413-22. PMID: 16939484.

62.   Hoggard TA, Chang F, Perry KR, Subramanian S, Kenworthy J, Chueng J, et al. Yeast

heterochromatin regulators Sir2 and Sir3 act directly at euchromatic DNA replication origins.

PLoS Genet. 2018;14(5):e1007418. PMID: 29795547.

63.   Utani K, Aladjem MI. Extra View: Sirt1 acts as A gatekeeper of replication initiation to

preserve genomic stability. Nucleus. 2018;9(1):261-7. PMID: 29578371.

64.   Yoshida K, Bacal J, Desmarais D, Padioleau I, Tsaponina O, Chabes A, et al. The histone

deacetylases sir2 and rpd3 act on ribosomal DNA to control the replication program in budding

yeast. Mol Cell. 2014;54(4):691-7. PMID: 24856221.

65.   Kwan EX, Foss EJ, Tsuchiyama S, Alvino GM, Kruglyak L, Kaeberlein M, et al. A

natural polymorphism in rDNA replication origins links origin activation with calorie restriction

and lifespan. PLoS Genet. 2013;9(3):e1003329. PMID: 23505383.

66.   Jessberger R. Age-related aneuploidy through cohesion exhaustion. EMBO Rep.

2012;13(6):539-46. PMID: 22565322.

67.   Lister LM, Kouznetsova A, Hyslop LA, Kalleas D, Pace SL, Barel JC, et al. Age-related

meiotic segregation errors in mammalian oocytes are preceded by depletion of cohesin and Sgo2.

Curr Biol. 2010;20(17):1511-21. PMID: 20817533.

68.   Perkins AT, Das TM, Panzera LC, Bickel SE. Oxidative stress in oocytes during

midprophase induces premature loss of cohesion and chromosome segregation errors. Proc Natl

Acad Sci USA. 2016;113(44):E6823-e30. PMID: 27791141.

69.   Baker DJ, Jeganathan KB, Cameron JD, Thompson M, Juneja S, Kopecka A, et al.

BubR1 insufficiency causes early onset of aging-associated phenotypes and infertility in mice.

Nat Genet. 2004;36(7):744-9. PMID: 15208629.

70.   Baker DJ, Dawlaty MM, Wijshake T, Jeganathan KB, Malureanu L, van Ree JH, et al.

Increased expression of BubR1 protects against aneuploidy and cancer and extends healthy

lifespan. Nat Cell Biol. 2013;15(1):96-102. PMID: 23242215.

71.   North BJ, Rosenberg MA, Jeganathan KB, Hafner AV, Michan S, Dai J, et al. SIRT2

induces the checkpoint kinase BubR1 to increase lifespan. EMBO J. 2014;33(13):1438-53.

PMID: 24825348.

72.   Uhlmann F. SMC complexes: from DNA to chromosomes. Nat Rev Mol Cell Biol.

2016;17(7):399-412. PMID: 27075410.

73.   Huang J, Brito IL, Villen J, Gygi SP, Amon A, Moazed D. Inhibition of homologous

recombination by a cohesin-associated clamp complex recruited to the rDNA recombination

enhancer. Genes Dev. 2006;20(20):2887-901. PMID: 17043313.

74.   Bitto A, Wang AM, Bennett CF, Kaeberlein M. Biochemical genetic pathways that

modulate aging in multiple species. Cold Spring Harb Perspect Med. 2015;5(11). PMID:

26525455.

75.   Ohkuni K, Kitagawa K. Endogenous transcription at the centromere facilitates

centromere activity in budding yeast. Curr Biol. 2011;21(20):1695-703. PMID: 22000103.

76.   Smith JS, Caputo E, Boeke JD. A genetic screen for ribosomal DNA silencing defects

identifies multiple DNA replication and chromatin-modulating factors. Mol Cell Biol.

1999;19(4):3184-97. PMID: 10082585.

77.   O'Sullivan JM, Sontam DM, Grierson R, Jones B. Repeated elements coordinate the

spatial organization of the yeast genome. Yeast. 2009;26(2):125-38. MID: 19235779.

78.   Nagaoka SI, Hassold TJ, Hunt PA. Human aneuploidy: mechanisms and new insights

into an age-old problem. Nat Rev Genet. 2012;13(7):493-504. PMID: 22705668.

79.   Lushnikova T, Bouska A, Odvody J, Dupont WD, Eischen CM. Aging mice have

increased chromosome instability that is exacerbated by elevated Mdm2 expression. Oncogene.

2011;30(46):4622-31. PMID: 21602883.

80.   Riesen M, Morgan A. Calorie restriction reduces rDNA recombination independently of

rDNA silencing. Aging Cell. 2009;8(6):624-32. PMID: 19732046.

81.   Smith DL, Jr., Li C, Matecic M, Maqani N, Bryk M, Smith JS. Calorie restriction effects

on silencing and recombination at the yeast rDNA. Aging Cell. 2009;8(6):633-42. PMID:

19732044.

82.   Matecic M, Smith DL, Pan X, Maqani N, Bekiranov S, Boeke JD, et al. A microarray-

based genetic screen for yeast chronological aging factors. PLoS Genet. 2010;6(4):e1000921.

PMID: 20421943.

83.   Buck SW, Sandmeier JJ, Smith JS. RNA polymerase I propagates unidirectional

spreading of rDNA silent chromatin. Cell. 2002;111(7):1003-14. PMID: 12507427.

84.   Steffen KK, Kennedy BK, Kaeberlein M. Measuring replicative life span in the budding

yeast. J Vis Exp. 2009;(28). PMID: 19556967.

85.   Ausubel FM, Brent R, Kingston RE, Moore DD, Seidman JG, Smith JA, et al., editors.

Current Protocols in Molecular Biology. New York: John Wiley & Sons, Inc.; 2000.

86.   Belton JM, Dekker J. Measuring Chromatin Structure in Budding Yeast. Cold Spring

Harb Protoc. 2015;2015(7):614-8. PMID: 26134912.

87.   Burton JN, Liachko I, Dunham MJ, Shendure J. Species-level deconvolution of

metagenome assemblies with Hi-C-based contact probability maps. G3. 2014;4(7):1339-46.

PMID: 24855317.

Figure captions

Fig 1. Cohesin distribution at the rDNA and centromeres in WT and *sir2*Δ cells. (A)

Composite plot of Mcd1-13xMyc ChIP-Seq binding across the rDNA array from WT and *sir2*Δ

log-phase cells binned at 50bp resolution. **(B)** ChIP-qPCR quantifying Mcd1 enrichment at the

rDNA IGS1 region from log-phase cells. **(C)** Composite plot of Mcd1-13xMyc enrichment at all

16 yeast point centromeres. **(D)** ChIP-qPCR quantifying Mcd1 binding at two centromeres and a

pericentromeric site 15kb away from *CEN4*. (* p<0.005, two-tailed student t-test).

Fig 2. Depletion of Sir2 complexes and cohesin in replicatively aging cells. (A) Western blot

of Sir2 protein levels in young and aged cells. Vma2 serves as a loading control. **(B)** Depiction

of limiting Sir2 shared between the SIR and RENT complexes. **(C)** Western blot of Sir4 proteins

in young and aged cells. **(D)** Western blot of Sir3 in young and aged cells. *(E)* Western blot of

13x-Myc tagged Net1 in young and aged cells. **(F)** Western blot of Mcd1-13xMyc in young and

aged cells. **(G)** ChIP-qPCR of Mcd1-13xMyc in young and aged cells normalized to background

signal at an intergenic site near *PDC1*. **(H)** ChIP-qPCR of Sir4-13xMyc in young and aged cells

normalized to the intergenic *PDC1* site. Bud scar counts indicate the average for each enriched

population used for the western blotting. Qty indicates the mean western signal of each protein in

aged cells relative to the signal in young cells, which is set at 1.0 (n=3 biological replicates).

Standard errors are provided in Supplementary Fig 1B.

Fig 3. Sister chromatid cohesion is weakened in old yeast cells. (A) Representative control

images of arm cohesion in an *mcd1-1* mutant at permissive (25°C) and non-permissive (37°C)

temperatures. **(B)** Quantification of cohesion maintenance (1-dot) or loss (2-dots) from 100 cells.

(C) Representative images of young (log-phase) or aged cells arrested with nocodazole (Nz) and

monitored for *CENIV* arm cohesion at the *LYS4* locus. **(D)** Images of cohesion at the *CEN4* locus

in nocodazole-arrested young and aged cells. **(E)** Images of centromeric (*CEN*4) cohesion in

young and aged cells not treated with Nz. **(F)** Quantifying cohesion loss (2-dots) young cells or

aged cells with at least 5 bud scars (100 cells of each population). *p<0.05, two-tailed Fischer's

exact test.

Fig 4. Chromosomal instability during replicative aging is linked to the rDNA. (A)

Schematic of artificial chromosome loss assay for ½ sectored colonies. **(B)** Quantification of

sectoring frequency in young or aged WT, *sir2Δ*, *hst1Δ*, and *sir2Δ hst1Δ* strains. *HMR* or *FOB1*

were also deleted where indicated. Approximately 10,000 colonies were analyzed for each strain,

in several biological replicates. *p<0.05, two-tailed student t-test. **(C)** Iteratively corrected (IC)

and read-normalized heatmap of ChrXII Hi-C data at 10 kb resolution revealing a driving

interaction between *CEN12* (bin 15) and unique sequence adjacent to the rDNA (bin 45). **(D)**

Observed/expected statistical heatmap of ChrXII Hi-C data at 10 kb resolution using HOMER.

Yellow bracket indicates a putative loop structure between bins 15 and 45. **(E)** IC/read-

normalized Hi-C heatmap of chromosomes XI, XII, and XIII showing centromere clustering.

Yellow arrows indicate examples of centromere alignment with *CEN12*.

Fig 5. Galactose induces CIN and shortens yeast RLS. (A) Representative images of

chromosome loss (sectoring) for WT (YPH278) cells grown continuously in 2% glucose,

galactose, or raffinose. **(B)** Quantification of half-sector colonies for WT (YPH278) cells grown

on each carbon source. *p<0.01, two-tailed student t-test. **(C)** RLS assay of YPH278 cells (n=64)

growing on rich YEP agar plates containing either 2% glucose, galactose, or raffinose. **(D)** RLS

of WT (BY4741) cells (n=32) growing on 2% glucose, galactose, or raffinose.

Fig 6. Modulation of RLS by manipulating *MCD1* and *SIR2* expression levels. (A) RLS

assays of WT (*MCD1*$^{+}$) and *mcd1L12STOP* strains containing an integrated empty tet^{on}

expression vector or the same vector overexpressing *MCD1* (n=96 cells each strain). **(B)** RLS

assay of WT and *mcd1L12STOP* strains containing an integrated empty pRS306 vector or the

SIR2 vector, pJSB186 (n=40 cells each). **(C)** RLS assay showing *MCD1* overexpression does not

rescue the short RLS of a *sir2Δhst1Δ* mutant. **(D)** rDNA recombination (marker loss) assay with

a strain bearing *ADE2* within the rDNA array. Either an empty tet^{on} vector (pRF10) or an *MCD1*

overexpressing version (pRF11) were integrated at the *leu2Δ1* locus of WT or *sir2Δ* versions of

W303AR (* p<0.05, two-tailed t-test).

Fig 7. CR effects are upstream of Sir2 and Mcd1 function in old cells. **(A)** RLS assay of WT

(JH5275b) and *mcd1L12STOP* (JH5276b) strains under normal 2% glucose and CR (0.5%

glucose) conditions (n=32 cells). **(B)** Western blot of Mcd1-13xMyc levels in young and aged

cells of the MEP strain RF10. Cultures were grown in YEP media containing either 2% or 0.5%

glucose. **(C)** Chromosome loss (sectoring) assay showing reduced sectoring in WT (YPH278),

sir2Δ (RF32), *hst1Δ* (RF33), and *sir2Δ hst1Δ* (RF43) strains when grown in media containing

0.5% glucose compared to 2%. * p<0.05, two-tailed t-test. **(D)** RLS assay of WT and *sir2Δhst1Δ*

mutant strains under normal and CR conditions. (n=32 cells).

Fig 8. Model of residual cohesin redistribution from rDNA to centromeres in aging yeast

cells. Centromeres of all 16 chromosomes cluster, and the rDNA remains isolated. The rDNA

may also contact the centromeres during anaphase, which could impact the fidelity of

chromosome segregation or promote recombination between the rDNA and other chromosomes.

Cohesin and Sir2-containing complexes are depleted in cells, but the remaining cohesin complex

preferentially associates with centromeres in a futile attempt to maintain sister chromatid

cohesion.

Supporting Figure and Table Captions

S1 Fig. Detailed western blot and bud scar count quantification. (A) Bar graph showing

average bud scar counts from young and aged cell populations used for western blotting. **(B)**

Detailed bud scar counts and western blot quantitation from the triplicate biological samples. **(C)**

Representative image of an enriched old cell population from the third Lrs4-13xMyc biological

replicate. Inset image allows counting of individual bud scars.

S2 Fig. Western blots of additionally tested proteins from young and aged cells. **(A)** A

13xMyc-tagged version of the Sir2 paralog, Hst1, is also depleted in aged cells. Hst1 is in a

complex with Sum1 and Rfm1. **(B)** The Myc-tagged Smc1 subunit of cohesin is depleted in aged

cells to similar levels as Mcd1. The yeast cohesin complex consists of Smc1, Smc3, Mcd1, and

Irr1. Smc1 and Smc3 form the coiled-coil loop structure. **(C)** A 13x-Myc tagged version of the

cohesin loading complex subunit, Scc2, also decreases in old cells. The cohesin loading complex

consists of Scc2 and Scc4.

S3 Fig. RLS of the cohesion visualization strains is normal and unaffected by position of

the lacO array. Strain 3349-1B contains a lacO array at the *LYS4* locus on Chr. IV and is used

as a proxy for arm cohesion, while strain 3460-2A is used to monitor centromeric cohesion 10 kb

away from the *CEN4* locus.

S4 Fig. Confirmation of doxycycline-induced *MCD1* overexpression in WT and

mcd1L12STOP strains. RT-qPCR of *MCD1* transcript levels relative to actin transcript levels

were quantified from the empty vector strains (RF146 and RF147) or the *MCD1* overexperssion

strains RF179 and RF180. Total RNA was isolated following 6 hours doxycycline induction

during log-phase growth.

S1 Table. Yeast Strains. This table lists each *Saccharomyces cerevisiae* strain used in the study,

along with its genotype and source.

S2 Table. Oligonucleotides used in the study.

A ChrIV Arm Cohesion (Nz)

mcd1-1 25C   *mcd1-1* 37C

B

C ChrIV Arm Cohesion (Nz)

Young   24hrs Aged

D ChrIV Cen. Cohesion (Nz)

Young   24hrs Aged

E ChrIV Cen. Cohesion

Young   24hrs Aged

F

Figure 7