1 # polyRAD: Genotype calling with uncertainty from
2 # sequencing data in polyploids and diploids
3

Lindsay V. Clark[1], Alexander E. Lipka, and Erik J. Sacks

Department of Crop Sciences, University of Illinois at Urbana-Champaign, Urbana, Illinois

61801, USA.

ORCID ID: 0000-0002-3881-9252 (L.V.C.)

[1]Corresponding author: Department of Crop Sciences, University of Illinois at Urbana-

Champaign, 1201 W. Gregory Dr., Urbana, IL 61801. Email: lvclark@illinois.edu

Keywords: next-generation DNA sequencing; polyploidy; Bayesian genotype calling; single

nucleotide polymorphism; genotype imputation; R

Abstract

Low or uneven read depth is a common limitation of genotyping-by-sequencing (GBS)

and restriction site-associated DNA sequencing (RAD-seq), resulting in high missing data rates,

heterozygotes miscalled as homozygotes, and uncertainty of allele copy number in heterozygous

polyploids. Bayesian genotype calling can mitigate these issues, but previously has only been

implemented in software that requires a reference genome or uses priors that may be

inappropriate for the population. Here we present several novel Bayesian algorithms that

estimate genotype posterior probabilities, all of which are implemented in a new R package,

polyRAD. Appropriate priors can be specified for mapping populations, populations in Hardy-

Weinberg equilibrium, or structured populations, and in each case can be informed by genotypes

at linked markers. The polyRAD software imports read depth from several existing pipelines,

and outputs continuous or discrete numerical genotypes suitable for analyses such as genome-

wide association and genomic prediction.

Introduction

Approximately 70% of vascular plant species are recent polyploids, yet genomic

resources and bioinformatics tools for polyploids typically lag behind those for diploids (Moghe

and Shiu 2014; Renny-Byfield and Wendel 2014; Bourke *et al.* 2018). Reduced representation

DNA sequencing methods, such as genotyping-by-sequencing (GBS) and restriction site-

associated DNA sequencing (RAD-seq), have made high-density genotyping considerably more

accessible and affordable (Poland and Rife 2012; Davey *et al.* 2013). However, the two most

popular pipelines for processing GBS and RAD-seq data, Stacks (Catchen *et al.* 2013) and

TASSEL (Glaubitz *et al.* 2014), do not output polyploid genotypes. Though pipelines for

polyploids are available, each have limitations that prevent their general application. For

example, the UNEAK pipeline is designed for diploidized polyploids only (Lu *et al.* 2013).

HaploTag is specialized for self-fertilizing polyploids (Tinker *et al.* 2016). FreeBayes and

GATK can output polyploid genotypes, but require a reference genome (McKenna *et al.* 2010;

Garrison and Marth 2012). The software EBG imports read depth from other pipelines to

estimate auto- or allopolyploid genotypes (Blischak *et al.* 2018) but requires allele frequency

estimations from the parent species for allopolyploids. Thus, important staple crops such as

wheat, potato, sweet potato, yam, and plantain are underserved by existing genotyping software,

limiting our ability to perform marker-assisted selection, while yield increases from breeding are

not keeping pace with projected food demands (Ray *et al.* 2013).

We present a new R package, polyRAD, for genotype estimation from read depth in

polyploids and diploids. The software polyRAD is designed on the principle originally proposed

by Li (2011) that it is not necessary to call genotypes with complete certainty in order to make

useful inferences from sequencing data. Initially, SNP discovery is performed by other software

such as TASSEL (Glaubitz *et al.* 2014) or Stacks (Catchen *et al.* 2013), with or without a

reference genome, then allelic read depth is imported into polyRAD from those pipelines or the

read counting software TagDigger (Clark and Sacks 2016). In polyRAD, one or several ploidies

can be specified, including any level of auto- and/or allopolyploidy, allowing inheritance modes

to vary across the genome. Genotype probabilities are estimated by polyRAD under a Bayesian

framework, where priors are based on mapping population design, Hardy-Weinberg equilibrium

(HWE), or population structure, with or without linkage disequilibrium (LD). Multi-allelic loci

(haplotypes) are allowed, and are in fact encouraged because LD within the span of one RAD tag

is not informative for genotype imputation. In addition to exporting the most probable genotype

for each individual and locus, continuous numerical genotypes can be exported reflecting the

relative probabilities of all possible allele copy numbers, and can then be used for genome-wide

association or genomic prediction in software such as GAPIT (Lipka *et al.* 2012), FarmCPU (Liu

et al. 2016b), TASSEL (Bradbury *et al.* 2007), or rrBLUP (Endelman 2011). polyRAD is the

first Bayesian genotype caller to incorporate population structure and multiple inheritance

modes, as well as the first with an option for mapping populations. It is available at

https://github.com/lvclark/polyRAD and https://CRAN.R-project.org/package=polyRAD.

Methods

Overview

polyRAD implements Bayesian genotype estimation, similar to that proposed and implemented by several other groups (Li 2011; Nielsen *et al.* 2011; Garrison and Marth 2012; Korneliussen *et al.* 2014; Maruki and Lynch 2017; Blischak *et al.* 2018). In all polyRAD pipelines, genotype prior probabilities ($P(G_i)$) are specified from population parameters, and optionally from genotypes at linked markers (see Supplementary Methods). For a given individual and locus, consider every sequencing read to be a Bernoulli trial, where the read either matches a given allele (success) or some other allele (failure). The probability of success is:

Eqn. 1: $\pi_i = (1 - c) * \frac{i}{k} + c * p,$

where c is the cross-contamination rate, i is the allele copy number in the genotype, k is the ploidy, and p is the allele frequency in the population. For every possible allele copy number at a given locus and individual, the following equation is used to estimate the likelihood of the observed read depth using the binomial probability mass function:

Eqn. 2: $L(a, b|G_i) = \binom{a+b}{a} * \pi_i^a * (1 - \pi_i)^b,$

where a is the number of reads for a given allele at a given locus, b is the number of reads for other alleles at that locus, and G_i is the state in which a locus has i copies of a given allele.

From the priors and likelihoods, a posterior probability can then be estimated for each possible allele copy number for each individual and allele using Bayes' theorem (Shiryaev 2011):

Eqn. 3: $P(G_i|a, b) = \frac{L(a, b|G_i) * P(G_i)}{\sum_{i=0}^{k} L(a, b|G_i) * P(G_i)},$

where all terms are as previously described.

Bayesian genotype estimation allows correction of genotyping errors in diploids and

polyploids, i.e. when an individual is truly heterozygous but only one allele was sequenced, or

when an individual appears heterozygous due to sequencing error or contamination but is truly

homozygous. It also enables estimation of allele dosage in heterozygous polyploid genotypes.

Moreover, genotype posterior probabilities are more influenced by priors when read depth is

low, and by genotype likelihoods derived from allelic read depth when read depth is high. When

read depth is zero for a given individual and locus, genotype posterior probabilities are equal to

priors, and thus, somewhat counter-intuitively, the concept of missing data does not apply. It is

therefore not necessary to impute missing genotypes in a second step if the priors are sufficiently

informative.

For export to other software, as well as iteration within the polyRAD pipelines, a given

allele's weighted mean genotype (*wmg*) is a mean of the number of copies of that allele, with the

posterior genotype probabilities (Eqn. 3) serving as weights. Thus, for an individual and allele,

the weighted mean genotype (*wmg*) is calculated as:

Eqn. 4: $wmg = \sum_{i=0}^{k} P(G_i|a, b) * i,$

where all terms are as previously described. Additional details and equations for specification of

prior genotype probabilities and estimation of other parameters are provided in Supplementary

Materials. A flow chart of how this Bayesian genotypic estimation is implemented into

polyRAD is displayed in Fig. 1. In brief, for mapping populations, genotype priors are specified

based on parental genotypes and progeny allele frequencies, and all parameters are estimated

once. For diversity panels, genotype priors are adjusted and parameters re-estimated iteratively

until allele frequencies converge. Source code is available at

https://github.com/lvclark/polyRAD, archived at Zenodo (doi: 10.5281/zenodo.1143744).

Fig. 1. Overview of polyRAD algorithms for genotype estimation. Genotype posterior

probabilities are estimated iteratively until allele frequencies converge, except in the case of

mapping populations, where allele frequencies are only estimated once. Purple boxes indicate

inputs to the pipeline (read depth, contamination rate, and optionally, genomic positions of loci).

Blue boxes indicate estimated parameters (allele frequencies, genotype likelihoods and prior and

posterior probabilities, linkage between alleles, and probability of sampling each allele). Green

boxes indicate alternative methodologies for genotype prior probability estimation (mapping,

HWE, and population structure). Priors for the HWE and population structure models can be

adjusted for self-fertilization according to de Silva et al. (2005). Orange boxes indicate sample $\times$

allele matrices indicating approximate allele copy number. Dashed arrows indicate steps that

happen only once at the beginning or end of the pipeline, whereas solid arrows indicate iterative

steps. Circular arrows highlight cycles of iteration. Eqns. 1-4 are provided in the main

manuscript, and Eqns. 5-19 are provided in Supplemental Materials.

Example use

Executable examples are provided in the vignette and manual distributed with polyRAD.

Here we provide an additional brief example. Box 1 illustrates the use of polyRAD on a

diversity panel of a generic tetraploid species with a reference genome. Tools from the

Bioconductor package VariantAnnotation (Obenchain *et al.* 2014) are used by the polyRAD

function *VCF2RADdata* for import of a VCF file to the polyRAD-specific "RADdata" format.

SNP filtering criteria are specified with the *min.ind.with.reads* and *min.ind.with.minor.allele*

arguments, respectively. The *possiblePloidies* argument indicates that the inheritance mode

could be allotetraploid (*c(2,2)*) or autotetraploid (*4*). Any ploidy may be specified with

possiblePloidies, for example *8* for auto-octoploid, with the only limitation that all subgenomes

in an allopolyploid must have the same ploidy. By default, *VCF2RADdata* groups SNP alleles

into haplotypes that appear to have come from the same RAD tag, the size of which is specified

by *tagsize*, in basepairs. The function *IteratePopStructLD* is then used for genotype estimation,

taking both population structure and LD into account. The probabilistic principal components

analysis method from the Bioconductor package pcaMethods (Stacklies *et al.* 2007) is used

internally by *IteratePopStructLD* in order to estimate population structure. The *LDdist* argument

indicates the distance in basepairs within which to search for alleles at other loci that can help

predict copy number of a given allele. Once genotype posterior probabilities are estimated, other

parameters are cleared from memory using the *StripDown* function. Continuous numerical

genotypes are then formatted for GAPIT (Lipka *et al.* 2012) using the *ExportGAPIT* function.

Alternative functions are listed in Table 1. A very similar script could be used for a species

without a reference genome, with *IteratePopStruct* in place of *IteratePopStructLD*, and a

different import function for the appropriate non-reference pipeline.

```
library(polyRAD)

library(VariantAnnotation)

# prepare the VCF file for import

myvcf <- "somegenotypes.vcf"

myvcfbg <- bgzip(myvcf)

indexTabix(myvcfbz, format = "vcf")

# import VCF into a RADdata object

myRAD <- VCF2RADdata(myvcfbg,

                     tagsize = 64,

                     min.ind.with.reads = 300,

                     min.ind.with.minor.allele = 15,

                     possiblePloidies = list(c(2,2), 4))

# genotype estimation with pop. structure pipeline

myRAD <- IteratePopStructLD(myRAD, LDdist = 5e4)

# free up memory

myRAD <- StripDown(myRAD)

# export for GAPIT

myGM_GD <- ExportGAPIT(myRAD)
```

148   Box 1. Example R script using polyRAD. Read depth is imported from a VCF file, genotypes

149   are estimated using population structure and LD, and continuous numerical genotypes are

150   formatted for GAPIT.

152   Table 1. Overview of main polyRAD functions.

Import functions	
VCF2RADdata	Imports any VCF with an allelic read depth (AD) field, such as those exported by TASSEL-GBSv2 or GATK.
readTagDigger	Imports CSV file of read depth output by TagDigger.
readStacks	Reads catalog and matches files from Stacks.
readHMC	Reads files output by UNEAK.
Genotype estimation functions	
PipelineMapping2Parents	For mapping populations with any number of generations of backcrossing, intermating, and/or selfing.
IterateHWE	For diversity panels without population structure.[*]
IterateHWE_LD	For diversity panels with LD and without population structure.[*]
IteratePopStruct	For diversity panels with population structure.[*]
IteratePopStructLD	For diversity panels with population structure and LD.[*]
Export functions	
ExportGAPIT	Format genotypes for the *GD* and *GM* arguments of GAPIT or FarmCPU.
Export_rrBLUP_Amat	Format genotypes for the *A.mat* function in rrBLUP.
Export_rrBLUP_GWAS	Format genotypes for the *GWAS* function in rrBLUP.
Export_TASSEL_Numeric	Write file formatted for TASSEL with continuous numeric genotypes.
GetWeightedMeanGenotypes	Create a matrix of continuous numeric genotypes.
GetProbableGenotypes	Create a matrix of discrete genotypes, indicating the most probable genotype for each individual and allele.

153   [*]The rate of self-fertilization can be specified for self-compatible plant species.

Testing

To test the accuracy of polyRAD, we used datasets from three previously studied

populations: 1) RAD-seq data and GoldenGate SNP genotypes from a diversity panel (n = 565)

of the outcrossing, diploidized allotetraploid grass *Miscanthus sinensis* (Clark *et al.* 2014), 2)

RAD-seq data and GoldenGate SNP genotypes from a bi-parental F_1 mapping population (n =

275) of *M. sinensis* (Liu *et al.* 2016a), and 3) SNP array genotypes from a biparental F_1 mapping

population of autotetraploid potato (n = 238) (da Silva *et al.* 2017). Allelic read depth at

simulated RAD-seq markers was generated from the GoldenGate or SNP array genotypes, with

overall locus depth drawn from a gamma distribution to resemble depth of actual RAD-seq

markers (shape = 2 and scale = 5). The read depth for an individual genotype was also sampled

from a gamma distribution, with the shape equal to the locus depth divided by 10, and scale = 10.

The *M. sinensis* diversity panel included 395 GoldenGate markers, plus real RAD-seq data for

those same individuals across 3290 tag locations within 20 kb of any GoldenGate markers, called

with the TASSEL GBS v2 pipeline (Glaubitz *et al.* 2014) using the *M. sinensis* v7.1 reference

genome (DOE-JGI, http://phytozome.jgi.doe.gov/). Additionally, to test the effect of ploidy

within the *M. sinensis* diversity panel, tetraploidy was simulated by summing GoldenGate

genotypes and RAD-seq read depth of each individual with one of the ten individuals most

closely related to it, selected at random. The *M. sinensis* mapping population included 241

GoldenGate markers genotyped across 83 individuals, plus 3062 RAD-seq markers called with

the UNEAK pipeline (Lu *et al.* 2013) across those 83 individuals plus an additional 192

individuals. The potato mapping population included genotypes at 2538 markers. Additional

simulations using data from diversity panels of soybean (Song *et al.* 2015), apple (Chagné *et al.*

2012), and potato (Hamilton *et al.* 2011) are presented in Figs. S1, S2, and S3. In each

population, the simulated and real RAD-seq data was used for genotype calling with polyRAD

and EBG (Blischak *et al.* 2018) and missing genotypes from the EBG output were imputed with

LinkImpute (Money *et al.* 2015) and/or rrBLUP (Endelman 2011) as appropriate. To estimate

the accuracy of genotype calling and imputation, the root mean squared error (RMSE) was

calculated between numeric genotypes (ranging from zero to the ploidy) at each simulated RAD-

seq marker and at the GoldenGate or SNP array marker from which it was derived. Data and

scripts for analysis are available at https://doi.org/10.13012/B2IDB-9729830_V1.

Results and discussion

Accuracy of polyRAD

In the *M. sinensis* diversity panel, polyRAD showed improved genotype accuracy over

the HWE, disequilibrium, and GATK methods implemented in EBG, particularly at low read

depths (Fig. 2). Under the HWE model in polyRAD with discrete genotypes output, errors in

genotypes with more than zero reads were similar to those from the HWE and "diseq" models of

EBG in both diploid and tetraploid systems (Fig. 2A). However, when priors in polyRAD were

based on population structure, errors decreased, particularly in tetraploids and at low read depth

(Fig. 2A). In diploids and tetraploids respectively using the polyRAD population structure

model, error (RMSE) was reduced by 12.4% (SE 1.0%) and 25.3% (SE 0.6%) relative to the

GATK model, and by 8.2% (SE 1.0%) and 16.3% (SE 0.6%) relative to the EBG HWE model.

Given the known population structure in *M. sinensis* (Clark *et al.* 2014), it is unsurprising that a

population structure-aware genotyping method would be more accurate than one based on HWE.

For genotypes with zero reads, imputation was most accurate when it accounted for population

structure, using either polyRAD or rrBLUP (Fig. 2B). Although modeling LD did not improve

accuracy in *M. sinensis* (Fig. 2), likely due to low LD as a result of outcrossing (Slavov *et al.*

2014), modeling LD did improve accuracy in wild soybean, apple, and a simulated inbreeding

allohexaploid (Figs. S1, S2, and S3, and Supporting Results).

Fig. 2. Genotyping error of polyRAD, EBG, LinkImpute, and rrBLUP in a diversity panel of

565 diploid *Miscanthus sinensis*. The benefits of incorporating population structure into the

genotyping model and using continuous rather than discrete genotypes are illustrated. Genotypes

were coded on a scale of 0 to 2 for diploids and 0 to 4 for tetraploids. Root mean squared error

(RMSE) was calculated between actual genotypes and genotypes ascertained from simulated

RAD-seq reads at 395 SNP markers (lower RMSE = higher accuracy). To simulate tetraploid

genotypes, genotypes were combined across closely-related individuals. (A) RMSE calculated

using only genotypes with more than zero reads. Median read depth is indicated, including

genotypes with zero reads. Each point represents one SNP. Smoothing curves are included for

each genotyping method and genotype type. (B) RMSE calculated using only genotypes with

zero reads, by genotyping or imputation method and genotype type. LinkImpute works for

diploids only and therefore was not tested on the tetraploid population.

218   In diploid *M. sinensis* and tetraploid potato F1 mapping populations, polyRAD

219 outperformed the GATK method, particularly when linked markers were used for informing the

220 priors in polyRAD (Fig. 3A). In diploids and tetraploids respectively using genotypes with non-

221 zero read depth, error (RMSE) was reduced by 38.2% (SE 2.4%) and 51.7% (SE 0.5%) using the

222 polyRAD linkage model with respect to the GATK model. For imputation, polyRAD using the

223 linkage model performed similarly to LinkImpute and rrBLUP (Fig. 3B). Although only F1

224 populations are presented here, many other population types are supported in polyRAD.

Fig. 3. Genotyping error of polyRAD, EBG, LinkImpute, and rrBLUP in F1 mapping

populations of diploid *Miscanthus sinensis* and tetraploid potato. The benefits of incorporating

population design and linkage into the genotyping model, as well as using continuous rather than

discrete genotypes, are illustrated. Genotypes were coded on a scale of 0 to 2 for diploids and 0

to 4 for tetraploids. Root mean squared error (RMSE) was calculated between actual genotypes

and genotypes ascertained from simulated RAD-seq reads (lower RMSE = higher accuracy).

The diploid population included 241 markers across 83 individuals, and the tetraploid population

included 2538 markers across 238 individuals. (A) RMSE calculated using only genotypes with

more than zero reads. Median read depth is indicated, including genotypes with zero reads.

Each point represents one SNP. Smoothing curves are included for each genotyping method and

genotype type. (B) RMSE calculated using only genotypes with zero reads, by genotyping or

imputation method and genotype type. Discrete genotypes from the polyRAD model without

linkage were not available in the diploid population for genotypes with zero reads because most

markers had even priors across two genotypes. LinkImpute works for diploids only and

therefore was not tested on the tetraploid population.

Genotyping error was also reduced 10-15% in most cases by exporting genotypes as

continuous numerical variables (weighted mean genotypes), rather than discrete values (Figs. 2

and 3). For example, in a diploid, a true heterozygote (numeric value of 1) with reads only for

the reference allele might erroneously be called as zero (homozygous for the reference allele) if

only the most probable genotype is exported. However, the genotype could be called 0.4 if

continuous genotypes are allowed, indicating that there is a 60% chance of it being a

homozygote and 40% chance of it being a heterozygote, and thereby reducing the error from 1.0

to 0.6. Similarly in polyploids, continuous numerical genotypes can correct for errors in allele

copy number estimation of heterozygotes.

Downstream applications and implications for sequencing strategies

The genotyping methods implemented in polyRAD will have the most benefit for marker

analysis where 1) the accuracy of individual genotypes is important, and 2) genotypes can be

treated as continuous rather than discrete variables. Analyses that seek to quantify marker-trait

associations in a population of individuals, such as genome-wide association studies, quantitative

trait locus mapping, and genomic prediction methods involving variable selection, could

especially benefit. By reducing genotyping error, polyRAD will increase the power of these

methods to detect true associations. Analyses that will benefit less from polyRAD genotyping

are those where an average is taken across many genotypes in order to estimate a statistic, such

as allele frequencies in a population or overall relatedness of individuals (including kinship-

based methods of genomic prediction), because genotyping errors generally are not biased

towards one allele or the other and tend to balance out over many individuals and loci (Buerkle

and Gompert 2013; Dodds *et al.* 2015).

The advantages of polyRAD for accurate genotyping at low sequence read depth alter the

economics of sequence-based genotyping, enabling researchers to purchase fewer sequencing

lanes, multiplex more samples per lane, and/or retain more markers during filtering. In

particular, for protocols using restriction enzymes where read depth varies considerably from

locus to locus depending on fragment size (Beissinger *et al.* 2013; Davey *et al.* 2013; Andrews *et*

al. 2016), there are diminishing returns on increasing the per-sample read depth, because some

loci receive far more reads than are needed for accurate genotyping while other loci remain poor

quality. Using population structure and linkage between loci, polyRAD uses information from

high-depth markers to improve genotyping accuracy of low-depth markers, helping to maximize

the useful information that is obtained from sequencing data. This advance is expected to

improve breeding efficiency and economics.

Acknowlegements

277   This material is based upon work supported by the National Science Foundation under

278   Grant No. 1661490. The authors thank Marcus Olatoye, Per McCord, and Joyce Njuguna for

279   testing the polyRAD software.

References

Andrews, K. R., J. M. Good, M. R. Miller, G. Luikart, and P. A. Hohenlohe, 2016 Harnessing

the power of RADseq for ecological and evolutionary genomics. Nat. Rev. Genet. 17: 81–

92.

Beissinger, T. M., C. N. Hirsch, R. S. Sekhon, J. M. Foerster, J. M. Johnson *et al.*, 2013 Marker

density and read depth for genotyping populations using genotyping-by-sequencing.

Genetics 193: 1073–1081.

Blischak, P. D., L. S. Kubatko, and A. D. Wolfe, 2018 SNP genotyping and parameter estimation

in polyploids using low-coverage sequencing data. Bioinformatics 34: 407–415.

Bourke, P. M., R. E. Voorrips, R. G. F. Visser, and C. Maliepaard, 2018 Tools for Genetic

Studies in Experimental Populations of Polyploids. Front. Plant Sci. 9: 513.

Bradbury, P. J., Z. Zhang, D. E. Kroon, T. M. Casstevens, Y. Ramdoss *et al.*, 2007 TASSEL:

software for association mapping of complex traits in diverse samples. Bioinformatics 23:

2633–2635.

Buerkle, C. A., and Z. Gompert, 2013 Population genomics based on low coverage sequencing:

How low should we go? Mol. Ecol. 22: 3028–3035.

Catchen, J., P. A. Hohenlohe, S. Bassham, A. Amores, and W. A. Cresko, 2013 Stacks: an

analysis tool set for population genomics. Mol. Ecol. 22: 3124–40.

Chagné, D., R. N. Crowhurst, M. Troggio, M. W. Davey, B. Gilmore *et al.*, 2012 Genome-Wide

SNP Detection, Validation, and Development of an 8K SNP Array for Apple (M.

Bendahmane, Ed.). PLoS One 7: e31745.

Clark, L. V, J. E. Brummer, K. Głowacka, M. C. Hall, K. Heo *et al.*, 2014 A footprint of past

climate change on the diversity and population structure of *Miscanthus sinensis*. Ann. Bot.

114: 97–107.

Clark, L. V, and E. J. Sacks, 2016 TagDigger: user-friendly extraction of read counts from GBS

and RAD-seq data. Source Code Biol. Med. 11: 11.

Davey, J. W., T. Cezard, P. Fuentes-Utrilla, C. Eland, K. Gharbi *et al.*, 2013 Special features of

RAD Sequencing data: implications for genotyping. Mol. Ecol. 22: 3151–3164.

Dodds, K. G., J. C. McEwan, R. Brauning, R. M. Anderson, T. C. van Stijn *et al.*, 2015

Construction of relatedness matrices using genotyping-by-sequencing data. BMC Genomics

16: 1047.

Endelman, J. B., 2011 Ridge regression and other kernels for genomic selection with R package

rrBLUP. Plant Genome J. 4: 250–255.

Garrison, E., and G. Marth, 2012 Haplotype-based variant detection from short-read sequencing.

arXiv 1207.3907.

Glaubitz, J. C., T. M. Casstevens, F. Lu, J. Harriman, R. J. Elshire *et al.*, 2014 TASSEL-GBS: A

High Capacity Genotyping by Sequencing Analysis Pipeline. PLoS One 9: e90346.

Hamilton, J. P., C. N. Hansey, B. R. Whitty, K. Stoffel, A. N. Massa *et al.*, 2011 Single

nucleotide polymorphism discovery in elite north American potato germplasm. BMC

Genomics 12:.

Korneliussen, T. S., A. Albrechtsen, and R. Nielsen, 2014 ANGSD: Analysis of Next Generation

Sequencing Data. BMC Bioinformatics 15: 356.

Li, H., 2011 A statistical framework for SNP calling, mutation discovery, association mapping

and population genetical parameter estimation from sequencing data. Bioinformatics 27:

2987–2993.

Lipka, A. E., F. Tian, Q. Wang, J. Peiffer, M. Li *et al.*, 2012 GAPIT: genome association and

prediction integrated tool. Bioinformatics 28: 2397–2399.

Liu, S., L. V. Clark, K. Swaminathan, J. M. Gifford, J. A. Juvik *et al.*, 2016a High density

genetic map of *Miscanthus sinensis* reveals inheritance of zebra stripe. GCB Bioenergy 8:

616–630.

Liu, X., M. Huang, B. Fan, E. S. Buckler, and Z. Zhang, 2016b Iterative Usage of Fixed and

Random Effect Models for Powerful and Efficient Genome-Wide Association Studies.

PLOS Genet. 12: e1005767.

Lu, F., A. E. Lipka, J. Glaubitz, R. Elshire, J. H. Cherney *et al.*, 2013 Switchgrass genomic

diversity, ploidy, and evolution: novel insights from a network-based SNP discovery

protocol. PLoS Genet. 9: e1003215.

Maruki, T., and M. Lynch, 2017 Genotype Calling from Population-Genomic Sequencing Data.

G3 7: 1393–1404.

McKenna, A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis *et al.*, 2010 The Genome

Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing

data. Genome Res. 20: 1297–1303.

Moghe, G. D., and S. H. Shiu, 2014 The causes and molecular consequences of polyploidy in

flowering plants. Ann. N. Y. Acad. Sci. 1320: 16–34.

Money, D., K. Gardner, Z. Migicovsky, H. Schwaninger, G.-Y. Zhong *et al.*, 2015 LinkImpute:

Fast and Accurate Genotype Imputation for Nonmodel Organisms. G3 5: 2383–90.

Nielsen, R., J. S. Paul, A. Albrechtsen, and Y. S. Song, 2011 Genotype and SNP calling from

next-generation sequencing data. Nat. Rev. Genet. 12: 443–451.

Obenchain, V., M. Lawrence, V. Carey, S. Gogarten, P. Shannon *et al.*, 2014 VariantAnnotation:

A Bioconductor package for exploration and annotation of genetic variants. Bioinformatics

30: 2076–2078.

Poland, J. A., and T. W. Rife, 2012 Genotyping-by-sequencing for plant breeding and genetics.

Plant Genome J. 5: 92–102.

Ray, D. K., N. D. Mueller, P. C. West, and J. A. Foley, 2013 Yield Trends Are Insufficient to

Double Global Crop Production by 2050. PLoS One 8: e66428.

Renny-Byfield, S., and J. F. Wendel, 2014 Doubling down on genomes: Polyploidy and crop

plants. Am. J. Bot. 101: 1711–1725.

Shiryaev, A. N., 2011 Bayes formula. Encycl. Math. Available at:

https://www.encyclopediaofmath.org//index.php?title=Bayes_formula&oldid=16075.

da Silva, W., J. Ingram, C. A. Hackett, J. J. Coombs, D. Douches *et al.*, 2017 Mapping Loci That

Control Tuber and Foliar Symptoms Caused by PVY in Autotetraploid Potato (*Solanum*

tuberosum L.). G3 7: 3587–3595.

De Silva, H. N., A. J. Hall, E. Rikkerink, M. A. McNeilage, and L. G. Fraser, 2005 Estimation of

allele frequencies in polyploids under certain patterns of inheritance. Heredity 95: 327–334.

Slavov, G. T., R. Nipper, P. Robson, K. Farrar, G. G. Allison *et al.*, 2014 Genome-wide

association studies and prediction of 17 traits related to phenology, biomass and cell wall

composition in the energy grass *Miscanthus sinensis*. New Phytol. 201: 1227–1239.

Song, Q., D. L. Hyten, G. Jia, C. V. Quigley, E. W. Fickus *et al.*, 2015 Fingerprinting Soybean

Germplasm and Its Utility in Genomic Research. G3 5: 1999–2006.

Stacklies, W., H. Redestig, M. Scholz, D. Walther, and J. Selbig, 2007 pcaMethods - A

bioconductor package providing PCA methods for incomplete data. Bioinformatics 23:

1164–1167.

Tinker, N. A., W. A. Bekele, and J. Hattori, 2016 Haplotag: Software for Haplotype-Based

Genotyping-by-Sequencing Analysis. G3 6: 857–863.