Manuscript title: Molecular Diversity of Intrinsically Photosensitive Ganglion Cells

Authors:

Daniel Berg[1,2], Katherine Kartheiser[2], Megan Leyrer[2], Alexandra Saali[2], David Berson[2]

1. Molecular Biology Program

2. Department of Neuroscience

Brown University, Providence, RI 02912

Correspondence to: David Berson (david_berson@brown.edu)

WORD COUNT: **Abstract** (148 words); **Intro** (784 words); **Results** (5159); **Discussion** (2657)

14 <u>Abstract</u>

Intrinsically photosensitive retinal ganglion cells (ipRGCs) are rare mammalian photoreceptors

essential for non-image-forming vision functions, such as circadian photoentrainment and the

pupillary light reflex. They comprise multiple subtypes distinguishable by morphology, physiology,

projections, and levels of expression of melanopsin (Opn4), their photopigment. The molecular

programs that differentiate ipRGCs from other ganglion cells and ipRGC subtypes from one

another remain elusive. Here, we present comprehensive gene expression profiles of early

postnatal and adult mouse ipRGCs purified from two lines of reporter mice marking different sets

of ipRGC subtypes. We find dozens of novel genes highly enriched in ipRGCs. We reveal that

Rasgrp1 and Tbx20 are selectively expressed in subsets of ipRGCs, though these molecularly

defined groups imperfectly match established ipRGC subtypes. We demonstrate that the ipRGCs

regulating circadian photoentrainment are unexpectedly diverse at the molecular level. Our

findings reveal unexpected complexity in gene expression patterns across mammalian ipRGC

subtypes.

<u>Introduction</u>

Many unique attributes distinguish intrinsically photosensitive RGCs from conventional

RGCs. Only ipRGCs express the blue-light sensitive photopigment melanopsin (OPN4), which

renders them autonomously light-sensitive. They violate the usual stratification rule in which ON-

type RGCs deploy their dendrites only in the inner (proximal) half of the inner plexiform layer; their

inputs from ON bipolar cells are atypical (Dumitrescu et al., 2009; Hoshi et al., 2009; Kim et al.,

2010). Whereas most RGCs direct their entire output through the optic nerve, some ipRGCs

modulate intraretinal processing, through dopaminergic (Zhang et al., 2008; Xue et al., 2011) and

other amacrine cells (Reifler et al., 2015; Sabbah et al., 2017) and spontaneous retinal waves

during the early postnatal period (Renna et al., 2011). Additionally, ipRGCs appear more resistant

than RGCs overall to various sorts of insults, including intraocular hypertension, optic nerve injury,

glutamate-induced excitotoxicity, and glaucoma (Cui et al., 2015). Functionally, ipRGCs are

unique among RGCs in their ability to encode overall light intensity for extended periods (Wong,

2012). This tonic luminance signal is transmitted to specific brain targets for a variety of functions.

Projections to the hypothalamus mediate photoentrainment of circadian rhythms, while those to

the midbrain to drive pupillary constriction. These outputs derive from molecularly distinct ipRGC

subtypes.

The distinctive structural and functional properties of ipRGCs must ultimately be traceable

to different patterns of gene expression. However, there is very little information on what these

differences might be. For example, the basic molecular framework of the melanopsin

phototransduction cascade, a major defining feature of ipRGCs, has only begun to be identified

(Hughes et al., 2012) and the precise phototransduction mechanisms remain poorly

characterized. More generally, very little is known about the developmental molecular

mechanisms that direct certain immature RGCs to an ipRGC fate, or that maintain the distinctive

features of ipRGCs throughout maturity. Thus, there is ample motivation for comparing the

transcriptional profiles of ipRGCs as compared to those of conventional RGCs.

The ipRGCs consist of at least six anatomically distinct retinal subtypes, termed M1-M6.

These which differ in their level of melanopsin expression, visual response properties, dendritic

stratification, axonal projections, and contributions to light-modulated behavioral responses

(Schmidt et al., 2011). Little is known about the gene regulatory programs that differentiate and

maintain the specialized properties of individual ipRGC subtypes. M1 ipRGCs have been further

subdivided based on their expression of the transcription factor Pou4f2 (Brn3b) (Chen et al., 2011;

Jain et al., 2012). M1 ipRGCs that innervate the suprachiasmatic nucleus (SCN) do not express

Brn3b, while those that project to other M1-cell targets, such as the olivary pretectal nucleus

(OPN), do express this transcription factor (Chen et al., 2011). Indeed, ablation of Brn3b-positive

ipRGCs severely impairs the pupillary light reflex, but leaves circadian photoentrainment intact

(Chen et al., 2011). There is surely additional molecular diversity among ipRGCs, both within and

between established ipRGC subtypes, and this further motivates the present study.

Previous attempts to develop a "molecular parts list" for ipRGCs through gene expression

profiling of adult ipRGCs have been limited by the extreme heterogeneity of retinal tissue, and the

fragility of mature retinal neurons, and the minuscule amount of genetic material from ipRGCs,

which comprise far fewer than 1% of all retinal neurons (Lobo et al., 2006; Heiman et al., 2008;

Sanes and Masland, 2015). Some progress has been made by purifying ipRGCs from

enzymatically dissociated retinas using either anti-melanopsin immuno-panning or fluorescence-

activated cell sorting (FACS) of genetically-labeled fluorescent ipRGCs. However, prior efforts

have been limited by low yield and inclusion of contaminating cell populations such as rods

(Hartwick et al., 2007; Peirson et al., 2007; Siegert et al., 2012).

Here we conducted a thorough unbiased transcriptomic analysis of ipRGCs by purifying

GFP-tagged ipRGCs through a combination of FACS and immunoaffinity and comparing this the

transcriptional profile of GFP-negative RGCs. We did this in two different mouse lines, marking

partially overlapping subsets of ipRGCs. One was a BAC transgenic reporter (*Opn4-GFP*), which

fluorescently labels M1-M3 ipRGCs (see Methods) and the other was a *Opn4-Cre;Z/EG* reporter

system, which labels all ipRGC subtypes, M1-M6 (Schmidt et al., 2008; Ecker et al., 2010;

Quattrochi et al., 2018; Stabio et al., 2018). The specificity and purity achieved by our approach

is validated by the substantial enrichment in the ipRGC samples of known molecular markers of

ipRGCs and by the fact that transcripts selectively expressed in potential contaminating cell types

are generally absent. We identified more than 75 new gene candidates expressed much more

highly in adult ipRGCs than in other RGCs. We validate two of the new molecular markers at the

protein level: the Ras GEF Rasgrp1 and the T-box transcription factor Tbx20 and relate these to

established ipRGC subtypes and patterns of central projection.

<u>Results</u>

We enzymatically dissociated retinas from melanopsin-reporter mice, selected for RGCs

by anti-Thy1 immunoaffinity, and sorted these into presumptive ipRGC and conventional

ganglion-cell (cRGC) pools by FACS based on the fluorescent labeling of ipRGCs (see Methods,

Figure 1A). We then compared gene expression in these ipRGC-enriched and cRGC-enriched

cell samples to identify genes differentially expressed in ipRGCs as compared to other ganglion

cells. We used two strains of melanopsin-reporter mice (see Methods). One of these was a BAC

transgenic mouse (here termed *Opn4-GFP*) in which retinal GFP expression is apparently

restricted to ipRGCs of subtypes M1, M2 and M3 (see Methods; personal communication R.

Maloney and L. Quattrochi). This is presumably because expression of the reporter is coupled to

expression of melanopsin, which is highest in this subset of ipRGCs. The other melanopsin

reporter mouse was obtained by crossing a knock-in mouse in which *Cre* replaces *Opn4*

(Opn4$^{cre/cre}$) with a Cre reporter strain (Z/EG). The resulting Cre-driven labeling of melanopsin-

expressing cells is more sensitive than in the other strain of reporter mice, and labels all known

types of ipRGCs, M1-M6, while labeling few if any conventional RGCs (cRGCs) (Ecker et al.,

2010; Estevez et al., 2012; Quattrochi et al., 2018; Stabio et al., 2018) (Figure 1A; see Methods).

Cell composition and purity of isolated ipRGCs and conventional RGCs

The relationship among the transcriptional profiles of ipRGC and cRGC samples across

replicates are illustrated in the multidimensional scaling (MDS) plots of Fig. 1B. These show the

relationship between all pairs of samples (one of ipRGCs, the other of cRGCs) based on a count-

specific pairwise distance measure (Anders et al., 2013) (Figure 1B). These sample pairs were

clearly separated along the first dimension, indicating pronounced differences in overall gene-

expression patterns between ipRGC and cRGC samples. Samples of ipRGCs and cRGCs derived

from the same retina and processed in parallel tended to be closely spaced along the second

dimension, indicating greater similarity within than across replicates. This may reflect slight

differences in overall genetic makeup of mice contributing to each pool, since both strains used

were on a mixed genetic background, or to slight technical differences in the acquisition and

processing of RNA from one run to the next.

Transcriptional data offer broad internal evidence for the efficacy of purification of cell

samples. As expected, the *Opn4* (melanopsin) gene was among the genes much more highly

expressed in ipRGCs than in cRGCs. For example, *Opn4* was enriched 40-fold in adult ipRGCs

purified from *Opn4-GFP* mice, and this was highly significant, at $q<1\times10^{-55}$ false discovery rate

(FDR). Though *Opn4* expression was detected in cRGCs at modest levels (Figure S1), this was

expected, because some ipRGCs lack GFP expression in both melanopsin reporter lines (Opn4-

GFP and Opn4-cre/GFP), and these would have been pooled with cRGCs during the FACS

procedure.

Transcripts of other genes known to be expressed in ipRGCs were also enriched in the

ipRGC pool relative to the cRGC pool (FDR < 0.05, significantly expressed in ipRGC samples,

and absent or weakly expressed in cRGC samples). Among these genes were *Adcyap1* (pituitary

adenylate-cyclase activating polypeptide; PACAP), *Tbr2*, (*Eomesodermin*), *Trpc7* and, to a lesser

extent, *Trpc6* (Hannibal et al., 2004; Xue et al., 2011; Sand et al., 2012; Mao et al., 2014; Sweeney

et al., 2014). Taken together, these results demonstrate that mRNA isolated from purified ipRGC

samples were enriched as expected for transcripts for genes that are known to be differentially

expressed in ipRGCs.

In the purified ipRGC samples, we found little or no evidence of contamination by

transcripts from other retinal cell types. For example, transcript levels were very low for rod and

cone opsins, for the amacrine-specific marker ChAT, for several bipolar markers (Otx2; Vsx2;

Grm6; Trpm1), and for markers of astrocytes, microglial and vascular cells (Figure S1). Several

transcripts suitable for assessing potential contamination from Müller glia (*Glul, Vim)* were present

at surprisingly high levels in the purified ipRGC samples, suggesting that these glial cells may

contaminate the transcriptional picture to some degree.

In general, the cRGC samples were relatively less pure than the ipRGCs samples by this

measure. A particularly informative transcript for assessing such contamination is that for the

rhodopsin gene (*Rho*), because rods are by far the most common neuronal type in the mouse

retina and express *Rho at* very high levels. Rhodopsin transcripts were significantly (150-fold)

more abundant in the cRGC samples than in ipRGC samples (Figure Supplement 1), whether

isolated from *Opn4-GFP* or *Opn4-Cre/GFP* adult reporter mice (FDR < 6 x 10^{-9}). Evidently, the

second isolation step in which GFP+ positive cells were isolated by FACS from the purified RGC

pool was a key factor in the greater purity of the ipRGC sample. Similarly, transcripts associated

with bipolar cells and Müller glial cells were generally more abundant in cRGC than ipRGC

samples. For example, the cRGCs had relatively high expression of the known Müller glia markers

Glul, *Apoe*, *Aqp4*, and *Vim*, generally higher than in the ipRGC pool (Figure S1). Contamination

of adult cRGC samples by other cell types may explain why most RGC markers, such as *Rbpms*

and *Sncg* (Soto et al., 2008; Rodriguez et al., 2014), were less abundant in the cRGC cell pool

than in the ipRGC pool. However, the data suggest that contamination in the cRGC pool was not

uniform across retinal cell types. Amacrine-specific transcripts were no more abundant overall in

cRGCs than in ipRGCs, and microglial and vascular markers were essentially absent, as in

ipRGCs.

In immature mice (P5; *Opn4-GFP*), contamination of cRGC samples by non-RGC

transcripts appeared more modest than in adults. The major sources of contamination (rods and

Müller glia) are still being born and undergoing early-stage differentiation at this age, and this

would presumably depress the abundance of their cell-type-specific transcripts (Young, 1985;

Morrow et al., 1998; Matsushima et al., 2011).

To summarize, this analysis suggests that all samples were relatively free of

contamination by most other retinal cell types, and that the ipRGC samples were particular pure.

Contamination of the cRGC samples appears to derive mainly from Müller cells and strongly

expressed genes in rods. Though this must be factored into the analysis, our primary focus was

on genes more strongly expressed in ipRGCs than in cRGCs, and this difference seems unlikely

to be affected by the modest contamination of the cRGC pool.

Genes differentially expressed in ipRGCs

Comparing the abundance of transcripts in the ipRGC and cRGC pools, we identified over

75 genes that were differentially elevated expression in ipRGCs (as marked by one or both

melanopsin-reporter lines) relative to cRGCs. Briefly, identification of differentially expressed

genes in ipRGCs relied on the following stringent criteria: 1) low false discovery rate with high

fold-change, 2) corroboration of differential expression across both ipRGC reporters, and 3)

absence of gene expression in cRGC samples (see Methods). The identified differentially

expressed genes are diverse, and most have not been previously identified as ipRGC-enriched

(Figure 2; see Methods). Here, we survey some of these genes, grouped by their functional

features (Figure 2).

1. *Transcription factors*

Transcription factors, by regulating other genes, help to generate and maintain ipRGC

identity. We noted above that the T-box transcription factor *Tbr2* was much more strongly

expressed in adult ipRGCs than in cRGCs, as expected (Sweeney et al., 2014). *Tbr2* is best

known for its key role in early retinal development. Its expression in adult retina is far more

restricted, but it remains expressed in the majority of ipRGCs. A second T-box transcription factor,

Tbx20, was similarly enriched (Figure 2). *Tbx20* has not been previously linked to adult retinal

function, but we will show that it too is quite selectively expressed in ipRGCs. Additionally, four

other transcription factors, *Irx6*, *Dmrtb1*, *Nr4a3*, and *Pou6f2,* were differentially expressed in adult

ipRGCs. Most of these genes serve as broad lineage determinants in early retinal development

(Zhou et al., 1996; Star et al., 2012). Other highly expressed genes included the neuron-derived

orphan receptor 1 *Nor1* (*Nr4a3*), which codes for a nuclear receptor, and *Elavl2* gene, which

codes for a RNA-binding protein important for mRNA metabolism and neuronal differentiation

(Fornaro et al., 2007; Hinman and Lou, 2008). Pathway analysis (DAVID) of differentially

expressed genes in ipRGCs suggested specialization in heparan sulfate biosynthesis, including

Hs3st4, *Hs3st2*, *Hs6st2*, *Ndst4*, and *Gpc5* (Figure 2).

2. *Synaptic transmission*

Some of the genes differentially expressed in ipRGCs have known roles in regulating

synaptic function, especially at presynaptic sites. Among others, these genes include *Sh3gl3*,

Entpd1, *Rab3b*, *Baiap3*, *Chl1*, *Sh3gl3*, and *Adra2a* (Figure 2).

3. *Growth factors and neuropeptides*

Multiple growth factors and neuropeptides were also differentially expressed in adult ipRGCs.

These include *Bmp7*, *Fgf1*, *Gal*, *Gdf6*, *Grem1*, *Nmb* and *Nppb*) (Figure 2).

4. ***Receptors and channels***

Multiple genes encoding diverse surface receptors were differentially expressed in ipRGCs

(Figure 2—figure supplement 2). For example, expression data suggest that ionotropic nicotinic

acetylcholine receptors in ipRGCs may be composed of α3, α4, α6, β2 and β3 subunits (Figure

2—figure supplement 2), although the α3 and α4 transcripts were borderline for differential

expression in ipRGCs. In agreement with previous studies, we found that ipRGCs expressed the

Drd1 dopamine receptor, but had low levels of Drd2 expression (Van Hook et al., 2012). Several

serotonin receptor genes (Htr1b, Htr1d, and Htr5a) were modestly enriched in ipRGCs. The

ipRGCs were also found to express many glutamate receptors subunits, but only one of these -

the NMDA receptor subunit 3A (GRIN3A) - was differentially expressed relative to other adult

RGCs. The mu opioid receptor gene *Oprm1* is differentially expressed in ipRGCs; it could regulate

their light responses interacting with L-type calcium channels, which carry the majority of light-

evoked inward calcium current in ipRGCs (Moises et al., 1994; Diaz et al., 1995; Doğrul et al.,

2001; Hartwick et al., 2007). Our data appear at odds with earlier reports that M1 and M4 ipRGCs

express the melatonin receptor genes Mtnr1a and Mtnr1b (Sengupta et al., 2011; Pack et al.,

2015; Sheng et al., 2015). Additionally, *Kcnh1*, also known as ether-a-go-go (*Eag1*), was

differentially expressed in ipRGCs (Figure 2). Kcnh1 is a voltage-gated K+ channel that has been

shown to be crucial for the generation of dark current in the inner segment of rods (Frings et al.,

1998), but may normally regulate other neuronal functions in ipRGCs (Martin et al., 2008).

5. ***Cell adhesion***

Genes encoding for several cell adhesion molecules were differentially expressed in

ipRGCs (Figure 2—figure supplement 3B). For example, the cell adhesion molecule *DscamL1*

was relatively low in ipRGCs during postnatal development, but the closely related genes

encoding the immunoglobulin superfamily (IgSF) adhesion molecules *Sidekick-1* and *Sidekick-2*

were enriched in developing ipRGCs. *Unc5a* and *Unc5d* were significantly differentially expressed

both in early postnatal and adult ipRGCs. In contrast, expression of *Unc5b* and *Unc5C* in ipRGCs

was low relative to that in cRGCs. As suggested previously, expression of the repulsive ligand

Sema6a was significantly lower in ipRGCs than cRGCs during postnatal development (Matsuoka

et al., 2011). However, its receptor *PlxnA4* was enriched in P5 ipRGCs. Another semaphorin,

Sema5a, was also significantly enriched in developing ipRGCs. Other differentially expressed

cell-adhesion molecules *Salm5* (*Lrfn5*), *Clstn2*, *Thbs1*, *Lrrtm2*, *Pcdh19*, *Ptprm*, and *Lrrc4c (Ngl1)*

could play significant roles in the formation of ipRGC synapses (Burden-Gulley and Brady-Kalnay,

1999; Lin et al., 2003; de Wit et al., 2009; Xu et al., 2010; Lipina et al., 2016; Pederick et al., 2016;

Lin et al., 2018). The cell surface glycoprotein *Mdga1* was also differentially expressed in

developing ipRGCs, and is known to influence the formation and maintenance of inhibitory

synapses (Pettem et al., 2013).

6.   ***Tolerance to stress***

There is increasing evidence that ipRGCs are resistant to stress and able to survive under

circumstances that are fatal for other retinal neurons (Li et al., 2008; de Sevilla Müller et al., 2014;

Cui et al., 2015; Duan et al., 2015). The harsh dissociation and FACS processing has the potential

of generating stress-induced gene expression changes (Figure 1 and 2, see methods). We

attempted to identify potential survival molecular programs that are specific to ipRGCs compared

to generic RGCs. The genes *Adcyap1* (PACAP), *Igf1*, and *Spp1* (*osteopontin*), all of which have

previously described roles in promoting ipRGC survival (Atlasz et al., 2010; Duan et al., 2015)

were differentially expressed in ipRGCs. We also identified a number of genes related to glial

function differentially expressed in ipRGCs, including *Gldn*, *Cntn2*, *Lama4*, and *Astn2*, and *Thbs1*

(Figure 2).

7.   ***Phototransduction***

Photoactivation of melanopsin photopigment typically triggers a phosphoinositide

signaling cascade resembling that in rhabdomeric (invertebrate) photoreceptors, involving G

proteins in the Gq family, phospholipase C, and canonical TRP channels. In ipRGCs, the

phototransduction cascade typically signals through Gq-family proteins and phospholipase C beta

4 (PLCB4) to open canonical TRP channels (Trpc7 and Trpc6) (Graham et al., 2008; Xue et al.,

2011; Hu et al., 2013; Emanuel and Do, 2015; Emanuel et al., 2017) (Figure 2—figure supplement

4A). We determined that the genes in this signaling cascade (*Opn4*, *Trpc7*, *Trpc6*, *Plcb4*, and

several Gq genes) were expressed at relatively high levels in all three ipRGC pools (i.e., selective-

postnatal; selective-adult; or pan-subtype adult). Moreover, two key genes - *Opn4* and *Trpc7* -

were more highly expressed in ipRGCs than in cRGCs in all three ipRGC pools. *Trpc6* was also

significantly overexpressed in ipRGCs in younger animals, with a trend in this direction also in

adult ipRGCs, but *Trpc7* was expressed at much higher levels than *Trpc6*. Plcb4 appears

essential for melanopsin phototransduction in some cells, and it was expressed at much higher

levels than Plcb1, 2 or 3. However, except in young mice, it was not more highly expressed in

ipRGCs than cRGCs.

Only recently has the precise identity of the Gα subunits combination in ipRGCs been

identified as redundantly expressing and signaling through the Gnaq, Gna11, or Gna14 subunits

(Hughes et al., 2015). Our studies suggest a similar expression pattern, including a lack of Gna15

expression (Figure 2—figure supplement 4A). Further, we determined that Gna14 was

differentially expressed in our P5 ipRGC samples, but it did not reach a statistical significant

difference in adult Opn4-GFP ipRGCs. Gnaq appears to be among the highest expressing Gq/11

subunits in our study, in contrast to the lack of Gnaq gene expression detected by Siegert and

colleagues (2012). To date, the G$\beta\gamma$ complex remains unknown. Our studies determined that the

beta subunit Gnb1 is by far the highest expressing, having a 15-fold higher expression than the

other subunits Gnb2, Gnb4, or Gnb5; while Gnb3 showed no expression in adult ipRGCs (Figure

2—figure supplement 4C). Additionally, we found that the gamma subunit Gng4 is differentially

expressed in ipRGCs.

Also differentially expressed in ipRGCs were two factors with known roles in

diacylglycerol (DAG) signaling, *Rasgrp1* and *Dgkg* (Figure 2—figure supplement 4B). Ras guanyl

nucleotide-releasing protein 1 (*Rasgrp1*) is a guanine nucleotide exchange factor (GEF) that

activates *Ras* by facilitating its GTP binding (Bivona et al., 2003). Rasgrp1 binds DAG and Ca^{2+},

both of which are elevated by melanopsin phototransduction. This provides a possible basis for

intrinsic photoresponses of ipRGCs to modulate Ras signaling and thus genes governing cell

growth, differentiation and survival. We will return to a more detailed consideration of Rasgrp1

later in this report. Diacylglycerol kinase gamma (*Dgkg*) (Bivona et al., 2003; Shulga et al., 2011)

converts DAG to phosphatidic acid, thus acting as a terminator of DAG signaling. Because DAG

appears to be a key link between early steps in phototransduction and gating of the light-activated

channels, Dgkg may regulate the kinetics of the photoresponse in ipRGCs. The protein products

of the two overexpressed genes may interact. Diacylglycerol kinases are also known to bind to

Rasgrp and modulate its activity (Topham and Prescott, 2001). Diacylglycerol and calcium are

also known to activate the protein kinase C (PKC) family members Prkcd and Prkcq (Oancea and

Meyer, 1998), which we determined to be differentially expressed in ipRGCs. Protein kinase C

(PKC) activity has been suggested to be important for deactivating TRPC activity in the

invertebrate photoreceptors and potentially also for the Opn4 phototransduction cascade

(Graham et al., 2008). Peirson and colleagues (2007) previously identified another PKC member,

Prkcz, as being important for ipRGC-mediated photoentrainment of circadian rhythms (Peirson et

al., 2007). However, Prkcz is only moderately expressed and similar to control samples in our

study.

In other photoreceptors, RGS (Regulator of G protein signaling) proteins play a key role

in terminating the photoresponse by accelerating the intrinsic GTPase activity of the cognate G-

protein (e.g., transducin in rods). Two RGS genes were overexpressed in all three ipRGC pools:

Rgs4 and *Rgs17*. At least one of these (Rgs17) regulates Gq signaling (Mao et al., 2004; Ji et al.,

2011) (Figure 2—figure supplement 4B).

The arrestins also contribute to response termination by binding to phosphorylated opsin.

ipRGCs exhibited strong expression of both beta arrestin genes (Arrb1, Arrb2) but low expression

of rod (*Sag*) and cone (*Arr3*) arrestin genes. This is consistent with earlier evidence that beta

arrestins rather than conventional retinal arrestins bind photoactivated melanopsin in ipRGCs

(Cameron and Robinson, 2014). Still, these beta arrestin transcripts are both at similarly high

levels in cRGCs as in ipRGCs, presumably because these arrestins regulate diverse GPCRs

(Figure 2—figure supplement 4B).

Many of the genes involved in rod and cone phototransduction had low expression (scarce

or no read alignment) and/or were present at much lower levels in ipRGCs than cRGCs. These

include the genes for opsins, transducin alpha, and arrestin in rods (*Rho*, *Gnat1, Sag*) and cones

(*Opn1mw, Opn1sw, Gnat2,* and *Arr3*; Figure 2—figure supplement 4C). Although Cngb1 was

differentially expressed in ipRGCs, the total reads aligning to the Cngb1 locus were low and

derived mainly from a limited region of the gene, and the obligatory alpha subunits were not

detected, so this could be a false positive (Figure 2—figure supplement 4C).

Genes differentially repressed in ipRGCs

The lack of contamination by non-RGC retinal neurons in the postnatal day 5 (P5) samples

allowed us to identify genes that were differentially repressed in ipRGCs compared to cRGCs in

early postnatal development. Our data suggested that the transcription factor *Jun* (Jun Proto-

Oncogene) and *Irx4* are differentially repressed in P5 ipRGCs samples (Figure 2—figure

supplement 3). Other genes that were differentially repressed in the P5 ipRGC samples included

Satb1, *Satb2*, and *Foxp2* showed, all of which are known to have restricted expression in the

abundant F-RGC type that is likely included in the cRGC samples (Rousso et al., 2016). The

Pou4f1 (*Brn3a)* and *Pou4f3* (*Brn3c*) transcription factors were both differentially repressed in P5

ipRGCs, consistent with their known lack of expression in ipRGCs (Jain et al., 2012) (Figure 2—

figure supplement 3). The transcriptional repressors *Bcl11b (CTIP2)*, *Irx4*, and *Tbr1* were all found

to be differentially repressed in ipRGC compared to cRGCs samples. Furthermore, the *Cdkn1c*

(*p57KIP2),* a gene known to be transcriptionally repressed by Bcl11b (Topark-Ngarm et al., 2006),

had relatively increased expression in ipRGCs (Figure 2—figure supplement 3).

Expression differences between the *Opn4-GFP* and *Opn4-Cre/GFP* reporter systems

To study gene expression differences across the different ipRGC subtypes, we compared

the expression patterns of *Opn4-Cre/GFP* (labels all M1-M5 subtypes) and *Opn4-GFP* (labels

only the M1-M3 subtypes) (Figure 2—figure supplement 1). In general, genes differentially

expressed in ipRGCs identified in the two reporter systems were both supportive and cross-

correlated. However, we identified 24 genes that were differentially expressed in the adult *Opn4-*

Cre/GFP reporter but had low or no apparent expression in the *Opn4-GFP* reporter, suggesting

selective expression in one or more of the M4-M6 ipRGC subtypes. The *Opn4-Cre/GFP* specific

genes included *Anxa2*, *Gem*, *Sema3d*, *Rbp4*, and *Rxrg*. Recently, an Rpb4 reporter (Rbp4-Cre)

was demonstrated to mark amacrine cells coupled to ipRGCs, although there was an apparent

lack of labeling in ipRGCs (Sabbah et al., 2017). The Kcnk4/TRAAK, another gene that was

differentially expressed in the *Opn4-Cre/GFP* reporter, encodes a two-pore potassium channel

subunit (Fink et al., 1998). Additionally, our data suggest that the Kcns3 electrically silent voltage-

gated potassium channel subunit has its expression restricted to the ipRGCs labeled by *Opn4-*

Cre/GFP, but this difference did not reach statistical significance (FDR 0.13). However, close

inspection of reads aligning to Kcns3 using the integrated genome viewer (IGV) confirmed weak

expression in ipRGCs and absent expression in cRGCs for *Opn4-Cre/GFP* samples (data not

shown). Lastly, the neurexophilins Nxph1 and Nxph3 were differentially expressed in the *Opn4-*

GFP and *Opn4-Cre/GFP* reporters, respectively (Figure 2—figure supplement 1). These proteins

are known to bind α-neurexins in mice and have restricted expression patterns (Missler et al.,

1998; Beglopoulos et al., 2005; Craig and Kang, 2007).

Rasgrp1 is selectively expressed in ipRGCs

We sought to test our transcript-level differential expression analysis at the protein level

and to determine whether their expression is selective for particular adult ipRGC subtypes.

Transcriptional profiling suggested that Rasgrp1 is expressed differentially, possibly even

selectively, in ipRGCs. However, differential mRNA expression does not guarantee a

correspondence with protein product (Koussounadis et al., 2015). Therefore, we used

immunofluorescence against Rasgrp1 (Puente et al., 2000) to label the Rasgrp1 protein in whole-

mount retinas from adult wild type mice. Rasgrp1-immunopositive somata were present in the

ganglion cell layer (GCL) and in the inner nuclear layer (INL). The latter likely represent amacrine

cells or displaced ganglion cells, judging by their close proximity to the inner plexiform layer (IPL)

(Figure 3). Immunolabeling marked the cytoplasm as well as the somatic plasma membrane of

these cells. Occasionally, particularly strongly Rasgrp1-labeled cells had some dendritic labeling.

Rasgrp1 immunostaining was also observed in a subset of photoreceptors in the outer retina (data

not shown).

To test whether the Rasgrp1-positive cells in the ganglion-cell layer were RGCs, we

carried out double immunofluorescence for both Rasgrp1 (antibody m199) and the RNA-binding

protein Rbpms, which selectively labels all and only RGCs (Rodriguez et al., 2014). About half of

Rasgrp1-immunopositive cells in the GCL were RGCs, as determined by co-labeling for Rbpms

(56%, $n = 708$ across three retinas, Figure 3A,B). The remainder can be assumed to be displaced

amacrine cells.

Most of these Rasgrp1-expressing RGCs were ipRGCs, as revealed by their

immunoreactivity for melanopsin (95.9$\pm$1.1%, $n = 412$, Figure 3A,B). In contrast, only a fraction of

Opn4-immunopositive ipRGCs were Rasgrp1 immunopositive (34%, *n* = 1169). Thus, Rasgrp1

expression in the GCL is apparently restricted to a subpopulation of ipRGCs.

We next tested whether the immunolabeling of RGCs represented endogenous Rasgrp1

protein expression. The antibody used in this study has been previously shown to specifically

label Rasgrp1 protein expression in hippocampal neurons (Pierret et al., 2000). As a further test

for the specificity of the antibody, we compared immunofluorescence labelling of whole mount

retinas from normal and Rasgrp1-knockout mice generated by inserting *LacZ* and a *Neo* cassette

into the Rasgrp1 gene to disrupt its expression (Dower et al., 2000). Our control experiments

showed that the GCL and INL cellular immunolabeling is absent in the Rasgrp1 knockout (Figure

3C). However, vasculature and photoreceptor cell labeling persisted in Rasgrp1-knockout mouse

retinas, suggesting cross-reactivity of antibody with other proteins.

Rasgrp1 expression is restricted to diverse ipRGC subtypes

We next determined which of the established morphological subtypes of ipRGCs express

Rasgrp1 into adulthood (Figure 3D). For this purpose, we used key characteristics such as relative

Opn4 expression, soma size, and dendritic morphology (see Methods). In the GCL, the majority

of M1/3 cells (71.6$\pm$3.9%, *n* = 300 across three retinas) but only a fraction of M2 cells (23.4$\pm$5.4%,

n = 389) and M5/6 cells (31.4$\pm$6.13%, *n* = 138) expressed Rasgrp1 (Figure 3D). In the INL,

displaced M1 cells express Rasgrp1 at a similar percentage as conventionally placed M1 cells

(70.7$\pm$8.0%, *n* = 3 retinas). We found no examples of Rasgrp1 immunoreactivity in M4 cells (0%,

n = 172, Figure 3D). Of the Rasgrp1-expressing ipRGCs, half were M1/3 cells (52.8$\pm$3.8%), nearly

a quarter were M2 cells (21.1$\pm$2.7%) and a small percent (10.4$\pm$1.8%) were M5/6 cells (*n* = 396

Rasgrp1[+]/Opn4[+] cells across three retinas). Therefore, Rasgrp1 is selectively expressed in a

diverse set of ipRGC subtypes.

Spatial Distribution of Rasgrp1-expressing ipRGCs and amacrine cells

Our data show neither a ventral-dorsal nor a naso-temporal density gradient of Rasgrp1-expressing M1-M3 ipRGC across the retina (Figure 3—figure supplement 1C). However, we did observe a minor ventral-dorsal gradient of Rasgrp1-expressing RGCs, with a higher density in the ventral (65%) compared to the dorsal (35%) retina (Figure 3—figure supplement 1E). As shown above, Rasgrp1-expressing RGCs are almost exclusively ipRGCs (96%).

Tbx20 is expressed in a diverse subset of ipRGCs

The T-box transcription factor Tbx20 was suggested from our gene expression analysis to be differentially expressed in ipRGCs. Immunofluorescence co-loclalization analysis of Tbx20 and Opn4 expression confirmed its high expression in a subset of ipRGCs (Figure 4D). Tbx20 was expressed in most M1 cells (82.6±1.8%, n=514 across four retinas), but only in a minority of M2/3 cells (30.2±6.5%, n=1305) and M5/6 cells (12.4±3.4%, n = 603). Half of the displaced M1 (dM1) cells expressed Tbx20 (46.0±7.1%, n = 153). Strikingly, however, Tbx20 was not expressed in M4 cells (0%, n = 283). These results demonstrate that Tbx20 is expressed in a diverse subset of ipRGCs.

Many Tbx20 cells were not detectably immunopositive for Opn4. Only 41% of Tbx20 immunopositive were also Opn4-immunoreactive (18.5±2.6% were M1 cells; 16.0±2.0% were M2/3 cells; and only 3.6±0.9% were M5/6 cells; n = 2184 across four retinas; Figure 4—figure supplement 1). The remaining Tbx20-immunopositive cells were RGCs, as confirmed by Rbpms-immunoreactivity (data not shown). Additionally, Tbx20-immunopositive RGCs that were also Opn4-immunonegative were topographically enriched in the ventral retina, with most Tbx20-positive cells in the dorsal retina being accounted for by Opn4-immunoreactivity.

Tbx20 expression in M5-M6 ipRGCs

To investigate whether some or all of the Tbx20-immunopositive RGCs that were Opn4-

immunonegative might be ipRGCs of the M5 and M6 subtypes that exhibit weak Opn4

immunostaining, we examined the co-localization of Tbx20-immunopositive cells, Opn4-

immunopositive cells, and all GFP-labeled cells in the *Opn4-Cre;Z/EG* mouse reporter, which

among other ipRGCs, labels M5 and M6 cells. We observed examples of Tbx20-immunopositive

cells that were GFP-positive (M1-M6 ipRGCs), but not Opn4-immunopositive (M1-M4),

suggesting that Tbx20 is expressed in at least a subset of M5 or M6 ipRGCs (Figure 4—figure

supplement 1A,B).

To test the implication that many Tbx20 cells were M5 or M6, we turned to Cdh3-GFP

mice. Recently, our group determined that essentially all GFP+ RGCs in the Cdh3-GFP reporter

mouse you used in your studies are either M6 cells (the great majority) or M5 cells (minority)

(Quattrochi et al., 2018). We tested Tbx20 immunoreactivity in the context of Opn4

immunofluorescence and *Cdh3-GFP* labeling (Figure 4B,C). For the purpose of this study, we

focused on GFP cells in the GCL that are Opn4-immunonegative (to distinguish from Opn4-

immunopositive M2 types). We found that at three weeks after birth, most *Cdh3-GFP* cells express

the Tbx20 protein (82.1±4.3%, *n* = 439 across four retinas, Figure 4C). Many, but not all, of the

Opn4-immunonegative Tbx20-positive cells were GFP$^+$ (27%, n=1277 Tbx20$^+$;Opn4$^-$ cells; 4

retinas). The dorsal-ventral gradient of Tbx20-positive cells that are Opn4-immunonegative was

broadly similar to the retinal labeling of the *Cdh3-GFP* reporter. A large portion of Tbx20-

immunopositive cells remained unclassified (43.3±3.8%, n=2184; Figure 4—figure supplement

1D).

Further, we determined whether Tbx20 expression correlates with the related T-box

transcription factor Tbr2, a gene previously described to be enriched in adult ipRGCs (Mao et al.,

2014; Sweeney et al., 2014). All Tbx20-expressing cells were strongly Tbr2-immunopositive (*n* =

328; 4 regions distributed across a single adult *Opn4-Cre/GFP* retina; Figure 4—figure

supplement 1C). Therefore, whereas Tbr2 is expressed in a broad range of types that includes

the entire ipRGC family, Tbx20 expression is confined to a diverse subset of ipRGC subtypes.

Molecular diversity of Rasgrp1 and Tbx20 expression in ipRGCs

Our expression studies revealed that Rasgrp1 and Tbx20 have a strikingly similar pattern

of expression among ipRGC subtypes. Both genes were expressed in the majority of M1 cells, a

minority of M2 cells, and a small population of M5/6 cells, but not in M4 cells (Figures 4 and 5).

To directly test for co-expression, we compared and contrasted the expression patterns of Tbx20-

and Rasgrp1-immunoreactivity in the context of the M1-M4 subtypes revealed by Opn4-

immunoreactivity (Figure 5). Rasgrp1 co-expression with Tbx20 was only observed in a fraction

of M1/3 cells (26.0±1.8%; n = 241 across two retinas; Figure 5). Further, M1 cells expressing

either Rasgrp1 or Tbx20 alone accounted for roughly similar fractions of M1 cells (37.3±4.0% and

31.1±3.5%, respectively). Only a small fraction of M1 cells were immunonegative for both Rasgrp1

and Tbx20 (5.3±0.7%). In contrast, half of M2 cells lacked Rasgrp1 and Tbx20 immunoreactivity

(57.7±3.3%; n = 388 across two retinas). Approximately a third of M2 cells expressed Tbx20

(31.0±0.7%), while only 11.3±5.1% expressed Rasgrp1. We did not observe any example of an

M2 cell expressing both Rasgrp1 and Tbx20.

Molecular diversity of SCN-projecting ipRGCs

We further examined the Rasgrp1- and Tbx20-expressing ipRGC subtypes to seek

intersectional expression patterns that would divide ipRGCs by their downstream visual pathways.

Earlier studies showed that M1 cells could be subdivided based on their level of expression of

Brn3b (Chen et al., 2011). We used quadruple immunolabeling to simultaneously test Brn3b

expression with Rasgrp1- and Tbx20-immunoreactivity in the context of Opn4-immunolabeled

ipRGCs (25 regions, three wild type retinas; Figure 5A,B). We determined that a minority of M1/3

cells express Brn3b (7.9±6.0%, n=241), which is similar to previous studies (Jain et al., 2012).

The Brn3b⁺ M1/3 cells expressed either Tbx20 or Rasgrp1 (91.0 and 9.0±10.1%, respectively;

n=30) (Figure 5C).

Further, we determined that most M2 cells expressed Brn3b (90.8±6.9, n=168). In contrast

to M1/3 ipRGCs, the majority of Brn3b⁺ M2 ipRGCs did not express either Rasgrp1 or Tbx20

(67.4.6±13.8, n=222). Most M2 cells expressing Tbx20 were also Brn3b immunopositive

(84.5±25.1, n=118). The small subset of M2 cells that express Rasgrp1 could be further divided

by Brn3b presence or absence (5.0±6.0% and 4.5±1.4%, respectively; n=168). Generally, we

found no cells co-expressing all three genes (n=729).

Our immunostaining study in the retina suggested that SCN-projecting M1 cells (Brn3b-

negative) might represent molecularly diverse cell populations (Figure 5C). We correlated gene

expression of Rasgrp1 and Tbx20 in the retina with retrograde labeling from the SCN (Figure 6A).

We injected rhodamine-conjugated retrobeads in the SCN, followed by immunofluorescence

labeling for Opn4, Rasgrp1, and Tbx20 (Figure 6A-C). All injection sites clearly involved the SCN,

as revealed by DAPI labeling, but did not spread to the optic chiasm or tract (Figure 6B).

Quantitative co-expression analysis (18 confocal images collected across the contralateral and

ipsilateral retinas) revealed that nearly all retrolabeled cells were Opn4-immunopositive (95%).

These cells exhibited variable patterns of staining for the other proteins. Most cells expressed

both Rasgrp1 and Tbx20 (76%), but equal minorities expressed either Rasgrp1 (12%) or Tbx20

(12%, Figure 6D). This expression pattern was consistent across the ipsi- and contralateral retina

(Figure 6D), as suggested by a bilateral input to the SCN (Hattar et al., 2006; Fernandez et al.,

2016). Therefore, we show that SCN-projecting ipRGCs have a complex pattern of Rasgrp1 and

Tbx20 gene expression. Together, these results provide evidence for previously unrecognized

molecular diversity in adult ipRGCs.

Discussion (2657 words)

Prior efforts to assess the distinctive genetic composition of ipRGCs have been complicated by

their rarity among diverse retinal cell types and the inherent difficulties of maintaining viability of

dissociated mature neurons (Lobo et al., 2006). Our approach was first to isolate RGCs by

immunoaffinity, then to further purify ipRGCs from these based on genetic labeling and FACS,

and to finally to compare the transcriptional profiles of the purified ipRGCs to those of the residual

cell pool, consisting mainly of conventional RGCs. The relative purity of our ipRGC sample is

supported by enrichment for transcripts of genes known to be differentially expressed in ipRGCs

and the low levels of transcripts selectively expressed in potentially contaminating populations,

including the abundant rod photoreceptors. Our isolation method and differential expression

analysis allowed us to identify more than 75 differentially expressed genes in ipRGCs relative to

conventional RGCs.

Genes differentially expressed in adult ipRGCs

There is limited knowledge of specific gene expression in ipRGCs generally and within

particular ipRGC subtypes, especially non-M1 ipRGCs. Many diverse genes appeared more

highly expressed in ipRGCs than in conventional RGCs. We confirmed differential protein

expression in ipRGCs immunohistochemically for two of these genes: Tbx20, a transcription factor

implicated in visual development; and Rasgrp1, a G-protein exchange factor that may interact

with the melanopsin phototransduction cascade. However, only a subset of ipRGCs appeared to

express detectable levels of these proteins, and such variable expression was apparent even

among ipRGCs of the same subtype. Some ipRGCs expressed both proteins, but many did not.

This diversity even extended to the M1 cells projecting to the SCN, which had been thought to

share the distinctive molecular feature of little or no expression of the transcription factor Brn3b.

These novel markers of molecularly distinctive ipRGC varieties open the way for cell-type-specific

manipulations through intersectional strategies.

What type(s) of adult ipRGCs express Rasgrp1?

Rasgrp1 expression has previously been detected in the hippocampus, striatum and

olfactory regions of the brain (Ebinu et al., 1998; Toki et al., 2001), but our study appears to be

the first to explore Rasgrp1 expression in the eye. Rasgrp1-like immunoreactivity marked a

diverse subpopulation of ipRGC subtypes, including the M1-M3 ipRGC subtypes but not the M4-

type. Either M5 or M6 ipRGCs, or both, also appear likely to express Rasgrp1, because some

Rasgrp1 immunoreactive cells had weak Opn4-immunoreactivity without the characteristic

dendritic labeling of M1-M3 ipRGCs and with somas too small to be M4 cells (Ecker et al., 2010;

Quattrochi et al., 2018; Stabio et al., 2018).

What is the function of Rasgrp1 in adult ipRGCs?

The function of Rasgrp1 in ipRGCs is unknown, but it could interact with the melanopsin

phototransduction cascade. The direct photoresponse of ipRGCs appears to increase levels of

both DAG and calcium. Both of these signaling molecules bind to and activate Rasgrp1, and

trigger its translocation to the Golgi apparatus (Bivona et al., 2003; Graham et al., 2008; Zhang

et al., 2010). However, ipRGC phototransduction Rasgrp1 signaling does not appear to be

essential for ipRGC phototransduction because more than a quarter of M1 ipRGCs and the great

majority of M2 cells are immunonegative for Rasgrp1. Even in ipRGCs, Rasgrp1 may be activated

by DAG and calcium signals unrelated to Opn4 phototransduction, and such signals are

presumably also responsible for modulating Rasgrp1 in cells (such as certain amacrine cells)

which express Rasgrp1 but not melanopsin.

Rasgrp1 has the potential to affect any number of neuronal signaling pathways. Ras

signaling pathways are enormously complex and the cross talk between pathways makes it even

harder to identify specific effects. One basic mechanism for specificity in Ras signaling is the

distinct subcellular targeting of downstream components of the signaling pathway. In

lymphocytes, localized Ras signaling of Rasgrp1 occurs preferentially on the Golgi apparatus,

which is a rare form of compartmentalized Ras signaling (Bivona et al., 2003; Zhang et al., 2010).

The Golgi apparatus in neurons provides the posttranslational protein modifications required for

organizing protein and organelle trafficking throughout the cell. Rasgrp1 could play a crucial role

in orchestrating a specific set of post-translational modifications at the Golgi.

An important survival mechanism in M1 ipRGCs is the maintenance of mTOR expression

by melanopsin phototransduction (Duan et al., 2015; Li et al., 2016). The DAG-activated Rasgrp1-

Ras-Mek1/2-Erk1/2 pathway contributes to mTOR activation in thymocytes (Gorentla et al., 2011).

The percentage of M1 cells expressing Rasgrp1 (72%) in this study is similar to the percentage

of M1 cells that survive optic nerve crush (~70%, Li et al., 2016). It would be interesting to test

whether Rasgrp1 is required for the maintained mTOR levels in these surviving M1 ipRGCs.

Recent studies have also clarified different mechanisms responsible for M1 survival and the

regeneration potential of alpha-RGCs such as M4 ipRGCs (Duan et al., 2015; Li et al., 2016).

Despite surviving well, M1 ipRGCs were not capable of regenerate their axons (Li et al., 2016).

In contrast, alpha-RGCs had a unique capability to regenerate their axons that was promoted by

their high levels of mTOR expression as well as osteopontin/Spp1 expression coupled with the

growth factor IGF-1(Duan et al., 2015). Whereas M1 ipRGCs were found to maintain mTOR

expression after axon injury, M4 ipRGCs and other alpha RGC types were demonstrated to have

diminished mTOR expression levels (Li et al., 2016). The absence of detectable Rasgrp1

expression in M4 ipRGCs may help to account for their failure to maintain mTOR expression after

injury.

Central brain targets of Rasgrp1-expressing ipRGCs

At the circuit level, Rasgrp1 is not anticipated to be an essential regulator of circadian

photoentrainment. The majority of M1 ipRGCs are known to provide the primary retinal input to

the SCN, while a subset of M1 ipRGCs send input to the OPN to regulate the pupillary reflex. Our

studies found that Rasgrp1 was expressed in the majority of M1 ipRGCs, which suggested to us

that it might correlate directly and completely with the SCN-projecting M1s. However, retrograde

tracing experiments from the SCN revealed that some SCN-projecting M1 cells were Rasgrp1-

immunonegative. Accumulating research suggests that the SCN is more compartmentalized than

previously recognized (Bedont and Blackshaw, 2015). The question of whether Rasgrp1 is

expressed in a subset of M1 ipRGCs that targets a specific compartment of the SCN remains to

be determined.

We find Rasgrp1 to be expressed not only in SCN-projecting M1 ipRGCs, but also in other

ipRGC subtypes, especially M2 cells and apparently M5 and/or M6 cells. Collectively, these types

project to various non-image-forming visual centers, including the olivary pretectal nucleus,

intergeniculate leaflet, and dorsal lateral geniculate nucleus (Quattrochi et al., 2018; Stabio et al.,

2018).

Tbx20 is expressed in a diverse set of ipRGC subtypes

The T-box transcription factor Tbx20 exhibited enriched expression relative to

conventional RGCs in postnatal and adult retinas. Double immunolabeling revealed that many

ganglion cells that expressed this protein also expressed melanopsin. As was true for Rasgrp1,

Tbx20 was determined to be expressed in most M1 ipRGCs (75%), a substantial minority of M2

cells (40%) and an additional population of RGCs whose identity was not immediately obvious.

We decided to then compare Tbx20 against other known molecular patterns in ipRGCs. Whereas

most RGCs follow a Brn3b-dependent developmental program, the M1 ipRGCs that project to the

SCN do not express Brn3b while OPN-projecting M1 ipRGCs express Brn3b. We found that

Brn3b-expressing M1 cells are also Tbx20-immunopositive. The Brn3b-negative M1 cells are split

between cells that express Tbx20 and those that do not. This finding suggests that ipRGCs are

more molecularly diverse than originally anticipated: Tbx20 is expressed in ipRGCs with differing

brain targets, Tbx20 is expressed across multiple morphologically defined subtypes, and Tbx20

is not expressed in all of any one type. The exploration of Tbx20 coexpression with Rasgrp1

revealed a complex coexpression pattern among M1-M3 ipRGCs.

What is the function of Tbx20 in adult ipRGCs?

Tbx20 has well-established roles in embryonic development and is continuously required

in mature neurons and other cell types to maintain their identities and functional properties during

adulthood (Naiche et al., 2005). Tbx20 functions as a repressor in early embryonic ocular

development (Carson et al., 2000; Pocock et al., 2008) and is required for the expansion of the

small pool of precursor cells in the optic vesicle (Carson et al., 2004). However, little is known

about the function of Tbx20 in the adult retina.

Tbx20 can function as a transcriptional activator in parallel with its repressor activity, with

these two roles impinging on distinct biological processes (Sakabe et al., 2012). In addition to its

key developmental roles, Tbx20 appears vital for maintaining the structure and function of cardiac

muscle cells in the adult mouse heart (Stennard et al., 2003; Shen et al., 2011). In adult

cardiomyocytes, Tbx20 is responsible for directly activating genes critical for normal adult cardiac

function such as those required for ion transport and heart contraction (Shen et al., 2011; Sakabe

et al., 2012). In contrast, genes directly repressed by Tbx20 have known roles in non-heart

developmental programs, cell cycle, proliferation, and immune response (Sakabe et al., 2012).

The transcriptional effects of Tbx20 shift during cardiac development, from early mediation of

proliferation of cardiac progenitors, to implementation of an anti-proliferative program in the adult

heart (Cai et al., 2005; Takeuchi et al., 2005). Therefore, Tbx20 cooperates with distinct cohorts

of transcription factors to either promote or repress distinct molecular programs in a context-

dependent manner (Sakabe et al., 2012). Similarly, binary cell fate specification in the retina is

driven by complex genetic programs that require the simultaneous activation and repression of

genes by transcription factors. Tbx20 may prove to have a similar reversal in its transcriptional

activity in the retina when transitioning from broad embryonic development program to regulating

adult neuron identity of a subset of ipRGCs. Further, Tbr2 is another Tbox family member that is

known to have a critical role in the development of retinal ganglion cells (Mao et al., 2008), which

later becomes essential to a restricted set of ipRGCs that participate in NIF visual circuits (Mao

et al., 2014; Sweeney et al., 2014). Our studies determined that Tbx20 and Tbr2 are coexpressed

in adult ipRGCs. They may work cooperatively to specify ipRGC subtype identity by regulating

cell-specific transcriptional programs and repressing alternate fates.

Characterization of ipRGC subtypes

Retinal cell types are generally classified using a combination of morphology, gene

expression, mosaic organization, light responses and synaptic connectivity (Sanes and Masland,

2015). By these criteria, intrinsically photosensitive RGCs comprise at least 6 distinct cell types

(Figure 7). Though all express melanopsin, they differ from one another in the strength of the

intrinsic response, their morphology, pattern of light responses, and projections to the brain.

However, the further subdivision may be in order. The M1 type appears subdivisible into at least

two subtypes, one expressing the transcription factor Brn3b and innervating the OPN and

geniculate complex, while the other lacks Brn3b expression and innervates the SCN (Chen et al.,

2011). Our study shows further diversity in the M1 and M2 types based on the expression of

Rasgrp1 and Tbx20. For example, we find molecular diversity among in the SCN-projecting

ipRGC subtypes (Figure 7). It is unclear to us whether this should be used to propose a further

formal subdivision of M1 and M2 cells. For example, the expression of these proteins could

fluctuate over time in individual cells and be uncorrelated across cells of the same type. One

would like to know that these patterns of expression are stable over time and correlated with other

cell features before proposing such further subdivision. The extensive overlap among dendritic

fields of M1 (and of M2) cells (Berson et al., 2010) means that either type could be subdivided

into two or perhaps three subtypes while still maintaining full retinal coverage (i.e., tiling), but it

seems likely that we will either have to accept that there is substantial molecular diversity with

single types (as there is substantial functional diversity among M1 cells (Emanuel et al., 2017))

or that the dogma of complete retinal tiling by single types must be abandoned.

In addition to Tbx20 expression in a subset of SCN-projecting M1s, Tbx20 may also

regulate the molecular program of ipRGCs projecting to the OPN and control the pupillary light

reflex. The Brn3b-expressing M1 ipRGCs, a subset of M2 ipRGCs and *Cdh3-GFP* labeled M6

ipRGCs are all known to project to the OPN, and all express Tbx20. However, this is not a direct

association and further retrograde or Tbx20 conditional knockout studies are required, especially

to determine the function of Tbx20-expressing M2s.

Comparison with other ipRGC gene expression profiles

Siegert and colleagues (2012) surveyed gene expression in many different sets of mouse

retinal neurons, using specific mouse reporters strains (including the *Opn4-Cre* reporter system

for ipRGCs), FACS isolation of labeled cells, and microarray analysis. Many of the genes they

found strongly expressed in ipRGCs were also among the genes we found differentially expressed

in ipRGCs. However, dozens of additional genes differentially expressed in ipRGCs emerged

from our analysis that were not detected in theirs (Siegert et al., 2012). Discrepancies between

their findings and ours may stem from technical factors such as differing degrees of contamination

of the starting material with rod photoreceptor transcripts, the use of internal control cell

populations for relative gene expression comparison in our study but not theirs, and differences

between microarray and RNA-sequencing methodologies.

Another recent studied used single-cell transcriptomic analysis of the mouse retina and

were able to identify ipRGCs from their cell suspensions (Macosko et al., 2015). Single cell

technology is ideal, in principle, for the precise identification of an individual neuron's molecular

identity despite the extreme heterogeneity of the nervous system. Macosko et al., 2015 could

definitively distinguish the main broad class of RGCs, but they required *post hoc* supervised

analysis to distinguish a limited number of genes attributed to Opn4-positive cell clusters. They

identified nine genes with a two-fold increase in expression compared to Opn4-negative cells.

Three genes (Tbr2, Igf1, and Tbx20) were also found to be enriched in our ipRGC samples. In

contrast, Tbx20 did not reach above threshold for Siegert et al., (2012), but it is among the highest

expressing ipRGC-enriched genes in our study. Single-cell gene expression profiling methods

such as Drop-Seq hold great promise and will certainly continue to be pursued more in future

studies of neuron subtype gene expression. Recently, single-cell studies of pre-enriched bipolar

cells were able to distinguish molecular markers for all previously recognized bipolar cell types

(Shekhar et al., 2016). Similarly, ipRGCs subtype-specific gene expression may become

deciphered in the future using single-cell analysis that incorporates a pre-enrichment step for

ganglion cells.

Conclusion

In conclusion, our results demonstrate a method to purify ipRGCs and identify an

extensive list of more than 75 genes that are differentially expressed compared to generic RGCs.

Of course, our identified differentially expressed genes in ipRGCs should not be considered a

complete account of all genes relevant to ipRGC function. Instead, we hope that it will provide a

beneficial resource that will generate hypothesis that lead to key insights into ipRGC function in

non-image forming vision. The more than 75 genes suggested to be differentially expressed in

ipRGCs will be useful for the identification of marker genes for ipRGC subtypes, comparison of

gene expression across types, understanding the intracellular gene networks underlying ipRGC

phenotypes, and the testing for conservation of ipRGC molecular programs across mammalian

species.

We are encouraged that our gene expression profiling data of ipRGCs lead us to the

identification of Tbx20 and Rasgrp1 as novel, selectively expressed genes in ipRGCs. These

results serve as a good proof of principal for the validity of our gene expression profiling results.

In addition, the stable, specific expression in adult ipRGCs suggests that the differential gene

expression of Rasgrp1 and Tbx20 were not simply due to transient, stress-induced molecular

program resulting from the harsh processing steps prior to sequencing. We determined that Tbx20

and Rasgrp1 are expressed across ipRGCs that belong to multiple morphological types, have

diverse molecular expression, differing physiology, and are involved with multiple visual brain

circuits.

Methods

Animals:

All experiments were conducted in accordance with NIH guidelines under protocols approved by

the Brown University (Providence, RI) Animal Care and Use Committee. Both male and female

adult mice (P30 to P90) were used unless otherwise stated. *Opn4$^{cre/cre}$* mice (Ecker et al. 2010)

crossed with floxed-stop reporter mice: "*Z/EG*" (*Jax#003920*); the offspring express GFP in *cre-*

expressing cells (M1-M6), as described by Ecker et al., 2010. *Opn4-GFP(ND100Gsat)* is a BAC

transgenic originated from the GENSAT project at Rockefeller University. Rasgrp1-KO

(Rasgrp1^{tm1Jstn}, Dower et al., 2000) was initially provided generously by Robert Barrington, U. of

Alabama for initial testing. A colony was created inhouse from stock at Jackson labs (Jax 022353).

Rasgrp1-Cre(PO1 founder line) was rederived from Jackson Labs (#34811-UCD). *Cdh3-GFP*

reporter is a BAC transgenic originally generated by the Gensat project (MMRRC,

BK102Gsat/MMNC) and has been used previously (Quattrochi et al., 2018). This mouse line has

been backcrossed in wild type background for at least 10 generations. *Cdh3-GFP* mice were three

weeks old or younger unless otherwise stated.

Gene expression analysis of purified mouse ipRGCs

For our transcriptomics studies, we used two ipRGC reporters available in the lab to

identify selective gene expression in ipRGCs compared to RGCs as a whole: 1) BAC transgenic

Opn4-EGFP Gensat mice from MMRRC (#033064-UCD) and 2) knock-in *Opn4-Cre* mice

(obtained from S. Hattar; Ecker et al., 2010) with Cre-dependent GFP reporter (Z/EG obtained

from Jackson labs; (Novak et al., 2000). The *Opn4-GFP* mice were maintained as heterozygotes

on *C57/BL6* background while *Opn4$^{Cre/+}$;Z/EG* mice were generated by breeding homozygous

Opn4-Cre mice with heterozygous *Z/EG* mice and maintained on mixed background. Although

Opn4-GFP reporter labeling in ipRGCs has not been reported previously, a similar BAC

transgenic strategy has been demonstrated to label M1-M3 ipRGCs (Schmidt et al., 2008). We

tested the correlation of the *Opn4-GFP* reporter expression and endogenous melanopsin in

RGCs. We used anti-melanopsin immunoreactivity label M1-M3 ipRGCs (Berson et al., 2010) and

anti-EGFP antibodies in a whole-mount retina from an adult *Opn4-EGFP* mouse. Importantly, all

Opn4-GFP$^+$ cells were Opn4-immunopositive (n=60 GFP$^+$ cells across seven regions of one

retina, Figure 8A). Unexpectedly, more than half of the Opn4-immunopositive M1-M3 ipRGCs

were not labeled by the reporter (55%, n=133). Therefore, the coexpression of EGFP expression

by the *Opn4-GFP* reporter is strongly correlated with M1-M3 ipRGCs, but only accounts for about

half of the population. The other reporter system used in this study, *Opn4$^{Cre/+}$; Z/EG* mice, has

been previously demonstrated to label with EGFP all six known morphological types of ipRGCs),

named M1–M6 (Ecker et al., 2010; Quattrochi et al., 2018; Stabio et al., 2018). However, Opn4-

immunofluorescence studies of four regions across an *Opn4-Cre/GFP* retina revealed that more

than one-fourth of M1, displaced M1, and M2 cells lacked discernable GFP-labeling (28%, n=81;

27%, n=15; 33%, n=132, respectively) (Figure 8C). There were many additional GFP$^+$ cells that

were Opn4-immunonegative, with large soma cells being designated as M4 ipRGCs (54%, n=67

M4 cells). The remaining identified M4 cells had somas with weak, but present, Opn4-

immunoreactivity and were only partially accounted for by GFP-labeling (27% Opn4$^+$;GFP$^{neg.}$,

n=67 M4 cells). Additionally, small soma GFP$^+$ cells with absent Opn4-immunoreactivity were

designated as M5/6 cells, since the lack of dendritic labeling made it impossible to distinguish

between the M5 and M6 types (72%, n=202 M5/6 cells). We observed small cell bodies with

weakly Opn4-immunlabeling that did not extend to the dendrites, which were also designated as

M5/6 types (28%).

Two ages were chosen for retina tissue collection for purification of ipRGC neurons:

Postnatal day 5 (+/- 1day) and young adult (P30 +/- 3 days). Three or more biological replicates

were used for each dataset (three for postnatal day 4-6 (P4-6) Gensat *Opn4::GFP*, six replicates

for Gensat *Opn4::GFP*, four replicates for *Opn4::Cre/GFP* reporter). Each adult replicate required

the pooled retinas from 15-20 transgenic mice to acquire suitable numbers of cells for the

transcriptomics. These steps are outlined in much more depth and detail in the following sections.

The choice of transcriptomics preparation strategies and final readout of processed

samples are interrelated and subject to a number of technical concerns vital to the success of

transcriptomics analysis. Seven steps in the development and completion of the cell type-specific

transcriptomics procedure of cells isolated from early postnatal and adult mouse retinas (Figure

1). **First**, dissociation of retinal tissue to a cell suspension; **Second**, purification of RGCs; **Third**,

FACS analysis and sorting; **Fourth**, extraction of RNA; **Fifth,** cDNA amplification because the

resulting RNA is typically low in abundance; **Sixth**, shear cDNA into sequenceable fragments that

are then sequenced with Illumina deep-sequencing; and **Finally**, the raw reads must be analyzed

to identify differential expression of genes in ipRGCs.

Retina dissociation

The intertwined nature and tight cell–cell adhesions of neural cells make it difficult to

separate cells without causing cellular damage and activating stress or cell death pathways.

Therefore, vigorous dissociation of tissues can lead to activation of stress or cell death pathways

and distort the resulting expression profile. However, poor dissociation can lead to a severe

decrease in isolated cells and make downstream expression studies an impossibility. As a first

step, we dissected retinal tissue, which we then digested in a protease solution to loosen and

disrupt the intertwined neural cells into single cell suspensions for subsequent cell-type isolation.

The essential components for proper dissociation included: the proteolytic enzyme papain L-

cysteine to promote enzymatic activity, DNase for destroying the extremely sticky free DNA

strands released by damaged cells, and an absence of calcium from solutions to promote

disruption of cell-cell adhesions (Barres et al., 1988). We further optimized cell viability and

recovery by replacing dPBS with HibernateA and including B27 throughout the cell-isolation

procedure (Brewer et al., 1993; Brewer, 1997; Brewer and Torricelli, 2007). The use of survival-

promoting media such as Hibernate-A and supplements such as B27 was particularly relevant for

adult neural dissociation since adult neurons have been demonstrated to be especially prone to

cell death (Eide et al., 2005; Brewer and Torricelli, 2007). We incubated the dissected retinas in

pre-activated protease solution and completed the cell dissociation by triturating the cells with a

1mL pipette tip.

RGC Pre-Enrichment

Pre-enrichment of ganglion cells prior to isolation of ipRGCs is necessary due to their

extraordinary rarity in the retina. Ganglion cells only make up 1% of mouse retinal cells and only

1-5% of ganglion cells are ipRGCs (0.01-0.05% of retina cells) (Hattar et al., 2002; Ecker et al.,

2010; Berson et al., 2010). In contrast, the classic photoreceptors rods and cones account for 75-

80% of all mouse retinal cells (Jeon et al., 1998). Therefore, we pre-enriched for ganglion cells

prior to positively selecting fluorescently labeled ipRGCs using fluorescence activated cell sorting.

We incorporated an immunopanning system that has proven effective at isolating a homogeneous

population of RGCs (Barres et al., 1988; Cahoy et al., 2008). Immunopanning takes advantage

of the cell surface protein Thy1 which is shared among retinal ganglion cells (Barres et al., 1988).

The classic immunopanning process uses antibodies raised against Thy1 to select the RGCs

from the heterogeneous retinal cell suspension. We improved upon viability and reproducibility of

the immunopanning procedure by adapting it to use a magnetic-activated cell sorting procedure

(MACS, Miltenyi Biotec), eliminating the need for using harsh lysis treatment (data not shown).

We incubated the dissociated retinal cell solution with Thy1.2-conjugated magnetic nanoparticles

which are retained in a magnetized column and the isolated RGCs can then be acquired (Haeryfar

and Hoskin, 2004). In preparation for FACS, the DNA intercalating dye, 7-AAD, was added to the

solution to discriminate dying cells that consequently had compromised cell membranes.

Fluoresence-Activated Cell Sorting (FACS)

We used a FACS Aria (BD Biosciences) electrostatic sorter to isolate a homogeneous

GFP-labeled cell population from the dissociated RGC-enriched cell suspension of the transgenic

melanopsin reporter mice (Figure 1). FACS has been successfully used previously to profile gene

expression in cell subtypes of the nervous system (Lobo et al., 2006; Cahoy et al., 2008; Siegert

et al., 2012). Live cell gating was achieved by excluding 7-AAD labeled cells (high fluorescence

signal in far red emission) (Figure 9). Fluorescently labeled ipRGCs were positively selected from

the solution of enriched ganglion cells by their relatively high level of GFP fluorescence (FITC

gating). In parallel, cells with relatively low GFP-fluorescence (low FITC) were also isolated and

designated as a "generic RGC" control population for direct comparison with correlated ipRGC

samples. The parallel isolation of generic RGCs was designed as an internal negative control for

comparison with isolated ipRGCs. Accordingly, the generic RGC populations were treated with

the same reagents, cytometer settings, centrifugation forces, and temperatures throughout the

procedure. This is especially important for isolating adult RGC populations, since they are

particularly susceptible to the stresses of FACS sorting(Lobo et al., 2006; Cahoy et al., 2008;

Heiman et al., 2008). The large amount of small cellular debris generated during the dissociation

process made it a challenge to keep the number of collected generic RGCs consistent with the

rare ipRGCs (Figure 9). Cellular debris registers as being essentially non-fluorescent, is smaller

and less complex than the generic RGC population that is of the same relative size and complexity

as the isolated GFP-positive cells. Therefore, we limited cell debris by acquiring the ipRGC

(GFP+) and generic RGC (GFP-negative) populations with the same relative cell complexity

(indicator of cell health) and cell size selection to exclude the relatively small cellular debris or

doublets.

RNA Extraction

The small volumes allowed by the electrostatic FACSAria sorter allowed us to lyse sorted

cells directly into Qiagen RLT buffer and directly proceed to RNA extraction using Minelute

columns (Qiagen). The enriched RNA was treated on-column with DNase to remove any residual

genomic DNA from the sample. RNA-processing was done in an enclosed RNase-free

environment to limit degradation of RNA throughout the extraction process. Additionally, RNA

integrity was analyzed using the Agilent 2100 Bioanalyzer and the PicoChip, which is able to

qualitatively test the low RNA recovery samples (Figure 10A). Initially, we proceeded immediately

with cDNA processing after RNA extraction. However, freezing at -80 degrees did not seem to

effect RNA integrity since the frozen RNA samples still received RIN score of 9.0 or greater

(Figure 10A). Therefore, most of the cDNA libraries were prepared after storage of extracted RNA

at -80 degrees Celsius.

cDNA Preparation

RNA-seq transcriptome analysis requires large amounts of RNA material using TruSeq,

ranging on the order of 100-1000ng of total RNA. However, our improved method for isolating

ipRGCs was able to isolate 12,000 GFP+ ipRGCs from nine postnatal day 5 (P5) transgenic

reporter mice. This was only expected to provide about 12ng of extracted total RNA by qualitative

estimates considering that a single cell holds 5-10pg of total RNA. Further, twice as many adult

mice of the same genotype were required to provide only 1,000 cells due the the relative fragility

of adult retinal ganglion cells described above. Therefore, we decided that some form of

amplification was necessary to study the molecular programs used by adult ipRGCs. The Nugen

Ovation RNA amplification system was successfully used in microarray studies with as little as

500pg of total RNA input (Caretti et al., 2008; Clément-Ziza et al., 2009; Morse et al., 2010)

(Figure 10B). Sequencing analysis using the Ovation system has previously been reported to

generate cDNA containing negligible rRNA reads (<4%), while providing a representative

transcriptome with sufficient biological replicates(Tariq et al., 2011).

RNA-seq library preparation

We determined that the Truseq system was the necessary platform for my cDNA samples

to produce the 10nM sequencing library concentration required at the on-site Genomics Core

facility in preparation for 50bp single-end Illumina sequencing. Before preparing the library, we

first sheared cDNA to the appropriate size, (200-300bp median) using the Covaris system (Figure

10C). Each sample was subsequently processed using a unique barcode adapter to allow for

multiplexing multiple samples on the HiSeq (commonly 200million 50bp reads divided among

sequencing samples). Excess adapter sequences was removed using Ampure bead isolation,

which removes all DNA fragments less than 200bp. Finally, the Genomics Core completed the

final quality control of the DNA library prior to sequencing: testing the library fragment size

distribution (High Sensitivity Bioanalyzer) and qPCR analysis using primers that match the library

adapters (Quail et al., 2008) (Figure 10D).

Initially, we processed a large number of samples at once with multiplexed sequencing

using HiSeq (8 samples per lane) to minimize high costs of sequencing at the expense of

sequencing depth. We later reran many of the sequencing libraries with less multiplexing,

enabling increased sequencing depth. The corresponding technical replicates were merged

together for differential expression analysis (Figure 10E). The final read counts of each sample is

shown in Figure 10F.

Differential gene and transcript expression analysis

The completion of sequencing generated tens of millions of reads that are used to

compare gene expression levels between isolated ipRGCs (GFP+) and generic RGCs (GFP-).

Well-established, powerful RNA-seq differential expression analysis pipelines have been

developed such as Cuffdiff, EdgeR, and DESeq (Anders and Huber, 2010; Trapnell et al., 2012;

Anders et al., 2013). The Cuffdiff pipeline prioritizes isoform quantification and diversity (Trapnell

et al., 2012). However, the short 50bp single-end reads that are generated in our study are not

well-suited for prioritizing isoform discovery and analysis. Further, Cuffdiff does not fully take

advantage of our purposeful pairwise-comparison between groups. In contrast, the EdgeR

package is better suited for our purpose; it is designed to count the number of reads that align to

an annotated gene (mouse reference genome, in our case) and subsequently performs statistical

analysis on a generated table of counts to identify quantitative changes in expression levels

between the two experimental samples. EdgeR compares and retains the relationship between

all pairs of experimental samples when calculating differential expression likelihood (Anders et

al., 2013). Our analysis filtered out genes with very low counts, less than 1 count per million (cpm),

in more than half of the samples used in the differential expression analysis. This is a common

cutoff and considers 1) that a gene must be expressed at a minimum threshold in order to become

biologically important and 2) that the inclusion of genes with very low counts may negatively affect

the statistical approximations used by the EdgeR pipeline.

To identify the set of differentially expressed genes in the ipRGC populations, we used the

following strict criteria. First, we identified genes with low false discovery rate (FDR < 0.05) and

high fold-change (greater than 2-fold) suggesting differential expression between ipRGCs and

generic RGCs. Second, we considered whether the differentially gene expression was

corroborated across both reporters (*Opn4::GFP* labeling M1-M3 cells and the *Opn4::Cre/GFP*

system that labels M1-M6 ipRGCs) and both ages (P5 and adult). Third, we identified whether

the differentially expressed genes have nearly absent gene expression in generic RGC samples

to distinguish potential for selective gene expression in ipRGCs. This was distinguished both at

the level of count-values and manual inspection of aligned raw reads using the Integrated

Genome Viewer (IGV) (Thorvaldsdóttir and Robinson, 2013). Using IGV, we verified that the

reads align with reference gene model for full-length coverage across multiple ipRGC replicates

and that there were absent or partial reads aligned across the generic RGC replicates.

Determining differentially *repressed* genes in adult ipRGCs was confounded by the high

amount of contaminants in generic RGC populations. We could not decipher whether a gene with

relatively low expression in ipRGCs was the result of non-RGC populations contaminating the

generic RGC control population. In contrast, the P5 generic RGC samples from the *Opn4-GFP*

reporter were determined to have greatly reduced levels of contamination and similar levels of

RGC marker expression (Figure 1—figure supplement 1). This made it possible to identify genes

more weakly expressed in ipRGCs than in generic RGCs in early postnatal development (Figure

2—figure supplement 1).

Count-based differential expression pipeline for RNA-seq data using edgeR and/or
DESeq.
# 1) Assess sequence quality control with FastQC)
# 2) remove adapters
$ fastx_clipper -Q33 -a adapter_sequence -l 15 -v -i DJB0005.fastq -o ad_DJB0005.fastq
943
944   # 3) remove low quality reads
945    $ fastq_quality_filter -v -Q33 -q 30 -p 90 -i ad_DJB0005.fastq -o adq_DJB0005.fastq
946
947   ##
948
949   # Align the reads (using tophat2) to the reference genome
950    $ tophat2 --no-coverage-search -o DJB05_th2out genome DJB05_merged.fastq
951
952   ##
953
954   # Sort by name, convert to SAM for htseq-count
955    $ samtools sort -n DJB05_th2out/accepted_hits.bam -o DJB05_sortname.bam
956    $ samtools view -o DJB05_sortname.sam DJB05_sortname.bam
957
958   ##
959
960   # COUNT READS USING HTSEQ-COUNT
961    $ htseq-count -s no -a 10 DJB05_sortname.sam genes.gtf > DJB05.count
962
963   ##
964

```
965    # Count-based differential analysis with edgeR
966    $ module load R
967    $ cd ~/data/2016_rerun_myrnaseq/
968    $ Rscript edgeR_filter1cpm_updated.R GSad_2016.filelist
969
970    ###################################################################
971
972    #Below is script within "edgeR_filter1cpm_updated.R"
973    #!/usr/bin/Rscript
974    #structure of file:1 col of batch names, 2 columns of sample names, with label at top of each
975    column, tab separated
976    #label1label2
977    #b1     samp1 samp4
978    #b2     samp2 samp5
979    #b3     samp3 samp6
980    #samples in same row are assumed to be in same batch
981    args <- commandArgs(TRUE)
982    filename=args[1]
983    bampath="~/data/2016_rerun_myrnaseq"
984    annotation="~/data/2016_rerun_myrnaseq/genes.gtf"
985    baseoutdir="~/data/2016_rerun_myrnaseq/"
986
987    library(edgeR)
988
989    x=read.table(filename,header=T)
990    label1=colnames(x[1])
991    label2=colnames(x[2])
992    samplelist=c(as.vector(x[,1]),as.vector(x[,2]))
993    conditions=c(rep(label1,nrow(x)),rep(label2,nrow(x)))
994    batch=rep(row.names(x),2)
995    names(conditions)=samplelist
996
997    #convert condition and batch to factors--prob not necessary for DESeq but edgeR likes it
998    condition=factor(conditions)
999    batch=factor(batch)
1000
1001   #set up output directory for this experiment, create if it doesn't exist
1002   outdir=sprintf("%s/%s-%s",baseoutdir,label1,label2)
1003   dir.create(outdir)
1004
1005   #read in count table
1006   count.table=read.table(sprintf("%s/%s-%s-rawcounts.txt",outdir,label1,label2),header=T)
1007
1008   # filtering--keep only reads with > 1cpm in at least half the samples
1009   keep_cpm <- rowSums(cpm(count.table)>1) >=nrow(x)
1010   keep_quantile <- rowSums(count.table)>quantile(rowSums(count.table), probs=.5)
1011
1012   #save output of cpm vs quantile filters to log file
1013   sink(sprintf("%s/edgeR.log",outdir),append=T,split=T)
1014   cat("Comparison table of cpm vs quantile filters. CPM>2 in half of samples, quantile at 50%.\n")
1015   addmargins(table(keep_cpm, keep_quantile))
```

```
1016    sink()
1017    count.table <- count.table[keep_cpm, ]
1018
1019    edesign=model.matrix(~batch+condition)
1020    e <- DGEList(counts=count.table)
1021    e <- calcNormFactors(e)
1022    e <- estimateGLMCommonDisp(e, edesign)
1023    e <- estimateGLMTrendedDisp(e, edesign)
1024    e <- estimateGLMTagwiseDisp(e, edesign)
1025
1026    #print size factors to log file
1027    sink(sprintf("%s/edgeR.log",outdir),append=T,split=T)
1028    cat("Normalization factors:\n")
1029    e$samples
1030    sink()
1031    #print dispersion and PCA plots to pdf
1032    pdf(sprintf("%s/%s-%s-edgeRdispersion.pdf",outdir,label1,label2))
1033    plotBCV(e, cex=0.4, main="edgeR: Biological coefficient of variation (BCV) vs abundance")
1034    dev.off()
1035    pdf(sprintf("%s/%s-%s-edgeRPCA.pdf",outdir,label1,label2))
1036    plotMDS(e, main="edgeR MDS Plot")
1037    dev.off()
1038
1039    #Fit curves to GLM
1040    efit <- glmFit(e, edesign)
1041    efit <- glmLRT(efit, coef=sprintf("condition%s",label1))
1042    #make results table and save
1043    stats <- topTags(efit, n=nrow(e))$table
1044    cpms <- cpm(e,)[rownames(stats),normalized.lib.sizes=T]
1045    etable=data.frame(stats,cpms)
1046    etable <- etable[order(etable$FDR), ]
1047    write.table(etable,file=sprintf("%s/%s-%s-edgeR-
1048    filtered.txt",outdir,label1,label2),quote=F,sep="\t")
1049    #copy sample file (argument) to outdir
1050    file.copy(filename,sprintf("%s/%s-%s-samplelist",outdir,label1,label2))
1051
1052    ################################################################
1053
1054    #Below is script within "GSad_2016.filelist"
1055    GSadPos      GSadNeg
1056    b1      DJB07 DJB08
1057    b2      DJB30 DJB31
1058    b3      DJB28 DJB29
1059    b4      DJB16 DJB21
1060    b5      DJB36 DJB37
1061    b6      DJB38 DJB39
1062
1063    ################################################################
1064
1065
```

Antibodies for i*mmunohistochemistry:*

For these studies, the following primary antibodies were used: rabbit anti-melanopsin (Advanced

Targeting Systems; 1:10,000), guinea pig anti-RBPMS (PhosphoSolutions 1832-RBPMS), rabbit

anti-green fluorescent protein (GFP; Invitrogen); Goat anti-Brn3b antibody (Santa Cruz #sc-

6026); mouse anti-Rasgrp1 (Santa Cruz sc-8430); guinea pig anti-Tbx20[1:8500] (Song et al.,

2006)Rabbit anti-Giantin (Abcam ab24586). Secondary antibodies consisted of Alexa Fluor 350,

488, 594 or 647 donkey anti-goat, Alexa Fluor 594 donkey anti-rabbit and Alexa Fluor 594 goat

anti-guinea pig.

1074

Retina Tissue Preparations and Solutions

Mice were euthanized by inhalation of CO_2. Prior to removing the eye, the dorsal margin of the

cornea was marked with a cautery and this was used to guide the placement of a large relieving

cut in the dorsal retina as a subsequent guide to retinal orientation. Eyes were removed

immediately after death and placed in Hibernate-A solution preheated to 37 °C. To keep track of

retinal orientation, the right and left eye were identified and processed separately.

1081

Immunohistochemistry

After the retina was removed from the eye, it was placed on Millipore nitrocellulose paper. The

retinas were fixed for 30 minutes at room temperature using 4% paraformaldehyde freshly

prepared in 0.1M phosphate buffered saline (PBS; pH 7.4). The tissue was then washed for 15

minutes in PBS three times. The tissue was then incubated in a blocking solution of 0.5% Triton-

X and 5% Goat Serum in PBS for two hours at room temperature. The tissue was incubated for

two nights at 4 °C while on a shaker in the primary antibodies diluted in this same blocking

solution. The following day, the samples were washed six times for 20 minutes in 0.1% Tween-

20 in PBS. The tissue was then incubated for two hours in the appropriate Invitrogen or Jackson

labs secondary antibodies diluted 1:1000 in the blocking solution at room temperature. The tissue

was then washed six times for 10 minutes in 0.1% Tween-20 in PBS. The retinas were then

mounted in Aquamount, coverslipped, and sealed with fingernail polish.

For Rasgrp1 immunofluorescence studies, an additional antigen retrieval step was included.

antigen retrieval the tissue was then placed in Tris-EDTA (pH 8.0) for 30 minutes at 80 °C. The

samples were then allowed to return to room temperatures (about 15-30 minutes) before they

were removed from the Tris-EDTA solution and washed three times for 15 minutes in PBS.

1098

Image Acquisition

Immunofluorescent images were captured on a Zeiss Confocal (LSM 510) and Nikon Eclipse

microscope (Micro Video Instruments, Inc. E614, Avon, MA) with a built in Spot Camera

(Diagnostic Instruments, Inc. HRD 100-NIK Sterling Heights, MI). Confocal images were taken

with a 20x objective (Plan Apochromat, WD 0.55 mm) at a resolution of 2048 pixels. To enhance

clarity, image files were pseudocolored and the brightness and contrast was adjusted using

ImageJ 1.47 (National Insistute of Heath, Bethesda, MD). All final images were constructed using

ImageJ and Powerpoint (Microsoft Corporation, Redmond, WA).

1107

Analysis of Rasgrp1 expression in ipRGC subtypes.

Because M1 and M2 cells have the highest levels of melanopsin expression of the ipRGC

population, their dendrites were clearly visible and decipherable with immunostaining. M1 cells

were identified by their dendritic projections to the OFF layer of the inner plexiform layer (IPL).

M2 cells were identified by their dendrites which monostratify the ON sublamina of the IPL (Berson

et al. 2010; Schmidt and Kofuji 2009). M3 cells bistratify the ON and OFF sublamina of the IPL

(Schmidt and Kofuji 2011) and because these cells have similar levels of melanopsin expression

and soma size as M2 cells, it is likely that M3 cells were included in the M2 population quantified

in this study.

M4-M6 cells have the lowest levels of melanopsin-expression of the ipRGC population and their

dendrites were not visible or decipherable with immunostaining. However, at least some members

of the M4/M5/M6 population had lightly immunoreactive somas (Ecker et al., 2010; Quattrochi et

al., 2018; Stabio et al., 2018). M4 cells were identified by their large soma size, low melanopsin

immunodetectability and the lack of dendritic labeling (Estevez et al., 2012). The M5 and M6 cells

we observed were identified by their low levels of melanopsin labeling and M2-sized somas (Ecker

et al., 2010; Quattrochi et al., 2018; Stabio et al., 2018).

1124

Analysis of Tbx20 expression in Cdh3-GFP mice.

ipRGC subtypes were identified using a combination of morphological clues and process of

elimination. In the case of M1 and M2 cells, which express the highest levels of melanopsin,

confocal images of their immunofluorescence reveal dendritic information. As a result, unique

dendritic features distinguish M1 and M2 cells. Cells with dendrites stratifying in the OFF layer of

the IPL were identified as M1 cells, whereas cells stratifying in the ON layer were identified as M2

cells (Berson et al. 2010; Schmidt and Kofuji 2009). Using this method, M3 cells, which stratify

in the ON and OFF sublamina of the IPL, were included in the M1 cell population unless otherwise

noted (Schmidt and Kofuji 2011). As a result of their low levels of melanopsin, M4 and M5/6 cells

are weakly labeled using anti-melanopsin immunohistochemistry with only some of their somata,

but no dendrites, visible (Ecker et al., 2010; Estevez et al., 2012; Quattrochi et al., 2018; Stabio

et al., 2018). Therefore, the method employed for deciphering M1 and M2 cells cannot be used

to identify M4 and M5/6 cells. Instead, M4 cells were categorized by their lack of dendritic labeling,

and their large soma size (Estevez et al., 2012). M5/6 cells were characterized by their lack of

dendritic labeling, M2-sized somata, and process of elimination (Ecker et al., 2010; Estevez et al.,

2012; Quattrochi et al., 2018; Stabio et al., 2018). In other words, cells that were not stained by

the melanopsin antibody (which would detect M1-M3), but were labeled by GFP (labeling M1-

M6), and had small somata, fell into the M5/6 category.

1143

Brain injection of retrobeads into suprachiasmatic nucleus

1144

Adult wild type mice (P30-P60) were anaesthetized with isofluorance and fluorescently labelled

rhodamine latex microspheres (RetroBeads, Lumafluor) were injected into the ipsilateral

suprachiasmatic nucleus to retrogradely label RGCs with axon terminals at the injections site.

Three to five days later, the brain was removed and immediately fixed overnight. The following

day, the brain was rinsed in 0.1M PBS and sectioned at 50um in the coronal plane. The slices

were incubated with DAPI staining for 10 minutes was done in order to provide a reference for

SCN location as indicated by concentrated cellular staining at the SCN. The slices were imaged

for DAPI in UV channel and overlayed with rhodamine channel to identify the injection site in

relation to SCN location. Special attention was paid to ensure that the injection did not extend into

optic nerve and confound results by introducing off-target RGC labeling of fibers of passage.

1155

Brain histology

1156

Animals were sacrificed via transcardial perfusion, and brains were removed and incubated in 4%

paraformaldehyde overnight. Brains were sectioned at 50μm in the coronal plane. To reveal

individual processes in viral tracing experiments, virallyexpressed EYFP was enhanced using

rabbit-anti-gfp (1:1000) and goat-anti-rabbit alexa 488 (1:500). To observe parvalbumin

expression in OPN neurons, parvalbumin immunohistochemistry was performed using mouse-

anti-parvalbumin (1:1000) followed by goat-anti-mouse alexa 488.

Stained and sliced brain slices were mounted on glass cover slides and imaged using a SPOT

RT Slider digital microscope camera mounted to a Nikon (Diagnostic Instruments) as described

previously (Berson et al., 2010; Estevez et al., 2012). Images were assembled in Adobe

Photoshop CS3.

1167

1168

1169

1170

1171

1172

1173   **Figure legends (see attached PDF for figure images)**

1174

Figure 1. Experimental design of gene expression profiling from purified ipRGCs and

comparison with generic RGCs.

A. Two transgenic reporters were used for gene expression profiling of ipRGCs. The BAC

transgenic Opn4-GFP labels M1-M3 ipRGCs while the Opn4-Cre crossed with a cre-dependent

GFP reporter labels M1-M6 ipRGCs. B. Schematic of the gene expression profiling procedure. 1)

Isolation of cell populations from enzymatically dissociated retinas. 2) The surface protein Thy-1

is enriched in RGCs, this high affinity of Thy1-conjugated magnetic beads to RGCs was used to

enrich the extracted cell populations with RGCs. 3) Fluorescence-activated cell sorting (FACS)

was used to isolate GFP-positive cells (ipRGCs) from GFP-negative cells (cRGCs). These two

populations were isolated in parallel to provide direct internal testing of ipRGCs versus cRGCs

under the same treatments, conditions, and genetic backgrounds. 4) The RNA of these two main

populations was subjected to mRNA extraction. 5) The RNA was converted to cDNA and amplified

using Nugen Ovation RNA amplification system. 6) Illumina Truseq sequencing libraries were

prepared by ligating adapters to the cDNA. Single-end 50 base pair sequencing was completed

using the Illumina HiSeq system. 7) Differentially expressed genes were determined using EdgeR

bioinformatics pipeline. See Methods for details.

B. EdgeR multi-dimensional scaling (MDS) plot illustrates the overall similarity between

expression profiles of different samples. Each sample is denoted by a letter ("i" for ipRGCs; c for

cRGCs) and a number, corresponding to particular replicate, comprising one pool of purified

RGCs then divided into the two pools. Numbering scheme represents paired ipRGCs (GFP+) and

cRGCs (GFP-) replicates (i.e., ipRGC sample 'i1' was processed in parallel with cRGC sample

'c1', sample 'i2' with 'c2', etc.). Distances are approximately the log2 fold changes between

samples. Green and gray ovals represent ipRGC (GFP+) and cRGC (GFP-) samples,

respectively. Adapted from EdgeR simple graphical output of individual samples in 2D space.

1199

Figure 1—figure supplement 1. Purity and cell composition assessment of ipRGC and

generic RGC samples.

Heat map of known cell type marker gene expression in the retina to assess purity and cell

composition of ipRGC and generic RGC samples. Shown are biological replicates tested for

Opn4-GFP (P5 and adult) and *Opn4-Cre/GFP* reporters. Relative expression levels, fold change,

and false discovery rate (FDR) are color-coded as indicated in the figure. White boxes indicate

high gene expression, while blue represents little or no detected expression. FDR is not available

("NA") in cases that our analysis filtered out genes with very low counts, less than 1 count per

million (cpm), in more than half of the samples used in the differential expression analysis.

1209

1210   **Figure 2. The expression pattern of candidate ipRGC-specific genes.**

1211   Heat map of 83 genes differentially expressed in ipRGCs that have functional links to GPCR

1212   signaling, regulation and maintenance of molecular programs, neuron communication and

1213   organization, neuron survival, and neuron-glia interactions. Relative expression levels, fold

1214   change, and FDR are color-coded as indicated in the figure.

1215

1216   **Figure 2—figure supplement 1.**

1217   Heat map of genes differentially expressed in adult ipRGCs labeled by the *Opn4-Cre/GFP*

1218   reporter (M1-M6 ipRGCs) compared to *Opn4-GFP* (M1-M3 ipRGCs). Relative expression levels,

1219   fold change, and FDR are color-coded as indicated in the figure.

1220

1221   **Figure 2—figure supplement 2.**

1222   Heat map of genes encoding for nicotinic acetylcholine, dopamine, serotonin, glycine, glutamate,

1223   and melatonin receptors. Relative expression levels, fold change, and FDR are color-coded as

1224   indicated in the figure.

1225

Figure 2—figure supplement 3. The expression pattern of developmentally regulated genes

in ipRGCs.

A. Heat map of genes encoding transcription factors that have a particular temporal pattern of

differential expression in ipRGCs (e.g., high gene expression in P5 ipRGCs relative to adult

expression). Relative expression levels, fold change, and FDR are color-coded as indicated in

the figure.

B. Heat map of genes relevant for development of ipRGCs. Relative expression levels, fold

change, and FDR are color-coded as indicated in the figure.

1234

Figure 2—figure supplement 4.

A. Distinct from rod and cone photoreceptors, the light-activation of Opn4 triggers a membrane-

bound signaling cascade including $G_{q/11}$ type G-proteins, the generation of 1,2-diacylglycerol

(DAG) by PLCβ4, the opening of downstream TRPC6 and TRPC7 channels, and ultimately leads

to the influx of calcium through L-type voltage-gated calcium channels.

B. Heat map of genes that are potentially relevant to the Opn4-mediated phototransduction

signaling cascade. Relative expression levels, fold change, and FDR are color-coded as indicated

in the figure.

C. Heat map of genes previously described to play a role in the light response, dark adaptation,

and chromophore regeneration of rod and cone photoreceptors. Relative expression levels, fold

change, and FDR are color-coded as indicated in the figure.

1246

Figure 3. Rasgrp1 is selectively expressed in ipRGCs.

A. Whole mount retina immunostained for Opn4, Rasgrp1, and the pan-RGC marker Rbpms

(gray-scale). Focal plane is at the ganglion cell layer (GCL). We quantified co-localization of the

three markers in confocal images of 49 regions that were topographically dispersed across three

whole-mount adult retinas.

B. Co-localization of Rasgrp1 (green), Rbpms (red), and Opn4 (magenta). Rasgrp1 is expressed

in a subpopulation of amacrine cells and RGCs (Rbpms-negative and -positive, respectively).

Scale bar, 20 μm.

C. Rasgrp1 immunolabeling (antibody sc-8430) of cell bodies in GCL of Rasgrp1$^{+/-}$ heterozygous

mice (left, yellow arrows). Absence of cell body immunolabeling in Rasgrp1$^{-/-}$ knockout mice (right)

suggests a lack of cellular off-target antibody staining.

D. Quantification of Rasgrp1-expression across Opn4-immunopositive ipRGC subtypes. 70% of

M1 cells were Rasgrp1 immunopositive while only 20-30% of either M2, M5 or M6 cells were

Rasgrp1 immunopositive. None of the identified M4 cells were Rasgrp1 immunopositive. M1 and

M3 types were combined during the process of co-expression analysis (designated "M1/M3").

Error bars represent standard error of the mean.

1263

Figure 3—figure supplement 1.

A. Distribution of all Rasgrp1-expressing RGCs (Rasgrp1^{+};Rbpms^{+}) that belong to specific RGC

types, to the extent that could be determined, including Opn4-immunoreactive ipRGC subtypes.

No examples of M4 cells were observed to express Rasgrp1. The vast majority (96%) of Rasgrp1-

RGCs are Opn4-immunopositive and therefore ipRGCs. The remaining "unknown" RGC types

expressing Rasgrp1 (Rasgrp1^{+} ; Rbpms^{+} ; Opn4$^{neg.}$) could be a low-expressing ipRGC type or

conventional RGCs. Error bars represent standard error of the mean.

B. Areas sampled for two wholemount wild type retinas immunostained for Rasgrp1, Opn4 and

RBPMS. Yellow squares represent the locations of confocal images used for cell quantification.

C. Analysis of the M1-M3 ipRGC population did not suggest a gradient of ipRGC spatial

distribution across the retina. Seven frames were used to represent each area of the retina to

maintain equal spatial contribution. Total M1-M3 represented in the dorsal-ventral and nasal-

temporal columns is 217.

E. Analysis of the M1 cell population revealed a slight naso-temporal gradient of M1 spatial

distribution across the retina. Seven frames were used to represent each area of the retina to

maintain equal spatial contribution. Total M1s represented in the dorsal-ventral column and nasal-

temporal columns is 113 and 110, respectively.

F. Analysis of the Rasgrp1-positive RGC population suggests a slight ventral-dorsal gradient of

Rasgrp1-positive RGCs spatial distribution across the retina. Seven frames were used to

represent each area of the retina to maintain equal spatial contribution. Total Rasgrp1-positive

RGCs represented in the dorsal-ventral column and nasal-temporal column is 117 and 118,

respectively.

F. The amacrine cells (presumed) expressing Rasgrp1 exhibited a dramatic center-peripheral

gradient of Rasgrp1-positive non-RGC spatial distribution across the retina. Four frames were

used to represent each area of the retina to maintain equal spatial contribution. The total Rasgrp1-

positive non-RGCs represented in the periphery and center was 314.

1290

Figure 4. Colocalization study of Tbx20-expression in ipRGC subtypes.

A. Triple immunofluorescence of Opn4, Tbx20, and *Cdh3-GFP* (gray scale).

B. Tbx20-expression in subset of M1-M3 ipRGCs as well as an additional population of Opn4-

immunonegative cells.

B,C. Tbx20 is concentrated in the nucleus of most *Cdh3-GFP*-cells.

D. Quantification of Tbx20-expression across Opn4-immunopositive ipRGC subtypes. Tbx20

immunofluorescence labels multiple ipRGC subtypes, including M1s, M2 cells and small soma,

low Opn4 expression cells (presumptive M5/6 ipRGCs), and Cdh3-GFP cells (M6-type enriched).

M2 and M3 types were combined during the process of co-expression analysis (designated

"M2/M3"). Error bars represent standard error of the mean.

1301

Figure 4—figure supplement 1. Coexpression study of Tbx20 with Opn4-Cre/GFP and Tbr2,

including distribution of Tbx20-expression across ipRGC subtypes.

A-C. Quadruple immunofluorescence of Opn4, Tbx20, Opn4-Cre/GFP, and Tbr2. Scale bar, 20

μm.

A. Gray scale of Opn4, Opn4-Cre/GFP, and Tbx20 immunofluorescence.

B. Co-expression study of Tbx20 (green) in the context of Opn4 (magenta) and *Opn4-Cre/GFP*

(red) labeling. GFP cells that are Opn4-immunonegative are inferred M4-M6 types.

C. Co-expression analysis of Tbr2 (magenta) with Tbx20 (green).

D. Distribution of Tbx20 expressing cells that belong to specific RGC types, to the extent that

could be determined, including Opn4-immunoreactive ipRGC subtypes and RGCs labeled by the

Cdh3-GFP transgenic reporter. Unaccounted Tbx20-expressing cells are designated as

"unknown" RGC types. Error bars represent standard error of the mean.

1314

1315

Figure 5. Complex pattern of Rasgrp1-Tbx20-Brn3b co-expression suggests further

diversity in ipRGC family.

A. Quadruple immunofluorescence study of Tbx20, Brn3b, Opn4, and Rasgrp1 (gray-scale).

B. Rasgrp1 and Opn4 (left panel) were initially quantified for ipRGC subtype expression prior to

comparison with Tbx20 (middle panel) and Brn3b (right panel) expression. Rasgrp1, Brn3b, and

Tbx20 expression are partially overlapping.

C. Integrated co-expression patterns of Brn3b, Rasgrp1, and Tbx20 with M1 and M2 ipRGC

subtypes. The M1 group includes displaced M1 and M3 types.

1324

Figure 5—figure supplement 1. Topographic distribution and ipRGC subtype-specific

quantification of Rasgrp1-Tbx20-Brn3b expression.

A. Comparison of co-expression patterns of Brn3b, Rasgrp1, and Tbx20 within group of combined

M1-M3 (M1+M2+M3) ipRGCs, the M1/3 ipRGCs (M1+M3), displaced M1s, and M2 ipRGCs.

B. Lack of major topographic variations in the fraction of M1/M3 ipRGCs immunoreactive for

Rasgrp1, Tbx20, and Brn3b.

C. Comparison of Rasgrp1-Tbx20-Brn3b expression pattern in M2 ipRGCs across dorsal, center,

and ventral regions of the retina.

D. Distribution of Tbx20-expression in M1-M3 ipRGCs compared to Opn4-immunonegative cells

in the context of topgraphic regions across the retina.

1335

Figure 6. The ipRGCs projecting to the suprachiasmatic nucleus (SCN) have a molecularly

diverse pattern of Rasgrp1 and Tbx20 expression.

A. Experimental design of fluorescent bead injection to SCN, followed by examination of Rasgrp1

and Tbx20 expression in retrograde labeled RGCs.

B. Neuroannatomical study to verify that retrograde injection is within the SCN, but not the optic

nerve.

B. Triple immunofluorescence of Opn4, Rasgrp1, and Tbx20 in combination with fluorescent

Retrobeads. Retrobeads were mostly observed in Opn4-immunopositive RGCs (M1-M3 ipRGCs).

Quantification of Rasgrp1 and Tbx20 in retrolabeled cells.

D. SCN-projecting ipRGCs in the ipsilateral and contralateral retina are molecularly diverse for

Tbx20 and Rasgrp1 expression.

1347

1348 **Figure 7. Current model of ipRGC family members integrating molecular, physiology, brain**

1349 **circuitry, and morphology (see text for details).**

1350

Figure 8. Coexpression of BAC transgenic Opn4-GFP reporter with Opn4

immunoreactivity.

A. Immunofluorescence of anti-Opn4 staining of whole mount retina from transgenic *Opn4-GFP*

mice with fluorescent protein expression in ipRGCs. Red, Opn4-immunolabeling; green,

fluorescently labeled cells; yellow, merged co-localized labeling pattern. Scale bar, 20 μm.

B. Co-expression of *Opn4-Cre/GFP* labeling with immunofluorescence of anti-Opn4 staining of

whole mount retina. Red, Opn4-immunolabeling; green, fluorescently labeled cells; yellow,

merged co-localized labeling pattern. Scale bar, 20 μm.

C. Quantification of labeling efficiency of Opn4-immunolabeled M1-M3 ipRGCs by *Opn4-*

Cre/GFP. Additional comparison of GFP-labeling in low Opn4-expressing ipRGC subtypes M4

(large soma) and M5/6 (small soma).

1362

Figure 9. Fluorescence activated cell sorting (FACS) gating strategy for isolating ipRGCs

(GFP+) in parallel with GFP-negative cells that are enriched for RGCs.

A-C. Healthy cells were selected against death marker 7-AAD (not G1). The ipRGCs (GFP$^+$) and

generic RGCs (GFP$^-$) cells were selected based on intensity and similar relative cell size ultimately

using gates G2A and G3A, respectively.

A. Example sort from retina of postnatal day 4 (P4) *Opn4-GFP* mouse.

C. Example sort from retina of young adult *Opn4-GFP* mouse, with noticeably higher debri and

cell death.

D. Microscopy testing of accurate sorting of GFP+ cells isolated from P4 *Opn4-Cre/GFP* mouse.

1372

1373 **Figure 10. Steps involved with processing mRNA extracted from purified cell populations**

1374 **and preparing for RNA-sequencing (see Methods for details).**

1375

1376 <u>**References**</u>

Anders S, Huber W (2010) Differential expression analysis for sequence count data. Genome Biology 2010 11:10 11:R106.

Anders S, McCarthy DJ, Chen Y, Okoniewski M, Smyth GK, Huber W, Robinson MD (2013) Count-based differential expression analysis of RNA sequencing data using R and Bioconductor. Nature Protocols 8:1765–1786.

Atlasz T, Szabadfi K, Kiss P, Racz B, Gallyas F, Tamas A, Gaal V, Marton Z, Gabriel R, Reglodi D (2010) Pituitary adenylate cyclase activating polypeptide in the retina: focus on the retinoprotective effects. Ann N Y Acad Sci 1200:128–139.

Barres BA, Silverstein BE, Corey DP, Chun LLY (1988) Immunological, morphological, and electrophysiological variation among retinal ganglion cells purified by panning. Neuron 1:791–803.

Bedont JL, Blackshaw S (2015) Constructing the suprachiasmatic nucleus: a watchmaker's perspective on the central clockworks. Front Syst Neurosci 9:74.

Beglopoulos V, Montag-Sallaz M, Rohlmann A, Piechotta K, Ahmad M, Montag D, Missler M (2005) Neurexophilin 3 is highly localized in cortical and cerebellar regions and is functionally important for sensorimotor gating and motor coordination. Mol Cell Biol 25:7278–7288.

Berson DM, Castrucci AM, Provencio I (2010) Morphology and mosaics of melanopsin-expressing retinal ganglion cell types in mice. Journal of Comparative Neurology 518:2405–2422.

Bivona TG, Pérez De Castro I, Ahearn IM, Grana TM, Chiu VK, Lockyer PJ, Cullen PJ, Pellicer A, Cox AD, Philips MR (2003) Phospholipase Cgamma activates Ras on the Golgi apparatus by means of RasGRP1. Nature 424:694–698.

Brewer GJ (1997) Isolation and culture of adult rat hippocampal neurons. Journal of Neuroscience Methods 71:143–155.

Brewer GJ, Torricelli JR (2007) Isolation and culture of adult neurons and neurospheres. Nature Protocols 2:1490–1498.

Brewer GJ, Torricelli JR, Evege EK, Price PJ (1993) Optimized survival of hippocampal neurons in B27-supplemented neurobasal™, a new serum-free medium combination. Journal of Neuroscience Research 35:567–576.

Burden-Gulley SM, Brady-Kalnay SM (1999) PTPmu regulates N-cadherin-dependent neurite outgrowth. J Cell Biol 144:1323–1336.

Cahoy JD, Emery B, Kaushal A, Foo LC, Zamanian JL, Christopherson KS, Xing Y, Lubischer JL, Krieg PA, Krupenko SA, Thompson WJ, Barres BA (2008) A transcriptome database for astrocytes, neurons, and oligodendrocytes: a new resource for understanding brain development and function. J Neurosci 28:264–278.

Cai C-L, Zhou W, Yang L, Bu L, Qyang Y, Zhang X, Li X, Rosenfeld MG, Chen J, Evans S (2005) T-box genes coordinate regional rates of proliferation and regional specification during cardiogenesis. Development 132:2475–2487.

Cameron EG, Robinson PR (2014) β-Arrestin-Dependent Deactivation of Mouse Melanopsin Craft CM, ed. PLoS ONE 9:e113138.

Caretti E, Devarajan K, Coudry R, Ross E, Clapper ML, Cooper HS, Bellacosa A (2008) Comparison of RNA amplification methods and chip platforms for microarray analysis of samples processed by laser capture microdissection. Journal of Cellular Biochemistry 103:556–563.

Carson CT, Kinzler ER, Parr BA (2000) Tbx12, a novel T-box gene, is expressed during early stages of heart and retinal development. Mechanisms of Development 96:137–140.

Carson CT, Pagratis M, Parr BA (2004) Tbx12 regulates eye development in Xenopus embryos. Biochemical and Biophysical Research Communications 318:485–489.

Chen SK, Badea TC, Hattar S (2011) Photoentrainment and pupillary light reflex are mediated by distinct populations of ipRGCs. Nature 476:92–95.

Clément-Ziza M, Gentien D, Lyonnet S, Thiery J-P, Besmond C, Decraene C (2009) Evaluation of methods for amplification of picogram amounts of total RNA for whole genome expression profiling. BMC Genomics 10:246.

Craig AM, Kang Y (2007) Neurexin-neuroligin signaling in synapse development. Current Opinion in Neurobiology 17:43–52.

Cui Q, Ren C, Sollars PJ, Pickard GE, So KF (2015) The injury resistant ability of melanopsin-expressing intrinsically photosensitive retinal ganglion cells. Neuroscience 284:845–853.

de Sevilla Müller LP, Sargoy A, Rodriguez AR, Brecha NC (2014) Melanopsin Ganglion Cells Are the Most Resistant Retinal Ganglion Cell Type to Axonal Injury in the Rat Retina Tosini G, ed. PLoS ONE 9:e93274.

de Wit J, Sylwestrak E, O'Sullivan ML, Otto S, Tiglio K, Savas JN, Yates JR, Comoletti D, Taylor P, Ghosh A (2009) LRRTM2 interacts with Neurexin1 and regulates excitatory synapse formation. Neuron 64:799–806.

Diaz A, Ruiz F, Flórez J, Hurlé MA, Pazos A (1995) Mu-opioid receptor regulation during opioid tolerance and supersensitivity in rat central nervous system. J Pharmacol Exp Ther 274:1545–1551.

Doğrul A, Yeşilyurt O, Işimer A, Güzeldemir ME (2001) L-type and T-type calcium channel blockade potentiate the analgesic effects of morphine and selective mu opioid agonist, but not to selective delta and kappa agonist at the level of the spinal cord in mice. Pain 93:61–68.

Dower NA, Stang SL, Bottorff DA, Ebinu JO, Dickie P, Ostergaard HL, Stone JC (2000) RasGRP is essential for mouse thymocyte differentiation and TCR signaling. Nat Immunol 1:317–321.

Duan X, Qiao M, Bei F, Kim I-J, He Z, Sanes JR (2015) Subtype-Specific Regeneration of
Retinal Ganglion Cells following Axotomy: Effects of Osteopontin and mTOR Signaling.
Neuron 85:1244–1256.

Dumitrescu ON, Pucci FG, Wong KY, Berson DM (2009) Ectopic retinal ON bipolar cell
synapses in the OFF inner plexiform layer: Contacts with dopaminergic amacrine cells and
melanopsin ganglion cells. Journal of Comparative Neurology 517:226–244.

Ebinu JO, Bottorff DA, Chan EY, Stang SL, Dunn RJ, Stone JC (1998) RasGRP, a Ras guanyl
nucleotide- releasing protein with calcium- and diacylglycerol-binding motifs. Science
280:1082–1086.

Ecker JL, Dumitrescu ON, Wong KY, Alam NM, Chen S-K, LeGates T, Renna JM, Prusky GT,
Berson DM, Hattar S (2010) Melanopsin-Expressing Retinal Ganglion-Cell Photoreceptors:
Cellular Diversity and Role in Pattern Vision. Neuron 67:49–60.

Eide EJ, Woolf MF, Kang H, Woolf P, Hurst W, Camacho F, Vielhaber EL, Giovanni A, Virshup
DM (2005) Control of mammalian circadian rhythm by CKIepsilon-regulated proteasome-
mediated PER2 degradation. Mol Cell Biol 25:2795–2807.

Emanuel AJ, Do MTH (2015) Melanopsin Tristability for Sustained and Broadband
Phototransduction. Neuron 85:1043–1055.

Emanuel AJ, Kapur K, Do MTH (2017) Biophysical Variation within the M1 Type of Ganglion
Cell Photoreceptor. Cell Rep 21:1048–1062.

Estevez ME, Fogerson PM, Ilardi MC, Borghuis BG, Chan E, Weng S, Auferkorte ON, Demb
JB, Berson DM (2012) Form and function of the M4 cell, an intrinsically photosensitive
retinal ganglion cell type contributing to geniculocortical vision. J Neurosci 32:13608–13620.

Fernandez DC, Chang Y-T, Hattar S, Chen S-K (2016) Architecture of retinal projections to the
central circadian pacemaker. Proc Natl Acad Sci USA 113:6047–6052.

Fink M, Lesage F, Duprat F, Heurteaux C, Reyes R, Fosset M, Lazdunski M (1998) A neuronal
two P domain K+ channel stimulated by arachidonic acid and polyunsaturated fatty acids.
EMBO J 17:3297–3308.

Fornaro M, Raimondo S, Lee JM, Giuseppina Giacobini-Robecchi M (2007) Neuron-specific Hu
proteins sub-cellular localization in primary sensory neurons. Annals of Anatomy -
Anatomischer Anzeiger 189:223–228.

Frings S, Brüll N, Dzeja C, Angele A, Hagen V, Kaupp UB, Baumann A (1998) Characterization
of ether-à-go-go channels present in photoreceptors reveals similarity to IKx, a K+ current in
rod inner segments. J Gen Physiol 111:583–599.

Gorentla BK, Wan C-K, Zhong X-P (2011) Negative regulation of mTOR activation by
diacylglycerol kinases. Blood 117:4022–4031.

Graham DM, Wong KY, Shapiro P, Frederick C, Pattabiraman K, Berson DM (2008) Melanopsin
Ganglion Cells Use a Membrane-Associated Rhabdomeric Phototransduction Cascade.
Journal of Neurophysiology 99:2522–2532.

Haeryfar SMM, Hoskin DW (2004) Thy-1: More than a Mouse Pan-T Cell Marker. The Journal of
Immunology 173:3581–3588.

Hannibal J, Hindersson P, Østergaard J, Georg B, Heegaard S, Larsen PJ, Fahrenkrug J (2004)
Melanopsin Is Expressed in PACAP-Containing Retinal Ganglion Cells of the Human
Retinohypothalamic Tract. Invest Ophthalmol Vis Sci 45:4202–4209.

Hartwick ATE, Bramley JR, Yu J, Stevens KT, Allen CN, Baldridge WH, Sollars PJ, Pickard GE
(2007) Light-evoked calcium responses of isolated melanopsin-expressing retinal ganglion
cells. J Neurosci 27:13468–13480.

Hattar S, Kumar M, Park A, Tong P, Tung J, Yau K-W, Berson DM (2006) Central projections of
melanopsin-expressing retinal ganglion cells in the mouse. J Comp Neurol 497:326–349.

Heiman M, Schaefer A, Gong S, Peterson JD, Day M, Ramsey KE, Suárez-Fariñas M, Schwarz
C, Stephan DA, Surmeier DJ, Greengard P, Heintz N (2008) A Translational Profiling
Approach for the Molecular Characterization of CNS Cell Types. Cell 135:738–748.

Hinman MN, Lou H (2008) Diverse molecular functions of Hu proteins. Cell Mol Life Sci
65:3168–3181.

Hoshi H, Liu W-L, Massey SC, Mills SL (2009) ON inputs to the OFF layer: bipolar cells that
break the stratification rules of the retina. J Neurosci 29:8875–8883.

Hu C, Hill DD, Wong KY (2013) Intrinsic physiological properties of the five types of mouse
ganglion-cell photoreceptors. Journal of Neurophysiology 109:1876–1889.

Hughes S, Hankins MW, Foster RG, Peirson SN (2012) Melanopsin phototransduction: slowly
emerging from the dark. Prog Brain Res 199:19–40.

Hughes S, Jagannath A, Hickey D, Gatti S, Wood M, Peirson SN, Foster RG, Hankins MW
(2015) Using siRNA to define functional interactions between melanopsin and multiple G
Protein partners. Cell Mol Life Sci 72:165–179.

Jain V, Ravindran E, Dhingra NK (2012) Differential expression of Brn3 transcription factors in
intrinsically photosensitive retinal ganglion cells in mouse. Journal of Comparative
Neurology 520:742–755.

Jeon C-J, Strettoi E, Masland RH (1998) The Major Cell Populations of the Mouse Retina. J
Neurosci 18:8936–8946.

Ji M, Zhao W-J, Dong L-D, Miao Y, Yang X-L, Sun X-H, Wang Z (2011) RGS2 and RGS4
modulate melatonin-induced potentiation of glycine currents in rat retinal ganglion cells.
Brain Research 1411:1–8.

Kim I-J, Zhang Y, Meister M, Sanes JR (2010) Laminar restriction of retinal ganglion cell
dendrites and axons: subtype-specific developmental patterns revealed with transgenic
markers. J Neurosci 30:1452–1462.

Koussounadis A, Langdon SP, Um IH, Harrison DJ, Smith VA (2015) Relationship between
differentially expressed mRNA and mRNA-protein correlations in a xenograft model system.
Sci Rep 5:10775.

Li S, Yang C, Zhang L, Gao X, Wang X, Liu W, Wang Y, Jiang S, Wong YH, Zhang Y, Liu K
(2016) Promoting axon regeneration in the adult CNS by modulation of the
melanopsin/GPCR signaling. Proc Natl Acad Sci USA 113:1937–1942.

Li S-Y, Yau S-Y, Chen B-Y, Tay DK, Lee VWH, Pu M-L, Chan HHL, So K-F (2008) Enhanced
Survival of Melanopsin-expressing Retinal Ganglion Cells After Injury is Associated with the
PI3 K/Akt Pathway. Cell Mol Neurobiol 28:1095–1107.

Lin JC, Ho W-H, Gurney A, Rosenthal A (2003) The netrin-G1 ligand NGL-1 promotes the
outgrowth of thalamocortical axons. Nature Neuroscience 6:1270–1276.

Lin Z, Liu J, Ding H, Xu F, Liu H (2018) Structural basis of SALM5-induced PTPδ dimerization
for synaptic differentiation. Nature Communications 9:268.

Lipina TV, Prasad T, Yokomaku D, Luo L, Connor SA, Kawabe H, Wang YT, Brose N, Roder
JC, Craig AM (2016) Cognitive Deficits in Calsyntenin-2-deficient Mice Associated with
Reduced GABAergic Transmission. Neuropsychopharmacology 41:802–810.

Lobo MK, Karsten SL, Gray M, Geschwind DH, Yang XW (2006) FACS-array profiling of striatal
projection neuron subtypes in juvenile and adult mouse brains. Nature Neuroscience 9:443–
452.

Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, Tirosh I, Bialas AR,
Kamitaki N, Martersteck EM, Trombetta JJ, Weitz DA, Sanes JR, Shalek AK, Regev A,
McCarroll SA (2015) Highly Parallel Genome-wide Expression Profiling of Individual Cells
Using Nanoliter Droplets. Cell 161:1202–1214.

Mao C-A, Kiyama T, Pan P, Furuta Y, Hadjantonakis A-K, Klein WH (2008) Eomesodermin, a
target gene of Pou4f2, is required for retinal ganglion cell and optic nerve development in
the mouse. Development 135:271–280.

Mao C-A, Li H, Zhang Z, Kiyama T, Panda S, Hattar S, Ribelayga CP, Mills SL, Wang SW
(2014) T-box transcription regulator Tbr2 is essential for the formation and maintenance of
Opn4/melanopsin-expressing intrinsically photosensitive retinal ganglion cells. J Neurosci
34:13083–13095.

Mao H, Zhao Q, Daigle M, Ghahremani MH, Chidiac P, Albert PR (2004) RGS17/RGSZ2, a
novel regulator of Gi/o, Gz, and Gq signaling. J Biol Chem 279:26314–26322.

Martin S, Lino de Oliveira C, Mello de Queiroz F, Pardo LA, Stühmer W, Del Bel E (2008) Eag1
potassium channel immunohistochemistry in the CNS of adult rat and selected regions of
human brain. Neuroscience 155:833–844.

Matsuoka RL, Nguyen-Ba-Charvet KT, Parray A, Badea TC, Chédotal A, Kolodkin AL (2011)
Transmembrane semaphorin signalling controls laminar stratification in the mammalian
retina. Nature 470:259–263.

Matsushima D, Heavner W, Pevny LH (2011) Combinatorial regulation of optic cup progenitor
cell fate by SOX2 and PAX6. Development 138:443–454.

Missler M, Hammer RE, Südhof TC (1998) Neurexophilin binding to alpha-neurexins. A single
LNS domain functions as an independently folding ligand-binding unit. J Biol Chem
273:34716–34723.

Moises HC, Rusin KI, Macdonald RL (1994) Mu- and kappa-opioid receptors selectively reduce
the same transient components of high-threshold calcium current in rat dorsal root ganglion
sensory neurons. J Neurosci 14:5903–5916.

Morrow EM, Belliveau MJ, Cepko CL (1998) Two Phases of Rod Photoreceptor Differentiation
during Rat Retinal Development. J Neurosci 18:3738–3748.

Morse AM, Carballo V, Baldwin DA, Taylor CG, McIntyre LM (2010) Comparison between
NuGEN's WT-Ovation Pico and one-direct amplification systems. J Biomol Tech 21:141–
147.

Naiche LA, Harrelson Z, Kelly RG, Papaioannou VE (2005) T-box genes in vertebrate
development. Annu Rev Genet 39:219–239.

Novak A, Guo C, Yang W, Nagy A, Lobe CG (2000) Z/EG, a double reporter mouse line that
expresses enhanced green fluorescent protein upon Cre-mediated excision. Genesis.

Oancea E, Meyer T (1998) Protein kinase C as a molecular machine for decoding calcium and
diacylglycerol signals. Cell 95:307–318.

Pack W, Hill DD, Wong KY (2015) Melatonin modulates M4-type ganglion-cell photoreceptors.
Neuroscience 303:178–188.

Pederick DT, Homan CC, Jaehne EJ, Piltz SG, Haines BP, Baune BT, Jolly LA, Hughes JN,
Gecz J, Thomas PQ (2016) Pcdh19 Loss-of-Function Increases Neuronal Migration In Vitro
but is Dispensable for Brain Development in Mice. Sci Rep 6:26765.

Peirson SN, Oster H, Jones SL, Leitges M, Hankins MW, Foster RG (2007) Microarray Analysis
and Functional Genomics Identify Novel Components of Melanopsin Signaling. Current
Biology 17:1363–1372.

Pettem KL, Yokomaku D, Takahashi H, Ge Y, Craig AM (2013) Interaction between autism-
linked MDGAs and neuroligins suppresses inhibitory synapse development. J Cell Biol
200:321–336.

Pierret P, Dunn RJ, Djordjevic B, Stone JC, Richardson PM (2000) Distribution of ras guanyl
releasing protein (RasGRP) mRNA in the adult rat central nervous system. J Neurocytol
29:485–497.

Pocock R, Mione M, Hussain S, Maxwell S, Pontecorvi M, Aslam S, Gerrelli D, Sowden JC,
Woollard A (2008) Neuronal function of Tbx20 conserved from nematodes to vertebrates.
Developmental Biology 317:671–685.

Puente LG, Stone JC, Ostergaard HL (2000) Evidence for protein kinase C-dependent and -
independent activation of mitogen-activated protein kinase in T cells: potential role of
additional diacylglycerol binding proteins. The Journal of Immunology 165:6865–6871.

Quail MA, Kozarewa I, Smith F, Scally A, Stephens PJ, Durbin R, Swerdlow H, Turner DJ
(2008) A large genome center's improvements to the Illumina sequencing system. Nat
Methods 5:1005–1010.

Quattrochi LE, Stabio ME, Kim I, Berson DM (2018) The M6 cell: A small-field bistratified
photosensitive ganglion cell. in press.

Reifler AN, Chervenak AP, Dolikian ME, Benenati BA, Li BY, Wachter RD, Lynch AM,
Demertzis ZD, Meyers BS, Abufarha FS, Jaeckel ER, Flannery MP, Wong KY (2015) All
Spiking, Sustained ON Displaced Amacrine Cells Receive Gap-Junction Input from
Melanopsin Ganglion Cells. Current Biology 25:2763–2773.

Renna JM, Weng S, Berson DM (2011) Light acts through melanopsin to alter retinal waves and
segregation of retinogeniculate afferents. Nature Neuroscience 14:827–829.

Rodriguez AR, Sevilla Müller LP, Brecha NC (2014) The RNA binding protein RBPMS is a
selective marker of ganglion cells in the mammalian retina. Journal of Comparative
Neurology 522:1411–1443.

Rousso DL, Qiao M, Kagan RD, Yamagata M, Palmiter RD, Sanes JR (2016) Two Pairs of ON
and OFF Retinal Ganglion Cells Are Defined by Intersectional Patterns of Transcription
Factor Expression. Cell Rep 15:1930–1944.

Sabbah S, Berg D, Papendorp C, Briggman KL, Berson DM (2017) A Cre Mouse Line for
Probing Irradiance- and Direction-Encoding Retinal Networks. eNeuro 4:ENEURO.0065–
17.2017.

Sakabe NJ, Aneas I, Shen T, Shokri L, Park S-Y, Bulyk ML, Evans SM, Nobrega MA (2012)
Dual transcriptional activator and repressor roles of TBX20 regulate adult cardiac structure
and function. Hum Mol Genet 21:2194–2204.

Sand A, Schmidt TM, Kofuji P (2012) Diverse types of ganglion cell photoreceptors in the
mammalian retina. Progress in Retinal and Eye Research 31:287–302.

Sanes JR, Masland RH (2015) The Types of Retinal Ganglion Cells: Current Status and
Implications for Neuronal Classification. http://dxdoiorg/101146/annurev-neuro-071714-
034120 38:221–246.

Schmidt TM, Do MTH, Dacey D, Lucas R, Hattar S, Matynia A (2011) Melanopsin-positive
intrinsically photosensitive retinal ganglion cells: from form to function. J Neurosci
31:16094–16101.

Schmidt TM, Taniguchi K, Kofuji P (2008) Intrinsic and Extrinsic Light Responses in
Melanopsin-Expressing Ganglion Cells During Mouse Development. Journal of
Neurophysiology 100:371–384.

Sengupta A, Baba K, Mazzoni F, Pozdeyev NV, Strettoi E, Iuvone PM, Tosini G (2011)
Localization of melatonin receptor 1 in mouse retina and its role in the circadian regulation
of the electroretinogram and dopamine levels. Yamazaki S, ed. PLoS ONE 6:e24483.

Shekhar K, Lapan SW, Whitney IE, Tran NM, Macosko EZ, Kowalczyk M, Adiconis X, Levin JZ,
Nemesh J, Goldman M, McCarroll SA, Cepko CL, Regev A, Sanes JR (2016)
Comprehensive Classification of Retinal Bipolar Neurons by Single-Cell Transcriptomics.
Cell 166:1308–1323.e1330.

Shen T, Aneas I, Sakabe N, Dirschinger RJ, Wang G, Smemo S, Westlund JM, Cheng H,
Dalton N, Gu Y, Boogerd CJ, Cai C-L, Peterson K, Chen J, Nobrega MA, Evans SM (2011)
Tbx20 regulates a genetic program essential to adult mouse cardiomyocyte function. J Clin
Invest 121:4640–4654.

Sheng W-L, Chen W-Y, Yang X-L, Zhong Y-M, Weng S-J (2015) Co-expression of two subtypes
of melatonin receptor on rat M1-type intrinsically photosensitive retinal ganglion cells.
Barnes S, ed. PLoS ONE 10:e0117967.

Shulga YV, Topham MK, Epand RM (2011) Regulation and Functions of Diacylglycerol Kinases.
Chemical Reviews 111:6186–6208.

Siegert S, Cabuy E, Scherf BG, Kohler H, Panda S, Le Y-Z, Fehling HJ, Gaidatzis D, Stadler
MB, Roska B (2012) Transcriptional code and disease map for adult retinal cell types.
Nature Neuroscience 15:487–495.

Soto I, Oglesby E, Buckingham BP, Son JL, Roberson EDO, Steele MR, Inman DM, Vetter ML,
Horner PJ, Marsh-Armstrong N (2008) Retinal Ganglion Cells Downregulate Gene
Expression and Lose Their Axons within the Optic Nerve Head in a Mouse Glaucoma
Model. J Neurosci 28:548–561.

Stabio ME, Sabbah S, Quattrochi LE, Ilardi MC, Fogerson PM, Leyrer ML, Kim MT, Kim I, Schiel
M, Renna JM, Briggman KL, Berson DM (2018) The M5 Cell: A Color-Opponent Intrinsically
Photosensitive Retinal Ganglion Cell. Neuron 97:251.

Star EN, Zhu M, Shi Z, Liu H, Pashmforoush M, Sauve Y, Bruneau BG, Chow RL (2012)
Regulation of retinal interneuron subtype identity by the Iroquois homeobox gene Irx6.
Development 139:4644–4655.

Stennard FA, Costa MW, Elliott DA, Rankin S, Haast SJP, Lai D, McDonald LPA, Niederreither
K, Dolle P, Bruneau BG, Zorn AM, Harvey RP (2003) Cardiac T-box factor Tbx20 directly
interacts with Nkx2-5, GATA4, and GATA5 in regulation of gene expression in the
developing heart. Developmental Biology 262:206–224.

Sweeney NT, Tierney H, Feldheim DA (2014) Tbr2 is required to generate a neural circuit
mediating the pupillary light reflex. J Neurosci 34:5447–5453.

Takeuchi JK, Mileikovskaia M, Koshiba-Takeuchi K, Heidt AB, Mori AD, Arruda EP,
Gertsenstein M, Georges R, Davidson L, Mo R, Hui C-C, Henkelman RM, Nemer M, Black
BL, Nagy A, Bruneau BG (2005) Tbx20 dose-dependently regulates transcription factor
networks required for mouse heart and motoneuron development. Development 132:2463–
2474.

Tariq MA, Kim HJ, Jejelowo O, Pourmand N (2011) Whole-transcriptome RNAseq analysis from
minute amount of total RNA. Nucleic Acids Res 39:e120–e120.

Thorvaldsdóttir H, Robinson JT (2013) Integrative Genomics Viewer (IGV): high-performance
genomics data visualization and exploration. Briefings in ….

Toki S, Kawasaki H, Tashiro N, Housman DE, Graybiel AM (2001) Guanine nucleotide
exchange factors CalDAG-GEFI and CalDAG-GEFII are colocalized in striatal projection
neurons. J Comp Neurol 437:398–407.

Topark-Ngarm A, Golonzhka O, Peterson VJ, Barrett B, Martinez B, Crofoot K, Filtz TM, Leid M
(2006) CTIP2 associates with the NuRD complex on the promoter of p57KIP2, a newly
identified CTIP2 target gene. J Biol Chem 281:32272–32283.

Topham MK, Prescott SM (2001) Diacylglycerol kinase zeta regulates Ras activation by a novel
mechanism. J Cell Biol 152:1135–1143.

Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL, Rinn JL,
Pachter L (2012) Differential gene and transcript expression analysis of RNA-seq
experiments with TopHat and Cufflinks. Nature Protocols 7:562–578.

Van Hook MJ, Wong KY, Berson DM (2012) Dopaminergic modulation of ganglion-cell
photoreceptors in rat. European Journal of Neuroscience 35:507–518.

Wong KY (2012) A retinal ganglion cell that can signal irradiance continuously for 10 hours. J
Neurosci 32:11478–11485.

Xu J, Xiao N, Xia J (2010) Thrombospondin 1 accelerates synaptogenesis in hippocampal
neurons through neuroligin 1. Nature Neuroscience 13:22–24.

Xue T, Do MTH, Riccio A, Jiang Z, Hsieh J, Wang HC, Merbs SL, Welsbie DS, Yoshioka T,
Weissgerber P, Stolz S, Flockerzi V, Freichel M, Simon MI, Clapham DE, Yau KW (2011)
Melanopsin signalling in mammalian iris and retina. Nature 479:67–73.

Young RW (1985) Cell differentiation in the retina of the mouse. The Anatomical Record
212:199–205.

Zhang M, Xia H, Li X, Wang X, Dong Y, Zhang T, Yu H (2010) C1 domain mediates CalDAGIII
localization to the Golgi. Mol Biol Rep 37:3481–3485.

Zhou H, Yoshioka T, Nathans J (1996) Retina-derived POU-domain factor-1: a complex POU-
domain gene implicated in the development of retinal ganglion and amacrine cells. J
Neurosci 16:2261–2274.

1706

Figure 1.

Figure 1—figure supplement 1.

Figure 2.

Figure 2—figure supplement 1.

Figure 2—figure supplement 2.

Figure 2—figure supplement 3

Figure 2—figure supplement 4.

Figure 3.

A. Opn4 Rasgrp1 Rbpms

B. Rasgrp1 Rbpms   Rasgrp1 Opn4

➜ **Rasgrp1+ ipRGC**
(Rasgrp1+ ; Rbpms+, Opn4+)

* **Rasgrp1+ amacrine**
(**Rasgrp1+** ; Rbpmsneg.)

▶ **Rasgrp1neg. ipRGC**
(Rasgrp1neg. ; Rbpms+ ; Opn4+)

C. Rasgrp1+/-   Rasgrp1-/-

Anti-Rasgrp1   Anti-Rasgrp1

D.

Fraction of ipRGC types that express Rasgrp1

M1/3   dM1   M2   M4   M5

Figure 3—figure supplement 1.

Figure 4.

Figure 4—figure supplement 1.

Figure 5.   **A.**

Figure 6.

Figure 7.

Circadian rhythm photoentrainment

Morphology	Mostly M1, rare M2

Molecular	Brn3b⁻

Rasgrp1 +/- Tbx20

Brain targets   SCN

Figure 8. A.

Tg *Opn4-GFP*

Opn4

merged

⇐ GFP ; anti-Opn4   ⇉ GFP$^{neg.}$; anti-Opn4   GFP$^+$; anti-Opn4$^{neg.}$

B.

Opn4-Cre/ZEG

Opn4

merged

⇒ GFP ; anti-Opn4   ⇉ GFP$^{neg.}$; anti-Opn4$^+$ * Opn4-GFP$^+$; anti-Opn4$^{neg.}$

C.

Percent ipRGCs expressing marker

- GFP
- Opn4
- Opn4 ; GFP

M1/3   dM1   M2   M4   M5/6

Figure 9.

A. Sample sort *Opn4::GFP* – Age P4

Count

7-AAD-A

~~G1~~

G4 G3 G2

Count

FITC-A

7-AAD⁺ dying cells

Debri

GFP⁻ RGCs

G3A

GFP⁺ M1+M2 ipRGCs

G2A

Rel. Cell Complexity

Relative Cell Size

B. Flow cytometry gating

- ■ All Events
  - ■ Gate 1 "G1" 7-AAD⁺ (apoptosis)
  - ■ NOT(G1) Low 7-AAD (viable)
    - ■ G2 GFP positive (ipRGCs)
      - ■ G2A Select similar RGC size
    - ■ G3 GFP⁻ (non-ipRGC RGCs)
      - ■ G3A Select similar RGC size
    - ■ G4 Cellular debris

C. Sample sort Adult *Opn4::GFP* (M1-M3 ipRGCs)

~~G1~~

G4 G3 G2

Count(x 1,000)

7-AAD-A

FITC-A

GFP⁻ RGCs

G3A P5

GFP⁺ M1+M2 ipRGCs

G2A

Rel. Cell Complexity

Relative Cell Size

D. Testing isolated cells from sorted *Opn4::Cre GFP* reporter (P4)

GFP DIC

50µm 100

Figure 10.

A. RNA extraction and quality test for degradation

Total RNA of P4 Opn4::GFP - high Quality RNA

B. cDNA processing of extracted RNA

C. RNA-seq library preparation (Truseq)

D. Test concentration and adapter length (qPCR & Bioanalyzer)

E. Multiplexed Illumina sequencing

Fully ligated templates participate in bridge amplification on flowcell surface

F.

Opn4-GFP (M1-M3 ipRGCs)				Opn4-Cre/ZEG (M1-M6 ipRGCs)			
raw reads	GFP+ sample	GFP- sample	raw reads	raw reads	GFP+ sample	GFP- sample	raw reads
95016286	DJB0007*	DJB0008*	90897302	109774976	DJB0005*	DJB0006*	96032720
44201103	DJB0030*	DJB0031*	43690608	98636762	DJB0022*	DJB0023*	110267207
45578821	DJB0028*	DJB0029*	49110355	64188702	DJB0032*	DJB0035*	68930442
49661171	DJB0016*	DJB0021*	47265858	21602002	DJB0040	DJB0041	22028459
10724088	DJB0036	DJB0037	11294631				
11377625	DJB0038	DJB0039	22789777	* concatenated technical replicates			